

Harnack inequality for stable processes on d -sets

by

KRZYSZTOF BOGDAN, ANDRZEJ STÓŚ and PAWEŁ SZTONYK (Wrocław)

Abstract. We investigate properties of functions which are harmonic with respect to α -stable processes on d -sets such as the Sierpiński gasket or carpet. We prove the Harnack inequality for such functions. For every process we estimate its transition density and harmonic measure of the ball. We prove continuity of the density of the harmonic measure. We also give some results on the decay rate of harmonic functions on regular subsets of the d -set. In the case of the Sierpiński gasket we even obtain the Boundary Harnack Principle.

1. Introduction. In the last two decades we observed a rapid development of analysis and probability theory on fractals; see [DSV], [Str], [Ba], [BB], [BP] and the references therein. [Ba], for example, presents probabilistic techniques in potential theory corresponding to the generator of the so-called fractional diffusion, the fractal analogue of the classical Brownian motion (see the next section for precise definitions).

Starting from the fractional diffusions, a class of *subordinated* processes on d -sets was introduced in [S]. By an analogy with the classical setting, we call these processes α -stable. The definition is briefly recalled in Section 3.

The present paper addresses several important problems of the potential theory of α -stable processes on d -sets. One of the results is the Harnack inequality for nonnegative harmonic functions of the process. The main results were announced earlier in [BSS]. For related recent developments we refer the reader to [FJ], [FU], [Kum], [HL].

The paper is organized as follows. In Section 3 we give estimates and some regularity results for the transition densities of the stable process. In Section 4 we obtain estimates for the expected exit time for subdomains of the d -set. We generally follow the approach designed in [Ba]; in particular

2000 *Mathematics Subject Classification*: Primary 60J45; Secondary 60J35.

Key words and phrases: stable processes, fractals, diffusions on fractals, Harnack inequality, Boundary Harnack Principle.

Research partially supported by KBN grant 2 P03A 041 22 and RTN Harmonic Analysis and Related Problems contract HPRN-CT-2001-00273-HARP.

Lemmas 4.1 and 4.3, and Proposition 4.4 have their analogues in the diffusion case. Here we give new and slightly generalized computations related to the α -stable process.

In Section 6 we prove the existence and joint continuity of the Poisson kernel $P_D(x, y)$ for an open ball D in the d -set. $P_D(x, y)$ is given by the Ikeda–Watanabe Formula [IW] describing a relation between the harmonic measure and the Lévy measure which is absent in the diffusion case. We derive optimal estimates for the Poisson kernel when x and y are away from the boundary (Proposition 6.4). The estimates turn out to be sufficient for the proof of the Harnack inequality when $\alpha \in (0, 2/d_w) \cup (d_s, 2)$. The latter is given in Section 7. In the recurrent case the proof of Lemma 7.3 employs an interesting formula (56) involving the hitting time for a point and the Green function. The transient case relies on estimates of the Poisson kernel for balls. We note here that the Harnack inequality has been recently established for fractional diffusions ([Ba], [BB1]) and for pure jump processes in \mathbb{R}^N ([BL]). In each case, including ours, the methods of proof are completely different.

Section 8 gives an estimate for the exit times of subdomains of the d -set (Theorem 8.3) which easily yields the decay rate of harmonic functions near the boundary (Theorem 8.4). The latter has an analogue in the theory of rotation invariant α -stable processes in \mathbb{R}^N (see [B, Lemma 3]). However, [B] makes an essential use of the exact formula for the Poisson kernel for a ball, which is not available in our case. Section 8 also contains a Carleson type estimate for $\alpha \in (0, 2/d_w)$ with a proof adapted from [BBy]. Our main contribution is in showing that the *weak scaling* of the process is sufficient for this proof to work. The restriction on α above is due to the fact that our proof depends on the polarity of the boundary of a ball. Finally, we give a proof of the Boundary Harnack Principle in the Sierpiński gasket case for $\alpha \in (0, 2/d_w) \cup (d_s, 2)$. Due to the simple geometry of this set the proof is an application of the Harnack inequality. We believe that the Boundary Harnack Principle holds more generally (e.g. for the Sierpiński carpet) but more complicated methods must be used to prove it.

2. Preliminaries. In this section we collect some notation and definitions adapted from [Ba] and [P].

Let F be a nonempty closed subset of \mathbb{R}^N , $N \geq 1$. Set $d \in (0, N]$. We say that a (positive) Borel measure μ is a d -measure on F if for some constants $c_1, c_2 > 0$ it satisfies

$$(1) \quad c_1 r^d \leq \mu(B(x, r)) \leq c_2 r^d, \quad x \in F, \quad 0 < r \leq r_0,$$

where r_0 is the diameter of F and $B(x, r)$ denotes the ball in \mathbb{R}^N with center x and radius r . We call F a d -set if $F = \text{supp}(\mu)$ for some d -measure μ . It is known that any d -measure is a regular Borel measure. Any two d -measures

on the same d -set F are equivalent and the d -dimensional Hausdorff measure restricted to a d -set F is a d -measure [JW].

We use c (with subscripts) to denote positive and finite constants which depend only on the d -measure μ , F (and d), the fractional diffusion on F and the stability index α (see below). Any additional dependence is indicated explicitly, e.g. $c_4 = c_4(D, \kappa)$. Constants are numbered consecutively within each proof. The value of c (without subscript) may change from place to place. We write (e.g.) $f(x) \asymp g(x)$, $x \in F$, to indicate that there are constants $c_1, c_2 > 0$ (independent of x) such that $c_1 f(x) \leq g(x) \leq c_2 f(x)$ for all $x \in F$. We denote by $|\cdot|$ the Euclidean distance in \mathbb{R}^N . From now on $B(x, r)$ denotes the *Euclidean* ball intersected with our d -set F . For a subset $D \subseteq F$ we always take complements in F , i.e. $D^c = F \setminus D$. Without losing generality, in what follows we assume that $0 \in F$, which often simplifies the notation.

The following lemma is a convenient replacement for integration in polar coordinates.

LEMMA 2.1. *Let F and μ be as introduced above. Then for every $x_0 \in F$, $r > 0$ and $\lambda > 0$ there is $c = c(\lambda)$ such that*

$$(2) \quad \int_{|x-x_0|>r} |x-x_0|^{-d-\lambda} d\mu(x) \leq cr^{-\lambda},$$

$$(3) \quad \int_{|x-x_0|<r} |x-x_0|^{-d+\lambda} d\mu(x) \leq cr^\lambda.$$

Proof. From (1) it follows that

$$\begin{aligned} \int_{|x-x_0|>r} |x-x_0|^{-d-\lambda} d\mu(x) &= \sum_{n=0}^{\infty} \int_{2^n r < |x-x_0| \leq 2^{n+1} r} |x-x_0|^{-d-\lambda} d\mu(x) \\ &\leq \sum_{n=0}^{\infty} (2^n r)^{-d-\lambda} \mu(B(x_0, 2^{n+1} r)) \\ &\leq cr^{-d-\lambda} \sum_{n=0}^{\infty} 2^{-n(d+\lambda)} (2^{n+1} r)^d \leq cr^{-\lambda}. \end{aligned}$$

This proves (2). The estimate (3) follows in a similar way. ■

From now on we let $F \subset \mathbb{R}^N$ be a connected d -set, $d \in (0, N]$, $N \geq 2$, and we let μ be its d -measure. We put $t_0 = \infty$ when $r_0 = \infty$ and $t_0 = r_0^{d_w}$ otherwise (see Definition 2.1). We often refer to the general theory of Markov processes in the setting of [BG] or [ChZ].

DEFINITION 2.1. A Markov process $X = (P^x, X_t)_{x \in F, t \geq 0}$ is called a *fractional diffusion on F* if

- (a) X is a diffusion with state space F ,

(b) X has a symmetric transition density $q(t, x, y) = q(t, y, x)$, $t > 0$, $x, y \in F$, which is jointly continuous for each $t > 0$ and satisfies, for some constants $c_1, \dots, c_4 > 0$, $d_w > 1$ and all $x, y \in F$ and $t \in (0, t_0)$,

$$(4) \quad c_1 t^{-d_s/2} \exp\left(-c_2 \left(\frac{|x-y|}{t^{1/d_w}}\right)^\gamma\right) \leq q(t, x, y) \leq c_3 t^{-d_s/2} \exp\left(-c_4 \left(\frac{|x-y|}{t^{1/d_w}}\right)^\gamma\right).$$

Here $d_s = 2d/d_w$, $\gamma = d_w/(d_w - 1)$.

It is also known (see [P]) that all fractional diffusions on a fixed d -set F must have the same value of the constant d_w , i.e. d_w depends only on the underlying geometry. We have $d_w = 2$ for $F = \mathbb{R}^N$ and if the heat kernel of the diffusion satisfies (4) then $d_w \geq 2$ (see [G]). We note here that the above definition differs from that given in [Ba]. Following [P] we use the Euclidean distance instead of the intrinsic shortest path metric (see [Ba]). Since the well known fractal diffusions were constructed in the shortest path metric setting (e.g. [Ba]), (4) is virtually tantamount to the assumption that the two metrics are equivalent.

3. Stable process. From now on we fix $\alpha \in (0, 2)$. We also assume that F is a *connected* d -set with d -measure μ and $r_0 = \infty$ in (1). In particular $d \geq 1$ and F is necessarily unbounded. We briefly recall the construction of the α -stable process from [S]. Suppose that there exists a fractional diffusion on F and let $q(u, x, y)$, $u > 0$, $x, y \in F$, denote its transition density with respect to μ . Let $(Y_t)_{t>0}$ be the $\alpha/2$ -stable subordinator given by the Laplace transform $E \exp(-uY_t) = \exp(-tu^{\alpha/2})$. Let $\eta_t(u)$, $t > 0$, $u \geq 0$, be its one-dimensional distribution density (see [Be] or [BG] for more details). For $t > 0$ and $x, y \in F$ we define

$$p(t, x, y) = \int_0^\infty q(u, x, y) \eta_t(u) du.$$

By the general theory $p(t, x, y)$ is the transition density of a Markov process called the subordinate process (see [BG, p. 18]), which we denote by $(X_t)_{t>0}$ and call α -stable.

To simplify the notation, for the rest of the paper we let $d_\alpha = d + \alpha d_w/2$. The main result of this section is the theorem below. It resembles a well known estimate for the rotation invariant α -stable process on \mathbb{R}^N and can be interpreted as *weak scaling* of our process.

THEOREM 3.1 (Weak scaling). *For $t > 0$, $x, y \in F$, $x \neq y$, we have*

$$(5) \quad p(t, x, y) \asymp \min\left(\frac{t}{|x-y|^{d_\alpha}}, t^{-d_s/\alpha}\right),$$

in particular

$$(6) \quad c_1 t^{-d_s/\alpha} \leq p(t, x, x) \leq c_2 t^{-d_s/\alpha}.$$

Proof. By Theorem 37.1 of [D],

$$(7) \quad \lim_{u \rightarrow \infty} \eta_1(u) u^{1+\alpha/2} = \alpha / (2\Gamma(1 - \alpha/2)).$$

This, boundedness of $\eta_1(\cdot)$ and the scaling property

$$(8) \quad \eta_t(u) = t^{-2/\alpha} \eta_1(t^{-2/\alpha} u), \quad t, u > 0,$$

yield the following estimates:

$$(9) \quad \eta_t(u) \leq c_3 t u^{-1-\alpha/2}, \quad t, u > 0,$$

$$(10) \quad \eta_t(u) \geq c_4 t u^{-1-\alpha/2}, \quad t > 0, u > u_0 t^{2/\alpha},$$

where u_0 depends only on α . Let $t > 0, x, y \in F, x \neq y$, and $d(t, x, y) = |x - y|^\gamma t^{-2/(\alpha(d_w-1))}$. By the definition of $p(t, x, y)$, (10) and substitution $v = c_2 |x - y|^\gamma u^{-1/(d_w-1)}$ (where c_2 is defined in (4)) we get

$$(11) \quad \begin{aligned} p(t, x, y) &\geq ct \int_{u_0 t^{2/\alpha}}^{\infty} u^{-d_s/2} \exp(-c_2 |x - y|^\gamma u^{-1/(d_w-1)}) u^{-1-\alpha/2} du \\ &\geq ct |x - y|^{-d_\alpha} \int_0^{c_5 d(t,x,y)} v^{(d_s+\alpha)(d_w-1)/2-1} e^{-v} dv \\ &\geq ct |x - y|^{-d_\alpha} e^{-c_5 d(t,x,y)} \int_0^{c_5 d(t,x,y)} v^{(d_s+\alpha)(d_w-1)/2-1} dv \\ &= ct^{-d_s/\alpha} e^{-c_5 d(t,x,y)}. \end{aligned}$$

If $t \geq |x - y|^{\alpha d_w/2}$ or $|x - y| t^{-2/(\alpha d_w)} \leq 1$, then $d(t, x, y) \leq 1$ and $\exp(-c_5 d(t, x, y)) \geq \exp(-c_5)$. Consequently, we then get

$$(12) \quad p(t, x, y) \geq c_6 t^{-d_s/\alpha}.$$

Thus

$$p(t, x, y) \geq c_6 \min\left(\frac{t}{|x - y|^{d_\alpha}}, t^{-d_s/\alpha}\right)$$

in this case. On the other hand, if $t < |x - y|^{\alpha d_w/2}$ then $d(t, x, y) > 1$ so that the integral in (11) is bounded away from 0 and $p(t, x, y) \geq c_7 t |x - y|^{-d_\alpha} \geq c_7 \min(t/|x - y|^{d_\alpha}, t^{-d_s/\alpha})$ again. By (4), (9) and the same substitution as in (11),

$$(13) \quad \begin{aligned} p(t, x, y) &\leq c_5 t \int_0^{\infty} u^{-d_s/2-\alpha/2-1} \exp(-c_6 |x - y|^\gamma u^{-1/(d_w-1)}) du \\ &= \frac{ct}{|x - y|^{d_\alpha}} \Gamma((d_s + \alpha)(d_w - 1)/2). \end{aligned}$$

This gives the upper bound of (5) with the first term under the minimum.

To finish the proof of (5) we will verify the estimate

$$(14) \quad \eta_t(u) \leq ctu^{-1-\alpha/2} \exp(-tu^{-\alpha/2}), \quad u > 0, t > 0.$$

Indeed, from Lemma 1 in [H] (see also the proof therein) we have

$$\eta_1(u) \leq cu^{(\alpha-4)/(4-2\alpha)} \exp(-c_8u^{-\alpha/(2-\alpha)})$$

as $u \rightarrow 0$. Since $\alpha/(2-\alpha) > \alpha/2$, we clearly have

$$u^{(\alpha-4)/(4-2\alpha)} e^{-c_8u^{-\alpha/(2-\alpha)}} = o(u^{-1-\alpha/2} e^{-u^{-\alpha/2}}) \quad \text{as } u \rightarrow 0.$$

From this and (9),

$$\eta_1(u) \leq cu^{-1-\alpha/2} \exp(-u^{-\alpha/2}), \quad u > 0.$$

(14) follows from this and (8). By (14) and the substitution $v = tu^{-\alpha/2}$, for any $x, y \in F$ we obtain

$$(15) \quad \begin{aligned} p(t, x, y) &\leq ct \int_0^\infty u^{-d_s/2} u^{-1-\alpha/2} \exp(-tu^{-\alpha/2}) du \\ &= ct^{-d_s/\alpha} \int_0^\infty v^{d_s/\alpha} e^{-v} dv = ct^{-d_s/\alpha}. \end{aligned}$$

This completes the proof of (5) and also gives the upper bound in (6). The lower bound in (6) follows from (5) by continuity; see Proposition 3.2 below. ■

REMARK 1. Note that

$$(16) \quad t \leq |x - y|^{\alpha d_w/2} \quad \text{if and only if} \quad \frac{t}{|x - y|^{d_\alpha}} \leq t^{-d_s/\alpha}.$$

Therefore, Theorem 3.1 can be reformulated in the following way:

$$p(t, x, y) \asymp t^{-d_s/\alpha} \varphi(|x - y|t^{-2/(\alpha d_w)}), \quad t > 0, x, y \in F,$$

with $\varphi(u) = (1 \vee |u|)^{-d_\alpha} \asymp (1 + |u|)^{-d_\alpha}$.

REMARK 2. For later convenience we note that given $\beta \in (0, 1)$ and $t \geq \beta|x - y|^{\alpha d_w/2}$, the inequality (12) still holds true with some $c_6 = c_6(\beta)$.

Consider a Markov process with state space being an open set $D \subseteq F$ and transition probability semigroup $P_t(x, E)$, $t > 0$, $x \in D$, where E is a Borel subset of D . We say that the semigroup has the *strong Feller property* if $P_t(L^\infty(D, \mu)) \subseteq C_b(D)$, $t > 0$, where $C_b(D)$ stands for the continuous bounded functions on D . By $C_0(D)$ we denote the set of continuous bounded functions on D that tend to zero at the boundary (and also when $x \rightarrow \infty$ if D is unbounded). The semigroup is said to be *strongly continuous* on a function space S if

$$\lim_{t \rightarrow 0} \|P_t f - f\| = 0, \quad f \in S,$$

where the norm is taken in S . In what follows, S will be equal to $C_0(D)$ or $L^p(D)$, $p \in [1, \infty)$. The semigroup is said to have the *Feller property* if $P_t(C_0(D)) \subseteq C_0(D)$ and P_t is strongly continuous on $C_0(D)$.

We now return to the study of our stable process on F .

PROPOSITION 3.2. (i) *The transition density $p(t, x, y)$ is jointly continuous in $(t, x, y) \in (0, \infty) \times F \times F$ for each $t > 0$.*

(ii) *The transition semigroup (P_t) generated by our stable process has both the Feller and strong Feller properties. In particular, (P_t) is strongly continuous on $C_0(F)$.*

Proof. One applies the same arguments as for the Brownian motion on \mathbb{R}^N using properties of the underlying fractional diffusion and the dominated convergence theorem together with the upper bounds from (5), (6) and (14). ■

By virtue of (ii) in the above proposition we may and do assume that path functions of our stable process are right-continuous with left hand limits and that the process is quasi-left-continuous (see [BG]).

4. Exit time. Let E be a Borel subset of F . We let $T_E = \inf\{t \geq 0 : X_t \in E\}$ and $\tau_E = T_{E^c}$.

LEMMA 4.1. *Let $\kappa > 1$. There exist constants c_1 and c_2 such that for every $x \in F$ and $r > 0, t > 0$,*

$$(17) \quad P^x[X_t \notin B(x, r)] \geq c_2 t r^{-\alpha d_w/2}, \quad 0 < t < r^{\alpha d_w/2},$$

and

$$(18) \quad P^y[X_t \notin B(x, r)] \leq c_1 (\kappa/(\kappa - 1))^{\alpha d_w/2} t r^{-\alpha d_w/2},$$

for all $y \in B(x, r/\kappa)$.

Proof. Let $a > 1$ and let x, r, t be as in (17). If $|y - x| \geq r$ then $|y - x|^{\alpha d_w/2} \geq t$, and by (5) and (16) we have

$$\begin{aligned} P^x[X_t \notin B(x, r)] &\geq \int_{r \leq |y-x| < ar} p(t, x, y) d\mu(y) \geq ct \int_{r \leq |y-x| < ar} \frac{d\mu(y)}{|y-x|^{d_\alpha}} \\ &\geq c t r^{-d_\alpha} a^{-d_\alpha} \mu(B(x, ar) \setminus B(x, r)) \\ &\geq c t r^{-\alpha d_w/2} a^{-d_\alpha} (c_1 a^d - c_2), \end{aligned}$$

where the constants c_1, c_2 come from (1). We can choose a large enough to make the last factor positive and (17) follows.

We now fix $\kappa > 1, x \in F, t > 0$, and $r > 0$. Let $r_0 = r(\kappa - 1)/\kappa$. Observe that if $y \in B(x, r/\kappa)$ then $B(y, r_0) \subseteq B(x, r)$. By Theorem 3.1 and (2) we

obtain

$$\begin{aligned} P^y[X_t \notin B(x, r)] &\leq P^y[X_t \notin B(y, r_0)] = \int_{B(y, r_0)^c} p(t, y, z) d\mu(z) \\ &\leq ct \int_{B(y, r_0)^c} |z - y|^{-d_\alpha} d\mu(z) \leq ctr_0^{-\alpha d_w/2}, \end{aligned}$$

and (18) follows. ■

The following simple lemma will be used without further mention (see [ChZ, Proposition 1.20]).

LEMMA 4.2. *Let B be a Borel set in F . For each $t > 0$ and $x \in F$ we have $P^x[\tau_B = t] = 0$.*

LEMMA 4.3. *For each $\kappa > 1$ there exists $c_1 = c_1(\kappa)$ such that for $x \in F$, $r > 0$, $y \in B(x, r/\kappa)$ we have*

$$P^y[\tau_{B(x, r)} < t] \leq c_1 tr^{-\alpha d_w/2}.$$

Proof. Let $1 < \kappa_1 < \kappa$ and $\lambda > 1$ be such that $\kappa_1 \lambda = \kappa$ (e.g. $\kappa_1 = \lambda = \sqrt{\kappa}$). Define $T = \tau_{B(x, \lambda r)}$. For $y \in B(x, r/\kappa_1)$ we have

$$\begin{aligned} P^y[T < t] &= P^y[X_t \notin B(x, r); T < t] + P^y[X_t \in B(x, r); T < t] \\ &\leq P^y[X_t \notin B(x, r)] + P^y[X_t \in B(x, r); T < t] = A + B. \end{aligned}$$

By (18), we obtain $A \leq c_2 tr^{-\alpha d_w/2}$ with $c_2 = c_2(\kappa_1)$.

By the strong Markov property we have

$$(19) \quad B = E^y[P^{X(T)}[X_{t-u} \in B(x, r)]|_{u=T}; T < t].$$

We now estimate the integrand in (19):

$$\begin{aligned} P^{X(T)}[X_{t-u} \in B(x, r)]|_{u=T} &\leq \sup_{z \in B(x, \lambda r)^c} P^z[X_{t-u} \in B(x, r)]|_{u=T} \\ &\leq \sup_{u \leq t} \sup_{z \in B(x, \lambda r)^c} P^z[X_u \in B(x, r)] \\ &\leq \sup_{u \leq t} \sup_{z \in B(x, \lambda r)^c} P^z[X_u \notin B(z, (\lambda - 1)r)] \\ &\leq c_3 tr^{-\alpha d_w/2}, \end{aligned}$$

where $c_3 = c_3(\lambda)$. Consequently,

$$(20) \quad P^y[T < t] \leq c_4 tr^{-\alpha d_w/2}, \quad y \in B(x, r/\kappa_1),$$

where $c_4 = c_4(\kappa)$. Apply (20) to r/λ instead of r and the assertion follows. ■

PROPOSITION 4.4. *There exists c_1 such that for $x \in F$, $r > 0$,*

$$(21) \quad \sup_{y \in B(x, r)} E^y \tau_{B(x, r)} \leq c_1 r^{\alpha d_w/2},$$

and for each $\kappa > 1$ there exists a constant $c_2 = c_2(\kappa) > 0$ such that

$$(22) \quad E^y \tau_{B(x,r)} \geq c_2(\kappa) r^{\alpha d_w/2}, \quad x \in F, y \in B(x, r/\kappa).$$

Proof. For each $y \in B(x, r)$, from (5) we have

$$\begin{aligned} P^y[\tau_{B(x,r)} > t] &\leq P^y[X_t \in B(x, r)] = \int_{B(x,r)} p(t, y, z) d\mu(z) \\ &\leq ct^{-d_s/\alpha} \mu(B(x, r)) \leq cr^d t^{-d_s/\alpha}. \end{aligned}$$

Hence,

$$P^y[\tau_{B(x,r)} > c_3 r^{\alpha d_w/2}] \leq cr^d c_3^{-d_s/\alpha} (r^{\alpha d_w/2})^{-d_s/\alpha} = 1/2$$

for a suitably chosen value of c_3 . Let $t_0 = c_3 r^{\alpha d_w/2}$. Then, by the Markov property, for $k = 1, 2, \dots$ we have

$$\begin{aligned} P^y[\tau_{B(x,r)} > (k+1)t_0] &= P^y[\tau_{B(x,r)} \circ \theta_{t_0} > kt_0, \tau_{B(x,r)} > t_0] \\ &= E^y[P^{X(t_0)}[\tau_{B(x,r)} > kt_0]; \tau_{B(x,r)} > t_0] \\ &\leq P^y[\tau_{B(x,r)} > t_0] \sup_{z \in B(x,r)} P^z[\tau_{B(x,r)} > kt_0] \end{aligned}$$

(here θ stands for the standard shift operator on the space of trajectories). By induction we get

$$P^y[\tau_{B(x,r)} > kt_0] \leq (1/2)^k, \quad y \in B(x, r), k = 0, 1, \dots$$

Thus,

$$E^y \tau_{B(x,r)} = \int_0^\infty P^y[\tau_{B(x,r)} > t] dt \leq \sum_{k=0}^\infty t_0 P^y[\tau_{B(x,r)} > kt_0] = 2c_3 r^{\alpha d_w/2},$$

which gives (21).

By Lemma 4.3 there exists $c_4 = c_4(\kappa)$ such that

$$P^y[\tau_{B(x,r)} < c_4 r^{\alpha d_w/2}] \leq 1/2, \quad y \in B(x, r/\kappa).$$

It follows that for $t_0 = c_4 r^{\alpha d_w/2}$ we have

$$E^y \tau_{B(x,r)} \geq t_0 P^y[\tau_{B(x,r)} > t_0] \geq (1/2) c_4 r^{\alpha d_w/2}, \quad y \in B(x, r/\kappa).$$

The proof is complete. ■

5. Killed process and Green function. Let $D \subseteq F$ be an open set. By (P_t^D) we denote the semigroup generated by the process killed on exiting D , that is (see [BG]),

$$P_t^D f(x) = E^x[f(X_t); t < \tau_D],$$

for, e.g., nonnegative or bounded Borel functions f on F .

The following proposition summarizes properties of P_t^D .

PROPOSITION 5.1. *Let $D \subseteq F$ be an open bounded set.*

- (i) *The semigroup (P_t^D) has both Feller and strong Feller properties.*
- (ii) *The semigroup is determined by a transition density $p^D(t, x, y)$, i.e. for any nonnegative or bounded Borel function f on F we have*

$$P_t^D f(x) = \int_D f(y)p^D(t, x, y) d\mu(y), \quad x \in F.$$

Moreover, for each $t > 0$, $p^D(t, x, y)$ is continuous in $(x, y) \in D \times D$ and

$$(23) \quad \begin{aligned} p^D(t, x, y) &= p^D(t, y, x), \quad x, y \in F \setminus \partial D, t > 0, \\ p^D(t, x, y) &= p(t, x, y) - r^D(t, x, y), \quad x, y \in F, t > 0, \end{aligned}$$

with $r^D(t, x, y) = E^x[p(t - \tau_D, X_{\tau_D}, y); t > \tau_D];$

$$(24) \quad p^D(t, x, y) > 0, \quad x, y \in D, t > 0;$$

$$(25) \quad p^D(t, x, y) = \int_F p^D(s, x, z)p^D(t - s, z, y) d\mu(z), \quad t > s > 0, x, y \in F;$$

$$p^D(t, x, y) = 0, \quad x \in D^c.$$

Proof. The standard arguments that can be found e.g. in [ChZ] (see also [Bs]) work also in the present setting with the exception of (24). We give a proof of (24) similar to but more direct than the one in [CS].

Let $K \subseteq D$ be a compact set and let $x, y \in K$. Define $\varrho = \text{dist}(K, D^c)$. By (5) and Lemma 4.3, we have

$$\begin{aligned} r^D(t, x, y) &= E^x[p(t - \tau_D, X_{\tau_D}, y); t > \tau_D] \\ &\leq cE^x[(t - \tau_D)|X_{\tau_D} - y|^{-d_\alpha}; t > \tau_D] \\ &\leq ct\varrho^{-d_\alpha} P^x[\tau_D < t] \leq ct\varrho^{-d_\alpha} P^x[\tau_{B(x, \varrho)} < t] \leq ct^2\varrho^{-d - \alpha d_w}. \end{aligned}$$

Also, by Theorem 3.1,

$$p(t, x, y)^{-1} \leq c \max(t^{d_s/\alpha}, t^{-1}|x - y|^{d_\alpha}),$$

so that

$$\frac{r^D(t, x, y)}{p(t, x, y)} \leq c \max(t^{2+d_s/\alpha}, t|x - y|^{d_\alpha})\varrho^{-d - \alpha d_w} \leq c_1 \max(t^{2+d_s/\alpha}, t),$$

where $c_1 = c_1(D, K)$. It follows from (23) that for $t < t_0 = t_0(K)$ and $x, y \in K$,

$$(26) \quad p^D(t, x, y) > \frac{1}{2}p(t, x, y) > 0.$$

Fix arbitrary $x_0, y_0 \in D$. Let $r > 0$ be such that $K = \overline{B(x_0, r) \cup B(y_0, r)} \subseteq D$. We then use $t_0 = t_0(K)$ as above, (26) and the semigroup property to obtain

$$p^D(t, x, y) \geq \int_K p^D(t/2, x, z)p^D(t/2, z, y) d\mu(z) > 0$$

for $x, y \in K$ and $t < 2t_0$. Similarly, by induction

$$p^D(t, x, y) > 0, \quad x, y \in K, \quad t < nt_0, \quad n = 1, 2, \dots,$$

and (24) follows. ■

The estimate (27) below is taken from [R] (Lemma 6). For the reader's convenience we include a version of the proof.

LEMMA 5.2. *We have*

$$(27) \quad p^D(t, x, y) \leq ct^{-1-d_s/\alpha} r^{\alpha d_w/2}, \quad x, y \in F, \quad t > 0.$$

Proof. From the semigroup property (25) and the estimate (5) we have

$$\begin{aligned} p^D(t, x, y) &= \int_D p^D(t/2, x, z) p^D(t/2, z, y) d\mu(z) \\ &\leq \sup_{z \in D} p^D(t/2, z, y) \int_D p^D(t/2, x, z) d\mu(z) \\ &\leq ct^{-d_s/\alpha} P^x[\tau_D > t/2]. \end{aligned}$$

Hence, by the elementary inequality

$$P^x[\tau_D > t/2] \leq 2E^x \tau_D/t$$

and (21) we get the assertion. ■

Let $D \subseteq F$ be an open set. We define the Green function of D by

$$(28) \quad G_D(x, y) = \int_0^\infty p^D(t, x, y) dt.$$

Proposition 5.1 implies that if D is an open bounded set in F then $G_D(x, y)$ has the following properties: it is symmetric and strictly positive on $D \times D$; it vanishes if $x \in D^c$ or, by the symmetry of $p^D(t, x, y)$, $y \in \text{int}(D^c)$. G_D is extended continuous on $D \times D$ for $\alpha \leq d_s$ as can be verified by an adaptation of the corresponding proof from [ChZ] ($G_D(x, x) = \infty$ by (26)). To show the continuity for $\alpha > d_s$ we note that by (6) and a version of (27),

$$(29) \quad p^D(t, x, y) \leq c \min(t^{-d_s/\alpha}, t^{-1-d_s/\alpha}), \quad t > 0, \quad x, y \in F.$$

Since this is integrable over $(0, \infty)$, the desired assertion follows from the bounded convergence theorem. We leave the details to the interested reader.

Here we give an expression for the potential kernel of the stable process X_t .

LEMMA 5.3. *If $\alpha < d_s$ then*

$$(30) \quad K_\alpha(x, y) := \int_0^\infty p(t, x, y) dt \asymp |x - y|^{-d+\alpha d_w/2}, \quad x, y \in F.$$

Proof. By Theorem 3.1 and (16),

$$(31) \quad \int_0^\infty p(t, x, y) dt \asymp |x - y|^{-d-\alpha d_w/2} \int_0^{|x-y|^{\alpha d_w/2}} t dt + \int_{|x-y|^{\alpha d_w/2}}^\infty t^{-d_s/\alpha} dt$$

$$(32) \quad \asymp |x - y|^{-d+\alpha d_w/2}. \blacksquare$$

COROLLARY 5.4. *The process is transient if and only if $\alpha < d_s$.*

LEMMA 5.5. *If $\alpha < d_s$ then for any open bounded $D \subseteq F$,*

$$(33) \quad G_D(x, y) = K_\alpha(x, y) - E^x K_\alpha(X_{\tau_D}, y)$$

(unless $x = y \in D^c$) and

$$(34) \quad G_D(x, y) \leq K_\alpha(x, y) \asymp |x - y|^{-d+\alpha d_w/2}, \quad x, y \in F.$$

Proof. From (23) by a simple change of variable we obtain

$$\begin{aligned} G_D(x, y) &= \int_0^\infty p(t, x, y) dt - E^x \int_{\tau_D}^\infty p(t - \tau_D, X_{\tau_D}, y) dt \\ &= \int_0^\infty p(t, x, y) dt - E^x \int_0^\infty p(t, X_{\tau_D}, y) dt, \end{aligned}$$

which is clearly (33).

Now, (34) follows immediately from (33) and Lemma 5.3. \blacksquare

PROPOSITION 5.6. *If $\alpha > d_s$ then*

$$G_{B(x,r)}(x, y) \leq cr^{-d+\alpha d_w/2}, \quad x, y \in F, r > 0.$$

Proof. Define $D = B(x, r)$. We have

$$\int_0^{r^{\alpha d_w/2}} p^D(t, x, y) dt \leq \int_0^{r^{\alpha d_w/2}} p(t, x, y) dt \leq c \int_0^{r^{\alpha d_w/2}} t^{-d_s/\alpha} dt.$$

Since $\alpha > d_s$, we get

$$(35) \quad \int_0^{r^{\alpha d_w/2}} p^D(t, x, y) dt \leq c_1 r^{-d+\alpha d_w/2}, \quad x, y \in F.$$

From (27) it follows that

$$(36) \quad \int_{r^{\alpha d_w/2}}^\infty p^D(t, x, y) dt \leq cr^{\alpha d_w/2} \int_{r^{\alpha d_w/2}}^\infty t^{-1-d_s/\alpha} dt = cr^{-d+\alpha d_w/2}.$$

Now, the assertion follows from (35), (36) and the definition of the Green function. \blacksquare

Let u be a Borel measurable function u on F , which is bounded from below (above). We say that u is α -harmonic in an open set $U \subseteq F$ if

$$u(x) = E^x u(X(\tau_B)), \quad x \in B,$$

for every bounded open set B with the closure \bar{B} contained in U . We say that u is regular α -harmonic in U if

$$u(x) = E^x u(X(\tau_U)), \quad x \in U.$$

By the strong Markov property of X , regular α -harmonic functions are α -harmonic. We give an elementary proof of α -harmonicity of $G_D(x, y)$.

PROPOSITION 5.7. *Let D be an open bounded set in F and $\alpha \neq d_s$. Then $G_D(\cdot, y)$ is α -harmonic in $D \setminus \{y\}$ for any $y \in D$.*

Proof. Fix $y \in D$ and let U be an arbitrary open set with $\bar{U} \subseteq D \setminus \{y\}$. (In fact it is enough to assume $U \subseteq D$ and $\text{dist}(U, y) > 0$; we will use this later.) For $x \in U$ and a nonnegative Borel measurable function ψ supported in U^c we have

$$\begin{aligned} (37) \quad E^x \int_0^{\tau_D} \psi(X_t) dt &= E^x \int_0^{\tau_U} \psi(X_t) dt + E^x \left[\int_{\tau_U}^{\tau_D} \psi(X_t) dt; \tau_U < \tau_D \right] \\ &= E^x \left[E^{X_{\tau_U}} \int_0^{\tau_D} \psi(X_t) dt; \tau_U < \tau_D \right], \end{aligned}$$

by the strong Markov property. In terms of the Green function of D this is

$$\begin{aligned} (38) \quad \int_D G_D(x, z) \psi(z) d\mu(z) &= E^x \left[\int_D G_D(X_{\tau_U}, z) \psi(z) d\mu(z); \tau_U < \tau_D \right] \\ &= \int_D E^x [G_D(X_{\tau_U}, z); \tau_U < \tau_D] \psi(z) d\mu(z). \end{aligned}$$

For almost all $z \in D \cap U^c$ (with respect to μ) we obtain

$$G_D(x, z) = E^x [G_D(X_{\tau_U}, z); \tau_U < \tau_D].$$

Now, let $\alpha > d_s$. The continuity of $G_D(x, \cdot)$ and boundedness over $D \cap U^c$ (see Proposition 5.6) yield

$$(39) \quad G_D(x, y) = E^x [G_D(X_{\tau_U}, y); \tau_U < \tau_D].$$

On the set $\{\tau_U = \tau_D\}$ we have $X_{\tau_U} \in D^c$ and hence $G_D(X_{\tau_U}, y) = 0$. Hence, from (39) it follows that

$$(40) \quad E^x G_D(X_{\tau_U}, y) = E^x [G_D(X_{\tau_U}, y); \tau_U < \tau_D] = G_D(x, y),$$

which completes the proof for $\alpha > d_s$.

If $\alpha < d_s$ then $K_\alpha(\cdot, y)$ is α -harmonic in $F \setminus \{y\}$. Indeed, by (33) applied to an open bounded set $V \subseteq F \setminus \{y\}$ with $\text{dist}(y, V) > 0$, we have

$$E^x K_\alpha(X_{\tau_V}, y) = K_\alpha(x, y) - G_V(x, y).$$

Since $y \in \text{int}(V^c)$, we get $G_V(x, y) = 0$ and consequently

$$E^x K_\alpha(X_{\tau_V}, y) = K_\alpha(x, y),$$

which gives our claim. Now, the assertion for $\alpha < d_s$ follows from (33) and the strong Markov property. ■

REMARK. It is also possible to derive (39) for $\alpha < d_s$ by modifying the method just applied for the case $\alpha > d_s$. Indeed, we take into account uniform integrability (see (34)) of the family $\{G_D(w, \tilde{y})\mathbf{1}_{B(y, \delta)}(w) : \tilde{y} \in B(y, \delta)\}$, where $\delta > 0$ is such that $B(y, 2\delta) \subseteq U^c$, and estimates of the Poisson kernel for U near y similar to those given in Section 6.

LEMMA 5.8. *There exist $a > 1$ and c such that for all $x, y \in F$,*

$$(41) \quad G_D(x, y) \geq c|x - y|^{-d+\alpha d_w/2},$$

where $D = B(x, a|x - y|)$.

Proof. Let $a > 1$. Define $r = |x - y|$. By the definition of $G_D(x, y)$, Theorem 3.1 and (16) we obtain

$$\begin{aligned} G_D(x, y) &\geq \int_0^{|x-y|^{\alpha d_w/2}} (p(t, x, y) - E^x[p(t - \tau_D, X_{\tau_D}, y); \tau_D < t]) dt \\ &\geq \int_0^{|x-y|^{\alpha d_w/2}} (c_1 t|x - y|^{-d_\alpha} - E^x[c_2(t - \tau_D)|X_{\tau_D} - y|^{-d_\alpha}; \tau_D < t]) dt, \end{aligned}$$

where c_1 and c_2 are defined by the lower and upper bound in (5), respectively. Observe that $|X_{\tau_D} - y| \geq (a - 1)|x - y|$. Now, let $a > 1$ be such that $c_2(a - 1)^{-d_\alpha} \leq c_1/2$. It follows that

$$\begin{aligned} G_D(x, y) &\geq \int_0^{|x-y|^{\alpha d_w/2}} (c_1 t|x - y|^{-d_\alpha} - c_2 t((a - 1)|x - y|)^{-d_\alpha}) dt \\ &\geq (c_1/2)|x - y|^{-d_\alpha} \int_0^{|x-y|^{\alpha d_w/2}} t dt = (c_1/4)|x - y|^{-d+\alpha d_w/2}, \end{aligned}$$

which gives (41). ■

6. Harmonic measure. For $x, y \in F$ define

$$(42) \quad N(x, y) = \lim_{t \rightarrow 0} \frac{p(t, x, y)}{t} = \lim_{t \rightarrow 0} \int_0^\infty q(u, x, y) \frac{\eta_t(u)}{t} du.$$

We claim that the limit exists everywhere and is finite off the diagonal. Indeed, from (7) and (8) we have

$$\begin{aligned} \lim_{t \rightarrow 0} t^{-1} \eta_t(u) &= \lim_{t \rightarrow 0} t^{-1-2/\alpha} \eta_1(ut^{-2/\alpha}) = u^{-1-\alpha/2} \lim_{s \rightarrow \infty} s^{1+\alpha/2} \eta_1(s) \\ &= [\alpha/(2\Gamma(1 - \alpha/2))]u^{-1-\alpha/2}, \quad u > 0. \end{aligned}$$

By (4) and (9), for $x \neq y$,

$$t^{-1}q(u, x, y)\eta_t(u) \leq c_2 u^{-d_s/2} \exp(-c_3|x - y|^\gamma u^{-\gamma/d_w})u^{-1-\alpha/2}, \quad u \in (0, \infty).$$

Since this is integrable our claim follows by the dominated convergence theorem. For later use we note that

$$(43) \quad N(x, y) = c_1 \int_0^\infty q(u, x, y)u^{-1-\alpha/2} du,$$

with $c_1 = \alpha/(2\Gamma(1 - \alpha/2))$.

Let E be a Borel subset of F and $x \in F$. Define

$$n(x, E) = \int_E N(x, y) d\mu(y),$$

Note that by Proposition 4.4, for a bounded Borel set $D \subseteq F$ we have

$$\sup_{x \in F} E^x \tau_D < \infty.$$

We have the following formula (see [IW]).

PROPOSITION 6.1. *Assume that $D \subseteq F$ is an open nonempty bounded set, $E \subseteq F$ is a Borel set and $\text{dist}(E, D) > 0$. Then*

$$(44) \quad P^x[X_{\tau_D} \in E] = \int_D G_D(x, y)n(y, E) d\mu(y).$$

Proof. We need to check the following assumptions (A1) and (A2) from [IW]. Let $\mathbb{M} = (S, P^x, x \in S)$ be a Markov process on a locally compact, separable metric space S which satisfies

(A1) The semigroup

$$T_t f(x) = \int_S f(y)P(t, x, dy)$$

maps $C_0(S)$ into $C_0(S)$ and is strongly continuous in $t \geq 0$.

(A2) There exists a positive kernel $n(x, E)$, $x \in S$, $E \subseteq S$ a Borel subset, such that

1° If $\text{dist}(x, E) > 0$ then

$$n(x, E) < \infty.$$

2° For $f \in C_0(S)$ and a bounded open set D with $\text{dist}(D, \text{supp } f) > 0$,

$$\sup_{x \in D, t > 0} t^{-1}T_t f(x) < \infty$$

and

$$\lim_{t \rightarrow 0^+} t^{-1}T_t f(x) = \lim_{t \rightarrow 0^+} t^{-1} \int_S f(y) P(t, x, dy) = \int_S f(y) n(x, dy)$$

for every $x \in D$.

Assumption (A1) is satisfied by Proposition 3.2. To establish (A2), fix D as above. Let $f \in C_0(D)$ be such that $\text{dist}(D, \text{supp } f) \geq \delta > 0$. Then, by (5) and Lemma 2.1,

$$\sup_{x \in D, t > 0} t^{-1} P_t f(x) \leq c \sup_{x \in D} \int \frac{f(y) d\mu(y)}{|x - y|^{d_\alpha}} \leq c \|f\|_\infty \delta^{-d_\alpha} < \infty$$

and

$$\lim_{t \rightarrow 0} t^{-1} P_t f(x) = \lim_{t \rightarrow 0} t^{-1} \int f(y) p(t, x, y) d\mu(y) = \int f(y) n(x, dy),$$

by bounded convergence. ■

COROLLARY 6.2. *Under the assumptions of Proposition 6.1 we have*

$$(45) \quad P^x[X_{\tau_D} \in E] \asymp \int_D \int_E \frac{G_D(x, y)}{|y - z|^{d_\alpha}} d\mu(z) d\mu(y).$$

Proof. By a substitution as in (11), from (43) for $x, y \in F$, $x \neq y$, it follows that

$$\begin{aligned} N(x, y) &\asymp \int_0^\infty u^{-1-\alpha/2-d_s/2} \exp(-c|x - y|^\gamma u^{-\gamma/d_w}) du \\ &= c|x - y|^{-d_\alpha} \int_0^\infty v^{-(d_w-1)(-1-\alpha/2-d_s/2)-d_w} e^{-v} dv. \end{aligned}$$

This yields

$$(46) \quad n(x, E) \asymp \int_E \frac{d\mu(y)}{|y - x|^{d_\alpha}}, \quad x \in F,$$

and (45) follows from (44). ■

In particular, the above corollary implies that the distribution of X_{τ_D} is absolutely continuous with respect to μ on $\text{int}(D^c)$. The corresponding density (*Poisson kernel*) is denoted by $P_D(x, y)$.

PROPOSITION 6.3. *Let $D \subseteq F$ be an open bounded set and $\alpha \neq d_s$. Then the Poisson kernel $P_D(\cdot, \cdot)$ admits a version which is jointly continuous on $D \times \text{int}(D^c)$.*

Proof. We claim that $N(x, y)$ is continuous on the set $S_a = \{(x, y) \in F \times F : |x - y| > a\}$ for each $a > 0$. Indeed, let $(x, y) \in S_a$ and $x_n \rightarrow x$ and $y_n \rightarrow y$ as $n \rightarrow \infty$. Then, for sufficiently large n , we have $(x_n, y_n) \in S_a$ and $u^{-1-\alpha/2} q(u, x_n, y_n) \leq c_1 u^{-d_s/2-1-\alpha/2} \exp(-c_2 a^\gamma u^{-\gamma/d_w})$, $u \in (0, \infty)$.

Consequently, by (43) and dominated convergence we obtain

$$N(x_n, y_n) = c_3 \int_0^\infty q(u, x_n, y_n) u^{-1-\alpha/2} du \rightarrow c_3 \int_0^\infty q(u, x, y) u^{-1-\alpha/2} du$$

as $n \rightarrow \infty$. This shows our claim.

We let

$$P_D(x, y) = \int_D G_D(x, z)N(z, y) d\mu(z), \quad x \in D, \quad y \in \text{int}(D^c).$$

Assume that $\alpha < d_s$. Fix $(x, y) \in D \times \text{int}(D^c)$ and let $x_n \rightarrow x, y_n \rightarrow y$. We may and do assume that $\text{dist}(y_n, D) \geq \frac{1}{2} \text{dist}(y, D), n = 1, 2, \dots$. Then, by (34) and (46), for $\varepsilon > 0$ such that $\varepsilon < (d_s/\alpha - 1)^{-1}$ and for R large enough we have

$$\begin{aligned} (47) \quad & \sup_n \int_D |G_D(x_n, z)N(z, y_n)|^{1+\varepsilon} d\mu(z) \\ & \leq c \sup_n \int_D |x_n - z|^{(-d+\alpha d_w/2)(1+\varepsilon)} |z - y_n|^{-d_\alpha(1+\varepsilon)} d\mu(z) \\ & \leq c(\text{dist}(y, D))^{-d_\alpha(1+\varepsilon)} \sup_n \int_D |x_n - z|^{(-d+\alpha d_w/2)(1+\varepsilon)} d\mu(z) \\ & \leq c(\text{dist}(y, D))^{-d_\alpha(1+\varepsilon)} \sup_n \int_{B(x_n, R)} |x_n - z|^{(-d+\alpha d_w/2)(1+\varepsilon)} d\mu(z) < \infty, \end{aligned}$$

by (3). It follows that $(G_D(x_n, \cdot)N(\cdot, y_n))_{n \in \mathbb{N}}$ are uniformly integrable and consequently

$$\begin{aligned} (48) \quad & \lim_{n \rightarrow \infty} P_D(x_n, y_n) = \lim_{n \rightarrow \infty} \int_D G_D(x_n, z)N(z, y_n) d\mu(z) \\ & = \int_D G_D(x, z)N(z, y) d\mu(z) = P_D(x, y). \end{aligned}$$

This completes the case $\alpha < d_s$. If $\alpha > d_s$ then the process is point-recurrent and the Green function for D is bounded (see Proposition 5.6). Therefore, a similar but simpler argument applies. ■

PROPOSITION 6.4. *There is a constant c such that for each $\kappa > 1, x_0 \in F, r > 0$ and for $c_1 = ((\kappa + 1)/(\kappa - 1))^{d_\alpha c}$ and $c_2 = ((\kappa - 1)/(\kappa + 1))^{d_\alpha c}$ we have*

$$(49) \quad P_{B(x_0, r)}(x, z) \leq c_1 r^{\alpha d_w/2} |x - z|^{-d_\alpha}, \quad x \in B(x_0, r), z \in B(x_0, \kappa r)^c,$$

$$(50) \quad P_{B(x_0, r)}(x, z) \geq c_2 r^{\alpha d_w/2} |x - z|^{-d_\alpha}, \quad x \in B(x_0, r/\kappa), z \in \text{int}(B(x_0, r)^c).$$

Proof. Let $x, y \in B = B(x_0, r)$. Then, for $z \in B(x_0, \kappa r)^c$,

$$|y - z| \geq |z - x_0| - |y - x_0| \geq \kappa r - r = (\kappa - 1)r$$

and consequently

$$\begin{aligned} |x - z| & \leq |y - z| + |y - x| \leq |y - z| + 2r \\ & \leq |y - z| + 2|y - z|/(\kappa - 1) = |y - z|(\kappa + 1)/(\kappa - 1). \end{aligned}$$

Thus, from Proposition 4.4 it follows that

$$(51) \quad P_B(x, z) \asymp \int_B G_B(x, y) |y - z|^{-d_\alpha} d\mu(y) \\ \leq ((\kappa + 1)/(\kappa - 1))^{d_\alpha} |x - z|^{-d_\alpha} E^x \tau_B \leq c_1 |x - z|^{-d_\alpha} r^{\alpha d_w/2},$$

with $c_1 = c_1(\kappa)$. This gives (49). Now, let $x \in B(x_0, r/\kappa)$, $z \in \text{int}(B(x_0, r)^c)$. Then $|y - z| \leq |x - z|(\kappa + 1)/(\kappa - 1)$, and from (22) we obtain

$$P_B(x, z) \geq c((\kappa - 1)/(\kappa + 1))^{d_\alpha} |x - z|^{-d_\alpha} E^x \tau_B \geq c_2 |x - z|^{-d_\alpha} r^{\alpha d_w/2},$$

with $c_2 = c_2(\kappa)$ (cf. (51)). This completes the proof. ■

LEMMA 6.5. *Let $\alpha < d_s$ and $D = B(x_0, r)$ where $x_0 \in F$ and $r > 0$ are arbitrary. Then for each $k \geq 2$ there exists a constant $c_1 = c_1(k)$ such that*

$$(52) \quad P_D(x, y) \leq c_1 r^{-d+\alpha d_w/2} \delta(y)^{-\alpha d_w/2}, \quad y \in B(x_0, kr) \cap D^c, x \in B(x_0, r/2),$$

where $\delta(y) = \text{dist}(y, D)$.

Proof. By (45) and (34) we obtain

$$(53) \quad P_D(x, y) \asymp \int_D G_D(x, z) |z - y|^{-d_\alpha} d\mu(z) \\ \leq c \left(\int_{B(x, r/4)} + \int_{D \cap B(x, r/4)^c} \right) |x - z|^{-d+\alpha d_w/2} |z - y|^{-d_\alpha} d\mu(z).$$

If $|x - z| < r/4$ then $|z - y| > r/4$, so that the first integral in (53) is not greater than

$$c(r/4)^{-d_\alpha} \int_{B(x, r/4)} |x - z|^{-d+\alpha d_w/2} d\mu(z) \\ \leq c(r/4)^{-d_\alpha} (r/4)^{\alpha d_w/2} = cr^{-d+\alpha d_w/2} r^{-\alpha d_w/2} \leq c_2 r^{-d+\alpha d_w/2} \delta(y)^{-\alpha d_w/2},$$

by Lemma 2.1 and the fact that $\delta(y) < (k-1)r$. Here $c_2 = c_2(k)$. The second integral in (53) does not exceed

$$cr^{-d+\alpha d_w/2} \int_{D \setminus B(x, r/4)} |z - y|^{-d-\alpha d_w/2} d\mu(z) \\ \leq cr^{-d+\alpha d_w/2} \int_{B(y, \delta(y))^c} |z - y|^{-d-\alpha d_w/2} d\mu(z) \leq cr^{-d+\alpha d_w/2} \delta(y)^{-\alpha d_w/2},$$

by Lemma 2.1. This completes the proof. ■

7. Harnack inequality. Recall that $F \subseteq \mathbb{R}^N$. The main result of this section can be stated as follows.

THEOREM 7.1 (Harnack inequality). *Let $\alpha \in (0, 2/d_w) \cup (d_s, 2)$. Then there exist c_1, c_2 such that for any $x_0 \in F$, $r > 0$ and any function $h \geq 0$*

(regular) α -harmonic in $B(x_0, c_1r)$, we have

$$(54) \quad h(x) \geq c_2h(y), \quad x, y \in B(x_0, r).$$

Since we can always take a smaller ball as the region of α -harmonicity of our function, we may and do assume *regular* α -harmonicity above. The method of proof depends on whether the process is point-recurrent or transient. Theorem 7.1 will be strengthened in Corollary 7.7 below.

7.1. Recurrent case. In this subsection we assume that $\alpha > d_s$, so that the process is point-recurrent (see Lemma 7.2 and Remark 3 below).

For $\lambda > 0$ define the λ -potential

$$G^\lambda(x, y) = \int_0^\infty e^{-\lambda t} p(t, x, y) dt, \quad x, y \in F.$$

LEMMA 7.2. *There exist c_1 such that*

$$(55) \quad G^\lambda(x, y) \leq c_1\lambda^{d_s/\alpha-1}, \quad x, y \in F.$$

Proof. This follows immediately from the definition of $G^\lambda(x, y)$ and our basic estimate $p(t, x, y) \leq ct^{-d_s/\alpha}$:

$$G^\lambda(x, y) \leq c \int_0^\infty e^{-\lambda t} t^{-d_s/\alpha} dt = c\Gamma(1 - d_s/\alpha)\lambda^{d_s/\alpha-1}. \blacksquare$$

REMARK 3. Since the λ -potential is bounded on $F \times F$, it follows that points have positive capacity and the process is point-recurrent (see e.g. [PS, Theorem 7.1]). In particular, $P^x[T_{\{y\}} < \infty] = 1$, $x, y \in F$ (in fact, points are *regular* for themselves, see [BG, Ch. 6, Proposition 4.11]).

For simplicity we write $T_y := T_{\{y\}}$.

LEMMA 7.3. *There exist constants $a > 1$ and p_0 such that*

$$P^x[T_y < \tau_{B(x, a|x-y|)}] > p_0, \quad x, y \in F.$$

Proof. First, we prove a general fact:

$$(56) \quad P^x[T_y < \tau_D] = G_D(x, y)/G_D(y, y).$$

Let $\delta > 0$ be such that $B(y, \delta) \subseteq D$. From (39) with $U = D \setminus B(y, \delta)$ and $x \in U$ we obtain

$$G_D(x, y) = E^x[G_D(X(\tau_{D \setminus B(y, \delta)}), y); T_{B(y, \delta)} < \tau_D].$$

Letting $\delta \rightarrow 0$, by the continuity and boundedness of $G_D(\cdot, y)$, we get (56).

Now, let a and $r = |x - y|$ be as in Lemma 5.8. Then, from (56) with $D = B(x, a|x - y|)$, (41) and Proposition 5.6, it follows that

$$P^x[T_y < \tau_D] \geq c \frac{|x - y|^{-d+\alpha d_w/2}}{(ar)^{-d+\alpha d_w/2}} \geq p_0,$$

which completes the proof. \blacksquare

Proof of Theorem 7.1 for $d_s < \alpha < 2$. Let a, p_0 be the constants of Lemma 7.3 and c_1 be such that $B(x, 2ar) \subseteq B(x_0, c_1r)$, $x \in B(x_0, r)$. Then $B(x, a|x - y|) \subseteq B(x_0, c_1r)$ for any $x, y \in B(x_0, r)$. It follows that

$$\begin{aligned} h(x) &= E^x h(X_{\tau_{B(x, a|x-y|)}}) \geq E^x [h(X_{\tau_{B(x, a|x-y|)}}); T_y < \tau_{B(x, a|x-y|)}] \\ &= E^x [E^{X(T_y)} [h(X_{\tau_{B(x, a|x-y|)}})]; T_y < \tau_{B(x, a|x-y|)}] \\ &= h(y) P^x [T_y < \tau_{B(x, a|x-y|)}] \geq p_0 h(y). \end{aligned}$$

This proves Theorem 7.1 for $\alpha > d_s$. ■

7.2. Transient case. Throughout this subsection we consider the transient case $\alpha < d_s$ (see Corollary 5.4). However, due to our restricted knowledge on the Poisson kernel and some geometrical reasons, in the proof of the remaining part of Theorem 7.1 we also assume the more restrictive condition $\alpha < 2/d_w$. We say that E is *polar* if $P^x [T_E < \infty] = 0$ for all $x \in F$. For a Borel set E let $\dim_H(E)$ denote its Hausdorff dimension.

LEMMA 7.4. *Let E be a Borel set in $F \subseteq \mathbb{R}^N$. If $\alpha < 2(d - \dim_H(E))/d_w$ then E is polar.*

Proof. The proof is an application of some general facts from potential theory. For $t > 0$ and a Borel measure m with compact support in \mathbb{R}^N , such that $0 < m(\mathbb{R}^N) < \infty$, let $\phi_{m,t}(x)$ be its t -potential:

$$\phi_{m,t}(x) = \int \frac{dm(y)}{|x - y|^t}.$$

For a compact set $K \subseteq \mathbb{R}^N$ define the t -capacity of E by

$$C_t(K) = \sup_m \left\{ \left(\int \phi_{m,t}(x) dx \right)^{-1} \right\},$$

where the supremum is taken over Borel measures m such that $\text{supp } m \subseteq K$ and $m(K) = 1$ (see [Fa1]). Equivalently (see [L]),

$$C_t(K) = \sup \{ m(K) : \phi_{m,t} \leq 1 \}.$$

For an arbitrary $E \subseteq \mathbb{R}^N$ define

$$C_t(E) = \sup \{ C_t(K) : K \text{ is compact, } K \subseteq E \}.$$

By Corollary 6.5 from [Fa1],

$$\dim_H(E) = \inf \{ t : C_t(E) = 0 \} = \sup \{ t : C_t(E) > 0 \}.$$

When $\alpha d_w/2 < d$ (i.e. $\alpha < d_s$), by [BG, Ch. 6, Section 4], E is polar if and only if $C_{d-\alpha d_w/2}(E) = 0$.

Now, by our assumption, $\dim_H(E) < d - \alpha d_w/2$ so that $C_{d-\alpha d_w/2}(E) = 0$. It follows that E is polar, which is the assertion of the lemma. ■

For $z \in B(0, r/2)$ and $s \in (r, 2r)$ let $K(z, s) = B(z(s-r)/r, (s+r)/2)$. Observe that $B(0, r) \subseteq K(z, s) \subseteq B(0, s)$. Indeed, if $y \in B(0, r)$ then

$$|y - z(s-r)/r| \leq |y| + |z|(s-r)/r \leq r + (s-r)/2 = (s+r)/2,$$

so $B(0, r) \subseteq K(z, s)$; in particular $F \cap K(z, s)$ is not empty ($0 \in F$, but $z(s-r)/r$ may belong to $\mathbb{R}^N \setminus F$). Also, if $y \in K(z, s)$ then

$$|y| \leq |z|(s-r)/r + (s+r)/2 \leq (s-r)/2 + (s+r)/2 = s,$$

and consequently $K(z, s) \subseteq B(0, s)$. Note also that if $|z| \leq r/2$ and $s_1 \leq s_2$ then $K(z, s_1) \subseteq K(z, s_2)$. Indeed, if $x \in K(z, s_1)$ then

$$\begin{aligned} \left| x - z \frac{s_2 - r}{r} \right| &\leq \left| x - z \frac{s_1 - r}{r} \right| + |z| \left| \frac{s_1 - r}{r} - \frac{s_2 - r}{r} \right| \\ &\leq \frac{s_1 + r}{2} + \frac{r}{2} \frac{s_2 - s_1}{r} = \frac{s_2 + r}{2}, \end{aligned}$$

which means that $x \in K(z, s_2)$.

LEMMA 7.5. *Let $r > 0$. There exists $z_0 \in B(0, r/12)$ (not necessarily in F) such that for almost all $s \in (r, 2r)$ the stable process does not hit the boundary of $K(z_0, s)$ in F .*

Proof. Let $s \in (r, 2r)$ and $\partial K(z, s) = \partial B(z(s-r)/r, (s+r)/2) \cap F$ (on the right hand side of this equality we consider the boundary of the ball in \mathbb{R}^N). From Theorem 8.1 and the Product Formula 7.3 in [Fa2] it follows that for almost all (with respect to the Lebesgue measure) $z \in B(0, r/2) \subseteq \mathbb{R}^N$ we have $\dim_{\mathbb{H}}(\partial K(z, s)) \leq (N-1+d) - N = d-1$. Therefore, $\alpha < 2/d_w \leq 2(d - \dim_{\mathbb{H}}(\partial K(z, s)))/d_w$, and so $\partial K(z, s)$ is polar by Lemma 7.4. For $x \in F$ and a Borel set A the mapping $(z, s) \mapsto P^x[X_{\tau_{K(z,s)}} \in A]$ is jointly measurable. We thus we obtain

$$\int_r^{2r} \int_{z \in B(0, r/2)} P^x[X_{\tau_{K(z,s)}} \in \partial K(z, s)] dz ds = 0.$$

Hence, by Fubini's theorem

$$(57) \quad \int_r^{2r} P^x[X_{\tau_{K(z,s)}} \in \partial K(z, s)] ds = 0$$

for almost all $z \in B(0, r/2)$ and the assertion follows. ■

REMARK 4. When the process does not hit the boundary of a region D , for every Borel $u \geq 0$,

$$E^x u(X_{\tau_D}) = \int_{D^c} P_D(x, y) u(y) d\mu(y).$$

This fact will be exploited in what follows.

Proof of Theorem 7.1 for $\alpha \in (0, 2/d_w)$. Without losing generality we assume that $x_0 = 0$. Let h be a positive function that is regular α -harmonic in $B(0, 2r)$. We will show that $h(x) \leq ch(\tilde{x})$, $x, \tilde{x} \in B(0, r/2)$. Let

$$\begin{aligned} h_1(x) &= E^x[h(X_{\tau_{B(0,2r)}}); X_{\tau_{B(0,2r)}} \in B(0, 3r)], \\ h_2(x) &= E^x[h(X_{\tau_{B(0,2r)}}); X_{\tau_{B(0,2r)}} \in B(0, 3r)^c]. \end{aligned}$$

Then $h = h_1 + h_2$ on F and, by definition, the functions h_1 and h_2 are regular α -harmonic in $B(0, 2r)$. Moreover, since $\text{supp } h_2 \cap B(0, 2r)^c \subseteq B(0, 3r)^c$, for $x, \tilde{x} \in B(0, r/2)$ we have, by Proposition 6.4,

$$\begin{aligned} h_2(x) &= \int_{B(0,3r)^c} h_2(y)P_{B(0,2r)}(x, y) d\mu(y) \\ &\leq cr^{\alpha d_w/2} \int_{B(0,3r)^c} h_2(y)|x - y|^{-d_\alpha} d\mu(y) \\ &\leq cr^{\alpha d_w/2} \int_{B(0,3r)^c} h_2(y)|\tilde{x} - y|^{-d_\alpha} d\mu(y) \\ &\leq c \int_{B(0,3r)^c} h_2(y)P_{B(0,2r)}(\tilde{x}, y) d\mu(y) = ch_2(\tilde{x}). \end{aligned}$$

Consequently, it is enough to show an analogous inequality for h_1 . Since we do not know whether the process hits the boundary of $B(0, 3r/4)$, we have the inequality

$$(58) \quad h_1(\tilde{x}) = E^{\tilde{x}}h_1(X_{\tau_{B(0,3r/4)}}) \geq \int_{B(0,3r/4)^c} P_{B(0,3r/4)}(\tilde{x}, y)h_1(y) d\mu(y).$$

Define $R = B(0, 3r) \setminus B(0, r)$. Note that $\text{supp } h_1 \subseteq B(0, 3r)$, and also, for $\tilde{x} \in B(0, r/2)$ and $y \in R$ we have $|\tilde{x} - y| \leq 4r$. From (58) and Proposition 6.4 it follows that

$$\begin{aligned} h_1(\tilde{x}) &\geq \int_R P_{B(0,3r/4)}(\tilde{x}, y)h_1(y) d\mu(y) \geq cr^{\alpha d_w/2} \int_R |\tilde{x} - y|^{-d_\alpha} h_1(y) d\mu(y) \\ &\geq cr^{-d} \int_R h_1(y) d\mu(y). \end{aligned}$$

Set

$$P(x, y) = \int_r^{2r} P_{K(z_0, s)}(x, y) ds, \quad x \in B(0, r/2), y \in R,$$

with z_0 of Lemma 7.5 and the usual convention: $P_D(x, y) = 0$ if $y \in D$. Then

$$h_1(x) = r^{-1} \int_R h_1(y)P(x, y) d\mu(y), \quad x \in B(0, r/2).$$

Indeed, by the fact that $\text{supp } h_1 \subseteq B(0, 3r)$, $B(0, r) \subseteq K(z_0, s) \subseteq B(0, 2r)$,

α -harmonicity of h_1 on $B(0, 2r)$, Lemma 7.5 and Remark 4 we obtain

$$\begin{aligned} \int_R h_1(y)P(x, y) d\mu(y) &= \int_R h_1(y) \int_r^{2r} P_{K(z_0, s)}(x, y) ds d\mu(y) \\ &= \int_r^{2r} \int_{y \in R} h_1(y)P_{K(z_0, s)}(x, y) d\mu(y) ds \\ &= \int_r^{2r} h_1(x) ds = rh_1(x), \end{aligned}$$

which gives our claim.

Next we prove that

$$(59) \quad P(x, y) \leq cr^{1-d}, \quad x \in B(0, r/2), y \in R.$$

Define $\delta_s(y) = \text{dist}(y, K(z_0, s))$. Let $s_0(y) = \inf\{s > 0 : y \in K(z_0, s)\}$ so that $y \in K(z_0, s)$ if $s > s_0(y)$. Recall that $|z_0| < r/12 < r/4$. Hence,

$$B(0, r/2) \subseteq B\left(\frac{z_0(s-r)}{r}, \frac{1}{2} \frac{s+r}{2}\right), \quad s \in (r, 2r).$$

By Lemma 6.5 and the substitution $s = r(u + 1)$, for $x \in B(0, r/2)$ and $y \in R$ we get

$$\begin{aligned} (60) \quad P(x, y) &\leq cr^{-d+\alpha d_w/2} \int_r^{2r \wedge s_0(y)} \delta_s(y)^{-\alpha d_w/2} ds \\ &\leq cr^{-d+\alpha d_w/2} \int_r^{2r \wedge s_0(y)} (|y - z_0(s-r)|/r - (s+r)/2)^{-\alpha d_w/2} ds \\ &= cr^{1-d+\alpha d_w/2} \int_0^{1 \wedge u_0(y)} (|y - uz_0| - (u+2)r/2)^{-\alpha d_w/2} du, \end{aligned}$$

where $u_0 = u_0(y) = -1 + s_0(y)/r \geq 0$. Moreover,

$$(61) \quad |y - uz_0| - (u+2)r/2 = \frac{|y - uz_0|^2 - (u+2)^2 r^2/4}{|y - uz_0| + (u+2)r/2} \asymp \frac{1}{r} g(u),$$

where g is the numerator of the last fraction:

$$g(u) = g_{y, z_0, r}(u) = u^2(|z_0|^2 - r^2/4) - u(2\langle z_0, y \rangle + r^2) + |y|^2 - r^2.$$

Observe that $g(0) > 0$ and $|z_0|^2 - r^2/4 < 0$ and so $u = u_0$ is the unique nonnegative solution of $g(u) = 0$. Hence

$$\begin{aligned} u_0 &= \frac{2\langle z_0, y \rangle + r^2 - \sqrt{(2\langle z_0, y \rangle + r^2)^2 + (r^2 - 4|z_0|^2)(|y|^2 - r^2)}}{2(|z_0|^2 - r^2/4)} \\ &= \frac{2(|y|^2 - r^2)}{\sqrt{(2\langle z_0, y \rangle + r^2)^2 + (r^2 - 4|z_0|^2)(|y|^2 - r^2)} + 2\langle z_0, y \rangle + r^2}. \end{aligned}$$

Since $|z_0| < r/12$, we have $2|\langle z_0, y \rangle| \leq 2|z_0| \cdot |y| < r^2/2$ and $r^2 - 4|z_0|^2 \asymp r^2$. Consequently, $u_0 \asymp (|y| - r)/r$ and $g(0)/u_0 \asymp r^2$. By concavity of $g(u)$ we have $g(u) \geq g(0)(1 - u/u_0)$. From (60), (61) and the fact that $u_0 \leq 2$ it follows that

$$\begin{aligned} P(x, y) &\leq cr^{1-d+\alpha d_w/2} r^{\alpha d_w/2} \int_0^{1 \wedge u_0} g(u)^{-\alpha d_w/2} du \\ &\leq cr^{1-d+\alpha d_w} (g(0)/u_0)^{-\alpha d_w/2} \int_0^{1 \wedge u_0} (u_0 - u)^{-\alpha d_w/2} du \\ &\leq cr^{1-d} [-(u_0 - u)^{1-\alpha d_w/2}]_0^{1 \wedge u_0} \leq cr^{1-d}, \end{aligned}$$

which gives our claim. It follows that

$$\begin{aligned} h_1(x) &= r^{-1} \int_R h_1(y) P(x, y) d\mu(y) \leq cr^{-1} \int_R h_1(y) r^{1-d} d\mu(y) \\ &= cr^{-d} \int_R h_1(y) d\mu(y) \leq ch_1(\tilde{x}). \end{aligned}$$

Thus, for functions α -harmonic in $B(0, 2r)$ we showed (54) except that $x, y \in B(x_0, r)$ is replaced by $x, y \in B(x_0, r/2)$. Hence, substituting $2r$ for r we get the assertion of Theorem 7.1 with $c_1 = 4$. The proof is complete. ■

The following lemma is motivated by [BBy]. It is a useful tool in the proof of the Boundary Harnack Principle and extends Theorem 7.1 slightly. For the reader’s convenience we give the proof, as the general context is much different than the one in [BBy]. We point out that the situation is completely different as compared to the diffusion case. This is due to the fact that the definition of α -harmonic functions is a global one. In general, a ball in F may be disconnected. These circumstances make the classical chain argument unavailable in our setting (compare also with Proposition 3.37 and Corollary 3.38 in [Ba]). A desired extension of (54) in the diffusion case is not even true, since on each component of a (disconnected) ball the harmonic function can be defined separately.

LEMMA 7.6. *Let $x_1, x_2 \in F$, $r > 0$ and $M \in \mathbb{R}$ satisfy $|x_1 - x_2| \leq Mr$. Let $u \geq 0$ be a function which is α -harmonic in $B(x_1, r) \cup B(x_2, r)$. Then α -harmonic*

$$c^{-1}M^{-d\alpha}u(x_2) \leq u(x_1) \leq cM^{d\alpha}u(x_2).$$

Proof. Without any loss of generality we assume $x_1 = 0$ and $|x_2| \geq 3r/2$, because otherwise we can consider smaller r . By Theorem 7.1 we obtain

$$u(x) \geq cu(x_2), \quad x \in B(x_2, r/4).$$

By (50), for $x \in B(x_2, r/4)$ we have

$$P_{B(0,r/2)}(0, x) \geq cr^{\alpha d_w/2} |x|^{-d_\alpha} \geq cr^{\alpha d_w/2} (Mr)^{-d_\alpha} = cM^{-d_\alpha} r^{-d}.$$

It follows that

$$u(0) \geq \int_{B(x_2,r/4)} P_{B(0,r/2)}(0, x)u(x) d\mu(x) \geq c\mu(B(x_2, r/4))M^{-d_\alpha}r^{-d}u(x_2),$$

and, by (1) and symmetry, the assertion follows. ■

COROLLARY 7.7. *Let $\alpha \in (0, 2/d_w) \cup (d_s, 2)$. For each $\kappa > 1$ there exists $c_1 = c_1(\kappa)$ such that for any $x_0 \in F$, $r > 0$ and any function $h \geq 0$ regular α -harmonic in $B(x_0, r)$, we have*

$$(62) \quad h(x) \geq c_1 h(y), \quad x, y \in B(x_0, r/\kappa).$$

Proof. As previously we assume $x_0 = 0$. It is enough to show

$$(63) \quad c_2 h(0) \leq h(x) \leq c_3 h(0), \quad x \in B(0, r/\kappa),$$

with $c_2, c_3 = c(\kappa)$. Let $\tilde{r} = (\kappa - 1)r/\kappa$. Clearly, $B(x, \tilde{r}) \subseteq B(0, r)$, so that h is α -harmonic in $B(x, \tilde{r}) \cup B(0, \tilde{r})$. From Lemma 7.6, with \tilde{r} and $M = 1/(\kappa - 1)$, we obtain (63), and the assertion follows. ■

8. Estimates near boundary. Let D be an open nonempty bounded subset of F . We will be interested in the decay rate of the exit time and α -harmonic functions near the boundary of D . The main results of this section are Theorems 8.3 and 8.4.

To measure the regularity of a set we make the following definition. We say that $D \subseteq F$ has the *outer fatness property* if there are constants $c_1 = c_1(D)$ and $R_0 = R_0(D)$ such that

$$(64) \quad \mu(D^c \cap B(x, r)) \geq c_1 r^d, \quad x \in \partial D, r \in (0, R_0).$$

It is clear that this holds for the interior of the *natural cells* and their finite unions in the Sierpiński gasket (see below for the definition). For the rest of this section we assume that D satisfies (64).

In Lemma 8.1 through Theorem 8.4 for simplicity we assume that $\text{diam}(D) = 1$. We set $D_n = \{x \in D : \delta(x) \leq k^{-n}\}$, $n = 0, 1, \dots$, where $\delta(x) = \text{dist}(x, D^c) \leq 1$ and k is a natural number whose value will be specified later. Observe that $D_0 = D$. Let $P_n = D_n \setminus D_{n+1}$, $n = 0, 1, \dots$

LEMMA 8.1. *There exists c_1 such that for each $a > 2$, $r > 0$ and $x \in F$,*

$$P^x[X_{\tau_{B(x,r)}} \notin B(x, ar)] \leq c_1 a^{-\alpha d_w/2}.$$

Proof. Since $a > 2$ we can apply Proposition 6.4 with $\kappa = 2$ and get the constant independent of a . Thus

$$\begin{aligned}
 P^x[X_{\tau_{B(x,r)}} \notin B(x, ar)] &\leq cr^{\alpha d_w/2} \int_{B(x, ar)^c} |x - y|^{-d_\alpha} d\mu(y) \\
 &= cr^{\alpha d_w/2} \sum_{m=0}^\infty \int_{ar2^m \leq |y-x| \leq ar2^{m+1}} |x - y|^{-d_\alpha} d\mu(y) \\
 &\leq cr^{\alpha d_w/2} \sum_{m=0}^\infty \mu(B(x, ar2^{m+1})) (ar2^m)^{-d - \alpha d_w/2} \\
 &\leq cr^{\alpha d_w/2} (ar)^{-\alpha d_w/2} \sum_{m=0}^\infty 2^{-m\alpha d_w/2}
 \end{aligned}$$

and the assertion follows. ■

REMARK. That the constant c_1 does not depend on r and $a > 2$ may be viewed as an instance of weak scaling for our process.

For the rest of this section let $B_n = B(x, k^{-n})$.

PROPOSITION 8.2. *There exists $p = p(D) > 0$ such that*

$$P^x[\tau_D \leq \tau_{B_n}] \geq p, \quad x \in D_n.$$

Proof. We only need to prove the inequality for $n \geq 1$. Fix $x \in D_n$ and let $x_0 \in D^c$ be such that $|x - x_0| \leq k^{-n}$. Define $r_n = 1/k^n$ and $A_n = B(x_0, r_n) \cap D^c$. Then we have

$$P^x[\tau_D \leq \tau_{B_n}] \geq P^x[X(\tau_{B_n}) \in D^c] \geq \int_{A_n} P_{B(x, r_n)}(x, y) d\mu(y).$$

We choose k large enough so that $1/k < R_0$ in (64). Observe that if (64) holds for all points in ∂D then, by (1), it holds for x_0 as well. Also, we have $|x - y| \leq |x - x_0| + |x_0 - y| \leq 2k^{-n}$, $y \in A_n$. Therefore, by (50), for $n \geq 1$ we have

$$P^x[\tau_D \leq \tau_{B_n}] \geq c \int_{A_n} r_n^{\alpha d_w/2} |x - y|^{-d_\alpha} d\mu(y) \geq ck^{nd} \mu(A_n) \geq cc_1,$$

where the constant c_1 comes from (64). This completes the proof. ■

THEOREM 8.3. *There exist $\beta = \beta(D) \in (0, 1)$ and $c_0 = c_0(D)$ such that*

$$E^x \tau_D \leq c_0 \delta(x)^\beta, \quad x \in D.$$

Proof. Fix $x \in D$. Let r_n be as in the proof of the preceding proposition. Define

$$u_n = \sup\{E^x \tau_D : x \in D_n\}.$$

Clearly, it is enough to show

$$u_n \leq c_3 c_4^n, \quad n = 0, 1, \dots,$$

with $c_3 = c_3(D)$, $c_4 = c_4(D) \in (0, 1)$. By the strong Markov property we have

$$\begin{aligned} E^x \tau_D &= E^x[\tau_D; \tau_{B_n} \geq \tau_D] + E^x[\tau_D; \tau_{B_n} < \tau_D] \\ &\leq E^x \tau_{B_n} + E^x[E^{X(\tau_{B_n})} \tau_D; \tau_{B_n} < \tau_D]. \end{aligned}$$

By Proposition 4.4, the first term is not greater than $cr_n^{\alpha d_w/2} = ck^{-n\alpha d_w/2}$. The second one is equal to

$$\sum_{j=0}^{n-2} E^x[E^{X(\tau_{B_n})} \tau_D; X(\tau_{B_n}) \in P_j] + E^x[E^{X(\tau_{B_n})} \tau_D; X(\tau_{B_n}) \in D_{n-1}] = A + B.$$

By Proposition 8.2,

$$B \leq u_{n-1} P^x[\tau_{B_n} \leq \tau_D] \leq (1 - p)u_{n-1}.$$

Suppose that $n \geq 2$, for otherwise the term A is absent. Then

$$A \leq \sum_{j=0}^{n-2} u_j P^x[X(\tau_{B_n}) \in P_j] = \sum_{j=2}^n u_{n-j} P^x[X(\tau_{B_n}) \in P_{n-j}].$$

Let $x_0 \in D^c$ be a point closest to x . Note that $|y - x| \leq 1/(2k^{n-j+1})$ yields

$$\delta(y) \leq |y - x_0| \leq |y - x| + |x - x_0| \leq 1/(2k^{n-j+1}) + 1/k^n \leq 1/k^{n-j+1},$$

provided $k \geq 2$ and $j \geq 2$. In other words, $P_{n-j} \subseteq B(x, r_n k^{j-1}/2)^c$. Consequently,

$$\begin{aligned} P^x[X(\tau_{B_n}) \in P_{n-j}] &\leq P^x[X(\tau_{B_n}) \notin B(x, r_n k^{j-1}/2)] \\ &\leq ck^{-(j-1)\alpha d_w/2} \leq ck^{-(j-1)\alpha d_w/2}, \end{aligned}$$

by Lemma 8.1. We conclude that

$$(65) \quad u_n \leq c_1 k_0^{-n} + (1 - p)u_{n-1} + c_2 \sum_{j=2}^n u_{n-j} k_0^{-(j-1)}, \quad n = 2, 3, \dots,$$

where $k_0 = k^{\alpha d_w/2}$. Let $c_4 = c_4(D)$ be such that $1 - p < c_4 < 1$. Fix any $n_0 \geq 3$. We now choose the value of k large enough to satisfy the following:

- (a) $k_0^{1/4} c_4 > 1$,
- (b) $c_2 < k_0^{1/4}$,
- (c) $(k_0 c_4)^{-n_0} + (1 - p)/c_4 + (k_0^{1/4} - 1)^{-1} < 1$.

Moreover, we may and do choose $c_3 \geq c_1$ such that for $m = 0, 1, \dots, n_0$,

$$(66) \quad u_m \leq c_3 c_4^m.$$

We now extend (66) to all m by induction. Assume that (66) holds for $m = 0, 1, \dots, n - 1$. From (65) it follows that

$$\begin{aligned}
 u_n &\leq c_1 k_0^{-n} + (1-p)c_3 c_4^{n-1} + c_2 \sum_{j=2}^n c_3 c_4^{n-j} k_0^{-j+1} \\
 &\leq c_3 c_4^n (1/(c_4 k_0)^n + (1-p)/c_4) + c_2 c_3 k_0^{-1/4} k_0^{5/4} \sum_{j=2}^n c_4^{n-j} k_0^{-j/4} k_0^{-3j/4} \\
 &\leq c_3 c_4^n (1/(c_4 k_0)^{n_0} + (1-p)/c_4) + c_3 c_4^n k_0^{5/4} \sum_{j=2}^{\infty} k_0^{-3j/4} \leq c_3 c_4^n,
 \end{aligned}$$

by our assumptions on k_0 . This ends the proof. ■

The above method also applies to harmonic functions. Analogous results are used in [B], [JK] in proofs of the Boundary Harnack Principle. In what follows we will assume the following *inner fatness property* of D .

There exist constants $\theta = \theta(D) \in (0, 1)$ and $R_0 = R_0(D)$ such that for every $r \in (0, R_0)$ and $Q \in \partial D$ there is a point $A = A(Q, r) \in D \cap B(Q, r)$ such that

$$(67) \quad B(A, \theta r) \subseteq D \cap B(Q, r).$$

Since we can always take a smaller *localization radius* R_0 , it is convenient to use the same symbol as in (64). It is clear that the interiors of a natural cell (or a finite sum of cells) in the Sierpiński gasket satisfy this condition (see below).

THEOREM 8.4. *There exist constants $\beta = \beta(D)$, $r_0 = r_0(D)$ and $c_0 = c_0(D)$ such that for all $Q \in \partial D$ and $r \in (0, r_0)$, and functions $u \geq 0$ regular α -harmonic in $D \cap B(Q, r)$ and satisfying $u(x) = 0$ on $D^c \cap B(Q, r)$, we have*

$$u(x) \leq c_0 (|x - Q|/r)^\beta c(u), \quad x \in D \cap B(Q, r),$$

where $c(u) = \sup\{u(y) : y \in D \cap B(Q, r)\}$.

Proof. Let $D_n = D \cap B(Q, r/k^n)$, $n = 0, 1, \dots$, with $k \geq 2$ to be specified later. Define $u_n = \sup_{x \in D_n} u(x)$. We fix $n \geq 1$ and $x \in D_n$. Define $r_n = r/(4k^n)$ and $B_n = B(x, r_n)$. We have

$$\begin{aligned}
 u(x) &= E^x u(X(\tau_{D_0})) \\
 &= E^x [u(X(\tau_{D_0})); \tau_{B_n} > \tau_{D_0}] + E^x [u(X(\tau_{D_0})); \tau_{B_n} \leq \tau_{D_0}].
 \end{aligned}$$

On the set $\{\tau_{B_n} > \tau_{D_0}\}$ we have

$$X(\tau_{D_0}) \in D_0^c \cap B_n \subseteq D_0^c \cap B(Q, r) = D^c \cap B(Q, r),$$

so that $u(X(\tau_{D_0})) = 0$. Hence, by the strong Markov property

$$\begin{aligned}
 u(x) &= E^x [E^{X(\tau_{B_n})} u(X(\tau_{D_0})); \tau_{B_n} \leq \tau_{D_0}] \\
 &\leq E^x u(X(\tau_{B_n})) \\
 &= E^x [u(X(\tau_{B_n})); X(\tau_{B_n}) \notin B(Q, r)] + E^x [u(X(\tau_{B_n})); X(\tau_{B_n}) \in B(Q, r)].
 \end{aligned}$$

By our assumption, there is a ball $B(A, \theta r) \subseteq D \cap B(Q, r)$. Let $B_A = B(A, \theta r/2)$. Then

$$(68) \quad E^x[u(X(\tau_{B_n})); X(\tau_{B_n}) \notin B(Q, r)] \\ = \int_{B(Q, r)^c} u(z) \frac{P_{B_n}(x, z)}{P_{B_A}(A, z)} P_{B_A}(A, z) d\mu(z).$$

For $x \in D_n$ and $z \in B(Q, r)^c$ we have $|x - z| \geq r - r/(4k^n) \geq r/2 \geq 2r_n$. Also $|A - z| \geq \theta r = 2r_A$, where $r_A = \theta r/2$. It follows that we can apply Proposition 6.4, which gives

$$(69) \quad \frac{P_{B_n}(x, z)}{P_{B_A}(A, z)} \leq c \frac{r_n^{\alpha d_w/2} |x - z|^{-d_\alpha}}{r_A^{\alpha d_w/2} |A - z|^{-d_\alpha}} \leq \frac{c}{k^{n\alpha d_w/2}} \frac{|x - z|^{-d_\alpha}}{|A - z|^{-d_\alpha}}.$$

If $|z - Q| \geq r$ then

$$|x - z| \geq |z - Q| - |Q - x| \geq |z - Q| - r/k^n \geq |z - Q|(1 - 1/k^n) \geq \frac{1}{2} |z - Q|$$

and

$$|A - z| \leq |z - Q| + |Q - A| \leq |z - Q| + r \leq 2|z - Q|.$$

Combining this with (68) and (69) we obtain

$$E^x[u(X(\tau_{B_n})); X(\tau_{B_n}) \notin B(Q, r)] \leq ck^{-n\alpha d_w/2} \int_{B(Q, r)^c} u(z) P_{B_A}(A, z) d\mu(z) \\ \leq ck^{-n\alpha d_w/2} u(A) \leq ck^{-n\alpha d_w/2} u_0.$$

We need to estimate $E^x[u(X(\tau_{B_n})); X(\tau_{B_n}) \in B(Q, r)]$. This can be done exactly as in the proof of Theorem 8.3. Consequently, we arrive at (65) and the same argument as before completes the proof. ■

Proposition 8.5 below is an analogue of the Carleson estimate. We adapt the proof from [BBY] (see also [B]). Our contribution is the control of the scale parameters in the computations. Since the process does not prefer any particular scale, the result cannot depend on it, and the weak scaling suffices to prove that independence.

PROPOSITION 8.5. *Let $\alpha < 2/d_w$. There exists a constant $c_1 = c_1(\theta)$ such that for all $Q \in \partial D$ and $r \in (0, R_0/2)$, and functions $u \geq 0$ regular α -harmonic in $D \cap B(Q, 2r)$ and satisfying $u(x) = 0$ on $D^c \cap B(Q, 2r)$, we have*

$$u(x) \leq c_1 u(A), \quad x \in D \cap B(Q, r),$$

where A is as in (67).

Proof. We assume $Q = 0$. Let $K(z_0, s)$ be as in Lemma 7.5 and for $\sigma \in (r, 2r)$ define

$$u_\sigma(z) = E^z u(X(\tau_{K(z_0, \sigma)})), \quad z \in F.$$

Then

$$(70) \quad u(z) \leq u_\sigma(z), \quad z \in F.$$

Indeed,

$$\begin{aligned} u(z) &= E^z u(X(\tau_{D \cap K(z_0, \sigma)})) = E^z [u(X(\tau_{D \cap K(z_0, \sigma)})); X(\tau_{D \cap K(z_0, \sigma)}) \notin K(z_0, \sigma)] \\ &= E^z [u(X_{\tau_{K(z_0, \sigma)}}); X(\tau_{D \cap K(z_0, \sigma)}) \notin K(z_0, \sigma)] \\ &\leq E^z u(X_{\tau_{K(z_0, \sigma)}}) = u_\sigma(z). \end{aligned}$$

We claim that there is σ_0 in $(7r/4, 2r)$ such that

$$(71) \quad u_{\sigma_0}(0) \leq cr^{\alpha d_w/2} \int_{|y|>r} u(y)|y|^{-d_\alpha} d\mu(y).$$

Indeed, by Lemma 7.5 and Remark 4, the process does not hit the boundary of $K(z_0, \sigma)$ for almost all $\sigma \in (7r/4, 2r)$. Thus, we have

$$\begin{aligned} (72) \quad \int_{7r/4}^{2r} u_\sigma(0) d\sigma &= \int_{7r/4}^{2r} \int_{|y|>r} P_{K(z_0, \sigma)}(0, y) u(y) d\mu(y) d\sigma \\ &= \left(\int_{|y| \in (r, 4r)} + \int_{|y| > 4r} \right) u(y) \int_{7r/4}^{2r} P_{K(z_0, \sigma)}(0, y) d\sigma d\mu(y) \\ &= A + B. \end{aligned}$$

We estimate the integral A . Since $|y| \asymp r$, similarly to (59) we obtain

$$\int_{7r/4}^{2r} P_{K(z_0, \sigma)}(0, y) d\sigma \leq cr^{1-d} \asymp r^{1+\alpha d_w/2} |y|^{-d_\alpha}.$$

It follows that

$$A \leq cr^{1+\alpha d_w/2} \int_{|y| \in (r, 4r)} u(y)|y|^{-d_\alpha} d\mu(y).$$

To estimate the integral B in (72), observe that for $|y| > 4r$ we have $|y - z_0(s - r)/r| > 3r > 2(\sigma + r)/2$ and by Proposition 6.4 we get

$$\int_{7r/4}^{2r} P_{K(z_0, \sigma)}(0, y) d\sigma \leq c \int_{7r/4}^{2r} ((\sigma + r)/2)^{\alpha d_w/2} |y|^{-d_\alpha} d\sigma \leq cr^{1+\alpha d_w/2} |y|^{-d_\alpha}.$$

It follows that

$$B \leq cr^{1+\alpha d_w/2} \int_{|y|>4r} u(y)|y|^{-d_\alpha} d\mu(y).$$

Consequently, we can estimate the mean value of $u_\sigma(0)$ over $\sigma \in (7r/4, 2r)$,

$$(4/r) \int_{7r/4}^{2r} u_\sigma(0) d\sigma \leq cr^{\alpha d_w/2} \int_{|y| \geq r} u(y)|y|^{-d_\alpha} d\mu(y),$$

which gives our claim (71).

By the assumption (67), we fix a point A such that $B(A, \theta r) \subseteq D \cap B(Q, r)$. Since u is regular α -harmonic in $B(A, \theta r)$ by (50) we obtain

$$\begin{aligned} u(A) &\geq \int_{|y-A|>\theta r} P_{B(A,\theta r)}(A, y)u(y) d\mu(y) \\ &\geq c(\theta r)^{\alpha d_w/2} \int_{|y-A|>\theta r} |y - A|^{-d_\alpha} u(y) d\mu(y) \\ &\geq c_2 r^{\alpha d_w/2} \int_{|y|>r} |y - A|^{-d_\alpha} u(y) d\mu(y), \end{aligned}$$

where $c_2 = c_2(\theta)$. If $|y| > r$ then $|A| < r < |y|$, and consequently $|y - A| \leq |y| + |A| \leq 2|y|$. Hence, by the above inequality and (71),

$$(73) \quad u(A) \geq cc_2 r^{\alpha d_w/2} \int_{|y|>r} |y|^{-d_\alpha} u(y) d\mu(y) \geq c_3 u_{\sigma_0}(0),$$

with $c_3 = c_3(\theta)$. From (70), Corollary 7.7 for the ball $B(0, \sigma_0)$ and $z \in B(0, r)$, and (73) it follows that

$$(74) \quad u(z) \leq u_{\sigma_0}(z) \leq cu_{\sigma_0}(0) \leq c_4 u(A), \quad z \in B(0, r),$$

where $c_4 = c_4(\theta)$. This ends the proof. ■

8.1. Boundary Harnack Principle. Below we present a proof of the Boundary Harnack Principle for the Sierpiński gasket. For the sake of convenience, we recall here briefly the construction of the set (we introduce an unbounded version). Let F_0 be the closed convex triangle with vertices at $(0, 0)$, $(1, 0)$ and $(1/2, \sqrt{3}/2)$. Let A be the interior of the triangle whose vertices are the midpoints of the edges of F_0 . Let $F_1 = F_0 \setminus A$. Then F_1 consists of three closed triangles of sides $1/2$. To obtain F_2 we apply the above procedure to the triangles in F_1 , and so on. Set

$$F_\infty = \bigcap_{n=0}^\infty F_n, \quad F = \bigcup_{n=0}^\infty 2^n F_\infty.$$

We call F the (unbounded) *Sierpiński gasket*. The collection of those triangles in $\bigcup_{k=0}^\infty 2^k F_{n+k}$ (of sides 2^{-n}) is denoted by \mathcal{S}_n , $n = 0, 1, \dots$. Note that F lies between the x -axis and the line $y = \sqrt{3}x$.

By a *natural cell* (or simply *cell*) we mean the intersection of F with a triangle from \mathcal{S}_n for some $n = 0, 1, \dots$.

We assume that our region D is the interior of the sum of a finite number of natural cells (possibly of different sizes). In other words, there exist $n_0 \in \mathbb{N} \cup \{0\}$, $n_i \in \mathbb{N} \cup \{0\}$, $i = 1, \dots, n_0$, and $S_i \in \mathcal{S}_{n_i}$ such that

$$D = \text{int} \left(F \cap \bigcup_{i=1}^{n_0} S_i \right).$$

Note that the interior is taken with respect to the topology of F (inherited from \mathbb{R}^N) and since S_i are closed, any two adjacent cells always make a connected set. It is clear that (64) and (67) hold for our D (with some $R_0 = R_0(D)$). Moreover, observe that the distance between any two disjoint cells in D is at least $R_1 = R_1(D) > 0$. We may and do assume that $R_0 < R_1$. Since in the proof of the Boundary Harnack Principle we consider local neighborhoods of a boundary point, it is enough to deal with a single cell.

The result can be stated as follows.

THEOREM 8.6 (Boundary Harnack Principle). *Let $0 < \alpha < 2/d_w$ or $d_s < \alpha < 2$. There exists a constant $c_1 = c_1(D)$ such that for all $Q \in \partial D$, $r \in (0, R_0)$, and functions $u, v \geq 0$ regular α -harmonic in $D \cap B(Q, 2r)$ which vanish on $D^c \cap B(Q, 2r)$ and satisfy $u(A_1) = v(A_1)$, where A_1 satisfying (67) is defined below, we have*

$$(75) \quad c_1^{-1}v(x) \leq u(x) \leq c_1v(x), \quad x \in D \cap B(Q, r/4).$$

Before the proof we make some clarifying remarks. Without losing generality we assume that our d -measure μ is the d -dimensional Hausdorff measure. As usual, we assume $Q = 0$ and let $\Omega = D \cap B(Q, r)$. For arbitrary $r \in (0, R_0)$ we can always find $\tilde{r} = 2^{-n}$, for some $n \in \mathbb{N}$, such that $r/2 \leq \tilde{r} < r$. Thus, it is enough to prove (75) for $x \in D \cap B(Q, \tilde{r}/2)$. Therefore, we may and do assume that r itself is of the form $r = 2^{-n}$ with n fixed. This implies that a ball with center at any vertex of any triangle from \mathcal{S}_n and of radius $2^{-i}r$, $i \in \mathbb{N}$, consists of triangles from \mathcal{S}_{n+i} (intersected with F). In particular, Ω is the intersection of D with a triangle from \mathcal{S}_n whose one vertex is at 0 (see Figure 1).




Fig. 1

Let $\Omega_1 = \frac{1}{2}\Omega = D \cap B(0, r/2)$ and set $\{A_1, A_2\} = \partial\Omega_1 \setminus \{0\}$ and $\{S_1, S_2\} = \partial\Omega \setminus \{0\}$, so that $S_i = 2A_i$, $i = 1, 2$. Define $R_i = B(S_i, r/2) \cap \Omega^c$, $i = 1, 2$, and $R = R_1 \cup R_2$. Moreover, let $T_i = B(S_i, 3r/4) \setminus (\Omega \cup R_i)$, $i = 1, 2$, and $T = T_1 \cup T_2$.

Proof of the Boundary Harnack Principle. Let u_1, u_2 be functions such that

$$u_1(y) = \begin{cases} u(y), & y \in R, \\ 0, & y \in \Omega^c \setminus R, \end{cases} \quad u_2(y) = \begin{cases} u(y), & y \in \Omega^c \setminus R, \\ 0, & y \in R, \end{cases}$$

and u_1 and u_2 are regular α -harmonic in Ω . Note that $u_1, u_2 \geq 0$ and $u_1 + u_2 = u$. We define v_1 and v_2 analogously. By the Harnack inequality for $B(S_i, r)$, $i = 1, 2$, we have

$$u(y) \asymp u(A_i), \quad y \in R_i \cup T_i, \quad i = 1, 2, \quad u(A_2) \asymp u(A_1).$$

Consequently,

$$(76) \quad u(y) \asymp u(A_1), \quad y \in R \cup T.$$

Fix $x \in \Omega$. From (76) it follows that

$$u_1(x) = E^x[u(X_{\tau_\Omega}); X_{\tau_\Omega} \in R] \asymp u(A_1)P^x[X_{\tau_\Omega} \in R].$$

From the analogous relation for v_1 and our assumption $u(A_1) = v(A_1)$ we get

$$(77) \quad u_1(x) \asymp v_1(x) \leq v(x), \quad x \in \Omega.$$

Since $v_2 = 0$ on R we have

$$(78) \quad \begin{aligned} v_2(A_1) &= E^{A_1}v_2(X_{\tau_\Omega}) \\ &= E^{A_1}[v(X_{\tau_\Omega}); X_{\tau_\Omega} \in T] \\ &\quad + E^{A_1}[v(X_{\tau_\Omega}); X_{\tau_\Omega} \in (\Omega \cup R \cup T)^c]. \end{aligned}$$

By the relation for v analogous to (76) we obtain

$$\inf_{z \in T} v(z) \geq \inf_{z \in T \cup R} v(z) \asymp v(A_1) \asymp \sup_{z \in T \cup R} v(z),$$

which yields

$$(79) \quad E^{A_1}[v(X_{\tau_\Omega}); X_{\tau_\Omega} \in T] \geq c \sup_{z \in T \cup R} v(z)P^{A_1}[X_{\tau_\Omega} \in T].$$

On the other hand,

$$\begin{aligned} v(A_1) &= E^{A_1}[v(X_{\tau_\Omega}); X_{\tau_\Omega} \in R \cup T] + E^{A_1}[v(X_{\tau_\Omega}); X_{\tau_\Omega} \in (\Omega \cup R \cup T)^c] \\ &\leq \sup_{z \in T \cup R} v(z)P^{A_1}[X_{\tau_\Omega} \in R \cup T] + E^{A_1}[v(X_{\tau_\Omega}); X_{\tau_\Omega} \in (\Omega \cup R \cup T)^c]. \end{aligned}$$

Observe that

$$P^{A_1}[X_{\tau_\Omega} \in R \cup T] \leq c_0P^{A_1}[X_{\tau_\Omega} \in T].$$

Indeed, for $y \in T_1$ we have $|A_1 - y| \leq 2r$ and by (50) we obtain

$$\begin{aligned} P^{A_1}[X_{\tau_\Omega} \in T] &\geq P^{A_1}[X_{\tau_{B(A_1, r/2)}} \in T_1] \geq cr^{\alpha d_w/2} \int_{T_1} |A_1 - y|^{-d_\alpha} d\mu(y) \\ &\geq cr^{\alpha d_w/2} (2r)^{-d_\alpha} \mu(T_1) \geq c_0^{-1}. \end{aligned}$$

Hence, we get

$$\begin{aligned} (80) \quad v(A_1) &\leq c \sup_{z \in T \cup R} v(z) P^{A_1}[X_{\tau_\Omega} \in T] \\ &\quad + E^{A_1}[v(X_{\tau_\Omega}); X_{\tau_\Omega} \in (\Omega \cup R \cup T)^c] \end{aligned}$$

Combining (78), (79) and (80) we get

$$(81) \quad v_2(A_1) \geq cv(A_1).$$

Let $K = \Omega \cup R \cup (D^c \cap B(0, 2r))$. Observe that for $z \in \Omega$ and $y \in K^c$ we have $|y - z| \asymp |y|$. Therefore, for $x \in \Omega$ we obtain

$$\begin{aligned} u_2(x) &= \int_{K^c} P_\Omega(x, y) u(y) d\mu(y) \\ &\asymp \int_{K^c} \left(\int_\Omega G_\Omega(x, z) |z - y|^{-d_\alpha} d\mu(z) \right) u(y) d\mu(y) \\ &\asymp \int_{K^c} \left(\int_\Omega G_\Omega(x, z) d\mu(z) \right) u(y) |y|^{-d_\alpha} d\mu(y) \\ &= E^x \tau_\Omega \int_{K^c} u(y) |y|^{-d_\alpha} d\mu(y) \end{aligned}$$

and the analogous relation for v_2 . It follows that

$$(82) \quad u_2(x)/u_2(A_1) \asymp E^x \tau_\Omega / E^{A_1} \tau_\Omega \asymp v_2(x)/v_2(A_1).$$

Denote the last quotient by q_0 . Then, by (82), the definition of u_2 , the assumption $u(A_1) = v(A_1)$ and (81),

$$u_2(x) \leq cq_0 u_2(A_1) \leq cq_0 u(A_1) = cq_0 v(A_1) \leq cq_0 v_2(A_1) = cv_2(x), \quad x \in \Omega.$$

Together with (77) and symmetry this completes the proof. ■

REMARK 5. We want to emphasize that in the particular case of the Sierpiński gasket, the Boundary Harnack Principle is a consequence of the Harnack inequality alone. It seems that a similar approach works for *nested fractals*. On the other hand the Boundary Harnack Principle for the Sierpiński carpet should be available by other, more complicated methods used in [B] for Lipschitz domains in \mathbb{R}^N .

Acknowledgements. The authors wish to thank T. Byczkowski, J. Dziubański and P. Graczyk for helpful discussions and comments. We are also grateful to an anonymous referee for useful comments and clarifying remarks.

Note added in proof. After the paper was submitted we learned that Z.-Q. Chen and T. Kumagai [CK] studied general stable-like processes on fractals defined by means of appropriate Dirichlet forms. Their paper contains very interesting results, which partially overlap ours; however, their methods of proof are completely different, based on tightness results and the parabolic Harnack inequality.

References

- [Ba] M. T. Barlow, *Diffusions on fractals*, in: Lectures on Probability Theory and Statistics, École d'Été de Probabilités de Saint-Flour XXV – 1995, Lecture Notes in Math. 1690, Springer, New York, 1998, 1–121.
- [BB] M. T. Barlow and R. F. Bass, *The construction of Brownian motion on the Sierpiński carpet*, Ann. Inst. H. Poincaré Probab. Statist. 25 (1989), 225–257.
- [BB1] —, —, *Brownian motion and harmonic analysis on Sierpiński carpets*, Canad. J. Math. 51 (1999), 673–744.
- [BP] M. T. Barlow and E. A. Perkins, *Brownian motion on the Sierpiński gasket*, Probab. Theory Related Fields 79 (1988), 543–623.
- [Bs] R. F. Bass, *Probabilistic Techniques in Analysis*, Springer, New York, 1995.
- [BL] R. F. Bass and D. A. Levin, *Harnack inequalities for jump processes*, Potential Anal. 17 (2002), 375–388.
- [Be] J. Bertoin, *Lévy Processes*, Cambridge Univ. Press, Cambridge, 1996.
- [BG] R. M. Blumenthal and R. K. Gettoor, *Markov Processes and Potential Theory*, Pure Appl. Math. 29, Academic Press, New York, 1968.
- [B] K. Bogdan, *The boundary Harnack principle for the fractional Laplacian*, Studia Math. 123 (1997), 43–80.
- [BBy] K. Bogdan and T. Byczkowski, *Probabilistic proof of boundary Harnack principle for α -harmonic functions*, Potential Anal. 11 (1999), 135–156.
- [BSS] K. Bogdan, A. Stós and P. Sztonyk, *Harnack inequality for symmetric stable processes on fractals*, C. R. Acad. Sci. Paris Sér. I 335 (2002), 59–63.
- [CK] Z.-Q. Chen and T. Kumagai, *Heat kernel estimates for stable-like processes on d -sets*, preprint, 2002.
- [CS] Z.-Q. Chen and R. Song, *Intrinsic ultracontractivity and conditional gauge for symmetric stable processes*, J. Funct. Anal. 159 (1998), 267–294.
- [ChZ] K. L. Chung and Z. Zhao, *From Brownian Motion to Schrödinger's Equation*, Springer, New York, 1995.
- [DSV] K. Darlymple, R. Strichartz and J. P. Vinson, *Fractal differential equations on the Sierpiński gasket*, J. Fourier Anal. Appl. 5 (1999), 203–284.
- [D] G. Doetsch, *Introduction to the Theory and Application of the Laplace Transformation*, Springer, New York, 1974.
- [Fa1] K. Falconer, *The Geometry of Fractal Sets*, Cambridge Univ. Press, Cambridge, 1992.
- [Fa2] K. Falconer, *Fractal Geometry. Mathematical Foundations and Applications*, Wiley, Chichester, 1990.
- [FJ] W. Farkas and N. Jacob, *Sobolev spaces on non-smooth domains and Dirichlet forms related to subordinate reflecting diffusions*, Math. Nachr. 224 (2001), 75–104.
- [FOT] M. Fukushima, Y. Oshima and M. Takeda, *Dirichlet Forms and Symmetric Markov Processes*, de Gruyter, Berlin, 1994.

- [FU] M. Fukushima and T. Uemura, *On Sobolev and capacity inequalities for contractive Besov spaces over d -sets*, Potential Anal. 18 (2003), 59–77.
- [G] A. Grigor'yan, J. X. Hu and K.-S. Lau, *Heat kernels on metric measure spaces and an application to semilinear elliptic equations*, Trans. Amer. Math. Soc. 355 (2003), 2065–2095.
- [H] J. Hawkes, *A lower Lipschitz condition for the stable subordinator*, Z. Wahrsch. Verw. Geb. 17 (1971), 23–32.
- [HL] J. X. Hu and K.-S. Lau, *Riesz potentials and Laplacians on fractals*, preprint.
- [IW] N. Ikeda and S. Watanabe, *On some relations between the harmonic measure and the Lévy measure for a certain class of Markov processes*, J. Math. Kyoto Univ. 2 (1962), 79–95.
- [JK] D. S. Jerison and C. E. Kenig, *Boundary value problems on Lipschitz domains*, in: W. Littman (ed.), Studies in Partial Differential Equations, MAA Stud. Math. 23, Math. Assoc. Amer., Washington, DC, 1982, 1–68.
- [J] A. Jonsson, *Brownian motion on fractals and function spaces*, Math. Z. 222 (1996), 495–504.
- [JW] A. Jonsson and H. Wallin, *Function Spaces on Subsets of \mathbb{R}^N* , Math. Rep. 2, no. 1, Harwood, London, 1984.
- [Kum] T. Kumagai, *Some remarks for stable-like jump processes on fractals*, in: Proc. Conf. held in Graz, 2001, Birkhäuser, 2002, 185–196.
- [L] N. S. Landkof, *Foundations of Modern Potential Theory*, Springer, Berlin, 1969.
- [P] K. Pietruska-Pałuba, *On function spaces related to fractional diffusions on d -sets*, Stochastics Stochastics Rep. 70 (2000), 153–164.
- [PS] S. C. Port and C. J. Stone, *Infinitely divisible processes and their potential theory*, Ann. Inst. Fourier (Grenoble) 21 (1971), no. 2, 157–275, and no. 4, 179–265.
- [R] M. Ryznar, *Estimates of Green function for relativistic α -stable process*, Potential Anal. 17 (2002), 1–23.
- [SW] R. Song and J. M. Wu, *Boundary Harnack principle for symmetric stable processes*, J. Funct. Anal. 168 (1999), 403–427.
- [S] A. Stós, *Symmetric stable processes on d -sets*, Bull. Polish Acad. Sci. Math. 48 (2000), 237–245.
- [Str] R. Strichartz, *Analysis on fractals*, Notices Amer. Math. Soc. 46 (1999), 1199–1208.

Institute of Mathematics
 Wrocław University of Technology
 Wybrzeże Wyspiańskiego 27
 50-370 Wrocław, Poland
 E-mail: bogdan@im.pwr.wroc.pl
 stos@im.pwr.wroc.pl
 sztonyk@im.pwr.wroc.pl

Received September 30, 2002
Revised version March 21, 2003

(5045)