

Health benefits of resveratrol administration

Sabina Galiniak[✉], David Aebisher and Dorota Bartusik-Aebisher

Faculty of Medicine, Rzeszów University, Rzeszów, Poland

Resveratrol is a polyphenol that is abundant in grape skin and seeds. Food sources of resveratrol include wine, berries, and peanuts. This compound has many properties, including activity against glycation, oxidative stress, inflammation, neurodegeneration, several types of cancer, and aging. Because resveratrol is generally well-tolerated, it is believed to be a promising compound in preventing many diseases, such as diabetes and its complications. Unfortunately, this compound exhibits low bioavailability and solubility. The aim of this review is to summarize the latest information on the multiple effects of resveratrol on health and the benefits of its intake, based on *in vitro* and *in vivo* studies in animals and humans.

Key words: resveratrol, glycation, oxidative stress, polyphenol


Received: 01 December, 2018; **revised:** 29 January, 2019; **accepted:** 01 February, 2019; **available on-line:** 28 February, 2019

[✉]e-mail: sgaliniak@ur.edu.pl

Abbreviations: AGEs, advanced glycation end products; MGO, methylglyoxal; GO, glyoxal; HSA, human serum albumin; BSA, bovine serum albumin; AOPP, advanced oxidation protein products; ROS, reactive oxygen species; MDA, malondialdehyde; CAT, catalase; SOD, superoxide dismutase; IL, interleukin; TNF- α , tumor necrosis factor α ; NF κ B, nuclear factor κ B; I κ B, inhibitor of κ B; JAK/STAT, Janus kinases/signal transducer and activator of transcription; PI3K, phosphatidylinositol-3-kinases; Akt, protein kinase B; Bcl-2, B-cell lymphoma 2; Bax, Bcl-2-associated X protein

INTRODUCTION

Resveratrol (3,5,4'-trans-trihydroxystilbene) is a polyphenolic phytoalexin belonging to the stilbene family. It is a natural dietary plant compound that occurs mainly in grape skin and seeds but is also found in wines and various other types of plant foods, especially peanuts, berries, and tea (Shrikanta *et al.*, 2015). Resveratrol is synthesized by more than 70 species of plants in response to infection, stress, injury, bacteria or fungal infections, and UV-irradiation (Hasan & Bae, 2017). Synthesis of this molecule in plants is catalysed by resveratrol synthase in the phenylpropanoid pathway in a process similar to that of flavonoids (Kapetanovic *et al.*, 2011). Resveratrol was first reported and isolated from white hellebore by a Japanese researcher Takaoka in 1939 (Takaoka, 1939). Resveratrol possesses two phenol rings (monophenol and diphenol) bonded together by a double styrene bond and it exists in both cis and trans isomeric forms. *Trans*-resveratrol (Scheme 1) appears to be the more abundant and stable natural form (Gambini *et al.*, 2015). This molecule has three hydroxyl groups which are involved in free radical scavenging and metal chelation (Caruso *et al.*, 2004; Gülçin, 2010). The presence of hydroxyl groups also facilitates interaction with macromolecules.




Scheme 1. Structure of *trans*-resveratrol.

Resveratrol reveals a wide range of biological properties including anti-glycation, antioxidant, anti-inflammation, neuroprotective, anti-cancer, and anti-aging activity in various *in vitro* and *in vivo* experimental models (Öztürk *et al.*, 2017; Rauf *et al.*, 2017; Jardim *et al.*, 2018; Li *et al.*, 2018; Salehi *et al.*, 2018). Currently, resveratrol research spans a large spectrum of areas and is a subject of considerable scientific attention, in particular, investigations of its biological activity and the effects of resveratrol administration in numerous diseases in both clinical and nonclinical trials (Scheme 2).

SELECTED BIOLOGICAL PROPERTIES OF RESVERATROL

Antiglycation activity

Glycation is a nonenzymatic reaction between proteins and reducing sugars, leading to the formation of advanced glycation end products (AGEs). Advanced glycation end products accumulate and cause damage at the tissue and cellular level, including lipid peroxidation, endothelial dysfunction, changes in protein structure, and



Scheme 2. The activities of resveratrol on health.

stimulation of inappropriate cellular activity (Singh *et al.*, 2001; Galiniak *et al.*, 2017). Glycation produces highly reactive dicarbonyl compounds such as methylglyoxal and glyoxal, which are key precursors to the formation of AGEs. Methylglyoxal (MGO) is also generated by glycolysis, glucose autooxidation, and lipid peroxidation, while glyoxal (GO) is formed during lipid peroxidation and degradation of monosaccharide and saccharide derivatives. Both MGO and GO are known to enhance oxidative stress in the human body. Recent reports suggest that increased glycation is associated with a higher prevalence of diabetes and related complications (Adamska *et al.*, 2018; Rhee & Kim, 2018), lung (Khan *et al.*, 2018) and breast cancer (Walter *et al.*, 2018), myasthenia gravis (Adamczyk-Sowa *et al.*, 2017), and neurodegenerative diseases (Pinkas & Aschner, 2016).

Increasingly, evidence indicates that compounds of natural origin, especially polyphenols, have antiglycation properties. Among effective glycation inhibitors are phenolic acids, naringin, genistein, rutin, quercitrin, and kaempferol (Sadowska-Bartosz *et al.*, 2014; Yeh *et al.*, 2017). Many reports have confirmed that resveratrol displays antiglycation activity as well.

A high percentage (up to 99%) of resveratrol-induced inhibition of AGE production was demonstrated by Shen and others (Shen *et al.*, 2017) in BSA/fructose, BSA/MGO, and arginine/MGO mixtures. Resveratrol inhibition of AGE and binding to toxic MGO and GO have demonstrated both anti-oxidative and pro-oxidative effects, as resveratrol adducts can lead to the oxidation of amino acids in human serum albumin (HAS) (Arcanjo *et al.*, 2018). Furthermore, resveratrol reveals antiglycation activity against the harmful effect of AGEs or α -dicarbonyls on porcine chondrocytes (Liu *et al.*, 2010), mouse oocytes (Liu *et al.*, 2013), and dendritic cells obtained from peripheral blood mononuclear cells (Buttari *et al.*, 2013).

Yilmaz and others (Yilmaz *et al.*, 2017) has shown that administration of resveratrol in drinking water to chronic MG-treated rats significantly reduces the level of advanced oxidation protein products (AOPP), AGEs, and protein carbonyl in plasma, as well as markers of oxidative stress in the liver. A significant decrease in urine albumin and creatinine and an increase in serum antioxidant enzymes were also observed in patients with type II diabetes and diabetic nephropathy who received resveratrol at the dose of 500 mg per day (Şattarinezhad *et al.*, 2018). Resveratrol prevents opacification and formation of polyols in the bovine lens, as well as ameliorates kidney function due to suppression of AGEs formation, suggesting that resveratrol may be considered as a protective agent against diabetic complications such as cataracts and nephropathy (Ciddi & Dodda, 2014; Hussein & Mahfouz, 2016). Furthermore, resveratrol is a prospective therapeutic agent against diabetic ototoxicity (Erkan *et al.*, 2018), renal fibrosis (He *et al.*, 2016), progression of cataracts (Higashi *et al.*, 2018), and bone density loss in patients with type 2 diabetes (Bo *et al.*, 2018). Likewise, resveratrol is also involved in protection against diabetic cardiomyopathy development *via* improvement of mitochondrial function and inhibition of apoptosis of cardiomyocytes (Diao *et al.*, 2018). Resveratrol preserves pancreatic tissue by reducing inflammatory factors and glucose levels in serum, and ultimately leads to the protection of cardiovascular tissues in diabetic rat models of coronary heart disease (Huo *et al.*, 2019).

Similarly, it has been shown that by modulating the switching between apoptosis and autophagy resveratrol has a beneficial effect on cardiomyopathy induced in car-

diac myoblast cells exposed to the combination of high glucose and palmitate (Xu *et al.*, 2018). Guzmán and others (Guzmán *et al.*, 2018) noted that resveratrol at a dose of 1, 10 and 100 μ M has a protective effect on acute high glucose-induced damage in endothelial cells. Moreover, resveratrol attenuates methylglyoxal induced endothelial damage by promoting the expression and activity of endothelial nitric oxide synthase in thoracic aorta in older rats (Tasatargil *et al.*, 2018). In diabetic mice receiving resveratrol, a decrease in the level of apoptosis of glomerular podocytes and renal tubular epithelial cells was noted (Zhang *et al.*, 2018). The diabetic rats treated with resveratrol had reduced levels of factors related to endoplasmic reticulum stress which underlies the progression of diabetic nephropathy (Yuan *et al.*, 2018).

In summary, the results indicate that resveratrol may be considered as a beneficial anti-glycation agent in *in vitro* and *in vivo* experiments as well as in the treatment of diseases associated with increased glycation.

Antioxidant activity

Oxidative stress is defined as an imbalance between generation of reactive oxygen species and the antioxidant defence in favor of oxidant production. Enhanced oxidative stress damages macromolecules and impairs their functions, which underlies many age-related diseases including cancer, diabetes, chronic kidney disease, cardiovascular and neurodegenerative diseases (Liguori *et al.*, 2018). Moreover, overproduction of ROS induces inflammation, dysregulation of mitochondria and cell death (Wu *et al.*, 2018). Also, it was shown that autooxidation of glucose may contribute to excessive ROS production and intensification of oxidative stress (Matough *et al.*, 2012). Resveratrol is confirmed to be a powerful antioxidant which activity is associated with presence of three hydroxyl groups in its structure. Resveratrol has an inhibitory effect on excessive ROS production, aberrant mitochondrial distribution, and lipid peroxidation (Liu *et al.*, 2013; Liguori *et al.*, 2018). Treatment of primary epidermal keratinocytes with resveratrol leads to a 1.3-fold increase of endogenously generated glutathione and quantitative reduction of the cellular redox environment and endogenous ROS production (Plauth *et al.*, 2016). In astroglial cells treated with ammonia, resveratrol prevents both an increase in ROS production and a decrease of mitochondrial membrane potential, which indicates a role in maintaining cellular redox homeostasis (Bobermin *et al.*, 2018). In fibroblasts exposed to rotenone, resveratrol decreases mitochondria fragmentation and maintains the potential of the mitochondrial membrane, as well as prevents the attenuation of oxidative phosphorylation, thus exerting a protective effect against the harmful impact of ROS (Sgarbi *et al.*, 2018). Recent study by Cheng and others (Cheng *et al.*, 2019) revealed that resveratrol prevents hepatic steatosis in obese mice fed a high-fat diet by reducing oxidative stress and inflammation. The protective effect of resveratrol against hepatic steatosis is supported by lowering the accumulation of lipid droplets in hepatocytes (Zhou *et al.*, 2018).

Furthermore, resveratrol treatment of the epithelial cells protects them from oxidative damage induced by hydrogen peroxide and promotes the expression and phosphorylation of occludin and other zonula occludens proteins. This protection is caused by reducing MDA and intracellular ROS concentration and increasing expression levels of antioxidant enzymes (Wang *et al.*, 2016). The protective effects of chronic administration of resveratrol against AGE-induced oxidative stress and

apoptosis, together with the improvement of glucose tolerance, are also observed in pancreatic cells in mice with type 2 diabetes (Lee *et al.*, 2012; Ginés *et al.*, 2017).

Resveratrol intake by diabetic rats at a dose of 5 mg/kg/day leads to normalization of antioxidant status, exacerbated by oxidative stress induced by hyperglycemia (Hussein & Mahfouz, 2016). Administration of resveratrol at a dose of 10 or 20 mg/kg for 4 weeks in rats with a streptozotocin-induced diabetes results in a reduction in the content of AOPP and MDA as well as catalase (CAT) and superoxide dismutase (SOD) activity in the lens compared to untreated diabetic rats (Sedlak *et al.*, 2018). Resveratrol-treated animals have reduced production of ROS, elevated membrane potential, and inhibition of cytochrome c release from the inner mitochondrial membrane (Zhang *et al.*, 2018). Wang and others (Wang *et al.*, 2018) observed that resveratrol counteracted the accumulation of 3-nitrotyrosine and generation of 4-hydroxynonenal during the development of diabetes mellitus-induced cardiomyopathy. Resveratrol-treated rats with diabetes have a significant decrease in the MDA level and total oxidant level, as well as an increase in total antioxidant capacity (Moridi *et al.*, 2015; Khazaei *et al.*, 2016) when compared to untreated groups. Moreover, resveratrol protects the spinal cord from ischemic damage in rats by reducing plasma levels of nitrite, AOPP and MDA, and increasing the enzymatic activity of SOD and CAT (Fu *et al.*, 2018). Resveratrol also attenuates oxidative stress in rats with experimental periodontitis (Corrêa *et al.*, 2018), induced early Alzheimer's disease (Lin *et al.*, 2018) and chronic obstructive pulmonary disease (Wang *et al.*, 2017).

The studies mentioned above indicate that resveratrol has a therapeutic effect in cell and animal experiments involving increased oxidative stress, which is associated to the limitation of ROS generation and stimulation of compounds that act as an antioxidant barrier.

OTHER BIOLOGICAL ACTIVITIES OF RESVERATROL

Anti-inflammatory activity

Resveratrol suppresses IL-6 transcription and translation, resulting in attenuation of its secretion by macrophages (Ohtsu *et al.*, 2017). Likewise, the administration of resveratrol to monocyte cultures leads to a reduction in the expression of inflammatory mediators: TNF- α and IL-8, without inducing cytotoxicity (Pinheiro *et al.*, 2018). Resveratrol significantly inhibits the production of extracellular matrix proteins by pancreatic stellate cells, which are involved in the development of pancreatic fibrosis (Xia *et al.*, 2018). Furthermore, resveratrol is involved in the inhibition of toll-like receptors, which in their active form can induce proinflammatory cytokines and chemokine expression and stimulate the activation of innate and adaptive immunity (Chen *et al.*, 2018). Resveratrol reduces matrix-metalloprotease expression and suppresses the production of IL-1, IL-6 and TNF- α in a dose dependent manner in chondrocytes with induced osteoarthritis (Li *et al.*, 2018). Moreover, resveratrol treatment in patients after oral implantology reduces serum levels of IL-1 β , IL-17A and TNF- α while the levels of IL-2, IL-6 and IL-10 are elevated (BaGen *et al.*, 2018). Ma and others (Ma *et al.*, 2015) observed that resveratrol effectively suppresses NF- κ B signaling through inhibiting the activities of NF- κ B and I κ B kinase, as well as by suppressing the phosphorylation of JAK/STAT signalling pathways. Resveratrol shows protective activity against intestinal

ischemia-reperfusion injury by inhibiting mast cells from degranulation and decreasing apoptosis of intestinal epithelial cells, which prevents overall organ dysfunction (Zhao *et al.*, 2018).

Anti-inflammatory activities of resveratrol are observed also in case of hyper-acute small intestinal inflammation (Bereswill *et al.*, 2010) as well as in immune-mediated diseases (Švajger & Jeras, 2012). Moreover, it prevents acceleration of cholesterol accumulation and disturbances of macrophage lipid homeostasis after induction by glycation products (Zhang *et al.*, 2010).

Neuroprotection

Resveratrol is also involved in the reduction of neuronal damage and apoptosis and the improvement of the central nervous system function. Resveratrol has been shown to reduce neurodegeneration in the murine cerebral cortex and enhance memory recovery after exposure to fluoride (Sharma *et al.*, 2018). Also, the administration of resveratrol improves cognition, learning and memory in rats with vascular dementia (Ma *et al.*, 2013). Report by Corpas and others (Corpas *et al.*, 2018) indicates that resveratrol also improves cognition and induces neuroprotection in amyloid and tau pathologies in mice models of Alzheimer's disease. However, results of resveratrol supplementation in human are inconsistent. A meta-analysis of randomized controlled trials suggests that resveratrol may be beneficial (Marx *et al.*, 2018) or may have no significant impact on the selected measures of cognitive performance (Farzaei *et al.*, 2018). Several neuroprotective properties of resveratrol have been suggested in the studies of its effects in the intracerebral hemorrhage (Bonsack *et al.*, 2017), cerebral neurodamage (Nalagani & Karnati, 2016), and central nervous system injuries such as stroke (Lopez *et al.*, 2015). Likewise, resveratrol shows a neuroprotective effects in cerebral ischemia/reperfusion injury in rat brain by reducing the cerebral infarct volume and stimulating the expression of components of the intracellular signaling pathway including kinases such as JAK2, PI3K or Akt and anti-apoptotic molecules, while down-regulating the expression of pro-apoptotic caspase-3 and Bax (Hou *et al.*, 2018).

Resveratrol also alleviates neuropathic pain in mice through repressing the expression of proinflammatory cytokines and increasing the expression of anti-inflammatory IL-10 (Tao *et al.*, 2016). Similarly, the observed beneficial effect of resveratrol on hyperalgesia in rats with chronic neuropathic pain is due to the inhibition of the expression of glial fibrillary acidic protein and the P2X7 receptor, a key player in nervous pathological pain (Xie *et al.*, 2017). In addition, resveratrol at the dose of 10-80 mg/kg per day may be an effective treatment for depression in animal models (Moore *et al.*, 2018). There is also evidence that the neuroprotective effect is also enhanced by the antioxidant and anti-inflammatory properties of resveratrol.

Anti-cancer activity

Many *in vitro* and *in vivo* studies suggest that resveratrol has anti-cancer properties due to its wide range of activities, including antioxidant effects and regulating the expression of pro-apoptotic proteins, as well as molecules underlying tumor development. Resveratrol is known to reduce the incidence and development of various types of cancer, such as cervical (Zhou *et al.*, 2018), pancreatic (Zhao *et al.*, 2018), gastric (Wu *et al.*, 2018), breast and colorectal (Lucas *et al.*, 2018), as well as thyroid cancer

(Zheng *et al.*, 2018). Research results indicate that resveratrol has protective effect on the normal cells, while inducing death in cancer cells. This can be associated with different cellular targets and metabolic pathways of resveratrol in healthy and cancerous cells. In addition, this dual pattern of resveratrol action depends on the dose. Lower concentrations increase expression of cell survival proteins, whereas higher doses stimulate cell apoptosis or necrosis regardless of whether the cell is healthy or pathological (Szende *et al.*, 2010; San Hipólito-Luengo *et al.*, 2017). Resveratrol at a high dose inhibits synthesis of nucleic acids and proteins, leads to impairment of chromatin structure, and finally, causes cell death (Mukherjee *et al.*, 2010).

Monteillier and others (Monteillier *et al.*, 2018) showed that intranasal administration of 60 mg/kg resveratrol to mice with induced lung cancer caused a notable decrease in the tumor multiplicity and volume via enhanced apoptosis. Moreover, treating gastric cancer cells with resveratrol elevated the levels of pro-apoptotic proteins such as Bax, while the levels of anti-apoptotic proteins such as Bcl-2 were decreased compared to untreated controls (Wu *et al.*, 2018). Treatment of osteosarcoma cells with 20 μ M resveratrol results in reduction of cell viability, decrease in self-renewing and tumorigenesis via inhibition of intracellular STAT3 signaling and cytokine synthesis (Peng & Jiang, 2018). In general, the anti-cancer activity of resveratrol is based on the suppression of the expression of proteins involved in carcinogenesis, such as phospholipid scramblase 1 (Zhou *et al.*, 2018), stimulation of caspase 3 cleavage (Lucas *et al.*, 2018) and activation of the mitochondrial ROS signaling pathway (Zheng *et al.*, 2018). Furthermore, resveratrol inhibits the cell cycle by inducing S-phase arrest in gastric cancer cells in a dose-dependent manner (Wu *et al.*, 2018). Resveratrol has also been found to increase the effect of anticancer drugs and decrease drug resistance in cancer cells (Halajian *et al.*, 2018; Pouyafar *et al.*, 2019). Taken together, *in vivo* and *in vitro* studies confirm the beneficial antitumor effect of resveratrol.

Anti-ageing activity

Resveratrol has been reported to extend life span in several different subjects including the fish species *Nothobranchius furzeri* (Valenzano *et al.*, 2006) and *Nothobranchius guentheri* (Liu *et al.*, 2015), as well as *Drosophila melanogaster* (Wang *et al.*, 2013), honey bee (Rascon *et al.*, 2012) and mice (Baur *et al.*, 2006). Roggerio and others (Roggerio *et al.*, 2018) observed that administration of resveratrol (500 mg/day) in healthy and slightly overweight subjects resulted in higher gene expression and serum concentration of sirtuin-1. Sirtuins exhibit a broad spectrum of activity, including anti-ageing and anti-inflammatory effects, inhibition of degenerative disorders such as liver steatosis, as well as improvement of endothelial function, and prevention of cancer (Wątroba & Szukiewicz, 2016). Recent report shows that short-term injection of resveratrol in postovulatory oocyte delayed the aging process of oocytes in middle-aged mice by promoting the expression of sirtuin-1, reducing ROS generation, and ameliorating mitochondrial function (Liang *et al.* 2018). Generally, the mechanisms of resveratrol longevity activity are similar to caloric restriction (Bass *et al.*, 2007).

Despite a number of studies that demonstrate the life-extending activity of resveratrol, its anti-ageing properties are still controversial due to the growing number of reports to the contrary. Recently, Ramos-Gomez and others

(Ramos-Gomez *et al.*, 2017) described that resveratrol caused mitochondrial dysfunction and reduction in the chronological life-span of *Saccharomyces cerevisiae*. Similarly, it is known that dietary resveratrol does not extend the life span of the mosquito *Anopheles stephensi* (Johnson & Riehle, 2015) nor *Drosophila melanogaster* and does not influence gene expression of longevity-associated and antioxidant enzymes (Staats *et al.*, 2018). Wang *et al.* (2013) suggested that the life-extending effect of resveratrol depends on dietary composition, dose and gender. Their results indicate that resveratrol at a dose of 400 μ M has a prolongevity effect on females of *Drosophila melanogaster* fed a high fat diet. However, a lower resveratrol (or fat?) concentration had no effect on the lifespan of female flies fed a (fat? calorie?) restricted diet or diet rich in sugar and protein. Similarly, it was indicated that the life-extending activities of resveratrol depended on the model organism and its genetic background (Pallauf *et al.*, 2016).

Besides the aforementioned effects of resveratrol, it exhibits additional beneficial activities on other targets at the cellular and tissue levels. A report by Hara and others (Hara *et al.*, 2018) suggests that resveratrol attenuates changes induced in bovine embryos composed of 8–12 cells by vitrification due to degradation of damaged mitochondria, without affecting ATP content and activation of further embryonic development. Gorga and others (Gorga *et al.*, 2018) observed that resveratrol decreased expression of cyclins and activated sirtuin 1, which lead to regulation of immature Sertoli cells proliferation. Resveratrol also acts as an activator of Notch signalling and an inhibitor of endothelial cell proliferation and migration (LaFoya *et al.*, 2019).

Resveratrol induces neuronal differentiation in murine neuroblastoma cells (Namsi *et al.*, 2018) and differentiation of monocytes to macrophages (Vasamsetti *et al.*, 2016). Recent reports show that resveratrol increases self-renewal and maintains the pluripotency of human and mouse embryonic stem cells (Li *et al.*, 2017; Safa-einejad *et al.*, 2017). Intra-gastric administration of resveratrol causes activation of cardiac stem cells, an increase of capillary density, and reduction of apoptosis of cardiomyocytes, which may be beneficial in myocardial regeneration after acute myocardial infarction (Ling *et al.*, 2017). Furthermore, by reducing the expression of perlipin 5, resveratrol accelerates lipid hydrolysis in brown adipose tissue, which may cause a decrease in weight and myocardial steatosis of heart tissue (Mehdi *et al.*, 2018). Resveratrol also exhibits antimicrobial activity against *Haemophilus influenzae* (Euba *et al.*, 2017), *Escherichia coli* (Subramanian *et al.*, 2014), *Propionibacterium acnes* (Taylor *et al.*, 2014), and *Staphylococcus aureus* (Wu & Huang, 2017). Toniolo *et al.* (2018) observed that supplementation with 0.04% resveratrol for six months also improves the resistance to fatigue and functional-mechanical properties of skeletal muscles in aged mice. Furthermore, resveratrol at a dose of 500 mg per day attenuates joint pain and improves functional activity of patients with knee osteoarthritis (Hussain *et al.*, 2018; Marouf *et al.*, 2018). Farrokhi and others (Farrokhi *et al.*, 2018) demonstrated that 120 μ M resveratrol reduced production of matrix metalloproteinase 9, which makes it a potential therapeutic agent in atherosclerosis. Resveratrol also has a beneficial effect in counteracting allergic asthma (Alharris *et al.*, 2018), as well as anxiety and depression disorders (Liu *et al.*, 2019). Moreover, resveratrol improves testosterone levels and sperm parameters by reducing apoptosis in testes (Shatti, 2018).

Unfortunately, despite numerous reports presenting a beneficial effect of resveratrol treatments on many diseases or pathological states, it has poor bioavailability and water solubility (less than 0.05 mg/ml). Reports indicate that a 25 mg intake of resveratrol resulted in plasma concentrations lower than 10 ng/ml, while concentrations of 500 ng/ml in plasma were measured after a high dose of 5000 mg (Walle, 2011). Sergides and others (Sergides *et al.*, 2016) showed that intake of 500 mg of resveratrol in the form of tablets was well-tolerated and led to plasma concentrations of about 70 ng/ml. Other studies also confirm that administration of resveratrol is generally well-tolerated and safe (Sergides *et al.*, 2016; Berman *et al.*, 2017), although some adverse effects were reported when resveratrol was given in high doses (Novelle *et al.*, 2015). Diarrhoea, nausea, anemia, vomiting, flatulence, abdominal discomfort as well as renal failure via cast and crystal nephropathy and acute tubular damage are among the most common side effects of resveratrol, however, the nephrotoxic effect of resveratrol occurred only in patients with multiple myeloma (Brown *et al.*, 2010; la Porte *et al.*, 2010; Popat *et al.*, 2013). A recent study showed that incubation of human placental explants with resveratrol at a dose up to 100 μ M led to impairment of fatty acid uptake and oxidation of placental tissue, which may negatively affect fetal development (Landau *et al.*, 2017). Kumaran and others (Kumaran *et al.*, 2018) also demonstrated that oral resveratrol tablet taken once a day induced thrombocytopenia in woman with melasma.

Resveratrol has a short half-life of approximately 1.5 h due to rapid absorption in the intestine and degradation in the liver (Marier *et al.*, 2002). After consumption, 77–80% of resveratrol is absorbed into the blood stream by active transport via intestinal epithelial cells, after which it binds to albumin and lipoproteins. This polyphenol is easily released from the complexes and can be transported into cells. About 49–61% of resveratrol is excreted in the urine (Soleas *et al.*, 2001).

To improve the bioavailability of this compound, more complex formulations have been prepared, including nanoparticles and nanostructured lipid carriers containing resveratrol (Gokce *et al.*, 2012; Peñalva *et al.*, 2018). Enhancing resveratrol bioavailability was also attempted in rats by treating them with 3,5,4'-tri-O-acetylresveratrol (TARÉS), an acetylated resveratrol prodrug that can be enzymatically hydrolyzed to free trans-resveratrol in cells (de Vries *et al.*, 2018).

CONCLUSION

Overall, resveratrol has potential benefits for human health and exhibits protective effects against glycation, free radicals' production, neurodegeneration, inflammation, and tumor development. Life-extending properties of resveratrol are still controversial, however, there is evidence suggesting anti-ageing activity. The main mechanisms of resveratrol activity are based on the prevention of apoptosis and ROS production by downregulation of expression of anti- or proapoptotic proteins involved in antioxidant barrier systems, and improvements in mitochondrial function.

Nonetheless, more research, especially clinical trials on large number of patients are still needed for unambiguous confirmation of its positive action. Future research should also focus on improving resveratrol bioavailability

and counteracting any adverse effects after administration.

REFERENCES

- Adamczyk-Sowa M, Bieszcza-Bedrejczuk E, Galiniak S, Rozmilowska I, Czyżewski D, Bartosz G, Sadowska-Bartosz I (2017) Oxidative modifications of blood serum proteins in myasthenia gravis. *J Neuroimmunol* **305**: 145–153. <https://doi.org/10.1016/j.jneuroim.2017.01.019>
- Adamska A, Araszkievicz A, Pilacinski S, Gandecka A, Grzelka A, Kowalska K, Malinska A, Nowicki M, Zozulinska-Ziolkiewicz D (2018) Dermal microvessel density and maturity is closely associated with atherogenic dyslipidemia and accumulation of advanced glycation end products in adult patients with type 1 diabetes. *Microvasc Res* **121**: 46–51. <https://doi.org/10.1016/j.mvr.2018.10.002>
- Alharris E, Alghetaa H, Seth R, Chatterjee S, Singh NP, Nagarkatti M, Nagarkatti P (2018) Resveratrol attenuates allergic asthma and associated inflammation in the lungs through regulation of miRNA-34a that targets FoxP3 in mice. *Front Immunol* **9**: 2992. <https://doi.org/10.3389/fimmu.2018.02992>
- Arcanjo NMO, Luna C, Madruga MS, Estévez M (2018) Antioxidant and pro-oxidant actions of resveratrol on human serum albumin in the presence of toxic diabetes metabolites: Glyoxal and methylglyoxal. *Biochim Biophys Acta Gen Subj* **1862**: 1938–1947. <https://doi.org/10.1016/j.bbagen.2018.06.007>
- BaGen H, Liu X, Han J (2018) The anti-inflammation effects of resveratrol for patients after oral implantology. *Biomed Res* **29**: 1841–1844. <https://doi.org/10.4066/biomedicalresearch.29-18-398>
- Bass TM, Weinkove D, Houthoofd K, Gems D, Partridge L (2007) Effects of resveratrol on lifespan in *Drosophila melanogaster* and *Caenorhabditis elegans*. *Mech Ageing Dev* **128**: 546–552. <https://doi.org/10.1016/j.mad.2007.07.007>
- Baur JA, Pearson KJ, Price NL, Jamieson HA, Lerin C, Kalra A, Prabhu VV, Allard JS, Lopez-Lluch G, Lewis K, Pistell PJ, Poosala S, Becker KG, Boss O, Gwinn D, Wang M, Ramaswamy S, Fishbein KW, Spencer RG, Lakatta EG, Le Couteur D, Shaw RJ, Navas P, Puigserver P, Ingram DK, de Cabo R, Sinclair DA (2006) Resveratrol improves health and survival of mice on a high-calorie diet. *Nature* **444**: 337–342. <https://doi.org/10.1038/nature05354>
- Bereswill S, Muñoz M, Fischer A, Plickert R, Haag LM, Otto B, Kühl AA, Lodenkemper C, Göbel UB, Heimesaat MM (2010) Anti-inflammatory effects of resveratrol, curcumin and simvastatin in acute small intestinal inflammation. *PLoS One* **5**: e15099. <https://doi.org/10.1371/journal.pone.0015099>
- Berman AY, Motechin RA, Wiesenfeld MY, Holz MK (2017) The therapeutic potential of resveratrol: a review of clinical trials. *NPJ Precis Oncol* **1**. <https://doi.org/10.1038/s41698-017-0038-6>
- Bo S, Gambino R, Ponzio V, Cioffi I, Goitre I, Evangelista A, Ciccone G, Cassader M, Procopio M (2018) Effects of resveratrol on bone health in type 2 diabetic patients. A double-blind randomized-controlled trial. *Nutr Diabetes* **8**: 51. <https://doi.org/10.1038/s41387-018-0059-4>
- Bobermin LD, Souza DO, Gonçalves CA, Quincozes-Santos A (2018) Resveratrol prevents ammonia-induced mitochondrial dysfunction and cellular redox imbalance in C6 astroglial cells. *Nutr Neurosci* **21**: 276–285. <https://doi.org/10.1080/1028415X.2017.1284375>
- Bonsack F, Alleyne CH Jr, Sukumari-Ramesh S (2017) Resveratrol attenuates neurodegeneration and improves neurological outcomes after intracerebral hemorrhage in mice. *Front Cell Neurosci* **11**: 228. <https://doi.org/10.3389/fncel.2017.00228>
- Brown VA, Patel KR, Viskaduraki M, Crowell JA, Perloff M, Booth TD, Vasilinin G, Sen A, Schinas AM, Piccirilli G, Brown K, Steward WP, Gescher AJ, Brenner DE (2010) Repeat dose study of the cancer chemopreventive agent resveratrol in healthy volunteers: safety, pharmacokinetics, and effect on the insulin-like growth factor axis. *Cancer Res* **70**: 9003–9011. <https://doi.org/10.1158/0008-5472.CAN-10-2364>
- Buttari B, Profumo E, Facchiano F, Ozturk EI, Segoni L, Saso L, Riganò R (2013) Resveratrol prevents dendritic cell maturation in response to advanced glycation end products. *Oxid Med Cell Longev* **2013**: 574029. <https://doi.org/10.1155/2013/574029>
- Caruso F, Tanski J, Villegas-Estrada A, Rossi M (2004) Structural basis for antioxidant activity of trans-resveratrol: ab initio calculations and crystal and molecular structure. *J Agric Food Chem* **52**: 7279–7285. <https://doi.org/10.1021/jf048794e>
- Chen CY, Kao CL, Liu CM (2018) The cancer prevention, anti-inflammatory and anti-oxidation of bioactive phytochemicals targeting the TLR4 signaling pathway. *Int J Mol Sci* **19**. <https://doi.org/10.3390/ijms19092729>
- Cheng K, Song Z, Zhang H, Li S, Wang C, Zhang L, Wang T (2019) The therapeutic effects of resveratrol on hepatic steatosis in high-fat diet-induced obese mice by improving oxidative stress, inflamma-

- tion and lipid-related gene transcriptional expression. *Med Mol Morphol* in press. <https://doi.org/10.1007/s00795-019-00216-7>
- Ciddi V, Dodda D (2014) Therapeutic potential of resveratrol in diabetic complications: *In vitro* and *in vivo* studies. *Pharmacol Rep* **66**: 799–803. <https://doi.org/10.1016/j.pharep.2014.04.006>
- Corpas R, Grinán-Ferré C, Rodríguez-Farré E, Pallás M, Sanfeliu C (2018) Resveratrol induces brain resilience against Alzheimer neurodegeneration through proteostasis enhancement. *Mol Neurobiol* in press. <https://doi.org/10.1007/s12035-018-1157-y>
- Corrêa MG, Absy S, Tenenbaum H, Ribeiro FV, Cirano FR, Casati MZ, Pimentel SP (2018) Resveratrol attenuates oxidative stress during experimental periodontitis in rats exposed to cigarette smoke inhalation. *J Periodontol Res* in press. <https://doi.org/10.1111/jre.12622>
- de Vries K, Strydom M, Steenkamp V (2018) Bioavailability of resveratrol: possibilities for enhancement. *J Herb Med* **11**: 71–79. <https://doi.org/10.1016/j.hermed.2017.09.002>
- Diao J, Wei J, Yan R, Fan G, Lin L, Chen M (2018) Effects of resveratrol on regulation on UCP2 and cardiac function in diabetic rats. *J Physiol Biochem* in press. <https://doi.org/10.1007/s13105-018-0648-7>
- Erkan SO, Tuhanioglu B, Grgen SG, zdař T, Tařtekin B, Pelit A, Grgl O (2018) The effect of resveratrol on the histologic characteristics of the cochlea in diabetic rats. *Laryngoscope* in press. <https://doi.org/10.1002/lary.27253>
- Euba B, Lpez-Lpez N, Rodrguez-Arce I, Fernndez-Calvet A, Barbern M, Caturla N, Mart S, Dez-Martnez R, Garmenda J (2017) Resveratrol therapeutics combines both antimicrobial and immunomodulatory properties against respiratory infection by nontypeable *Haemophilus influenzae*. *Sci Rep* **7**: 12860. <https://doi.org/10.1038/s41598-017-13034-7>
- Farrokhi E, Ghatreh-Samani K, Salehi-Vanani N, Mahmoodi A (2018) The effect of resveratrol on expression of matrix metalloproteinase 9 and its tissue inhibitors in vascular smooth muscle cells. *ARYA Atheroscler* **14**: 157–162. <https://doi.org/10.22122/arya.v14i4.1484>
- Farzaei MH, Rahimi R, Nikfar S, Abdollahi M (2018) Effect of resveratrol on cognitive and memory performance and mood: A meta-analysis of 225 patients. *Pharmacol Res* **128**: 338–344. <https://doi.org/10.1016/j.phrs.2017.08.009>
- LaFoya B, Munroe JA, Albig AR (2019) A comparison of resveratrol and other polyphenolic compounds on Notch activation and endothelial cell activity. *PLoS One* **14**: e0210607. <https://doi.org/10.1371/journal.pone.0210607>
- Fu S, Lv R, Wang L, Hou H, Liu H, Shao S (2018) Resveratrol, an antioxidant, protects spinal cord injury in rats by suppressing MAPK pathway. *Saudi J Biol Sci* **25**: 259–266. <https://doi.org/10.1016/j.sjbs.2016.10.019>
- Galiniak S, Krawczyk-Marc I, Sk-Mastej A, Leksa N, Biesiadecki M, Orkisz S (2017) Clinical aspects of protein glycation. *Eur J Clin Exp Med* **15**: 263–267. <https://doi.org/10.15584/ejcem.2017.3.10>
- Gambini J, Ingls M, Olaso G, Lopez-Gruoso R, Bonet-Costa V, Gimeno-Mallench L, Mas-Bargues C, Abdelaziz KM, Gomez-Cabrera MC, Vina J, Borrs C (2015) Properties of resveratrol: *in vitro* and *in vivo* studies about metabolism, bioavailability, and biological effects in animal models and humans. *Oxid Med Cell Longev* **2015**: 837042. <https://doi.org/10.1155/2015/837042>
- Gins C, Cuesta S, Kireev R, Garca C4 Rancan L, Paredes SD, Vara E, Tresguerres JAF (2017) Protective effect of resveratrol against inflammation, oxidative stress and apoptosis in pancreas of aged SAMP8 mice. *Exp Gerontol* **90**: 61–70. <https://doi.org/10.1016/j.exger.2017.01.021>
- Gokce EH, Korkmaz E, Dellera E, Sandri G, Bonferoni MC, Ozer O (2012) Resveratrol-loaded solid lipid nanoparticles versus nanostructured lipid carriers: Evaluation of antioxidant potential for dermal applications. *Int J Nanomed* **7**: 1841–1850. <https://doi.org/10.2147/IJN.S29710>
- Gorga A, Rindone GM, Regueira M, Pellizzari EH, Camberos MC, Cigorraga SB, Riera MF, Galarido MN (2018) Effect of resveratrol on Sertoli cell proliferation. *J Cell Biochem* **119**: 10131–10142. <https://doi.org/10.1002/jcb.27350>
- Glin I (2010) Antioxidant properties of resveratrol: A structure–activity insight. *Innov Food Sci Emerg Technol* **11**: 210–218. <https://doi.org/10.1016/j.ifset.2009.07.002>
- Guzmn L, Balada C, Flores G, Alvarez R, Knox M, Vinet R, Martnez JL (2018) t-Resveratrol protects against acute high glucose damage in endothelial cells. *Plant Foods Hum Nutr* **73**: 235–240. <https://doi.org/10.1007/s11130-018-0683-0>
- Hallajian F, Ghasmi M, Abedi SM, Behzadi R, Hayati E, Sadeghzadeh N, Rezazadeh F, Karimi H (2018) Evaluation of the effect of resveratrol and doxorubicin on 99mTc-MIBI uptake in breast Cancer cell xenografts in mice. *Cancer Biother Radiopharm* **33**: 403–410. <https://doi.org/10.1089/cbr.2018.2523>
- Hara T, Kin A, Aoki S, Nakamura S, Shirasuna K, Kuwayama T, Iwata H (2018) Resveratrol enhances the clearance of mitochondrial damage by vitrification and improves the development of vitrified-warmed bovine embryos. *PLoS One* **13**: e0204571. <https://doi.org/10.1371/journal.pone.0204571>
- Hasan M, Bae H (2017) An overview of stress-induced resveratrol synthesis in grapes: perspectives for resveratrol-enriched grape products. *Molecules* **22**. <https://doi.org/10.3390/molecules22020294>
- He T, Xiong J, Nie L, Yu Y, Guan X, Xu X, Xiao T, Yang K, Liu L, Zhang D, Huang Y, Zhang J, Wang J, Sharma K, Zhao J (2016) Resveratrol inhibits renal interstitial fibrosis in diabetic nephropathy by regulating AMPK/NOX4/ROS pathway. *J Mol Med (Berl)* **94**: 1359–1371. <https://doi.org/10.1007/s00109-016-1451-y>
- Higashi Y, Higashi K, Mori A, Sakamoto K, Ishii K, Nakahara T (2018) Anti-cataract effect of resveratrol in high-glucose-treated streptozotocin-induced diabetic rats. *Biol Pharm Bull* **41**: 1586–1592. <https://doi.org/10.1248/bpb.b18-00328>
- Hou Y, Wang K, Wan W, Cheng Y, Pu X, Ye X (2018) Resveratrol provides neuroprotection by regulating the JAK2/STAT3/PI3K/AKT/mTOR pathway after stroke in rats. *Genes Dis* **5**: 245–255. <https://doi.org/10.1016/j.gendis.2018.06.001>
- Huo X, Zhang T2, Meng Q1, Li C3, You B (2019) Resveratrol effects on a diabetic rat model with coronary heart disease. *Med Sci Monit* **25**: 540–546. <https://doi.org/10.12659/MSM.910996>
- Hussain SA, Marouf BH, Ali ZS, Ahmad RS (2018) Efficacy and safety of co-administration of resveratrol with meloxicam in patients with knee osteoarthritis: a pilot interventional study. *Clin Interv Aging* **13**: 1621–1630. <https://doi.org/10.2147/CIA.S172758>
- Hussein MM, Mahfouz MK (2016) Effect of resveratrol and rosuvastatin on experimental diabetic nephropathy in rats. *Biomed Pharmacother* **82**: 685–692. <https://doi.org/10.1016/j.biopha.2016.06.004>
- Jardim FR, de Rossi FT, Nascimento MX, da Silva Barros RG, Borges PA, Prescilio IC, de Oliveira MR (2018) Resveratrol and brain mitochondria: a review. *Mol Neurobiol* **55**: 2085–2101. <https://doi.org/10.1007/s12035-017-0448-z>
- Johnson AA, Riehle MA (2015) Resveratrol fails to extend life span in the mosquito *Anopheles stephensi*. *Rejuvenation Res* **18**: 473–478. <https://doi.org/10.1089/rej.2015.1670>
- Kapetanovic IM, Muzzio M, Huang Z, Thompson TN, McCormick DL (2011) Pharmacokinetics, oral bioavailability, and metabolic profile of resveratrol and its dimethylether analog, pterostilbene, in rats. *Cancer Chemother Pharmacol* **68**: 593–601. <https://doi.org/10.1007/s00280-010-1525-4>
- Khan H, Khan MS, Ahmad S (2018) The *in vivo* and *in vitro* approaches for establishing a link between advanced glycation end products and lung cancer. *J Cell Biochem* **119**: 9099–9109. <https://doi.org/10.1002/jcb.27170>
- Khazaei M, Karimi J, Sheikh N, Goodarzi MT, Saidijam M, Khodadadi I, Moridi H (2016) Effects of resveratrol on receptor for advanced glycation end products (rage) expression and oxidative stress in the liver of rats with type 2 diabetes. *Phytother Res* **30**: 66–71. <https://doi.org/10.1002/ptr.5501>
- Kumaran MS, Dabas G, Kapadia AB, Parsad D (2018) Resveratrol-induced thrombocytopenia: Inadvertent side-effect of a commonly used antioxidant. *Dermatol Ther* **31**. <https://doi.org/10.1111/dth.12578>
- Landau D, Haghiac M, Calabuig-Navarro V, Minimum J, O'Tierney-Ginn PF (2017) Potential adverse effects of resveratrol: findings in human placental explants. *Am J Obstet Gynecol* **216**: S317. <https://doi.org/10.1016/j.ajog.2016.11.273>
- la Porte C, Voduc N, Zhang G, Seguin I, Tardiff D, Singhal N, Cameron DW (2010) Steady-State pharmacokinetics and tolerability of trans-resveratrol 2000 mg twice daily with food, quercetin and alcohol (ethanol) in healthy human subjects. *Clin Pharmacokinet* **49**: 449–454. <https://doi.org/10.2165/11531820-000000000-00000>
- Lee YE, Kim JW, Lee EM, Ahn YB, Song KH, Yoon KH, Kim HW, Park CW, Li G, Liu Z, Ko SH (2012) Chronic resveratrol treatment protects pancreatic islets against oxidative stress in db/db mice. *PLoS One* **7**: e50412. <https://doi.org/10.1371/journal.pone.0050412>
- Li C, Wu W, Jiao G, Chen Y, Liu H (2018) Resveratrol attenuates inflammation and reduces matrix-metalloprotease expression by inducing autophagy via suppressing the Wnt/b-catenin signalling pathway in IL-1b-induced osteoarthritis chondrocytes. *RSC Adv* **8**: 20202. <https://doi.org/10.1039/c8ra00993g>
- Li N, Du Z, Shen Q, Lei Q, Zhang Y, Zhang M, Hua J (2017) Resveratrol enhances self-renewal of mouse embryonic stem cells. *J Cell Biochem* **118**: 1928–1935. <https://doi.org/10.1002/jcb.25942>
- Li YR, Li S, Lin CC (2018) Effect of resveratrol and pterostilbene on aging and longevity. *Biofactors* **44**: 69–82. <https://doi.org/10.1002/biof.1400>
- Liang QX, Lin YH, Zhang CH, Sun HM, Zhou L, Schatten H, Sun QY, Qian WP (2018) Resveratrol increases resistance of mouse oocytes to postovulatory aging *in vivo*. *Aging (Albany NY)* **10**: 1586–1596. <https://doi.org/10.18632/aging.101494>
- Liguori I, Russo G, Curcio F, Bulli G, Aran L, Della-Morte D, Gargiulo G, Testa G, Cacciatore F, Bonaduce D, Abete P (2018) Oxidative stress, aging, and diseases. *Clin Interv Aging* **13**: 757–772. <https://doi.org/10.2147/CIA.S158513>
- Lin YT, Wu YC, Sun GC, Ho CY, Wong TY, Lin CH, Chen HH, Yeh TC, Li CJ, Tseng CJ, Cheng PW (2018) Effect of resveratrol on reactive oxygen species-induced cognitive impairment in rats with an-

- giotensin II-induced early Alzheimer's disease. *J Clin Med* 7. <https://doi.org/10.3390/jcm7100329>
- Ling L, Gu S, Cheng Y (2017) Resveratrol activates endogenous cardiac stem cells and improves myocardial regeneration following acute myocardial infarction. *Mol Med Rep* 15: 1188–1194. <https://doi.org/10.3892/mmr.2017.6143>
- Liu FC, Hung LF, Wu WL, Chang DM, Huang CY, Lai JH, Ho LJ (2010) Chondroprotective effects and mechanisms of resveratrol in advanced glycation end products-stimulated chondrocytes. *Arthritis Res Ther* 12: R167. <https://doi.org/10.1186/ar3127>
- Liu T, Ma Y, Zhang R, Zhong H, Wang L, Zhao J, Yang L, Fan X (2019) Resveratrol ameliorates estrogen deficiency-induced depression- and anxiety-like behaviors and hippocampal inflammation in mice. *Psychopharmacology (Berl)* in press. <https://doi.org/10.1007/s00213-018-5148-5>
- Liu T, Qi H, Ma L, Liu Z, Fu H, Zhu W, Song T, Yang B, Li G (2015) Resveratrol attenuates oxidative stress and extends life span in the annual fish *Nothobranchius guentheri*. *Rejuvenation Res* 18: 225–233. <https://doi.org/10.1089/rej.2014.1618>
- Liu Y, He XQ, Huang X, Ding L, Xu L, Shen YT, Zhang F, Zhu MB, Xu BH, Qi ZQ, Wang HL (2013) Resveratrol protects mouse oocytes from methylglyoxal-induced oxidative damage. *PLoS One* 8: e77960. <https://doi.org/10.1371/journal.pone.0077960>
- Lopez MS, Dempsey RJ, Vemuganti R (2015) Resveratrol neuroprotection in stroke and traumatic CNS injury. *Neurochem Int* 89: 75–82. <https://doi.org/10.1016/j.neuint.2015.08.009>
- Lucas J, Hsieh TC, Halicka HD, Darzynkiewicz Z, Wu JM (2018) Up-regulation of PDL1 expression by resveratrol and piceatannol in breast and colorectal cancer cells occurs via HDAC3/p300mediated NF κ B signaling. *Int J Oncol* 53: 1469–1480. <https://doi.org/10.3892/ijo.2018.4512>
- Ma C, Wang Y, Dong L, Li M, Cai W (2015) Anti-inflammatory effect of resveratrol through the suppression of NF- κ B and JAK/STAT signalling pathways. *Acta Biochim Biophys Sin (Shanghai)* 47: 207–213. <https://doi.org/10.1093/abbs/gmu135>
- Ma X, Sun Z, Liu Y, Jia Y, Zhang B, Zhang J (2013) Resveratrol improves cognition and reduces oxidative stress in rats with vascular dementia. *Neural Regen Res* 8: 2050–2059. <https://doi.org/10.3969/j.issn.1673-5374.2013.22.004>
- Marier JF, Vachon P, Gritsas A, Zhang J, Moreau JP, Ducharme MP (2002) Metabolism and disposition of resveratrol in rats: extent of absorption, glucuronidation, and enterohepatic recirculation evidenced by a linked-rat model. *J Pharmacol Exp Ther* 302: 369–373. PMID: 12065739
- Marouf BH, Hussain SA, Ali ZS, Ahmmad RS (2018) Resveratrol supplementation reduces pain and inflammation in knee osteoarthritis patients treated with meloxicam: A randomized placebo-controlled study. *J Med Food* in press. <https://doi.org/10.1089/jmf.2017.4176>
- Marx W, Kelly JT, Marshall S, Cutajar J, Annois B, Pipingas A, Tierney A, Iatsopoulos C (2018) Effect of resveratrol supplementation on cognitive performance and mood in adults: a systematic literature review and meta-analysis of randomized controlled trials. *Nutr Rev* 76: 432–443. <https://doi.org/10.1093/nutrit/nuy010>
- Matough FA, Budin SB, Hamid ZA, Alwahaibi N, Mohamed J (2012) The role of oxidative stress and antioxidants in diabetic complications. *Sultan Qaboos Univ Med J* 12: 5–18. PMID: PMC3286717
- Mehdi F, Keihan GS, Asadollah AS, Effat F (2018) The effects of resveratrol, metformin, cold and strength training on the level of perilipin 5 in the heart, skeletal muscle and brown adipose tissues in mouse. *Cell Biochem Biophys* 76: 471–476. <https://doi.org/10.1007/s12013-018-0860-7>
- Monteillal A, Voisin A, Furrer P, Allémann E, Cuendet M (2018) Intranasal administration of resveratrol successfully prevents lung cancer in A/J mice. *Sci Rep* 8: 14257. <https://doi.org/10.1038/s41598-018-32423-0>
- Moore A, Beidler J, Hong MY (2018) Resveratrol and depression in animal models: a systematic review of the biological mechanisms. *Molecules* 23. <https://doi.org/10.3390/molecules23092197>
- Moridi H, Karimi J, Sheikh N, Goodarzi MT, Saidijam M, Yadegarari R, Khazaei M, Khodadadi I, Tavilani H, Piri H, Asadi S, Zarei S, Rezaei A (2015) Resveratrol-dependent down-regulation of receptor for advanced glycation end-products and oxidative stress in kidney of rats with diabetes. *Int J Endocrinol Metab* 13: e23542. <https://doi.org/10.5812/ijem.23542>
- Mukherjee S, Dudley JJ, Das DK (2010) Dose-dependency of resveratrol in providing health benefits. *Dose Response* 8: 478–500. <https://doi.org/10.2203/dose-response.09-015.Mukherjee>
- Nalagoni CSR, Karnati PR (2016) Protective effect of resveratrol against neuronal damage through oxidative stress in cerebral hemisphere of aluminum and fluoride treated rats. *Interdiscip Toxicol* 9: 78–82. <https://doi.org/10.1515/intox-2016-0009>
- Namsi A, Nury T, Hamdouni H, Yammine A, Vejux A, Vervandier-Passeur D, Latruffe N, Masmoudi-Kouki O, Lizard G (2018) Induction of neuronal differentiation of murine N2a cells by two polyphenols present in the Mediterranean diet mimicking neurotrophins activities: resveratrol and apigenin. *Diseases* 6. <https://doi.org/10.3390/diseases6030067>
- Novelle MG, Wahl D, Diéguez C, Bernier M, de Cabo R (2015) Resveratrol supplementation: Where are we now and where should we go? *Ageing Res Rev* 21: 1–15. <https://doi.org/10.1016/j.arr.2015.01.002>
- Ohtsu A, Shibutani Y, Seno K, Iwata H, Kuwayama T, Shirasuna K (2017) Advanced glycation end products and lipopolysaccharides stimulate interleukin-6 secretion via the RAGE/TLR4-NF- κ B-ROS pathways and resveratrol attenuates these inflammatory responses in mouse macrophages. *Exp Ther Med* 14: 4363–4370. <https://doi.org/10.3892/etm.2017.5045>
- Öztürk E, Arslan AKK, Yerer MB, Bishayee A (2017) Resveratrol and diabetes: A critical review of clinical studies. *Biomed Pharmacother* 95: 230–234. <https://doi.org/10.1016/j.biopha.2017.08.070>
- Pallauf K, Rimbach G, Rupp PM, Chin D, Wolf IM (2016) Resveratrol and lifespan in model organisms. *Curr Med Chem* 23: 4639–4680. <https://doi.org/10.2174/0929867323666161024151233>
- Peñalva R, Morales J, González-Navarro CJ, Larrañeta E, Quincoces G, Peñuelas I, Irache JM (2018) Increased oral bioavailability of resveratrol by its encapsulation in casein nanoparticles. *Int J Mol Sci* 19. <https://doi.org/10.3390/ijms19092816>
- Peng L, Jiang D (2018) Resveratrol eliminates cancer stem cells of osteosarcoma by STAT3 pathway inhibition. *PLoS One* 13: e0205918. <https://doi.org/10.1371/journal.pone.0205918>
- Pinheiro DML, de Oliveira AHS, Coutinho LG, Fontes FL, de Medeiros Oliveira RK, Oliveira TT, Faustino ALF, Lira V, de Melo Campos JTA, Lajus TBP, de Souza SJ, Agnez-Lima LF (2018) Resveratrol decreases the expression of genes involved in inflammation through transcriptional regulation. *Free Radic Biol Med* 130: 8–22. <https://doi.org/10.1016/j.freeradbiomed.2018.10.432>
- Pinkas A, Aschner M (2016) Advanced glycation end-products and their receptors: related pathologies, recent therapeutic strategies, and a potential model for future neurodegeneration studies. *Chem Res Toxicol* 29: 707–714. <https://doi.org/10.1021/acs.chemrestox.6b00034>
- Plauth A, Geikowski A, Cichon S, Wowro SJ, Liedgens L, Rousseau M, Weidner C, Fuhr L, Kliem M, Jenkins G, Lotito S, Wainwright IJ, Sauer S (2016) Hormetic shifting of redox environment by pro-oxidative resveratrol protects cells against stress. *Free Radic Biol Med* 99: 608–622. <https://doi.org/10.1016/j.freeradbiomed.2016.08.006>
- Popat R, Plesner T, Davies F, Cook G, Cook M, Elliott P, Jacobson E, Gumbleton T, Oakervee H, Cavenagh J (2013) A phase 2 study of SRT501 (resveratrol) with bortezomib for patients with relapsed and/or refractory multiple myeloma. *Br J Haematol* 160: 714–717. <https://doi.org/10.1111/bjh.12154>
- Pouyafar A, Zadi Heydarabad M, Aghdam SB, Khaksar M, Azimi A, Rahbarghazi R, Talebi M (2019) Resveratrol potentially increased the tumoricidal effect of doxorubicin on SKOV3 cancer stem cells *in vitro*. *J Cell Biochem* in press. <https://doi.org/10.1002/jcb.28129>
- Ramos-Gomez M, Olivares-Marin IK, Canizal-García M, González-Hernández JC, Nava GM, Madrigal-Perez LA (2017) Resveratrol induces mitochondrial dysfunction and decreases chronological life span of *Saccharomyces cerevisiae* in a glucose-dependent manner. *J Bioenerg Biomembr* 49: 241–251. <https://doi.org/10.1007/s10863-017-9709-9>
- Rascon B, Hubbard BP, Sinclair DA, Amdam GV (2012) The lifespan extension effects of resveratrol are conserved in the honey bee and may be driven by a mechanism related to caloric restriction. *Ageing (Albany NY)* 4: 499–508. <https://doi.org/10.18632/aging.100474>
- Rauf A, Imran M, Suleria HAR, Ahmad B, Peters DG, Mubarak MS (2017) A comprehensive review of the health perspectives of resveratrol. *Food Funct* 8: 4284–4305. <https://doi.org/10.1039/c7fo01300k>
- Rhee SY, Kim YS (2018) The role of advanced glycation end products in diabetic vascular complications. *Diabetes Metab J* 42: 188–195. <https://doi.org/10.4093/dmj.2017.0105>
- Rivière C, Pawlus AD, Mérillon JM (2012) Natural stilbenoids: distribution in the plant kingdom and chemotaxonomic interest in Vitaceae. *Nat Prod Rep* 29: 1317–1333. <https://doi.org/10.1039/c2np20049j>
- Roggerio A, Strunz CMC, Pacanaro AP, Leal DP, Takada JY, Avakian SD, Mansur AP (2018) Gene expression of sirtuin-1 and endogenous secretory receptor for advanced glycation end products in healthy and slightly overweight subjects after caloric restriction and resveratrol administration. *Nutrients* 10. <https://doi.org/10.3390/nu10070937>
- Sadowska-Bartosz I, Galiniak S, Bartosz G (2014) Kinetics of glycoxidation of bovine serum albumin by glucose, fructose and ribose and its prevention by food components. *Molecules* 19: 18828–18849. <https://doi.org/10.3390/molecules191118828>
- Safaiejad Z, Nabiuni M, Peymani M, Ghaedi K, Nasr-Esfahani MH, Baharvand H (2017) Resveratrol promotes human embryonic stem cells self-renewal by targeting SIRT1-ERK signaling pathway. *Eur J Cell Biol* 96: 665–672. <https://doi.org/10.1016/j.ejcb.2017.08.002>
- Salehi B, Mishra AP, Nigam M, Sener B, Kilic M, Sharifi-Rad M, Fokou PVT, Martins N, Sharifi-Rad J (2018) Resveratrol: A dou-

- ble-edged sword in health benefits. *Biomedicines* **6**. <https://doi.org/10.3390/biomedicines6030091>
- San Hipolito-Luengo A, Alcaide A, Ramos-Gonzalez M, Cercas E, Vallejo S, Romero A, Talero E, Sanchez-Ferrer CF, Motiva V, Peiro C (2017) Dual effects of resveratrol on cell death and proliferation of colon cancer cells. *Nutr Cancer* **69**: 1019–1027. <https://doi.org/10.1080/01635581.2017.1359309>
- Sattarinezhad A, Roozbeh J, Shirazi Yeganeh B, Omrani GR, Shams M (2018) Resveratrol reduces albuminuria in diabetic nephropathy: A randomized double-blind placebo-controlled clinical trial. *Diabetes Metab* in press. <https://doi.org/10.1016/j.diabet.2018.05.010>
- Sedlak L, Wojnar W, Zych M, Wyględowska-Promieńska D, Mrukwa-Kominek E, Kaczmarczyk-Sedlak I (2018) Effect of resveratrol, a dietary-derived polyphenol, on the oxidative stress and polyol pathway in the lens of rats with streptozotocin-induced diabetes. *Nutrients* **10**: 1423. <https://doi.org/10.3390/nu10101423>
- Sergides C, Chirilă M, Silvestro L, Pitta D, Pittas A (2016) Bioavailability and safety study of resveratrol 500 mg tablets in healthy male and female volunteers. *Exp Ther Med* **11**: 164–170.
- Sgarbi G, Liuzzi F, Baracca A, Solaini G Resveratrol preserves mitochondrial function in a human post-mitotic cell model. *J Nutr Biochem* **62**: 9–17. <https://doi.org/10.1016/j.jnutbio.2018.07.017>
- Sharma C, Suhalka P, Bhatnagar M (2018) Curcumin and resveratrol rescue cortical-hippocampal system from chronic fluoride-induced neurodegeneration and enhance memory retrieval. *Int J Neurosci* **13**: 1–15. <https://doi.org/10.1080/00207454.2018.1458727>
- Shati AA (2018) Resveratrol improves sperm parameter and testicular apoptosis in cisplatin-treated rats: effects on ERK1/2, JNK, and Akt pathways. *Syst Biol Reprod Med* **3**: 1–14. <https://doi.org/10.1080/19396368.2018.1541114>
- Shen Y, Xu Z, Sheng Z (2017) Ability of resveratrol to inhibit advanced glycation end product formation and carbohydrate-hydrolyzing enzyme activity, and to conjugate methylglyoxal. *Food Chem* **216**: 153–160. <https://doi.org/10.1016/j.foodchem.2016.08.034>
- Shrikanta A, Kumar A, Govindaswamy V (2015) Resveratrol content and antioxidant properties of underutilized fruits. *J Food Sci Technol* **52**: 383–390. <https://doi.org/10.1007/s13197-013-0993-z>
- Singh R, Barden A, Mori T, Beilin L (2001) Advanced glycation end-products: a review. *Diabetologia* **44**: 129–146. <https://doi.org/10.1007/s001250051591>
- Soleas GJ, Angelini M, Grass L, Diamandis EP, Goldberg DM (2001) Absorption of trans-resveratrol in rats. *Methods Enzymol* **335**: 145–154. PMID: 11400363
- Staats S, Wagner AE, Kowalewski B, Rieck FT, Soukup ST, Kulling SE, Rimbach G (2018) Dietary resveratrol does not affect life span, body composition, stress response, and longevity-related gene expression in *Drosophila melanogaster*. *Int J Mol Sci* **19**: 223. <https://doi.org/10.3390/ijms19010223>
- Subramanian M, Goswami M, Chakraborty S, Jawali N (2014) Resveratrol induced inhibition of *Escherichia coli* proceeds via membrane oxidation and independent of diffusible reactive oxygen species generation. *Redox Biol* **2**: 865–872. <https://doi.org/10.1016/j.redox.2014.06.007>
- Szende B, Tyihak E, Kiraly-Veghely Z (2010) Dose-dependent effect of resveratrol on proliferation and apoptosis in endothelial and tumor cell cultures. *Exp Mol Med* **32**: 88. <https://doi.org/10.1038/emmm.2000.16>
- Švaiger U, Jeras M (2012) Anti-inflammatory effects of resveratrol and its potential use in therapy of immune-mediated diseases. *Int Rev Immunol* **31**: 202–222. <https://doi.org/10.3109/08830185.2012.665108>
- Takaoka M (1939) Resveratrol, a new phenolic compound from *Veratrum grandiflorum*. *Nippon Kagaku Kaishi* **60**: 1090–1100.
- Tao L, Ding Q, Gao C, Sun X (2016) Resveratrol attenuates neuropathic pain through balancing pro-inflammatory and anti-inflammatory cytokines release in mice. *Int Immunopharmacol* **34**: 165–172. <https://doi.org/10.1016/j.intimp.2016.02.033>
- Tasatargil A, Tanriover G, Barutçigil A, Turkmen E (2018) Protective effect of resveratrol on methylglyoxal-induced endothelial dysfunction in aged rats. *Aging Clin Exp Res* in press. <https://doi.org/10.1007/s40520-018-0986-x>
- Taylor EJ, Yu Y, Champer J, Kim J (2014) Resveratrol demonstrates antimicrobial effects against *Propionibacterium acnes* in vitro. *Dermatol Ther (Heidelb)* **4**: 249–257. <https://doi.org/10.1007/s13555-014-0063-0>
- Toniolo L, Fusco P, Formoso L, Mazzi A, Canato M, Reggiani C, Giacomello E (2018) Resveratrol treatment reduces the appearance of tubular aggregates and improves the resistance to fatigue in aging mice skeletal muscles. *Exp Gerontol* **111**: 170–179. <https://doi.org/10.1016/j.exger.2018.07.012>
- Valenzano DR, Terzibas E, Genade T, Cattaneo A, Domenici L, Cellerino A (2006) Resveratrol prolongs lifespan and retards the onset of age-related markers in a short-lived vertebrate. *Curr Biol* **16**: 296–300. <https://doi.org/10.1016/j.cub.2005.12.038>
- Vasamsetti SB, Karnewar S, Gopoju R, Gollavilli PN, Narra SR, Kumar JM, Kotamraju S (2016) Resveratrol attenuates monocyte-to-macrophage differentiation and associated inflammation via modulation of intracellular GSH homeostasis: Relevance in atherosclerosis. *Free Radic Biol Med* **96**: 392–405. <https://doi.org/10.1016/j.freeradbiomed.2016.05.003>
- Walle T (2011) Bioavailability of resveratrol. *Ann N Y Acad Sci* **1215**: 9–15. <https://doi.org/10.1111/j.1749-6632.2010.05842.x>
- Walter KR, Ford ME, Gregoski MJ, Kramer RM, Knight KD, Spruill L, Nogueira LM, Krisanits BA, Phan V, La Rue AC, Lilly MB, Ambs S, Chan K, Turner TF, Varner H, Singh S, Uribarri J, Garrett-Mayer E, Armeson KE, Hilton EJ, Clair MJ, Taylor MH, Abbott AM, Findlay VJ, Peterson LL, Magwood G, Turner DP (2018) Advanced glycation end products are elevated in estrogen receptor-positive breast cancer patients, alter response to therapy, and can be targeted by lifestyle intervention. *Breast Cancer Res Treat* in press. <https://doi.org/10.1007/s10549-018-4992-7>
- Wang C, Wheeler CT, Alberico T, Sun X, Seeberger J, Laslo M, Spangler E, Kern B, de Cabo R, Zou S (2013) The effect of resveratrol on lifespan depends on both gender and dietary nutrient composition in *Drosophila melanogaster*. *Age (Dordr)* **35**: 69–81. <https://doi.org/10.1007/s11357-011-9332-3>
- Wang G, Song X, Zhao L, Li Z, Liu B (2018) Resveratrol prevents diabetic cardiomyopathy by increasing NRF2 expression and transcriptional activity. *Biomed Res Int* **2018**: 2150218. <https://doi.org/10.1155/2018/2150218>
- Wang N, Han Q, Wang G, Ma WP, Wang J, Wu WX, Guo Y, Liu L, Jiang XY, Xie XL, Jiang HQ (2016) Resveratrol protects oxidative stress-induced intestinal epithelial barrier dysfunction by upregulating heme oxygenase-1 expression. *Dig Dis Sci* **61**: 2522–2534. <https://doi.org/10.1007/s10620-016-4184-4>
- Wang XL, Li T, Li JH, Miao SY, Xiao XZ (2017) The effects of resveratrol on inflammation and oxidative stress in a rat model of chronic obstructive pulmonary disease. *Molecules* **22**. <https://doi.org/10.3390/molecules22091529>
- Wątroba M, Szukiewicz D (2016) The role of sirtuins in aging and age-related diseases. *Adv Med Sci* **61**: 52–62. <https://doi.org/10.1016/j.advms.2015.09.003>
- Wu MY, Yiang GT, Lai TT, Li CJ (2018) The oxidative stress and mitochondrial dysfunction during the pathogenesis of diabetic retinopathy. *Oxid Med Cell Longev* **2018**: 3420187. <https://doi.org/10.1155/2018/3420187>
- Wu S, Huang J (2017) Resveratrol alleviates *Staphylococcus aureus* pneumonia by inhibition of the NLRP3 inflammasome. *Exp Ther Med* **14**: 6099–6104. <https://doi.org/10.3892/etm.2017.5337>
- Wu X, Xu Y, Zhu B, Liu Q, Yao Q, Zhao G (2018) Resveratrol induces apoptosis in SGC-7901 gastric cancer cells. *Oncol Lett* **16**: 2949–2956. <https://doi.org/10.3892/ol.2018.9045>
- Xia Y, Xiao HT, Liu K, Zhang HJ, Tsang SW (2018) Resveratrol ameliorates the severity of fibrogenesis in mice with experimental chronic pancreatitis. *Mol Nutr Food Res* **62**: e1700561. <https://doi.org/10.1002/mnfr.201700561>
- Xie J, Liu S, Wu B2 Li G, Rao S, Zou L, Yi Z, Zhang C, Jia T, Zhao S, Schmalzing G, Hausmann R, Nie H, Li G, Liang S (2017) The protective effect of resveratrol in the transmission of neuropathic pain mediated by the P2X7 receptor in the dorsal root ganglia. *Neurochem Int* **103**: 24–35. <https://doi.org/10.1016/j.neuint.2016.12.006>
- Xu K, Liu XF, Ke ZQ, Yao Q, Guo S, Liu C (2018) Resveratrol modulates apoptosis and autophagy induced by high glucose and palmitate in cardiac cells. *Cell Physiol Biochem* **46**: 2031–2040. <https://doi.org/10.1159/000489442>
- Yeh WJ, Hsia SM, Lee WH, Wu CH (2017) Polyphenols with anti-glycation activity and mechanisms of action: A review of recent findings. *J Food Drug Anal* **25**: 84–92. <https://doi.org/10.1016/j.jfda.2016.10.017>
- Yilmaz Z, Kalaz EB, Aydın AF, Olgaç V, Doğru-Abbasoğlu S, Uysal M, Koçak-Toker N (2018) The effect of resveratrol on glycation and oxidation products in plasma and liver of chronic methylglyoxal-treated rats. *Pharmacol Rep* **70**: 584–590. <https://doi.org/10.1016/j.pharep.2017.12.005>
- Yuan D, Liu XM, Fang Z, Du LL, Chang J, Lin SH (2018) Protective effect of resveratrol on kidney in rats with diabetic nephropathy and its effect on endoplasmic reticulum stress. *Eur Rev Med Pharmacol Sci* **22**: 1485–1493. https://doi.org/10.26355/eurrev_201803_14497
- Zhang T, Chi Y, Kang Y, Lu H, Niu H, Liu W, Li Y (2018) Resveratrol ameliorates podocyte damage in diabetic mice via SIRT1/PGC-1 α mediated attenuation of mitochondrial oxidative stress. *J Cell Physiol* in press. <https://doi.org/10.1002/jcp.27306>
- Zhang Y, Luo Z, Ma L, Xu Q, Yang Q, Si L (2010) Resveratrol prevents the impairment of advanced glycosylation end products (AGE) on macrophage lipid homeostasis by suppressing the receptor for AGE via peroxisome proliferator-activated receptor gamma activation. *Int J Mol Med* **25**: 729–734. PMID: 20372816
- Zhao W, Huang X, Han X, Hu D, Hu X, Li Y, Huang P, Yao W (2018). Resveratrol suppresses gut-derived NLRP3 inflammasome partly through stabilizing mast cells in a rat model. *Mediators Inflamm* **2018**: 6158671. <https://doi.org/10.1155/2018/6158671>
- Zhao Y, Yuan X, Li X, Zhang Y (2018) Resveratrol significantly inhibits the occurrence and development of cervical cancer by regu-

- lating phospholipid scramblase 1. *J Cell Biochem* in press. <https://doi.org/10.1002/jcb.27335>
- Zheng X, Jia B, Tian XT, Song X, Wu ML, Kong QY, Li H, Liu J (2018) Correlation of reactive oxygen species levels with resveratrol sensitivities of anaplastic thyroid cancer cells. *Oxid Med Cell Longev* **2018**: 6235417. <https://doi.org/10.1155/2018/6235417>
- Zhou C, Qian W, Ma J, Cheng L, Jiang Z, Yan B, Li J, Duan W, Sun L, Cao J, Wang F, Wu E, Wu Z, Ma Q, Li X (2018) Resveratrol enhances the chemotherapeutic response and reverses the stemness induced by gemcitabine in pancreatic cancer cells *via* targeting SREBP1. *Cell Prolif* **19**: e12514. <https://doi.org/10.1111/cpr.12514>
- Zhou R, Yi L, Ye X, Zeng X, Liu K, Qin Y, Zhang Q, Mi M (2018) Resveratrol ameliorates lipid droplet accumulation in liver through a SIRT1/ATF6-dependent mechanism. *Cell Physiol Biochem* **51**: 2397–2420. <https://doi.org/10.1159/000495898>