

Review Article

Health-Promoting Properties of *Eucommia ulmoides*: A Review

Tarique Hussain,^{1,2} Bi'e Tan,^{1,3} Gang Liu,² Oso Abimbola Oladele,⁴ Najma Rahu,⁵
M. C. Tossou,^{1,2} and Yulong Yin¹

¹Observation and Experiment Station of Animal Nutrition and Feed Science in South-Central China, Ministry of Agriculture, Hunan Provincial Engineering Research Center for Healthy Livestock and Poultry Production, Key Laboratory of Agro-Ecological Processes in Subtropical Region, Institute of Subtropical Agriculture, Chinese Academy of Sciences, Changsha, Hunan 410125, China

²University of the Chinese Academy of Sciences, Beijing 10008, China

³Hunan Collaborative Innovation Center for Utilization of Botanical Functional Ingredients, Changsha, Hunan 410000, China

⁴Department of Animal Nutrition, College of Animal Science and Livestock Production, Federal University of Agriculture, Abeokuta 110101, Nigeria

⁵Department of Veterinary Microbiology, Faculty of Animal Husbandry and Veterinary Sciences, Sindh Agriculture University, Tando Jam, Sindh 70050, Pakistan

Correspondence should be addressed to Bi'e Tan; bietan@isa.ac.cn

Received 4 November 2015; Accepted 31 January 2016

Academic Editor: Il-Moo Chang

Copyright © 2016 Tarique Hussain et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Eucommia ulmoides (EU) (also known as “Du Zhong” in Chinese language) is a plant containing various kinds of chemical constituents such as lignans, iridoids, phenolics, steroids, flavonoids, and other compounds. These constituents of EU possess various medicinal properties and have been used in Chinese Traditional Medicine (TCM) as a folk drink and functional food for several thousand years. EU has several pharmacological properties such as antioxidant, anti-inflammatory, antiallergic, antimicrobial, anticancer, antiaging, cardioprotective, and neuroprotective properties. Hence, it has been widely used solely or in combination with other compounds to treat cardiovascular and cerebrovascular diseases, sexual dysfunction, cancer, metabolic syndrome, and neurological diseases. This review paper summarizes the various active ingredients contained in EU and their health-promoting properties, thus serving as a reference material for the application of EU.

1. Introduction

Eucommia ulmoides (EU) (commonly called “Du Zhong” in Chinese language) belong to the family of Eucommiaceae, a genus of the small tree native to Central China [1]. This plant is widely cultivated in China on a large scale because of its medicinal importance. About 112 compounds have been isolated from EU which include lignans, iridoids, phenolics, steroids, and other compounds. Complementary herbs formula of this plant (such as delicious tea) has shown some medicinal properties. The leaf of EU has higher activity related to cortex, flower, and fruit [2, 3]. The leaves of EU have been reported to enhance bones strength and body muscles [4], thus leading to longevity and promoting fertility in humans [5]. Delicious tea formula made from

the leaf of EU was reported to reduce fattiness and enhance energy metabolism. Flavonoid compounds (such as rutin, chlorogenic acid, ferulic acid, and caffeic acid) have been reported to exhibit antioxidants activity in the leaves of EU [6].

Although there has been enough literature on phytochemical properties of EU, few studies however existed on the pharmacological properties of the various compounds extracted from the barks, seeds, stems, and leaves of EU. This review paper will elucidate detailed information regarding different compounds extracted from the various parts (barks, seeds, stem, and leaf) of EU and the prospective uses of these compounds in health-promoting properties with scientific lines of evidence and thus provide a reference material for the application of EU.

2. Chemical Composition of *Eucommia ulmoides*

Various compounds isolated from different parts of EU are shown in Table 1.

2.1. Lignans and Iridoids. Lignans and their derivatives are the key components of EU [7]. To date, 28 lignans (such as bisepoxy lignans, monoepoxy lignans, neolignans, and sesquiolignans) have been isolated from bark, leaves, and seeds of EU. Iridoid glycoside, a class of secondary metabolites, is the second main component of EU. Iridoids are typically found in plants known as glycosides. Twenty-four iridoids have been isolated and identified from EU (Table 1). These isolated compounds include geniposidic acid, aucubin, and asperuloside which have been reported to have wide pharmacological properties [8–10]. Two new compounds of iridoids, Eucommides-A and -C, have recently been isolated. These two natural compounds are considered as conjugates of iridoid and amino acids. However, the mechanism underlying their activity is not available [11].

2.2. Phenolic Compounds. Phenolic compounds which are derived from the foods have been reported to have positive impact on human health [12, 13]. About 29 phenolic compounds have been isolated and identified from EU [14]. Total content of phenolic compounds (in gallic acid equivalents of all the extracts) was analyzed using the Folin-Ciocalteu phenol reagent. Effects of seasonal variation on the contents of some compounds and antioxidants have been reported. Within the same year, higher contents of phenolics and flavonoids were discovered in the leaves of EU in August and May, respectively. Rutin, quercetin, geniposidic acid, and aucubin existed in higher concentration in May or June [15]. Moreover, higher activity of 1,1-diphenyl-2-picrylhydrazyl (DPPH) radical scavenging activity and metal ion chelating ability were found in the leaves of EU harvested in August. Increased content of food antioxidants was also reported in May when compared to other periods of the year [15]. The leaf of EU has been found to be a rich source of amino acids, vitamins, minerals, and flavonoids such as quercetin, rutin, and geniposidic acid [11, 16]. A total of 7 flavonoids have been isolated from *Eucommia* plants [17]. Rutin and quercetin are the most important flavonoids [18]. Flavonoids are important compounds which are common in nature and are considered as secondary metabolites and function as chemical messengers, physiological regulators, and cell cycle inhibitors.

2.3. Steroids and Terpenoids. Six steroids and five terpenoids have been extracted and categorized from EU. These include β -sitosterol, daucosterol, ulmoprenol, betalin, betulinic acid, ursolic acid, eucommidiol, rehmaglutin C, and 1,4 α ,5,7 α -tetrahydro-7-hydroxymethyl-cyclopenta[c]pyran-4-carboxylic methyl ester which was specifically isolated from the bark of EU [19]. Loliolide has also been isolated from the leaves [20].

2.4. Polysaccharides. Polysaccharides from EU for 15 days at the concentrations of 300–600 mg/kg were reported to

exhibit protective effects on kidneys as observed by malonaldehyde and glutathione levels after renal perfusions [21]. Histological examination also showed evidence of antioxidative properties. Extracts from the bark of EU using 70% ethanol also showed protective effects against cadmium at 125–500 mg/kg [22]. Histological examination also showed that EU in combination with *Panax pseudoginseng* at 25% and 50% weight, respectively, for six weeks at a dose rate of 35.7–41.6 mg/kg exerted light protective effects on glomerular filtration rate [8]. Two new polysaccharides have been separated from EU, which are eucommian A and B [23].

2.5. Other Ingredients and Chemicals. Amino acids, microelements, vitamins, and fatty acids have also been isolated from EU [17, 21–23]. Sun et al. also discovered new compounds such as n-octacosanoic acid, and tetracosanoic-2,3-dihydroxypropylester from EU [24].

Fatty acid composition of oil extracted from the seed of EU showed different concentrations of polyunsaturated fatty acids such as linoleic acid, linolenic acid (56.51% of total fatty acids, TFAs), and linoleic acid (12.66% of TFAs). Meanwhile, the main monounsaturated fatty acid isolated from the seed was found to be isoleic acid (15.80% of TFAs). Dominant saturated fatty acids isolated include palmitic acid and stearic acid which represent 9.82% and 2.59% of TFAs, respectively [25].

3. Health-Promoting Compounds of *Eucommia ulmoides*

3.1. Protective Effects on Cardiovascular System. In Chinese traditional medicines, *Eucommia* is considered as a major herbal tonic for cardiac patients. *Eucommia* bark extract is an active component used for antihypertensive formulations. It has been confirmed in many human as well as animal models as a vasorelaxant. Lignan from EU when administered to rats of the Okamoto strain (SHR) at the dose rate of 300 mg/kg for 16 weeks resulted in improved vascular remodeling and reduced mean arterial blood pressure. EU minimizes blood pressure at the dose of 500–1000 mg/kg. However, in high fructose fed diet, it develops insulin resistance and hypertension [26–28]. Supplementation of 500 and 1000 mg of EU for 8 weeks and thrice daily for 2 weeks showed minimal reduction in blood pressure and reduction in systolic and diastolic blood pressure [29]. Antihypertensive effect on the parasympathetic nervous system has been reported following the application of EU [30]. EU also serves as a vasorelaxative agent depending on nitric oxide and assumed to be linked with potassium channels [31]. EU has beta blocking potential which at 0.5% w/v reduces isoproterenol-stimulated lipolysis from 2.67 to 1.4 times the buffer control [29]. EU has been demonstrated to prevent hypertensive remodeling which is associated with aldose reductase inhibition [32]. The application of lignans from EU under condition of hypertension due to vascular remodeling was reported to serve as a new therapeutic agent [33].

EU also showed antihyperlipidemic properties by suppressing hepatic fatty acid and cholesterol biosynthesis [34]. In hyperlipidemic hamsters, dietary supplementation with

TABLE 1: Compounds isolated from various parts of *Eucommia ulmoides*.

Category	Compounds	References
	<i>Bark of Eucommia ulmoides</i>	
Lignans	(+)-1-Hydroxypinoresinol-4',4''-di-O- β -D-glucopyranoside	[35]
	(+)-1-Hydroxypinoresinol-4'-O- β -D-glucopyranoside	[36]
	(+)-1-Hydroxypinoresinol-4''-O- β -D-glucopyranoside	[36]
	(+)-Epipinoresinol	[37]
	(+)-1-Hydroxypinoresinol	[37]
	(+)-Medioresinol	[37]
	(+)-Medioresinol-di-O- β -D-glucopyranoside	[38]
	(+)-Pinoresinol	[37]
	(+)-Pinoresinol-4'-O- β -D-glucopyranoside	[38]
	(+)-Pinoresinol-di-O- β -D-glucopyranoside	[39]
	(+)-Pinoresinol-4-O- β -D-glucopyranosyl(1-6)- β -D-glucopyranoside	[40]
	(+)-Syringaresinol	[37]
	(+)-Syringaresinol-O- β -D-glucopyranoside	[35]
	Eucommin A (+)-medioresinol-4'- β -D-glucopyranoside	[35]
	Liriodendrin (+)-syringaresinol-di-O- β -D-glucopyranoside	[38]
Phenolics	Astragalinal	[11, 41]
	Isoquercetin	[11, 42]
	Quercetin	[41]
	Quercetin-3-O-galactoside (hyperin)	[41]
	Quercetin-3-O-xyloglucoside	[42]
	Rutin	[11-24]
	Wogonin	[43]
	(-)-Epicatechin	[24]
	(\pm)-Threo-guaiacyl glycerol	[36]
	Caffeic acid	[44]
	Catechin	[24]
	Chlorogenic acid	[11, 24]
	Coniferol	[37]
	Erythro-guaiacylglycerol- β -coniferyl aldehyde ether	[37]
	Eucophenoside	[45]
	Methyl chlorogenate	[37]
	Protocatechuic acid	[46]
Threo-guaiacylglycerol- β -coniferyl aldehyde ether	[37]	
Vanillic acid	[47]	
Iridoids	Deoxyeucommiol	[48]
	Eucommiol-II	[48]
	Eucommiside-I	[48]
	Genipin	[37]
	Geniposide	[48]
	Geniposidic acid	[11, 48]
Monoepoxyignans	(-)-Olivil-4',4''-di-O- β -D-glucopyranoside	[35]
	(+)-Cycloolivil	[36]
	(+)-Olivil	[36]
	(+)-Olivil-4'-O- β -D-glucopyranoside	[36, 49]
	(+)-Olivil-4''-O- β -D-glucopyranoside	[36]
Neolignans	Citrusin B	[50]
	Dehydrodiconiferylalcohol-4, γ '-di-O- β -D-glucopyranoside	[50]
	Dihydroxydehydrodiconiferyl alcohol	[37]
	Erythro-dihydroxydehydrodiconiferyl alcohol	[37]
	Threo-dihydroxydehydrodiconiferyl alcohol	[37]

TABLE 1: Continued.

Category	Compounds	References
Sesquilignans	(-)-HedyotolC-4',4'''-di-O- β -D-glucopyranoside	[48]
	Syringylglycerol- β -syringaresinol ether-4''-4'''-O- β -D-glucopyranoside	[50]
	Syringylglycerol- β -syringaresinol ether-4''-4'''-O- β -D-glucopyranoside	[50]
Steroid and terpenoid	1,4 α ,5,7 α -Tetrahydro-7-hydroxymethyl-cyclopenta[c]pyran-4-carboxylicmethyl ester	[19]
	Betalin	[47]
	Betulic acid	[47]
	Daucosterol	[51]
	Eucommidiol	[19]
	Rehmaglutin C	[19]
Others	Ursolic acid	[47]
	(α R)- α ,4,2',4'-Tetrahydroxydihydrochalcone	[43]
	(α R)- α -O- β -D-Glucopyranosyl-4,2',4'-trihydroxydihyd	[43]
	4,2',4'-Trihydroxychalcone	[43]
	Eucomman A	[52]
	Eucomman B	[23]
	n-Oetaeosanoic acid	[24]
Quercetin-3-O- α -L-arabinopyranosyl-(1-2)- β -D-glucopyranoside	[53, 54]	
Tetraeosanoie-2,3-dihydroxypropyl ester	[24]	
<i>Leaves of Eucommia ulmoides</i>		
Phenolics	Astragalin	[11, 41]
	Hirsutin	[17]
	Isoquercetin	[11, 42]
	Kaempferol	[17]
	Kaempferol-3-O-6''-acetyl-glucoside	[55]
	Kaempferol-3-O-rutinoside	[11]
	Quercetin-3-O- α -L-arabinopyranosyl-(1-2)- β -D-glucopyranoside	[53, 54, 56]
	Rutin	[11, 24]
	Ajugoside	[57]
	Asperuloside	[11]
	Asperulosidic acid	[58]
	Aueubin or aueuboside	[57]
	Deacetyl asperulosidic acid	[58]
	Eucommiol	[57]
	Eucommioside	[59]
	Eucomoside A	[11]
	Eucomoside B	[11]
	Eucomoside C	[11]
	Geniposidic acid	[11, 36]
	Harpagide acetate	[57]
	Reptoside	[57]
	Scandoside-10-O-acetate	[11]
	Ulmoside	[59, 60]
	3-(3,4-Dihydroxyphenyl) propionic acid	[60]
	3(3-Hydroxyphenyl-propionic acid)	[60]
	3,4-Dihydrobenzonic acid	[17]
	Caffeic acid	[37, 61]
	Chlorogenic acid methylester	[62]
	Eatechol	[60]
	Isochlorogenic acid A	[42]
	Isochlorogenic acid C	[42]
	p-trans-Coumaric acid	[58]
	Pyrogallol	[58]

TABLE 1: Continued.

Category	Compounds	References
Monoepoxy lignans	(+)-Olivil-4'-O- β -D-glucopyranoside	[36, 49]
Steroid and terpenoid	Loliolide	[18]
	Ulmoidol	[63]
Others	Ethyl glucopyranoside	[21]
	Seeds of <i>Eucommia ulmoides</i>	
Iridoids	Ulmoidoside A	[64]
	Ulmoidoside B	[64]
	Ulmoidoside C	[64]
	Ulmoidoside D	[64]
Phenolics	Dihydrocaffeic acid	[64]
	Stems of <i>Eucommia ulmoides</i>	
Phenolics	Coniferin	[44]
	Koaburaside	[44]
	Syringin	[44]

leaf extract of EU at the dose of 0.175 g/100 g for 10 weeks reduced the concentrations of triglycerides, total cholesterol, low-density lipoprotein cholesterol (LDL-C), non-high-density lipoprotein cholesterol (non-HDL-C), and free acids in plasma and hepatic lipids compared to control group (fed 10 g, coconut oil, 0.2% cholesterol, w/w) [34]. In a similar manner, 1 mg or 5 mg intraduodenal injection of EU leaf extract reduced plasma triglyceride levels [65].

3.2. Antioxidant Effects. Antioxidant compounds from *Eucommia* plant reduced the level of free radicals [66, 67] and improved the disease condition caused by oxidative stress [68, 69]. Strong antioxidant properties of EU have been established under *in vivo* and *in vitro* studies [70, 71]. Extracts from EU reduced the level of hydrogen peroxide which expresses some caspase proteins by MC3T3E1 cells up to half concentration from 12.5 to 25 μ g/mL [71]. Extract of EU was reported to increase the actions of erythrocyte, superoxide dismutase, and catalase and glutathione peroxidase and reduce the concentration of hydrogen peroxide and lipid peroxide in erythrocytes, liver, and kidney [70]. Studies on diabetic rats indicated that superoxide dismutase (SOD) can be enhanced by *Eucommia* bark. *Eucommia* also increases the level of other antioxidant enzymes in the blood to neutralize free radicals [70].

Phenolics and flavonoids of medicinal herbs contributed significantly to oxidative activities in EU [34, 69–72]. Phenolics and flavonoids safely react with free radicals by donating a hydrogen atom or an electron and terminate chain reaction before the vital organs are damaged [73]. Antioxidant properties from leaves of the EU roasted cortex and seeds were analyzed by calculating radical scavenging activity of 2,2-diphenyl-1-picrylhydrazyl and ferric reducing antioxidant power and lipid peroxidation inhibition capacity in a β -carotene/linoleic acid system. Results indicated that leaf of the extract showed maximum DPPH radical scavenging activity with reducing rate and inhibition rate of 81.40%, followed by butylated hydroxytoluene (BHT) (76.60%) and the roasted cortex extract (16.72%). However, the seed extract

had the lowest activity of 7.65%. In ferric reducing antioxidant power assays, the order of ferric reducing activities of EU extracts from leaf, seed, and roasted cortex was compared with positive control. In the β -carotene/linoleic acid emulsion system, the leaf extract showed better antioxidant capacity (43.58%) than the roasted cortex extract (26.71%) or seed extract (25.10%) [74].

In addition, aucubin compounds of EU have been demonstrated to exhibit photoprotective effects against oxidative stress. Ultraviolet (UV) B radiation produces free radicals in the skin which induce the synthesis of metalloproteinases (MMPs) causing photoaging in the skin, wrinkling, and discoloration which are prone to cancer. Aucubin played a vital role in defense mechanism against free radicals caused by UV irradiation [75].

3.3. Antibacterial, Antiviral, and Anti-Inflammatory Activity. EU have been reported to inhibit the growth of bacteria and reduce the secretion of proinflammatory cytokines in few studies. Ethanol extracts of EU at the dose rate of 0.1 and 1.0 mg/mL of 95% (v/v) were reported to exhibit some antibacterial (against *Acinetobacter baumannii* and *Staphylococcus aureus*) and antifungal (against *Aspergillus fumigatus*) effects [15–34, 65–76]. Furthermore, it has been reported that the same concentration of 0.1 mg/mL EU extracts reduces the secretion of proinflammatory cytokines including tumor necrosis factor- α (TNF- α), interleukin-8 (IL-8), and IL-1 β by human monocytic (THP-1) cells pretreated with heat-killed *P. acnes*. Aqueous extract of EU significantly decreased cyclooxygenase-2 (COX-2) enzyme with IC₅₀ = 9.92 mg/mL, although the effects were lowered compared with nonsteroid anti-inflammatory drugs [54]. Cortex of EU at the concentration of 0.1 and 0.5 mg/mL decreased production of (TNF- α , IL-6, and COX-2) prostaglandin E and nitric oxide [77].

Suppression of HIV infection has also been reported with daily intake of EU extracts or its alkaline extracts in tea formula. Alkaline extract of EU leaf in combination with 22% uronic acid, 27% reducing sugars, and 46% neutral sugars reduced HIV-induced cytopathicity (HTLV-III) with

extremely low cytotoxicity in infected MT-4 cells (EC₅₀) [78]. Lv et al. also demonstrated that the samples from EU Oliver had potent inhibitory activity against the HIV gp41 six-helix bundle formation [79].

3.4. Antiobesity Effects. Previous studies have shown that EU has antiobesity and antimetabolic syndrome properties [8, 26, 34, 80, 81]. It has been demonstrated that both *Eucommia* leaf extract (ELE) and *Eucommia* green leaf powder (EGLP) markedly suppressed body weight and white adipose tissue (WAT) in female ICR mice fed high-fat diets (HFD). The antiobesity effect of *Eucommia* green leaf extract (EGLE) has been linked to various compounds such as geniposidic acid, asperuloside, and chlorogenic acid which was isolated from the extract [8]. Application of water extract from the leaf of EU at the rate of 5% diet was reported to reduce fat accumulation rate in osteoporotic mice [4] although application of 500–1000 mg/kg EU leaf extract beyond 4 weeks showed no effect on fat accumulation in fructose overfed rats [26].

Antiobesity and antimetabolic syndrome activity in rat fed with a 35% high-fat diet could be maintained through secretion and regulation of adipocytokines that depend on the accumulation of visceral fat to improve insulin resistance or hyperlipidemia [80]. Administration of EU extracts at the concentration of 300–1600 mg/kg intake has been reported to enhance gene expressions for fat oxidation [81]. Administration of the extract was confirmed to increase fat oxidation in liver [34, 54, 65–81]. This increased fat oxidation in liver following administration of EU extract was attributed to the rate limiting stages of β -oxidation (CPT1A, ACOX1, and ACADVL), α -oxidation, and ω -oxidation (CYP4A1) [81].

3.5. Neuroprotective Effects. The stem bark extract of EU exhibited acetylcholinesterase inhibition properties *in vitro* (172 μ g/mL) IC₅₀ and neuroprotective effects against beta-amyloid proteins [4]. It also inhibits 30–70% of cytotoxicity and efficacy of oxidative biomarkers when applied at a concentration of 2.5 μ g/mL [10]. Stem bark extract of EU showed higher protection activity against memory dysfunctions at the dose of 10–20 mg/kg with intracerebral injection of beta-amyloid proteins in rats [4].

3.6. Metabolic Modulation and Bones. *Eucommia* cortex extract can be used in the control of osteoporosis. This is because *Eucommia* extract is actively involved in mechanisms which initiate osteoblast, enhance osteogenesis, decrease osteoclast, and thus prevent osteolysis [82]. Total glycosides from *Eucommia ulmoides* seed (TGEUS) have been shown to improve bone density and femur strength in rats [83]. Daily administration of TGEUS at the rate of 400 mg/kg body weight/day to normal and Dawley rats was reported to significantly increase bone mineral density and showed improvements in microarchitecture structure of the femur bone [83].

Eucommia cortex extract was reported to induce the release of growth hormone (GH) responsible for bone maturation and bone remodeling. Products of alcoholic extraction from *Eucommia* bark were reported to be very potent in the

release of growth hormone secretagogue. Increasing signals of estrogen receptor alpha has been shown to increase the growth of bone [84]. An exception to this effect was noticed in ovariectomized rats which showed no effect on the growth of bone [47, 82]. In menopausal research model, 5% diet of the EU was observed to minimize the bone loss in ovariectomized rats [61]. *Eucommia* cortex fed at the dosage of 300–500 mg/kg showed reduced bone mass which is not significantly different from group fed with estradiol drug [61]. Antioxidant properties of *Eucommia* leaf extract were also reported to contribute positively to the promotion of bone growth by improving cell integrity during oxidative stress when applied at a reduced dosage (6.25 μ g/mL) [71]. Therefore, *Eucommia* extract can be established as a therapeutic agent under conditions of osteoporosis [85].

3.7. Phytoestrogenic Properties. EU was reported to exhibit phytoestrogenic and androgenic properties [84]. *Eucommia* bark contains isoflavonoids, with estrogen like properties, which bind to human estrogen receptors. None of these isoflavonoids has male hormone like effect that interacts with human androgen receptor. *Eucommia* bark has been reported to show bimodal phytoandrogenic and hormone enhancing effects [84]. Androgen receptors play a key role in male as well as in female physiology such as skeletal muscle development, bone density, and sex drive [86, 87]. Ethanol extracts of *Eucommia* bark were reported to attach in a weak manner to activated androgen receptors with high affinity and produce testosterone at the rate of 5–25 ng/mL in mammalian COS-7 cells [84]. Oral induction of the ethanol extract showed no increase in prostatic weights at the dose of 1–50 mg. However, 20% increases in prostatic weights were observed by increasing the dosage up to 5000 μ g injection [84]. Application of EU at a concentration of 50 ng/mL enhanced the signals of estradiol in a manner similar to androgen receptors [84]. However, the promoting effect of EU on the cortisol and progesterone receptors was not observed [84].

In vivo animal studies conducted using oral administration of EU extracts potentiated androgenic and hormonal effects. A form of tripartite synergism between sex steroid receptors, sex hormones, and lipidic augmenters isolated from EU was found by Ong and Tan [84]. It has been shown that the activities of sex hormone in the body are optimized with the application of EU [84].

3.8. Hepatoprotective Effects. Study was conducted on different doses of *Eucommia ulmoides* and carbon tetrachloride on Sprague Dawley male rats to investigate the protective effects of EU in response to CCl₄ induced acute liver lipid accumulation. Results demonstrated that *Eucommia ulmoides* Oliv. cortex extracts (EUCE) significantly decreased the hepatic lipid accumulation induced by CCl₄. EU enhances lysosomal enzyme activity relieving protein folding requirement which turns into attenuation of ER stress. ApoB secretion was improved by effects of ER stress; along this, it regulates biotransformation of CCl₄ and its resultant inhibition of ROS accumulation [88].

4. Future Perspective and Conclusion

This review paper discusses health-promoting properties of EU on cardiovascular system and antioxidant, antibacterial, antiviral, anti-inflammatory, antiobesity, and neuroprotective effects and metabolic modulation on bones and phytoestrogenic properties. These health-promoting properties have attracted much interest in the extraction and functional development of active ingredients of EU. In further studies, molecular mechanisms underlying certain health-promoting properties of EU need to be explored.

Conflict of Interests

The authors declare that there is no conflict of interests.

Acknowledgments




This work was supported by the National Natural Science Foundation of China (nos. 31330075, 31560640, 31372326, and 31301989), the Science and Technology Department of Hunan province (2015JC3126), and the State Key Laboratory of Animal Nutrition (2004DA125184F1401). The authors are also thankful to CAS-TWAS President's Fellowship and UCAS financial and infrastructure support.

References

- [1] V. B. Call and D. L. Dilcher, "The fossil record of *Eucommia* (Eucommiaceae) in North America," *American Journal of Botany*, vol. 84, no. 6, pp. 798–814, 1997.
- [2] G.-C. Yen and C.-L. Hsieh, "Antioxidant activity of extracts from Du-zhong (*Eucommia ulmoides*) toward various lipid peroxidation models *in vitro*," *Journal of Agricultural and Food Chemistry*, vol. 46, no. 10, pp. 3952–3957, 1998.
- [3] Q. Zhang, Y.-Q. Su, F.-X. Yang, J.-N. Peng, X.-H. Li, and R.-C. Sun, "Antioxidative activity of water extracts from leaf, male flower, raw cortex and fruit of *Eucommia ulmoides* Oliv," *Forest Products Journal*, vol. 57, no. 12, pp. 74–78, 2007.
- [4] S.-H. Kwon, H.-K. Lee, J.-A. Kim et al., "Neuroprotective effects of *Eucommia ulmoides* Oliv. Bark on amyloid beta_{25–35}-induced learning and memory impairments in mice," *Neuroscience Letters*, vol. 487, no. 1, pp. 123–127, 2011.
- [5] Y. Nakazawa, T. Takeda, N. Suzuki et al., "Histochemical study of trans-polyisoprene accumulation by spectral confocal laser scanning microscopy and a specific dye showing fluorescence solvatochromism in the rubber-producing plant, *Eucommia ulmoides* Oliver," *Planta*, vol. 238, no. 3, pp. 549–560, 2013.
- [6] A. Kulomaa, H. Sirén, and M.-L. Riekkola, "Identification of antioxidative compounds in plant beverages by capillary electrophoresis with the marker index technique," *Journal of Chromatography A*, vol. 781, no. 1–2, pp. 523–532, 1997.
- [7] X. He, J. Wang, M. Li et al., "*Eucommia ulmoides* Oliv.: ethnopharmacology, phytochemistry and pharmacology of an important traditional Chinese medicine," *Journal of Ethnopharmacology*, vol. 151, no. 1, pp. 78–92, 2014.
- [8] T. Hirata, T. Kobayashi, A. Wada et al., "Anti-obesity compounds in green leaves of *Eucommia ulmoides*," *Bioorganic & Medicinal Chemistry Letters*, vol. 21, no. 6, pp. 1786–1791, 2011.
- [9] R. Tundis, M. R. Loizzo, F. Menichini, G. A. Statti, and F. Menichini, "Biological and pharmacological activities of iridoids: recent developments," *Mini-Reviews in Medicinal Chemistry*, vol. 8, no. 4, pp. 399–420, 2008.
- [10] Y. Zhou, M. Liang, W. Li et al., "Protective effects of *Eucommia ulmoides* Oliv. bark and leaf on amyloid β -induced cytotoxicity," *Environmental Toxicology and Pharmacology*, vol. 28, no. 3, pp. 342–349, 2009.
- [11] C. Takamura, T. Hirata, T. Ueda et al., "Iridoids from the green leaves of *Eucommia ulmoides*," *Journal of Natural Products*, vol. 70, no. 8, pp. 1312–1316, 2007.
- [12] L.-Z. Lin and J. M. Harnly, "A screening method for the identification of glycosylated flavonoids and other phenolic compounds using a standard analytical approach for all plant materials," *Journal of Agricultural and Food Chemistry*, vol. 55, no. 4, pp. 1084–1096, 2007.
- [13] P. Mattila and J. Hellström, "Phenolic acids in potatoes, vegetables, and some of their products," *Journal of Food Composition and Analysis*, vol. 20, no. 3–4, pp. 152–160, 2007.
- [14] X. Chen, X. Sang, S. Li, S. Zhang, and L. Bai, "Studies on a chlorogenic acid-producing endophytic fungi isolated from *Eucommia ulmoides* Oliver," *Journal of Industrial Microbiology & Biotechnology*, vol. 37, no. 5, pp. 447–454, 2010.
- [15] L. Zhang, A. S. Ravipati, S. R. Koyyalamudi et al., "Anti-fungal and anti-bacterial activities of ethanol extracts of selected traditional Chinese medicinal herbs," *Asian Pacific Journal of Tropical Medicine*, vol. 6, no. 9, pp. 673–681, 2013.
- [16] Z. Chen, J. He, D. Tang, Q. Shi, and L. Qunxue, "Nutrient distribution and uptake characteristics in *Fucommia ulmoides* Oliv. plant," *Journal of Northwest Forestry College*, vol. 19, no. 3, pp. 15–17, 2004.
- [17] J. Cheng, Y. Y. Zhao, Y. X. Cui, and T. M. Cheng, "Studies on flavonoids from leave of *Eucommia ulmoides* Oliv," *Zhongguo Zhong Yao Za Zhi*, vol. 25, no. 5, pp. 284–286, 2000 (Chinese).
- [18] L. Chen and Z. Yang, "Study on dynamic variation of three flavonoids contents in folium eucommiae harvested from different seasons," *Strait Pharmaceutical Journal*, vol. 4, pp. 29–30, 2007.
- [19] H. M. Hua, H. Q. Yin, B. Q. Li et al., "Study on the constituents of the bark of *Eucommia ulmoides*," *Molecular Plant Breeding*, vol. 1, pp. 801–803, 2003.
- [20] N. Okada, K. Shirata, M. Niwano, H. Koshino, and M. Uramoto, "Immunosuppressive activity of a monoterpene from *Eucommia ulmoides*," *Phytochemistry*, vol. 37, no. 1, pp. 281–282, 1994.
- [21] Y. K. Hong, W. J. Liu, T. Li, and S. She, "Optimization of extraction of *Eucommia ulmoides* polysaccharides by response surface methodology," *Carbohydrate Polymers*, vol. 92, no. 2, pp. 1761–1766, 2013.
- [22] E. Liu, L. Han, J. Wang et al., "*Eucommia ulmoides* bark protects against renal injury in cadmium-challenged rats," *Journal of Medicinal Food*, vol. 15, no. 3, pp. 307–314, 2012.
- [23] M. Tomoda, R. Gonda, and N. Shimizu, "A reticuloendothelial system-activating glycan from the barks of *Eucommia ulmoides*," *Phytochemistry*, vol. 29, no. 10, pp. 3091–3094, 1990.
- [24] Y. R. Sun, J. X. Dong, and S. G. Wu, "Studies on chemical constituents from *Eucommia ulmoides* Oliv," *Journal of Chinese Medicinal Materials*, vol. 27, no. 5, pp. 33–35, 2004 (Chinese).
- [25] L. X. Zhang, X. Y. Ji, B. B. Tan, Y. Z. Liang, and N. N. Liang, "Identification of the composition of fatty acids in *Eucommia ulmoides* seed oil by fraction chain length and mass spectrometry," *Food Chemistry*, vol. 121, no. 3, pp. 815–819, 2010.

- [26] X. Jin, K. Amitani, Y. Zamami et al., "Ameliorative effect of *Eucommia ulmoides* Oliv. leaves extract (ELE) on insulin resistance and abnormal perivascular innervation in fructose-drinking rats," *Journal of Ethnopharmacology*, vol. 128, no. 3, pp. 672–678, 2010.
- [27] C. Lang, Z. Liu, H. W. Taylor, and D. G. Baker, "Effect of *Eucommia ulmoides* on systolic blood pressure in the spontaneous hypertensive rat," *American Journal of Chinese Medicine*, vol. 33, no. 2, pp. 215–230, 2005.
- [28] L.-F. Luo, W.-H. Wu, Y.-J. Zhou, J. Yan, G.-P. Yang, and D.-S. Ouyang, "Antihypertensive effect of *Eucommia ulmoides* Oliv. extracts in spontaneously hypertensive rats," *Journal of Ethnopharmacology*, vol. 129, no. 2, pp. 238–243, 2010.
- [29] F. Greenway, Z. Liu, Y. Yu, and A. Gupta, "A clinical trial testing the safety and efficacy of a standardized *Eucommia ulmoides* oliver bark extract to treat hypertension," *Alternative Medicine Review*, vol. 16, no. 4, pp. 338–347, 2011.
- [30] T. Deyama, S. Nishibe, and Y. Nakazawa, "Constituents and pharmacological effects of *Eucommia* and *Siberian ginseng*," *Acta Pharmacologica Sinica*, vol. 22, no. 12, pp. 1057–1070, 2001.
- [31] C. Y. Kwan, C. X. Chen, T. Deyama, and S. Nishibe, "Endothelium-dependent vasorelaxant effects of the aqueous extracts of the *Eucommia ulmoides* Oliv. leaf and bark: implications on their antihypertensive action," *Vascular Pharmacology*, vol. 40, no. 5, pp. 229–235, 2004.
- [32] Z.-Y. Li, J. Gu, J. Yan et al., "Hypertensive cardiac remodeling effects of lignan extracts from *Eucommia ulmoides* Oliv. bark—a famous traditional Chinese medicine," *The American Journal of Chinese Medicine*, vol. 41, no. 4, pp. 801–815, 2013.
- [33] J. Gu, J.-J. Wang, J. Yan et al., "Effects of lignans extracted from *Eucommia ulmoides* and aldose reductase inhibitor epalrestat on hypertensive vascular remodeling," *Journal of Ethnopharmacology*, vol. 133, no. 1, pp. 6–13, 2011.
- [34] M.-S. Choi, U. J. Jung, H.-J. Kim et al., "Du-zhong (*Eucommia ulmoides* Oliver) leaf extract mediates hypolipidemic action in hamsters fed a high-fat diet," *The American Journal of Chinese Medicine*, vol. 36, no. 1, pp. 81–93, 2008.
- [35] T. Deyama, T. Ikawa, and S. Nishibe, "The constituents of *Eucommia ulmoides* OLIV. II. Isolation and structures of three new lignan glycosides," *Chemical & Pharmaceutical Bulletin*, vol. 33, no. 9, pp. 3651–3657, 1985.
- [36] T. Deyama, T. Ikawa, S. Kitagawa, and S. Nishibe, "The constituents of *Eucommia ulmoides* Oliv. III. Isolation and structure of a new lignan glycoside," *Chemical and Pharmaceutical Bulletin*, vol. 34, no. 2, pp. 523–527, 1986.
- [37] T. Deyama, T. Ikawa, S. Kitagawa, and S. Nishibe, "The constituents of *Eucommia ulmoides* Oliv. V. Isolation of dihydroxydehydrodiconiferyl alcohol isomers and phenolic compounds," *Chemical & Pharmaceutical Bulletin*, vol. 35, no. 5, pp. 1785–1789, 1987.
- [38] T. Deyama, "The constituents of *Eucommia ulmoides* Oliv. I. Isolation of (+)-medioresinol Di-O- β -D-glucopyranoside," *Chemical and Pharmaceutical Bulletin*, vol. 31, no. 9, pp. 2993–2997, 1983.
- [39] C. J. Sih, P. R. Ravikumar, F. C. Huang, C. Buckner, and H. Whitlock Jr., "Isolation and synthesis of pinoresinol diglucoside, a major antihypertensive principle of Tu-Chung (*Eucommia ulmoides*, Oliver)," *Journal of the American Chemical Society*, vol. 98, no. 17, pp. 5412–5413, 1976.
- [40] S.-Y. Shi, M.-J. Peng, Y.-P. Zhang, and S. Peng, "Combination of preparative HPLC and HSCCC methods to separate phosphodiesterase inhibitors from *Eucommia ulmoides* bark guided by ultrafiltration-based ligand screening," *Analytical and Bioanalytical Chemistry*, vol. 405, no. 12, pp. 4213–4223, 2013.
- [41] J.-X. Wang, X.-H. Hao, H.-M. Lei, X.-J. You, and J.-S. Li, "Study on the constituents of the bark of *Eucommia ulmoides* Oliv," *Northwest Pharmaceutical Journal*, vol. S1, pp. 13–15, 1996 (Chinese).
- [42] C. L. Si, X. J. Deng, D. Wang, P. T. Liu, and Z. Liu, "Study on chemical compositions of *Eucommia ulmoides* Oliv. inner bark and its extractives," *Chemistry and Industry of Forest Products*, vol. 28, no. 5, pp. 7–10, 2008 (Chinese).
- [43] L. N. Yao, *Studies on the chemical constituents of Eucommia ulmoides Oliv [The Master's Degree Thesis]*, Tianjin University, Tianjin, China, 2010 (Chinese).
- [44] M. B. Gewali, M. Hattori, and T. Namba, "Constituents of the stems of *Eucommia ulmoides* Oliv," *Shoyakugaku Zasshi*, vol. 42, pp. 247–248, 1988 (Japanese).
- [45] L. N. Yao, "Studies on the chemical constituents of *Eucommia ulmoides* Oliv," *Zhong Yao Cai*, vol. 27, no. 5, pp. 341–343, 2004 (Chinese).
- [46] C. J. Chen, S. Shen, T. S. Xiao, Q. Wang, J. Q. Jiang, and Y. Li, "Chemical constituents from the bark of *Eucommia ulmoides*," *Asia-Pacific Traditional Medicine*, vol. 8, pp. 25–26, 2012 (Chinese).
- [47] D. Li, H. L. Wang, and J. M. Chen, "The chemical constituents of *Eucommia ulmoides* Oliv.," *Chinese Bulletin of Botany*, vol. 28, pp. 528–534, 1986.
- [48] T. Deyama, T. Ikawa, S. Kitagawa, and S. Nishibe, "The constituents of *Eucommia ulmoides* OLIV. IV. Isolation of a new sesquiliglan glycoside and iridoids," *Chemical & Pharmaceutical Bulletin*, vol. 34, no. 12, pp. 4933–4938, 1986.
- [49] A. Shimoyama, M. Yamadaki, and Y. Nakazawa, "Constituents of the leaf of *Eucommia ulmoides* Oliv," *Shoyakugaku Zasshi*, vol. 47, p. 56, 1993 (Japanese).
- [50] T. Deyama, T. Ikawa, S. Kitagawa, and S. Nishibe, "The constituents of *Eucommia ulmoides* Oliv. VI. Isolation of a new sesquiliglan and neolignan glycosides," *Chemical & Pharmaceutical Bulletin*, vol. 35, no. 5, pp. 1803–1807, 1987.
- [51] J. W. Xu, D. Li, and P. Zhao, "The constituents of *Eucommia ulmoides* Oliv," *Botanical Bulletin*, vol. 31, p. 132, 1989.
- [52] R. Gonda, M. Tomoda, N. Shimizu, and M. Kanari, "An acidic polysaccharide having activity on the reticuloendothelial system from the bark of *Eucommia ulmoides*," *Chemical & Pharmaceutical Bulletin*, vol. 38, no. 7, pp. 1966–1969, 1990.
- [53] L. P. Sun, L. Ma, B. Zhang, and H. Xu, "Research progress of flavonoids in *Eucommia ulmoides* Oliv," *Food Science and Technology International*, vol. 30, pp. 359–363, 2009 (Chinese).
- [54] B. H. Kim, K. S. Park, and I.-M. Chang, "Elucidation of anti-inflammatory potencies of *Eucommia ulmoides* bark and *Plantago asiatica* seeds," *Journal of Medicinal Food*, vol. 12, no. 4, pp. 764–769, 2009.
- [55] Q. Wei, D. M. Wang, X. H. Ma, and K. J. Zang, "A study on the measurement of flavonoids in the leaves of *Eucommia ulmoides* Oliv," *Journal of Northwest Sci-Tech University Agriculture and Forestry*, vol. 29, pp. 119–123, 2001 (Chinese).
- [56] H. Y. Kim, B. H. Moon, H. J. Lee, and D. H. Choi, "Flavonol glycosides from the leaves of *Eucommia ulmoides* O. with glycation inhibitory activity," *Journal of Ethnopharmacology*, vol. 93, no. 2–3, pp. 227–230, 2004.
- [57] A. Bianco, C. Iavarone, and C. Trogolo, "Structure of eucommiol, a new cyclopentenoid-tetrol from *Eucommia ulmoides*," *Tetrahedron*, vol. 30, no. 23–24, pp. 4117–4121, 1974.

- [58] T. Nakamura, Y. Nakazawa, S. Onizuka et al., "Antimutagenicity of Tochu tea (an aqueous extract of *Eucommia ulmoides* leaves) I, the clastogen-suppressing effects of Tochu tea in CHO cells and mice," *Mutation Research*, vol. 388, no. 1, pp. 7–20, 1997.
- [59] A. Bianeo, C. C. Bonini, C. Iavarone, and C. Trogolo, "Structure elucidation of eucommioside (2''-O- β -D-glucopyranosyl eucommiol) from *Eucommia ulmoides*," *Phytochemistry*, vol. 21, pp. 201–203, 1982.
- [60] M. Hattori, Q. M. Che, and M. B. Gewali, "Constituents of the leaves of *Eucommia ulmoides* Oliv.," *Shoyakugaku Zasshi*, vol. 42, pp. 76–80, 1988 (Japanese).
- [61] W. Zhang, T. Fujikawa, K. Mizuno et al., "*Eucommia* leaf extract (ELE) prevents OVX-induced osteoporosis and obesity in rats," *American Journal of Chinese Medicine*, vol. 40, no. 4, pp. 735–752, 2012.
- [62] Y. Z. Peng, M. R. Deng, F. Zhou, and S. Liu, "Chemical constituents from leaves of *eucommia ulmoides* collected from Zhangjiajie," *Central South Pharmacy*, vol. 11, pp. 179–181, 2013 (Chinese).
- [63] C. Tanaka, T. Nakamura, Y. Nakazawa, and T. Nohara, "A new triterpenoid from the leaves of *Eucommia ulmoides* OLIV.," *Chemical & Pharmaceutical Bulletin*, vol. 45, no. 8, pp. 1379–1380, 1997.
- [64] S. Yahara, K. Kato, Y. Nakazawa, Y. Toda, and T. Nohara, "New iridoid trimers and tetramers from seeds of *Eucommia ulmoides*," *Chemical and Pharmaceutical Bulletin*, vol. 38, no. 1, pp. 267–269, 1990.
- [65] Y. Horii, M. Tanida, J. Shen et al., "Effects of *Eucommia* leaf extracts on autonomic nerves, body temperature, lipolysis, food intake, and body weight," *Neuroscience Letters*, vol. 479, no. 3, pp. 181–186, 2010.
- [66] Y. Cai, Q. Luo, M. Sun, and H. Corke, "Antioxidant activity and phenolic compounds of 112 traditional Chinese medicinal plants associated with anticancer," *Life Sciences*, vol. 74, no. 17, pp. 2157–2184, 2004.
- [67] S. Dragland, H. Senoo, K. Wake, K. Holte, and R. Blomhoff, "Several culinary and medicinal herbs are important sources of dietary antioxidants," *Journal of Nutrition*, vol. 133, no. 5, pp. 1286–1290, 2003.
- [68] A. C. Akinmoladun, E. M. Obuotor, and E. O. Farombi, "Evaluation of antioxidant and free radical scavenging capacities of some Nigerian indigenous medicinal plants," *Journal of Medicinal Food*, vol. 13, no. 2, pp. 444–451, 2010.
- [69] R. Hendra, S. Ahmad, E. Oskoueian, A. Sukari, and M. Y. Shukor, "Antioxidant, anti-inflammatory and cytotoxicity of *Phaleria macrocarpa* (Boerl.) Scheff fruit," *BMC Complementary and Alternative Medicine*, vol. 11, article 110, 2011.
- [70] S. A. Park, M.-S. Choi, U. J. Jung et al., "*Eucommia ulmoides* oliver leaf extract increases endogenous antioxidant activity in type 2 diabetic mice," *Journal of Medicinal Food*, vol. 9, no. 4, pp. 474–479, 2006.
- [71] J. Lin, Y. J. Fan, C. Mehl et al., "*Eucommia ulmoides* Oliv. Antagonizes H₂O₂-induced rat osteoblastic MC3T3-E1 apoptosis by inhibiting expressions of caspases 3, 6, 7, and 9," *Journal of Zhejiang University Science B*, vol. 12, no. 1, pp. 47–54, 2011 (Chinese).
- [72] L. Zhang, A. S. Ravipati, S. R. Koyyalamudi et al., "Antioxidant and anti-inflammatory activities of selected medicinal plants containing phenolic and flavonoid compounds," *Journal of Agricultural and Food Chemistry*, vol. 59, no. 23, pp. 12361–12367, 2011.
- [73] S. Petti and C. Scully, "Polyphenols, oral health and disease: a review," *Journal of Dentistry*, vol. 37, no. 6, pp. 413–423, 2009.
- [74] Z. Xu, M. Tang, Y. Li, F. Liu, X. Li, and R. Dai, "Antioxidant properties of du-zhong (*Eucommia ulmoides* Oliv.) extracts and their effects on color stability and lipid oxidation of raw pork patties," *Journal of Agricultural and Food Chemistry*, vol. 58, no. 12, pp. 7289–7296, 2010 (Chinese).
- [75] J. N. Ho, Y. H. Lee, J. S. Park et al., "Protective effects of aucubin isolated from *Eucommia ulmoides* against UVB-induced oxidative stress in human skin fibroblasts," *Biological and Pharmaceutical Bulletin*, vol. 28, no. 7, pp. 1244–1248, 2005.
- [76] T.-H. Tsai, T.-H. Tsai, W.-H. Wu, J. T.-P. Tseng, and P.-J. Tsai, "In vitro antimicrobial and anti-inflammatory effects of herbs against *Propionibacterium acnes*," *Food Chemistry*, vol. 119, no. 3, pp. 964–968, 2010.
- [77] M.-C. Kim, D.-S. Kim, S.-J. Kim et al., "Eucommiae cortex inhibits TNF- α and IL-6 through the suppression of caspase-1 in lipopolysaccharide-stimulated mouse peritoneal macrophages," *The American Journal of Chinese Medicine*, vol. 40, no. 1, pp. 135–149, 2012.
- [78] M. Nakano, H. Nakashima, and Y. Itoh, "Anti-human immunodeficiency virus activity of oligosaccharides from rooibos tea (*Aspalathus linearis*) extracts in vitro," *Leukemia*, vol. 11, supplement 3, pp. 128–130, 1997.
- [79] L. Lv, Y. R. Sun, W. Xu, S. W. Liu, J. J. Rao, and S. G. Wu, "Isolation and characterization of the anti-HIV active component from *Eucommia ulmoides*," *Zhong Yao Cai*, vol. 31, no. 6, pp. 847–850, 2008 (Chinese).
- [80] T. Fujikawa, T. Hirata, A. Wada et al., "Chronic administration of *Eucommia* leaf stimulates metabolic function of rats across several organs," *British Journal of Nutrition*, vol. 104, no. 12, pp. 1868–1877, 2010.
- [81] Y. Kobayashi, T. Hiroi, M. Araki et al., "Facilitative effects of *Eucommia ulmoides* on fatty acid oxidation in hypertriglyceridaemic rats," *Journal of the Science of Food and Agriculture*, vol. 92, no. 2, pp. 358–365, 2012.
- [82] R. Zhang, Z. G. Liu, C. Li et al., "Du-Zhong (*Eucommia ulmoides* Oliv.) cortex extract prevent OVX-induced osteoporosis in rats," *Bone*, vol. 45, no. 3, pp. 553–559, 2009.
- [83] Y. Li, M.-J. Wang, S. Li et al., "Effect of total glycosides from *Eucommia ulmoides* seed on bone microarchitecture in rats," *Phytotherapy Research*, vol. 25, no. 12, pp. 1895–1897, 2011.
- [84] V. Y. C. Ong and B. K. H. Tan, "Novel phytoandrogens and lipidic augmenters from *Eucommia ulmoides*," *BMC Complementary and Alternative Medicine*, vol. 7, article 3, 2007.
- [85] H. Ha, J. Ho, S. Shin et al., "Effects of *Eucommiae* cortex on osteoblast-like cell proliferation and osteoclast inhibition," *Archives of Pharmacal Research*, vol. 26, no. 11, pp. 929–936, 2003.
- [86] S. L. Davison and R. Bell, "Androgen physiology," *Seminars in Reproductive Medicine*, vol. 24, no. 2, pp. 71–77, 2006.
- [87] C. A. Quigley, A. De-Bellis, K. B. Marschke, M. K. El-Awady, E. M. Wilson, and F. S. French, "Androgen receptor defects: historical, clinical, and molecular perspectives," *Endocrine Reviews*, vol. 16, no. 3, pp. 271–321, 1995.
- [88] C.-F. Jin, B. Li, S.-M. Lin, R.-K. Yadav, H.-R. Kim, and H.-J. Chae, "Mechanism of the inhibitory effects of *Eucommia ulmoides* Oliv. cortex extracts (EUCE) in the CCl₄-induced acute liver lipid accumulation in rats," *International Journal of Endocrinology*, vol. 2013, Article ID 751854, 11 pages, 2013.



Hindawi
Submit your manuscripts at
<http://www.hindawi.com>

