

REVIEW

Open Access

'Hearts and minds': association, causation and implication of cognitive impairment in heart failure

Jane A Cannon¹, John JV McMurray¹ and Terry J Quinn^{2*}

Abstract

The clinical syndrome of heart failure is one of the leading causes of hospitalisation and mortality in older adults. An association between cognitive impairment and heart failure is well described but our understanding of the relationship between the two conditions remains limited. In this review we provide a synthesis of available evidence, focussing on epidemiology, the potential pathogenesis, and treatment implications of cognitive decline in heart failure. Most evidence available relates to heart failure with reduced ejection fraction and the syndromes of chronic cognitive decline or dementia. These conditions are only part of a complex heart failure-cognition paradigm. Associations between cognition and heart failure with preserved ejection fraction and between acute delirium and heart failure also seem evident and where data are available we will discuss these syndromes. Many questions remain unanswered regarding heart failure and cognition. Much of the observational evidence on the association is confounded by study design, comorbidity and insensitive cognitive assessment tools. If a causal link exists, there are several potential pathophysiological explanations. Plausible underlying mechanisms relating to cerebral hypoperfusion or occult cerebrovascular disease have been described and it seems likely that these may coexist and exert synergistic effects. Despite the prevalence of the two conditions, when cognitive impairment coexists with heart failure there is no specific guidance on treatment. Institution of evidence-based heart failure therapies that reduce mortality and hospitalisations seems intuitive and there is no signal that these interventions have an adverse effect on cognition. However, cognitive impairment will present a further barrier to the often complex medication self-management that is required in contemporary heart failure treatment.

Definitions and burden of heart failure

The term 'heart failure' (HF) is used to describe a condition wherein cardiac output is insufficient to meet metabolic requirements [1]. Clinically, it is defined as a syndrome where patients have typical signs and symptoms resulting from an abnormality of cardiac structure or function [2]. Contemporary terminology used to describe HF is based on left ventricular ejection fraction (EF). This is considered important not only because of prognosis (the lower the EF the poorer the survival) but also because the major trials that inform the evidence base have almost exclusively focussed on patients who have HF with reduced ejection fraction (HF-REF) [2]. A

subgroup of patients also present with classical signs and symptoms but in the context of preserved ejection fraction (HF-PEF). These patients often have evidence of diastolic dysfunction and this is considered by many as the cause of HF symptoms.

It is estimated that 1 to 2% of the adult population in developed countries have HF with the prevalence increasing to $\geq 10\%$ among patients aged over 70 years; more than half of these patients have HF-REF [3]. The most common underlying aetiology in HF-REF is coronary artery disease (CAD) resulting in myocardial damage. Other common causes include hypertension, valvular pathology, viral infection and alcohol excess [2]. HF-PEF is more common in older, female patients. It is less frequently due to CAD and more often linked to hypertension and atrial fibrillation (AF), with the diagnosis being

* Correspondence: Terry.Quinn@glasgow.ac.uk

²Department of Academic Geriatric Medicine, Institute of Cardiovascular and Medical Sciences, New Lister Building, Glasgow Royal Infirmary, Glasgow UK G4 0SF, UK

Full list of author information is available at the end of the article

one of exclusion of other non-cardiac causes of breathlessness [2].

HF admissions account for 5% of all medical admissions (making it the commonest cause of unscheduled admission in older adults) and 2% of the total UK National Health Service budget [4]. Societal and demographic changes, including aging of the general population and improved survival from CAD, will increase HF prevalence (Figure 1) with a potential doubling in HF prevalence within the next 40 years [2].

Heart failure and cognitive impairment – strength of association

The co-existence of symptomatic 'heart failure' and 'brain failure' has been recognised for decades, with a description of 'cardiogenic dementia' first introduced in the 1970s. While the co-occurrence of HF and cognitive problems will be familiar to most clinicians, the topic has received relatively little research interest compared with other aspects of cardiac disease. In collating and offering a synthesis of the available literature describing the association of HF and cognition, we have found a disparate and inconsistent literature, characterised by small sample sizes, heterogeneity and multiple potential biases. We provide a brief narrative overview of the field and have tabulated a more detailed summary of findings from available cross-sectional and prospective studies (Tables 1 to 3).

Studies describing cognitive impairment (CI) in HF-REF have estimated prevalence at anywhere between 30

and 80% of patients (Table 1). This heterogeneity results from differences in study designs, case mix and cognitive assessments employed. Accepting the limitations of the evidence, even at the more conservative estimates of prevalence, the literature would suggest that CI frequently co-exists with HF-REF (Table 1).

Cross-sectional studies of cognition in HF have value in quantifying the burden of prevalent disease but give no clues as to temporal relationship or causation. To describe the incidence and 'natural history' of cognition in HF ideally requires prospective follow-up of a cohort free from CI at inception. Few studies have utilised this design and, where data are available, the validity is limited by small sample sizes, limited follow-up with substantial attrition and use of cognitive assessment tools that may not be sensitive to modest but clinically meaningful change (Table 2). Inherent in this study design is the assumption that CI follows or is a consequence of the HF pathology [16]. A literature around 'reverse causation' in heart disease has been described. In brief, early studies describing association of psychological or 'personality' factors and heart disease assumed that the neuropsychological traits pre-dated and were probably causative in the development of the cardiac condition. Subsequent data have questioned this temporality and suggest that subclinical (undiagnosed) vascular disease may cause psychological distress phenotypes [44]. Such arguments may also hold for HF and neuropsychological disease, where both cognitive change and psychological




Table 1 Studies examining the prevalence of cognitive impairment in patients with heart failure

Study	Sample	Population	Median age in years (SD)	Study methodology	Inclusion criteria	Exclusion criteria	CV measures/criteria	Cognitive assessment tool (s) used	Results	
Zuccalà 1997 [6]	57 HF pts	Consecutive admissions to hospital	77	Cross-sectional	Not specified	Co-morbid psychiatric or physical illness and previous diagnosis of CI	LVEF (mean EF 45%) NYHA II-III	MMSE, MDBandRCPM	53% of HF pts showed global CI with MMSE less than 24	
Callegari 2002 [7]	64 HF pts, 321 healthy controls	Age <65 years and consecutive admissions to hospital	52 (8)	Cross-sectional	Not specified	Co-morbid psychiatric or neurological illness. Previous diagnosis of CI and female sex	LVEF <50% NYHA I-III Cardiopulmonary testing with treadmill Right heart catheterisation	Multidomain neuropsychiatric battery	HF pts scored lower than control group in short-term verbal memory, short-term visuospatial memory and visual spatial logical ability	
Trojano 2003 [8]	149 HF NYHA II pts	Age >65 years and consecutive admissions to hospital	HF NYHA II: 75 (7)	Cross-sectional	Not specified	Co-morbid psychiatric, neurological or physical illness. Previous diagnosis of CI	No measure of LV function	Multidomain neuropsychiatric battery	HF pts scored worse than those without HF in domains of: attention, verbal fluency, verbal learning	
	159 HF NYHA III/IV		HF NYHA III/IV: 77 (7)				NYHA II-IV			No significant difference between pts with moderate or severe HF
	207 non-HF controls		Non-HF controls: 74 (7)							
Zuccalà 2005 [9]	1,511 HF pts, 11,790 control patients	All geriatric or general medical hospital admissions	79 (9)	Cross-sectional	Not specified	Not specified	HF diagnosis based on discharge documentation	Hodkinson abbreviated mental test	35% of HF pts showed global CI 29% of non-HF pts showed global CI	
Feola 2007 [10]	60 HF-REF	Consecutive admissions to hospital	66	Cross-sectional	HF-REF: clinical HF, NYHA II-IV, LVEF ≤50%	Not specified	LVEF	Multidomain neuropsychiatric battery	23% of HF pts showed global CI	
	12 HF-PEF				HF-PEF: diagnosed based on E/A ratio, deceleration time and LV dilatation		NYHA II-IV BNP			
Debette 2007 [11]	83 HF pts	Consecutive admissions to hospital	62	Cross-sectional	Not specified	Hearing/visual impairment	LVEF <45% NYHA I-IV	MMSE	61% of HF pts showed global CI	

Table 1 Studies examining the prevalence of cognitive impairment in patients with heart failure (Continued)

Dodson 2013 [12]	282 decompensated HF pts	Age >65 years and non-consecutive admissions to hospital	80 (8)	Prospective	English speaking	Co-morbid psychiatric illness	HF diagnosis based on documentation in medical records	MMSE	25% of HF pts showed evidence of mild CI 22% of HF pts showed moderate to severe CI
Schmidt 1991 [13]	20 iDCM pts 20 healthy controls	Age <50 years and ambulatory outpatients only	iDCM: 38 (5) Healthy controls: 41 (8)	Cross-sectional	Not specified	Co-morbid psychiatric, neurological or physical illness	LVEF 14-45% NYHA II-IV	LGT-3 and ALID	Systolic HF pts performed worse than the control group in domains of attention, learning and memory and reaction time
Grubb 2000 [14]	20 HF pts with CADs 20 CAD control	Ambulatory outpatients only	HF: 68 CAD controls: 67	Cross-sectional	Not specified	Co-morbid psychiatric or neurological illness. Previous hospital admission within 6 months	HF: LVEF <40%, NYHA III/IV CAD controls: LVEF >55%, no CHF	RBMT and WMS	No difference between HF pts and control group
Riegel 2002 [15]	42 HF pts	Ambulatory outpatients only	75 (12)	Cross-sectional	English speaking	Co-morbid physical or psychiatric illness	No measure of LV function NYHA I-IV	MMSE and CIMS	29% of HF pts showed evidence of global CI
Vogels 2007 [16]	62 HF pts 53 CAD controls 42 healthy controls	Age >50 years and ambulatory outpatients only	HF: 69 (9) CAD controls: 69 (10) Healthy controls: 67 (9)	Case control	HF pts: diagnosis of HF >6 months and stable on medication >4 weeks CAD controls: IHD but no clinical CHF and EF >40%	Co-morbid physical, neurological or psychiatric illness. Previous diagnosis of CI	LVEF <40% NYHA II-IV	Multidomain neuropsychiatric battery	HF pts scored lower than the healthy control group in all domains HF pts scored lower than the IHD control group in domains of memory and mental speed IHD control group scored lower than the healthy control group in language only
Hoth 2008 [17]	31 HF pts 31 CAD controls	Age >55 years and ambulatory outpatients only	HF: 69 (9) CAD controls: 69 (9)	Cross-sectional	English speaking Minimum of 8th grade education CAD controls: angina/previous MI/PCI/PVD and HF excluded on basis of clinical examination	Co-morbid physical, neurological or psychiatric illness. Previous diagnosis of CI	LVEF <40% NYHA II-IV	Multidomain neuropsychiatric battery	Systolic HF pts scored lower than the IHD control group in domains of executive function and cognitive flexibility

Table 1 Studies examining the prevalence of cognitive impairment in patients with heart failure (Continued)

Beer 2009 [18]	31 HF pts 24 healthy controls	Ambulatory outpatients only	HF: 54 (11) Healthy controls: 56 (8)	Case control	Not specified	Co-morbid neurological illness or previous diagnosis of CI	LVEF <40% NYHA II LWHFQ	Block design, CVLT and 'F,A,S test'	Systolic HF pts scored lower than control group in all cognitive domains
Stanek 2009 [19]	40 HF pts, 35 CAD controls	Ambulatory outpatients only	70 (8)	Prospective	English speaking CAD controls: history of MI, CAD, cardiac surgery, hypertension	Co-morbid psychiatric or neurological illness. Previous diagnosis of CI	NYHA II-III CO <4 L/minute on echo	DRS	No difference between systolic HF pts and CAD control patients in all domains
Sauvé 2009 [20]	50 HF pts 50 healthy controls	Age >30 years in HF pts and >55 years in controls. Ambulatory outpatients only	HF: 63 (14) Healthy controls: 63 (14)	Case control	Diagnosis of HF >6 months	Co-morbid psychiatric or neurological illness	LVEF ≤40% NYHA II-IV	Multidomain neuropsychiatric battery	Systolic HF pts scored lower than control group in domain of verbal memory
Pressler 2010 [21]	249 HF pts 63 healthy controls 102 general medical pts	Ambulatory outpatients only	HF: 63 (15) Healthy controls: 53 (17) Medical group: 63 (12)	Cross-sectional	HF: LVEF ≤40% and clinical HF Healthy controls: absence of any medical condition or controlled CV risk factors Medical group: major chronic disorder other than HF	Co-morbid psychiatric, neurological or physical illness. Previous diagnosis of CI	NYHA LVEF	Multidomain neuropsychiatric battery	HF group performed worse than healthy and general medical groups in domains of memory, executive function and psychomotor speed
Bauer 2012 [22]	51 HF-REF, 29 HF-PEF	Age >21 years and ambulatory outpatients only	72 (12)	Cross-sectional	HF-REF: history of HF-REF >6 months, stable on medication >4 weeks, LVEF ≤40% HF-PEF: history of HF-PEF >6 months, stable on medication >4 weeks, LVEF >41%	Co-morbid psychiatric, neurological or physical illness. Previous diagnosis of CI	LVEF NYHA	Multidomain neuropsychiatric battery	HF-REF and HF-PEF pts performed worse than age- and educated- adjusted healthy control groups in executive function, attention, language, memory and psychomotor speed

Table 1 Studies examining the prevalence of cognitive impairment in patients with heart failure (Continued)

Festa 2011 [23]	169 HF-REF, 38 HF-PEF	Age >17 years and ambulatory outpatients only	69	Retrospective	On medical treatment for HF	Co-morbid neurological illness	LVEF	Multidomain neuropsychiatric battery	Low EF was associated with poor memory in pts over 63 years old Haemodynamically stable
Not receiving	mechanical circulatory support				Pts <63 years old had preserved memory function regardless of EF.				
	Steinberg 2011 [24]	55 HF pts	Ambulatory outpatients only	55 (8)	Cross-sectional	Stable clinical status	Co-morbid neurological or physical illness. Previous diagnosis of CI	LVEF ≤45% 44% of HF pts showed evidence of global CI	Multidomain neuropsychiatric battery
NYHA I-III									
6 minute walk test									
Jefferson 2011 [25]	1,114 pts from Framingham Heart Study	Age >40 and <89 years and ambulatory outpatients only	67 (9)	Cross-sectional	Not specified	Co-morbid neurological illness or previous diagnosis of CI	LVEF Cardiac MRI	Multidomain neuropsychiatric battery	U-shaped association between LVEF and cognitive performance
Miller 2012 [26]	140 HF pts	Age >50 and <85 years and ambulatory outpatients only	69 (9)	Cross-sectional	English speaking	Co-morbid psychiatric or neurological illness	No measure of LV function No NYHA classification 2 minute step test	Multidomain neuropsychiatric battery	62% of HF pts showed evidence of global CI
Almeida 2012 [27]	35 HF pts	Age >45 years and ambulatory outpatients only	HF: 69 (9)	Cross-sectional	HF: EF <40%, clinical HF ≥6 months, English speaking, NYHA I-III	Co-morbid psychiatric, neurological or physical illness. Previous diagnosis of CI	LVEF	Multidomain neuropsychiatric battery	HF pts scored lower than the healthy control group in domains of immediate/long-term memory and psychomotor speed
	56 CAD controls		CAD controls: 67 (10)		CAD controls: previous MI, English speaking, EF ≥60%, no clinical HF		NYHA		No difference between the HF group and IHD control group in cognition
	64 healthy controls		Healthy controls: 69 (11)		Healthy controls: English speaking, no previous MI/angina, EF ≥60%				

Table 1 Studies examining the prevalence of cognitive impairment in patients with heart failure (Continued)

Hawkins 2012 [28]	251 HF pts	Ambulatory outpatients only	66 (10)	Cross-sectional	English speaking	Co-morbid psychiatric illness. Previous diagnosis of CI	LVEF ≤40%	Multidomain neuropsychiatric battery	58% of HF pts had CI with poor scores in the domains of verbal learning and verbal memory
Bratzke- Bauer 2013 [29]	47 HF-REF	Age >50 years and ambulatory outpatients only	HF-REF: 75 (9)	Cross-sectional	History of HF >6 months	Co-morbid psychiatric, neurological or physical illness. Previous diagnosis of CI	LVEF	Multidomain neuropsychiatric battery	23% of the HF-REF cohort showed evidence of CI
	33 HF-PEF		HF-PEF: 68 (15)		Stable on medication ≥4 weeks		NYHA		3% of the HF-PEF cohort showed evidence of CI
Huijts 2013 [30]	491 HF-REF	Age >60 years and ambulatory outpatients only	77 (8)	Prospective	HF-REF: hospitalization within past year	Co-morbid physical illness	HF-REF: LVEF <45%, NYHA II-IV, NT-proBNP >400 pg/ml	AMT	8% of HF-REF group showed evidence of severe CI (AMT ≤7)
	120 HF-PEF		HF-PEF: NT- proBNP ≥400 pg/ml if pt <75 years or ≥800 pg/ml if pt ≥75 years		HF-PEF: LVEF ≥45%		13% of HF-PEF group showed evidence of severe CI (AMT ≤7)		
Kindermann 2012 [31]	20 decompensated HF pts	Decompensated HF: non- consecutive admissions to hospital	Decompensated HF: 60 (16)	Prospective	Decompensated HF: 60 (16)	Prospective	Decompensated HF: caused by ischaemic or DCM, symptomatic HF for ≥6 months, clinical signs of decompensation, for example, raised JVP	Co-morbid psychiatric, neurological or physical illness. Previous diagnosis of CI	LVEF <45%

Table 1 Studies examining the prevalence of cognitive impairment in patients with heart failure (Continued)

	20 stable HF pts	Stable HF: outpatients	Stable HF: 61 (17)	Stable HF pts: CHF of ischaemic or DCM, NYHA III-IV, no clinical signs/ history of decompensation for ≥ 3 months	NYHA III/IV
Stable HF group scored lower than the healthy control group in domains of intelligence and episodic memory	20 healthy controls	Healthy controls: 62 (15)			

AHA, American Heart Association; ALID, adjective list of Janke and Debus; AMT, Abbreviated Mental Test; BNP, brain natriuretic peptide; CAD, coronary artery disease; CHF, congestive heart failure; CI, cognitive impairment; CIMS, complex ideational material subset; CO, cardiac output; CV, cardiovascular; CVLT, California Verbal Learning Test; DCM, dilated cardiomyopathy; DRS, Disability Rating Scale; E/A ratio, ratio of mitral peak velocity of early filling (E) to mitral peak velocity of late filling (A); EF, ejection fraction; HF, heart failure; HF-REF, heart failure-reduced ejection fraction; HF-PEF, heart failure-preserved ejection fraction; iDCM, idiopathic dilated cardiomyopathy; IHD, ischaemic heart disease; JVP, jugular venous pressure; LGT-3, Lern und Gedachtnistest; LV, left ventricular; LVEF, left ventricular ejection fraction; LWHFQ, Living With Heart Failure Questionnaire; MDB, mental deterioration battery; MI, myocardial infarction; MMSE, Mini-Mental State Examination; MRI, magnetic resonance imaging; NT-pro BNP, N-terminal prohormone brain natriuretic peptide; NYHA, New York Heart Association; pts, patients; PCI, percutaneous coronary intervention; PVD, peripheral vascular disease; RBMT, Rivermead Behavioural Memory Test; RCPM, raven coloured progressive matrices; SD, standard deviation; WMS, Weschler Memory Scale.

Table 2 Studies examining cognitive changes over time in the heart failure population

Study	Sample	Population	Median age in years (SD)	Study methodology	Inclusion criteria	Exclusion criteria	CV measures	Cognitive assessment tool used	Follow-up period	Results
Karlsson 2005 [32]	146 CHF pts	Age >60 years and outpatients	76 (8)	Prospective	EF <45%	Co-morbid psychiatric, neurological or physical illness. Previous diagnosis of CI	LVEF NYHA	MMSE	6 months	12% of HF patients had MMSE scores <24 at baseline And 4% had MMSE scores <24 at 6 months
Tanne 2005 [33]	20 CHF underwent exercise programme 5 CHF pts as control pts	Outpatients	63 (13)	Prospective	EF ≤35% NYHA III	Co-morbid psychiatric, neurological or physical illness	LVEF NYHA Mod-Bruce ETT 6 minute walk test	Multidomain neuropsychiatric battery	18 weeks	Improvement in executive function post-exercise programme No change in cognition in control group with time
Stanek 2009 [19]	40 HF pts, 35 CAD controls	Age >53 and <84 years. Outpatients	70 (8)	Prospective	HF: English speaking NYHA II or III CO <4 L/minute CAD controls: CO ≥4 L/minute, history of MI/ CAD	Co-morbid psychiatric or neurological illness. Previous diagnosis of CI	NYHA CO	DRS	12 months	HF patients improved at 12 months, particularly in attention Cardiac controls stable at 12 months
Almeida 2013 [34]	77 HF pts 73 CAD controls 81 healthy controls	Age >45 years and outpatients	HF: 68 (10) CAD controls: 68 (10) Healthy controls: 69 (11)	Prospective	HF: EF <40%, English speaking CAD controls: previous MI and EF >60%, English speaking Healthy controls: no history of CAD, English speaking	Co-morbid psychiatric or neurological illness. Previous diagnosis of CI	NYHA LVEF 6 minute walk test	Multidomain neuropsychiatric battery	2 years	CHF group showed cognitive decline compared with CAD and healthy controls
Hjelm 2011 [35]	95 HF pts 607 non-CHF controls	Age >80 years and outpatients	84 (3)	Prospective	Not specified	Not specified	HF diagnosis based on documentation in medical records	Multidomain neuropsychiatric battery	10 years	HF patients showed significant decline in episodic memory and spatial performance compared with controls

Table 2 Studies examining cognitive changes over time in the heart failure population (Continued)

Riegel 2012 [36]	279 consecutive HF pts (HF-REF and HF-PEF)	Age <80 years and outpatients	62 (12)	Prospective	Stage C HF and English speaking	Co-morbid psychiatric or physical illness. Previous diagnosis of CI	NYHA I-IV LVEF	Multidomain neuropsychiatric battery	6 months	No significant change in cognition over 6 months (HF-REF and HF-PEF) Minimal improvement in DSST in both groups (likely due to learned effect) Higher LVEF associated with lower DSST score
Huijts 2013 [30]	491 HF-REF 120 HF-PEF	Age >60 years and outpatients	77 (8)	Prospective	HF-REF: hospitalization within past year HF-PEF: NT- proBNP ≥400 pg/ml if pt <75 years or ≥800 pg/ml if pt ≥75 years	Co-morbid physical illness	HF-REF: LVEF <45%, NYHA II- IV, NT-proBNP >400 pg/ml 120 HF-PEF: LVEF ≥45%	AMT	18 months	HF-REF: 23% of HF pts showed decline of ≥1 point in AMT over 18 months HF-PEF: 24% of HF pts showed improvement of ≥1 point in AMT over 18 months

AMT, Abbreviated Mental Test; CAD, coronary artery disease; CHF, congestive heart failure; CI, cognitive impairment; CO, cardiac output; CV, cardiovascular; DRS, Disability Rating Scale; DSST, digit symbol substitution test; EF, ejection fraction; ETT, exercise tolerance test; HF, heart failure; HF-REF, heart failure-reduced ejection fraction; HF-PEF, heart failure-preserved ejection fraction; LVEF, left ventricular ejection fraction; MI, myocardial infarction; MMSE, Mini-Mental State Examination; NT-pro BNP, N-terminal prohormone brain natriuretic peptide; NYHA, New York Heart Association; pts, patients; SD, standard deviation.

distress may be the cause or effect of HF. Investigating reverse causation is challenging but possible; to avoid biases from early mortality, large datasets with sufficient prospective follow-up are required [44].

Association does not imply causation and we must be mindful that both HF and CI are diseases of older age with many shared pathologies. Recognising this, many HF studies have defined an age-related inclusion criterion. With all the caveats that come with the heterogeneity of the available data, it would seem that association of CI and HF is present at all ages (Table 1). Studies that have attempted more sophisticated adjustment for confounders illustrate the inherent difficulty in teasing out what is contributory to cognitive decline and what is association or epi-phenomenon. In general, HF patients tend to have poorer scores on cognitive tests when compared with a 'healthy' (no cardiac disease) control group [34], but this comparator is still potentially confounded by cardiovascular comorbidity in the HF group. Inclusion of a cohort with common vascular risk factors but no HF may allow determination of whether HF *per se* is associated with CI. Where attempts have been made to utilise this design, studies have been modest in size and results contradictory [16,19]. Some authors have described little difference between groups and others have described increased rates of CI in HF-REF groups, particularly in 'executive function' domains.

A direct 'dose response' relationship between severity of HF and severity of CI would strengthen arguments for a causal link. HF-REF can be quantified in terms of EF or symptom burden. For both measures there is an independent association with increasing prevalence of CI [6,8,13,16,17,20] and the poorest scores on cognitive testing are most often seen in those with the severest disease [23]. Interestingly, an association with CI is also seen in those with echocardiographic evidence of reduced EF but without symptoms of HF (that is, patients with asymptomatic left ventricular systolic dysfunction) [7].

Few studies have described cognitive function in patients with HF-PEF [10,22,23,29,30,45], but the pattern seems to be that CI is a substantial problem in all HF regardless of EF. Whether the prevalence or phenotype of cognitive change differs between HF-PEF and HF-REF is not clear as there have been few comparative studies. In keeping with much of the HF and cognition literature, where data are available, there is substantial potential for bias and results are contradictory. Some authors have described higher proportion of cognitive problems in HF-REF [29], while secondary analyses of clinical trials have suggested either an equal proportion of CI across the groups or an excess of CI in those with HF-PEF [30,45].

Heart failure and delirium

Two patterns of cognitive problems in HF are recognised: a chronic, progressive decline in cognitive ability and a more acute change in cognition often in association with decompensated disease. The acute delirium and HF relationship has not been well described. Delirium is a common sequela of decompensated HF; one study estimated that 17% of unscheduled HF hospitalisations had features of delirium [46]. Where delirium accompanies HF, outcomes are generally poor with increased mortality and length of stay [46]. However, delirium is a frequent complication of most medical emergencies in older adults and the delirium of decompensated HF may be no more or less frequent than the delirium that accompanies other medical conditions such as stroke or pneumonia.

Impact of cognitive impairment in heart failure

There is a literature describing the relationship between CI and 'classical cardiovascular trial' outcomes (Table 3). In general the presence of CI in HF is associated with poorer clinical outcomes, including longer hospital admissions, increased inpatient mortality and increased 1-year mortality [37]. However, as CI seems to be associated with more severe HF and with other medical comorbidities, we should not assume that poorer outcomes are directly attributable to the cognitive state. Several other important metrics have been described in HF cohorts and all seem to be worsened by the presence of CI, including functional ability, medication adherence and institutionalisation (Table 3). Cognitive decline tends not to occur in isolation and, as with other diseases of older age, the presence of impaired cognition in HF is often associated with concomitant functional decline and poor levels of self-care [32,37,38,40-43,47].

Potential pathophysiological explanations of cognitive impairment in heart failure

Historically, research describing the pathology of the dementias has been polarised, with vocal proponents for 'amyloid' and 'cerebral small vessel disease' aetiologies. Increasingly these processes are recognised as co-existing with complex biological interactions [48]. The same is likely true of the pathogenesis of CI in HF. Chronic cerebral hypoperfusion and occult cardioembolic disease are exemplar mechanistic explanations that have dominated the literature on cognition in HF. Both processes have face validity, have strong supporting scientific and observational data and yet have traditionally been studied in isolation [49]. For ease of understanding, we will keep this dichotomy and discuss the potential pathological mechanisms separately; however, it seems likely that both processes frequently coexist in patients with HF and may exert pathological synergy.

Table 3 Studies examining the relationship between cognitive impairment and outcomes in patients with heart failure

Study	Sample	Population	Median age in years (SD)	Study methodology	Inclusion criteria	Exclusion criteria	Measures	Results
Zuccalà 2003 [37]	1511 HF pts 11,790 controls	All geriatric or general medical admissions	79 (9)	Prospective	Not specified	Not specified	Hodkinson abbreviated mental test Length of hospital stay 1 year mortality	Mean length of hospital stay: pts with CI = 15 ± 10 days; pts without CI = 15 ± 9 days Inpatient mortality: pts with CI, 18%; pts without CI, 3% 1-year mortality: pts with CI, 27%; pts without CI, 15%
Karlsson 2005 [32]	146 CHF pts	Age >60 years and outpatients	76 (8)	Prospective	LVEF <45% NYHA II–IV	Co-morbid psychiatric, neurological or physical illness. Previous diagnosis of CI	HF self-care questionnaire MMSE	Self-care scores were significantly higher in those with MMSE >24 compared to those ≤24
Riegel 2007 [38]	29 CHF pts	Outpatients	64 (10)	Cross-sectional	LVSD on echo Clinical HF English speaking	Co-morbid psychiatric or physical illness. Previous diagnosis of CI	Self-care of HF index DSST Probed memory recall	CI was worse in the poor self-care group compared to the good and expert self-care groups but did not reach level of significance
Cameron 2009 [39]	50 CHF pts	Age >45 years and consecutive hospital admissions	73 (11)	Cross-sectional	Clinical CHF LVSD on echo English speaking	Co-morbid neurological illness. Previous diagnosis of CI	Self-care of HF index Cardiac depression scale MMSE	CI was not a predictor of self-care
Cameron 2010 [40]	93 CHF pts	Age >45 years and consecutive hospital admissions	73 (11)	Cross-sectional	Clinical CHF LVSD on echo English speaking	Co-morbid neurological illness. Previous diagnosis of CI	Self-care HF index MMSE MoCA	CI and self-care management were significantly associated (t = 2.7; P < 0.01)
Pulignano 2010 [41]	93 CHF pts	Consecutive outpatients	77 (6)	Cross-sectional	Not specified	Not specified	The European heart failure self-care behaviour scale MMSE	MMSE was negatively correlated with self-care behavioural scores (r = 0.58, P < 0.001)
Alosco 2013 [42]	110 CHF pts	Age >50 years and <85 years. Outpatients	70 (9)	Prospective	NYHA II–IV English speaking	Co-morbid psychiatric, neurological or physical illness. Previous diagnosis of CI	Lawton-Brody instrumental activities of daily living Modified MMSE (3MS)	Poorer performance on 3MS was associated with worse total activities of daily living performance
Harkness 2013 [43]	100 CHF pts	Age >55 years and outpatients	72 (10)	Cross-sectional	Confirmed HF using the Boston criteria LVEF ≤45% Change in symptoms on previous 3 months English speaking	Co-morbid psychiatric illness or previous diagnosis of CI	MoCA Self-care in HF index Geriatric Depression Scale	MoCA score of <26 was significantly associated with worse self-care management

Table 3 Studies examining the relationship between cognitive impairment and outcomes in patients with heart failure (Continued)

Alosco 2013 [42]	175 CHF pts	Age >50 years and <85 years. Outpatients	68 (10)	Cross- sectional	NYHA II-IV English speaking	Co-morbid psychiatric, neurological or physical illness. Previous diagnosis of CI	Lawton-Brody instrumental activities of daily living Executive function assessed by FAB and LNS	Poorer executive function was independently associated with poorer total activities of daily living performance
---------------------	----------------	---	---------	---------------------	------------------------------------	---	--	---

CHF, congestive heart failure; CI, cognitive impairment; DSST, digit symbol substitution test; FAB, frontal assessment battery; HF, heart failure; LNS, letter number sequencing; LVEF, left ventricular ejection fraction; LVSD, left ventricular systolic dysfunction; MMSE, Mini-Mental State Examination; MoCA, Montreal Cognitive Assessment Tool; NYHA, New York Heart Association; SD, standard deviation.

Although most of the postulated mechanisms we will discuss have been described in the context of HF-REF, issues of cerebral hypoperfusion, thrombotic disease and concomitant cardiovascular disease are also seen in HF-PEF [2] and it seems likely they will factor in the pathogenesis of any cognitive decline seen in this syndrome.

Confounding from other diseases

Co-existence of dementia and CI has been reported in a variety of cardiovascular disorders, including CAD, myocardial infarction and valvular heart disease. Midlife exposure to the common vascular risk factors of diabetes, hypertension and smoking is associated with later life cognitive decline [16]. This background is relevant to the study of patients with HF as many have a history of one or more of these co-morbidities. As discussed previously, dissecting the contribution of HF from concomitant vascular risk and disease is challenging but is essential for future studies that wish to describe the cognitive component of HF.

AF is a potential confounding condition worthy of separate discussion. The association of AF with cognitive decline is compelling [50]. Much of the CI associated with AF will be driven by cardioembolic stroke. However, cognitive decline is also seen in patients with AF and no history of clinical stroke, possibly representing occult embolic disease [50]. AF is common in HF and prevalence increases with severity of disease. Up to 50% of patients with 'end-stage' HF have AF [51]. Increasing use of ambulatory monitors is discovering substantial undetected paroxysmal AF and so these figures may be underestimates. While AF will be a factor in the pathogenesis of some HF-related CI, it is probably not the sole explanation. Where studies have controlled for the presence of AF in their HF patient population, there remains substantial prevalent CI [10,11,13,16,31].

Any discussion of cognition in cardiac disease has to consider the effect of invasive and instrumental procedures. The interventional toolkit available to cardiologists is increasingly sophisticated, with new indications emerging. Acute and chronic neurological deficits associated with cardiac surgery are well described [52] while interventions such as cardiac catheterisation and transcatheter aortic valve replacement have also been associated with post-procedure CI [53]. The mechanism of neurological insult associated with these procedures is likely a combination of reduced cerebral perfusion and embolic disease.

As well as 'physical' conditions, mood disorder may also represent an important confounder of association between HF and CI. Clinically important depression and anxiety are common in patients with HF. Depression is found in nearly 30% of HF patients and is associated with poor outcomes [54]. There is a complex interplay

between cognitive decline (particularly in the context of 'small vessel disease'), mood disorder and systemic vascular disease that is poorly understood but likely to be relevant to HF. Mood disorders are particularly important to detect as they can respond to intervention, making mood disorder in HF a potentially treatable form of cognitive decline.

Shared pathophysiology (systemic inflammation and amyloid)

Several recent studies have demonstrated the formation of tangle and plaque-like structures and fibrillar deposits (that is, the 'hallmark' lesions of Alzheimer's disease (AD) dementia) within the myocardium of patients with hypertrophic cardiomyopathy and idiopathic dilated cardiomyopathy [55]. Mis-folded proteins in the form of intermediate oligomers have also been described in cardiac tissue, with a distribution similar to that observed in the brain of patients with AD [55], raising the possibility of a common myocardial and cerebral pathology in a subset of patients with HF.

The systemic inflammatory state recognised in patients with HF may also contribute to CI [56]. It is postulated that inflammatory mediators influence cognition via diverse cytokine-mediated interactions between neurons and glial cells. *In vitro* and animal models support the inflammation and cognitive decline hypothesis and studies in humans with HF are emerging, although data are far from definitive at present [56].

Acute and chronic hypoperfusion

A mechanistic link between hypotension and CI, mediated via chronic cerebral hypoperfusion and loss of the normal autoregulation of cerebral perfusion pressures, has been postulated. Many diseases, including diabetes mellitus and depression, are associated with impaired reactivity of cerebrovascular perfusion autoregulatory systems and this state seems to confer a higher risk of cognitive decline [57]. HF patients often have systemic hypotension and in the context of disordered autoregulation this could lead to further insults to cerebral perfusion. Cerebral perfusion abnormalities have been demonstrated in HF patients, with reactivity more impaired in patients with greater severity of HF.

These hypoperfusion cognitive problems are not necessarily 'vascular' dementia. In animal models, reduced cerebral blood flow triggers a neurotoxic cascade that culminates in accumulation of amyloid and hyperphosphorylated tau proteins, the classical precursors of AD. If chronic hypoperfusion is causative, then improving cerebral blood flow should reduce cognitive decline. There is some evidence to support this view in patients with severe HF who have undergone cardiac transplant, pacemaker or cardiac resynchronisation therapy, and in

whom measures of cognition have stabilised or improved post-procedure [58].

Thrombosis and cerebral infarction

The potential importance of AF-related cardioembolism has been discussed. Cardioembolism is also seen in HF with sinus rhythm where ventricular function is the most important determinant of thrombus formation and potential embolic cerebral infarction [59] (Figure 2). Downregulation of thrombomodulin, structural changes in the cardiac chambers and potential blood stasis in the context of reduced myocardial contractility are associated with thrombus formation that may in turn lead to arterial events of clinical stroke or occult cerebral infarction [59]. This systemic prothrombotic phenotype increases risk of all thrombo-embolic diseases and HF is also associated with venous thromboembolism [60,61]. This is not surprising, as abnormalities in all three constituents of Virchow's Triad (abnormal blood constituents, abnormal vessel wall and abnormal blood flow) are present in HF. Neurohormonal activation seen in HF is associated with increased production of thrombogenic factors such as von Willebrand factor, thromboxane A2 and endothelin. The end result is a hypercoagulable state with increased serum levels of circulating fibrinogen, fibrinopeptide A and D-dimer (amongst others) resulting in platelet and thrombin activation and ultimately leading to plasma hyperviscosity and thrombosis [1]. A relationship between all these circulating markers of thrombosis and haemostasis and cognitive decline, particularly 'vascular dementia', has been described [62]. It

would seem intuitive that anticoagulation may prevent sequelae of thrombosis; however, studies of formal anticoagulation in HF with sinus rhythm have been equivocal. To date, no large study of anticoagulation in HF describing cognitive outcomes has been published.

Cognitive screening in heart failure services

Given the prevalence and potential impact of CI in HF, a case could be made for routine cognitive screening of HF patients. This is a controversial area with strongly held views on both sides. Recent observational data suggest that informal assessment of cognition by a cardiologist is insufficiently sensitive, with around three in four HF patients with important cognitive problems not recognised as such in routine consultations [63]. To date, routine screening for CI has not been incorporated into HF clinical guidelines; this may be due in part to the lack of a standardised screening technique that is feasible and acceptable for use in the cardiology outpatient setting. A recent systematic review of cognitive screening questionnaires utilised in HF studies concluded that the accuracy of traditional cognitive assessment measures is questionable in HF populations and appropriate thresholds/normative values need to be established [64]. In this regard we welcome ongoing work by the Cochrane Dementia and Cognitive Improvement Group to offer synthesis of test accuracy of cognitive assessments in various healthcare contexts [65].

Treatment implications of cognitive impairment in heart failure

There is an impressive evidence base to support pharmacological interventions in HF-REF. Historically HF trials have described clinical outcomes such as death, vascular events and hospitalisation with decompensated HF. There has been little focus on cognition or dementia as trial outcome or as a case mix adjuster. In fact for many of the trials that inform the HF evidence base, dementia or CI will have been an exclusion criterion. Where trialists have attempted to describe cognitive effects of HF treatment, results have been neutral [30].

Central to the treatment of HF is relatively complex multi-drug pharmacological treatment with attendant need for careful biochemical surveillance and self-monitoring. To achieve optimal outcomes requires strict adherence to prescribed evidence-based therapy [2]. Poor adherence is linked to an elevated risk of hospitalisation and death, whereas appropriate self-management may reduce these risks [2]. It seems intuitive that ensuring adherence and self-management would be especially challenging in the context of CI.

Interventions with angiotensin converting enzyme inhibitors (ACE-is), which have effects on the renin-




Figure 2 Magnetic resonance imaging of brain (diffusion weighted imaging sequences) in a patient with severe left ventricular systolic dysfunction and acute cognitive change.

The initial images were felt to represent a multi-infarct state, presumed cardioembolic and 'watershed' (hypoperfusion) infarction. Subsequent investigations revealed that the patient had 'shared' cardiac and cerebral pathology caused by a systemic and cerebral vasculitic process.

angiotensin-aldosterone system (RAAS), have been a mainstay of HF-REF therapy for decades. ACE is also important in neurotransmitter modulation and there are theoretical reasons to believe that ACE-is may have an effect on cognitive decline. Cognitive sub-studies of the Cardiovascular Health Study and the Italian Longitudinal Study on Ageing [66,67] both reported that subjects treated with ACE-is had equivalent rates of incident dementia compared with those treated with other antihypertensives. However, there were intriguing within-class differences in cognitive outcomes - for example, between centrally and non-centrally active agents and between differing drug potencies [67]. The other pillars of HF-REF therapy, beta-blockers and mineralocorticoid receptor antagonists, may also influence cognition. Although no studies specific to HF are available, there is hypertension literature suggesting theoretical cognitive effects of beta-blockade but inconclusive evidence that this is clinically important [68]. Cognitive effects of mineralocorticoid receptor antagonists have been demonstrated in animal models but human data are limited [69].

Novel approaches to pharmacological intervention in HF are being developed, with the natriuretic peptide system a key therapeutic target. These peptides possess differing degrees of haemodynamic, neurohormonal, renal and cardiac effects which may be favourable in the HF setting and may augment the effects of RAAS blockade. Preliminary studies using inhibitors of neprilysin (also known as neutral endopeptidase), an enzyme involved in the breakdown of endogenous natriuretic peptides, have yielded encouraging results [70]. Based on this experience a phase III trial comparing the angiotensin receptor neprilysin inhibitor molecule LCZ696 to the ACE-i enalapril was undertaken in chronic HF-REF (PARADIGM-HF). This trial was recently stopped for benefit of LCZ696 over enalapril [71]. However, cardiac optimism must be tempered by caution regarding potential non-cardiac, cognitive adverse effects. Mutations in the neprilysin gene have been associated with familial forms of AD and neprilysin-deficient mice show an AD phenotype [72].

In the light of non-definitive data, how should we treat a patient with HF and CI? Cognitive enhancing medication such as acetylcholinesterase inhibitors have recognised effects on the cardiac conduction system, occasionally causing bradycardia, sick sinus syndrome or other arrhythmias (including torsades de pointes) resulting from QT prolongation through excessive cholinergic stimulation. One recent study showed donepezil to be safe in patients without symptomatic heart disease and actually reduced levels of plasma brain natriuretic peptide in patients with subclinical HF [73].

Although there are no data to suggest cognitive benefits of standard HF therapy, there are equally no signals of harm. Given the beneficial effects of pharmacological therapy on mortality and hospitalisation, it would seem sensible to consider these evidence-based medical interventions for all HF patients, tailoring the intervention to suit the patient. A multidisciplinary approach with frequent review and medication titration seems to work well. Prescribers need to be alert to the potential effects of CI on concordance with sometimes complex drug regimens. Early use of compliance aids and involvement of family or carers may help in this regard. The goal of management of HF is to provide 'seamless care' in both the community and hospital to ensure the treatment of every patient is optimal. Despite the plethora of publications and guidelines, the data consistently show a lower uptake of evidence-based investigations and therapies in older patients with consequent higher rates of HF hospitalizations and mortality [43]. The current shift away from concentration on individual drug therapies to a focus on systems of care that allow effective treatment delivery is welcomed.

Conclusion

Recurrent themes in our synthesis of the literature regarding CI and HF are a lack of primary data, methodological limitations in available research, and conflicting results. To progress our understanding we recommend increasing use of cognitive assessment using standardised screening tools in all future HF studies. Although we found numerous studies assessing prevalence, there is a dearth of studies investigating the incidence of CI in HF. Once the incidence and prevalence of CI in HF are better defined we need to evaluate the consequences of CI in HF. Identifying underlying mechanisms for CI in HF may present targets for intervention, the 'holy grail' of cognitive research. A number of processes have been postulated, and we now need confirmatory studies using new developments in neuroimaging and biomarkers in representative populations of HF patients. All of this will require a multidisciplinary approach between HF and dementia research teams. Such collaborative activity is urgently needed given the projected increases in both CI and HF.

Note: This article is part of a series on *The impact of acute and chronic medical disorders on accelerated cognitive decline*, edited by Carol Brayne and Daniel Davis. Other articles in this series can be found at <http://alres.com/series/medicaldisorders>.

Abbreviations

ACE-(i): Angiotensin converting enzyme-(inhibitor); AD: Alzheimer's disease; AF: Atrial fibrillation; CAD: Coronary artery disease; CI: Cognitive impairment; EF: Ejection fraction; HF: Heart failure; HF-PEF: Heart failure-preserved ejection fraction; HF-REF: Heart failure-reduced ejection fraction; RAAS: Renin-angiotensin-aldosterone system.

Competing interests

JC has no competing interests. JJVM has no competing interests. TJQ has received modest honoraria, research funding and travel support from: Astra-Zeneca; Bayer; Boehringer Ingelheim; Bristol Meyers Squibb; Merck; Pfizer. He holds grants relating to cognitive assessment from British Geriatric Society; Chest Heart and Stroke Scotland; Chief Scientists Office Scotland; The Stroke Association.

Author details

¹British Heart Foundation Glasgow, Institute of Cardiovascular and Medical Sciences, University of Glasgow, University Avenue, Glasgow G12 8TA, UK.

²Department of Academic Geriatric Medicine, Institute of Cardiovascular and Medical Sciences, New Lister Building, Glasgow Royal Infirmary, Glasgow UK G4 0SF, UK.

Published online: 27 February 2015

References

- Braunwald E. Heart Disease: A Textbook of Cardiovascular Medicine, vol. 2. Philadelphia, US: WB Saunders; 2005.
- McMurray JJ, Adamopoulos S, Anker SD, Auricchio A, Böhm M, Dickstein K, et al. ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure 2012: The Task Force for the Diagnosis and Treatment of Acute and Chronic Heart Failure 2012 of the European Society of Cardiology. Developed in collaboration with the Heart Failure Association (HFA) of the ESC. *Eur Heart J*. 2012;33:1787–847.
- Jhund PS, Macintyre K, Simpson CR, Lewsey JD, Stewart S, Redpath A, et al. Long-term trends in first hospitalization for heart failure and subsequent survival between 1986 and 2003: a population study of 5.1 million people. *Circulation*. 2009;119:515–23.
- Stewart S, MacIntyre K, Capewell S, McMurray JJ. Heart failure and the aging population: an increasing burden in the 21st century? *Heart*. 2003;89:49–53.
- Ho KKL, Pinsky JL, Kannel WB, Levy D, Pitt B. The epidemiology of heart failure: The Framingham Study. *J Am Coll Cardiol*. 1993;Suppl 4:A6–A13.
- Zuccalà G, Cattel C, Manes-Gravina E, Di Niro MG, Cocchi A, Bernabei R. Left ventricular dysfunction: a clue to cognitive impairment in older patients with heart failure. *J Neurol Neurosurg Psychiatry*. 1997;63:509–12.
- Callegari S, Majani G, Giardini A, Pierobon A, Opasich C, Cobelli F, et al. Relationship between cognitive impairment and clinical status in chronic heart failure patients. *Monaldi Arch Chest Dis*. 2002;58:19–25.
- Trojano L, Antonelli Inc, Acanfora D, Picone C, Mecocci P, Rengo F. Cognitive impairment: a key feature of congestive heart failure in the elderly. *J Neurol*. 2003;250:1456–63.
- Zuccalà G, Marzetti E, Cesari M, Lo Monaco MR, Antonica L, Cocchi A, et al. Correlates of cognitive impairment among patients with heart failure: results of a multicenter survey. *Am J Med*. 2005;118:496–502.
- Feola M, Rosso GL, Peano M, Agostini M, Aspromonte N, Carena G, et al. Correlation between cognitive impairment and prognostic parameters in patients with congestive heart failure. *Arch Med Res*. 2007;38:234–9.
- Debette S, Bauters C, Leys D, Lamblin N, Pasquier F, de Groote P. Prevalence and determinants of cognitive impairment in chronic heart failure patients. *Congest Heart Fail*. 2007;13:205–8.
- Dodson J. Cognitive impairment in older adults with heart failure: prevalence, documentation, and impact on outcomes. *Am J Med*. 2013;126:12026.
- Schmidt R, Fazekas F, Offenbacher H, Dusleag J, Lechner H. Brain magnetic resonance imaging and neuropsychologic evaluation of patients with idiopathic dilated cardiomyopathy. *Stroke*. 1991;22:195–9.
- Grubb NR, Simpson C, Fox KA. Memory function in patients with stable, moderate to severe cardiac failure. *Am Heart J*. 2000;140:E1–5.
- Riegel B, Bennett JA, Davis A, Carlson B, Montague J, Robin H, et al. Cognitive impairment in heart failure: issues of measurement and etiology. *Am J Crit Care*. 2002;11:520–8.
- Vogels RL, Oosterman JM, van Harten B, Scheltens P, van der Flier WM, Schroeder-Tanka JM, et al. Profile of cognitive impairment in chronic heart failure. *J Am Geriatr Soc*. 2007;55:1764–70.
- Hoth KF, Poppas A, Moser DJ, Paul RH, Cohen RA. Cardiac dysfunction and cognition in older adults with heart failure. *Cogn Behav Neurol*. 2008;21:65–72.
- Beer C, Ebenezer E, Fenner S, Lautenschlager NT, Arnolda L, Flicker L, et al. Contributors to cognitive impairment in congestive heart failure: a pilot case-control study. *Intern Med J*. 2009;39:600–5.
- Stanek KM, Gunstad J, Paul RH, Poppas A, Jefferson AL, Sweet LH, et al. Longitudinal cognitive performance in older adults with cardiovascular disease: evidence for improvement in heart failure. *J Cardiovasc Nurs*. 2009;24:192–7.
- Sauve MJ, Lewis WR, Blankenbiller M, Rickabaugh B, Pressler SJ. Cognitive impairments in chronic heart failure: a case controlled study. *J Card Fail*. 2009;15:1–10.
- Pressler S. Cognitive deficits and health-related quality of life in chronic heart failure. *J Cardiovasc Nurs*. 2010;25:189–98.
- Bauer L, Pozehl B, Hertzog M, Johnson J, Zimmerman L, Filipi M. A brief neuropsychological battery for use in the chronic heart failure population. *Eur J Cardiovasc Nurs*. 2012;11:223–30.
- Festa JR, Jia X, Cheung K, Marchidann A, Schmidt M, Shapiro PA, et al. Association of low ejection fraction with impaired verbal memory in older patients with heart failure. *Arch Neurol*. 2011;68:1021–6.
- Steinberg G, Lossnitzer N, Schellberg D, Mueller-Tasch T, Krueger C, Haass M, et al. Peak oxygen uptake and left ventricular ejection fraction, but not depressive symptoms, are associated with cognitive impairment in patients with chronic heart failure. *Int J Gen Med*. 2011;4:879–87.
- Jefferson AL, Himali JJ, Au R, Seshadri S, Decarli C, O'Donnell CJ, et al. Relation of left ventricular ejection fraction to cognitive aging (from the Framingham Heart Study). *Am J Cardiol*. 2011;108:1346–51.
- Miller LA, Spitznagel MB, Alosco ML, Cohen RA, Raz N, Sweet LH, et al. Cognitive profiles in heart failure: a cluster analytic approach. *J Clin Exp Neuropsychol*. 2012;34:509–20.
- Almeida OP, Garrido GJ, Beer C, Lautenschlager NT, Arnolda L, Flicker L. Cognitive and brain changes associated with ischaemic heart disease and heart failure. *Eur Heart J*. 2012;33:1769–76.
- Hawkins LA, Kilian S, Firek A, Kashner TM, Firek CJ, Silvet H. Cognitive impairment and medication adherence in outpatients with heart failure. *Heart Lung*. 2012;41:572–82.
- Bratzke-Bauer L. Neuropsychological patterns differ by type of left ventricular dysfunction in heart failure. *Arch Clin Neuropsychol*. 2013;28:114–24.
- Huijts M, van Oostenbrugge RJ, Duits A, Burkard T, Muzzarelli S, Maeder MT, et al. Cognitive impairment in heart failure: results from the Trial of Intensified versus standard Medical therapy in Elderly patients with Congestive Heart Failure (TIME-CHF) randomized trial. *Eur J Heart Fail*. 2013;15:699–707.
- Kindermann I, Fischer D, Karbach J, Link A, Walenta K, Barth C, et al. Cognitive function in patients with decompensated heart failure: the Cognitive Impairment in Heart Failure (CogImpair-HF) study. *Eur J Heart Fail*. 2012;14:404–13.
- Karlsson MR, Edner M, Henriksson P, Mejhert M, Persson H, Grut M. Billing, A nurse-based management program in heart failure patients affects females and persons with cognitive dysfunction most. *Patient Educ Couns*. 2005;58:146–53.
- Tanne D. Cognitive functions in severe congestive heart failure before and after an exercise training program. *Int J Cardiol*. 2005;103:145–9.
- Almeida O. Brain and mood changes over 2 years in healthy controls and adults with heart failure and ischaemic heart disease. *Eur J Heart Failure*. 2013;15:850–8.
- Hjelm C, Dahl A, Brostrom A, Martensson J, Johansson B, Stromberg A. The influence of heart failure on longitudinal changes in cognition among individuals 80 years of age and older. *J Clin Nurs*. 2012;21:994–1003.
- Riegel B, Lee CS, Glaser D, Moelter ST. Patterns of change in cognitive function over six months in adults with chronic heart failure. *Cardiol Res Pract*. 2012;2012:631075.
- Zuccalà G, Pedone C, Cesari M, Onder G, Pahor M, Marzetti E, et al. The effects of cognitive impairment on mortality among hospitalized patients with heart failure. *Am J Med*. 2003;115:97–103.
- Riegel B, Vaughan Dickson V, Goldberg LR, Deatrick JA. Factors associated with the development of expertise in heart failure self-care. *J Nurs Res*. 2007;56:235–43.

39. Cameron J. Testing a model of patient characteristics, psychological status, and cognitive function as predictors of self-care in persons with chronic heart failure. *J Acute Crit Care*. 2009;38:410–8.
40. Cameron J, Worrall-Carter L, Page K, Riegel B, Lo SK, Stewart S. Does cognitive impairment predict poor self-care in patients with heart failure. *Eur J Heart Fail*. 2010;12:508–15.
41. Pulignano G, Del Sindaco D, Minardi G, Tarantini L, Cioffi G, Bernardi L, et al. Translation and validation of the Italian version of the European Heart Failure Self-Care Behaviour Scale. *J Cardiovasc Med*. 2010;11:493–8.
42. Alosco ML, Spitznagel MB, Cohen R, Sweet LH, Colbert LH, Josephson R, et al. Reduced cognitive function predicts functional decline in patients with heart failure over 12 months. *Eur J Cardiovasc Nurs*. 2013;13:304–10.
43. Harkness K, Heckman GA, Akhtar-Danesh N, Demers C, Gunn E, McKelvie RS. Cognitive function and self-care management in older patients with heart failure. *Eur J Cardiovasc Nurs*. 2013;13:277–84.
44. Russ TC, Stamatakis E, Hamer M, Starr JM, Kivimaki M, Batty GD. Association between psychological distress and mortality: individual participant pooled analysis of 10 prospective cohort studies. *BMJ*. 2012;345:e4933.
45. van den Hurk K, Reijmer YD, van den Berg E, Alsema M, Nijpels G, Kostense PJ, et al. Heart failure and cognitive function in the general population: the Hoorn Study. *Eur J Heart Fail*. 2011;13:1362–9.
46. Uthamalingham S. Usefulness of acute delirium as a predictor of adverse outcomes in patients >65 years of age with acute decompensated heart failure. *Am J Cardiol*. 2011;108:402–8.
47. Alosco ML, Spitznagel MB, Raz N, Cohen R, Sweet LH, Colbert LH, et al. Executive dysfunction is independently associated with reduced functional independence in heart failure. *J Clin Nurs*. 2014;23:829–36.
48. Jicha GA, Parisi JE, Dickson DW, Johnson K, Cha R, Ivnik RJ, et al. Neuropathologic outcome of mild cognitive impairment following progression to clinical dementia. *Arch Neurol*. 2006;63:674–81.
49. Georgiadis D, Sievert M, Cencetti S, Uhlmann F, Krivokuca M, Zierz S, et al. Cerebrovascular reactivity is impaired in patients with cardiac failure. *Eur Heart J*. 2000;21:407–13.
50. Kalantarian S, Stern TA, Mansour M, Ruskin JN. Cognitive impairment associated with atrial fibrillation: a meta-analysis. *Ann Intern Med*. 2013;158:338–46.
51. Neuberger HR, Mewis C, van Veldhuisen DJ, Schotten U, van Gelder IC, Allessie MA, et al. Management of atrial fibrillation in patients with heart failure. *Eur Heart J*. 2007;28:2568–77.
52. Caplan LR. Translating what is known about neurological complications of coronary artery bypass graft surgery into action. *Arch Neurol*. 2009;66:1062–4.
53. Kahlert P, Knipp SC, Schlamann M, Thielmann M, Al-Rashid F, Weber M, et al. Silent and apparent cerebral ischemia after percutaneous transfemoral aortic valve implantation: a diffusion-weighted magnetic resonance imaging study. *Circulation*. 2010;121:870–8.
54. Diez-Quevedo C, Lupón J, González B, Urrutia A, Cano L, Cabanes R, et al. Depression, antidepressants, and long-term mortality in heart failure. *Int J Cardiol*. 2013;167:1217–25.
55. Willis MS, Patterson C. Proteotoxicity and cardiac dysfunction - Alzheimer's disease of the heart? *N Engl J Med*. 2013;368:455–64.
56. Athilingam P, Moynihan J, Chen L, D'Aouat R, Groer M, Kip K. Elevated Levels of Interleukin 6 and C-reactive protein associated with cognitive impairment in heart failure. *Congest Heart Fail*. 2013;19:92–8.
57. Tchistiakova E, Anderson ND, Greenwood CE, MacIntosh BJ. Combined effects of type 2 diabetes and hypertension associated with cortical thinning and impaired cerebrovascular reactivity relative to hypertension alone in older adults. *Neuroimage Clin*. 2014;5:36–41.
58. Conti JB, Sears SF. Cardiac resynchronization therapy: can we make our heart failure patients smarter? *Trans Am Clin Climatol Assoc*. 2007;118:153–64.
59. Kalaria VG, Passannante MR, Shah T, Modi K, Weisse AB. Effect of mitral regurgitation on left ventricular thrombus formation in dilated cardiomyopathy. *Am Heart J*. 1998;135:215–20.
60. Al-Khadra AS, Salem DN, Rand WM, Udelson JE, Smith JJ, Konstam MA. Antiplatelet agents and survival: a cohort analysis from the Studies of Left Ventricular Dysfunction (SOLVD) trial. *J Am Coll Cardiol*. 1998;31:419–25.
61. Freudenberger RS, Hellkamp AS, Halperin JL, Poole J, Anderson J, Johnson G, et al. Risk of thromboembolism in heart failure: an analysis from the Sudden Cardiac Death in Heart Failure Trial (SCD-HeFT). *Circulation*. 2007;115:2637–41.
62. Quinn TJ, Gallacher J, Deary IJ, Lowe GD, Fenton C, Stott DJ. Association between circulating hemostatic measures and dementia or cognitive impairment: systematic review and meta-analyses. *J Thromb Haemost*. 2011;9:1475–82.
63. Hanon O, Vidal JS, de Groote P, Galinier M, Isnard R, Logeart D, et al. Prevalence of memory disorders in ambulatory patients aged >=70 years with chronic heart failure (from the EFICARE Study). *Am J Cardiol*. 2014;113:1205–10.
64. Davis KK, Allen JK. Identifying cognitive impairment in heart failure: a review of screening measures. *Heart Lung*. 2013;42:92–7.
65. Quinn TJ, Fearon P, Noel-Storr AH, Young C, McShane R, Stott DJ. Informant Questionnaire on Cognitive Decline in the Elderly (IQCODE) for the diagnosis of dementia within community dwelling populations. *Cochrane Database Syst Rev*. 2014;4:CD010079.
66. Sink KM, Leng X, Williamson J, Kritchevsky SB, Yaffe K, Kuller L, et al. Angiotensin-converting enzyme inhibitors and cognitive decline in older adults with hypertension: results from the Cardiovascular Health Study. *Arch Intern Med*. 2009;169:1195–202.
67. Solfrizzi V, Scafato E, Frisardi V, Seripa D, Logroscino G, Kehoe PG, et al. Angiotensin-converting enzyme inhibitors and incidence of mild cognitive impairment. The Italian Longitudinal Study on Aging. *Age (Dordr)*. 2013;35:441–53.
68. Levi MN, Macquin-Mavier I, Tropeano AI, Bachoud-Levi AC, Maison P. Antihypertensive classes, cognitive decline and incidence of dementia: a network meta-analysis. *J Hypertens*. 2013;31:1073–82.
69. Korte SM, Korte-Bouws GA, Koob GF, De Kloet ER, Bohus B. Mineralocorticoid and glucocorticoid receptor antagonists in animal models of anxiety. *Pharmacol Biochem Behav*. 1996;54:261–7.
70. Packer M, Califf RM, Konstam MA, Krum H, McMurray JJ, Rouleau JL, et al. Comparison of omapatrilat and enalapril in patients with chronic heart failure: the Omapatrilat Versus Enalapril Randomized Trial of Utility in Reducing Events (OVERTURE). *Circulation*. 2002;106:920–6.
71. McMurray JJV, Packer M, Desai AS, Gong J, Lefkowitz MP, Rizkala AR, et al. Angiotensin-neprilysin inhibition versus enalapril in heart failure. *N Engl J Med*. 2014;371:993–1004.
72. Madani R, Poirier R, Wolfer DP, Welzl H, Groscurth P, Lipp HP, et al. Lack of neprilysin suffices to generate murine amyloid-like deposits in the brain and behavioral deficit in vivo. *J Neurosci Res*. 2006;84:1871–8.
73. Kubo T, Sato T, Noguchi T, Kitaoka H, Yamasaki F, Kamimura N, et al. Influences of donepezil on cardiovascular system - possible therapeutic benefits for heart failure - donepezil cardiac test registry (DOCTER) study. *J Cardiovasc Pharmacol*. 2012;60:310–4.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

