

# HEAT-SHOCK PROTEINS, MOLECULAR CHAPERONES, AND THE STRESS RESPONSE: Evolutionary and Ecological Physiology

*Martin E. Feder*

Department of Organismal Biology and Anatomy and Committee on Evolutionary Biology, University of Chicago, 1027 East 57th Street, Chicago, Illinois 60637;  
e-mail: m-feder@uchicago.edu

*Gretchen E. Hofmann*

Department of Biology, University of New Mexico, Albuquerque, NM 87131;  
e-mail: ghofmann@unm.edu

KEY WORDS: hsp, temperature, protein denaturation and folding, inducible tolerance, environmental gradients

---

## ABSTRACT

Molecular chaperones, including the heat-shock proteins (Hsps), are a ubiquitous feature of cells in which these proteins cope with stress-induced denaturation of other proteins. Hsps have received the most attention in model organisms undergoing experimental stress in the laboratory, and the function of Hsps at the molecular and cellular level is becoming well understood in this context. A complementary focus is now emerging on the Hsps of both model and nonmodel organisms undergoing stress in nature, on the roles of Hsps in the stress physiology of whole multicellular eukaryotes and the tissues and organs they comprise, and on the ecological and evolutionary correlates of variation in Hsps and the genes that encode them. This focus discloses that (*a*) expression of Hsps can occur in nature, (*b*) all species have *hsp* genes but they vary in the patterns of their expression, (*c*) Hsp expression can be correlated with resistance to stress, and (*d*) species' thresholds for Hsp expression are correlated with levels of stress that they naturally undergo. These conclusions are now well established and may require little additional confirmation; many significant questions remain

unanswered concerning both the mechanisms of Hsp-mediated stress tolerance at the organismal level and the evolutionary mechanisms that have diversified the *hsp* genes.

---

## INTRODUCTION

Although heat-shock proteins (Hsps) first achieved notoriety as gene products whose expression is induced by heat and other stresses (1, 2), discoveries of the past decade have shifted the focus of research to understanding the roles of Hsps as molecular chaperones (3–5). As a result, Hsps, their close relatives, their molecular partners, and many newly discovered proteins are now known to play diverse roles, even in unstressed cells, in successful folding, assembly, intracellular localization, secretion, regulation, and degradation of other proteins (6); failure of these activities is thought to underlie numerous and important human diseases (7). Nonetheless, many of the questions of the past either remain unanswered, awaiting the development of appropriate experimental tools, or can now be revisited with new insights gleaned from the emerging understanding of molecular chaperones. This review seeks to facilitate the examination or re-examination of Hsps as responses to natural stress in diverse organisms inhabiting environments outside the laboratory, the function of Hsps in tolerance of natural stresses, and ecological and evolutionary variation in the heat-shock system. The review sequentially considers (a) the principal implications of laboratory-based studies for ecological and evolutionary research on Hsps, (b) expression of Hsps in nature, (c) covariation of Hsp expression with environmental and biological gradients of stress intensity, (d) the consequences of Hsp expression for fitness, and (e) evolutionary variation in Hsps and the genes that encode them. The primary objective of this review is to redirect the focus of evolutionary and ecological research on Hsps beyond the conclusions that are now well-established and onto the many important questions that remain unanswered.

## STATE OF THE LITERATURE

### *Established Conclusions*

The relevant literature on Hsps and molecular chaperones is huge, now comprising more than 12,000 references. Even a review of the relevant reviews is difficult. For this reason, we begin by describing several well established conclusions and cite a few of the many excellent recent reviews at diverse levels of sophistication (4–6, 8–11).

The genes encoding Hsps (*hsps*) are highly conserved and occur in every species in which they have been sought. Many of these genes and their products can be assigned to families on the basis of sequence homology and typical molecular weight (6): *hsp110*, *hsp100*, *hsp90*, *hsp70*, *hsp60*, *hsp40*, *hsp10*, and small *hsp* families. Gething (6) recognizes 7 additional families and 12 genes/proteins for which families have not yet been described. In eukaryotes, many families comprise multiple members that differ in inducibility, intracellular localization, and function.

Hsps function as molecular chaperones; i.e. they interact with other proteins and, in so doing, minimize the probability that these other proteins will interact inappropriately with one another. Hsps recognize and bind to other proteins when these other proteins are in non-native conformations, whether due to protein-denaturing stress or because the peptides they comprise have not yet been fully synthesized, folded, assembled, or localized to an appropriate cellular compartment. Binding and/or release of these other proteins is often regulated by association with and/or hydrolysis of nucleotides. Typically, Hsps function as oligomers, if not as complexes of several different chaperones, co-chaperones, and/or nucleotide exchange factors. Interaction with chaperones is variously responsible for (a) maintaining Hsps' partner proteins in a folding-competent, folded, or unfolded state; (b) organellar localization, import, and/or export; (c) minimizing the aggregation of non-native proteins; and (d) targeting non-native or aggregated proteins for degradation and removal from the cell. Presumably, the last two functions are most important in coping with environmental stress.

Not all Hsps are stress-inducible, but those that are respond to a variety of stresses, including extremes of temperature, cellular energy depletion, and extreme concentrations of ions, other osmolytes, gases, and various toxic substances. Activation of various intracellular signaling pathways results in Hsp expression. All known stresses, if sufficiently intense, induce Hsp expression. Accordingly, Hsps are equally well termed stress proteins, and their expression is termed the stress response. A common aspect of these inducing stresses is that they result in proteins having non-native conformations (12), which is consistent with the function of Hsps as molecular chaperones.

### *Implications of the Published Literature for Ecological and Evolutionary Studies of Hsps*

Space limitations necessitate that we choose among numerous equally valuable references in preparing this review. To present both the breadth and depth of research relevant to the evolutionary and ecological physiology of the heat-shock response, we have compiled a near-comprehensive bibliography of

that literature, which is available electronically (13) on the World Wide Web in the Supplemental Materials Section of the main Annual Reviews site (<http://www.AnnualReviews.org>).

A first implication of this massive literature is that the Hsp field has long ago concluded its exploratory phase. Showing that an as-yet-unexamined species expresses Hsps in response to heat or other stresses no longer has any particular novelty.

Second, much of the work on Hsps outside the laboratory or in nonmodel organisms was undertaken before the molecular diversity of Hsps and their function as molecular chaperones was obvious. In the interim, the experimental tools for examining Hsps and the standards for such examinations have both advanced considerably. As a result, much of the earlier work on evolutionary and ecological physiology of Hsps regrettably either does not withstand current scrutiny or contributes little to issues of current interest. Several issues are obvious:

1. Many of the apparently singular Hsps of previous years, often detected by one-dimensional electrophoresis and autoradiography, are now known to represent entire families of Hsps, often with (a) discrete distributions within the cell (e.g. cytoplasmic-nuclear, mitochondrial, chloroplast, or endoplasmic reticulum), (b) different degrees of inducibility (constitutively expressed, constitutively expressed but increasing during or after stress, exclusively inducible), (c) differing kinetics of induction and removal from the cell, and (d) differing tissue specificity. Representing this diversity as a single Hsp or two Hsps ("constitutive" and "inducible") through use of nonspecific probes or lysates of whole organisms and organs can obscure phenomena of great significance (e.g. compare Refs. 14 and 15 with 16). This problem is sometimes remediable only with great difficulty. Often, highly specific probes are available only for standard model organisms, particularly at the level of proteins, and great care must be taken in applying these probes to non-standard organisms (17).
2. Inducible stress tolerance is increasingly understood to result from numerous molecular mechanisms, of which Hsps are collectively only one. Other mechanisms include synthesis of osmotic stress protectants such as polyols and trehalose, modifications of the saturation of cell membrane lipids (homeoviscous adaptation), compensatory expression of isozymes or allozymes of significant enzymes, metabolic arrest, radical scavengers (superoxide dismutase, glutathione system, cytochrome P450), and so on. Accordingly, the unambiguous attribution of stress tolerance to Hsps in general or to any specific Hsp requires more than correlative evidence (18–21). Increasingly, proof resulting from genetic or direct experimental manipulation is

becoming the standard for establishing the functional or evolutionary significance of Hsps. Again, this rising standard is often met only with great difficulty in ecological and evolutionary physiological studies, for many of the techniques for genetic and experimental manipulation are not readily applicable to the more ecologically and evolutionarily interesting species.

## HSP-INDUCING STRESS IN NATURE AND NATURAL INDUCTION OF HSPS

Depending on their geographic locale, organisms in nature risk exposure to temperatures ranging from  $-100^{\circ}$  to more than  $100^{\circ}\text{C}$ , and comparable extremes of chemical and gas concentration, food and water availability, hydrostatic pressure, radiation, and toxic substances of human origin. Seemingly, Hsp expression should be a common occurrence in nature. In reality, however, movement and/or other behaviors may often enable organisms to avoid Hsp-inducing stress in nature by exploiting equable microhabitats in otherwise stressful environments (22–24). Also, biochemical specializations other than Hsps may stabilize many organisms (or particular stages of their life cycles) so that environmental extremes are not particularly stressful.

Even equable environments can contain Hsp-inducing microhabitats, and even mild stresses can induce Hsps when multiple stresses act in combination. For these reasons, we can assume neither the presence nor the absence of Hsp expression in nature; for that matter, we still do not know in any comprehensive sense whether wild organisms routinely, occasionally, or seldom express inducible Hsps. A growing body of evidence, however, establishes that at least in some circumstances and taxa, Hsp induction is not solely a laboratory phenomenon.

One caution in evaluating the subsequent account is that organisms in nature seldom undergo only one stress at a time. For example, an insect larva undergoing natural heat stress in a rotting fruit may simultaneously experience intense ultraviolet radiation, desiccation, and diverse alcohols and aldehydes, among other stresses. This situation differs from that in most laboratory experimentation, which involves one or a few stresses and makes attribution of Hsp expression to a particular stress in nature more complicated.

### *Aquatic Temperature Stress*

Due to the physical characteristics of water, the aquatic environment can be extremely stressful to its inhabitants. In general, the high specific heat and thermal conductivity of water ensure that the majority of aquatic organisms will have body temperatures equivalent to that of their surroundings. Furthermore, the relative thermal homogeneity of aquatic environments can frustrate behavioral

avoidance of thermal extremes. Some aquatic ectotherms nonetheless inhabit thermally equable habitats or waters with enough thermal diversity to enable behavioral thermoregulation; our focus is on those species that do not.

In the aquatic environment, habitual exposure to Hsp-inducing thermal stress may be most common in sessile organisms that occur in shallow, stagnant water (e.g. ponds, tidepools, swamps, tidal flats) or in the intertidal zone. Corals, for example, routinely undergo thermal stress that results in bleaching, during which the corals' endosymbionts die. Even modest increases in water temperature of 1-2°C can bleach corals; these temperatures also induce Hsp expression in several species (25, 26). Marine intertidal invertebrates undergo even larger increases in body temperatures during tidal emersion (27-30). For example, during aerial exposure intertidal mussels' body temperatures exceed seawater temperatures by more than 20°C (31), resulting in Hsp expression (29). A similar phenomenon occurs in encysted brine shrimp (*Artemia*) embryos (32, 33). Even relatively mobile aquatic ectotherms such as fish may undergo heat shock in nature (34, 35). For example, gobiid fishes of the genus *Gillichthys* can become trapped in shallow water, which is heated by the sun. Summer-acclimatized fish have higher levels of Hsp90 in brain tissue than do winter-acclimatized fish (34). In addition, the threshold Hsp induction temperature for one species, *G. mirabilis*, is significantly higher in summer than in winter. These data suggest that seasonal variation in water temperature can alter the heat-shock response. More exotic venues for aquatic thermal stress include thermal effluents of power plants, hydrothermal vents, and thermal hot springs, in which temperatures can exceed 100°C (see *Hsps of Archaea*).

### *Terrestrial Temperature Stress*

Unlike aquatic environments, terrestrial environments often offer diverse heat sources and sinks and retreats that organisms can exploit to avoid thermal stress. Thus, natural thermal stress and accompanying Hsp expression in terrestrial environments typically involve limitations in mitigating thermal extremes by movement and conflicts between thermoregulation and other needs. Salamanders, for example, which ordinarily maintain cool temperatures in nature, can inadvertently retreat beneath small sunlit rocks that become warm enough to induce *hsp70* mRNA expression (36); by abandoning these rocks to find cooler retreats they may risk immediate desiccation or even warmer temperatures.

The least equivocal case for routine exposure to Hsp-inducing temperature stress is for plants, which cannot change location except as seeds or pollen and can be limited in their ability to adjust heat exchange with the environment (37). Thus, plants in nature can become extremely hot (38). By inference, the entire range of plant heat-shock responses (8, 39) should manifest themselves in nature. Indeed, a small number of case studies document natural Hsp expression

(40–45), which can be greatest at times of day or in regions of an individual plant at which temperatures are highest (46). Plant species can differ dramatically, however, in both the magnitude and diversity of the particular Hsps that are expressed during days with especially warm weather (41). Plants should also be prone to natural cold stress (47), which ought to induce expression of Hsps (48–51).

Not surprisingly, therefore, many of the cases of natural thermal stress in animals on land involve animals that live inside or on plants (e.g. 52). *Drosophila* larvae and pupae encounter temperatures exceeding 40°C if the necrotic fruit they infest is in the sun, and express Hsp70 in response (53, 54). Presumably, other animals that cannot escape or offset intense solar heat loads will also express Hsps in nature; this hypothesis awaits systematic study. A unique case concerns desert ants, *Cataglyphis*, which voluntarily undergo body temperatures of >50°C, presumably to escape predators (55). The concentration of Hsp70 family members increases in this species before it naturally encounters high temperatures, as if in anticipation (56).

Terrestrial vertebrates are often especially effective in escaping heat stress, but both they and invertebrates are occasionally hyperthermic during intense physical activity, fever, or to conserve water. In birds and mammals, such hyperthermia activates HSF (the heat-shock transcription factor) and increases the level of Hsps (Hsp cognates, constitutively-expressed Hsps) and Hsps (57–59). Natural hypothermia of animals can be far more conspicuous than natural heat stress, involving diapause, overwintering in exposed sites, hibernation, and sometimes outright freezing. Diverse insects express Hsps in response to cold shock or during overwintering in diapause, although the identity of these Hsps, their tissue specificity, and their developmental regulation vary greatly (60–64). Some euthermic rodents express 70-kDa Hsps in response to cold ambient temperatures, possibly in tandem with nonshivering thermogenesis (65), and ground squirrels (*Spermophilus*) increase Hsp70 family members and ubiquitin-protein conjugates during hibernation (66).

In summary, laboratory studies of the heat-shock response often have proceeded far in advance of fieldwork that establishes an ecological context for their interpretation. Documentation of both natural thermal stress and Hsp expression in nature can provide this context, and a small but growing number of field studies demonstrate that such documentation is feasible.

### *Inducing Stresses Other than Temperature*

Virtually every nonthermal stress can induce Hsps (10, 67). Rarely, however, are these nonthermal stresses ecologically relevant; the literature in this area typically focuses on chemical stressors, and the corresponding data are essentially pharmacological. Even when the stress in question is ecologically

relevant, few studies of multicellular eukaryotes examine it in the field or in intact tissues and organisms. Some exceptional work, however, concerns plants and brine-shrimp (*Artemia*). The resurrection plant, a desert species, expresses Hsps in vegetative tissues during water stress; this expression is thought to contribute to desiccation tolerance (68). Similarly, rice seedlings express two proteins in the Hsp90 family upon exposure to water stress and elevated salinity (69). Embryos of the brine shrimp, one of the most hypoxia-tolerant metazoans, contain large quantities of p26, a molecular chaperone hypothesized to stabilize proteins during long bouts of anaerobic dormancy (see *Development*). Clearly, additional evolutionary physiological research in this area is sorely needed.

A recurrent theme is that thermal stress and these alternative stressors often result in different patterns of Hsp expression, indicating a diversity of regulatory mechanisms. Examples include variation in the expression of Hsp70 and ubiquitin in the *Drosophila* central nervous system under anoxia (70), and in protein expression during osmotic shock in isolated fish gill cells (71).

### *Bioindicators*

Owing to its responsiveness to diverse forms of stress, the heat-shock response has undergone widespread application in biomonitoring and environmental toxicology (72–75). In many cases, Hsps are especially useful biomarkers because their induction is much more sensitive to stress than traditional indices such as growth inhibition. The use of Hsps as biomarkers is most widespread in aquatic toxicology. Most of the literature demonstrates elevated Hsp levels or induction of Hsps under laboratory conditions and then proposes Hsps as a potential indicator of pollutants or toxins in the environment. For example, exposing freshwater sponges to pollutants extracted from river water elevates Hsp70 levels, which increase still further when thermal stress is also imposed (76). Additional examples of Hsp expression in aquatic toxicology concern rotifers, (77), marine sponges (78), amphipods (79), polychaetes (80), mollusks (81–84), and fish (85–87). Other applications purposefully deploy organisms in potentially polluted aquatic systems as biosensors (88, 89).

In the terrestrial environment, where heavy metal contamination and pesticide or herbicide accumulation can be critical problems, common soil organisms such as invertebrates (90) are useful Hsp-biomonitors of toxicants. For example, centipedes (*Lithobius*) collected from near a smelter had higher Hsp70 levels than those collected from unpolluted areas (91). Potentially, combinations of heavy metals can induce such distinctive patterns of Hsp expression in soil nematodes that these patterns can become diagnostic fingerprints for specific toxicants in soils (92).

Some aspects of the stress response, however, present problems for the use of Hsps as biomarkers in environmental toxicology. Because so many different

stresses induce Hsps, investigators may be unable to attribute changes in Hsp expression to any particular environmental stress. Organisms in the field often undergo multiple stresses simultaneously, the interaction of which can yield significant Hsp expression even when no single monitored toxicant is at harmful levels. Conversely, Hsps induced by another stress can enhance tolerance of a toxicant whose presence is being monitored. Laboratory studies support the difficulty of teasing apart environmental factors and attributing Hsp induction to a single stressor. For example, freshwater sponges exhibit greater tolerance of pollutants following a sublethal heat stress (76). Among the vertebrates, diseased fish have elevated levels of Hsps in their tissues, and disease-related expression may interfere with the use of Hsps as a biomarker (93). Thus, because numerous factors can induce Hsp expression and stress tolerance, the utility of Hsps as biomarkers of environmental toxins may be limited.

## ENVIRONMENTAL AND BIOLOGICAL CORRELATES OF THE HEAT-SHOCK RESPONSE AND HEAT-SHOCK PROTEINS

Many investigators view correlations of organismal traits (e.g. Hsp expression) and environmental or biological variables (e.g. level of environmental stress, developmental stage, distinctive role in a parasitic or symbiotic relationship with another species) as *prima facie* evidence of biological adaptation, and thus have actively sought such correlations in terms of the heat-shock response. While the probative value of such evidence in establishing adaptation has met with skepticism (94), in this section we consider the evidence for such correlations, whatever their meaning.

### *Variation in the Stress Response Along Environmental Gradients of Stress*

To understand how Hsps result in stress tolerance at the organismal level, many investigators have characterized the stress response along gradients that occur in nature. One central question is whether organisms from environments with little stress have a different or reduced stress response compared with organisms from environments with much stress. Little and much stress might correspond to the center and edge of a species' range, low versus high elevation, xeric and hot versus mesic and cool climate, temperate versus tropical/polar latitude, low versus high intertidal, and so on. In general, the resulting data support a correlation among Hsp expression, stress tolerance, and gradients of environmental stress. These gradients have received uneven attention, however, and their study has yielded mixed results. Comparative studies across many degrees of latitude have not produced the same results as studies of gradients on smaller scales

(e.g. diurnal or microclimatic variation in stress). Currently, not a single study has examined the stress response over the entire geographical distribution of a species; thus, whether species at the extremes of their distributions have an augmented heat-shock response is yet to be determined.

The majority of multi-species comparative studies focus on three aspects of the stress response: the minimum (threshold) and maximum temperatures at which Hsps are expressed and/or are present in cells, Hsp concentrations in cells, and the diversity of the specific Hsps that are expressed. Except for the work on threshold and maximum temperatures, much of this literature is a hodgepodge of disconnected studies that are seldom comparable because of methodological differences and permit few conclusions other than that species vary in the details of their stress response. Whether this variation has environmental correlates is uncertain. A rare and exemplary exception is the work of Bosch and colleagues on species of *Hydra* (95, 96); below we discuss this and other similar work.

In general, the threshold temperature for Hsp induction is correlated with the typical temperatures at which species live, with thermophilic species having a higher threshold than psychrophilic species. For example, a relatively cold-water, northern species of mussel (*Mytilus trossulus*) has a lower threshold for Hsp70 expression than its congener, *M. galloprovincialis*, a warm-water species with a more southern distribution (97). Limpet species that occur in the upper regions of the intertidal (*Lottia digitalis* and *Lottia pelta*) induce Hsps at 3–5°C higher than the threshold for limpets that occur lower in the intertidal (*Tectura scutum*) (AL Haag & GE Hofmann, unpublished data). Subtidal species of the marine snail *Tegula* exhibit much the same pattern (98). Aggregate expression of Hsp70 family members (17) occurs at 3–4°C higher in *Drosophila melanogaster* than in *D. ambigua*, a fruit fly of Palearctic origin (56). The same study reports a similar pattern for the desert ant *Cataglyphis* and *Formica polyctena*, a red wood ant from a temperate climate. One remarkable outcome of the *Cataglyphis* study is that Hsp synthesis in the desert ants continues at temperatures up to 45°C, whereas temperatures above 39°C inhibit Hsp synthesis in the temperate species. A similar pattern (although not as extreme) is evident for desert and non-desert *Drosophila* (22). These results suggest that translation itself may have an upper thermal maximum that varies among species adapted to different temperature environments.

Antarctic organisms represent a special case of psychrophily because the temperatures they experience are both extremely cold and extremely stable. In combination, do these conditions result in the evolutionary loss of a heat-shock response? In the subtidal alga, *Plocamium cartilagineum*, heat-inducible *hsp70* and *ubiquitin* transcription still occur, although the threshold is a spectacularly cold 5°C (99). Antarctic yeast species express Hsps at much lower temperatures than does *Saccharomyces*, and at least one species lacks inducible

thermotolerance (100, 101). In Antarctic fish, the picture is not as clear. Although a broadly cross-reactive anti-Hsp70 antibody can detect isoforms of Hsp70 in various tissues of the fish *Trematomus bernacchii*, heat shock temperatures from 6 to 10°C do not induce additional Hsp70 accumulation (GE Hofmann, unpublished data). A member of the *hsp70* gene family is present in two Antarctic fish species, *T. bernacchii* and *Notothenia coriiceps* (AC Whitmer & GE Hofmann, personal communication), and has been sequenced in Antarctic fish species (102). At the other extreme, some hyperthermophilic Archaea require temperatures in excess of 100°C to induce Hsp expression (see *Hsps of Archaea*).

Thermal stress gradients can be seasonal as well as geographic. In some cases, both Hsp expression and thermotolerance increase during warm seasons. The intertidal mussel *Mytilus californianus* displays significantly different Hsp induction profiles in summer than in winter, and summer-acclimatized mussels induce Hsps at a threshold temperature that is 6°C higher than the threshold in winter-acclimatized mussels (103). However, whether the accentuated Hsp expression in mussels in summer results in greater thermotolerance at the organismal level is unknown. Fish (34, 35) and intertidal invertebrates (31) also vary seasonally in Hsp expression.

In addition to work we cite elsewhere, other studies examine geographical gradients in fish (104, 105, 105a), maize (106), reptiles (107), and *Drosophila* (108); intertidal gradients in limpets (27); diurnal temperature change in *Drosophila* (109) and intertidal mussels (29); diurnal variation in spruce trees (46); and seasonal variation in insects (110).

One issue for future consideration is whether Hsps in general are specialized to function at higher temperatures than other proteins (especially enzymes), and whether homologous Hsps of species from various thermal environments have corresponding variation in thermostability of function (111). For example, that an Hsp's resistance to thermal denaturation varies according to the thermal niche of the species that expresses it has been demonstrated for only a single Hsp, alpha-crystallin (112). Another issue is how differing thresholds of Hsp expression have evolved, whether through mutations in HSF, general thermostability of proteins and cells (113), or some other mechanism (114).

### *The Parasitic Environment*

The roles of Hsps in host-parasite interactions have received considerable attention from both clinical and biological perspectives, with the majority of the research in two general categories. First, from the perspective of the host, Hsps expressed by invading parasites are potent antigens that elicit an immune response (115–117); parasites' Hsps are thus potentially useful in generating vaccines (118). From the perspective of the parasite, the synthesis of Hsps is a

cellular defense mechanism that enables the parasite to live in different thermal environments throughout its life-cycle (119). Parasites that infect mammalian and avian hosts can undergo profound changes in temperature during the transition to these hosts (with internal temperatures of 37°C or above) from ectotherm hosts or free-living stages. Induction of Hsps commonly accompanies this transition. Numerous studies have demonstrated developmentally regulated expression of Hsps in parasites; expression differs throughout the life-cycle both quantitatively and in the types of Hsps that temperature change induces. For example, mRNA transcripts for *hsp70* and *hsp83* homologues increase up to 100-fold as *Trypanosoma brucei* leaves the tsetse fly and enters a mammalian host (120). Aquatic snails release cercariae of the parasitic helminth, *Schistosoma mansoni*, into freshwater; cercariae penetrate human skin and develop into adult worms, eventually causing liver cirrhosis. The cercariae express two heat-inducible proteins that are not present in other stages (121).

Parasites that have an insect as the invertebrate vector have received much attention with regard to the developmentally regulated expression of Hsps. Examples include parasitic nematodes (122); cestode parasites (123); the malarial organism *Plasmodium* (124); *Borrelia burgdorferi*, the etiological agent of Lyme disease (125); the protist *Leishmania* (126); *Trypanosoma cruzi* (127); and *Theileria* (128).

Some parasite life cycles do not involve an animal vector; a free-living stage of the parasite occurs in water or soil and enters the host. In several cases, induction of Hsps accompanies the transition from the environment into the host. In the fungal parasite *Histoplasma capsulatum*, the temperature shift upon infection of a mammalian host cues both the transformation from a mycelial form to a budding yeast morphology and the expression of Hsps (129, 130). *Eimeria*, an intestinal parasite of numerous animals, expresses Hsp90 during infective life stages. *Eimeria* parasites are particularly interesting because this genus infects diverse hosts with correspondingly diverse body temperatures (e.g. marine fish, poultry, and cattle). However, specificity of infection is high at the species level, e.g. cattle are the exclusive host for *Eimeria bovis* (131). Whether the heat-shock response of *Eimeria* co-evolved with its speciation into these hosts is an open question.

Finally, the heat stress that infective life cycle stages of parasites experience is as diverse as their hosts. In nature, parasites of ectotherms can encounter dramatic shifts in temperature when their hosts' body temperature varies, as has been reported for parasites of reptiles, fish, and intertidal organisms (132, 133).

### *Symbiosis*

Just as Hsps may play an important role in parasitism, in which one species maintains a close but antagonistic relationship with others, they also function

in symbiosis, in which interspecific relationships can be equally close but not adversarial. Perhaps the most general example of this point concerns mitochondria and chloroplasts, which evolved from endosymbionts that colonized other cells early in the history of life. These organelles often require proteins that are encoded in the nuclear genome and synthesized by the host cell, and hence must be imported into the organelle. Hsps play diverse roles in this process in mitochondria. A cytoplasmic Hsp70 family member maintains peptides in an unfolded conformation, which enables the peptides to pass through pores in the mitochondrial membrane; a mitochondrial Hsp70 is part of the protein machinery that imports the peptide; and the Hsp60/Hsp10 apparatus participates in the folding of the imported protein (134). Several groups of primitive eukaryotes contain still other endosymbiotically derived organelles, the hydrogenosome and the nucleomorph, whose Hsps share a characteristic sequence with those of mitochondria and proteobacteria (135–138). The Hsp sequence similarities have been used to suggest that hydrogenosomes may derive from mitochondria, share a common origin with mitochondria, or represent independent colonizations of early eukaryotic cells (135–137, 139).

Aside from endosymbiotically derived organelles, the best-studied symbioses concern bacterial endosymbionts that infect insects, including aphids, flies, ants, and cockroaches. Aphids, for example, harbor the bacterium *Buchnera* in specialized cells (bacteriocytes) within a distinctive structure in the body cavity, the bacteriome (140). The bacteria express a protein, symbionin, at especially high levels, and this protein is a member of the GroEL (Hsp60) family. Other bacterial chaperones, including GroES (Hsp10) and DnaK (Hsp70), are also present at high levels (140). A similar phenomenon is evident in tsetse flies (141).

The function and significance of these high Hsp levels is enigmatic. The Hsps apparently are not a response of the endosymbionts to a novel (and therefore stressful) host environment, as the *Buchnera*/aphid symbiosis is 150–250 million years old. The endosymbionts, however, have been evolving at an especially high rate; thus, the elevated molecular chaperones could be compensating for decreased protein stability due to the accumulation of numerous amino acid substitutions (142, 143). Nonetheless, the bacteria themselves can mount a strong heat-shock response when their host undergoes stress (144). Other relevant symbioses include X-bacteria in the symbiosomes of *Amoeba* (e.g. 145), *Bradyrhizobium* and *Rhizobium* in the root nodules of nitrogen-fixing plants (146), and the zooxanthellae component of corals (26, 30). A recurrent theme is that the endosymbionts modify the amount and/or diversity of Hsps present in the symbiosis. In some cases, this modification is thought to contribute to the maintenance of the endosymbionts within the host, and in others to the augmentation of the heat-shock response of the symbiosis as a whole. Finally,

*Wolbachia*, a bacterial endosymbiont that infects millions of arthropod species, both interferes with the mating of infected and uninfected hosts and can alter their constitutive expression of Hsp70 and Hsp90 family members (147). Simulated natural heat stress can diminish this reproductive interference, possibly by overriding the symbiont's effect on the host Hsps.

### *Development*

Many species exhibit characteristic and distinctive patterns of Hsp expression (or nonexpression) during the various stages of development, including gametogenesis, embryogenesis, and metamorphosis (e.g. 148–151). These patterns are often consistent with enhanced stress resistance in developmental stages that encounter unusual levels of environmental stress or during circumstances such as dormancy and diapause (see below). In other cases, developmental programs of Hsp expression ensue even in the absence of any obvious environmental stress. One common pattern is that one or more Hsps are not expressed in the initial phases of embryogenesis (152–156) or late in gametogenesis (157–161), possibly because Hsps can be harmful to developing cells (see *Deleterious Aspects of Hsps*). Parental provision of Hsps or *hsp* mRNAs can sometimes override gametic or embryonic absence of Hsp expression (162, 163); in other cases this absence presumably poses a significant problem for continued development in the face of stress (164). Stress not only can kill vulnerable developmental stages outright, but also can produce lasting damage to surviving organisms, such as the phenocopying of genetic defects; Hsps may minimize such defects (165, 166).

Adaptational analyses of the developmental expression of Hsps are diverse. Some plant seeds presumably must endure extremes of heat, desiccation, and other stresses before germinating, and some must germinate under especially challenging conditions. However, although seeds clearly undergo developmentally regulated expression of Hsps and embryos can express Hsps in response to environmental stress (167–169), few investigators have considered whether these patterns of expression are amplified or modified in species and ecotypes that naturally encounter especially challenging stress regimes (68, 170). Our state of knowledge is similar for fungal spores, which express particular Hsps in a developmentally regulated program (e.g. 171, 172). Several interesting case studies are available for animals, although a general pattern is yet to emerge. In the most spectacular example, encysted brine-shrimp (*Artemia*) embryos undergo developmental arrest, in which they may survive for years without environmental water or oxygen. The encysted embryos accumulate enormous concentrations (15% of total protein) of a small Hsp (173–177) and trehalose (178), and suppress ubiquitination of damaged proteins (179). Non-adult *D. melanogaster* infest necrotic fruit, which can become extremely warm if it is sunlit (53, 54); this species mounts a massive heat-shock response, which

is greatest in the developmental stages that presumably undergo the most exposure to natural heat stress (18, 19, 180). Other flies overwinter while at a particular developmental stage, and undergo considerable Hsp expression in response to cold (60–63). Later in development, ubiquitin may assist in the degeneration of flight muscles that are no longer needed after nuptial flights of insects (181). Finally, the temperature threshold for expression of Hsps may itself undergo modification; e.g. the threshold decreases in mammalian testis, in which gametogenesis normally occurs at lower temperatures than in the core of the body (182, 183).

### *Aging and Senescence*

As mammals age, damage to proteins progressively accumulates, and both the ability to express Hsps (e.g. Hsp or *hsp* mRNA levels after a standard exposure to heat or other stress) and stress tolerance deteriorate (184, 185). Moreover, individual Hsps can become less able to mitigate the effects of stress on proteins as mammals age (186). In ecological and evolutionary terms, whether similar Hsp-aging relationships are important or even evident in wild organisms is unknown, although these relationships occur in diverse species in the laboratory: *Drosophila* (187, 188), nematodes (189–191), and *Daphnia* (192).

These findings have provoked great interest in how Hsps potentially affect senescence and lifespan. A unifying hypothesis in the foregoing work is that protein damage, due primarily to oxidation/free radical activity, gradually accumulates during the life of a cell or organism and can lead to death if unabated; Hsps and other molecular stress responses ordinarily can mitigate this damage to some extent, and the decreasing expression of Hsps with age therefore contributes to mortality. If this hypothesis is correct, then treatments that both reduce damage to protein and increase Hsp expression (e.g. heat shock) should prolong life. In nematodes (*Caenorhabditis elegans*), some single-gene mutations that increase lifespan are associated with increased thermotolerance, but through as-yet-undescribed mechanisms (189–191). In *Drosophila*, heat shock extends lifespan (193), and this extension is enhanced in flies transformed with additional copies of the *hsp70* gene (194). Nutrient deprivation can also extend life in rodents, presumably by reducing the metabolic rate and consequently, oxidative damage to proteins; starvation, however, variously increases, decreases, or has no effect on Hsp expression (195–199).

## FITNESS CONSEQUENCES OF HSP EXPRESSION

### *General Issues and Beneficial Aspects of Hsps*

Understanding the consequences of variation in Hsps and the stress response for Darwinian fitness requires a detailed appreciation of the mechanisms by which Hsps mitigate the impact of stress on individuals in natural populations. These

mechanisms are becoming well understood at the level of model proteins with which Hsps can interact, but are progressively less well understood at the level of the cell, tissue, organ, and whole organism. At the level of the model protein, various stresses clearly either directly or indirectly result in conformational change, and Hsps typically promote the reacquisition or maintenance of the native structure and function by minimizing the tendency of non-native proteins to interact inappropriately (200, 201). In cells, stress-induced conformational change, protein aggregation, and molecular chaperoning of model proteins are also well established (200–202), and many cellular components differ in stress-tolerant and stress-intolerant cells (67).

Two primary issues impede the linkage of variation in these well-established mechanisms and phenomena to variation in the fitness of individual complex multicellular eukaryotes. First, is the variation in sensitivity to stress among cells, cell types, tissues, organs, and organisms attributable to a small number of critical lesions, especially sensitive targets of stress and functions of specific Hsps in protecting or repairing these lesions/targets? Or is variation in sensitivity to stress an aggregate function of a widespread and diverse impact of stress on cellular structures, with Hsps mitigating multiple lesions in diverse ways (21)? The former alternative may be more analytically tractable than the latter. Second, given that cells and organisms may have multiple Hsps in each Hsp family, multiple Hsp families, and multiple non-Hsp mechanisms of stress mitigation, how can we unambiguously establish the contribution or importance of any particular Hsp, Hsp family, or mechanism in the complex cell, tissue, organ, or organism? Much of the published literature on the functional consequences of Hsp expression for whole organisms or the cells they comprise runs afoul of these issues. Literally thousands of studies report correlations between Hsp expression, diverse biological functions in the face of stress, and stress tolerance, but these typically conclude that their findings are at best consistent with a role of one or more Hsps in stress tolerance. Evaluating the roles of single factors in complex systems is an ongoing challenge in most areas of the biological sciences, and the heat-shock field largely has not yet deployed counterparts of the solutions that other fields have developed.

One conspicuous and major exception includes techniques and approaches, primarily drawn from molecular biology and genetics, that allow the manipulation of individual Hsps or specific genes that encode them. In rare instances, a species naturally may have an unusual genetic system (203) or a diminished suite of Hsps (95, 204) that accomplishes the same end; also, several chemical compounds may specifically inhibit one or more Hsps (e.g. 205, 206). The general implication of the resulting work (Table 1) is that, even in whole organisms or the cells they comprise, variation in single Hsps can be consequential for fitness. Some specific implications are as follows: Individual Hsps can have

**Table 1** Phenotypes of multicellular eukaryotes, and the cells and tissues that they comprise, for which Hsps are necessary and/or sufficient<sup>a</sup>

Protein	Phenotype
Hsp10	Cellular: tolerance of ischemia (no phenotype) (308); tolerance of ischemia when co-expressed with Hsp60 (308)
Hsp27	Cellular: resistance to chemotherapeutic drugs (309); resistance to hydrogen peroxide (310, 311); resistance to hydrogen peroxide (no phenotype) (312); resistance to ultraviolet radiation (no phenotype) (312); resistance of tumor cells to monocytes (311); sensitivity to lymphokine-activated killer cells (no phenotype) (311); tolerance of hyperthermia (312–316); resistance to tumor necrosis factor (317) (310); tolerance of ischemia (318); resistance of actin polymers to cytochalasin (314); accelerated nuclear protein aggregation (319); accelerated decline of thermal radiosensitization (319)
Crystallin	Cellular: tolerance of hyperthermia (320, 321); tolerance of ischemia (318); resistance to tumor necrosis factor (310); resistance to hydrogen peroxide (310)
Hsp60	Cellular: tolerance of hyperthermia (no phenotype) (322, 323); tolerance of ischemia (no phenotype) (308, 322, 323); tolerance of ischemia when co-expressed with Hsp10 (308)
Hsp65	Cellular: tumor regression (324); loss of tumorigenicity (325) Tissue/organ: regression of malignant tumors (324)
Hsp70	Cellular: tolerance of hyperthermia (326) (316, 322, 323, 327–342); tolerance of ischemia/hypoxia (322, 323, 340, 343–345); recovery from translational and transcriptional inhibition following heat shock (335); regulation of heat-shock response (331, 346, 347); tolerance of endotoxin (348); reduced protein denaturation upon heat exposure (349); tumorigenicity (350); cell proliferation (351, 164); resistance to hydrogen peroxide (311); resistance of tumor cells to monocytes (311); sensitivity to lymphokine-activated killer cells (no phenotype) (311); escape from drug-induced cell cycle arrest (352); protein glycosylation (353); tolerance of ultraviolet radiation (354); apoptosis (351, 355, 356); resistance to apoptosis (no phenotype) (328, 329) Tissue/organ: recovery of contractility after ischemia (345, 357–359); reduction in myocardial infarct size (345, 359); reduction of hyperthermic damage to midgut (221); resistance of heart to ischemic injury (357–359); resistance of hippocampus to ischemic injury (360) Organismal: tolerance of hyperthermia (18–20, 109, 156, 203, 221, 224, 278, 361); growth and development (222); regulation of heat-shock response (361); persistence in nature (no phenotype) (277)
Hsc70	Organismal: tolerance of hyperthermia (203)
Hsp72	Cellular: apoptosis (no phenotype) (362); protection against heat-induced nuclear protein aggregation (319); protection against hypoxia (363); protection against thermal radiosensitization (319) Tissue/organ: reduction in myocardial infarct size (364)
Grp78	Cellular: protein secretion (229–231)

*(Continued)*

**Table 1** (Continued)

Protein	Phenotype
Hsp90	Cellular: tolerance of hyperthermia (322, 323, 328, 329, 340); tolerance of ischemia (no phenotype) (322, 323, 340); apoptosis (362); apoptosis (no phenotype) (328, 329); cell proliferation and cell cycle control (365); glucocorticoid receptor function (205)
Hsp100	Organismal: host infection in <i>Leishmania</i> (126)
Hsp101	Cellular: tolerance of hyperthermia (366)
Many Hsps	Cellular: recovery of cell proliferation after heat shock (367); recovery from chromosome damage after heat shock (367–369); tolerance of hyperthermia (370, 371); tolerance of ischemia (336)
HSF	Organismal: oogenesis and development (372); thermotolerance (372, 373)

<sup>a</sup>In all cited work, specific Hsps have undergone experimental or natural manipulation.

pleiotropic effects, interacting with multiple systems in diverse ways. Findings from manipulations of individual Hsps usually (but not always) are consistent with the outcome of correlative studies (see above). Finally, despite the huge body of work on Hsps and the growing use of manipulative techniques, we have remarkably little physiological insight into exactly how the activity of Hsps culminates in the enhanced stress tolerance of multicellular eukaryotes and the cells and tissues that they comprise.

Interestingly, one clear conclusion that correlative studies have yielded is that Hsps cannot account for the entirety of inducible stress tolerance (207–217). Indeed, some component of inducible stress tolerance may be unrelated to protein synthesis in general (214, 218, 219).

### *Deleterious Aspects of Hsps*

The many advantages of the heat-shock response suggest that natural selection should maximize the expression of Hsps. By contrast, the genes encoding Hsps have not undergone unlimited amplification in the genome, and the Hsps themselves are subject to strict autoregulation by multiple molecular mechanisms (220). These contrary findings suggest that Hsps can have both positive and negative impacts on fitness, and that natural selection may have acted to balance these impacts in setting the level of Hsp expression. For example, while small to moderate increases in Hsp70 levels enhance inducible thermotolerance in *Drosophila*, large increases in Hsp70 levels actually decrease thermotolerance (221); evolution thus may favor an intermediate level of Hsp70. A common theme in related work is that high levels of Hsps may be especially detrimental to cells or developmental stages in which cell growth and division proceed at high rates. *Drosophila* larvae transformed with extra copies of the *hsp70* gene have greater larva-to-adult mortality and slower development than do control

larvae; these strain differences are proportional to the number of Hsp-inducing heat shocks administered to the larvae (222). Larvae naturally varying in Hsp70 expression display a similar pattern (223). *Drosophila* cells engineered to express Hsp70 constitutively at first grow more slowly than control cells, but subsequently resume control growth rates once the Hsp70 is sequestered from the cytoplasm (164); indeed, *Drosophila* embryos remove Hsp70 from their cells rapidly after heat shock (224). A yeast strain that cannot express Hsp104 grows faster than its wild-type counterpart on some media (171). More generally, most animal species that have been studied do not mount a heat-shock response during early stages of embryogenesis (see *Development*), when protein synthesis may be especially intense.

These negative effects may have at least two nonexclusive explanations (222, 225, 226): First, Hsps at high concentration could be toxic, directly interfere with ongoing processes in the cell, or otherwise alter function to the detriment of the cell (220). Second, the synthesis and degradation of Hsps could consume an intolerably large fraction of a cell's or organism's nutrient and energy stores, and/or occupy so large a fraction of the synthetic/catabolic apparatus that the processing of other essential biomolecules suffers (226–228). Consistent with the first explanation, cellular sequestration of Hsp70 is correlated with the resumption of proliferation in cells constitutively expressing this protein (164). Also, overexpression of an Hsp70 family member inhibits protein secretion and reduction increases secretion in mammalian cells in culture (229–231); excess amounts of another Hsp70 family member can promote protein aggregation in vitro (M Borrelli & J Lepock, personal communication); and Hsp70 can perturb the normal structures of nascent polypeptides (232).

Tests of the second explanation have manipulated the costs of or resources for Hsp expression. Growth of corn in nitrogen-rich soil increases the synthesis of Hsps in response to a standard heat shock (233); in plants grown in nitrogen-poor soil, other proteins may be catabolized to supply amino acids for synthesis of Hsps (234). These findings suggest that Hsp synthesis can be nitrogen-limited in plants. Starvation reduces the expression of Hsp 70 family members in mice (195). In *Drosophila* larvae, by contrast, co-expression of  $\beta$ -galactosidase and Hsps has no greater cost for growth and development than does expression of Hsps alone (225). Further study of this apparent trade-off of the benefits and disadvantages of Hsp expression, moreover, has the potential to link evolutionary and mechanistic views of this problem that heretofore have been separate (222, 225).

## MICROEVOLUTIONARY VARIATION IN HSPS

Hsps are routinely touted as adaptations that arose and are maintained via natural selection for stress resistance. Origin and maintenance of a trait by selection

require that it vary within populations, and that this intra-population variation have a genetic basis and affect the Darwinian fitness of individuals. Here we ask whether Hsps, the genes that encode them, and the factors that modify their expression display such patterns of variation and undergo stabilizing or directional selection in response to environmental stress.

First, not all intrapopulation and intraspecific variation results from genetic differences. For example, seasonal acclimatization and temperature acclimation in the laboratory can alter the minimum temperature at which *Gillichthys*, a gobiid fish, expresses an Hsp90 family member (34, 235). Seasonal acclimatization likewise affects Hsp70 levels in mussels (*Mytilus*) (103), and routine culture temperature affects the magnitude and temporal pattern of Hsp expression in HeLa cells (236). Such changes may stem from alterations in the cellular environment that modify the activation of HSF (113, 182, 236, 237). These changes, however, are not universal; laboratory thermal acclimation does not alter the thermal sensitivity of Hsp expression in fish hepatocytes in culture (238), *Drosophila* larvae (19), and mussels (103).

Even when acclimation and seasonal change are controlled, however, individuals within a population or species may vary in Hsp expression and/or the genes that determine it. Relevant research has examined this issue on two levels: direct sequence variation and restriction fragment length polymorphisms (RFLPs) (240). The sequence of *hsp70* varies among strains of the parasite *Trypanosoma* (241, 242) and the nematode *C. elegans* (243), and among conspecifics for some but not all of the mammalian *hsp70* family members (240), as does that of the 3' untranslated region of *hsp27* in normotensive and hypertensive rats (244). RFLPs consistent with intraspecific variation either in the *hsp* genes or flanking regions are detectable in the *hsp60* and *hsp70* of the spirochete *Borrelia* (245), in multiple *hsp70* family members of mammals (240, 246, 247), and in several plant species (248–250). One putative instance of intraspecific variation in *hsp* copy number concerns *D. melanogaster*, in which at least five nearly identical copies of *hsp70* occur at two chromosomal loci. At locus 87A7, two copies are arranged as an inverted repeat (251, 252); at 87C1, two copies flank a region containing at least one additional copy (252, 253) plus numerous  $\alpha/\beta$  repeats, which encode heat-inducible mRNAs of no proven function (254–257). Up to five additional *hsp70* copies have been reported, with copy number varying among strains and time of year (253, 255, 258–261). However, these reports either cannot exclude that such variation is actually in intergenic regions or that the reports are for *Drosophila* cells in culture or mutagenized laboratory strains rather than wild or even wild-type strains. The organization of the two chromosomal loci reportedly varies among natural populations (253). A less equivocal instance of evolutionary change in gene copy number concerns *Arabidopsis*, in which ecotypes vary in the number of ubiquitin-encoding repeats (262). For all

of the foregoing reports, the functional significance of intraspecific variation awaits elucidation or direct verification.

Hsp expression also exhibits genetic variation among individuals of a species; often, this variation is correlated with stress resistance (248, 250, 263–271). For example, isofemale lines of *Drosophila* founded from a single wild population differ more than twofold in Hsp70 expression; this variation is correlated with thermotolerance and is heritable (180, 223). Similarly, in the pathogenic fungus *Histoplasma*, naturally temperature-insensitive strains express more *hsp70* mRNA and do so at lower temperatures than in a temperature-sensitive strain (272). In humans, fibroblasts isolated from desert-dwelling Turkmen express more Hsps and have greater thermotolerance than fibroblasts from residents of more equable climates (273); presumably, however, these peoples do not differ in body temperature.

Given that the patterns of variation necessary for natural selection occur within species, that selection can alter Hsp expression is not surprising. Laboratory evolution at high temperatures paradoxically lowers Hsp70 expression and inducible thermotolerance in *Drosophila* (19, 274), and selection for resistance to hyperthermic paralysis alters both the *hsp68* promoter and the *hsp-omega* locus in *Drosophila* (254, 275). Additional findings relevant to natural selection and its underlying genetic basis come from closely related species, some so similar that they hybridize. In the fish species *Poeciliopsis monacha* and *P. lucida*, an unusual genetic system permits the generation of hemiclinal lines in which the paternal genome varies against a constant maternal genome. Hemiclinal thermotolerance was most strongly related to Hsc70 and only secondarily to Hsp70 levels, in a pattern consistent with straightforward Mendelian inheritance of parental genotypes and adaptation to the local thermal environment. By contrast, the heat-shock response of interspecific hybrids of tomato (*Lycopersicon*) is not intermediate to the parental responses (276). Non-hybridizing congeners often exhibit a correlation among the actual or inferred incidence of thermal stress in their environment, heat-shock response, and stress tolerance. Such data are now available for diverse animals (see *Variation in the Stress Response Along Environmental Gradients of Stress*).

A particular problem with such species comparisons is that the interpretation of the observed patterns is readily confounded by phylogenetic and statistical issues. A more general problem with both laboratory evolution and species comparisons is that they describe only a supposed correlation of the heat-shock genes/proteins of interest with evolution and seldom can establish the importance of the genes/proteins of interest to evolutionary process and outcome (19, 20). Study of free-ranging organisms (277) with *hsp* transgenes (e.g. 109, 278) may contribute much to resolving these problems.

## THE EVOLUTIONARY HISTORY OF HSPTS AND THE GENES THAT ENCODE THEM

Hsps are among the most ancient and highly conserved of all proteins. Homologues of Hsps occur in every species in which they have been sought, and in all kingdoms of living things. Thus, Hsps represent a remarkable example of molecular “descent with modification” at the levels of gene sequence, genomic organization, regulation of gene expression, and protein structure and function. So clear are the patterns of descent and modifications that they can be used to establish the evolutionary origins and the phylogenetic affinities of the major groups of organisms.

### *Hsps of Archaea and Exceptional Prokaryotes*

The Archaea or archaebacteria are the most extremophilic and most primitive organisms. The heat-shock response of extremophilic Archaea and nonarchaeal extremophiles occurs at remarkably high temperatures (279), e.g. 88°C in *Sulfolobus* (279) and >100°C in the hyperthermophilic species designated ES4, a heterotrophic sulfur reducer isolated from a deep-sea hydrothermal vent (280). The archaeal genome encodes homologues of most Hsps represented in other prokaryotes and eukaryotes (279), as well as their consensus promoter sequences (281). Notably, the archaeal Hsp60 homologues assemble into a dual ring-like structure, termed a rosettasome or thermosome, that resembles the structure that the chaperones GroEL and GroES form in bacteria (279, 282, 283). The archaeal structures have ATPase activity and can bind denatured proteins (282). At least some Archaea and Eubacteria differ in the number of monomers that comprise these structures (284). Surprisingly, the archaeal Hsp60s (e.g. TF55 of *Sulfolobus*) most closely resemble not a bacterial homologue, but the eukaryotic protein TCP1, which assembles into the t-complex polypeptide 1 ring complex (TRiC) in the cytosol (285, 286). Previously, Hsp60 homologues were thought to be absent from the eukaryotic cytosol. A growing body of evidence suggests that TCP1/TRiC and GroEL/GroES play comparable roles in their respective organisms and cellular compartments (287–289). Meanwhile, Trent (290) has suggested that the primary function of TF55 may be cytoskeletal, with molecular chaperoning a secondary or derived function.

Genes encoding Hsps are present even in the smallest of genomes. These include the genomes of mycoplasmas (291) and the nucleomorph, the vestigial nucleus of a phototrophic eukaryotic endosymbiont in cryptomonad algae (138). The section on symbiosis (see above) reviews the distribution of Hsps in various other organelles of endosymbiotic origin. Apparently, the problem of protein folding is ancient and ubiquitous, necessitating molecular chaperones in these diverse cases.

### *Large-Scale Evolution of hsp Genes*

The extraordinarily conserved nature of *hsp* genes (292) has facilitated their cloning, sequencing, and comparison in diverse organisms; their evolution is now becoming understood in detail. Gupta and colleagues have undertaken the most extensive surveys of *hsp* sequences, with a particular focus on organisms deemed critical to understanding the relationships of major taxa (292–297). The interpretations resulting from these comparisons relate to hypotheses about (a) the origin of eukaryotic cells, the eukaryotic nucleus, and endoplasmic reticulum (292, 296); (b) polyphyletic versus monophyletic origin of the major bacterial groups (292–294); and (c) the validity of the three-domain (Archaea, bacteria, and eukaryotes) dogma (292, 294). Whereas these interpretations are controversial, if not revolutionary, and therefore have not received universal acceptance, they nonetheless clearly illustrate how comparative analyses of *hsp* genes may address fundamental issues in evolutionary history.

On a less grand scale, *hsp* gene families represent superb case studies of how one or a small number of primitive genes can diversify to encode a suite of compartment- and function-specific proteins. One of many examples is *dnaK*, a single gene in Archaea and bacteria that has become the complex multi-gene *hsp70* families of *Saccharomyces* (298), *Drosophila* (253, 299), and *Homo* (300). Another example concerns the small Hsps, which evolution recruited to become a major component of the lens of the eye: alpha crystallin (301). A growing body of work examines the discrete evolutionary events by which these changes may have occurred, including gene duplication/conversion events (302), retrotransposition, horizontal exchange of genomes, and others. New technologies promise to advance this work exponentially.

### *Discrete Examinations of Molecular Evolution*

Ideally, a complete study of the evolutionary physiology of Hsps might examine how the following co-evolve as populations or how closely related species enter environments in which they face novel stresses: sequence (both coding and non-coding) of the gene(s) for a particular Hsp, regulation of *hsp* gene expression, the role and importance of the Hsp in stress resistance, and the intensities and durations of stress that the populations and species actually face. Much of the evolutionary physiological investigation of Hsps fails to attain this admittedly ambitious goal for one or more reasons: (a) The species under study are too distantly related to reconstruct the functional, environmental, and genetic events during their divergence; (b) molecular biology, manipulative genetics, physiology, and environmental assessment are not all possible for the species in question; or (c) the breadth of the techniques and approaches necessary to perform such research is too daunting for a single research program. Two case studies exemplify both how this goal could be approached and how far the field has yet to go to attain it.

The coelenterate *Hydra oligactis* and several of its congeners are the only multicellular eukaryotes reported not to express Hsps in response to heat shock and other stresses. Other congeners (e.g. *Hydra attenuata* and *H. magnipapillata*) have a well-developed stress response; these and other data for putative ancestors of *Hydra* suggest an evolutionary loss of Hsp expression in *H. oligactis* (95, 96). Physiologically and ecologically, *H. oligactis* is deficient in inducible stress tolerance and disappears from certain habitats in nature during periods of stress (95). Subsequent work suggests that, at least for Hsp70, the loss has occurred due to mutations that affect the stability of *hsp70* mRNA, as *H. oligactis* has an *hsp70* gene and expresses a heat-inducible *hsp70* mRNA in quantities similar to that in the heat-tolerant *H. magnipapillata* (96).

Dipteran insects and their ancestors have undergone an evolutionary proliferation of *hsp70* genes. Mosquitoes and *Drosophila* share a distinctly arranged duplication of the inducible *hsp70* gene (303), suggesting that this proliferation predates the original diversification of the Diptera. Within the genus *Drosophila*, all groups other than the *melanogaster* subgroup of species apparently retain the primitive copy number of two (108, 304, 305). Within the *melanogaster* subgroup, all species examined to date have four copies except for *D. melanogaster* (253), which has at least five *hsp70* copies in its haploid genome (see Microevolutionary Variation in Hsps). Curiously, *D. melanogaster* expresses no Hsp100 family members, which are critical for thermotolerance in other organisms (204, 306). The proliferation of *hsp70* copies is correlated with the ecological and biogeographic distribution of *Drosophila* species (19). Whereas most *Drosophila* species have small geographic ranges or narrow ecological niches, two of the *melanogaster* subgroup species (*simulans* and *melanogaster*) have cosmopolitan distributions, and a third (*yakuba*) is ecologically diverse throughout sub-Saharan Africa.

## CONCLUSION

Ecological and evolutionary physiological analysis of heat-shock proteins may be nearing the end of its initial descriptive phase. Although accounts of spectacular levels of Hsp-mediated stress resistance and exceptional consequences of Hsp expression will continue to be newsworthy, the major patterns of Hsp expression in multicellular eukaryotes are becoming so obvious that additional descriptive work is becoming increasingly difficult to justify. Clearly, however, major questions remain unanswered. How the activities of Hsps at the molecular level culminate in organismal stress tolerance and how the *hsp* genes, their regulation, the function of the proteins they encode, and the environments faced by the organisms in which they occur all co-evolve are but two of the

unresolved issues reviewed here. The perspective of evolutionary physiology can make significant contributions to the resolution of these and other issues. By placing results in actual environmental contexts, by assessing phenotypes of Hsps in the context of whole multicellular organisms, and by characterizing extant and historical variation in Hsps in natural populations and taxa, evolutionary physiologists can complement and extend a spectacular area of research that has been largely restricted to the molecular/cellular levels in the laboratory. By the same token, insights and techniques that laboratory-based investigators provide promise to continue to revolutionize the ecological and evolutionary study of Hsps. These approaches are both logical partners and necessary complements to one another (22, 307). Our understanding of Hsps has much to gain from the continued if not expanded synergy of these approaches.

#### ACKNOWLEDGMENTS

We thank BR Bettencourt, K Chavez, UT D'OBradior, AP Nguyen, J Meredith-Patla, and AC Whitmer for bibliographic assistance and/or editorial advice, and Susan Lindquist for founding the bibliographic database that made this review possible. Research was supported by National Science Foundation grants 97-23298 and 97-23063. Finally and most importantly, we beg the indulgence of numerous investigators whom space limitations precluded us from citing directly here; their contributions have made the field what it is today. We urge all readers contemplating research in this area to scan the complete bibliography (13) available online at <http://www.AnnualReviews.org>.

Visit the *Annual Reviews* home page at  
<http://www.AnnualReviews.org>

#### Literature Cited

1. Ritossa F. 1996. Discovery of the heat shock response. *Cell Stress Chaperones* 1:97-98
2. Lindquist S. 1986. The heat-shock response. *Annu. Rev. Biochem.* 55:1151-91
3. Gething MJ, Sambrook J. 1992. Protein folding in the cell. *Nature* 355:33-45
4. Morimoto RI, Tissieres A, Georgopoulos C, eds. 1994. *Heat Shock Proteins: Structure, Function and Regulation*. Cold Spring Harbor, NY: Cold Spring Harbor Lab. Press
5. Hartl FU. 1996. Molecular chaperones in cellular protein folding. *Nature* 381:571-80
6. Gething MJ, ed. 1997. *Guidebook to Molecular Chaperones and Protein-Folding Catalysts*. Oxford, UK: Oxford Univ. Press
7. Thomas PJ, Qu BH, Pedersen PL. 1995. Defective protein folding as a basis of human disease. *Trends Biochem. Sci.* 20:456-59
8. Boston RS, Viitanen PV, Vierling E. 1996. Molecular chaperones and protein folding in plants. *Plant Mol. Biol.* 32:191-222
9. Feige U, Morimoto RI, Yahara I, Polla BS, eds. 1996. *Stress-Inducible Cellular Responses*. Basel: Birkhäuser
10. Feder ME, Parsell DA, Lindquist SL. 1995. The stress response and stress proteins. In *Cell Biology of Trauma*, ed. JJ Lemasters, C Oliver, pp. 177-91. Boca Raton, FL: CRC

11. Nover L, ed. 1991. *Heat Shock Response*. Boca Raton, FL: CRC
12. Somero GN. 1995. Proteins and temperature. *Annu. Rev. Physiol.* 57:43–68
13. Feder ME, Hofmann GE. 1998. Evolutionary and ecological physiology of heat-shock proteins and the heat-shock response: a comprehensive bibliography. <http://www.AnnuReviews.org>
14. Stephanou G, Alahiotis SN, Christodoulou C, Marmaras VJ. 1983. Adaptation of *Drosophila melanogaster* to temperature. Heat-shock proteins and survival in *Drosophila melanogaster*. *Dev. Genet.* 3:299–308
15. Alahiotis SN, Stephanou G. 1982. Temperature adaptation of *Drosophila* populations. The heat shock proteins system. *Comp. Biochem. Physiol.* 73B:529–33
16. Palter KB, Watanabe M, Stinson L, Mahowald AP, Craig EA. 1986. Expression and localization of *Drosophila melanogaster* hsp70 cognate proteins. *Mol. Cell Biol.* 6:1187–203
17. Hightower LE. 1995. Desert ants. *Science* 268:1417
18. Feder ME. 1998. Engineering candidate genes in studies of adaptation: the heat-shock protein Hsp70 in *Drosophila melanogaster*. *Am. Nat.* In press
19. Feder ME, Krebs RA. 1998. Natural and genetic engineering of thermotolerance in *Drosophila melanogaster*. *Am. Zool.* 38:503–17
20. Feder ME, Krebs RA. 1997. Ecological and evolutionary physiology of heat-shock proteins and the stress response in *Drosophila*: complementary insights from genetic engineering and natural variation. In *Stress, Adaptation, and Evolution*, ed. R Bijlsma, V Loeschcke, pp. 155–73. Basel: Birkhäuser
21. Feder ME. 1996. Ecological and evolutionary physiology of stress proteins and the stress response: the *Drosophila melanogaster* model. In *Animals and Temperature: Phenotypic and Evolutionary Adaptation*, ed. IA Johnston, AF Bennett, pp. 79–102. Cambridge, UK: Cambridge Univ. Press
22. Huey RB, Bennett AF. 1990. Physiological adjustments to fluctuating thermal environments: an ecological and evolutionary perspective. In *Stress Proteins in Biology and Medicine*, ed. RIMorimoto, A Tissieres, C Georgopoulos, pp. 37–59. Cold Spring Harbor, NY: Cold Spring Harbor Lab. Press
23. Huey RB. 1991. Physiological consequences of habitat selection. *Am. Nat.* 137:S91–115
24. Bartholomew GA. 1964. The roles of physiology and behaviour in the maintenance of homeostasis in the desert environment. In *Homeostasis and Feedback Mechanisms*, ed. GM Hughes, pp. 7–29. Cambridge, UK: Cambridge Univ. Press
25. Sharp VA, Brown BE, Miller D. 1997. Heat shock protein (HSP 70) expression in the tropical reef coral *Goniopora djiboutiensis*. *J. Therm. Biol.* 22:11–19
26. Hayes RL, King CM. 1995. Induction of 70-kD heat shock protein in scleractinian corals by elevated temperature: significance for coral bleaching. *Mol. Mar. Biol. Biotechnol.* 4:36–42
27. Sanders BM, Hope C, Pascoe VM, Martin LS. 1991. Characterization of the stress protein response in two species of *Collisella* limpets with different temperature tolerances. *Physiol. Zool.* 64:1471–89
28. Sanders BM, Pascoe VM, Nakagawa PA, Martin LS. 1992. Persistence of the heat-shock response over time in a common *Mytilus* mussel. *Mol. Mar. Biol. Biotechnol.* 1:147–54
29. Hofmann GE, Somero GN. 1996. Protein ubiquitination and stress protein synthesis in *Mytilus trossulus* occurs during recovery from tidal emersion. *Mol. Mar. Biol. Biotechnol.* 5:175–84
30. Sharp VA, Miller D, Bythell JC, Brown BE. 1994. Expression of low molecular weight HSP 70 related polypeptides from the symbiotic sea anemone *Anemonia viridis* Forskal in response to heat shock. *J. Exp. Mar. Biol. Ecol.* 179:179–93
31. Hofmann GE, Somero GN. 1995. Evidence for protein damage at environmental temperatures: seasonal changes in levels of ubiquitin conjugates and hsp70 in the intertidal mussel *Mytilus trossulus*. *J. Exp. Biol.* 198:1509–18
32. Miller D, McLennan AG. 1988. The heat shock response of the cryptobiotic brine shrimp *Artemia*. I. A comparison of the thermotolerance of cysts and larvae. *J. Therm. Biol.* 13:119–24
33. Miller D, McLennan AG. 1988. The heat shock response of the cryptobiotic brine shrimp *Artemia*. II. Heat shock proteins. *J. Therm. Biol.* 13:125–34
34. Dietz TJ, Somero GN. 1992. The threshold induction temperature of the 90-kDa heat shock protein is subject to acclimatization in eurythermal goby fishes (genus *Gillichthys*). *Proc. Natl. Acad. Sci. USA* 89:3389–93
35. Fader SC, Yu Z, Spotila JR. 1994. Seasonal variation in heat shock proteins

- (hsp70) in stream fish under natural conditions. *J. Therm. Biol.* 19:335–41
36. Near JC, Easton DP, Rutledge PS, Dickinson DP, Spotila JS. 1990. Heat shock protein 70 gene expression in intact salamanders *Eurycea bislineata* in response to calibrated heat shocks and to high temperatures encountered in the field. *J. Exp. Zool.* 256:303–14
  37. Gates DM. 1980. *Biophysical Ecology*. New York: Springer-Verlag
  38. Kee SC, Nobel PS. 1986. Concomitant changes in high-temperature tolerance and heat-shock proteins in desert succulents. *Plant Physiol.* 80:596–98
  39. Nagao RT, Kimpel JA, Key JL. 1990. Molecular and cellular biology of the heat-shock response. *Adv. Genet.* 28:235–74
  40. Nguyen HT, Joshi CP, Klueva N, Weng J, Hendershot KL, Blum A. 1994. The heat-shock response and expression of heat-shock proteins in wheat under diurnal heat stress and field conditions. *Aust. J. Plant Physiol.* 21:857–67
  41. Hamilton EW, Heckathorn SA, Downs CA, Schwarz TE, Coleman JS, Hallberg RL. 1996. Heat shock proteins are produced by field-grown naturally occurring plants in the summer in the temperate northeast U.S. *Bull. Ecol. Soc. Am.* 77, Suppl. Part 2:180 (Abstr.)
  42. Kimpel JA, Key JL. 1985. Presence of heat shock mRNAs in field grown soybeans. *Plant Physiol.* 79:672–78
  43. Hernandez LD, Vierling E. 1993. Expression of low molecular weight heat-shock proteins under field conditions. *Plant Physiol.* 101:1209–16
  44. Hendershot KL, Weng J, Nguyen HT. 1992. Induction temperature of heat-shock protein synthesis in wheat. *Crop Sci.* 32:256–61
  45. Burke JJ, Hatfield JL, Klein RP, Mullet JE. 1985. Accumulation of heat shock proteins in field-grown cotton. *Plant Physiol.* 78:394–98
  46. Colombo SJ, Timmer VR, Colclough ML, Blumwald E. 1995. Diurnal variation in heat tolerance and heat shock protein expression in black spruce (*Picea mariana*). *Can. J. Forest Res.* 25:369–75
  47. Morris GJ, Coulson G, Meyer MA, McLellan MR, Fuller BJ, et al. 1983. Cold shock—a widespread cellular reaction. *Cryo-Letters* 4:179–92
  48. Danyluk J, Rassart E, Sarhan F. 1991. Gene expression during cold and heat shock in wheat. *Biochem. Cell Biol.* 69:383–91
  49. Krishna P, Sacco M, Cherutti JF, Hill S. 1995. Cold-induced accumulation of hsp90 transcripts in *Brassica napus*. *Plant Physiol.* 107:915–23
  50. Neven LG, Haskell DW, Guy CL, Denslow N, Klein PA, et al. 1992. Association of 70-kilodalton heat-shock cognate proteins with acclimation to cold. *Plant Physiol.* 99:1362–69
  51. Van Berkel J, Salamini F, Gebhardt C. 1994. Transcripts accumulating during cold storage of potato (*Solanum tuberosum* L.) tubers are sequence related to stress-responsive genes. *Plant Physiol.* 104:445–52
  52. Layne JR. 1991. Microclimate variability and the eurythermic natural of goldenrod gall fly (*Eurosta solidaginis*) larvae (Diptera: Tephritidae). *Can. J. Zool.* 69:614–17
  53. Feder ME, Blair N, Figueras H. 1997. Natural thermal stress and heat-shock protein expression in *Drosophila* larvae and pupae. *Funct. Ecol.* 11:90–100
  54. Feder ME. 1997. Necrotic fruit: a novel model system for thermal ecologists. *J. Therm. Biol.* 22:1–9
  55. Wehner R, Marsh AC, Wehner S. 1992. Desert ants on a thermal tightrope. *Nature* 357:586–87
  56. Gehring WJ, Wehner R. 1995. Heat shock protein synthesis and thermotolerance in *Cataglyphis*, an ant from the Sahara desert. *Proc. Natl. Acad. Sci. USA* 92:2994–98
  57. Locke M, Noble EG. 1995. Stress proteins: the exercise response. *Can. J. Appl. Physiol.* 20:155–67
  58. Brown IR, Rush SJ. 1996. In vivo activation of neural heat shock transcription factor HSF1 by a physiologically relevant increase in body temperature. *J. Neurosci. Res.* 44:52–57
  59. Di YP, Repasky EA, Subjeck JR. 1997. Distribution of HSP70, protein kinase C, and spectrin is altered in lymphocytes during a fever-like hyperthermia exposure. *J. Cell. Physiol.* 172:44–54
  60. Joplin KH, Denlinger DL. 1990. Developmental and tissue specific control of the heat shock induced 70 kDa related proteins in the flesh fly, *Sarcophaga crassipalpis*. *J. Insect Physiol.* 36:239–49
  61. Joplin KH, Yocum GD, Denlinger DL. 1990. Cold shock elicits expression of heat shock proteins in the flesh fly *Sarcophaga crassipalpis*. *J. Insect Physiol.* 36:825–34
  62. Yocum GD, Joplin KH, Denlinger DL. 1991. Expression of heat shock proteins

- in response to high and low temperature extremes in diapausing pharate larvae of the gypsy moth *Lymantria dispar*. *Arch. Insect Biochem. Physiol.* 18:239–50
63. Lee RE, Dommel RA, Joplin KH, Denlinger DL. 1995. Cryobiology of the freeze-tolerant gall fly *Eurosta solidaginis*: overwintering energetics and heat shock proteins. *Climate Res.* 5:61–67
  64. Denlinger DL, Lee RE, Yocum GD, Kukal O. 1992. Role of chilling in the acquisition of cold tolerance and the capacitation to express stress proteins in diapausing pharate larvae of the gypsy moth *Lymantria dispar*. *Arch. Insect Biochem. Physiol.* 21:271–80
  65. Matz JM, LaVoi KP, Moen RJ, Blake MJ. 1996. Cold-induced heat shock protein expression in rat aorta and brown adipose tissue. *Physiol. Behav.* 60:1369–74
  66. Sills NS, Gorham DA, Carey HV. 1998. Stress protein expression in a mammalian hibernator. *FASEB J.* 12:A379
  67. Li GC, Nussenzweig A. 1996. Thermotolerance and heat shock proteins: possible involvement of Ku autoantigen in regulating Hsp70 expression. See Ref. 9, pp. 121–37
  68. Alamillo J, Almoguera C, Bartels D, Jordano J. 1995. Constitutive expression of small heat shock proteins in vegetative tissues of the resurrection plant *Craterostigma plantagineum*. *Plant Mol. Biol.* 29:1093–99
  69. Pareek A, Singla SL, Kush AK, Grover A. 1997. Distribution patterns of HSP90 protein in rice. *Plant Sci.* 125:221–30
  70. Ma E, Haddad GG. 1997. Anoxia regulates gene expression in the central nervous system of *Drosophila melanogaster*. *Brain Res. Mol. Brain Res.* 46:325–28
  71. Kultz D. 1996. Plasticity and stressor specificity of osmotic and heat shock responses of *Gillichthys mirabilis* gill cells. *Am. J. Physiol.* 271:C1181–93
  72. de Pomerai D. 1996. Heat-shock proteins as biomarkers of pollution. *Hum. Exp. Toxicol.* 15:279–85
  73. Sanders BM, Dyer SD. 1994. Cellular stress response. *Environ. Toxicol. Chem.* 13:1209–10
  74. Sanders BM. 1993. Stress proteins in aquatic organisms: an environmental perspective. *Crit. Rev. Toxicol.* 23:49–75
  75. Ryan JA, Hightower LE. 1996. Stress proteins as molecular biomarkers for environmental toxicology. See Ref. 9, pp. 411–24
  76. Mueller WEG, Koziol C, Kurelec B, Dapper J, Batel R, Rinkevich B. 1995. Combinatory effects of temperature stress and nonionic organic pollutants on stress protein (hsp70) gene expression in the freshwater sponge *Ephydatia fluviatilis*. *Environ. Toxicol. Chem.* 14:1203–8
  77. Cochrane BJ, Irby RB, Snell TW. 1991. Effects of copper and tributyltin on stress protein abundance in the rotifer *Brachionus plicatilis*. *Comp. Biochem. Physiol.* 98C:385–90
  78. Krasko A, Scheffer U, Koziol C, Pancer Z, Batel R, et al. 1997. Diagnosis of sublethal stress in the marine sponge *Geodia cydonium*: application of the 70 kDa heat-shock protein and a novel biomarker, the Rab GDP dissociation inhibitor, as probes. *Aquat. Toxicol.* 37:157–68
  79. Werner I, Nagel R. 1997. Stress proteins HSP60 and HSP70 in 3 species of amphipods exposed to cadmium, diazinon, dieldrin and fluoranthene. *Environ. Toxicol. Chem.* 16:2393–403
  80. Ruffin P, Demuynck S, Hilbert JL, Dhainaut A. 1994. Stress protein in the polychaete annelid *Nereis diversicolor* induced by heat shock or cadmium exposure. *Biochimie* 76:423–27
  81. Steinert SA, Pickwell GV. 1988. Expression of heat shock proteins and metallothionein in mussels exposed to heat stress and metal ion challenge. *Mar. Environ. Res.* 24:211–14
  82. Veldhuizen Tsoerkan MB, Holwerda DA, van der Mast CA, Zandee DI. 1991. Synthesis of stress proteins under normal and heat shock conditions in gill tissue of sea mussels (*Mytilus edulis*) after chronic exposure to cadmium. *Comp. Biochem. Physiol.* 100C:699–706
  83. Nascimento IA, Dickson KL, Zimmerman EG. 1996. Heat shock protein response to thermal stress in the Asiatic clam, *Corbicula fluminea*. *J. Aquat. Ecosystem Health* 5:231–38
  84. Sanders BM, Martin LS, Howe SR, Nelson WG, Hegre ES, Phelps DK. 1994. Tissue-specific differences in accumulation of stress proteins in *Mytilus edulis* exposed to a range of copper concentrations. *Toxicol. Appl. Pharmacol.* 125:206–13
  85. Ryan JA, Hightower LE. 1994. Evaluation of heavy-metal ion toxicity in fish cells using a combined stress protein and cytotoxicity assay. *Environ. Toxicol. Chem.* 13:1231–40
  86. Dyer SD, Brooks GL, Dickson KL, Sanders BM, Zimmerman EG. 1993. Synthesis and accumulation of stress

- proteins in tissues of arsenite-exposed fathead minnows *Pimephales promelas*. *Environ. Toxicol. Chem.* 12:913–24
87. Vijayan MM, Pereira C, Forsyth RB, Kennedy CJ, Iwama GK. 1997. Handling stress does not affect the expression of hepatic heat shock protein 70 and conjugation enzymes in rainbow trout treated with beta-naphthoflavone. *Life Sci.* 61:117–27
  88. Van Dyk TK, Majarian WR, Konstantinov KB, Young RM, Dhurjati PS, LaRossa RA. 1994. Rapid and sensitive pollutant detection by induction of heat shock gene-bioluminescence gene fusions. *Appl. Environ. Microbiol.* 60:1414–20
  89. Veldhuizen Tsoerkan MB, Holwerda DA, de Bont AM, Smaal AC, Zandee DI. 1991. A field study on stress indices in the sea mussel, *Mytilus edulis*: application of the "stress approach" in biomonitoring. *Arch. Environ. Contam. Toxicol.* 21:497–504
  90. Kohler HR, Triebkorn R, Stocker W, Kloetzel PM, Alberti G. 1992. The 70 kD heat shock protein (hsp 70) in soil invertebrates: a possible tool for monitoring environmental toxicants. *Arch. Environ. Contam. Toxicol.* 22:334–38
  91. Pyza E, Mak P, Kramarz P, Laskowski R. 1997. Heat-shock proteins (Hsp70) as biomarkers in ecotoxicological studies. *Ecotoxicol. Environ. Safety* 38:244–51
  92. Stringham EG, Candido EPM. 1994. Transgenic hsp16-lacZ strains of the soil nematode *Caenorhabditis elegans* as biological monitors of environmental stress. *Environ. Toxicol. Chem.* 13:1211–20
  93. Forsyth RB, Candido EPM, Babich SL, Iwama GK. 1997. Stress protein expression in coho salmon with bacterial kidney disease. *J. Aquat. Anim. Health* 9:18–25
  94. Rose MR, Lauder GV, eds. 1996. *Adaptation*. New York/London: Academic
  95. Bosch TC, Krylow SM, Bode HR, Steele RE. 1988. Thermotolerance and synthesis of heat shock proteins: These responses are present in *Hydra attenuata* but absent in *Hydra oligactis*. *Proc. Natl. Acad. Sci. USA* 85:7927–31
  96. Gellner K, Praetzel G, Bosch TC. 1992. Cloning and expression of a heat-inducible hsp70 gene in two species of *Hydra* which differ in their stress response. *Eur. J. Biochem.* 210:683–91
  97. Hofmann GE, Somero GN. 1996. Interspecific variation in thermal denaturation of proteins in the congeneric mussels *Mytilus trossulus* and *M. galloprovincialis*: evidence from the heat-shock response and protein ubiquitination. *Mar. Biol.* 126:65–75
  98. Tomanek L, Somero GN. 1997. The effect of temperature on protein synthesis in snails of the genus *Tegula* from the sub- and intertidal zone. *Am. Zool.* 37:188A
  99. Vayda ME, Yuan ML. 1994. The heat shock response of an Antarctic alga is evident at 5 degrees C. *Plant Mol. Biol.* 24:229–33
  100. Berg GR, Inniss WE, Heikkila JJ. 1987. Stress proteins and thermotolerance in psychrotrophic yeasts from Arctic environments. *Can. J. Microbiol.* 33:383–89
  101. Deegenars ML, Watson K. 1997. Stress proteins and stress tolerance in an Antarctic, psychrophilic yeast, *Candida psychrophila*. *FEMS Microbiol. Lett.* 151:191–96
  102. Carratu L, Maresca B. 1997. Evolutionary adaptation of hsp70 gene in Antarctic fish. *Exp. Biol. Online* 2:C5.1 (Abstr.)
  103. Roberts DA, Hofmann GE, Somero GN. 1997. Heat-shock protein expression in *Mytilus californianus*: acclimatization (seasonal and tidal-height comparisons) and acclimation effects. *Biol. Bull.* 192:309–20
  104. Norris CE, diIorio PJ, Schultz RJ, Hightower LE. 1995. Variation in heat shock proteins within tropical and desert species of poeciliid fishes. *Mol. Biol. Evol.* 12:1048–62
  105. White CN, Hightower LE, Schultz RJ. 1994. Variation in heat-shock proteins among species of desert fishes (Poeciliidae, *Poeciliopsis*). *Mol. Biol. Evol.* 11:106–19
  - 105a. Maresca B, Patriarcha E, Goldenberg C, Sacco M. 1988. Heat shock and cold adaptation in Antarctic fishes: a molecular approach. *Comp. Biochem. Physiol.* 90B:623–29
  106. Ristic Z, Williams G, Yang G, Martin B, Fullerton S. 1996. Dehydration, damage to cellular membranes, and heat-shock proteins in maize hybrids from different climates. *J. Plant Physiol.* 149:424–32
  107. Ulmasov KA, Shammakov S, Karaev K, Evgenev MB. 1992. Heat shock proteins and thermoresistance in lizards. *Proc. Natl. Acad. Sci. USA* 89:1666–70
  108. Konstantopoulou I, Drosopoulou E, Scouras ZG. 1997. Variations in the heat-induced protein pattern of several *Drosophila montium* subgroup species (Diptera: Drosophilidae). *Genome* 40:132–37

109. Feder ME, Cartaño NV, Milos L, Krebs RA, Lindquist SL. 1996. Effect of engineering *Hsp70* copy number on Hsp70 expression and tolerance of ecologically relevant heat shock in larvae and pupae of *Drosophila melanogaster*. *J. Exp. Biol.* 199:1837–44
110. Nath BB, Lakhotia SC. 1989. Heat-shock response in a tropical *Chironomus*: seasonal variation in response and the effect of developmental stage and tissue type on heat shock protein synthesis. *Genome* 32:676–86
111. Hofmann GE. 1996. Molecular chaperone activity of the stress protein Hsc70 purified from an eurythermal goby, *Gillichthys mirabilis*. *Am. Zool.* 36:36A
112. McFall-Ngai M, Horwitz J. 1990. A comparative study of the thermal stability of the vertebrate eye lens: Antarctic fish to the desert iguana. *Exp. Eye Res.* 50:703–9
113. Clos J, Rabindran S, Wisniewski J, Wu C. 1993. Induction temperature of human heat shock factor is reprogrammed in a *Drosophila* cell environment. *Nature* 364:252–55
114. Carratu L, Franceschelli S, Pardini CL, Kobayashi GS, Horvath I, et al. 1996. Membrane lipid perturbation modifies the set point of the temperature of heat shock response in yeast. *Proc. Natl. Acad. Sci. USA* 93:3870–75
115. Maresca B, Kobayashi GS. 1994. Hsp70 in parasites: as an inducible protective protein and as an antigen. *Experientia* 50:1067–74
116. Kaufmann SH. 1992. The cellular immune response to heat shock proteins. *Experientia* 48:640–43
117. Polla BS. 1991. Heat shock proteins in host-parasite interactions. *Immunol. Today* 12:A38–41
118. Newport GR. 1991. Heat shock proteins as vaccine candidates. *Semin. Immunol.* 3:17–24
119. Tsuji N, Ohta M, Fujisaki K. 1997. Expression of a 70-kDa heat-shock-related protein during transformation from free-living infective larvae to the parasitic stage in *Strongyloides venezuelensis*. *Parasitol. Res.* 83:99–102
120. Van der Ploeg LH, Giannini SH, Cantor CR. 1985. Heat shock genes: regulatory role for differentiation in parasitic protozoa. *Science* 228:1443–46
121. Neumann S, Ziv E, Lantner F, Schechter I. 1993. Regulation of HSP70 gene expression during the life cycle of the parasitic helminth *Schistosoma mansoni*. *Eur. J. Biochem.* 212:589–96
122. van Leeuwen MA. 1995. Heat-shock and stress response of the parasitic nematode *Haemonchus contortus*. *Parasitol. Res.* 81:706–9
123. Ernani FP, Teale JM. 1993. Release of stress proteins from *Mesocostoides corti* is a brefeldin A-inhibitable process: evidence for active export of stress proteins. *Infect. Immun.* 61:2596–601
124. Syin C, Goldman ND. 1996. Cloning of a *Plasmodium falciparum* gene related to the human 60-kDa heat shock protein. *Mol. Biochem. Parasitol.* 79:13–19
125. Cluss RG, Boothby JT. 1990. Thermoregulation of protein synthesis in *Borrelia burgdorferi*. *Infect. Immun.* 58:1038–42
126. Hubel A, Krobitch S, Horauf A, Clos J. 1997. The *Leishmania major* Hsp100 is required chiefly in the mammalian stage of the parasite. *Mol. Cell. Biol.* 17:5987–95
127. Giambiagi-de Marval M, Souto-Pradon T, Rondinelli E. 1996. Characterization and cellular distribution of heat-shock proteins HSP70 and HSP60 in *Trypanosoma cruzi*. *Exp. Parasitol.* 83:335–45
128. Daubenberger C, Heussler V, Gobright E, Wijngaard P, Clevers HC, et al. 1997. Molecular characterisation of a cognate 70 kDa heat shock protein of the protozoan *Theileria parva*. *Mol. Biochem. Parasitol.* 85:265–69
129. Shearer GJ, Birge CH, Yuckenberg PD, Kobayashi GS, Medoff G. 1987. Heat-shock proteins induced during the mycelial-to-yeast transitions of strains of *Histoplasma capsulatum*. *J. Gen. Microbiol.* 133:3375–82
130. Maresca B. 1995. Unraveling the secrets of *Histoplasma capsulatum*. A model to study morphogenic adaptation during parasite host/host interaction. *Verh. K. Acad. Geneesk. Belg.* 57:133–56
131. Clark TG, Abrahamson MS, White MW. 1996. Developmental expression of heat shock protein 90 in *Eimeria bovis*. *Mol. Biochem. Parasitol.* 78:259–63
132. Tirard CT, Grossfeld RM, Volety AK, Chu FLE. 1995. Heat shock proteins of the oyster parasite *Perkinsus marinus*. *Dis. Aquat. Organisms* 22:147–51
133. Ulmasov KA, Ovezmukhammedov A, Karaev KK, Evgenev MB. 1988. Molecular mechanisms of adaptation to hyperthermia in higher organisms. III. Induction of heat-shock proteins in two *Leishmania* species. *Mol. Biol.* 22:1583–89

134. Ryan MT, Naylor DJ, Hoj PB, Clark MS, Hoogenraad NJ. 1997. The role of molecular chaperones in mitochondrial protein import and folding. *Int. Rev. Cytol.* 174:127–93
135. Germot A, Philippe H, Le Guyader H. 1997. Evidence for loss of mitochondria in microsporidia from a mitochondrial-type HSP70 in *Nosema locustae*. *Mol. Biochem. Parasitol.* 87:159–68
136. Bui ET, Bradley PJ, Johnson PJ. 1996. A common evolutionary origin for mitochondria and hydrogenosomes. *Proc. Natl. Acad. Sci. USA* 93:9651–56
137. Sogin ML. 1997. Organelle origins: energy-producing symbionts in early eukaryotes? *Curr. Biol.* 7:R315–17
138. Hofmann CJ, Rensing SA, Hauber MM, Martin WF, Muller SB, et al. 1994. The smallest known eukaryotic genomes encode a protein gene: towards an understanding of nucleomorph functions. *Mol. Gen. Genet.* 243:600–4
139. Germot A, Philippe H, Le Guyader H. 1996. Presence of a mitochondrial-type 70-kDa heat shock protein in *Trichomonas vaginalis* suggests a very early mitochondrial endosymbiosis in eukaryotes. *Proc. Natl. Acad. Sci. USA* 93:14614–17
140. Baumann P, Moran NA, Baumann L. 1997. The evolution and genetics of aphid endosymbionts. *BioScience* 47:12–20
141. Aksoy S. 1995. Molecular analysis of the endosymbionts of tsetse flies: 16S rDNA locus and over-expression of a chaperonin. *Insect Mol. Biol.* 4:23–29
142. Moran NA, Von Dohlen CD, Baumann P. 1995. Faster evolutionary rates in endosymbiotic bacteria than in cospeciating insect hosts. *J. Mol. Evol.* 41:727–31
143. Moran NA. 1996. Accelerated evolution and Muller's ratchet in endosymbiotic bacteria. *Proc. Natl. Acad. Sci. USA* 93:2873–78
144. Morioka M, Ishikawa H. 1992. Mutualism based on stress: selective synthesis and phosphorylation of a stress protein by an intracellular symbiont. *J. Biochem.* 111:431–35
145. Hong HK, Choi JY, Ahn TI. 1994. Molecular biological studies on the heat-shock responses in *Amoeba proteus*: I. Detection of heat-shock proteins. *Korean J. Zool.* 37:554–64
146. Choi EY, Ahn GS, Jeon KW. 1991. Elevated levels of stress proteins associated with bacterial symbiosis in *Amoeba proteus* and soybean root nodule cells. *Biosystems* 25:205–12
147. Feder ME, Karr TL. 1997. Evolutionarily significant consequences of the heat shock response for *Drosophila* and its endosymbiont *Wolbachia*. *Am. Zool.* 37:8A
148. Dix DJ. 1997. Hsp70 expression and function during gametogenesis. *Cell Stress Chaperones* 2:73–77
149. Lin JC, Song CW. 1993. Heat shock gene expression and development. I. An overview of fungal, plant, and poikilothermic animal developmental systems. *Dev. Genet.* 14:1–5
150. Winter J, Sinibaldi R. 1991. The expression of heat shock protein and cognate genes during plant development. *Results Prob. Cell Differ.* 17:83–105
151. Mosser DD, Duchaine J, Bourget L, Martin LH. 1993. Heat shock gene expression and development. II. An overview of mammalian and avian developmental systems. *Dev. Genet.* 14:87–91
152. Heikkila JJ, Ohan N, Tam Y, Ali A. 1997. Heat shock protein gene expression during *Xenopus* development. *Cell. Mol. Life Sci.* 53:114–21
153. Muller WU, Li GC, Goldstein LS. 1985. Heat does not induce synthesis of heat shock proteins or thermotolerance in the earliest stage of mouse embryo development. *Int. J. Hypertherm.* 1:97–102
154. Edwards JL, Ealy AD, Monterroso VH, Hansen PJ. 1997. Ontogeny of temperature-regulated heat shock protein 70 synthesis in preimplantation bovine embryos. *Mol. Reprod. Dev.* 48:25–33
155. Dura JM. 1981. Stage dependent synthesis of heat shock induced proteins in early embryos of *Drosophila melanogaster*. *Mol. Gen. Genet.* 184:381–85
156. Welte MA, Tetrault JM, Dellavalle RP, Lindquist SL. 1993. A new method for manipulating transgenes: engineering heat tolerance in a complex, multicellular organism. *Curr. Biol.* 3:842–53
157. Gagliardi D, Breton C, Chaboud A, Vergne P, Dumas C. 1995. Expression of heat shock factor and heat shock protein 70 genes during maize pollen development. *Plant Mol. Biol.* 29:841–56
158. Hendrey J, Kola I. 1991. Thermolability of mouse oocytes is due to the lack of expression and/or inducibility of Hsp70. *Mol. Reprod. Dev.* 28:1–8
159. Curci A, Bevilacqua A, Mangia F. 1987. Lack of heat-shock response in pre-ovulatory mouse oocytes. *Dev. Biol.* 123:154–60

160. Curci A, Bevilacqua A, Fiorenza MT, Mangia F. 1991. Developmental regulation of heat-shock response in mouse oogenesis: identification of differentially responsive oocyte classes during Graafian follicle development. *Dev. Biol.* 144:362-68
161. Zakeri ZF, Welch WJ, Wolgemuth DJ. 1990. Characterization and inducibility of hsp 70 proteins in the male mouse germ line. *J. Cell Biol.* 111:1785-92
162. Bedard PA, Brandhorst BP. 1986. Translational activation of maternal mRNA encoding the heat-shock protein hsp90 during sea urchin embryogenesis. *Dev. Biol.* 117:286-93
163. Gordon S, Bharadwaj S, Hnatov A, Ali A, Ovsenek N. 1997. Distinct stress-inducible and developmentally regulated heat shock transcription factors in *Xenopus* oocytes. *Dev. Biol.* 181:47-63
164. Feder JH, Rossi JM, Solomon J, Solomon N, Lindquist S. 1992. The consequences of expressing Hsp70 in *Drosophila* cells at normal temperatures. *Genes Dev.* 6:1402-13
165. Mitchell HK, Moller G, Petersen NS, Lipps-Sarmiento L. 1979. Specific protection from phenocopy induction by heat shock. *Dev. Genet.* 1:181-92
166. Welte MA, Duncan I, Lindquist S. 1995. The basis for a heat-induced developmental defect: defining crucial lesions. *Genes Dev.* 9:2240-50
167. Wehmeyer N, Hernandez LD, Finkelstein RR, Vierling E. 1996. Synthesis of small heat-shock proteins is part of the developmental program of late seed maturation. *Plant Physiol.* 112:747-57
168. Duck N, McCormick S, Winter J. 1989. Heat shock protein Hsp70 cognate gene expression in vegetative and reproductive organs of *Lycopersicon esculentum*. *Proc. Natl. Acad. Sci. USA* 86:3674-78
169. Coca MA, Almoguera C, Jordano J. 1994. Expression of sunflower low-molecular-weight heat-shock proteins during embryogenesis and persistence after germination: localization and possible functional implications. *Plant Mol. Biol.* 25:479-92
170. Helm KW, Petersen NS, Abernethy RH. 1989. Heat-shock response of germinating embryos of wheat: effects of imbibition time and seed vigor. *Plant Physiol.* 90:598-605
171. Sanchez Y, Taulien J, Borkovich KA, Lindquist S. 1992. Hsp104 is required for tolerance to many forms of stress. *EMBO J.* 11:2357-64
172. Silva AM, Juliani MH, da Costa JJ, Bonato MC. 1987. Acquisition of thermotolerance during development of *Blastocladiella emersonii*. *Biochem. Biophys. Res. Commun.* 144:491-98
173. Liang P, Amons R, Clegg JS, MacRae TH. 1997. Molecular characterization of a small heat shock/alpha-crystallin protein in encysted *Artemia* embryos. *J. Biol. Chem.* 272:19051-58
174. Liang P, Amons R, Macrae TH, Clegg JS. 1997. Purification, structure and in vitro molecular-chaperone activity of *Artemia* p26, a small heat-shock/alpha-crystallin protein. *Eur. J. Biochem.* 243:225-32
175. Jackson SA, Clegg JS. 1996. Ontogeny of low molecular weight stress protein p26 during early development of the brine shrimp, *Artemia franciscana*. *J. Exp. Biol.* 200:467-75
176. Clegg JS, Jackson SA, Liang P, MacRae TH. 1995. Nuclear-cytoplasmic translocations of protein p26 during aerobic-anoxic transitions in embryos of *Artemia franciscana*. *Exp. Cell Res.* 219:1-7
177. Clegg JS, Jackson SA, Warner AH. 1994. Extensive intracellular translocations of a major protein accompany anoxia in embryos of *Artemia franciscana*. *Exp. Cell Res.* 212:77-83
178. Clegg JS, Jackson SA. 1992. Aerobic heat shock activates trehalose synthesis in embryos of *Artemia franciscana*. *FEBS Lett.* 303:45-47
179. Anchordoguy TJ, Hand SC. 1994. Acute blockage of the ubiquitin-mediated proteolytic pathway during invertebrate quiescence. *Am. J. Physiol.* 267:R895-900
180. Krebs RA, Feder ME, Lee J. 1998. Heritability of expression of the 70-kD heat-shock protein in *Drosophila melanogaster* and its relevance to the evolution of thermotolerance. *Evolution* 52:841-47
181. Davis WL, Jacoby BH, Goodman DB. 1994. Immunolocalization of ubiquitin in degenerating insect flight muscle. *Histochem. J.* 26:298-305
182. Sarge KD, Bray AE, Goodson ML. 1995. Altered stress response in testis. *Nature* 374:126
183. Sarge KD. 1995. Male germ cell-specific alteration in temperature set point of the cellular stress response. *J. Biol. Chem.* 270:18745-48
184. Holbrook NJ, Udelsman R. 1994. Heat shock protein gene expression in response to physiologic stress and aging. See Ref. 4, pp. 577-93
185. Lee YK, Manalo D, Liu AY. 1996. Heat shock response, heat shock transcrip-

- tion factor and cell aging. *Biol. Signals* 5:180–91
186. Shpund S, Gershon D. 1997. Alterations in the chaperone activity of HSP70 in aging organisms. *Arch. Gerontol. Geriatr.* 24:125–31
  187. Wheeler JC, Bieschke ET, Tower J. 1995. Muscle-specific expression of *Drosophila* hsp70 in response to aging and oxidative stress. *Proc. Natl. Acad. Sci. USA* 92:10408–12
  188. Marin R, Valet JP, Tanguay RM. 1993. Heat shock induces changes in the expression and binding of ubiquitin in senescent *Drosophila melanogaster*. *Dev. Genet.* 14:78–86
  189. Lithgow GJ, White TM, Hinerfeld DA, Johnson TE. 1994. Thermotolerance of a long-lived mutant of *Caenorhabditis elegans*. *J. Gerontol.* 49B:270–76
  190. Lithgow GJ, White TM, Melov S, Johnson TE. 1995. Thermotolerance and extended life-span conferred by single-gene mutations and induced by thermal stress. *Proc. Natl. Acad. Sci. USA* 92:7540–44
  191. Lithgow GJ. 1996. Invertebrate gerontology: the age mutations of *Caenorhabditis elegans*. *BioEssays* 18:809–15
  192. Bond JA, Gonzalez CRM, Bradley BP. 1993. Age-dependent expression of proteins in the cladoceran *Daphnia magna* under normal and heat-stress conditions. *Comp. Biochem. Physiol.* 106B:913–17
  193. Khazaeli AA, Tatar M, Pletcher SD, Curtsinger JW. 1997. Heat-induced longevity extension in *Drosophila*. I. Heat treatment, mortality, and thermotolerance. *J. Gerontol.* 52A:B48–52
  194. Tatar M, Khazaeli AA, Curtsinger JW. 1997. Chaperoning extended life. *Nature* 390:30
  195. Dhahbi JM, Mote PL, Tillman JB, Walford RL, Spindler SR. 1997. Dietary energy tissue-specifically regulates endoplasmic reticulum chaperone gene expression in the liver of mice. *J. Nutr.* 127:1758–64
  196. Heydari AR, Conrad CC, Richardson A. 1995. Expression of heat shock genes in hepatocytes is affected by age and food restriction in rats. *J. Nutr.* 125:410–18
  197. Pahlavani MA, Harris MD, Moore SA, Richardson A. 1996. Expression of heat shock protein 70 in rat spleen lymphocytes is affected by age but not by food restriction. *J. Nutr.* 126:2069–75
  198. Lu Q, Wallrath LL, Granok H, Elgin SC. 1993. Expression of heat shock protein 70 is altered by age and diet at the level of transcription. *Mol. Cell Biol.* 13:2909–18
  199. Masoro EJ. 1996. Possible mechanisms underlying the antiaging actions of caloric restriction. *Toxicol. Pathol.* 24:738–41
  200. Parsell DA, Lindquist S. 1993. The function of heat-shock proteins in stress tolerance: degradation and reactivation of damaged proteins. *Annu. Rev. Genet.* 27:437–96
  201. Parsell DA, Lindquist S. 1994. Heat shock proteins and stress tolerance. See Ref. 9, pp. 457–94
  202. Bensaude O, Bellier S, Dubois MF, Giannoni F, Nguyen VT. 1996. Heat-shock induced protein modifications and modulation of enzyme activities. See Ref. 9, pp. 199–219
  203. dilorjo PJ, Holsinger K, Schultz RJ, Hightower LE. 1996. Quantitative evidence that both Hsc70 and Hsp70 contribute to thermal adaptation in hybrids of the livebearing fishes *Poeciliopsis*. *Cell Stress Chaperones* 1:139–47
  204. Feder ME, Lindquist SL. 1992. Evolutionary loss of a heat shock protein. *Am. Zool.* 32:51A (Abstr.)
  205. Whitesell L, Cook P. 1996. Stable and specific binding of heat shock protein 90 by geldanamycin disrupts glucocorticoid receptor function in intact cells. *Mol. Endocrinol.* 10:705–12
  206. Elia G, Santoro MG. 1994. Regulation of heat shock protein synthesis by quercetin in human erythroleukaemia cells. *Biochem. J.* 300:201–9
  207. Bonham-Smith PC, Kapoor M, Bewley JD. 1987. Establishment of thermotolerance in maize by exposure to stresses other than a heat shock does not require heat shock protein synthesis. *Plant Physiol.* 85:575–80
  208. Xiao CM, Mascarenhas JP. 1985. High temperature-induced thermotolerance in pollen tubes of *Tradescantia* and heat-shock proteins. *Plant Physiol.* 78:887–90
  209. VanBogelen RA, Acton MA, Neidhardt FC. 1987. Induction of the heat shock regulon does not produce thermotolerance in *Escherichia coli*. *Genes Dev.* 1:525–31
  210. Yocum GD, Denlinger DL. 1992. Prolonged thermotolerance in the flesh fly *Sarcophaga crassipalpis* does not require continuous expression or persistence of the 72 kDa heat-shock protein. *J. Insect Physiol.* 38:603–9
  211. Boon-Niermeijer EK, Tuyl M, van de Scheur H. 1986. Evidence for two states

- of thermotolerance. *Int. J. Hypertherm.* 2:93-105
212. Smith BJ, Yaffe MP. 1991. Uncoupling thermotolerance from the induction of heat shock proteins. *Proc. Natl. Acad. Sci. USA* 88:11091-94
  213. Easton DP, Rutledge PS, Spotila JR. 1987. Heat shock protein induction and induced thermal tolerance are independent in adult salamanders. *J. Exp. Zool.* 241:263-67
  214. Dingley F, Maynard Smith J. 1968. Temperature acclimatization in the absence of protein synthesis of *Drosophila subobscura*. *J. Insect Physiol.* 14:1185-94
  215. Finnell RH, Van Waes M, Bennett GD, Eberwine JH. 1993. Lack of concordance between heat shock proteins and the development of tolerance to teratogen-induced neural tube defects. *Dev. Genet.* 14:137-47
  216. Fisher B, Kraft P, Hahn GM, Anderson RL. 1992. Thermotolerance in the absence of induced heat shock proteins in a murine lymphoma. *Cancer Res.* 52:2854-61
  217. Watson K, Dunlop G, Cavicchioli R. 1984. Mitochondrial and cytoplasmic protein syntheses are not required for heat shock acquisition of ethanol and thermotolerance in yeast. *FEBS Lett.* 172:299-302
  218. Widelitz RB, Magun BE, Gerner EW. 1986. Effects of cycloheximide on thermotolerance expression, heat shock protein synthesis, and heat shock protein mRNA accumulation in rat fibroblasts. *Mol. Cell Biol.* 6:1088-94
  219. Jozwiak Z, Leyko W. 1992. Role of membrane components in thermal injury of cells and development of thermotolerance. *Int. J. Radiat. Biol.* 62:743-56
  220. Lindquist S. 1993. Autoregulation of the heat-shock response. In *Translational Regulation of Gene Expression 2*, ed. J Ilan, pp. 279-320. New York: Plenum
  221. Krebs RA, Feder ME. 1998. Hsp70 and larval thermotolerance in *Drosophila melanogaster*: How much is enough and when is more too much? *J. Insect Physiol.* 44:1091-1101
  222. Krebs RA, Feder ME. 1997. Deleterious consequences of Hsp70 overexpression in *Drosophila melanogaster* larvae. *Cell Stress Chaperones* 2:60-71
  223. Krebs RA, Feder ME. 1997. Natural variation in the expression of the heat-shock protein Hsp70 in a population of *Drosophila melanogaster*, and its correlation with tolerance of ecologically relevant thermal stress. *Evolution* 51:173-79
  224. Welte MA. 1994. *Thermotolerance in Drosophila embryos: the role of hsp70 and the basis for a specific phenocopy*. Ph.D. thesis. Univ. Chicago. 230 pp.
  225. Krebs RA, Feder ME. 1998. Experimental manipulation of the cost of thermal acclimation in *Drosophila melanogaster*. *Biol. J. Linn. Soc.* 63:593-601
  226. Hoffmann AA. 1995. Acclimation: increasing survival at a cost. *Trends Ecol. Evol.* 10:1-2
  227. Calow P. 1991. Physiological costs of combating chemical toxicants: ecological implications. *Comp. Biochem. Physiol.* 100C:3-6
  228. Koehn RK, Bayne BL. 1989. Towards a physiological and genetical understanding of the energetics of the stress response. *Biol. J. Linn. Soc.* 37:157-71
  229. Dorner AJ, Krane MG, Kaufman RJ. 1988. Reduction of endogenous GRP78 levels improves secretion of a heterologous protein in CHO cells. *Mol. Cell Biol.* 8:4063-70
  230. Dorner AJ, Wasley LC, Kaufman RJ. 1992. Overexpression of GRP78 mitigates stress induction of glucose regulated proteins and blocks secretion of selective proteins in Chinese hamster ovary cells. *EMBO J.* 11:1563-71
  231. Dorner AJ, Kaufman RJ. 1994. The levels of endoplasmic reticulum proteins and ATP affect folding and secretion of selective proteins. *Biologicals* 22:103-12
  232. Ryan C, Stevens TH, Schlesinger MJ. 1992. Inhibitory effects of HSP70 chaperones on nascent polypeptides. *Protein Sci.* 1:980-85
  233. Heckathorn SA, Poeller GJ, Coleman JS, Hallberg RL. 1996. Nitrogen availability alters patterns of accumulation of heat stress-induced proteins in plants. *Oecologia* 105:413-18
  234. Heckathorn SA, Poeller GJ, Coleman JS, Hallberg RL. 1996. Nitrogen availability and vegetative development influence the response of ribulose 1,5-bisphosphate carboxylase/oxygenase, phosphoenolpyruvate carboxylase, and heat-shock protein content to heat stress in *Zea mays* L. *Int. J. Plant Sci.* 157:588-95
  235. Dietz TJ, Somero GN. 1993. Species- and tissue-specific synthesis patterns for heat-shock proteins hsp70 and hsp90 in several marine teleost fishes. *Physiol. Zool.* 66:863-80

236. Abravaya K, Phillips B, Morimoto RI. 1991. Attenuation of the heat shock response in HeLa cells is mediated by the release of bound heat shock transcription factor and is modulated by changes in growth and in heat shock temperatures. *Genes Dev.* 5:2117-27
237. Sarge KD, Cullen KE. 1997. Regulation of hsp expression during rodent spermatogenesis. *Cell. Mol. Life Sci.* 53:191-97
238. Koban M, Graham G, Prosser CL. 1987. Induction of heat-shock protein synthesis in teleost hepatocytes: effects of acclimation temperature. *Physiol. Zool.* 60:290-96
239. Deleted in proof
240. Favatier F, Bornman L, Hightower LE, Gnther E, Polla BS. 1997. Variation in *hsp* gene expression and Hsp polymorphism: Do they contribute to differential disease susceptibility and stress tolerance? *Cell Stress Chaperones* 2: 141-55
241. Engman DM, Sias SR, Gabe JD, Donelson JE, Dragon EA. 1989. Comparison of HSP70 genes from two strains of *Trypanosoma cruzi*. *Mol. Biochem. Parasitol.* 37:285-87
242. Engman DM, Reddy LV, Donelson JE, Kirchoff LV. 1987. *Trypanosoma cruzi* exhibits inter- and intra-strain heterogeneity in molecular karyotype and chromosomal gene location. *Mol. Biochem. Parasitol.* 22:115-23
243. Snutch TP, Baillie DL. 1984. A high degree of DNA strain polymorphism associated with the major heat shock gene in *Caenorhabditis elegans*. *Mol. Gen. Genet.* 195:329-35
244. Hamet P, Kaiser MA, Sun Y, Page V, Vincent M, et al. 1996. HSP27 locus cosegregates with left ventricular mass independently of blood pressure. *Hypertension* 28:1112-17
245. Wallich R, Helmes C, Schaible UE, Lobet Y, Moter SE, et al. 1992. Evaluation of genetic divergence among *Borrelia burgdorferi* isolates by use of OspA, fla, HSP60, and HSP70 gene probes. *Infect. Immun.* 60:4856-66
246. Hamet P, Kong D, Pravenec M, Kunes J, Kren V, et al. 1992. Restriction fragment length polymorphism of *hsp70* gene, localized in the RT1 complex, is associated with hypertension in spontaneously hypertensive rats. *Hypertension* 19:611-14
247. Grosz MD, Skow LC, Stone RT. 1994. An AluI polymorphism at the bovine 70 kD heat-shock protein-1 (HSP70-1) locus. *Anim. Genet.* 25:196
248. Jorgensen JA, Nguyen HT. 1995. Genetic analysis of heat shock proteins in maize. *Theor. Appl. Genet.* 91:38-46
249. Ottaviano E, Sari Gorla M, Pe E, Frova C. 1991. Molecular markers, RFLPs and Hsps for the genetic dissection of thermotolerance in maize. *Theor. Appl. Genet.* 81:713-19
250. Dimascio JA, Sweeney PM, Danneberger TK, Kamalay JC. 1994. Analysis of heat shock response in perennial ryegrass using maize heat shock protein clones. *Crop Sci.* 34:798-804
251. Goldschmidt-Clermont M. 1980. Two genes for the major heat-shock protein of *Drosophila melanogaster* arranged as an inverted repeat. *Nucleic Acids Res.* 8:235-52
252. Ish-Horowicz D, Pinchin SM. 1980. Genomic organization of the 87A7 and 87C1 heat-induced loci of *Drosophila melanogaster*. *J. Mol. Biol.* 142:231-45
253. Leigh-Brown AJ, Ish-Horowicz D. 1981. Evolution of the 87A and 87C heat-shock loci in *Drosophila*. *Nature* 290:677-82
254. McKechnie SW, Halford MM, McColl G, Hoffmann AA. 1998. Both allelic variation and expression of nuclear and cytoplasmic transcripts of *hsp-omega* are closely associated with thermal phenotype in *Drosophila*. *Proc. Natl. Acad. Sci. USA* 95:2423-28
255. Ish-Horowicz D, Pinchin SM, Schedl P, Artavanis-Tsakonas S, Mirault ME. 1979. Genetic and molecular analysis of the 87A7 and 87C1 heat-inducible loci of *D. melanogaster*. *Cell* 18:1351-58
256. Lis JT, Ish-Horowicz D, Pinchin SM. 1981. Genomic organization and transcription of the alpha beta heat shock DNA in *Drosophila melanogaster*. *Nucleic Acids Res.* 9:5297-310
257. Lis JT, Prestidge L, Hogness DS. 1978. A novel arrangement of tandemly repeated genes at a major heat shock site in *D. melanogaster*. *Cell* 14:901-19
258. Mirault ME, Goldschmidt-Clermont M, Artavanis-Tsakonas S, Schedl P. 1979. Organization of the multiple genes for the 70,000-dalton heat-shock protein in *Drosophila melanogaster*. *Proc. Natl. Acad. Sci. USA* 76:5254-58
259. Craig EA, McCarthy BJ, Wadsworth SC. 1979. Sequence organization of two recombinant plasmids containing genes for the major heat shock-induced protein of *D. melanogaster*. *Cell* 16:575-88
260. Artavanis-Tsakonas S, Schedl P, Mirault ME, Moran L, Lis J. 1979. Genes for the 70,000 dalton heat shock protein in two

- cloned *D. melanogaster* DNA segments. *Cell* 17:9–18
261. Holmgren R, Livak K, Morimoto R, Freund R, Meselson M. 1979. Studies of cloned sequences from four *Drosophila* heat shock loci. *Cell* 18:1359–70
  262. Sun CW, Griffen S, Callis J. 1997. A model for the evolution of polyubiquitin genes from the study of *Arabidopsis thaliana* ecotypes. *Plant Mol. Biol.* 34:745–58
  263. Ristic Z, Gifford DJ, Cass DD. 1991. Heat shock proteins in two lines of *Zea mays* L. that differ in drought and heat resistance. *Plant Physiol.* 97:1430–34
  264. Colombo SJ, Colclough ML, Timmer VR, Blumwald E. 1992. Clonal variation in heat tolerance and heat shock protein expression in black spruce. *Silvae Genet.* 41:234–39
  265. Fender SE, O'Connell MA. 1989. Heat shock protein expression in thermotolerant and thermosensitive lines of cotton. *Plant Cell Rep.* 8:37–40
  266. Malayer JR, Hansen PJ. 1990. Differences between Brahman and Holstein cows in heat-shock induced alterations of protein synthesis and secretion by oviducts and uterine endometrium. *J. Anim. Sci.* 68:266–80
  267. Otsuka Y, Takano TS, Yamazaki T. 1997. Genetic variation in the expression of the six hsp genes in the presence of heat shock in *Drosophila melanogaster*. *Genes Genetic Syst.* 72:19–24
  268. Jorgensen JA, Weng J, Ho THD, Nguyen HT. 1992. Genotype-specific heat shock proteins in two maize inbreds. *Plant Cell Rep.* 11:576–80
  269. Frova C, Gorla MS. 1993. Quantitative expression of maize Hsps: genetic dissection and association with thermotolerance. *Theor. Appl. Genet.* 86:213–20
  270. Weng J, Nguyen HT. 1992. Differences in the heat-shock response between thermotolerant and thermosusceptible cultivars of hexaploid wheat. *Theor. Appl. Genet.* 84:941–46
  271. Brown DC, Bradley BP, Tedengren M. 1995. Genetic and environmental regulation of HSP70 expression. *Mar. Env. Res.* 39:181–84
  272. Caruso M, Sacco M, Medoff G, Maresca B. 1987. Heat shock 70 gene is differentially expressed in *Histoplasma capsulatum* strains with different levels of thermotolerance and pathogenicity. *Mol. Microbiol.* 1:151–58
  273. Lyashko VN, Vikulova VK, Chernikov VG, Ivanov VI, Ulmasov KA, et al. 1994. Comparison of the heat shock response in ethnically and ecologically different human populations. *Proc. Natl. Acad. Sci. USA* 91:12492–95
  274. Bettencourt BR, Feder ME, Cavicchi S. 1997. Laboratory evolution of Hsp70 expression in *Drosophila melanogaster*: functional consequences and molecular bases. *Am. Zool.* 37:189A (Abstr.)
  275. McColl G, Hoffmann AA, McKechnie SW. 1996. Response of two heat shock genes to selection for knockdown heat resistance in *Drosophila melanogaster*. *Genetics* 143:1615–27
  276. Fender SE, O'Connell MA. 1990. Expression of the heat shock response in a tomato interspecific hybrid is not intermediate between the two parental responses. *Plant Physiol.* 93:1140–46
  277. Gaugler R, Wilson M, Shearer P. 1997. Field release and environmental fate of a transgenic entomopathogenic nematode. *Biol. Control* 9:75–80
  278. Hashmi S, Hashmi G, Glazer I, Gaugler R. 1998. Thermal response of *Heterorhabditis bacteriophora* transformed with the *Caenorhabditis elegans* hsp70 encoding gene. *J. Exp. Zool.* 281:164–70
  279. Trent JD. 1996. A review of acquired thermotolerance, heat-shock proteins, and molecular chaperones in Archaea. *FEMS Micro. Rev.* 18:249–58
  280. Holden JF, Baross JA. 1993. Enhanced thermotolerance and temperature-induced changes in protein composition in the hyperthermophilic archaeon ES4. *J. Bacteriol.* 175:2839–43
  281. Hamilton PT, Reeve JN. 1985. Structure of genes and an insertion element in the methane producing archaeobacterium *Methanobrevibacter smithii*. *Mol. Gen. Genet.* 200:47–59
  282. Waldmann T, Nimmesgern E, Nitsch M, Peters J, Pfeifer G, et al. 1995. The thermosome of *Thermoplasma acidophilum* and its relationship to the eukaryotic chaperonin TRiC. *Eur. J. Biochem.* 227:848–56
  283. Phipps BM, Typke D, Heger R, Volker S, Hoffmann A, et al. 1993. Structure of a molecular chaperone from a thermophilic archaeobacterium. *Nature* 361:475–77
  284. Nitsch M, Klumpp M, Lupas A, Baumeister W. 1997. The thermosome: alternating alpha and beta-subunits within the chaperonin of the archaeon *Thermoplasma acidophilum*. *J. Mol. Biol.* 267:142–49
  285. Lewis VA, Hynes GM, Zheng D, Saibil H, Willison K. 1992. T-complex polypeptide-1 is a subunit of a het-

- eromeric particle in the eukaryotic cytosol. *Nature* 358:249–52
286. Trent JD, Nimmegern E, Wall JS, Hartl FU, Horwich AL. 1991. A molecular chaperone from a thermophilic archaeobacterium is related to the eukaryotic protein t-complex polypeptide-1. *Nature* 354:490–93
  287. Hendrick JP, Hartl FU. 1995. The role of molecular chaperones in protein folding. *FASEB J.* 9:1559–69
  288. Eggers DK, Welch WJ, Hansen WJ. 1997. Complexes between nascent polypeptides and their molecular chaperones in the cytosol of mammalian cells. *Mol. Biol. Cell* 8:1559–73
  289. Burston SG, Clarke AR. 1995. Molecular chaperones: physical and mechanistic properties. *Essays Biochem.* 29:125–36
  290. Trent JD, Kagawa HK, Yaoi T, Olle E, Zaluzec NJ. 1997. Chaperonin filaments: the archaeal cytoskeleton? *Proc. Natl. Acad. Sci. USA* 94:5383–88
  291. Dascher CC, Poddar SK, Maniloff J. 1990. Heat shock response in mycoplasmas, genome-limited organisms. *J. Bacteriol.* 172:1823–27
  292. Gupta RS, Singh B. 1994. Phylogenetic analysis of 70 kD heat shock protein sequences suggests a chimeric origin for the eukaryotic cell nucleus. *Curr. Biol.* 4:1104–14
  293. Bustard K, Gupta RS. 1997. The sequences of heat shock protein 40 (DnaJ) homologs provide evidence for a close evolutionary relationship between the *Deinococcus thermus* group and Cyanobacteria. *J. Mol. Evol.* 45:193–205
  294. Gupta RS, Bustard K, Falah M, Singh D. 1997. Sequencing of heat shock protein 70 (DnaK) homologs from *Deinococcus proteolyticus* and *Thermomicrobium roseum* and their integration in a protein-based phylogeny of prokaryotes. *J. Bacteriol.* 179:345–57
  295. Gupta RS. 1995. Phylogenetic analysis of the 90 kD heat shock family of protein sequences and an examination of the relationship among animals, plants, and fungi species. *Mol. Biol. Evol.* 12:1063–73
  296. Gupta RS, Aitken K, Falah M, Singh B. 1994. Cloning of *Giardia lamblia* heat shock protein HSP70 homologs: implications regarding origin of eukaryotic cells and of endoplasmic reticulum. *Proc. Natl. Acad. Sci. USA* 91:2895–99
  297. Gupta RS, Singh B. 1992. Cloning of the HSP70 gene from *Halobacterium marismortui*: relatedness of archaeobacterial HSP70 to its eubacterial homologs and a model for the evolution of the HSP70 gene. *J. Bacteriol.* 174:4594–605
  298. Boorstein WR, Ziegelhoffer T, Craig EA. 1994. Molecular evolution of the HSP70 multigene family. *J. Mol. Evol.* 38:1–17
  299. Rubin DM, Mehta AD, Zhu J, Shoham S, Chen X, et al. 1993. Genomic structure and sequence analysis of *Drosophila melanogaster* HSC70 genes. *Gene* 128:155–63
  300. Tavaría M, Gabriele T, Kola I, Anderson RL. 1996. A hitchhiker's guide to the human Hsp70 family. *Cell Stress Chaperones* 1:23–28
  301. de Jong WW, Leunissen JA, Voorter CE. 1993. Evolution of the alpha-crystallin/small heat-shock protein family. *Mol. Biol. Evol.* 10:103–26
  302. Ohta T. 1994. Further examples of evolution by gene duplication revealed through DNA sequence comparisons. *Genetics* 138:1331–37
  303. Benedict MQ, Cockburn AF, Seawright JA. 1993. The Hsp70 heat-shock gene family of the mosquito *Anopheles albimanus*. *Insect Mol. Biol.* 2:93–102
  304. Drosopoulou E, Konstantopoulou I, Scouras ZG. 1996. The heat shock genes in the *Drosophila montium* subgroup: chromosomal localization and evolutionary implications. *Chromosoma* 105:104–10
  305. Pardali E, Feggou E, Drosopoulou E, Konstantopoulou I, Scouras ZG, Mavragani-Tsipidou P. 1996. The Afro-tropical *Drosophila montium* subgroup: Balbiani ring 1, polytene chromosomes, and heat shock response of *Drosophila vulcana*. *Genome* 39:588–97
  306. Sanchez Y, Lindquist SL. 1990. HSP104 required for induced thermotolerance. *Science* 248:1112–15
  307. Feder ME, Block BA. 1991. On the future of physiological ecology. *Funct. Ecol.* 5:136–44
  308. Lau S, Patnaik N, Sayen MR, Mestri R. 1997. Simultaneous overexpression of two stress proteins in rat cardiomyocytes and myogenic cells confers protection against ischemia-induced injury. *Circulation* 96:2287–94
  309. Huot J, Roy G, Lambert H, Chretien P, Landry J. 1991. Increased survival after treatments with anticancer agents of Chinese hamster cells expressing the human Mr 27,000 heat shock protein. *Cancer Res.* 51:5245–52

310. Mehlen P, Preville X, Chareyron P, Briolay J, Klemenz R, Arrigo AP. 1995. Constitutive expression of human hsp27, *Drosophila* hsp27, or human alpha B-crystallin confers resistance to TNF- and oxidative stress-induced cytotoxicity in stably transfected murine L929 fibroblasts. *J. Immunol.* 154:363-74
311. Jaattela M, Wissing D. 1993. Heat-shock proteins protect cells from monocyte cytotoxicity: possible mechanism of self-protection. *J. Exp. Med.* 177:231-36
312. Trautinger F, Kokesch C, Herbacek I, Knobler RM, Kindas-Mugge I. 1997. Overexpression of the small heat shock protein, hsp27, confers resistance to hyperthermia, but not to oxidative stress and UV-induced cell death, in a stably transfected squamous cell carcinoma cell line. *J. Photochem. Photobiol.* 39B:90-95
313. Rollet E, Lavoie JN, Landry J, Tanguay RM. 1992. Expression of *Drosophila's* 27 kDa heat shock protein into rodent cells confers thermal resistance. *Biochem. Biophys. Res. Commun.* 185:116-20
314. Lavoie JN, Gingras-Breton G, Tanguay RM, Landry J. 1993. Induction of Chinese hamster HSP27 gene expression in mouse cells confers resistance to heat shock. HSP27 stabilization of the microfilament organization. *J. Biol. Chem.* 268:3420-29
315. Landry J, Chretien P, Lambert H, Hickey E, Weber LA. 1989. Heat shock resistance conferred by expression of the human HSP27 gene in rodent cells. *J. Cell Biol.* 109:7-15
316. Wissing D, Jaattela M. 1996. HSP27 and HSP70 increase the survival of WEHI-S cells exposed to hyperthermia. *Int. J. Hypertherm.* 12:125-38
317. Wang G, Klostergaard J, Khodadadian M, Wu J, Wu TW, et al. 1996. Murine cells transfected with human Hsp27 cDNA resist TNF-induced cytotoxicity. *J. Immunother. Emphasis Tumor Immunol.* 19:9-20
318. Martin JL, Mestril R, Hilal-Dandan R, Brunton LL, Dillmann WH. 1997. Small heat shock proteins and protection against ischemic injury in cardiac myocytes. *Circulation* 96:4343-48
319. Stege GJ, Kampinga HH, Konings AW. 1995. Heat-induced intranuclear protein aggregation and thermal radiosensitization. *Int. J. Radiat. Biol.* 67:203-9
320. Blackburn R, Galoforo S, Berns CM, Ireland M, Cho JM, et al. 1996. Thermal response in murine L929 cells lacking alpha B-crystallin expression and alpha B-crystallin expressing L929 transfectants. *Mol. Cell. Biochem.* 155:51-60
321. van den Ijssel PR, Overkamp P, Knauf U, Gaestel M, de Jong WW. 1994. Alpha A-crystallin confers cellular thermoresistance. *FEBS Lett.* 355:54-56
322. Heads RJ, Yellon DM, Latchman DS. 1995. Differential cytoprotection against heat stress or hypoxia following expression of specific stress protein genes in myogenic cells. *J. Mol. Cell. Cardiol.* 27:1669-78
323. Cumming DV, Heads RJ, Watson A, Latchman DS, Yellon DM. 1996. Differential protection of primary rat cardiocytes by transfection of specific heat stress proteins. *J. Mol. Cell. Cardiol.* 28:2343-49
324. Lukacs KV, Nakakes A, Atkins CJ, Lowrie DB, Colston MJ. 1997. In vivo gene therapy of malignant tumours with heat shock protein-65 gene. *Gene Ther.* 4:346-50
325. Lukacs KV, Lowrie DB, Stokes RW, Colston MJ. 1993. Tumor cells transfected with a bacterial heat-shock gene lose tumorigenicity and induce protection against tumors. *J. Exp. Med.* 178:343-48
326. Wischmeyer PE, Musch MW, Madonna MB, Thisted R, Chang EB. 1997. Glutamine protects intestinal epithelial cells: role of inducible HSP70. *Am. J. Physiol.* 272:G879-84
327. Angelidis CE, Lazaridis I, Pagoulatos GN. 1991. Constitutive expression of heat-shock protein 70 in mammalian cells confers thermoresistance. *Eur. J. Biochem.* 199:35-39
328. Mailhos C, Howard MK, Latchman DS. 1994. Heat shock proteins hsp90 and hsp70 protect neuronal cells from thermal stress but not from programmed cell death. *J. Neurochem.* 63:1787-95
329. Wyatt S, Mailhos C, Latchman DS. 1996. Trigeminal ganglion neurons are protected by the heat shock proteins hsp70 and hsp90 from thermal stress but not from programmed cell death following nerve growth factor withdrawal. *Brain Res. Mol. Brain Res.* 39:52-56
330. Uney JB, Staley K, Tyers P, Sofroniew MV, Kew JN. 1994. Transfection with hsp70i protects rat dorsal root ganglia neurones and glia from heat stress. *Gene Ther.* 1:S65
331. Solomon JM, Rossi JM, Golic K, McGarry T, Lindquist S. 1991. Changes in Hsp70 alter thermotolerance and heat-

- shock regulation in *Drosophila*. *New Biol.* 3:1106–20
332. Sato K, Saito H, Matsuki N. 1996. HSP70 is essential to the neuroprotective effect of heat-shock. *Brain Res.* 740:117–23
  333. Riabowol KT, Mizzen LA, Welch WJ. 1988. Heat shock is lethal to fibroblasts microinjected with antibodies against hsp70. *Science* 242:433–36
  334. Mestril R, Giordano FJ, Conde AG, Dillmann WH. 1996. Adenovirus-mediated gene transfer of a heat shock protein 70 (hsp 70i) protects against simulated ischemia. *J. Mol. Cell. Cardiol.* 28:2351–58
  335. Liu RY, Li X, Li L, Li GC. 1992. Expression of human hsp70 in rat fibroblasts enhances cell survival and facilitates recovery from translational and transcriptional inhibition following heat shock. *Cancer Res.* 52:3667–73
  336. Nakata N, Kato H, Kogure K. 1993. Inhibition of ischaemic tolerance in the gerbil hippocampus by quercetin and anti-heat shock protein-70 antibody. *NeuroReport* 4:695–98
  337. Lee YJ, Kim D, Hou ZZ, Curetty L, Borrelli MJ, Corry PM. 1993. Alteration of heat sensitivity by introduction of hsp70 or anti-hsp70 in CHO cells. *J. Therm. Biol.* 18:229–36
  338. Johnston RN, Kucey BL. 1988. Competitive inhibition of hsp70 gene expression causes thermosensitivity. *Science* 242:1551–54
  339. Heads RJ, Latchman DS, Yellon DM. 1994. Stable high level expression of a transfected human HSP70 gene protects a heart-derived muscle cell line against thermal stress. *J. Mol. Cell. Cardiol.* 26:695–99
  340. Amin V, Cumming DV, Latchman DS. 1996. Over-expression of heat shock protein 70 protects neuronal cells against both thermal and ischaemic stress but with different efficiencies. *Neurosci. Lett.* 206:45–48
  341. Khan NA, Sotelo J. 1989. Heat shock stress is deleterious to CNS cultured neurons microinjected with anti-HSP70 antibodies. *Biol. Cell* 65:199–202
  342. Li GC, Li LG, Liu YK, Mak JY, Chen LL, Lee WM. 1991. Thermal response of rat fibroblasts stably transfected with the human 70-kDa heat shock protein-encoding gene. *Proc. Natl. Acad. Sci. USA* 88:1681–85
  343. Williams RS, Thomas JA, Fina M, German Z, Benjamin IJ. 1993. Human heat shock protein 70 (hsp70) protects murine cells from injury during metabolic stress. *J. Clin. Invest.* 92:503–8
  344. Mestril R, Chi SH, Sayen MR, O'Reilly K, Dillmann WH. 1994. Expression of inducible stress protein 70 in rat heart myogenic cells confers protection against simulated ischemia-induced injury. *J. Clin. Invest.* 93:759–67
  345. Dillmann WH, Mestril R. 1995. Heat shock proteins in myocardial stress. *Z. Kardiol.* 4:87–90
  346. Jacobs M, Andersen JB, Kontinen V, Sarvas M. 1993. The DNA-binding activity of the human heat shock transcription factor is regulated in vivo by Hsp70. *Mol. Cell Biol.* 13:5427–38
  347. Ding XZ, Tsokos GC, Smallridge RC, Kiang JG. 1997. Heat shock gene-expression in HSP-70 and HSF1 gene-transfected human epidermoid A-431 cells. *Mol. Cell. Biochem.* 167:145–52
  348. Chi SH, Mestril R. 1996. Stable expression of a human HSP70 gene in a rat myogenic cell line confers protection against endotoxin. *Am. J. Physiol.* 270:C1017–21
  349. Han MY, Park YM. 1997. Reduced protein denaturation in thermotolerant cells by elevated levels of HSP70. *Korean J. Pharmacol.* 32:433–44
  350. Jaattela M. 1995. Over-expression of hsp70 confers tumorigenicity to mouse fibrosarcoma cells. *Int. J. Cancer* 60:689–93
  351. Wei YQ, Zhao X, Kariya Y, Teshigawara K, Uchida A. 1995. Inhibition of proliferation and induction of apoptosis by abrogation of heat-shock protein (HSP) 70 expression in tumor cells. *Cancer Immunol. Immunother.* 40:73–78
  352. Karlseder J, Wissing D, Holzer G, Orel L, Sliutz G, et al. 1996. Hsp70 over-expression mediates the escape of a doxorubicin-induced G2 cell cycle arrest. *Biochem. Biophys. Res. Commun.* 220:153–59
  353. Henle KJ, Jethmalani SM, Li L, Li GC. 1997. Protein glycosylation in a heat-resistant rat fibroblast cell model expressing human HSP70. *Biochem. Biophys. Res. Commun.* 232:26–32
  354. Simon MM, Reikerstorfer A, Schwarz A, Krone C, Luger TA, et al. 1995. Heat shock protein 70 overexpression affects the response to ultraviolet light in murine fibroblasts. Evidence for increased cell viability and suppression of cytokine release. *J. Clin. Invest.* 95:926–33
  355. Lioussis SN, Ding XZ, Kiang JG, Tsokos GC. 1997. Overexpression of

- the heat shock protein 70 enhances the TCR/CD3- and Fas/Apo-1/CD95-mediated apoptotic cell death in Jurkat T cells. *J. Immunol.* 158:5668-75
356. Mosser DD, Caron AW, Bourget L, Denis-Larose C, Massie B. 1997. Role of the human heat shock protein hsp70 in protection against stress-induced apoptosis. *Mol. Cell. Biol.* 17:5317-27
  357. Suzuki K, Sawa Y, Kaneda Y, Ichikawa H, Shirakura R, Matsuda H. 1997. In vivo gene transfection with heat shock protein 70 enhances myocardial tolerance to ischemia-reperfusion injury in rat. *J. Clin. Invest.* 99:1645-50
  358. Plumier JC, Ross BM, Currie RW, Angelidis CE, Kazlaris H, et al. 1995. Transgenic mice expressing the human heat shock protein 70 have improved post-ischemic myocardial recovery. *J. Clin. Invest.* 95:1854-60
  359. Marber MS, Mestrl R, Chi SH, Sayen MR, Yellon DM, Dillmann WH. 1995. Overexpression of the rat inducible 70-kD heat stress protein in a transgenic mouse increases the resistance of the heart to ischemic injury. *J. Clin. Invest.* 95:1446-56
  360. Plumier JC, Krueger AM, Currie RW, Kontoyiannis D, Kollias G, Pagoulatos GN. 1997. Transgenic mice expressing the human inducible Hsp70 have hippocampal neurons resistant to ischemic injury. *Cell Stress Chaperones* 2:162-67
  361. Lee JH, Schoffl F. 1996. An Hsp70 antisense gene affects the expression of HSP70/HSC70, the regulation of HSF, and the acquisition of thermotolerance in transgenic *Arabidopsis thaliana*. *Mol. Gen. Genet.* 252:11-19
  362. Galea-Lauri J, Richardson AJ, Latchman DS, Katz DR. 1996. Increased heat shock protein 90 (hsp90) expression leads to increased apoptosis in the monoblastoid cell line U937 following induction with TNF-alpha and cycloheximide: a possible role in immunopathology. *J. Immunol.* 157:4109-18
  363. Nakano M, Mann DL, Knowlton AA. 1997. Blocking the endogenous increase in HSP 72 increases susceptibility to hypoxia and reoxygenation in isolated adult feline cardiocytes. *Circulation* 95:1523-31
  364. Hutter JJ, Mestrl R, Tam EK, Sievers RE, Dillmann WH, Wolfe CL. 1996. Overexpression of heat shock protein 72 in transgenic mice decreases infarct size in vivo. *Circulation* 94:1408-11
  365. Galea-Lauri J, Latchman DS, Katz DR. 1996. The role of the 90-kDa heat shock protein in cell cycle control and differentiation of the monoblastoid cell line U937. *Exp. Cell Res.* 226:243-54
  366. Schirmer EC, Lindquist S, Vierling E. 1994. An *Arabidopsis* heat shock protein complements a thermotolerance defect in yeast. *Plant Cell* 6:1899-909
  367. Kutsikova IUA, Mamon LA. 1996. Consequences of exposure to extreme conditions in somatic cells of *Drosophila melanogaster* under conditions of disturbed synthesis of heat shock proteins. *Genetika* 32:1406-16
  368. Mamon LA, Kutsikova YA. 1993. The role of the heat-shock proteins in recovery of high temperature induced damages of mitotic chromosomes in *Drosophila melanogaster*. *Genetika* 29:604-12
  369. Mamon LA, Kutsikova YA. 1993. The role of the heat-shock proteins in recovery of cell proliferation following high temperature treatment of *Drosophila melanogaster*. *Genetika* 29:791-98
  370. Koishi M, Hosokawa N, Sato M, Nakai A, Hirayoshi K, et al. 1992. Quercetin, an inhibitor of heat shock protein synthesis, inhibits the acquisition of thermotolerance in a human colon carcinoma cell line. *Jpn. J. Cancer Res.* 83:1216-22
  371. Lee YJ, Currey L, Hou ZZ, Kim SH, Kim JH, Corry PM. 1992. Effect of pH on quercetin-induced suppression of heat shock gene expression and thermotolerance development in HT-29 cells. *Biochem. Biophys. Res. Commun.* 186:1121-28
  372. Jedlicka P, Mortin MA, Wu C. 1997. Multiple functions of *Drosophila* heat shock transcription factor in vivo. *EMBO J.* 16:2452-62
  373. Lee JH, Hubel A, Schoffl F. 1995. Derepression of the activity of genetically engineered heat shock factor causes constitutive synthesis of heat shock proteins and increased thermotolerance in transgenic *Arabidopsis*. *Plant J.* 8:603-12