



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Hemin pre-conditioning upregulates heme-oxygenase 1 in deceased donor renal transplant recipients and may offer protection: a randomised, controlled, phase IIB trial

Citation for published version:

Thomas, RAB, Czopek, A, Bellamy, COC, McNally, SJ, Kluth, DC & Marson, LP 2015, 'Hemin pre-conditioning upregulates heme-oxygenase 1 in deceased donor renal transplant recipients and may offer protection: a randomised, controlled, phase IIB trial', *Transplantation*.
<https://doi.org/10.1097/TP.0000000000000770>

Digital Object Identifier (DOI):

[10.1097/TP.0000000000000770](https://doi.org/10.1097/TP.0000000000000770)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Transplantation

Publisher Rights Statement:

This is the author's final manuscript as accepted for publication.

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Hemin pre-conditioning upregulates heme-oxygenase 1 in deceased donor renal transplant recipients and may offer protection: a randomised, controlled, phase IIB trial.

Rachel AB Thomas¹, Alicja Czopek¹, Christopher OC Bellamy², Stephen McNally^{1,3}, David C Kluth^{1*}, Lorna P Marson^{1*}.

¹Centre for Inflammation Research, Queen's Medical Research Institute, University of Edinburgh, 47 Little France Crescent, Edinburgh, EH16 4TJ

²Dept. Pathology, Edinburgh Royal Infirmary, 51 Little France Crescent, Edinburgh, EH16 4SA

³Freeman Hospital, High Heaton, Newcastle Upon Tyne NE7 7DN

*These authors contributed equally as senior authors.

Address for correspondence:

Rachel Thomas

rab_thomas@hotmail.co.uk

215 Lanark Road West, Currie, Midlothian, EH14 5NT, Scotland, UK

Funding

The study was funded by NHS Blood and Transplant.

Registration of clinical trial

EudraCT number: **2011-004311-23**

ClinicalTrials.gov number: **NCT01430156**

Disclosure

The authors declare no conflicts of interest.

Footnotes

Author's specific contributions

¹RT, DK and LM designed the trial with scientific research input from CB. RT was Chief Investigator for the HOT study and ran the day-to-day research and lab analysis. DK, SM and LM were Principle Investigators guiding the trial and approving changes to protocols and analysis. CB performed the histological scoring of renal tissue and provided laboratory advice. AC provided lab advice and assisted with specimen analysis. RT, DK and LM participated in data analysis.

RT wrote the first draft of this paper with significant input by DK and LM. All authors reviewed the manuscripts, made changes and approved it.

During the HOT study, RT was funded by NHS Blood and Transplant. No other financial support was provided. There were no conflicts of interests.

Author e-mails

[²rab_thomas@hotmail.co.uk](mailto:rab_thomas@hotmail.co.uk)

aczopek@staffmail.ed.ac.uk

Stephen.mcnally@gmail.com

Lorna.marson@ed.ac.uk

David.kluth@ed.ac.uk

c.bellamy@ed.ac.uk

Abbreviations:

CI	Chief Investigator
CIT	Cold ischemic time
CO	Carbon monoxide
DGF	Delayed graft function
ERI	Edinburgh Royal Infirmary
ESRD	End stage renal disease
HA	Heme arginate
HO-1	Heme-oxygenase 1
IQR	Interquartile range
IRI	Ischaemia reperfusion injury
KIM-1	Kidney injury marker 1
NGAL	Neutrophil gelatinase-associated lipocalin
MHRA	Medicines and Healthcare products Regulatory Agency (UK drug safety agency)
PBMC	Peripheral blood mononuclear cell
PCR	Polymerase chain reaction
WIT	Warm ischaemic time

Word count:

Abstract 250

Text including abstract 3288

Abstract

Background:

1 The enzyme heme oxygenase-1 (HO-1) degrades heme and protects against ischaemia-reperfusion
2 injury (IRI), which is inevitable in transplantation. Monocytes/macrophages (MΦ) are the major
3 source of HO-1 and higher levels improve renal transplant outcomes. Heme Arginate (HA) can safely
4 induce HO-1 in humans. This randomised controlled trial evaluated the effect of HA on HO-1
5 upregulation and renal function in recipients of deceased donor kidneys.
6
7
8
9
10
11
12

Methods:

13 40 recipients were randomised to either active (3mg kg⁻¹ HA: pre-operatively and day 2) or placebo
14 (NaCl: same schedule). Recipient blood was taken daily for peripheral blood mononuclear cells
15 (PBMC) extraction. Urine was also collected. Graft biopsies were taken pre-op and day 5.
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Results:

HA upregulated PBMC HO-1 protein at 24 hours more than placebo: HA 11.1ng/ml vs. placebo
0.14ng/ml (p<0.0001). PBMC HO-1 mRNA was also increased: HA 2.73 fold vs. placebo 1.41 fold
(p=0.02). HA increased day 5 tissue HO-1 protein immunopositivity compared with placebo: HA
0.21 vs. placebo -0.03 (p=0.02) and % HO-1 positive renal MΦ also increased: HA 50.8 cells per hpf
vs. placebo 22.3 (p=0.012). Urinary biomarkers were reduced after HA but not significantly so.
Histological injury and renal function were similar but the study was not powered to these endpoints.
Adverse events were equivalent between groups.

Discussion:

The primary outcome was achieved and demonstrated for the first time that HA safely induces HO-1
in renal transplant recipients. Larger studies are planned to determine the impact of HO-1
upregulation on clinical outcomes and evaluate the benefit to patients at risk of IRI.

Introduction

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Renal transplantation is the optimal management for end-stage renal disease (ESRD) because it improves quality of life and survival (1). Deceased donor renal transplantation inevitably results in ischemia-reperfusion injury (IRI), which is characterised by the production of reactive oxygen free radicals, alterations in blood flow, leucocyte infiltration and significant cell dysfunction (2). IRI increases the risk of delayed graft function (DGF), necessitating post-transplant dialysis and leading to an increased risk of rejection and reduced long-term graft survival (3). Apart from limiting cold ischemic time, few methods exist to minimise the effects of IRI. One potential technique is preconditioning, using either ischemia or pharmacology (4).

The enzyme heme-oxygenase 1 (HO-1) has emerged as a potential approach to attenuate IRI (5). HO-1 is the rate-limiting enzyme in heme catabolism and degrades the pro-inflammatory, oxidative free heme molecules into biliverdin (further degraded to bilirubin), carbon monoxide (CO) and free iron(6). Bilirubin is an anti-oxidant. CO has vasodilatory and anti-apoptotic properties and inhibits a number of pro-inflammatory signalling pathways and platelet aggregation. Free iron is sequestered by ferritin to limit its toxic effect (7).

Experimental models have shown that following renal IRI, macrophages have roles in both injury and repair. They are also the primary HO-1 expressing leucocyte (8). Macrophage infiltration into the kidney occurs within 24 hours of IRI and these initial cells have a pro-inflammatory phenotype (M1), contributing to various injury processes. As a result, depletion of macrophages at early time-points post-IRI protects renal function (9). By contrast, renal macrophages present on days 3-5 following IRI have an alternative activation phenotype (M2) and contribute to tissue repair and recovery of renal function (9). In a renal IRI model, increased macrophage HO-1 expression was associated with reduced injury (10). Improved renal function was also seen after infusion of HO-1 expressing macrophages in a similar model (11). Our laboratory has demonstrated in a mouse IRI model that heme arginate (HA) pre-conditioning upregulates renal HO-1, principally in interstitial macrophages, and results in a significant reduction in subsequent renal injury. This protective effect was lost following macrophage depletion (12). Thus, it was hypothesised that amplified HO-1 expression in recipient macrophages and renal grafts might reduce IRI and improve outcomes after clinical

transplantation. HA, a form of hemin, is suitable for clinical studies, as it has been used safely in the treatment of acute porphyria for over 30 years with few reported side effects. In healthy volunteers, both hemin and HA have been shown to increase HO-1 concentration in peripheral blood mononuclear cells (PBMC) and plasma (13, 14).

The HOT study was a randomised, placebo controlled, blinded, single centre study with the primary objective of determining whether pre-treatment of renal transplant recipients with HA upregulated HO-1 in PBMCs. Secondary objectives explored the effect of HA on graft tissue, urinary biomarkers and renal function.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Materials and methods

1
2 All patients on the East of Scotland renal transplant waiting list were informed about the study by
3 post before recruitment started. When patients were admitted to Edinburgh Royal Infirmary (ERI) for
4 potential deceased donor renal transplant, they were assessed for eligibility.
5

6 A patient was ineligible for inclusion if:
7

- 8 1. they were unable to receive the standard immunosuppressive regimen
- 9
- 10 2. they were unable to give informed consent
- 11
- 12 3. they had hypersensitivity reactions to HA
- 13
- 14 4. it was a combined transplant or their 3rd or subsequent kidney transplant
- 15
- 16 5. they were to be anticoagulated post-operatively or on combined anti-platelet agents due to
17 biopsy risks
18
19
20
21
22

23 Study protocol

24
25 This was a single-centre, randomised, blinded, placebo-controlled trial sponsored by ACCORD (a
26 joint company from University of Edinburgh and NHS Lothian) and funded by NHS Blood and
27 Transplant. The trial protocol was registered at the European Clinical Trials database (EudraCT no:
28 2011-004311-23) and ClinicalTrials.gov (NCT01430156) and adhered to the Declaration of Helsinki
29 and CONSORT guidelines. The Scottish Regional Ethics Committee and the Medicines and
30 Healthcare products Regulatory Agency (MHRA) approved the study (approval number
31 2011/R/TR/03 and protocol number HOTstudy_Thomas11).
32
33
34
35
36
37
38
39
40
41
42

43 The sample size calculation was performed on the observed heme oxygenase-1 (HO-1) upregulation
44 in cultured human macrophages following HA treatment. Using a two-sided two-sample test with a
45 5% level of significance and 80% power, assuming a mean baseline to 24-hour change of 7.47 and a
46 common standard deviation of 3.87, the minimum detectable difference in means would be 3.8 with a
47 sample size of 17 per group. The final sample size of 20 per group allowed for dropouts.
48
49
50
51
52
53

54
55 The randomisation was done by random, random block, with stratification by donor type. The trial
56
57
58
59
60
61
62
63
64
65

Investigator (CI) after consent had been given and participant number had been assigned. Subjects were randomised to receive either 100ml 0.9% NaCl (placebo) or 3mg kg⁻¹ HA (Normosang, Orphan Europe, France) diluted in 100ml 0.9% NaCl. In both groups, participants received the first infusion pre-operatively (D0) and this was repeated on day 2 (D2). To ensure maximum post-conditioning exposure, the D0 infusion was given as early as possible post-dialysis (if required). All staff (except the CI giving the infusion) and participants were blinded.

Prior to infusion, venous blood was taken for PBMC extraction and analysis of baseline HO-1 expression (D0). Venepuncture was repeated 24 hours after the first infusion (D1) and also on days 2 (D2, pre-2nd infusion), D3 and D5. Urine was collected on the same days for analysis of KIM-1 and NGAL. A needle-biopsy of the graft was taken at the back table (D0) for routine pathology and study analysis. A repeat biopsy was requested for all participants on D5. Laboratory samples were analysed by the CI after blinding by an independent lab member. All samples were stored in accordance with guidelines.

Otherwise all pre-operative and post-operative care was typical, including standard immunosuppression, which comprised of induction therapy with basiliximab, tacrolimus, mycophenolate mofetil (MMF) and prednisolone.

HO-1 mRNA and protein levels in PBMCs

PBMCs were isolated using the previously described Dextran-Percoll method and stored in appropriate buffers for subsequent batch analysis of mRNA and protein expression (15). PBMC RNA was extracted using the ISOLATE kit (Bioline, UK) and cDNA was synthesised using TaqMan, RT-PCR core kit (Invitrogen, USA). The cDNA was analysed in duplicate by quantitative real-time PCR using fluorogenic probes for HO-1 and 18s (housekeeping gene) on an ABI Prism 7900 PCR machine (Applied Biosystems, USA). For quantification, the threshold cycle of HO-1 (C_T) was correlated to the constant expression of the housekeeping gene; ΔC_T was defined as C_T (HO-1) - C_T (18s). The differences in expression pre and post infusion were plotted as $\Delta\Delta C_T$ and $2^{-\Delta\Delta C_T}$ calculation was performed to determine the fold increase and the two groups were compared.

Each sample protein concentration was determined using protein assay (Biorad, USA). All samples were standardised to 1mg/ml of total protein and this was analysed in duplicate by the HO-1 ELISA kit to determine the amount of HO-1 protein at each time point (ENZO, USA).

Analysis of renal tissue

The D0 biopsy and D5 biopsy samples were handled and analysed as below:

- RNA extracted from fresh tissue and analysed in the same manner as PBMC RNA
- a blinded expert (CB) examined a fixed sample to assess for renal injury using a 13 point system.
- a fixed section of all the tissue samples underwent immunohistological staining for HO-1 (ENZO, USA). ImageJ (NIH) was used to quantify the amount of HO-1 protein as percentage of total protein.
- another section of tissue underwent dual immunofluorescence staining for HO-1 (ENZO, USA) and CD68 (macrophage marker (Abcam, UK)) and the numbers of dual positive cells were counted per high power field.

Urinary biomarkers

Commercially available ELISA kits from R&D Systems measured urinary KIM-1 and NGAL. KIM-1 and NGAL values are recorded as a ratio to urinary creatinine to allow for variations in urine concentrations.

Clinical outcomes

Clinical blood results were recorded daily. Need for dialysis, adverse events and urine volumes were also recorded. In this study, DGF was determined by change in creatinine in the first week after transplantation. Renal function was recorded at day 30 and day 90.

Data and statistical analysis

The anonymised data was held in a secure, auditable database. PBMC and renal HO-1 expression, urine biomarkers and renal macrophage data were not normally distributed; a non-parametric Mann-Whitney statistical test was used. A comparison of proportions test was used to compare rates of DGF between groups. Results are presented as median [IQR] and % values (95% confidence

intervals) as appropriate. Statistical analysis was performed on SDSS and graphs created in Prism. A p-value of <0.05 was considered statistically significant.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Results

Trial participation

1 Patient recruitment began in January 2012 and continued until 40 patients had been randomised in
2
3 May 2013 (Figure 1). Baseline characteristics were similar between the groups (Table 1). No adverse
4
5 reactions and no deaths occurred during the seven-day trial period. There was one graft loss due to
6
7 technical reasons on day seven (placebo group). Acute rejection occurred in one participant in
8
9 placebo group and two in HA group. Five serious adverse events were reported including one ITU
10
11 admission for pulmonary oedema (placebo group), and one further procedure was required in each
12
13 group (placebo: radiological embolization for renal arterio-venous fistula, HA group: nephrectomy
14
15 for graft bleeding). All participants were followed up to 90 days post transplant.
16
17
18
19
20

21 Three potential recipients received the infusion but were not transplanted following back-table
22
23 examination of the kidney at day 0 (D0). All 37 participants who were transplanted provided a blood
24
25 sample at 24 hours (D1) post infusion for primary analysis. One participant in each group refused the
26
27 second infusion and subsequent venepuncture but consented to clinical follow-up. Thus 17 in the
28
29 placebo group and 18 in HA group received both infusions. Five participants in the placebo group
30
31 and three in the HA group declined the day 5 (D5) biopsy and two in each group were not considered
32
33 fit for biopsy. Therefore, there were 25 paired renal tissue samples available for analysis. All
34
35 participants with a transplant *in situ* at day seven were followed-up. Results are expressed as median
36
37 values with interquartile range [IQR].
38
39
40
41
42
43

HA upregulated HO-1 in peripheral blood monocytes

44 Expressed as the difference in HO-1 expression on D1 compared with D0, HA upregulated HO-1
45
46 concentration by 11.1ng/ml [1.0-37.0], compared with placebo -0.14ng/ml [-0.7-0.3] ($p < 0.0001$,
47
48 Figure 2a). This effect was confirmed with increased expression of HO-1 mRNA: HA treatment
49
50 upregulated HO-1 mRNA expression 2.73 fold [1.8-3.2] compared with placebo 1.41 fold [1.2-2.2]
51
52 ($p = 0.02$, Figure 2b). There was a peak in HO-1 PBMC protein concentration after each HA infusion
53
54 (Figure 2c) but mRNA upregulation did not significantly follow this pattern (Figure 2d).
55
56
57
58
59
60
61
62
63
64
65

HA upregulated HO-1 in renal macrophages

1 The number of CD68-positive macrophages per high power field (hpf) was similar between the
2
3 treatment groups at D0 (HA: 3.50 cells per hpf [1.50- 5.25], placebo: 3.00 [2.25- 4.25]) (p=0.956, Figure
4
5 3a). At D5 there were more CD68-positive macrophages in HA group (HA: 7.38 cells per hpf [4.80-
6
7 9.13], placebo: 11.0 [4.5- 25.0]) but this was not significant (p=0.13, Figure 3a). However, there was a
8
9 significant increase in the number of HO-1-positive CD68 macrophages following HA treatment at D5
10
11 (HA: 50.8 cells per hpf [40.0- 59.8], placebo: 22.3 [0.0- 34.8]) (p=0.012, Figure 3b and images Figure
12
13 3c-f).
14
15
16
17
18

HA upregulated HO-1 in renal tissue

19 There was significant increased expression of renal HO-1 protein in the HA-treated group over time,
20
21 which was not seen with placebo (change in protein expression [D5-D0]: HA: 0.21 [0.1- 0.6], placebo:
22
23 -0.03 [-1.3- 0.1]) (p=0.03, Figure 4a and images 4c-f). HO-1 mRNA expression was not significantly
24
25 affected by HA treatment: HA: 1.68 [0.20- 4.03], placebo 2.02 [0.75- 10.39] (p=0.45, Figure 4b).
26
27
28
29
30
31

HA had limited effect on urinary biomarkers

32 There was no significant difference in urinary KIM-1 and NGAL levels between patients treated with
33
34 HA versus placebo at each time point (Figures 5a and 5b).
35
36
37
38
39
40

HA did not alter histological injury or renal function

41 There were no significant differences in the severity of tubular injury or necrosis between HA and
42
43 placebo (data not shown).
44
45

46 There were 12 cases (67%) of DGF in the placebo group and 10 cases (53%) in HA group, a
47
48 difference of 14% (95% CI; -17.2 – 45.3), (p=0.38). When DGF was redefined as dialysis within the
49
50 first seven days post-transplant except for hyperkalaemia, the results were the same. At day 90, there
51
52 was one participant on dialysis in placebo group and none in HA group. There was no improvement
53
54 in creatinine after HA treatment at day 30 and day 90 (data not shown).
55
56
57
58
59
60
61
62
63
64
65

Discussion

1 The study achieved its principal objective: pre-treatment of deceased donor renal transplant recipients
2 with HA was feasible, safe and well tolerated in immunosuppressed ESRD patients. It also led to
3 significant upregulation of HO-1 protein and mRNA in recipient PBMCs. HO-1 is induced in
4 response to multiple stimuli including hypoxia and ischaemic cytokines so it was anticipated that all
5 participants would experience an upregulation in HO-1 after transplantation. Nevertheless, HA
6 treatment increased macrophage HO-1 more than placebo treatment and reiterated results from
7 cellular, pre-clinical and healthy volunteer studies (12-14). Figure 2c confirmed that, as expected,
8 there is a distinct peak in protein concentration after each HA infusion with a return to near baseline
9 expression 48 hours later. This replicates the findings of previous healthy volunteer studies (14). The
10 pattern of mRNA upregulation did not mirror this because although there was a significant peak at 24
11 hours, this was not seen again after the second infusion. This anomaly may be due to the $\Delta\Delta C_T$
12 method used because all samples were referenced to baseline (D0) and it may be more appropriate to
13 normalise D3 to D2 to identify a second upregulation.
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29

30 This is the first clinical study to show that HA treatment increases HO-1 protein immunopositivity
31 and the number of HO-1 positive macrophages in renal graft tissue. Renal HO-1 mRNA did not have
32 the same response which may be related to timing of the biopsy. Earlier research has shown that the
33 half-life of HO-1 mRNA *in vitro* renal cells after hemin treatment is only four hours (16) and
34 therefore the D5 biopsy may be too late to detect a difference in the more transient mRNA.
35
36
37
38
39
40
41
42
43

44 The present study was powered for a cellular outcome but given its importance, the study attempted
45 to establish whether pre-treatment with HA was associated with a reduction in renal injury or
46 function post-transplant. Serum creatinine is an unreliable marker of acute kidney injury and does not
47 accurately reflect the degree of damage until it reaches a nadir, which may be several days post-op.
48 The urinary biomarkers KIM-1 and NGAL have been validated as independent predictors of graft
49 survival after renal transplantation (17, 18). In this study, the maximum NGAL and KIM-1 levels
50 were lower after HA treatment on all days indicating reduced renal injury but the differences were
51 not statistically significant. There was also no evidence of histological protection after biopsy
52 analysis.
53
54
55
56
57
58
59
60
61
62
63
64
65

1 One of the challenges in clinical trials of renal transplantation is the lack of a single clear definition
2 of DGF (19). In the present study the ‘functional’ definition was used, defined as increased or stable
3 serum creatinine, or a decrease of <10% per day for three consecutive days in the first week after
4 transplantation (20). This was preferred over the definition of DGF as the requirement for dialysis
5 within the first seven days following transplantation, because the decision to dialyse a patient may be
6 subjective with variation in practice between clinicians. Also a recent study found functional DGF
7 but not dialysis-defined DGF to be associated independently with subsequent transplant failure (20).
8
9

10
11
12
13
14
15
16
17
18
19 Despite clear evidence that HA upregulated HO-1 in this study, it did not translate into the structural
20 or functional cytoprotection seen in pre-clinical and observational studies (12, 21-24). In laboratory
21 studies, induction of HO-1 in macrophages (10), and renal tissue (12, 23) offers protection against
22 renal IRI and the presence of HO-1 expressing renal macrophages has been shown to safeguard renal
23 function in the face of significant structural injury (11). There are a few possible explanations why
24 HO-1 upregulation did not confer protection; firstly, there was a modest protection but the sample
25 size was too small to detect it. This provisional study was not powered to this endpoint and larger
26 studies are required. This is a common issue in renal transplant research and few interventions have
27 impacted on DGF rates despite promising pre-clinical results.
28
29

30 Secondly, two doses of HA were insufficient. Human studies have shown that HO-1 expression
31 increases with additional HA doses and more may be required to maximise and sustain the effect on
32 macrophages and renal tissue for clinical benefits (14).
33
34

35 Thirdly, it may be that the equivocal structural and functional results are genuine because HA has
36 other effects in clinical IRI that inhibits the anticipated positive consequences of HO-1 upregulation.
37
38

39 Fourthly, it has also been proposed that the protective response of HO-1 may be limited and
40 susceptible to being overwhelmed. It may be that once a level of injury has been reached, further
41 HO-1 induction cannot prevent the damage (25). Clinical renal transplantation delivers sizable,
42 diverse insults to the kidney, which are not fully modelled in animal experimentation, and may
43 explain why preclinical findings do not equate to human studies.
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Given the unique nature of this study, further research is required to determine whether the amount of HO-1 upregulation seen here is sufficient for protection. This study establishes a sound basis for undertaking a clinical trial of HA vs. placebo powered to detect clinical differences in graft function and this is planned. There is also scope to increase the number of HA infusions within the UK prescribing licence and this is also currently under investigation in our centre. HA is a safe alternative to hemin because the effects of ferric heme are reduced when hemin is liganded to arginine as in HA (26). HA may offer an alternative to other renal transplant conditioning strategies that have not fulfilled their potential when translated to clinical studies such as erythropoietin (27, 28), statins (29) and ischaemic pre-conditioning (30). HA is well suited for such a trial in renal transplant recipients because it was well tolerated and has demonstrated induction of HO-1 in PBMCs and renal tissue when administered pre and post-renal transplant surgery.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Acknowledgements

The HOT study was only possible thanks to the generous funding from NHS Blood and Transplant and with help from the laboratory staff at QMRI and staff in the Radiology department, Pathology department and the Transplant unit in Royal Infirmary of Edinburgh. The East of Scotland recipient transplant coordinators were crucial to the study and our grateful thanks to them. Cat Graham and Sharon Harden at University of Edinburgh provided statistical support.

References

1. Oniscu GC, Brown H, Forsythe JL. Impact of cadaveric renal transplantation on survival in patients listed for transplantation. *J Am Soc Nephrol*. 2005;16(6):1859-65.
2. Chatauret N, Badet L, Barrou B, Hauet T. Ischemia-reperfusion: From cell biology to acute kidney injury. *Progres en urologie : journal de l'Association francaise d'urologie et de la Societe francaise d'urologie*. 2014;24 Suppl 1:S4-12.
3. Yarlagadda SG, Coca SG, Formica RN, Jr., Poggio ED, Parikh CR. Association between delayed graft function and allograft and patient survival: a systematic review and meta-analysis. *Nephrol Dial Transplant*. 2009;24(3):1039-47.
4. Selzner N, Boehnert M, Selzner M. Preconditioning, postconditioning, and remote conditioning in solid organ transplantation: basic mechanisms and translational applications. *Transplant Rev (Orlando)*. 2012;26(2):115-24.
5. Ferenbach DA, Kluth DC, Hughes J. Hemeoxygenase-1 and renal ischaemia-reperfusion injury. *Nephron Exp Nephrol*. 2010;115(3):e33-7.
6. Kirkby KA, Adin CA. Products of heme oxygenase and their potential therapeutic applications. *Am J Physiol Renal Physiol*. 2006;290(3):F563-71.
7. Nath KA. Heme oxygenase-1: a provenance for cytoprotective pathways in the kidney and other tissues. *Kidney Int*. 2006;70(3):432-43.
8. Roach JP, Moore EE, Partrick DA, Damle SS, Silliman CC, McIntyre RC, Jr., et al. Heme oxygenase-1 induction in macrophages by a hemoglobin-based oxygen carrier reduces endotoxin-stimulated cytokine secretion. *Shock*. 2009;31(3):251-7.
9. Lee S, Huen S, Nishio H, Nishio S, Lee HK, Choi BS, et al. Distinct macrophage phenotypes contribute to kidney injury and repair. *J Am Soc Nephrol*. 2011;22(2):317-26.
10. Gueler F, Park JK, Rong S, Kirsch T, Lindschau C, Zheng W, et al. Statins attenuate ischemia-reperfusion injury by inducing heme oxygenase-1 in infiltrating macrophages. *Am J Pathol*. 2007;170(4):1192-9.
11. Ferenbach DA, Ramdas V, Spencer N, Marson L, Anegon I, Hughes J, et al. Macrophages expressing heme oxygenase-1 improve renal function in ischemia/reperfusion injury. *Mol Ther*. 2010;18(9):1706-13.
12. Ferenbach DA, Nkejabega NC, McKay J, Choudhary AK, Vernon MA, Beesley MF, et al. The induction of macrophage hemeoxygenase-1 is protective during acute kidney injury in aging mice. *Kidney Int*. 2011;79(9):966-76.
13. Bharucha AE, Kulkarni A, Choi KM, Camilleri M, Lempke M, Brunn GJ, et al. First-in-human study demonstrating pharmacological activation of heme oxygenase-1 in humans. *Clin Pharmacol Ther*. 2010;87(2):187-90.
14. Doberer D, Haschemi A, Andreas M, Zapf TC, Clive B, Jeitler M, et al. Haem arginate infusion stimulates haem oxygenase-1 expression in healthy subjects. *Br J Pharmacol*. 2010;161(8):1751-62.
15. de Almeida MC, Silva AC, Barral A, Barral Netto M. A simple method for human peripheral blood monocyte isolation. *Memorias do Instituto Oswaldo Cruz*. 2000;95(2):221-3.
16. Hill-Kapturczak N, Sikorski E, Voakes C, Garcia J, Nick HS, Agarwal A. An internal enhancer regulates heme- and cadmium-mediated induction of human heme oxygenase-1. *Am J Physiol Renal Physiol*. 2003;285(3):F515-23.
17. Hollmen ME, Kyllonen LE, Inkinen KA, Lalla ML, Salmela KT. Urine neutrophil gelatinase-associated lipocalin is a marker of graft recovery after kidney transplantation. *Kidney Int*. 2011;79(1):89-98.
18. van Timmeren MM, Vaidya VS, van Ree RM, Oterdoom LH, de Vries AP, Gans RO, et al. High urinary excretion of kidney injury molecule-1 is an independent predictor of graft loss in renal transplant recipients. *Transplantation*. 2007;84(12):1625-30.
19. Yarlagadda SG, Coca SG, Garg AX, Doshi M, Poggio E, Marcus RJ, et al. Marked variation in the definition and diagnosis of delayed graft function: a systematic review. *Nephrol Dial Transplant*. 2008;23(9):2995-3003.

20. Sharif A, Borrows R. Delayed graft function after kidney transplantation: the clinical perspective. *Am J Kidney Dis.* 2013;62(1):150-8.
21. Blydt-Hansen TD, Katori M, Lassman C, Ke B, Coito AJ, Iyer S, et al. Gene transfer-induced local heme oxygenase-1 overexpression protects rat kidney transplants from ischemia/reperfusion injury. *J Am Soc Nephrol.* 2003;14(3):745-54.
22. Lemos FB, Ijzermans JN, Zondervan PE, Peeters AM, van den Engel S, Mol WM, et al. Differential expression of heme oxygenase-1 and vascular endothelial growth factor in cadaveric and living donor kidneys after ischemia-reperfusion. *J Am Soc Nephrol.* 2003;14(12):3278-87.
23. Tullius SG, Nieminen-Kelha M, Buelow R, Reutzel-Selke A, Martins PN, Pratschke J, et al. Inhibition of ischemia/reperfusion injury and chronic graft deterioration by a single-donor treatment with cobalt-protoporphyrin for the induction of heme oxygenase-1. *Transplantation.* 2002;74(5):591-8.
24. Ollinger R, Kogler P, Biebl M, Sieb M, Sucher R, Bosmuller C, et al. Protein levels of heme oxygenase-1 during reperfusion in human kidney transplants with delayed graft function. *Clin Transplant.* 2008;22(4):418-23.
25. Tsuchihashi S, Livhits M, Zhai Y, Busuttill RW, Araujo JA, Kupiec-Weglinski JW. Basal rather than induced heme oxygenase-1 levels are crucial in the antioxidant cytoprotection. *J Immunol.* 2006;177(7):4749-57.
26. Tenhunen R, Tokola O, Linden IB. Haem arginate: a new stable haem compound. *The Journal of pharmacy and pharmacology.* 1987;39(10):780-6.
27. Martinez F, Kamar N, Pallet N, Lang P, Durrbach A, Lebranchu Y, et al. High dose epoetin beta in the first weeks following renal transplantation and delayed graft function: Results of the Neo-PDGF Study. *Am J Transplant.* 2010;10(7):1695-700.
28. Aydin Z, Mallat MJ, Schaapherder AF, van Zonneveld AJ, van Kooten C, Rabelink TJ, et al. Randomized trial of short-course high-dose erythropoietin in donation after cardiac death kidney transplant recipients. *Am J Transplant.* 2012;12(7):1793-800.
29. Fellstrom B, Holdaas H, Jardine AG, Holme I, Nyberg G, Fauchald P, et al. Effect of fluvastatin on renal end points in the Assessment of Lescol in Renal Transplant (ALERT) trial. *Kidney Int.* 2004;66(4):1549-55.
30. Huang Y, Shan J, Wang C, Ma J, Li D, Li L, et al. Can ischemic preconditioning alone really protect organs from ischemia reperfusion injury in transplantation. *Transplant immunology.* 2009;20(3):127-31.

Tables for HOT document

Table 1. Baseline characteristics of the 40 randomised participants by treatment group

Characteristic	Placebo (n=19)	HA (n=21)
Mean age (range)	52.4 (25- 79)	52.1 (21- 69)
Male (%)	11 (58)	12 (57)
1 st transplant (%)	19 (100)	20 (95)
2 nd transplant (%)	0	1 (5)
Dialysis patient (%)	17 (89)	20 (95)
Drug history; Statin (%)	11 (58)	13 (62)
Drug history; any type antiplatelet agent (%)	8 (39)	6 (29)
Time from infusion to reperfusion in minutes (range)	420.6 (39.0- 1586.0)	249.0 (48.0- 803.0)
Donor characteristics		
Mean age (range)	45.2 (21- 73)	46.8 (14- 69)
Male (%)	13 (72)	10 (53)
DBD (%)	9 (50)	9 (47)
Mode of organ perfusion- hypothermic (%)	3 (17)	2 (11)
Mode of organ perfusion- cold static (%)	15 (83)	17 (89)
HLA match		
0 HLA-DRB1 mismatch	3 (17)	1 (5)
1 HLA- DRB1 mismatch	9 (50)	13 (69)
2 HLA-DRB1 mismatch	6 (33)	5 (27)
Number with virtual crossmatch ¹	12 (67)	15 (79)
Mean CIT in minutes ² (range)	691.7 (351- 1287)	731.5 (450- 1213)
Mean WIT in minutes ³ (range)	36.2 (28- 61)	37.6 (23- 59)

Mean values are given for continuous variables, while numbers of patients (percentages) are given otherwise.

¹Virtual crossmatch= crossmatch that is performed before the organ arrives at RIE. This reduces the ischaemic time and is possible when the recipient's antibody specificities have been identified and the donor HLA type is known.

²CIT= Cold ischaemic time= time from storage in ice at recovery to removal from ice during transplant operation

³WIT= Warm ischaemic time (2nd)= time from removal from ice for transplant to reperfusion

Figure legends for HOT_article document

Figure 1. Flow chart showing recruitment and study characteristics

Figure 2: Treatment with HA upregulated HO-1 protein and mRNA expression in peripheral blood mononuclear cells (PBMCs).

a) HO-1 protein upregulation: expressed as the difference in HO-1 expression on D1 compared with D0. HA upregulated HO-1 expression by 11.1ng/ml [1.0- 37.0], compared with placebo -0.14ng/ml [-0.7- 0.3]; $p < 0.0001$ (n=36, 1 sample could not be analysed). b) HA upregulated HO-1 mRNA expression 2.73 fold [1.8- 3.2] compared with placebo 1.41 fold [1.2- 2.2]; $p = 0.02$ (n=37). c) Change in PBMC HO-1 protein expression over 5 days, demonstrating significantly increased HO-1 expression on D1 and D3 after HA treatment ($p < 0.005$). d) Change in HO-1 mRNA expression over 5 days, expressed as fold increase compared with D0. No significant difference at D5.

Figure 3: There was significant upregulation of HO-1 expression in CD68-positive macrophages after HA treatment compared with placebo at day 5.

a) Number of CD68-positive macrophages at D0 and D5 with HA treatment compared with placebo: D0 $p = 0.956$, D5 $p = 0.13$ (n=21, some tissue not suitable for staining). b) Percentage CD68-positive macrophages expressing HO-1 at D0 and D5 with HA compared with placebo. At D5, HA: 50.81 cells per hpf [40.0- 59.8], placebo: 22.3 [0- 34.8]; $p = 0.012$ (n=21) c) Dual stain immunofluorescent images of renal tissue (green: CD68, red: HO-1, blue: nuclear marker), white arrows highlight dual positive cells from D0 HA d) D0 placebo e) D5 HA and f) D5 placebo (x20 magnification)

Figure 4: HO-1 protein but not mRNA was upregulated in renal tissue.

a) HO-1 protein upregulation: expressed as the difference in % HO-1 expression on D5 compared with D0, HA upregulated protein by 0.21% [0.1- 0.6] compared to placebo of -0.03% [-1.3- 0.1], $p = 0.017$ (n=22, three samples insufficient for protein analysis).

b) HO-1 mRNA upregulation at D5 expressed as fold increase from D0 showed no significant difference between the groups $p=0.45$ ($n=25$). Images of immunohistological staining for HO-1 (DAB) in renal tissue at c) D0 HA, d) D0 placebo, e) D5 HA and f) D5 placebo (x20 magnification)

Figure 5: Change in urinary biomarkers over 5 days after HA treatment and placebo.

There was no significant difference between the HA and placebo treatment at any time point in a) KIM-1 and b) NGAL-1. All p-values $= >0.05$.




Figure 1
[Click here to download high resolution image](#)




Figure 2: Treatment with HA upregulated HO-1 protein and mRNA expression in peripheral blood mononuclear cells (PBMCs).

Figure 3
[Click here to download high resolution image](#)




Figure 3: There was significant upregulation of HO-1 expression in CD68-positive macrophages after HA treatment compared with placebo at day 5.

Figure 4
[Click here to download high resolution image](#)




Figure 4: HO-1 protein but not mRNA was upregulated in renal tissue

Figure 5
[Click here to download high resolution image](#)




Figure 5: Change in urinary biomarkers over 5 days after HA treatment and placebo.



CONSORT 2010 checklist of information to include when reporting a randomised trial*

Section/Topic	Item No	Checklist item	Reported on page No
Title and abstract	1a	Identification as a randomised trial in the title	1
	1b	Structured summary of trial design, methods, results, and conclusions (for specific guidance see CONSORT for abstracts)	4
Introduction Background and objectives	2a	Scientific background and explanation of rationale	5/6
	2b	Specific objectives or hypotheses	5/6
Methods Trial design	3a	Description of trial design (such as parallel, factorial) including allocation ratio	7/8
	3b	Important changes to methods after trial commencement (such as eligibility criteria), with reasons	--
Participants	4a	Eligibility criteria for participants	7
	4b	Settings and locations where the data were collected	7
Interventions	5	The interventions for each group with sufficient details to allow replication, including how and when they were actually administered	7/8
Outcomes	6a	Completely defined pre-specified primary and secondary outcome measures, including how and when they were assessed	6/8/9
Sample size	6b	Any changes to trial outcomes after the trial commenced, with reasons	--
	7a	How sample size was determined	7
Randomisation:	7b	When applicable, explanation of any interim analyses and stopping guidelines	--
	8a	Method used to generate the random allocation sequence	7
Sequence generation	8b	Type of randomisation; details of any restriction (such as blocking and block size)	7
Allocation concealment mechanism	9	Mechanism used to implement the random allocation sequence (such as sequentially numbered containers), describing any steps taken to conceal the sequence until interventions were assigned	7/8
Implementation	10	Who generated the random allocation sequence, who enrolled participants, and who assigned participants to interventions	7/8
Blinding	11a	If done, who was blinded after assignment to interventions (for example, participants, care providers, those	8

	assessing outcomes) and how	
11b	If relevant, description of the similarity of interventions	8
12a	Statistical methods used to compare groups for primary and secondary outcomes	9/10
12b	Methods for additional analyses, such as subgroup analyses and adjusted analyses	—
Results		
13a	Participant flow (a diagram is strongly recommended)	11/ figure 1
13b	For each group, the numbers of participants who were randomly assigned, received intended treatment, and were analysed for the primary outcome	11/ figure 1
14a	Recruitment	11
14b	For each group, losses and exclusions after randomisation, together with reasons	11
15	Baseline data	table 1
16	Numbers analysed	11/ figure 1
17a	Outcomes and estimation	11/12
17b	For each primary and secondary outcome, results for each group, and the estimated effect size and its precision (such as 95% confidence interval)	—
18	Ancillary analyses	—
19	Harms	11
Discussion		
20	Limitations	14/15
21	Generalisability	14
22	Interpretation	13/14/15
Other information		
23	Registration	1/7
24	Protocol	1/7
25	Funding	1/7

*We strongly recommend reading this statement in conjunction with the CONSORT 2010 Explanation and Elaboration for important clarifications on all the items. If relevant, we also recommend reading CONSORT extensions for cluster randomised trials, non-inferiority and equivalence trials, non-pharmacological treatments, herbal interventions, and pragmatic trials. Additional extensions are forthcoming: for those and for up to date references relevant to this checklist, see www.consort-statement.org.



CONSORT 2010 Flow Diagram for HOT study

