

# Heterotrophic microbial colonization of the interior of impact-shocked rocks from Haughton impact structure, Devon Island, Nunavut, Canadian High Arctic

David A. Fike<sup>1</sup>, Charles Cockell<sup>2</sup>, David Pearce<sup>2</sup> and Pascal Lee<sup>3</sup>

<sup>1</sup>Department of Earth, Atmospheric, and Planetary Sciences, Massachusetts Institute of Technology, 77 Massachusetts Ave., 54-812, Cambridge, MA 01239, USA

<sup>2</sup>British Antarctic Survey, High Cross, Madingley Road, Cambridge CB3 0ET, UK

<sup>3</sup>SETI Institute, NASA Ames Research Center, Moffett Field, CA 94035-1000, USA

**Abstract:** The polar desert is one of the most extreme environments on Earth. Endolithic organisms can escape or mitigate the hazards of the polar desert by using the resources available in the interior of rocks. We examined endolithic communities within crystalline rocks that have undergone shock metamorphism as a result of an asteroid or comet impact. Specifically, we present a characterization of the heterotrophic endolithic community and its environment in the interior of impact-shocked gneisses and their host polymict breccia from the Haughton impact structure on Devon Island, Nunavut, Canadian High Arctic. Microbiological colonization of impact-shocked rocks is facilitated by impact-induced fissures and cavities, which occur throughout the samples, the walls of which are lined with high abundances of biologically important elements owing to the partial volatilization of minerals within the rock during the impact. 27 heterotrophic bacteria were isolated from these shocked rocks and were identified by 16S rDNA sequencing. The isolates from the shocked gneiss and the host breccia are similar to each other, and to other heterotrophic communities isolated from polar environments, suggesting that the interiors of the rocks are colonized by microorganisms from the surrounding country rocks and soils. Inductively coupled plasma–atomic emission spectroscopy (ICP-AES), scanning electron microscopy (SEM) and energy dispersive X-ray (EDX) analysis were used to identify the chemical composition of the shocked materials and to document the *in situ* growth of microbes in their interiors. The identification of these heterotrophic communities within impact-shocked crystalline rocks extends our knowledge of the habitable biosphere on Earth. The colonization of the interiors of these samples has astrobiological applications both for considering terrestrial, microbiological contamination of meteorites from the Antarctic ice sheet and for investigating possible habitats for microbial organisms on the early Earth, and more speculatively, on Mars and other planetary bodies.

Accepted 14 November 2002

**Key words:** endolith, Haughton impact structure, microbial colonization, shock-metamorphism, shocked gneiss.

## Introduction

The polar desert is one of the least hospitable biomes on Earth, as it is characterized by ablative winds, low growth temperatures, freeze–thaw cycles, short growing seasons, high ultraviolet-B (UVB) flux, desiccation and oligotrophic (low nutrient) conditions (e.g. Wynn-Williams 2000). In response to these conditions, many microorganisms, known as endoliths or ‘cryptoendoliths’, inhabit the interior pore spaces of rocks (Friedmann & Ocampo 1976; Friedmann 1982). For endoliths, rocks serve two primary functions. First, they provide protection from many of the environmental stresses (fluctuating temperatures, wind ablation, intense UVB radiation) that threaten microbial existence in the polar environment (Nienow & Friedmann 1993; Wynn-Williams

2000; Cockell *et al.* 2001). Secondly, the rocks act as a reservoir for water, nutrients and heat, giving endoliths an advantage over organisms trying to survive outside the lithic environment (Hirsch *et al.* 1988; Vincent 1988; Nienow & Friedmann 1993). As a result, endolithic organisms are able to remain active for significantly longer periods than those living on the surface (Nienow & Friedmann 1993).

Despite inhabiting the interior of a rock, endolithic organisms are not completely isolated from the outside world. Atmospheric circulation entrains microbes and nutrients and subsequently redistributes them globally (Burckle 1995a, b; Burckle & Wasell 1995; Vincent 2000). Wind, flowing water or meltwater from ice and windblown snow can transport and deposit microorganisms and nutrients into cracks and crevices in a rock surface, allowing endolithic communities

to grow and develop (Wynn-Williams 2000). For example, examination of microfossils within four meteorites from the Allan Hills and Queen Alexandra Range, Antarctica has revealed the presence of modern terrestrial freshwater and saltwater diatoms, as well as specimens representing extinct diatom species (Burckle & Delaney 1999). The high degree of exogenous organic matter and microorganisms present in Antarctic samples, despite the relative isolation of Antarctica from global patterns of atmospheric circulation (Vincent 2000), suggests that there would be even greater deposition of nutrients and organisms in other, less isolated environments, including the Arctic.

Since polar endoliths were first characterized (Friedmann & Ocampo 1976), porous sedimentary rocks, such as the Beacon sandstone formation in Antarctica, have been identified as particularly suitable for endolithic communities, because their high porosity favours colonization by endolithic bacteria (Nienow & Friedmann 1993). Most studies have focused on phototrophic communities inhabiting the near-surface environment (Friedmann & Ocampo 1976; Friedmann 1982; Vincent 1988; Nienow & Friedmann 1993), although a few have noted the importance of heterotrophic bacteria in these near-surface endolithic communities (Hirsch *et al.* 1988; Siebert & Hirsch 1988; Nienow & Friedmann 1993). While crystalline rocks, such as gneiss and granite, have also been studied in Antarctica (Nienow & Friedmann 1993), their typically low porosities make them poor hosts for endolithic bacteria. However, the Arctic presents a unique environment in which endolithic organisms can readily colonize crystalline rocks: impact-shocked gneisses and their host polymict breccias from the Haughton impact structure on Devon Island in the Canadian High Arctic. The colonization of the heavily shocked gneiss by cyanobacteria in preference to unshocked or low-shocked gneisses at Haughton has been described previously (Cockell & Lee 2000; Cockell *et al.* 2002) and previous studies have determined the chemical compositions and shock levels of these rocks (Metzler *et al.* 1988; Osinski & Spray 2001). These rocks have undergone impact-induced shock metamorphism, one of the many effects of an asteroid or comet impact, are heavily fractured, and some of their constituent minerals have been partially volatilized (Dressler & Reimold 2001). This has generated habitats, which can be exploited by significant populations of endolithic organisms (Cockell & Lee 2000; Cockell *et al.* 2001).

In this study we describe the characterization of the heterotrophic communities within impact-shocked gneisses and their host breccia from Haughton and we discuss the implications of our findings for the contamination of highly shocked meteorites in Antarctica and the search for signatures of life on Mars and other planetary bodies.

## Materials and methodology

### Study site

The Haughton impact structure is located at 75° 22' N, 89° 41' W on Devon Island, Nunavut in the Canadian High



**Fig. 1.** Geographical location. (a) Devon Island, Nunavut, in the Canadian High Arctic; (b) the Haughton Crater is located at 75° 22' N, 89° 41' W in the northwestern portion of Devon Island; (c) synthetic aperture radar image of the Haughton crater. The crater is approximately 24 km in diameter. Image obtained from Grieve (1988).

Arctic (Fig. 1). The impact structure is approximately 24 km in diameter (Grieve 1988) and was created by the impact of an asteroid or comet  $23.4 \pm 1.0$  Ma during the Early Miocene (Jessberger 1988). At the time of the impact, the Precambrian gneisses forming the crystalline basement on Devon Island were overlain by ~1750 m of Paleozoic sedimentary rocks, predominantly dolomite and limestone (Frisch & Thorsteinsson 1978). The presence of significant amounts of gneiss cropping out in the melt breccia deposits indicates that the excavation depth for the impact structure exceeded 1750 m.

Haughton is set in arid polar desert, with less than 5% vegetation cover (Cockell *et al.* 2001). On Devon Island, the summer growing season is short, with an average number of 188 degree-days in July (Gold 1988). The soils at and around

the Haughton impact structure are primarily dolomitic and oligotrophic (Bliss *et al.* 1994). Bacterial numbers within the soil average  $3\text{--}4 \times 10^6$  colony-forming units per gram of dry-weight soil (Cockell *et al.* 2001).

#### Sample collection


Samples of unshocked and shocked gneiss and host melt breccia considered in this study were aseptically collected from the exposed surface of the Haughton impact structure during the 1999–2001 summer field seasons of the NASA Haughton-Mars Project. All samples were between 7–15 cm in size and were retrieved from two locations within the impact structure: an isolated hill of impact melt rocks at  $75^\circ 24.53' \text{N}$ ,  $89^\circ 49.76' \text{W}$  ('Wyle Labs Hill', at 'Lake Trinity'); and an escarpment of melt breccia deposits at  $75^\circ 23.9' \text{N}$ ,  $89^\circ 31.6' \text{W}$  ('Bruno Escarpment'). In this study, the term 'unshocked' gneiss corresponds to those samples exposed to a maximum shock pressure of less than 5 GPa and 'shocked' gneiss to those samples exposed to between 25 and 45 GPa. Samples with shock levels in the 5–25 GPa range were not considered in this study. All samples were collected in sterile bags, transported in an ice cooler at approximately  $0^\circ \text{C}$ , and curated at the British Antarctic Survey (BAS) where they were stored at  $-20^\circ \text{C}$ .

#### Physical characterization of samples

Inductively coupled plasma–atomic emission spectroscopy (ICP-AES) was applied to samples to determine their bulk composition using the Optima 3300RL inductively coupled plasma–atomic emission spectrometer (Perkin Elmer, Shelton, CT) at Royal Holloway, University of London. Three samples of unshocked gneiss, three samples of breccia and five samples of shocked gneiss were analysed in duplicate. Samples were prepared following the methods of Totland *et al.* (1992), which include a potassium hydroxide fusion to determine major element concentrations and a hydrofluoric acid (HF) digest to determine trace element concentrations.

Scanning electron microscopy (SEM) was used to examine interior fragments of samples to identify both the micron-scale effects of the impact-shock process on the samples and microorganisms living within fissures and cavities of the samples. Hand samples were broken open and interior fragments showing no sign of surface weathering were chosen for study. These fragments (with dimensions of approximately  $0.5 \times 0.5 \times 0.5 \text{ cm}^3$ ) were mounted on SEM stubs with epoxy adhesive after aseptic preparation in the laboratory. These were then coated with gold for secondary electron SEM imaging using a Stereoscan S360 scanning electron microscope (Cambridge Instruments (LEO), Cambridge, UK) fitted with a PCIT image capture system and stored as standard computer TIFF files.

Energy-dispersive X-ray (EDX) microanalysis was applied to samples of both unshocked and shocked gneiss to determine their average surface composition (over  $10^4 \mu\text{m}^2$ ) and its micron-scale variations (transects of ten points, each covering  $1 \mu\text{m}^2$  and separated by  $5 \mu\text{m}$ ). Hand samples of unshocked



**Fig. 2.** Isolation of bacteria from interior fragments of shocked gneiss. (a) Cultivation of bacteria from the interior of shocked gneiss samples; (b) close-up showing colony formation only around fragments of the shocked gneiss.

and shocked gneiss were cut to expose their interiors. Samples for EDX analysis were taken from the centre of the samples, away from any sign of surface weathering. These samples were then thin-sectioned, polished and carbon coated for X-ray analysis using an Oxford Instruments EDX/INCA energy X-ray microanalysis system (Oxford Instruments, Oxford, UK) fitted with a germanium detector.


#### Isolation and identification of heterotrophic bacteria

12 samples of shocked gneiss and of host breccia were broken open inside sterile bags using a rock hammer. Fragments from the interior of the sample were carefully removed under sterile conditions in a microbiological flow cabinet and scraped using a sterile blade into a Petri dish containing 4% tryptone-soy agar (TSA) (Difco Ltd, UK). After 2 days incubation at  $20^\circ \text{C}$ , colonies had begun to grow around the pieces of rock fragments (Fig. 2). The colonies were subsequently streaked onto TSA plates until single isolates had been obtained. These colonies were maintained on TSA sloped agar tubes.

Polymerase chain reaction (PCR) was employed, using primers 8F (AGAGTTTGATCCTGGCTCAG) and 1500R







**Fig. 3.** SEM images. (a) Fragment of unshocked gneiss. Scale bar = 10  $\mu\text{m}$ . (b) Fragment of shocked gneiss. Note the surface variation, particularly the abundant vesicles, which can serve as microbial habitats. Scale bar = 100  $\mu\text{m}$ . (c) Fragment of breccia. Note the surface complexity drastically hinders visual identification of microbes. Scale bar = 20  $\mu\text{m}$ . (d) Cluster of three rod-shaped bacteria. Scale bar = 5  $\mu\text{m}$ . (e) Small group of coccoid-shaped bacteria. Note the EPS covering that seems to connect them. Scale bar = 5  $\mu\text{m}$ . (f) Biofilm of coccoid bacteria completely covering the interior surface of an open cavity. Scale bar = 20  $\mu\text{m}$ .


to the surface of unshocked gneiss (Fig. 4a) and the surface of shocked gneiss distant from surface cavities. The slides with isolate G20 had clearly biological spectra, for which the intensity relative to the silicate background increased in proportion to the thickness of the bacteria on the slide. The slide with the lowest bacterial density (Fig. 4d) and the surface cavities of shocked gneiss are observed to have approximately the same density of bacteria. The occurrence of chlorine in the spectra, observed even under the thinnest layer of bacteria, was the clearest and most sensitive indicator of the presence of bacteria using EDX.

EDX transects conducted along the surface of unshocked (Fig. 5a) and shocked (Fig. 5b) gneiss reveal impact-induced changes in surface composition and illustrate the microbial

habitats created by micro-scale variations in the surface of the shocked gneiss. The surface of unshocked gneiss showed relatively little variation in surface composition, even on a micron scale. However, the surface composition of the shocked gneiss varied considerably even over a few microns, and dramatically near or inside a surface cavity (right of Fig. 5b), which again was characterized by high abundances of biologically important elements, including P, Mg, Cl and S.

#### *Isolated heterotrophic bacteria*

A total of 27 bacteria were isolated from the interiors of samples of shocked rocks (14 from shocked gneiss, 13 from breccia). These isolates were selected, along with three isolates from Antarctic soil as a comparison, to undergo sequencing




**Fig. 4.** X-ray spectra. (a) Surface of unshocked gneiss. The spectra are dominated by Si, O, Fe and Al peaks. (b) Interior of a cavity on the surface of shocked gneiss. The spectra contain the biologically important elements Ca, Cl, Mg and K, in addition to those found in (a). For comparison (c) the surface of a sterile glass slide and (d) bacterial isolate G20 streaked onto a glass slide. The elemental signature of the glass slide is still visible. Note the similarity between unshocked gneiss (a) and the glass slide (c) and between the shocked gneiss (b) and isolate G20 (d).

of their 16S rDNA genes. This sequencing process was successful for 13 out of 14 isolates from the shocked gneiss samples, for all 13 isolates from the breccia samples and for all three of the Antarctic soil isolates. Isolate G20 from the gneiss sample did not amplify properly during the PCR stage, preventing a proper sequence from being obtained. This isolate was observed to possess a thick waxy coating when observed under an optical microscope. This waxy coating is the likely cause of the failure of the amplification process because excess polysaccharides have been shown to inhibit the PCR reaction (Demeke & Adams 1992; Do & Adams 1991). Of the 30 isolates successfully sequenced, 24 were significantly similar to sequences found in the NCBI nucleotide database and have been identified to the species level (see Table 3). The sequences for the remaining six isolates (four from shocked gneiss (G11, G0, G13, G14) and two from breccia (B8, B11)) showed no significant similarity to any sequence within the

database (having matches that, at best, were less than 20 base pairs in length).

The identification of our isolates with known species in the nucleotide database provides for the characterization of bacteria according to phylogeny, habitat and metabolism. Bacteria isolated from the shocked gneiss and host breccia were phylogenetically similar to each other, and to those isolated from Antarctic soil (Fig. 6). These isolates are prominently soil, ice or freshwater bacteria (Table 4). Members of the genera *Arthrobacter*, *Bacillus*, *Janthinobacter* and *Pseudomonas* are frequently found as soil bacteria. *Arthrobacter* species are common soil bacteria and all *Arthrobacter* species identified in this study have previously been isolated from soil as well as from glacial ice (Keddie *et al.* 1984). In addition, *A. nicotiana* has been found in cave silts, glacial silts, sewage, air and tobacco plants, and *A. sulfurous* has previously been isolated from oil brines (Keddie *et al.*



**Fig. 5.** X-ray transects across gneiss samples. Major elements are shown on the top and trace elements on the bottom of each graph. Each point sampled an area of approximately  $1 \mu\text{m}^2$ . Average values were obtained by integrating over a  $100 \mu\text{m}$  square centred on the transect and are shown for each element (where no average is shown, the integration did not detect that element). (a) Unshocked gneiss: the composition is primarily of silicon and iron oxides and shows little variation. (b) Shocked gneiss: the composition varies significantly across the transect, particularly within a surface cavity (located at  $x=40 \mu\text{m}$  on the distance axis). On flat surface areas the composition is primarily silicon, iron and aluminium oxides. Within the exposed surface cavity, peaks in the concentrations of sodium, chlorine, phosphorus and sulphur are observed. These elemental signatures are consistent with the X-ray spectra for biological samples (bacterial isolate G20 in Fig. 4d).

1984). *Bacillus* species are widely distributed in nature and *B. psychrophilus* has been isolated from soil and river waters (Claus & Berkeley 1984). *Janthinobacter lividum* is a common soil and freshwater organism, although it is most common in temperate climates (Sneath 1984). *Pseudomonas* species are common components of soil and freshwater bacterial communities. All *Pseudomonas* spp. isolated in this study have been found in freshwater habitats and *P. borealis*, *P. sp. IC038* and *P. psychrophilia* have also been isolated from soil environments (Palleroni 1984). Three bacteria were found that are not characteristically isolated from soil: *Caulobacter bacteroides*, typically isolated from freshwater habitats

(Abraham et al. 1999); *Planococcus citreus*, a marine organism, frequently isolated from polar waters (Kocur 1984); and *Stenotrophomonas maltophilia*, which was isolated from both the shocked gneiss and the breccia, customarily isolated from clinical specimens and from milk, water and frozen foods (Palleroni 1984).

The three isolates from Antarctic soil samples (two species of *Arthrobacteria* and one species of *Bacillus*) were similar to those isolated from shocked gneiss and breccia (see Table 4). *Arthrobacter* sp. CAB1 (Morikawa et al. 2002) had previously been isolated from 200 year old glacial ice in China (Christner 2002) and from agricultural soils (Lukow 1999). *Arthrobacter*




Table 3. *Phylogenetic Identification of Isolates from Shocked Gneiss, Breccia, and Antarctic Soil. The nearest phylogenetic species to our isolates is shown down to the species level, with the percent match between the sequence for our isolates and that found in the GenBank database. Novel indicates that there are no significant matches between the isolate sequence and any species in the database. N/A = sample G20 did not amplify properly in PCR*

	Phylum	Class	Genus	Species	% Match	Matches (bp)
<b>Shocked Gneiss</b>						
G4	<i>Actinobacteria</i>	<i>Actinobacteria</i>	<i>Arthrobacter</i>	<i>nicotiana</i>	98.5%	460
G50	<i>Firmicutes</i>	<i>Bacilli</i>	<i>Planococcus</i>	<i>citreus</i>	99.3%	456
G21	<i>Firmicutes</i>	<i>Bacilli</i>	<i>Bacillus</i>	<i>psychrophilus</i>	97.2%	447
G23	<i>Proteobacteria</i>	$\beta$ - <i>Proteobacteria</i>	<i>Janthinobacter</i>	<i>lividum</i>	99.4%	464
G10	<i>Proteobacteria</i>	$\gamma$ - <i>Proteobacteria</i>	<i>Pseudomonas</i>	<i>rhodesiae</i>	99.8%	473
G22	<i>Proteobacteria</i>	$\gamma$ - <i>Proteobacteria</i>	<i>Pseudomonas</i>	<i>borealis</i>	98.9%	465
G28	<i>Proteobacteria</i>	$\gamma$ - <i>Proteobacteria</i>	<i>Pseudomonas</i>	<i>grimontii</i>	99.6%	781
G16	<i>Proteobacteria</i>	$\gamma$ - <i>Proteobacteria</i>	<i>Stenotrophomonas</i>	<i>maltophilia</i>	99.3%	761
G15	<i>Proteobacteria</i>	$\gamma$ - <i>Proteobacteria</i>	<i>Stenotrophomonas</i>	<i>maltophilia</i>	99.6%	465
G11			Novel			
G0			Novel			
G13			Novel			
G14			Novel			
G20			N/A			
<b>Breccia</b>						
B10	<i>Actinobacteria</i>	<i>Actinobacteria</i>	<i>Arthrobacter</i>	<i>globiformis</i>	100.0%	474
B24	<i>Actinobacteria</i>	<i>Actinobacteria</i>	<i>Arthrobacter</i>	<i>nicotiana</i>	98.9%	470
B16	<i>Actinobacteria</i>	<i>Actinobacteria</i>	<i>Arthrobacter</i>	<i>nicotiana</i>	98.9%	470
B4	<i>Actinobacteria</i>	<i>Actinobacteria</i>	<i>Arthrobacter</i>	<i>sulfureus</i>	99.2%	730
B20	<i>Proteobacteria</i>	$\alpha$ - <i>Proteobacteria</i>	<i>Caulobacter</i>	<i>bacteriodes</i>	96.8%	477
B12	<i>Proteobacteria</i>	$\gamma$ - <i>Proteobacteria</i>	<i>Pseudomonas</i>	<i>IC038</i>	99.8%	479
B15	<i>Proteobacteria</i>	$\gamma$ - <i>Proteobacteria</i>	<i>Pseudomonas</i>	<i>psychrophilia</i>	97.1%	706
B17	<i>Proteobacteria</i>	$\gamma$ - <i>Proteobacteria</i>	<i>Pseudomonas</i>	<i>rhodesiae</i>	98.1%	471
B21	<i>Proteobacteria</i>	$\gamma$ - <i>Proteobacteria</i>	<i>Pseudomonas</i>	<i>rhodesiae</i>	93.3%	431
B3	<i>Proteobacteria</i>	$\gamma$ - <i>Proteobacteria</i>	<i>Stenotrophomonas</i>	<i>maltophilia</i>	99.0%	489
B6	<i>Proteobacteria</i>	$\gamma$ - <i>Proteobacteria</i>	<i>Stenotrophomonas</i>	<i>maltophilia</i>	96.8%	479
B8			Novel			
B11			Novel			
<b>Antarctic Soil</b>						
S13	<i>Actinobacteria</i>	<i>Actinobacteria</i>	<i>Arthrobacter</i>	<i>CAB1</i>	95.9%	772
S16	<i>Actinobacteria</i>	<i>Actinobacteria</i>	<i>Arthrobacter</i>	<i>S23H2</i>	95.8%	668
S21	<i>Firmicutes</i>	<i>Bacilli</i>	<i>Bacillus</i>	<i>psychrophilus</i>	98.0%	448

*sp.* S23H2 had previously been isolated from Antarctic sea ice brine and cryoconite holes (Junge *et al.* 1998). *Bacillus psychrophilus*, also isolated from shocked gneiss, had previously been isolated from soil and freshwater, including from a fiord in Greenland (Suzuki & Yamasato 1994).

The bacteria identified based on their sequence information all possess relatively similar metabolisms and physiological properties. All of the species identified are heterotrophs (chemoorganotrophs), although some species of *Pseudomonas* and *Bacillus* are known to be facultative chemolithoautotrophs (Claus & Berkeley 1984; Palleroni 1984). It has not been determined whether the *Pseudomonas spp.* or *Bacillus psychrophilus* identified in this study possess the ability to be facultative chemolithoautotrophs. All of the identified bacteria are motile, with the exception of the *Arthrobacter spp.*, which are non-motile (Palleroni 1984). The remaining bacteria are motile through a combination of polar (*Pseudomonas*, *Stenotrophomonas*, *Planococcus*), polar and subpolar/lateral (*Janthinobacter*) or peritrichous (*Bacillus*) flagella (Claus & Berkeley 1984; Keddie *et al.* 1984; Kocur 1984;

Palleroni 1984; Sneath 1984). Most of the bacteria isolated are obligate aerobes, notable exceptions include select species of *Pseudomonas*, which can replace oxygen with nitrate as a terminal electron acceptor, allowing anaerobic growth (Palleroni 1984), and species of *Bacillus*, which can replace oxygen with a variety of different terminal electron receptors, again allowing for anaerobic growth (Claus & Berkeley 1984). Facilities were not available to determine whether the *Pseudomonas spp.* or *Bacillus psychrophilus* identified in this study can function as facultative anaerobes. All of the isolated bacteria have strictly respiratory and never fermentative metabolisms (Claus & Berkeley 1984; Keddie *et al.* 1984; Kocur 1984; Palleroni 1984; Sneath 1984). All of the species are nutritionally non-exacting, either requiring no growth factors (*Pseudomonas spp.*, *Janthinobacter lividum*, *Bacillus psychrophilus*, *Planococcus citreus*) or possessing basic requirements (*Stenotrophomonas maltophilia* usually requires methionine or cystine, *Arthrobacter spp.* require biotin) (Claus & Berkeley 1984; Keddie *et al.* 1984; Kocur 1984; Palleroni 1984; Sneath 1984).



**Fig. 6.** Phylogenetic distribution of isolated bacteria. (a) Breccia; (b) shocked gneiss; (c) Antarctic soil. Bacteria of the genus *Arthrobacter* were isolated from all three types of samples. *Bacillus* and *Pseudomonas* species make up the majority of the remaining bacteria identified. *Stenotrophomonas* species, while found in both the shocked gneisses and breccia from Houghton, were not isolated from the Antarctic soil.

## Discussions

### Geological interpretation

An examination of the composition of gneiss yields insights into the effects of shock metamorphism. Overall, the surface composition as determined by EDX shows a significant disparity when compared with the bulk composition identified by ICP-AES. The ICP-AES bulk composition indicates that there are significantly higher concentrations of  $\text{Fe}_2\text{O}_3$ ,  $\text{Al}_2\text{O}_3$ ,  $\text{MgO}$ ,  $\text{K}_2\text{O}$ ,  $\text{Na}_2\text{O}$ ,  $\text{TiO}_2$  and  $\text{P}_2\text{O}_5$ , as well as of all trace elements, in samples of unshocked gneiss. The elevated concentration of  $\text{MgO}$  (9%) and  $\text{CaO}$  (23%) in the breccia and the high volatile concentration (up to 40%) agree with those measured by Metzler *et al.* (1988), and probably reflect the incorporation of dolomite into the breccia.

EDX spectra reveal the surface composition of unshocked gneiss to be composed almost entirely of silicon, oxygen, iron and aluminium, whereas the shocked samples have surface compositions containing less iron and aluminium and significant amounts of calcium, magnesium, phosphorus, potassium, sulphur and chlorine. This disparity may be explained by impact-induced shock metamorphism. First, as the shock wave resulting from the impact propagates through samples, darker, more refractory minerals are preferentially volatilized as they absorb a greater fraction of the energy incident upon them. This preferential volatilization tends to

remove darker metallic oxides, particularly  $\text{Fe}_2\text{O}_3$  and  $\text{Al}_2\text{O}_3$  (Bunch *et al.* 1998). This process creates vesicles and fissures as near- and sub-surface pockets of these minerals volatilize. Furthermore, as magnesium, phosphorus, potassium and sodium oxides are shock-volatilized, they may have left behind thin and/or patchy coatings on the interior of these vesicles. This might account for the abundance of these latter minerals localized around vesicles walls of the shocked samples and their absence from surfaces of unshocked gneiss, despite the actual bulk percentage of these minerals being significantly less in shocked samples than in unshocked samples. Thus, the impact-induced shock metamorphism has a 'chemical gardening' effect on interior composition, resulting in the enrichment of biologically important elements in vesicle walls and fissures of the shocked gneiss. The relative bulk abundance of biologically important elements such as sulphur, calcium and chlorine, could not be determined between unshocked and shocked gneiss because the ICP-AES analysis was not sensitive to them. However, we suspect that such nutrients are deposited from meltwater into impact-generated cavities, which would encourage endolithic heterotrophic bacteria to inhabit them.

### Biological interpretations

The heterotrophic bacteria isolated from samples in this study are similar to heterotrophic communities identified from other polar environments, including Antarctic soil samples (Roser *et al.* 1983), marine and sea ice samples (Delille 1992; Bowman *et al.* 1997), and Antarctic lacustrine samples (Franzmann *et al.* 1990; Nichols *et al.* 1995). Specifically, representatives of the genera *Pseudomonas*, *Arthrobacter*, *Bacillus* and *Planococcus*, which together constitute 70% of all bacteria identified in this study, were identified as major constituents of each of these other polar microbial communities. Two species, *Arthrobacter nicotiana* and *Planococcus citreus*, were found both in this study and in marine and sea ice samples (Bowman *et al.* 1997).

The similarity between bacteria identified from these diverse environments raises the question of the ultimate origin of the communities. Are these bacteria truly cosmopolitan, or are they endemic to a particular environment and have been transported into the others? The identification of the bacterial isolates from the shocked gneiss reveals that the majority are species associated with soil, ice or freshwater and are commonly isolated from the Antarctic. The ready transport of soil bacteria to marine or sea ice environments has been demonstrated by studies of Antarctic soils (Delille 1987, 1990; Roser *et al.* 1993; Rotert *et al.* 1993; Bowman *et al.* 1997). Similarly, studies have also documented ice bacteria isolated from soil samples (Roser *et al.* 1993; Bowman *et al.* 1997). The common occurrence of genera and even species in ecologically and geographically diverse locations supports the notion of global transport and distribution of bacteria, by vectors such as atmospheric circulation and meltwater flow. As mentioned earlier, precipitation, wind and snowmelt have been identified as means of inoculating rock interiors with bacteria from the surrounding environment (Vincent 2000),

Table 4. *Known Habitats of Identified Bacteria. This lists the species identified in this study through the sequencing of the genes for their 16S rRNA and whether they were found (+) or absent (–) in the samples of shocked gneiss, breccia, and Antarctic soil, and provides a list of other known habitats of the identified organisms*

Genus	Species	Shocked		Antarctic Soil	Other Known Habitats*
		Gneiss	Breccia		
<i>Arthrobacter</i>	<i>nicotiana</i>	+	+	–	Soil/Glacial Ice
<i>Arthrobacter</i>	<i>globiformis</i>	–	+	–	Soil/Glacial Silts
<i>Arthrobacter</i>	<i>sulfurous</i>	–	+	–	Soil/Glacial Ice
<i>Arthrobacter</i>	<i>CAB1</i>	–	–	+	Soil/Glacial Ice
<i>Arthrobacter</i>	<i>S23H2</i>	–	–	+	Sea Ice Brine
<i>Bacillus</i>	<i>psychrophilus</i>	+	–	+	Soil/Freshwater
<i>Caulobacter</i>	<i>bacteriodes</i>	–	+	–	Freshwater
<i>Janthinobacterium</i>	<i>lividum</i>	+	–	–	Soil/Freshwater
<i>Planococcus</i>	<i>citreus</i>	+	–	–	Sea Ice/Glacial Ice
<i>Pseudomonas</i>	<i>borealis</i>	+	–	–	Soil/Freshwater
<i>Pseudomonas</i>	<i>grimontii</i>	+	–	–	Freshwater
<i>Pseudomonas</i>	<i>IC038</i>	–	+	–	Soil/Freshwater
<i>Pseudomonas</i>	<i>psychrophilia</i>	–	+	–	Soil/Freshwater
<i>Pseudomonas</i>	<i>rhodesiae</i>	+	+	–	Freshwater
<i>Stenotrophomonas</i>	<i>maltophilia</i>	+	+	–	Clinical/Frozen Foods

\* See text for references.

and it is believed that a combination of these three factors is responsible for delivering the isolated bacteria to the interior of the samples discussed in this study.

The similarity between the bacteria isolated from within the breccia and the shocked gneiss, and to those isolated from soil, strengthens the argument that the bacteria inhabiting these rocks are derived from the surrounding soil. It is estimated that  $3-4 \times 10^6$  colony-forming units of heterotrophic bacteria inhabit the soil in the area from which the samples were collected (Cockell *et al.* 2001). While the argument could be made that the bacteria isolated are evidence of surface contamination in the process of extracting the interior fragments, thereby explaining both the presence of complex heterotrophic communities in the interior of these rocks and the similarities between the bacteria inhabiting the breccia and the shocked gneiss, we believe this not to be the case for several reasons. First, careful attention was given to selecting interior fragments to be used for obtaining the isolates. The original rocks (approximately 10 cm in diameter) were broken into several centimetre-sized pieces, from which only fragments showing fresh break marks and with no evidence of surface weathering were chosen. These fragments were then broken open again, with millimetre or sub-millimetre fragments from the centre used to isolate bacteria from the samples. During the isolation process, growth was observed to occur only in those areas that surrounded these fragments (see Fig. 2). As a result, the likelihood that surface materials contaminated these fragments is deemed to be insignificant. A stronger argument against surface contamination comes from the lack of phototrophic organisms isolated in our study. The outer surfaces of both shocked gneiss and breccia were observed to have visible coatings of cyanobacteria and other photosynthetic organisms. Further, Cockell *et al.* (2002) identified endolithic phototrophs living in the upper few millimetres of similar rocks. Phototrophic bacteria would be present in any surface and near-surface

contamination, and their complete absence in the isolates obtained from both the shocked gneiss and breccia samples indicates that surface contamination is not likely to have had a significant impact upon our results.

#### *Relevance to meteorites*

The microbial colonization of impact-shocked crystalline rocks has particular relevance to the study of meteorites, which are often highly shocked and in many cases covered by a fusion crust that is formed during entry into the Earth's atmosphere (e.g. Melosh 1989; Steele *et al.* 2000; Dressler & Reimold 2001). This fragile crust is frequently fractured and weathered, and may serve to trap dust particles (including microfossils) that have been transported into the proximity of the meteorites, as has been observed with Antarctic meteorites (Burckle & Delaney 1999; Steele *et al.* 2000). In addition to microfossils and microorganisms found on meteorite surfaces (Burckle 1995a, b; Burckle & Wasell 1995), detailed microscopic investigations of the interior of the Allan Hills 84001 Martian meteorite suggest that it may be contaminated by endolithic microorganisms of terrestrial origin (Steele *et al.* 2000). The entrapment of microorganisms by meteorites is likely to be a ubiquitous process in Antarctica (Burckle & Delaney 1999) and possibly elsewhere.

The presence of microfossils in the surface and near-surface environments of meteorites clearly demonstrates their susceptibility to terrestrial contamination. Our examination of shocked rocks from the Haughton impact structure provides insight into the extent to which, and the mechanism by which, their interiors have been colonized by heterotrophic bacteria. The prime factor facilitating this colonization appears to be the impact-induced porosity of the shocked rocks. The alteration of the chemical composition of vesicle walls to include higher abundances of biologically important elements, as a result of the partial volatilization of the constituent minerals, is also believed to favour colonization.

As many meteorites present evidence of shock, often to levels well in excess of those exhibited in the samples examined in this study, they would be natural hosts for heterotrophic colonization. The source for microbial contamination in Antarctic meteorites could be either deposition of wind-transported microorganisms or, more indirectly, microorganisms derived from the ice immediately surrounding the meteorite. It has been observed that the Antarctic ice sheet contains a variety of heterotrophic bacteria (Hirsh *et al.* 1988; Franzmann 1996; Kellogg & Kellogg 1996). Furthermore, the albedo difference between the meteorite and the surrounding ice can cause the surrounding ice to melt when the meteorite is heated by solar radiation during the Antarctic summer. Low-albedo rocks, debris and organic material (cryoconite) have been observed to sink into Antarctic blue ice as a result of seasonal thawing and refreezing, becoming trapped in regelation portholes (Lee *et al.* 1999). Here, the seasonal availability of meltwater could help concentrate any heterotrophs present in the nearby ice, providing a mechanism to inoculate the interior of meteorites with heterotrophs.

### Life on Mars

The observed impact-induced generation of habitats in crystalline rocks also has relevance to astrobiological questions of life on other planetary bodies with impact-shocked surface materials. This is particularly true of Mars, where as a result of the lack of global plate tectonics, the age of exposed surface materials could possibly exceed 4.5 Gyr (radiometric crystallization age of the ALH-84001 meteorite), a time when the impact rate was significantly higher than its present value. Were life to have evolved on Mars, impact-shocked rocks would have provided ready lithic habitats. As liquid water became increasingly scarce at the Martian surface, the interior of impact-shocked rocks could have served as a reservoir for water, heat and nutrients for any microbial colonies present. Thus, the interior of impact-shocked rocks in the Martian regolith, particularly that portion included at present in the Martian permafrost, might be among the best places to search for fossilized remnants of any past Martian life. Continuous permafrost, such as is found approximately 40 cm below the surface at Haughton and throughout much of the high Arctic, offers an environment that helps preserve an extended record of microbial endolithic colonization. Such preservation has been observed in Siberian permafrost, where ground ice and soils have been found to contain viable microorganisms that are probably 2–3 Myr old (Shi *et al.* 1997).

Recent observations from Mars Odyssey reveal concentrations of hydrogen in the upper few metres of large portions of the Martian subsurface, which may be water ice-rich layers on the scale of tens of centimetres (Mitrofanov *et al.* 2002). In such regions, ice may constitute  $35\% \pm 15\%$  by weight of the subsurface layer (Boynton *et al.* 2002). If these regions of high hydrogen concentration represent water ice content, then they would be ideal places to search for organic preservation. On Mars and beyond, as impacts are a universal process of planetary formation and evolution, we speculate that the interior of impact-shocked rocks might

offer a common environmental setting favourable for the development, survival and preservation of microbial life were it to arise on planetary bodies outside the Earth.

### References

- Abraham, W. *et al.* (1999). Phylogeny and polyphasic taxonomy of *Caulobacter* species. *Int. J. Systemic Bacteriol.* **49**, 1053–1073.
- Altschul, S., Madden, T., Schäffer, A., Zhang, J., Zhang, Z., Miller, W. & Lipman, D. (1997). Gapped BLAST and PSI-BLAST: new generation of protein database search programs. *Nucleic Acids Res.* **25**, 3389–3402.
- Bliss, L., Henry, G., Svoboda, J. & Bliss, D. (1994). The patterning of plant communities and edaphic factors along a High Arctic coastline: implications for succession. *Can. J. Botany* **72**, 1095–1107.
- Bowman, J., McGammon, S., Brown, M., Nichols, D. & McMeekin, T. (1997). Diversity and association of psychrophilic bacteria in Antarctic sea ice. *Appl. Environ. Microbiol.* **63**, 3068–3078.
- Boynton, W.V., Feldman, W.C., Squyres, S.W., Prettyman, T.H., Bruckner, J., Evans, L.G., Reedy, R.C., Starr, R., Arnold, J.R., Drake, D.M., Englert, P.A.J., Metzger, A.E., Mitrofanov, I., Trombka, J.I., d'Uston, C., Wanke, H., Gasnault, O., Hamara, D.K., Janes, D.M., Marcialis, R.L., Maurice, S., Mikheeva, I., Taylor, G.J., Tokar, R. & Shinohara, C. (2002). Distribution of hydrogen in the near surface of Mars: evidence for subsurface ice deposits. *Science* **297**(5578), 81–85.
- Bunch, T.E., Grieve, R.A.F., Lee, P., McKay, C.P., Rice, J.W., Schutt, J.W. & Zent, A. (1998). Haughton-Mars 97 II, preliminary observations on highly shocked crystalline basement rocks from the Haughton impact crater. *Lunar Planetary Science Conf. XXIX*, pp. 1307–1308. Houston, TX: Lunar and Planetary Institute.
- Burckle, L. (1995a). Pliocene-Pleistocene diatoms in Devonian to Cretaceous sedimentary and igneous rocks from Antarctica: a paradigm reclaimed. *Antarctic J.* **30**, 2–3.
- Burckle, L. (1995b). Diatoms in igneous and metamorphic rocks from Queen Maud Land, East Antarctica: improbable places, improbable surfaces. *Antarctic J.* **30**, 67–68.
- Burckle, L. & Delaney, J. S. (1999). Terrestrial microfossils in antarctic ordinary chondrites. *Meteorit. Planet. Sci.* **34**, 475–478.
- Burckle, L. & Wasell, A. (1995). Marine and freshwater diatoms in sediment pockets in igneous rocks from James Ross Island, Antarctica. *Antarctic J.* **30**, 8–9.
- Christner, B. (2002). 16S rRNA Sequence, GAPPED-BLAST Database, #AF479340.
- Claus, D. & Berkeley, R. (1984). *Genus Bacillus*. In *Bergey's Manual of Systematic Bacteriology*, ed. Sneath, P., Vol. 2, pp. 1105–1139. Baltimore: Williams & Wilkins.
- Cockell, C. & Lee, P. (2000). *Impact-Induced Formation of a Microbial Habitat*. 1st *Astrobiology Science Conf.*, NASA Ames Research Center, April 3–5, 2000. Reston, VA: Reston Communications.
- Cockell, C., Lee, P., Schruerger, A., Hidalgo, L., Jones, J. & Stokes, M. (2001). Microbiology and vegetation of micro-oases and polar desert, Haughton impact crater, Devon Island, Nunavut, Canada. *Arctic Antarctic Alpine Res.* **33**, 306–318.
- Cockell, C., Lee, P., Osinski, G., Horneck, G. & Broady, P. (2002). Impact-induced formation of microbial lithic habitats. *Meteorit. Planet. Sci.* **37**, 1287–1298.
- Colwell, R., MacDonell, M., & Swartz, D. (1989). Identification of an Antarctic endolithic microorganism by 5S rRNA sequence analysis. *Syst. Appl. Microbiol.* **11**, 182–186.
- Delille, D. (1987). Spatial distribution of coastal Antarctic seawater bacteria. *Polar Biol.* **11**, 41–45.
- Delille, D. (1990). Factors affecting the horizontal patchiness of coastal antarctic seawater bacteria. *Polar Biol.* **8**, 55–60.
- Delille, D. (1992). Marine bacterioplankton at the Weddell Sea ice edge, distribution of psychrophilic and psychrotrophic populations. *Polar Biol.* **12**, 205–210.
- Demeke, T. & Adams, R. (1992). The effects of plant polysaccharides and buffer additives on PCR. *Biotechniques* **12**, 333–334.

- Do, N. & Adams, R. (1991). A simple technique for removing plant polysaccharide contaminants from DNA. *Biotechniques* **10**, 162–166.
- Dressler, B.O. & Reimold, W.U. (2001). Terrestrial impact melt rocks and glasses. *Earth Sci. Rev.* **56**, 205–284.
- Egli, T. (2000). Nutrition of microorganisms. In *Encyclopedia of Microbiology*, ed. Lederberg, J., Vol. 3, pp. 431–447. San Diego: Academic Press.
- Franzmann, P. (1996). Examination of Antarctic prokaryotic diversity through molecular comparisons. *Biodiversity Conservation* **5**, 1295–1305.
- Franzmann, P., Deprez, P., McGuire, A., McMeekin, T. & Burton, H. (1990). The heterotrophic, bacterial microbiota of Burton Lake, Antarctica. *Polar Biol.* **10**, 261–264.
- Friedmann, E.I. (1982). Endolithic microorganisms in the Antarctic Cold Desert. *Science* **215**, 1045–1053.
- Friedmann, E.I. & Ocampo, R. (1976). Endolithic blue-green algae in the dry valleys: primary producers in the Antarctic Desert ecosystem. *Science* **193**, 1247–1249.
- Frisch, T. & Thorsteinsson, R. (1978). Houghton astrobleme: a Mid-Cenozoic impact crater Devon Island, Canadian Arctic archipelago. *Arctic* **31**, 108–124.
- Gold, W.G. (1988). The influence of cryptogamic crusts on the thermal environment and temperature relations of plants in a high Arctic polar desert, Devon Island, N.W.T., Canada. *Arctic and Alpine Research* **30**, 108–120.
- Gosink, J. & Staley, J. (1995). Biodiversity of gas vacuolate bacteria from Antarctic Sea ice and water. *Appl. Environ. Microbiol.* **61**, 3486–3489.
- Grieve, R.A. (1988). The Houghton impact structure – summary and synthesis of the results of the HISS Project. *Meteoritics Planet. Sci.* **23**, 249–254.
- Hirsch, P., Hoffman, B., Gallikowski, C.C., Mevs, U., Siebert, J. & Sittig, M. (1988). Diversity and identification of heterotrophs from Antarctic rocks of the McMurdo Dry Valleys (Ross Desert). *Polarforschung* **58**, 261–269.
- Jessberger, E.K. (1988).  $^{40}\text{Ar}$ – $^{39}\text{Ar}$  dating of the Houghton impact structure. *Meteorit. Planet. Sci.* **23**, 233–234.
- Junge, K., Gosink, J., Hoppe, H. & Staley, J. (1998). *Arthrobacter*, *Brachybacterium*, and *Planococcus* isolates identified from Antarctic sea ice brine. *System. Appl. Microbiol.* **21**, 306–314. (including: 16S rRNA Sequence, GAPPED-BLAST Database, #AF041789).
- Keddie, R., Collins, M. & Jones, D. (1984). Genus *Arthrobacter*. In *Bergey's Manual of Systematic Bacteriology*, ed. Sneath, P., Vol. 2, pp. 1288–1301. Baltimore: Williams & Wilkins.
- Kellogg, D. & Kellogg, T. (1996). Diatoms in Antarctic ice: implications for diatom contamination of glacial deposits. *Geology* **24**, 115–118.
- Kocur, M. (1984). Genus *Planococcus*. In *Bergey's Manual of Systematic Bacteriology*, ed. Sneath, P., Vol. 2, pp. 1011–1013. Baltimore: Williams & Wilkins.
- Lee, P. *et al.* (1999). Search for meteorites at Martin Hills and Pirrit Hills, Antarctica. *Lunar Planetary Science Conf. XXX*, March 1999. Houston, TX: Lunar and Planetary Institute.
- Lukow, T. (1998). 16S rRNA Sequence, GAPPED-BLAST Database, #AS0252579.
- Melosh, J. (1989). *Impact Cratering: a Geologic Process*. Clarendon Press: Oxford.
- Metzler, A., Ostertag, R., Redeker, H.-J. & Stoeffler, D. (1988). Composition of the crystalline basement and shock metamorphism of crystalline and sedimentary target rocks at the Houghton Impact Crater, Devon Island, Canada. *Meteorit. Planet. Sci.* **23**, 197–207.
- Mitrofanov, I., Anfimov, D., Kozyrev, A., Litvak, M., Sanin, A., Tret'yakov, V., Krylov, A., Shvetsov, V., Boynton, W., Shinohara, C., Hamara, D. & Saunders, R.S. (2002). Maps of subsurface hydrogen from the high energy neutron detector, Mars Odyssey. *Science* **297**(5578), 78–81.
- Morikawa, N., Kanaya, S. & Kato, T. (2002). 16S rRNA Sequence, GAPPED-BLAST Database, #AB039736.
- Nichols, D., Nichols, P. & McMeekin, T. (1995). Ecology and physiology of psychrophilic bacteria from Antarctic saline lakes and sea-ice. *Sci. Progr.* **78**, 311–347.
- Nienow, J. & Friedmann, E.I. (1993). Terrestrial lithophytic (rock) communities. In *Antarctic Microbiology*, Ch. 9. Wiley-Liss., New York.
- Osinski, G. & Spray, J. (2001). Impact-generated carbonate melts: evidence from the Houghton Structure, Canada. *Earth Planet. Sci. Lett.* **194**, 17–29.
- Palleroni, N. (1984). Genus *Pseudomonas*. In *Bergey's Manual of Systematic Bacteriology*, ed. Krieg, N., Vol. 1, pp. 141–199. Baltimore: Williams & Wilkins.
- Palmer, C. 2000. Polymerase chain reaction (PCR). In *Encyclopedia of Microbiology*, ed. Lederberg, J., Vol. 3, pp. 787–791. San Diego: Academic press.
- Roser, D., Seppelt, R. & Ashbolt, N. (1993). Microbiology of ornithogenic soils from the Windmill Islands, Budd Coast, continental Antarctica: microbial biomass distribution. *Soil Biol. Biochem.* **25**, 165–175.
- Rotert, K., Toste, A. & Steiert, J. (1993). Membrane fatty acid analysis of Antarctic bacteria. *FEMS Microbiol. Lett.* **114**, 253–258.
- Shi, T., Reeves, R.H., Gilichinsky, D.A. & Friedmann, E.I. (1997). Characterization of viable bacteria from Siberian permafrost by 16S rDNA sequencing. *Microbial Ecol.* **33**, 169–179.
- Siebert, J. & Hirsch, P. (1988). Characterization of 15 selected coccal bacteria isolated from Antarctic rock and soil samples from the McMurdo-Dry Valleys (South Victoria Land). *Polar Biol.* **9**, 37–42.
- Sneath, P. (1984). Genus *Janthinobacterium*. In *Bergey's Manual of Systematic Bacteriology*, ed. Krieg, N., Vol. 1, pp. 376–377. Baltimore: Williams & Wilkins.
- Steele, A., Goddard, D., Stapleton, D., Toporski, J., Peters, V., Bassinger, V., Sharples, G., Wynn-Williams, D. & McKay, D. (2000). Investigations into an unknown organism on the Martian Meteorite Allan Hills 84001. *Meteoritics Planet. Sci.* **35**, 237–241.
- Suzuki, T. & Yamasato, K. (1994). Phylogeny of spore-forming lactic acid bacteria based on 16S rRNA. *FEMS Microbiol. Lett.* **115**, 13–17.
- Totland, M., Jarvis, I. & Jarvis, K. E. (1992). An assessment of dissolution techniques for the analysis of geological samples by plasma spectroscopy. *Chem. Geol.* **95**, 35–62.
- Vincent, W.F. (1988). Lithic ecosystems: the rock environments. In Ch. 9, *Microbial Ecosystems in Antarctica*. Cambridge University Press: Cambridge.
- Vincent, W.F. (2000). Evolutionary origins of Antarctic microbiota: invasion, selection, and endemism. *Antarctic Sci.* **12**, 374–385.
- Wierzbos, J. & Ascaso, C. (2002). Microfossil record of rocks from the Ross Desert, Antarctica: implications in the search for past life on Mars. *Int. J. Astrobiol.* **1**, 51–60.
- Wynn-Williams, D.D. (2000). Cyanobacteria in deserts – life at the limit? In *The Ecology of Cyanobacteria*, eds Whitton, B.A. & Potts, M., pp. 341–343. Kluwer, Dordrecht.