

Hidden Charge $2e$ Boson in Doped Mott Insulators

Robert G. Leigh, Philip Phillips, and Ting-Pong Choy

Department of Physics, University of Illinois, 1110 West Green Street, Urbana, Illinois 61801 USA

(Received 5 December 2006; published 25 July 2007)

We construct the low-energy theory of a doped Mott insulator, such as the high-temperature superconductors, by explicitly integrating over the degrees of freedom far away from the chemical potential. For either hole or electron doping, a charge $2e$ bosonic field emerges at low energy. The charge $2e$ boson mediates dynamical spectral weight transfer across the Mott gap and creates a new charge e excitation by binding a hole. The result is a bifurcation of the electron dispersion below the chemical potential as observed recently in angle-resolved photoemission on Pb-doped $\text{Bi}_2\text{Sr}_2\text{CaCu}_2\text{O}_{8+\delta}$ (Pb2212).

DOI: [10.1103/PhysRevLett.99.046404](https://doi.org/10.1103/PhysRevLett.99.046404)

PACS numbers: 71.27.+a

Two problems beset the construction of a proper low-energy theory (explicit integration of the high-energy scale) of doped Mott insulators. First, the high-energy degrees of freedom are neither fermionic nor bosonic. To illustrate, in a Mott insulator, the chemical potential lies in a charge gap between two bands that represent electron motion on empty (lower Hubbard band, LHB for short) and singly occupied sites (upper Hubbard band, hereafter UHB). Since the latter involves double occupancy, the gap between the bands is set by the on-site repulsion energy U . Nonetheless, both double occupancy and double holes represent high-energy excitations in the half-filled insulating state as each is equally far from the chemical potential. As neither of these is fermionic, standard fermionic path integral procedures are of no use.

Second, unlike the static bands in band insulators, the UHB and LHB are not rigid, thereby permitting spectral weight transfer. When x holes are placed in a Mott insulator, at least $2x$ [1] single particle addition states are created just above the chemical potential. The deviation from x , as would be the case in a band insulator, is intrinsic to the strong correlations that mediate the Mott insulating state in a half-filled band, thereby distinguishing Mottness from ordering. The states in excess of x arise from two distinct effects. Each hole reduces the number of ways of creating a doubly occupied site by one, thereby reducing the spectral weight at high energy. As the x empty sites can be occupied by either spin up or spin down electrons, the $2x$ sum rule is exact [1] in the atomic limit. Further, in the presence of hybridization (with matrix element t), virtual excitations between the LHB and UHB increase the loss of spectral weight at high energy thereby leading to a faster than $2x$ growth [1–3] of the low-energy spectral weight, a phenomenon confirmed [4–6] widely in the high-temperature copper-oxide superconductors.

Because some of the low-energy degrees of freedom of doped Mott insulators derive from the high-energy scale, low-energy descriptions must either (C1) abandon Fermi statistics or (C2) generate new degrees of freedom [1] which ultimately leads to electron number nonconservation. Current proposals for the low-energy physics of

doped Mott insulators are based either on perturbation theory [7] followed by projecting out the high-energy sector or slaved [8] operators designed to exclude double occupancy. As projection is not integration, neither permits an explicit integration of the high-energy scale, and both miss relevant physical aspects.

We show that exact integration of the high-energy scale results in a low-energy theory that possesses a charge $2e$ bosonic mode. Such an excitation might have been anticipated in light of the mixing between high- and low-energy multiply charged states [9]. Our theory is an explicit example of proposal (C2) as the conserved charge involves both the boson and electron number. Note that the *emergence* of new degrees of freedom in a low-energy theory, not directly built out of elementary excitations, is not without precedent. Indeed, we believe that there are useful lessons to draw for Mott insulators from analogies with confining theories or other strongly coupled theories. A simple theoretical model which bears some resemblance to the theory that we develop below is the nonlinear σ model (NL σ M), in which an initially nondynamical field develops correlations and in fact determines the phase structure of the theory.

While our starting point is the one-band Hubbard model,

$$H_{\text{Hubb}} = -t \sum_{i,j,\sigma} g_{ij} c_{i,\sigma}^\dagger c_{j,\sigma} + U \sum_{i,\sigma} c_{i,\uparrow}^\dagger c_{i,\downarrow}^\dagger c_{i,\downarrow} c_{i,\uparrow} \quad (1)$$

our scheme is completely general and is applicable to the n -band case as well. Here i, j label lattice sites, g_{ij} is equal to one iff i, j are nearest neighbors and $c_{i\sigma}$ annihilates an electron with spin σ on lattice site i . The Hilbert space of this model is a product of Fock spaces, $\otimes_i (\mathcal{F}_\uparrow \otimes \mathcal{F}_\downarrow)$. We are concerned in the limit when the Hubbard bands are well separated, $U \gg t$. Given that the chemical potential lies in the gap between such well-separated bands at half-filling, the high-energy degree of freedom is ambiguous at half-filling. Both double occupancy (UHB) and double holes (LHB) are equally costly. Doping removes this ambiguity. Hole-doping jumps the chemical potential to the top of the LHB thereby defining double occupancy to be the high-energy scale. For electron doping, the chemical

potential lies at the bottom of the upper Hubbard band and it is the physics associated with double holes in the lower Hubbard band that should be coarse grained. A generalized particle-hole transformation relates the two theories.

Hole doping.—The basic idea of our construction is to rewrite the Hubbard model in such a way as to isolate the high-energy degrees of freedom so that they can be simply integrated out. We do this by first introducing a new oscillator that represents the degrees of freedom at high energy and including a constraint which ensures that the extended theory is *equivalent* to the Hubbard model. If we simply solve this constraint we return to the description (1) of the Hubbard model, while if instead, we integrate out the high-energy degrees of freedom, we will obtain the low-energy effective theory. To this end, we extend the Hilbert space $\otimes_i(\mathcal{F}_\uparrow \otimes \mathcal{F}_\downarrow \otimes \mathcal{F}_D)$. We associate D^\dagger with the creation of double occupation. In order to limit the Hilbert space to single occupation in the D sector, we will take D to be fermionic. Integrating over the high-energy scale will be accomplished by integrating over D . In particular, we will formulate a Hamiltonian for the extended theory in such a way that if we were to solve the constraint, precisely the Hubbard Hamiltonian (1) would be recovered. The action of the standard electron creation operator $c_{i\sigma}^\dagger$ and the new fermionic operator D^\dagger to create the allowed states on a single site are shown in Fig. 1. We now formulate a Euclidean Lagrangian

$$L = \int d^2\eta \left[\bar{\eta}\eta \sum_{i\sigma} (1 - n_{i\bar{\sigma}}) c_{i\sigma}^\dagger \dot{c}_{i\sigma} + \sum_i D_i^\dagger \dot{D}_i + U \sum_j D_j^\dagger D_j - t \sum_{i,j,\sigma} g_{ij} [C_{ij\sigma} c_{i,\sigma}^\dagger c_{j,\sigma} + D_i^\dagger c_{j,\sigma}^\dagger c_{i,\sigma} D_j + (D_j^\dagger \eta c_{i,\sigma} V_\sigma c_{j,\bar{\sigma}} + \text{H.c.})] + H_{\text{con}} \right], \quad (2)$$

in the extended Hilbert space in such a way as to include the hopping terms (see right panel in Fig. 1) present in the Hubbard model (where we have replaced doubly occupied sites by D occupation). Here, η is a formal complex Grassmann constant which we have inserted in order to




FIG. 1 (color online). Extended Hilbert space (left) which allows an explicit integration of the high-energy scale. Hopping processes (right) included in the Lagrangian.

keep track of statistics, and $d^2\eta$ denotes Grassmann integration. The parameter V_σ has values $V_\uparrow = 1$, $V_\downarrow = -1$, and simply ensures that D couples to the spin singlet. The operator $C_{ij\sigma}$ is of the form $C_{ij\sigma} \equiv \bar{\eta}\eta\alpha_{ij\sigma} \equiv \bar{\eta}\eta(1 - n_{i,\bar{\sigma}})(1 - n_{j,\bar{\sigma}})$ with number operators $n_{i,\sigma} \equiv c_{i,\sigma}^\dagger c_{i,\sigma}$. Note that the dynamical terms that appear in the Lagrangian are nontraditional because the dynamics with the $c_{i\sigma}$ operators must exclude those sites which contain the occupancy $c_{i\uparrow}^\dagger c_{i\downarrow}^\dagger |0\rangle$. Finally, the constraint H_{con} is taken to be

$$H_{\text{con}} = s\bar{\eta} \sum_j \varphi_j^\dagger (D_j - \eta c_{j,\uparrow} c_{j,\downarrow}) + \text{H.c.} \quad (3)$$

where φ_j is a charge $2e$ bosonic field. The constant s will be determined shortly. To see how this constraint removes unphysical states that arise from the extended Hilbert space, we compute the partition function

$$Z = \int [Dc Dc^\dagger D D D^\dagger D \varphi D \varphi^\dagger] \exp^{-\int_0^T L dt}. \quad (4)$$

The integration over φ_i yields a series of δ functions which makes the integral over D trivial. The resultant Lagrangian given by $\int d^2\eta \bar{\eta}\eta L_{\text{Hubb}} = \sum_{i\sigma} c_{i\sigma}^\dagger \dot{c}_{i\sigma} + H_{\text{Hubb}}$ is identical to that of the Hubbard model. This constitutes the ultraviolet (UV) limit of our theory. As is evident, in this limit the extended Hilbert space contracts, unphysical states such as $|1, 0, 1\rangle$, $|0, 1, 1\rangle$, $|1, 1, 1\rangle$ are set to zero, and we identify $|1, 1, 0\rangle$ with $|0, 0, 1\rangle$. Note there is no contradiction between treating D as fermionic and the constraint in Eq. (3). The constraint never governs the commutation relation for D . The value of D is determined by Eq. (3) only when φ is integrated over. This is followed immediately by an integration over D at which point D is eliminated from the theory.

The theory given above permits us to coarse grain the system cleanly for $U \gg t$. The energy scale associated with D is the large on-site energy U . Hence, it makes sense, instead of solving the constraint, to integrate out D . This will result in the low-energy (IR) theory. Such an integration may be done exactly as the theory is Gaussian in D . This is not possible in previous theories. Because of the dynamical term in the action, integration over D will yield a theory that is frequency dependent. We identify the corresponding low-energy theory by setting the frequency to zero. Since the theory is Gaussian, it suffices to complete the square in the D field. To accomplish this, we define the matrix

$$\mathcal{M}_{ij} = \left(\delta_{ij} - \frac{t}{(\omega + U)} g_{ij} \sum_\sigma c_{j,\sigma}^\dagger c_{i,\sigma} \right) \quad (5)$$

and $b_i = \sum_j b_{ij} = \sum_{j\sigma} g_{ij} c_{j,\sigma} V_\sigma c_{i,\bar{\sigma}}$. At zero frequency the Hamiltonian [10] is

$$H_h^{\text{IR}} = -t \sum_{i,j,\sigma} g_{ij} \alpha_{ij\sigma} c_{i,\sigma}^\dagger c_{j,\sigma} + H_{\text{int}} - \frac{1}{\beta} \text{Tr} \ln \mathcal{M}$$

where

$$H_{\text{int}} = -\frac{t^2}{U} \sum_{j,k} b_j^\dagger (\mathcal{M}^{-1})_{jk} b_k - \frac{s^2}{U} \sum_{i,j} \varphi_i^\dagger (\mathcal{M}^{-1})_{ij} \varphi_j - s \sum_j \varphi_j^\dagger c_{j,\uparrow} c_{j,\downarrow} + \frac{st}{U} \sum_{i,j} \varphi_i^\dagger (\mathcal{M}^{-1})_{ij} b_j + \text{H.c.}, \quad (6)$$

which constitutes the true (IR) limit as the high-energy scale has been removed. The energy scale s is set by noting that the fourth term entering our Hamiltonian can mediate spin exchange. As the energy scale for this process is t^2/U , we make the identification $s \simeq t$. Hence, appearing at low energy is a charge $2e$ bosonic field which can either annihilate or create doubly occupied sites or nearest-neighbor singlets. That the energy cost for double occupancy in the IR is t^2/U and not U underscores the fact that the UHB and LHB are not orthogonal. If they were, integrating out the high-energy scale would not result in new charge $2e$ degrees of freedom at low energy. While electron number conservation is broken in the IR, a conserved low-energy charge does exist, however [11]: $Q = \sum_{i\sigma} c_{i\sigma}^\dagger c_{i\sigma} + 2 \sum_i \varphi_i^\dagger \varphi_i$. As Eq. (6) lays plain, the bosons acquire dynamics only through electron motion. Further, they lack a Fock space of their own since all operators in the extended space have been integrated out. Indeed, on phenomenological grounds, weakly interacting Bose-Fermi models have been advanced [12,13]. What the current analysis clarifies is that there is a rigorous connection between a strongly coupled Bose-Fermi model and the low-energy

physics of doped Mott insulator. As we will see, φ_i 's role is to provide internal structure to the electron by mediating composite excitations.

In the limit $U \rightarrow \infty$, the theory reduces to the restricted hopping term and the third term in Eq. (6). Performing the φ integration in the partition function, we arrive at the constraint $\delta(c_{i\uparrow} c_{i\downarrow})$. This leads to a vanishing of double occupancy, the correct result for $U = \infty$. Second, for $\varphi = 0$, we have the restricted hopping term and second term in Eq. (6). Approximating \mathcal{M}_{ij} by its leading term δ_{ij} , we reduce the second term to $\sum_i b_i^\dagger b_i = \sum_{ij\ell\sigma\sigma'} g_{ij} g_{j\ell} c_{i,\sigma}^\dagger V_{\sigma} c_{j,\bar{\sigma}}^\dagger c_{\ell,\sigma'} V_{\sigma'} c_{j,\bar{\sigma}'}$, which contains the spin-spin interaction $-(S_i \cdot S_j - n_i n_j/4)$ as well as the three-site hopping term. Hence, the $\varphi = 0$ limit contains the t - J model, thereby establishing that the physics contained in φ_i is nonprojective.

Electron doping.—For electron doping, the chemical potential jumps to the bottom of the UHB and hence the degrees of freedom that lie far away from the chemical potential no longer correspond to double occupancy, but rather double holes. We proceed as before by extending the Hilbert space and constructing a new Lagrangian

$$L = \int d^2\eta \left[\bar{\eta} \eta \sum_{i\sigma} n_{i\bar{\sigma}} c_{i\sigma}^\dagger \dot{c}_{i\sigma} + \sum_i \tilde{D}_i^\dagger \dot{\tilde{D}}_i - U \sum_j \tilde{D}_j \tilde{D}_j^\dagger - t \sum_{i,j,\sigma} g_{ij} [c_{i,\sigma}^\dagger n_{i\bar{\sigma}} c_{j,\sigma} n_{j\bar{\sigma}} + \tilde{D}_j^\dagger c_{j,\sigma}^\dagger c_{i,\sigma} \tilde{D}_i - (\eta c_{i,\sigma} V_{\sigma} c_{j,\bar{\sigma}} \tilde{D}_i + \text{H.c.})] + \tilde{H}_{\text{con}} \right], \quad (7)$$

that preserves the distinct hops in the Hubbard model where the operator \tilde{D}_i is a fermion associated with double holes. In this case, the constraint is given by $\tilde{H}_{\text{con}} = s \bar{\eta} \sum_i \varphi_i (\tilde{D}_i - \eta c_{i\uparrow}^\dagger c_{i\downarrow}^\dagger) + \text{H.c.}$ Two differences to note are that (i) because the chemical potential resides in the UHB, the electron hopping term now involves sites that are at least singly occupied and (ii) the order of the \tilde{D}_i and c_i operators is important. If we integrate over φ_i and then \tilde{D}_i , all the unphysical states are removed and we obtain as before precisely L_{Hubb} . Hence, both theories yield the Hubbard model in their UV limits. They differ, however, in the IR as can be seen by performing the integration over \tilde{D}_i . The corresponding integral is again Gaussian and yields

$$H_e^{\text{IR}} = -t \sum_{i,j,\sigma} g_{ij} c_{i,\sigma}^\dagger n_{i\bar{\sigma}} c_{j,\sigma} n_{j\bar{\sigma}} + \tilde{H}_{\text{int}} - \frac{1}{\beta} \text{Tr} \ln \mathcal{M}$$

where

$$\begin{aligned} \tilde{H}_{\text{int}} = & -\frac{t^2}{U} \sum_{j,k} b_j^\dagger (\mathcal{M}^{-1})_{jk} b_k - \frac{s^2}{U} \sum_{i,j} \varphi_i (\mathcal{M}^{-1})_{ij} \varphi_j^\dagger \\ & - s \sum_j \varphi_j c_{j,\uparrow} c_{j,\downarrow} - \frac{st}{U} \sum_{i,j} \varphi_i (\mathcal{M}^{-1})_{ij} b_j + \text{H.c.}, \end{aligned} \quad (8)$$

as the IR limit of the electron-doped theory. In addition to the hopping term, the last term also differs from the hole-doped theory as it enters with the opposite sign. The generalized particle-hole transformation (GPHT) that leads to the hole-doped theory is $c_{i\sigma} \rightarrow e^{i\mathbf{Q} \cdot \mathbf{r}_i} c_{i\sigma}^\dagger$ [$\mathbf{Q} = (\pi, \pi)$] augmented with $\varphi_i \rightarrow -\varphi_i^\dagger$. As φ_i is a complex field, the GPHT interchanges the creation operators of opposite charge. We again make the identification $s \simeq t$ because the last term can also mediate spin exchange. When the boson vanishes, we do recover the exact particle-hole symmetric analogue of the hole-doped theory. Because the field φ now couples to double holes, the relevant creation operator has charge $-2e$ and the conserved charge [11] is $\tilde{Q} = \sum_{i\sigma} c_{i\sigma}^\dagger c_{i\sigma} - 2 \sum_i \varphi_i^\dagger \varphi_i$. This sign change in the conserved charge will manifest itself as a sign change in the chemical potential as long as $\langle \varphi_i^\dagger \varphi_i \rangle \neq 0$. Likewise, the correct $U \rightarrow \infty$ limit is obtained as before.

To uncloak how the spectrum changes upon single-electron addition or subtraction, we derive an *exact* expression for the electron operator in the new low-energy theory. To this end, we translate the Lagrangian for the hole-doped theory by a source term, that generates the canonical electron operator when the constraint is solved.




FIG. 2 (color online). (a) Spectral function along the nodal direction for filling $n = 0.9$ and $U = 10t$ at $T = 0$. The lower energy branch below the chemical potential arises from a binding [which opens a pseudogap in the density of states (c)] of a hole with the charge $2e$ boson. (b) Two kinks occur, one at $0.15t \approx 70$ meV and a higher-energy kink at $0.5t \approx 250$ meV at which the bifurcation appears.

The appropriate source term that yields the canonical electron operator in the UV region is $\sum_{i\sigma} J_{i,\sigma} [\bar{\eta}\eta(1 - n_{i,\bar{\sigma}})c_{i,\sigma}^\dagger + V_\sigma D_i^\dagger c_{i,\bar{\sigma}}\eta] + \text{H.c.}$ However, in the IR region in which we only integrate over the heavy degree of freedom D_i , the electron creation operator

$$(1 - n_{i,\bar{\sigma}})c_{i,\sigma}^\dagger + V_\sigma \frac{t}{U} b_i^\dagger \mathcal{M}_{ij}^{-1} c_{j,\bar{\sigma}} - V_\sigma \frac{s}{U} \varphi_i^\dagger \mathcal{M}_{ij}^{-1} c_{j,\bar{\sigma}} \quad (9)$$

contains the standard term for motion in the LHB, $(1 - n_{i,\bar{\sigma}})c_{i,\sigma}^\dagger$ with a renormalization from spin fluctuations (second term) and a new charge e excitation, $c_{i,\bar{\sigma}}\mathcal{M}_{ij}^{-1}\varphi_j^\dagger$, the IR analogue of the UHB excitation $n_{i,\bar{\sigma}}c_{i,\sigma}^\dagger$. Consequently, we predict that an electron at low energies is in a coherent superposition of the standard LHB state (modified with spin fluctuations) and a new charge e state described by $c_{i,\bar{\sigma}}\mathcal{M}_{ij}^{-1}\varphi_j^\dagger$. It is the presence of these two distinct excitations that preserves the $2x$ sum rule [1].

To illustrate this physics, we offer an approximate calculation of the electron spectral function. For the sake of the following discussion, we consider φ_i to be spatially independent and thereby compute the electron Green function by evaluating $\int d\phi^* d\phi \mathcal{F}[\int [Dc_i^* Dc_i] c_i(t) c_j(0)^* \times \exp(-\int L_{\text{IR}} dt)]/Z$, where \mathcal{F} denotes the Fourier transform. Further, we ignore the four-fermion term $b_i^\dagger b_i$, as this term simply renormalizes the standard LHB band as Eq. (9) indicates. The key results summarized in Fig. 2 are the following: (i) Below the chemical potential the electron spectral function consists of two branches. The inner curve corresponds to the standard LHB of the t - J model while the outer curve the new excitation arising from the nonprojective physics in the true low-energy theory. The bifurcation

persists for a wide range of doping ($1.3 > n > 0.7$) and cannot be captured by mean-field or saddle-point approximations to the Green function, which yield only one of the branches. (ii) Bifurcation occurs at the second higher-energy kink (roughly $0.5t$ approximately 250 meV for the cuprates). (iii) The difference between the two branches is largest at the momentum $(0,0)$ and scales as t . (iv) The presence of the two branches opens a pseudogap in the spectrum as shown in the density of states (panel c). The bifurcation, intensity of each branch, and energy of the kink are in excellent agreement with recent experiments [14] on Pb2212. However, the bifurcation has been interpreted as evidence for spin-charge separation. Although our method is approximate, it is sufficient to capture the essence of Eq. (8): two distinct excitations (as seen experimentally) constitute the removal of an electron at low energies in a doped Mott insulator.

Whether or not $c_{i\sigma}\mathcal{M}_{ij}^{-1}\varphi_j^\dagger$ constitutes a true bound state remains a conjecture at this point, though the presence of two states [Fig. 2(a)] in the excitation spectrum and Eq. (9) are consistent with such physics. An analysis based on Bethe-Salpeter equations is necessary. Consequently, the physics of doped Mott insulators turns on precisely the same kinds of problems that arise in other instances of strong coupling such as nuclear structure and confinement.

We thank Tudor Stanescu for comments, and the NSF No. DMR-0605769 for partial support.

-
- [1] M.B.J. Meinders, H. Eskes, and G.A. Sawatzky, Phys. Rev. B **48**, 3916 (1993).
 - [2] A.B. Harris and R.V. Lange, Phys. Rev. **157**, 295 (1967).
 - [3] H. Eskes *et al.*, Phys. Rev. B **50**, 17980 (1994).
 - [4] S.L. Cooper *et al.*, Phys. Rev. B **41**, 11605 (1990).
 - [5] S. Uchida *et al.*, Phys. Rev. B **43**, 7942 (1991).
 - [6] C.T. Chen *et al.*, Phys. Rev. Lett. **66**, 104 (1991).
 - [7] A.H. MacDonald, S.M. Girvin, and D. Yoshioka, Phys. Rev. B **41**, 2565 (1990); A.L. Chernyshev *et al.*, Phys. Rev. B **70**, 235111 (2004); R.B. Laughlin, arXiv:cond-mat/0209269; E. Altman and A. Auerbach, Phys. Rev. B **65**, 104508 (2002); T.C. Ribeiro and X.-G. Wen, Phys. Rev. Lett. **95**, 057001 (2005).
 - [8] S.E. Barnes, J. Phys. F **6**, 1375 (1976); **7**, 2637 (1977); P. Coleman, Phys. Rev. B **29**, 3035 (1984); G. Kotliar and A.E. Ruckenstein, Phys. Rev. Lett. **57**, 1362 (1986).
 - [9] P. Phillips, D. Galanakis, and T.D. Stanescu, Phys. Rev. Lett. **93**, 267004 (2004).
 - [10] The leading term of $k_B T \ln \mathcal{M}$ scales as $k_B T(t/U)^2$ and hence vanishes at $T = 0$. In all calculations, we replaced \mathcal{M}_{ij}^{-1} with its leading term in t/U , namely δ_{ij} .
 - [11] In this expression for Q , φ^\dagger should be interpreted as a creation operator with the appropriate charge.
 - [12] R. Friedberg and T.D. Lee, Phys. Rev. B **40**, 6745 (1989).
 - [13] J. Ranninger, J.M. Robin, and M. Eschrig, Phys. Rev. Lett. **74**, 4027 (1995).
 - [14] J. Graf *et al.*, Phys. Rev. Lett. **98**, 067004 (2007).