

Higgs singlet as a diphoton resonance in a vector-like quark model

Takaaki Nomura (KIAS)

Based on : Chuan-Hung Chen, T. N. arXiv:1512.06028 and work in progress

1. Introduction

2. A model & diphoton excess



3. Checking constraint

4. VLQ production

5. Summary

1. introduction

Diphoton excess at 750 GeV




ATLAS-CONF-2015-081, CMS-PAS-EXO-15-004

Both ATLAS and CMS observed bump on diphoton invariant mass distribution

3.6 σ : ATLAS

2.6 σ : CMS

(Local significance)



How we can interpret the diphoton excess?

- It could be new particle : spin 0 or 2

❖ **Let us consider scalar particle S with $m_S = 750$ GeV**

- Cross section to produce a new particle S

$$\sigma(pp \rightarrow S)BR(S \rightarrow \gamma\gamma) \approx 3 - 10 \text{ fb}$$

- Width of S?

Best fit value by ATLAS : $\Gamma \sim 45$ GeV

CMS : Narrow width is preferred

- S \rightarrow other modes : not observed

➔ BRs of S are constrained

Properties of the diphoton excess

final state f	σ at $\sqrt{s} = 8 \text{ TeV}$			implied bound on $\Gamma(S \rightarrow f)/\Gamma(S \rightarrow \gamma\gamma)_{\text{obs}}$
	observed	expected	ref.	
$\gamma\gamma$	$< 1.5 \text{ fb}$	$< 1.1 \text{ fb}$	[6, 7]	$< 0.8 (r/5)$
$e^+e^- + \mu^+\mu^-$	$< 1.2 \text{ fb}$	$< 1.2 \text{ fb}$	[8]	$< 0.6 (r/5)$
$\tau^+\tau^-$	$< 12 \text{ fb}$	15 fb	[9]	$< 6 (r/5)$
$Z\gamma$	$< 4.0 \text{ fb}$	$< 3.4 \text{ fb}$	[10]	$< 2 (r/5)$
ZZ	$< 12 \text{ fb}$	$< 20 \text{ fb}$	[11]	$< 6 (r/5)$
Zh	$< 19 \text{ fb}$	$< 28 \text{ fb}$	[12]	$< 10 (r/5)$
hh	$< 39 \text{ fb}$	$< 42 \text{ fb}$	[13]	$< 20 (r/5)$
W^+W^-	$< 40 \text{ fb}$	$< 70 \text{ fb}$	[14, 15]	$< 20 (r/5)$
$t\bar{t}$	$< 550 \text{ fb}$	-	[16]	$< 300 (r/5)$
invisible	$< 0.8 \text{ pb}$	-	[17]	$< 400 (r/5)$
$b\bar{b}$	$\lesssim 1 \text{ pb}$	$\lesssim 1 \text{ pb}$	[18]	$< 500 (r/5)$
jj	$\lesssim 2.5 \text{ pb}$	-	[5]	$< 1300 (r/5)$

$$r = \sigma_{13\text{TeV}} / \sigma_{8\text{TeV}}$$

$$\Gamma / M \approx 0.06$$

From Table 1 of arXiv:1512.04933 (Franceschini et. al.)

➤ $S \rightarrow$ other modes : not observed

➔ BRs of S are constrained

We consider a simple scenario to explain the excess

SM+Vector-like triplet quarks(VLTQ) + scalar singlet(S)
SU(2)

Our strategy

- ❖ S does not mix with SM Higgs
- ❖ $gg \rightarrow S$ and $S \rightarrow \gamma\gamma$ are induced by VLTQ loop
- ❖ VLTQs are heavy as $O(1)$ TeV : S does not decay into them
- ❖ Sizable Yukawa coupling of S and VLTQ enhance the process
- ❖ Two triplet give # of quark = 6 \rightarrow enhance S_{gg} and $S_{\gamma\gamma}$ coupling

1. Introduction

2. Model & diphoton excess

3. Checking constraint

4. VLQ production

5. Summary

Our Model

SM + vector-like triplet quarks (F_1, F_2) + singlet scalar (S)

VLTQ

$$F_1 : (3,3)(2/3), \quad F_2 : (3,3)(-1/3) \quad \{(SU(3), SU(2))(U(1)_Y)\}$$

$$F_1 = \begin{pmatrix} U_1 / \sqrt{2} & X \\ D_1 & -U_1 / \sqrt{2} \end{pmatrix}, \quad F_2 = \begin{pmatrix} D_2 / \sqrt{2} & U_2 \\ Y & -D_2 / \sqrt{2} \end{pmatrix} \quad \left[\begin{array}{l} Q_X = 5/3 \\ Q_Y = -4/3 \end{array} \right]$$

(Y.Okada, L.Panizzi (2013))

Yukawa couplings of VLTQ

$$L_{VLTQ}^{Yukawa} = Y_1 \bar{Q}_L F_{1R} \tilde{H} + Y_2 \bar{Q}_L F_{2R} H + y_1 \text{Tr}(\bar{F}_{1L} F_{1R}) S + y_2 \text{Tr}(\bar{F}_{2L} F_{2R}) S + h.c.$$

Our Model

SM + vector-like triplet quarks (F_1, F_2) + singlet scalar (S)

VLTQ


$$F_1 : (3,3)(2/3), \quad F_2 : (3,3)(-1/3) \quad \{(SU(3), SU(2))(U(1)_Y)\}$$

$$F_1 = \begin{pmatrix} U_1 / \sqrt{2} & X \\ D_1 & -U_1 / \sqrt{2} \end{pmatrix}, \quad F_2 = \begin{pmatrix} D_2 / \sqrt{2} & U_2 \\ Y & -D_2 / \sqrt{2} \end{pmatrix} \quad \left[\begin{array}{l} Q_X = 5/3 \\ Q_Y = -4/3 \end{array} \right]$$

(Y.Okada, L.Panizzi (2013))

Yukawa couplings of VLTQ

$$L_{VLTQ}^{Yukawa} = \underbrace{Y_1 \bar{Q}_L F_{1R} \tilde{H} + Y_2 \bar{Q}_L F_{2R} H}_{\text{blue line}} + \underbrace{y_1 \text{Tr}(\bar{F}_{1L} F_{1R}) S + y_2 \text{Tr}(\bar{F}_{2L} F_{2R}) S}_{\text{red line}} + h.c.$$



It induces decay of F_i & mixing of VLQ and SM quarks



$gg \rightarrow S$ and $S \rightarrow \gamma\gamma$

*We assume $Z_2: S \rightarrow -S, F_{iL} \rightarrow -F_{iL}$ to forbid S-H mixing (softly broken by F mass term)

Gauge interactions of VLTQs

$$L_{VFF} = -g \left[(\bar{X}\gamma^\mu U_1 + \bar{U}_1\gamma^\mu D_1 + \bar{D}_2\gamma^\mu Y + \bar{U}_2\gamma^\mu D_2) W_\mu^+ + h.c. \right] \\ - \left[\frac{g}{c_W} \bar{F}_1\gamma^\mu (T^3 - s_W^2 Q_1) F_1 + e \bar{F}_1\gamma^\mu Q_1 F_1 A_\mu + (F_1 \rightarrow F_2, Q_1 \rightarrow Q_2) \right]$$

$$F_1^T = (X, U_1, D_1); \quad F_2^T = (U_2, D_2, Y)$$

❖ Isospin of VLTQ is different from the SM quarks

➔ Z-mediated FCNC is induced since CKM matrix is not unitary

$$\mathcal{L}_{Wud} = -\frac{g}{\sqrt{2}} \bar{u}_L \gamma^\mu V_{CKM}^L d_L W_\mu^+ - \frac{g}{\sqrt{2}} \bar{u}_R \gamma^\mu V_{CKM}^R d_R W_\mu^+ + h.c.$$

$$\mathcal{L}_{Zqq} = -\frac{g}{c_W} C_{ij}^{qL} \bar{q}_{iL} \gamma^\mu q_{jL} Z_\mu - \frac{g}{c_W} C_{ij}^{qR} \bar{q}_{iR} \gamma^\mu q_{jR} Z_\mu$$

$$C_{ij}^{qL} = (I_3 - s_W^2 Q_q) \delta_{ij} + \frac{1}{2} (-V_{Li4}^q V_{Lj4}^{q*} + V_{Li5}^q V_{Lj5}^{q*})$$

$$C_{ij}^{qR} = -s_W^2 Q_q \delta_{ij} + \epsilon_q (V_R^q)_{i\alpha_q} (V_R^{q*})_{\alpha_q j}$$

$$V_{CKM}^L = V_L^u \begin{pmatrix} (V_{CKM})_{3 \times 3} & \mathbf{0}_{3 \times 2} \\ \text{---} & \text{---} \\ \mathbf{0}_{2 \times 3} & \sqrt{2} \mathbf{1}_{2 \times 2} \end{pmatrix} V_L^{d\dagger} \quad V_{CKM}^R = V_R^u \begin{pmatrix} \mathbf{0}_{3 \times 3} & \mathbf{0}_{3 \times 2} \\ \text{---} & \text{---} \\ \mathbf{0}_{2 \times 3} & \sqrt{2} \mathbf{1}_{2 \times 2} \end{pmatrix} V_R^{d\dagger}$$

Gluon fusion and decay modes of S

Gluon fusion and decay of S via VLTQ loop

$$gg \rightarrow S \rightarrow VV \quad L_{sgg} = \frac{\alpha_s}{8\pi} \left(\sum_{F_i} \frac{3y_i}{2m_{F_i}} A_{1/2}(\tau_{F_i}) \right) \phi G^{a\mu\nu} G_{\mu\nu}^a$$

Decay widths

$$\Gamma(S \rightarrow gg) = \frac{\alpha_s^2 m_S^3}{32\pi^3} \left| \sum_i \frac{y_i}{2m_{F_i}} A_{1/2}(\tau_i) \right|^2$$


$$\left(\tau_i = \frac{4m_{F_i}^2}{m_S^2} \right)$$

$$\Gamma(S \rightarrow \gamma\gamma) = \frac{\alpha^2 m_S^3}{256\pi^3} \left| \sum_i \frac{3y_i Q_{F_i}^2}{m_{F_i}} A_{1/2}(\tau_i) \right|^2$$

$$\Gamma(S \rightarrow Z\gamma) = \frac{9m_S^3}{32\pi^3} |A_F|^2 \left(1 - \frac{m_Z^2}{m_S^2} \right)^3 \quad \Gamma(S \rightarrow WW) = \frac{\alpha^2 m_S^3}{256\pi^3} \left| \sum_i \frac{6y_i}{m_{F_i} s_W^2} A_{1/2}(\tau_i) \right|^2$$

$$\Gamma(S \rightarrow ZZ) = \frac{\alpha^2 m_S^3}{256\pi^3} \left| \sum_i \frac{3y_i (T_{F_i}^3 - s_W^2 Q_{F_i})^2}{m_{F_i} s_W^2 c_W^2} A_{1/2}(\tau_i) \right|^2$$

Total decay width of S and BR



$$BR(S \rightarrow gg) \approx 0.67$$

$$BR(S \rightarrow W^+W^-) \approx 0.15$$

$$BR(S \rightarrow ZZ) \approx 0.095$$

$$BR(S \rightarrow Z\gamma) \approx 0.039$$

$$BR(S \rightarrow \gamma\gamma) \approx 0.017$$



$$\left(\begin{array}{l} BR(S \rightarrow hh) \approx 0.007 \\ BR(S \rightarrow t\bar{t}) \approx 0.018 \end{array} \right)$$

❖ The width is mostly less than 1 GeV : narrow width

❖ BRs are almost independent of VLTQ mass

$$K_{S \rightarrow gg} = 1.35$$

Gluon fusion production cross section at 13 TeV



❖ The VLTQ masses are universal for simplicity

$$K_{gg \rightarrow s} = 1.35$$


❖ Cross section is estimated with CalcHEP

(Djouadi Phys. Rep. 457, 1)

❖ The cross section can be O(1) pb with O(1) Yukawa

2. Model & diphoton excess

$$\sigma(gg \rightarrow S) \times \text{BR}(S \rightarrow \gamma\gamma)$$



~5 fb cross section is possible with O(1) TeV m_F and O(1) Yukawa coupling

1. Introduction

2. Model & diphoton excess

3. Checking constraint

4. VLQ production

5. Summary

3. Checking constraints

Checking constraints: 8TeV data

final state f	σ at $\sqrt{s} = 8 \text{ TeV}$		
	observed	expected	ref.
$\gamma\gamma$	$< 1.5 \text{ fb}$	$< 1.1 \text{ fb}$	[6, 7]
$e^+e^- + \mu^+\mu^-$	$< 1.2 \text{ fb}$	$< 1.2 \text{ fb}$	[8]
$\tau^+\tau^-$	$< 12 \text{ fb}$	15 fb	[9]
$Z\gamma$	$< 4.0 \text{ fb}$	$< 3.4 \text{ fb}$	[10]
ZZ	$< 12 \text{ fb}$	$< 20 \text{ fb}$	[11]
Zh	$< 19 \text{ fb}$	$< 28 \text{ fb}$	[12]
hh	$< 39 \text{ fb}$	$< 42 \text{ fb}$	[13]
W^+W^-	$< 40 \text{ fb}$	$< 70 \text{ fb}$	[14, 15]
$t\bar{t}$	$< 550 \text{ fb}$	-	[16]
invisible	$< 0.8 \text{ pb}$	-	[17]
$b\bar{b}$	$\lesssim 1 \text{ pb}$	$\lesssim 1 \text{ pb}$	[18]
jj	$\lesssim 2.5 \text{ pb}$	-	[5]

$$\sigma(gg \rightarrow S)_{13\text{TeV}} BR(S \rightarrow \gamma\gamma) \approx 6 \text{ fb}$$

$$BR(S \rightarrow \gamma\gamma) \approx 0.017$$

$$\rightarrow \sigma(gg \rightarrow S)_{13\text{TeV}} \approx 350 \text{ fb}$$

$$\left[\sigma(gg \rightarrow S)_{13\text{TeV}} / \sigma(gg \rightarrow S)_{8\text{TeV}} \approx 5 \right]$$

$$\sigma(gg \rightarrow S)_{8\text{TeV}} BR(S \rightarrow gg) \approx 49 \text{ fb}$$

$$\sigma(gg \rightarrow S)_{8\text{TeV}} BR(S \rightarrow W^+W^-) \approx 10 \text{ fb}$$

$$\sigma(gg \rightarrow S)_{8\text{TeV}} BR(S \rightarrow ZZ) \approx 7.0 \text{ fb}$$

$$\sigma(gg \rightarrow S)_{8\text{TeV}} BR(S \rightarrow Z\gamma) \approx 2.8 \text{ fb}$$

$$\sigma(gg \rightarrow S)_{8\text{TeV}} BR(S \rightarrow \gamma\gamma) \approx 1.2 \text{ fb}$$

From Table 1 of arXiv:1512.04933

Our scenario can satisfy all constraints!

$$BR(S \rightarrow gg) \approx 0.67$$

$$BR(S \rightarrow W^+W^-) \approx 0.15$$

$$BR(S \rightarrow ZZ) \approx 0.095$$

$$BR(S \rightarrow Z\gamma) \approx 0.039$$

$$BR(S \rightarrow \gamma\gamma) \approx 0.017$$

Checking constraints: FCNC

❖ $t \rightarrow ch$

$$\mathcal{L}_{hQq} = \frac{Y_{1i}}{\sqrt{2}}(v+h) \left(\frac{1}{\sqrt{2}} \bar{u}_{Li} U_{1R} + \bar{d}_{Li} D_{1R} \right) + \frac{Y_{2i}}{\sqrt{2}}(v+h) \left(\bar{u}_{Li} U_{2R} - \frac{1}{\sqrt{2}} \bar{d}_{Li} D_{2R} \right)$$

- SM Higgs mediated FCNC induces $t \rightarrow (\mathbf{u}, \mathbf{c})h$
- Y_{11}, Y_{21} contribute to $D-\bar{D}$, $K-\bar{K}$, $B_d-\bar{B}_d$ mixing \rightarrow we assume $Y_{11}=Y_{21}=0$
- $Y_{12}, Y_{13}, Y_{22}, Y_{23}$: constrained by $B_s-\bar{B}_s$ mixing
- When we assume $Y_{12} \sim Y_{13} \sim Y_{22} \sim Y_{23}$

$$\mathcal{L} = -C_{sb} \bar{s} P_R b h - \frac{m_t}{m_b} C_{sb} \bar{c} P_R t h + H.c. \quad \left(C_{sb} = \frac{m_b}{4v} (2\zeta_{12}\zeta_{13} + \zeta_{22}\zeta_{23}) \quad \zeta_{ij} = \frac{v Y_{ij}}{m_F} \right)$$

- Assuming $\Delta m_{B_s} = 1.1688 \times 10^{-11} \text{ GeV}$, we get $|C_{sb}| < 5.2 \times 10^{-4} \rightarrow \zeta^2 < 0.036$

$$\Gamma(t \rightarrow ch) = \frac{m_t}{32\pi} \left| \frac{m_t}{m_b} C_{bs} \right|^2 \left(1 - \frac{m_h^2}{m_t^2} \right)^2 \quad \Rightarrow \quad BR(t \rightarrow ch) < 1.1 \times 10^{-4}$$

$$\Gamma_t = 1.41 \text{ GeV}$$

3. Checking constraints

Checking constraints: $h \rightarrow \gamma\gamma$

Flavor mixing effect \mathbf{Y}_{1j} and \mathbf{Y}_{2j} contribute $\sigma(gg \rightarrow h) \text{BR}(h \rightarrow \gamma\gamma)$

$$\mu_{\gamma\gamma} = \frac{\sigma(pp \rightarrow h)_{\text{VLTQ}} \text{BR}(h \rightarrow \gamma\gamma)_{\text{VLTQ}}}{\sigma(pp \rightarrow h)_{\text{SM}} \text{BR}(h \rightarrow \gamma\gamma)_{\text{SM}}} \\ \approx \left| 1 + \frac{3}{4} \zeta_{gg} \right|^2 \left| 1 + \frac{N_c A_{1/2}(x_F) \zeta_{\gamma\gamma}}{A_1(x_W) + 4/3 A_{1/2}(x_t)} \right|^2$$

$$\zeta_{gg} = \zeta_{12}^2 + \zeta_{13}^2 + \zeta_{22}^2 + \zeta_{23}^2,$$

We assume $\zeta_{12} = \zeta_{13} = \zeta_{22} = \zeta_{23} = \zeta$

$$\zeta_{\gamma\gamma} = \frac{Q_u^2 + 2Q_d^2}{4} (\zeta_{12}^2 + \zeta_{13}^2) + \frac{2Q_u^2 + Q_d^2}{4} (\zeta_{22}^2 + \zeta_{23}^2)$$

$$\left(\begin{array}{l} (*) \\ m_t, m_F \gg m_h, x_W = 4m_W^2 / m_h^2, x_t = 4m_t^2 / m_h^2 \\ x_F = 4m_F^2 / m_h^2, A_1(x_W) \approx -8.3, A_{1/2}(x_t) \approx 1.38 \end{array} \right)$$

For $\zeta^2 < 0.036 \rightarrow \mu_{\gamma\gamma} < 1.18$

➔ Consistent with LHC data $\left\{ \begin{array}{l} \mu_{\gamma\gamma} < 1.17 \pm 0.27 \text{ (ATLAS)} \\ \mu_{\gamma\gamma} < 1.13 \pm 0.24 \text{ (CMS)} \end{array} \right.$

1. Introduction

2. Model & diphoton excess

3. Checking constraint

4. VLQ production

5. Summary

4. VLQ signature

Production of VLTQs at LHC 13 TeV

(work in progress)

Signature of our model: VLTQ production

VLTQ components: $U_{1,2}, D_{1,2}, X, Y$

➔ X, Y production would be interesting

$$\zeta_{11} = \zeta_{21} = 0.02$$

$$\zeta_{12} = \zeta_{13} = \zeta_{22} = \zeta_{23} \equiv \zeta = 0.2$$

❖ X, Y pair production cross section


$$Q = X_{5/3}, Y_{-4/3}.$$

m_F [GeV]	800	900	1000	1100	1200
$\sigma(pp \rightarrow Q\bar{Q})$ [fb]	88	42	22	11	6

❖ X, Y single production cross section via QCD process

TABLE III: Production cross section for $X_{5/3}W^-$ and $Y_{-4/3}W^+$ with various values of m_F , where $\sqrt{s} = 13$ TeV, $\zeta_{11,21} = 0.02$, and $\zeta = 0.2$ are used.

m_F [GeV]	800	900	1000	1100	1200
$\sigma(pp \rightarrow X_{5/3}W^-)$ [fb]	0.72	0.38	0.21	0.12	0.07
$\sigma(pp \rightarrow Y_{-4/3}W^+)$ [fb]	1.4	0.73	0.40	0.23	0.13




4. VLQ signature



Production of VLTQs at LHC 13 TeV

(work in progress)

❖ X, Y single production through electroweak process



Preliminary




s, c initial states are also important

We find large cross section can be obtained



Production of VLTQs at LHC 13 TeV

(work in progress)

❖ X, Y single production through electroweak process



Preliminary



s, c initial states are also important
We find large cross section can be obtained

Summary and Discussions

- ✧ **Diphoton excess can be explained by singlet scalar + VLTQ**
- ✧ **Production and decay processes from one-loop**
- ✧ **Specific pattern of BRs**
- ✧ **It can be consistent with other experimental constraints**
- ✧ **Width of S is narrow in our model**
- ✧ **The model would be tested by searching for VLTQs**

Thank you !

Loop functions in the partial decay widths

$$A_{1/2}(\tau) = 2\tau[1 + (1 - \tau)(\sin^{-1}(1/\sqrt{\tau}))^2] \quad (\tau \geq 1)$$

$$A_F = \frac{\alpha}{2\pi s_W c_W} \sum \frac{-4y_F Q_F}{m_F} (T_F^3 - s_W^2 Q_F) [I_1(\tau, \lambda) - I_2(\tau, \lambda)]$$

$$\tau_i = \frac{4m_{F_i}^2}{m_S^2}, \quad \lambda_i = \frac{4m_{F_i}^2}{m_Z^2}$$

$$I_1(a, b) = \frac{ab}{2(a-b)} + \frac{a^2 b^2}{2(a-b)^2} [f(a)^2 - f(b)^2] + \frac{a^2 b}{(a-b)^2} [g(a) - g(b)],$$

$$I_2(a, b) = -\frac{ab}{2(a-b)} [f(a)^2 - f(b)^2],$$

$$g(t) = \sqrt{t-1} \sin^{-1}(1/\sqrt{t}).$$