

High and novel genetic diversity of *Francisella tularensis* in Germany and indication of environmental persistence

C. SCHULZE¹†, K. HEUNER²†, K. MYRTENNÄS³, E. KARLSSON³, D. JACOB²,
P. KUTZER¹, K. GROßE⁴, M. FORSMAN³ AND R. GRUNOW²*

¹Landeslabor Berlin-Brandenburg, Frankfurt (Oder), Germany

²Robert Koch Institute, Centre for Biological Threats and Special Pathogens, Division 2 (ZBS 2), Highly Pathogenic Microorganisms, Berlin, Germany

³CBRN Defence and Security, Swedish Defence Research Agency (FOI), Umeå, Sweden

⁴Stadt Brandenburg an der Havel, Veterinär- und Lebensmittelüberwachungsamt, Germany

Received 20 January 2016; Final revision 11 May 2016; Accepted 22 May 2016;
first published online 30 June 2016

SUMMARY

In Germany tularemia is a re-emerging zoonotic disease. Therefore, we investigated wild animals and environmental water samples for the presence and phylogenetic diversity of *Francisella tularensis* in the poorly studied Berlin/Brandenburg region. The phylogenomic analysis of three isolates from wild animals revealed three new subclades within the phylogenetic tree of *F. tularensis* [B.71 from a raccoon dog (*Nyctereutes procyonoides*); B.74 from a red fox (*Vulpes vulpes*), and B.75 from a Eurasian beaver (*Castor fiber albicus*)]. The results from histological, PCR, and genomic investigations on the dead beaver showed that the animal suffered from a systemic infection. Indications were found that the bacteria were released from the beaver carcass into the surrounding environment. We demonstrated unexpectedly high and novel phylogenetic diversity of *F. tularensis* in Germany and the fact that the bacteria persist in the environment for at least one climatic season. These findings support a broader host species diversity than previously known regarding Germany. Our data further support the assumption derived from previous serological studies of an underestimated frequency of occurrence of the pathogen in the environment and in wild animals. *F. tularensis* was isolated from animal species not previously reported as natural hosts in Germany.

Key words: Ecology, *Francisella*, genotyping, Germany, tularemia, wild animals.

INTRODUCTION

The highly infectious pathogen *Francisella tularensis* is the causative agent of tularemia, a zoonotic disease with an unusually wide range of host species [1].

Doses as low as 10 bacteria of *F. tularensis* may cause disease [2]. Outbreaks in humans may be associated with exposure to different arthropod vectors, infected animals, food, water, fomites, or contaminated aerosols. The clinical manifestations depend mainly on the route of infection. The onset of the disease is typically sudden and influenza-like, with symptoms including high fever, lymph node enlargement, chills, fatigue, headache, and nausea [3]. Two main subspecies of *F. tularensis* are clinically relevant, namely *F. tularensis* subsp. *tularensis* which is found

* Author for correspondence: Professor R. Grunow, Centre for Biological Threats and Special Pathogens, Division 2 (ZBS 2), Highly Pathogenic Microorganisms, Robert Koch Institute, Seestr. 10, 13353 Berlin, Germany.
(Email: grunowr@rki.de)

† These authors contributed equally to this work.

only in North America and *F. tularensis* subsp. *holarctica* which occurs in most of Eurasia and North America [1, 3]. Three biovars have been described for *F. tularensis* subsp. *holarctica*: an erythromycin-sensitive biovar I (Ery^{Sen}), an erythromycin-resistant biovar II (Ery^{Res}), and biovar *japonica* which is distinguished by its ability to ferment glycerol [1]. More recently, whole genome sequencing and canonical single-nucleotide polymorphism (canSNP) genotyping methods have revealed multiple *F. tularensis* subsp. *holarctica*-type clades, including four basal clades B.4, B.6, B.12, and B.16 [4–6]. However, at present it remains unclear how the biovar classification system of *F. tularensis* subsp. *holarctica* relates to the phylogeny and current canSNP classification system of the bacterium, except for the B.16 subclade which correlates with biovar *japonica* and the B.12/B.13 clades correlating with erythromycin susceptibility [4, 7, 8].

In recent years, both multiple tularemia outbreaks and single human cases have been reported from almost the entire area of Germany, and tularemia is now considered a re-emerging zoonotic disease. The knowledge about human clinical cases as well as seroprevalence in humans has increased in recent years [9–14]. Indeed, an unexpectedly high rate (7.5%) of wild animals seropositive for *F. tularensis* was detected in the Berlin/Brandenburg region [15] and in other regions of Germany [12–14]. The reason for the relatively high (sero)prevalence seen in humans and wildlife is unknown. Therefore, we hypothesized that analysis of the genomic diversity of strains circulating in wild animals in the Berlin/Brandenburg region might indicate whether the studied region is affected with a single genotype of *F. tularensis* or whether a broader diversity may suggest repeated introductions of the pathogen into this area. Hence, we screened wild animals within the Berlin/Brandenburg region for *F. tularensis* and investigated the genomic diversity of the isolates.

METHODS

Samples

All samples from wild animals were obtained within the framework of the German rabies surveillance programme [16]. In total, we gathered 30 animal samples from two investigations. First, we collected 29 animals from 11 locations in the Berlin/Brandenburg region during 2012–2014 (Table 1). Our investigation was prompted by the finding of one dead Eurasian beaver

(*Castor fiber albicus*) close to the town of Brandenburg in February 2012. The beaver was found by a voluntary guard of the nature reserve and the local veterinary health authority of the town of Brandenburg (Fig. 1). The region is called the ‘Erdlöcher von Wust’, a small man-made lake close to the river Havel (Fig. 1c, red circle). This lake is part of a nature reserve. Second, we also included an uncharacterized *F. tularensis* isolate (isolate A63/63) recovered from the liver of a red fox (*Vulpes vulpes*; one of 305 animals analysed) found in the same region in 2008.

To investigate further the presence of *F. tularensis* bacteria in the immediate surroundings, we collected water samples from the site where the dead beaver was found (Fig. 1c, white circles) at ‘Erdlöcher von Wust’ during 2012 and 2013. In 2013, eight sites, but seven additional sites [in addition to Erdlöcher von Wust (site 1)] the Berlin/Brandenburg area of were sampled to screen for the presence of *F. tularensis* (Fig. 1b; Erdlöcher von Wust/Brandenburg [1], Great Plessower lake [2], Great Wannsee lake [3], Stölpchensee lake [4], boar puddle/Wannsee [5], Teltow sewer [6], Nordufer sewer/Berlin [7], and Spree river [8]). The five dates of sampling were 1 March 2012, 4 September 2012, 3 March 2013, 6 August 2013, and 28 November 2013.

Organ samples (kidney, liver, spleen, lymph node, urethra, intestine, lung) were stored at -80°C . We took 200 mg of each organ sample for homogenization using FastPrep tubes containing 1.4 mm ceramic spheres (MP Biomedicals, Germany) immediately after defrosting, and 1 ml sterile PBS was added to each tube. Homogenization was performed three times for 10 s using Precellys 24 (PEQLab Biotechnologie GmbH, Germany). The homogenates were used for cultivation and DNA extraction.

Histology

Tissue samples taken at necropsy from the beaver included the parotid lymph nodes, lung, heart, spleen, liver, kidneys, urethra, and ileum. These were fixed in 10% neutral-buffered formalin and embedded in paraffin wax. Native tissues from the same organs were additionally preserved by freezing at -20°C . For histopathological examination haematoxylin-eosin (HE) stain was used according to standard protocols.

Culture and erythromycin susceptibility

We cultured 100 μl of each organ homogenate in medium T [17] [1% brain heart infusion broth (Difco

Table 1. Selected wild animals examined for the presence of tularemia by qPCR

No.	Date	Species	Origin	Cause of death	<i>Ft</i> PCR
1	9 Feb. 2012	Eurasian beaver	Wust	Found dead	Positive*
2	21 Aug. 2012	Raccoon dog	Wust	Hunted	Positive*
3	30 Aug. 2012	Wild boar	Wust	Hunted	Positive
4	4 Sept. 2012	Raccoon dog	Kirchmöser	Hunted	Negative
5	4 Sept. 2012	Raccoon dog	Kirchmöser	Hunted	Negative
6	18 Sept. 2012	Nutria	Mötzow	Run over	Negative
7	6 Nov. 2012	Raccoon dog	Wust	Hunted	Negative
8	6 Nov. 2012	Raccoon dog	Wust	Hunted	Negative
9–12	13 Nov. 2012	Muskrats	Wustmittelbruch	Hunted	Negative
13	13 Nov. 2012	Raccoon	Wust	Hunted	Negative
14	13 Nov. 2012	Muskkrat	Wuster luch	Hunted	Negative
15	27 Nov. 2012	Raccoon	Wust	Hunted	Negative
16	27 Nov. 2012	Raccoon	Gollwitz	Hunted	Negative
17	8 Jan. 2013	Red fox	Gollwitz	Hunted	Negative
18	11 Feb. 2013	Wild boar	Gollwitz	Hunted	Negative
19	9 Mar. 2013	Wild boar	Gollwitz	Hunted	Negative
20	26 Mar. 2013	American mink	Brandenburg	Hunted	Negative
21	7 May 2013	Red fox	Gollwitz	Hunted	Negative
22	14 May 2013	Eurasian beaver	Brielow	Run over	Negative
23	14 May 2013	Red fox	Wust	Run over	Negative
24	12 Sept. 2013	Raccoon dog	Mötzower grenze	Hunted	Negative
25	4 Feb. 2014	Wild boar	Gollwitz	Hunted	Negative
26	4 Mar. 2014	Red fox	Gollwitz	Hunted	Negative
27	1 Apr. 2014	Eurasian beaver	'Havelumfluter'	Found dead	Negative
28	17 Apr. 2014	European otter	Zachow/Ketzin	Found dead	Negative
29	3 June 2014	Eurasian beaver	Malge	Found dead	Negative

* *F. tularensis* (*Ft*) subsp. *holarctica* was also isolated.

Laboratories Inc., USA), 1% bacto tryptone (Difco), 1% technical casamino acids (Difco), 0.005 g MgSO₄ 0.01% FeSO₄, 0.12% sodium citrate, 0.02% KCl, 0.04% K₂HPO₄, 0.06% L-cysteine, 1.5% glucose], on CHAB agar plates [CHA (Difco; Bestbion, Germany), 1% brain heart infusion broth, 1% proteose-peptone, 1% D-glucose, 0.5% NaCl, 0.1% L-cystine, 1.5% agar, 9% sheep blood], or on *Neisseria* selective medium Plus (Oxoid, Germany) at 37 °C with 5% CO₂ for up to 3 days. Picked clones from water samples were cultivated on CHAB-PACCV (CHABplus 8 × 10⁴ U/l polymyxin B, 2.5 mg/l amphotericin B, 20 mg/l sulfamethoxazol, 100 mg/l cycloheximide, 4 mg/l vancomycin), and on GVPC agar plates (Heipha Dr. Müller GmbH, Germany): ACES-buffered charcoal yeast extract (BCYE) agar plates supplemented with 80 000 IE polymyxin B, 1 mg/l vancomycin, and 80 mg/l cycloheximide. *Francisella* isolates were grown on CHAB agar plates as described above.

Antimicrobial susceptibility was determined using MIC Evaluator Strips (Oxoid). Isolates were grown on CHAB at 37 °C with 5% CO₂ for 48 h. Four to

five single colonies were inoculated in 0.9% NaCl until a turbidity equivalent to a 0.5 McFarland standard occurred. We then transferred 100 µl of the solution onto CHAB agar plates. The test strips were applied onto the inoculated agar surface, and the minimal inhibition concentration of erythromycin was determined after 48 h of incubation.

Molecular analyses

DNA preparation

DNA was extracted from isolates or organ homogenates using the DNeasy Blood and Tissue kit (Qiagen, Germany). Colonies grown on CHAB or 200 µl of the organ homogenates were directly transferred into lysis buffer containing proteinase K. Further steps were performed according to manufacturer's protocol. The purity and concentration of DNA were then determined by using either a Nanodrop spectrophotometer (PEQ-Lab Biotechnologie GmbH) or a Qubit 2.0 fluorometer using the dsDNA BR Assay kit (Life Technologies, USA).




Fig. 1. Location where the carcass of the tularemic beaver was found. (a) Map of Germany showing the Berlin/Brandenburg region and the town Brandenburg at the Havel (●). (b) Location of sample sites 1–6 in Berlin/Brandenburg. 1, 'Erdlöcher von Wust/Brandenburg'; 2, Great Plessower lake; 3, Great Wannsee lake; 4, Stölpchensee lake; 5, boar puddle/Wannsee; 6, Teltow sewer (sample sites 7 and 8, not shown, see Methods section). (c) 'Erdlöcher von Wust/Brandenburg' – location of the beaver carcass (A, red circle, sample site 1) and the additional water sampling sites (B, C, white circles) of this area are indicated. [(b) and (c), Google Maps/Earth images.]

DNA preparation from water samples. A total of 510 ml water was collected at each site and filtered through a 40 µm filter to remove possible debris. Of this sample 500 µl were plated onto two CHAB-PACCV and two GVPC agar plates, respectively. Half of the plates were incubated at 30 °C and the other half at 37 °C for 4–7 days and inspected daily for putative isolates of *Francisella* sp. Isolates were picked on CHAB_PACCV agar plates and cultivated further for 3–5 days. Next, 500 ml of the sample were centrifuged at 5000 *g* at room temperature for 30 min. The pellet was resuspended in ~5 ml of sample water and DNA was isolated by using the 'soil-DNA isolation kit' from MoBio (Dianova, Germany). For negative control we performed DNA isolation procedure from tap water. We stored 100 µl of total DNA extraction volume at –20 °C until polymerase chain reaction (PCR) analysis.

PCR screening

PCR screening of bacterial colonies from water samples and organ samples for *Francisella*. A loop of

bacteria from picked colonies was suspended in 250 µl water and incubated at 99 °C for 10 min. After centrifugation (10 min, 2500 *g*), 10 µl of the supernatant (exhibiting chromosomal DNA) was investigated by PCR using 16s rDNA *Francisella*-specific primers, F5/F11 [18] to detect putative *Francisella* isolates. PCR was carried out using a Thermocycler TRIO-Thermoblock (Biometra, Germany) and the HotStar *Taq* DNA polymerase (Qiagen). Initial denaturation was performed at 95 °C for 5 min with a final extension at 72 °C for 10 min. The cycling conditions (35 cycles) were 94 °C for 1 min, 60 °C for 45 s, and 72 °C for 2 min. PCR amplicons were purified with the QIAquick PCR purification kit according to the manufacturer's instructions (Qiagen). Sequence determination was performed by applying the dye terminator chemistry (Applied Biosystems, Germany), and chromatograms were analysed with Lasergene Software (DNASTar, USA) followed by sequence comparison by BLAST search (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>).

Quantitative real-time PCR (qPCR) and subtyping

Multiplex real-time PCR (5' nuclease assay) targeting *fopA* and *tul4* were used to analyse all suspected *F. tularensis* isolates for confirmation and DNA isolated from water samples. Primer and probe sequences for the detection of *fopA* were Ft-fopA-F: TTGGGCA AATCTAGCAGGTCA; Ft-fopA-R: ATCTGTAGT CAACACTTGCTTGAACA; and Ft-fopA-TM: FAM-AAGACCACCACCAACATCCCAAGCA-BHQ-1, and for the detection of *tul4*, Ft-tul4-F: AGATTACAATGGCAGGCTCC; Ft-tul4-R: AGC TGTCCTACTT ACCGCTACA; and Ft-tul4-TM: Cy5-TTCTAAGTGCCATGATACAAGCTTCCCAA-BHQ-2. The real-time PCR assays were run in a total volume of 25 µl including 5 µl sample volume or DNA. The reaction mix components were TaqMan Environmental MasterMix 2.0 (6.25 µl), 10 pmol/µl primers (0.75 µl each) and probes (0.25 µl each). Amplification was performed in an ABI7500 (Applied Biosystems, Life Technologies, Germany), each run with an initial denaturation step at 95 °C for 10 min, followed by 35 cycles containing a denaturation step at 95 °C for 15 s, and a combined primer annealing and elongation step at 55 °C for 60 s. An internal amplification control (KoMa) was added [19].

F. tularensis subspecies differentiation was performed by targeting region of difference 1 (RD1) according to protocols described by Broekhuijsen *et al.* [20]. PCR was carried out using the high fidelity triple master mix system (Eppendorf, Germany) with 15–100 ng template DNA.

Variable number of tandem repeat (VNTR) typing

VNTR markers Ft-M3, Ft-M6, Ft-M20, Ft-M21, Ft-M22, and Ft-M24 were analysed as described by Johansson *et al.* [21]. PCR was performed using Eppendorf Mastercyclers (Eppendorf) with 50–100 ng bacterial DNA in the presence of 10 pmol/µl primers (1 µl), 2 mM dNTPs (2.5 µl), 25 mM MgCl (1.5 µl), and 1 U/µl *Taq* DNA polymerase (1 µl) (all by Fermentas, Germany), in a total volume of 25 µl. The cycling conditions were an initial denaturation at 94 °C for 4 min, followed by 35 cycles comprising denaturation at 95 °C for 30 s, annealing at 56 °C or 61 °C for 30 s depending on the primer used, and extension at 72 °C for 30 s with a final extension step at 72 °C for 4 min. Each PCR product was subjected to agarose gel electrophoresis (PEQLab Biotechnologie GmbH).

DNA sequencing

For genome sequencing, indexed sequencing libraries were prepared from 50 ng DNA with the Nextera XT DNA Library Preparation kit (cat. no. FC-131-1024) according to the manufacturer's recommendations (Nextera XT DNA Library Preparation Guide, no. 15031942 rev C, Illumina, USA). A 500 cycle (2 × 250 Paired End) MiSeq Reagent kit v. 2 was used to sequence the libraries on the MiSeq sequencing platform. The genomes were assembled using ABySS v. 1.3.5 [22]. All sequence reads were deposited and are freely downloadable from the Sequence Reads Archive at the National Center for Biotechnology Information (NCBI) under BioProject PRJNA285142 with BioSample accession nos. SAMN03773881: A63/63 (FDC407); SAMN03773882: A271_1 (FDC408); and SAMN03773883: A317 (FDC409).

Phylogenetic analysis

The three draft genomes were aligned with seven representative genomes from previously identified main clades of subspecies *holarctica* using progressive Mauve with standard parameters. The GenBank assembly accession numbers for the seven published genomes were GCA_000168775-2 (FSC200); GCA_000009245-1 (LVS); GCA_000017785-1 (FTNF0012-00); and GCA_000014605-1 (OSU18), and NCBI biosample SAMN00000315 (OR96-0246); SAMN 01047935 (FSC162); and SAMN00092746 (Tul7). The phylogenetic tree was constructed in MEGA6 [23] using the neighbour-joining method, where all positions containing gaps and missing data were eliminated, and the distances were computed using the number-of-differences method [24]. The tree was midpoint rooted. A specific study of the *cas-9* gene was made by comparison with the 10 genomes above, and 25 other genomes from Poland, Finland, and Sweden (genomes to be published elsewhere) within the B.39 subclade.

Genotyping of genomes

The three draft genomes were genotyped using canSNPer which is a fast and lightweight tool written in Python for genotype classification in draft genomes of clonal pathogens based on canSNPs [25]. The markers included in the analysis have been described previously: B.1–B.14 by Vogler *et al.* and Birdsell *et al.* [6, 26]; B.15–B.25 by Svensson *et al.* [5]; B.26–B.32

by Chanturia *et al.* [27]; B.33–B.38 by Gyuranecz *et al.* [28]; and B.39–B.43 by Karlsson *et al.* [29]. Newly identified clades were named according to this canSNP classification system for *F. tularensis* that to date, in addition to the 43 canSNPs used above by canSNPer, also include the subspecies *holarctica* canSNPs B.67–B.68 [30], and B.69–B.70 [31].

RESULTS

Our investigation in the Berlin/Brandenburg region during 2012–2014 revealed that three (10%) out of a total of 29 examined animals – one found dead (a beaver) and two hunted [a raccoon dog (*Nyctereutes procyonoides*) and a wild boar (*Sus scrofa*)] – tested positive by PCR analysis (Table 1). The species distribution of the 29 animals examined for the presence of *F. tularensis* by qPCR and culture was as follows: six raccoon dogs, five muskrats (*Ondatra zibethicus*), four red foxes, four beavers, three raccoons (*Procyon lotor*), four wild boars, one European otter (*Lutra lutra*), one American mink (*Neovison vison*), and one nutria (*Myocastor coypus*) (Table 1). In addition, *F. tularensis* was isolated from two out of the three PCR-positive animals. One isolate (A317) was obtained from the raccoon dog and three (A271_1-3) from different organs of the beaver. To our knowledge, this is the first report of a *F. tularensis* subsp. *holarctica* isolated from a beaver in Germany. We further investigated this case in detail.

The tularemia-positive beaver was an adult male of 17.5 kg weight and 119 cm length found dead in February 2012 (Fig. 1) at the ‘Erdlöcher von Wust’ in a nature reserve close to the river Havel (Fig. 1c, red circle). The carcass was submitted for necropsy to the Berlin-Brandenburg State Laboratory in Frankfurt (Oder). At necropsy, the animal was found to be emaciated and uraemic. The left parotid lymph node contained a granulomatous and necrotizing lesion of ~8 mm in diameter (Fig. 2a). Multiple microscopic (pyo-)granulomas and necroses were additionally present in this lymph node (Fig. 2b, c) and the spleen (Fig. 2d). Randomly distributed microscopic foci of fresh lytic necrosis were distributed throughout the liver. Cause of post-renal uraemia and death was a rupture of the severely dilated urinary bladder due to multifocal suppurative nephritis and subsequent occlusion of the urethra by a cast composed of a mixture of urinary gravel, cellular debris, and exudate. At necropsy, the abdominal cavity was filled

with bloody urine after rupture of the urinary bladder and the urethra was morphologically occluded by a cast. However, a correlation of morphology and function is not always possible in cases of urolithiasis, especially when the urinary bladder is already ruptured at necropsy. Functionally, the occlusion seemed to be only partial at the time of the highest internal pressure shortly before final rupture of the urinary bladder, allowing some urine to leak from the carcass into the surrounding snow. *F. tularensis*-specific qPCR analysis was positive for the left parotid lymph node, kidneys, lung, and spleen. *F. tularensis* was cultured from the parotid lymph node, spleen, and lung (isolates A271_1-3).

A qPCR specific for *F. tularensis* was conducted on water samples collected during 2012–2013 in the immediate surroundings of the site where the dead beaver had been found (Fig. 1c, white circles), and seven more distant sites within the Berlin/Brandenburg region (Fig. 1b). We detected *F. tularensis* DNA at sample site A (Fig. 1c) in March 2012 and at all three sample sites (A–C) in September 2012 (Fig. 1c, Supplementary Table S1). We could also detect *F. tularensis* DNA in snow that we collected around the dead beaver which was contaminated with blood-containing urine (data not shown). In contrast, all samples taken 1 year later (2013) were negative for the presence of *F. tularensis* by qPCR analysis (Supplementary Table S1). Cultures for *F. tularensis* were negative for all water samples.

For further comparative analysis of the *F. tularensis* isolates, we also included an uncharacterized *F. tularensis* isolate (isolate A63/63) recovered from the liver of a red fox (one of 305 animals analysed) from the same region in 2008, hunted in the context of the German rabies surveillance programme [16]. All isolates were assigned to the erythromycin-resistant biovar II type (data not shown) but differed regarding their VNTR profiles (Table 2). In addition, VNTR analysis demonstrated that all three isolates from the beaver (A271_1-3) were identical (data not shown). The genomes of the lymph node isolate A271_1 (beaver), A317 (raccoon dog) and A63/63 (red fox) were sequenced on the Illumina sequencing platform producing draft genome sequences of good quality and sequence depth (Table 3). The canSNPer program [25] using the draft genomes as input, assigned all three isolates to the basal B.12 clade confirming their assignment to biovar II Ery^{Res}. The beaver isolate A271_1 was assigned to the B.33/34 subclade; the A317 isolate from the raccoon dog to the B.12/




Fig. 2. Macroscopic and microscopic lesions in the dead Eurasian beaver. (a) Focally extensive necrosis in the left parotid lymph node (arrows). (b) Left parotid lymph node; granulomatous (arrows) and necrotizing (arrowheads) lymphadenitis; H&E stain. (c) Left parotid lymph node; focal pyogranuloma; H&E stain. (d) Spleen; focal pyogranuloma (arrows) adjacent to an intact lymph follicle (arrowheads); H&E stain.

Table 2. *F. tularensis* *subsp.* *holarctica* isolates from Brandenburg region, Germany, investigated in this study

RKI ID*	FOI ID	Year	Source	VNTR typing (bp/no. of repeats)†				B.12 subclade	
				Ft-M3	Ft-M6	Ft-M20	Ft-M24	canSNPer‡	New§
A63/63	FDC407	2008	Red fox (<i>Vulpes vulpes</i>), liver	369/18	311/4	255/3	480/2	B.39	B.74
A271_1	FDC408	2012	Eurasian beaver (<i>Castor fiber albicus</i>), lymph node	297/10	353/6	255/3	459/1	B.33/34	B.75
A317	FDC409	2012	Raccoon dog (<i>Nyctereutes procyonoides</i>), lymph node	333/14	311/4	243/2	480/2	B.12/13	B.71

RKI, Robert Koch Institute; FOI, Swedish Defence Research Agency; VNTR, variable number tandem repeat.

* All three strains showed resistance to erythromycin.

† All three strains showed an identical PCR fragment size for VNTR markers Ft-M21 (396 bp/2 repeats), and Ft-M22 (254 bp/4 repeats).

‡ canSNPer version 1 [25] (<http://github.com/adrlar/CanSNPer>) (freely available).

§ See Figure 3, identified here.

13 clade; and the A63/63 isolate from the red fox to B.39. A phylogenetic analysis of the three new genomes and seven previously published reference genomes revealed branch lengths (number of SNPs) for the new branches and enabled the placement of the new genomes in their phylogenetic context (Fig. 3).

New subclade names were assigned to the new branches and one representative canSNP per new branch was extracted (Fig. 3). The phylogenetic tree was based on a total of 1 707 622 bp in common between the ten genomes and covered 95% of the total length of the genomes of ~1.89 Mbp. The genomes

Table 3. Statistics on the three new genome sequences*

RKI ID	FOI ID	No. contigs (>200 bp)†	Length (contigs >200 bp)	Identity SCHUS4-2	Identity FSC200	Expected sequence depth‡	Total no. pairs (1 pair = 2 × 250 bp)	N50§	BioProject	BioSample
A63/63	FDC407	97	1 823 965	92·03%	94·23%	398	1 428 416	26 808	PRJNA 285142	SAMN03773881
A271	FDC408	100	1 825 110	91·93%	93·91%	419	1 501 284	26 116	PRJNA285142	SAMN03773882
A317	FDC409	98	1 829 934	92·13%	94·58%	413	1 479 734	27 025	PRJNA285142	SAMN03773883

RKI, Robert Koch Institute; FOI, Swedish Defence Research Agency.

* Sequenced on the Illumina MiSeq sequencing platform at FOI with 251 bp read length and paired-end read type.

† Assembled with ABySS v. 1.3.5 [22].

‡ Calculated by (total no. pairs × read length × 2)/1 800 000.

§ The average length of a set of sequences. The N50 length is defined as the length N for which 50% of all bases in the sequences are in a sequence of length $L < N$.

were highly identical with 1537 base differences (0·09%) in this alignment, thus following a clonal inheriting pattern. In total, we extended the number of canSNPs with five markers that were designated B.71–B.75 according to the canSNP classification system. The branch leading to the beaver isolate A217_1 was designated B.75. The previous B.12/13 clade was divided into two sister clades: the relatively long branch leading to the raccoon dog isolate (A317) was denoted as B.71 and the sister branch was denoted as B.72, leading to all previously known branches within the basal clade B.12, the dominating basal clade in Eastern Europe from Scandinavia to the Black Sea (Fig. 3). The branch leading to the red fox isolate A63/63 was assigned to B.74, while its sister branch was designated as B.73. Examples of canSNP assays that could be developed for typing purposes are B.71 (position 1 330 179, ancestral SNP state G/derived SNP state A), B.72 (337 274, G/T), B.73 (198 537, C/T), B.74 (1 304 315 G/A), and B.75 (1 827 247 G/A) in FSC200 (GenBank accession no. CP003862·1) reference genome. Furthermore, genome comparisons revealed that isolate A63/63 contained a B.39 subclade-specific deletion (9 bp, TAGTGATCT) within the 3' site of the pseudo-cas9 gene [32], near the region Ft-M22 used for VNTR analysis. Preliminary analysis indicated that this was a canonical indel (insertion and deletion) for subclade B.39, based on comparison with 25 draft genomes from three countries that could be targeted in typing assays.

Altogether, VNTR-typing, erythromycin susceptibility testing, and full genome sequencing of the three *F. tularensis* isolates from the beaver, red fox and the new branch B.71 (of basal clade B.12) isolate from the raccoon dog were of the erythromycin-resistant biovar II type (Fig. 3, Table 2)

DISCUSSION

Our investigation in the Berlin/Brandenburg region during 2012–2014 revealed that three (10%) out of 29 animals examined were *F. tularensis* positive by PCR analysis. Despite the limited number of animals, our results are in good agreement with the findings of Kuehn *et al.* [15], indicating a seroprevalence of 7·5% on average for the Brandenburg area. This is the first report of *F. tularensis* isolated from a beaver in Germany, and we could corroborate the presence of *F. tularensis* in raccoon dogs and red foxes previously suggested by serological studies [15]. The unexpectedly diverse animal species found *F. tularensis* positive




Fig. 3. Neighbour-joining phylogenetic tree showing the relationship between the three new genomes and seven published genomes. The evolutionary distances are in the units of the number of base differences per sequence. The tree is based on 1537 differences identified in an alignment of a total of 1 707 622 bp in common between the ten genomes. CanSNPs B.71–B.75 (circled in green) and strains FDC407, FDC408, and FDC409 are new to this study and are indicated by an asterisk (* this study). B.I, Biovar I; B.II, biovar II.

indicate that tularemia in wildlife is probably underdiagnosed.

We investigated the case of a dead, tularemia-positive beaver in detail. Culturing from environmental samples from locations close to the site where the dead beaver was found did not identify living *F. tularensis* in any of the environmental samples during 2012 or 2013. One explanation for this might be that cultivation of *F. tularensis* from environmental samples is difficult, mainly because the organism grows slowly and fastidiously, and its isolation can be prevented by overgrowth of the background flora despite using selective antibiotics. Another explanation might be the presence of a very low concentration of culturable *Francisella* bacteria within the environment. In contrast, qPCR analysis indicates that the *F. tularensis* bacteria were persistent in the aquatic environment during one climatic season, but apparently no further than that (see below). We could not detect significant levels of *F. tularensis* DNA by qPCR analysis of the distant sites or in the following year, indicating that the entire area investigated is not highly ‘contaminated’ by *F. tularensis* DNA.

The results of the pathological examination of the beaver (Fig. 2) indicated a chronic systemic infection with final exacerbation leading to massive replication of *F. tularensis* subsp. *holarctica* and subsequent contamination of the surrounding environment. We detected *F. tularensis* DNA in the bloody urine-contaminated snow collected around the dead beaver (data not shown). Contamination of the aquatic

environment with *F. tularensis* by infected voles and beavers (carcasses) has been reported from Finland and the United States [33–36]. In addition, chronic shedding of bacteria by infected voles has been suggested [33]. In the case of *F. tularensis* contamination of the environment caused by beaver carcasses in the United States, the bacteria were not able to persist in the aquatic environment for a long time [34]. However, the source of infection and the route of transmission in the case of the beaver which died from tularemia remain unclear.

Altogether, this implicates a mechanism where animals with high bacterial load may act as local amplifiers [35] and contaminate the immediate environment, resulting in local hotspots of *F. tularensis* subsp. *holarctica*. Thus, mammals would act as local amplifiers to facilitate the spread of the disease, e.g. through contaminated water or by vectors [30, 37, 38].

The three recovered *F. tularensis* isolates from the beaver, a raccoon dog, and a red fox were of the erythromycin-resistant biovar II type (Fig. 3, Table 2) and were members of the B.33/34 subclade (beaver), B.12/13 subclade (raccoon dog), and B.39 subclade (red fox). Members of the B.33 subclade (i.e. branches B.33–B.38) have previously been found in Germany, Hungary, Austria, and Scandinavia [28, 39], whereas the beaver isolate constitutes a new subclade B.75. The B.39 subclade has not been described previously in Germany but has been found in Sweden, Finland, and Poland [6, 29, 31].

Altogether it is remarkable that our analysis of a small set of animals indicates an unexpectedly high diversity of strains present in the Berlin/Brandenburg area. The finding of a new B.12/13 subclade designated B.71 (Fig. 3) that was distantly related to all other strains previously found in the basal clade B.12 reveals gaps in the knowledge of the phylogeography of *F. tularensis* subsp. *holarctica* and shows that the diversity of tularemia is high also in non-Scandinavian countries. Previous assumptions that the high genomic diversity of *F. tularensis* in Scandinavia was indicative of Scandinavia as the source of the historical spread of tularemia to the rest of Europe [29] could be owing, at least in part, to a sampling bias in Scandinavia and insufficient sampling in other regions of Europe, including Germany. Recent studies of German *F. tularensis* subsp. *holarctica* isolates from brown hares (*Lepus europaeus*) and from an outbreak in non-human primates in animal facilities in Germany also revealed a high genetic and phenotypic diversity of the isolates investigated [13, 40]. In the study of Antwerpen *et al.*, using isolates from animal facilities in Germany, only two strains belong to basal clade B.12, whereas 12 strains belong to the erythromycin-sensitive basal clade B.6 [40]. In the study of Müller *et al.*, 14/31 isolates were assigned to basal clade B.12, 16 to basal clade B.6, and one to basal clade B.4. The identified basal clade B.12 strains all exhibit the same PCR fragment size (480 bp) for VNTR marker Ft-M24 and were localized primarily in Eastern Germany [13]. This is in agreement with our results, showing that all three animal isolates were members of three different B.12 subclades (B.39, B.33/34, and B.12/13). However, the beaver isolate A271_1 in subclade B.33/34 showed a PCR fragment size for VNTR marker Ft-M24 of 459 bp (Table 2) underlining that the exclusive use of VNTR markers for typing may give rise to incongruences in the reconstruction of phylogenetic relationships between strains, leading to potentially wrong interpretation of strain relationships in *F. tularensis*. However, the VNTR analysis of the three different isolates from the beaver (A271_1-3) demonstrated that all three were identical (data not shown), indicating absence within host diversity. Other recent examples of high *Francisella* diversity in Germany include an emerging subclade of *F. tularensis* subsp. *holarctica* in *Ixodes ricinus* ticks from south-western Germany [12] and the first German *Francisella* isolate (W12-1067) not belonging to the species *F. tularensis* [41].

In conclusion, based on the assumption that high genetic diversity indicates longer presence of a pathogen in a region, one could hypothesize from our data a long presence of tularemia in the study region within East Germany. Moreover, our findings demonstrate that further investigations on *Francisella* isolates from different animal species, the environment, and from patients are necessary to better understand the occurrence, persistence, and distribution of *F. tularensis* subsp. *holarctica* and also the presence of putative further *Francisella* species in Germany.

SUPPLEMENTARY MATERIAL

For supplementary material accompanying this paper visit <http://dx.doi.org/10.1017/S0950268816001175>.

ACKNOWLEDGEMENTS

We acknowledge the technical contribution from Elin Nilsson, Kerstin Rydzewski, Silke Becker, Petra Lochau, Iris Klein, Adrian Lärkeryd, and Caroline Öhrman. We would like to thank Ursula Erikli for copy-editing. This work was supported in part by the Swedish Ministry of Defence (No. A404015) and by the Robert Koch Institute.

DECLARATION OF INTEREST

None.

REFERENCES

1. Ellis J, *et al.* Tularemia. *Clinical Microbiology Reviews* 2002; **15**: 631–646.
2. Saslaw S, *et al.* Tularemia vaccine study. I. Intracutaneous challenge. *Archives of Internal Medicine* 1961; **107**: 689–701.
3. Penn R. *Francisella tularensis* (tularemia). In: Mandell GL, Bennet JE, Dolin R (eds). *Mandell, Douglas and Bennett's Principles and Practice of Infectious Diseases*. Elsevier/Churchill Livingstone, Philadelphia, 2010, pp. 2927–2937.
4. Svensson K, *et al.* Landscape epidemiology of tularemia outbreaks in Sweden. *Emerging Infectious Diseases* 2009; **15**: 1937–1947.
5. Svensson K, *et al.* A real-time PCR array for hierarchical identification of *Francisella* isolates. *PLoS ONE* 2009; **4**: e8360.
6. Vogler AJ, *et al.* Phylogeography of *Francisella tularensis*: global expansion of a highly fit clone. *Journal of Bacteriology* 2009; **191**: 2474–2484.
7. Vogler AJ, *et al.* An optimized, multiplexed multi-locus variable-number tandem repeat analysis system for

- genotyping *Francisella tularensis*. *Letters in Applied Microbiology* 2009; **48**: 140–144.
8. **Oraggi FC, Frey J, Pilo P.** Characterisation of a new group of *Francisella tularensis* subsp. *holarctica* in Switzerland with altered antimicrobial susceptibilities, 1996 to 2013. *European Surveillance* 2014; **24**: 19.
 9. **Jenzora A, et al.** Seroprevalence study of *Francisella tularensis* among hunters in Germany. *FEMS Immunology and Medical Microbiology* 2008; **53**: 183–189.
 10. **Kaysser P, et al.** Re-emergence of tularemia in Germany: presence of *Francisella tularensis* in different rodent species in endemic areas. *BMC Infectious Diseases* 2008; **8**: 157.
 11. **Spletstoeser WD, et al.** Tularemia in Germany: the tip of the iceberg? *Epidemiology and Infection* 2009; **137**: 736–743.
 12. **Gehring H, et al.** Presence of an emerging subclone of *Francisella tularensis holarctica* in *Ixodes ricinus* ticks from south-western Germany. *Ticks and Tick-Borne Diseases* 2013; **4**: 93–100.
 13. **Muller W, et al.** German *Francisella tularensis* isolates from European brown hares (*Lepus europaeus*) reveal genetic and phenotypic diversity. *BMC Microbiology* 2013; **13**: 61.
 14. **Otto P, et al.** Serological investigation of wild boars (*Sus scrofa*) and red foxes (*Vulpes vulpes*) as indicator animals for circulation of *Francisella tularensis* in Germany. *Vector-Borne and Zoonotic Diseases* 2014; **14**: 46–51.
 15. **Kuehn A, et al.** Tularaemia seroprevalence of captured and wild animals in Germany: the fox (*Vulpes vulpes*) as a biological indicator. *Epidemiology and Infection* 2013; **141**: 833–840.
 16. **Nattermann H, et al.** First isolation of *Francisella tularensis* subspecies *holarctica* from foxes (*Vulpes vulpes*) in Germany. 6th International Conference on Tularemia 2009, Berlin, 13–16 September 2009 (<http://www.tularaemia-network.com/>), P1–43.
 17. **Becker S, et al.** Successful re-evaluation of broth medium T for the growth of *Francisella tularensis* ssp. and other highly pathogenic bacteria. *Journal of Microbiological Methods* 2016; **121**: 5–7.
 18. **Forsman M, Sandstrom G, Sjostedt A.** Analysis of 16S ribosomal DNA sequences of *Francisella* strains and utilization for determination of the phylogeny of the genus and for identification of strains by PCR. *International Journal of Systematic Bacteriology* 1994; **44**: 38–46.
 19. **Kirchner S, et al.** Pentaplexed quantitative real-time PCR assay for the simultaneous detection and quantification of botulinum neurotoxin-producing clostridia in food and clinical samples. *Applied and Environmental Microbiology* 2010; **76**: 4387–4395.
 20. **Broekhuijsen M, et al.** Genome-wide DNA microarray analysis of *Francisella tularensis* strains demonstrates extensive genetic conservation within the species but identifies regions that are unique to the highly virulent *F. tularensis* subsp. *tularensis*. *Journal of Clinical Microbiology* 2003; **41**: 2924–2931.
 21. **Johansson A, et al.** Worldwide genetic relationships among *Francisella tularensis* isolates determined by multiple-locus variable-number tandem repeat analysis. *Journal of Bacteriology* 2004; **186**: 5808–5818.
 22. **Simpson JT, et al.** ABySS: a parallel assembler for short read sequence data. *Genome Research* 2009; **19**: 1117–1123.
 23. **Tamura K, et al.** MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Molecular Biology and Evolution* 2011; **28**: 2731–2739.
 24. **Nei M, Kumar S.** *Molecular Evolution and Phylogenetics*. Oxford University Press, New York, 2000.
 25. **Lärkeryd A, et al.** CanSNPer: a hierarchical genotype classifier of clonal pathogens. *Bioinformatics* 2014; **30**: 1762–1764.
 26. **Birdsell DN, et al.** *Francisella tularensis* subsp. *novicida* isolated from a human in Arizona. *BMC Research Notes* 2009; **2**: 223.
 27. **Chanturia G, et al.** Phylogeography of *Francisella tularensis* subspecies *holarctica* from the country of Georgia. *BMC Microbiology* 2011; **11**: 139.
 28. **Gyuranecz M, et al.** Phylogeography of *Francisella tularensis* subsp. *holarctica*, Europe. *Emerging Infectious Diseases* 2012; **18**: 290–293.
 29. **Karlsson E, et al.** The phylogeographic pattern of *Francisella tularensis* in Sweden indicates a Scandinavian origin of Eurosiberian tularaemia. *Environmental Microbiology* 2013; **15**: 634–645.
 30. **Karadenizli A, et al.** Genomic analyses of *Francisella tularensis* strains confirm disease transmission from drinking water sources, Turkey, 2008, 2009 and 2012. *Eurosurveillance* 2015; **20**.
 31. **Sissonen S, et al.** Phylogeography of *Francisella tularensis* subspecies *holarctica* in Finland, 1993–2011. *Infectious Diseases (London)* 2015; **47**: 705–710.
 32. **Schunder E, et al.** First indication for a functional CRISPR/Cas system in *Francisella tularensis*. *International Journal of Medical Microbiology* 2013; **303**: 51–60.
 33. **Bell JF, Stewart SJ.** Chronic shedding tularaemia nephritis in rodents: possible relation to occurrence of *Francisella tularensis* in lotic waters. *Journal of Wildlife Diseases* 1975; **11**: 421–4230.
 34. **Parker RR, et al.** Contamination of natural waters and mud with *Pasteurella tularensis* and tularemia in beavers and muskrats in the northwestern United States. *Bulletin of the National Institute of Health* 1951; **193**: 1–161.
 35. **Rossow H, et al.** Experimental infection of voles with *Francisella tularensis* indicates their amplification role in tularaemia outbreaks. *PLoS ONE* 2014; **9**: e108864.
 36. **Rossow H, et al.** Detection of *Francisella tularensis* in voles in Finland. *Vector-Borne and Zoonotic Diseases* 2014; **14**: 193–198.
 37. **Backman S, et al.** Transmission of tularemia from a water source by transstadial maintenance in a mosquito vector. *Scientific Report* 2015; **5**: 7793.
 38. **Boone I, et al.** Tularaemia in southwest Germany: Three cases of tick-borne transmission. *Ticks and Tick-borne Diseases* 2013; **6**: 611–614.
 39. **Gyuranecz M, et al.** Factors influencing emergence of tularemia, Hungary, 1984–2010. *Emerging Infectious Diseases* 2012; **18**: 1379–1381.

- 40 **Antwerpen MH, et al.** Rapid high resolution genotyping of *Francisella tularensis* by whole genome sequence comparison of annotated genes (MLST⁺). *PLoS ONE* 2015; **10**: e0123298.
41. **Rydzewski K, et al.** Genome sequence and phenotypic analysis of a first German *Francisella* sp. isolate (W12-1067) not belonging to the species *Francisella tularensis*. *BMC Microbiology* 2014; **14**: 169.