

Highly Efficient Isolation of *Populus* Mesophyll Protoplasts and Its Application in Transient Expression Assays

Jianjun Guo, Jennifer L. Morrell-Falvey, Jessy L. Labbé, Wellington Muchero, Udaya C. Kalluri, Gerald A. Tuskan, Jin-Gui Chen*

Biosciences Division, Oak Ridge National Laboratory, Oak Ridge, Tennessee, United States of America

Abstract

Background: *Populus* is a model woody plant and a promising feedstock for lignocellulosic biofuel production. However, its lengthy life cycle impedes rapid characterization of gene function.

Methodology/Principal Findings: We optimized a *Populus* leaf mesophyll protoplast isolation protocol and established a *Populus* protoplast transient expression system. We demonstrated that *Populus* protoplasts are able to respond to hormonal stimuli and that a series of organelle markers are correctly localized in the *Populus* protoplasts. Furthermore, we showed that the *Populus* protoplast transient expression system is suitable for studying protein-protein interaction, gene activation, and cellular signaling events.

Conclusions/Significance: This study established a method for efficient isolation of protoplasts from *Populus* leaf and demonstrated the efficacy of using *Populus* protoplast transient expression assays as an *in vivo* system to characterize genes and pathways.

Citation: Guo J, Morrell-Falvey JL, Labbé JL, Muchero W, Kalluri UC, et al. (2012) Highly Efficient Isolation of *Populus* Mesophyll Protoplasts and Its Application in Transient Expression Assays. PLoS ONE 7(9): e44908. doi:10.1371/journal.pone.0044908

Editor: Carl Ng, University College Dublin, Ireland

Received: April 30, 2012; **Accepted:** August 9, 2012; **Published:** September 13, 2012

Copyright: © 2012 Guo et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by the Plant-Microbe Interfaces Scientific Focus Area in the Genomic Science Program, United States Department of Energy, Office of Science, Biological and Environmental Research. Oak Ridge National Laboratory is managed by UT-Battelle, LLC, for the United States Department of Energy under contract DE-AC05-00OR22725. JG was supported by a postdoctoral fellowship from the Natural Sciences and Engineering Research Council of Canada. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: chenj@ornl.gov

Introduction

The establishment of renewable energy sources is essential for energy security and reduction of CO₂ emissions. Among all of the options, production of lignocellulosic ethanol from plant feedstock has become one of the favorable approaches [1,2,3,4]. *Populus* is characterized by a rapid growth rate, high productivity and wide adaption to environment and therefore, has become one of the promising bioenergy feedstocks [1,5,6]. With the availability of its full genome sequence [7] and the establishment of transgenic techniques [8,9,10], *Populus* is also emerging as a model organism for studying woody plants [11].

One of the rate-limiting steps in studying the function of genes in *Populus* is its long life cycle (e.g., it takes a minimum 6 to 12 years to reach reproductive maturity [7]). Furthermore, it typically takes four to eight months from inoculation to generate rooted seedlings via *Agrobacterium*-mediated transformation approach [12]. Therefore, the capacity of conventional genetic manipulation in *Populus*, one of the most commonly used approaches for studying gene function, is largely limited by the lengthy time required to obtain suitable genetic materials. This limitation is more pronounced when dealing with cellular signaling events that often involve multiple gene products.

Leaf mesophyll protoplasts have been developed as an informative system in *Arabidopsis* and a few other plant species for rapid dissection of signaling events that lead to the regulation of gene expression, as well as for many other molecular characterizations, such as protein-protein interaction, subcellular localization, and protein-DNA interaction [13,14]. *Arabidopsis* leaf mesophyll protoplasts have also been used to study the function of *Populus* genes [15,16]. However, the inherently different genomic, physiological and developmental characteristics between a herbaceous plant and a perennial tree species may render *Arabidopsis* protoplasts a less ideal system for dissecting the signaling events that regulate the development of *Populus*.

In this paper, we described the micro-propagated *Populus* grown on MS media as an excellent alternative starting material for the efficient isolation of leaf mesophyll protoplasts. We also optimized enzyme concentrations and digestion time for efficient protoplast isolation. We then established an efficient *Populus* protoplast transient expression system and demonstrated the efficacy of using this *in vivo* system to study the activity of *Populus* genes.

Results and Discussion

Isolation of Protoplasts from *Populus* Leaves

A protocol for the isolation of *Arabidopsis* mesophyll protoplasts has been well established [14,17,18]. There have been three reports on isolation of protoplasts from *Populus*. Two of them reported the isolation of protoplasts from *Populus* cell suspension culture [19] or leaves [20], and the other used *Populus* protoplasts to study protein subcellular localization of *Populus* salt overly sensitive genes [21]. In the latter study, it was briefly mentioned that an essentially same procedure was used as that in *Arabidopsis* [21]. Because this is no *Populus* cell suspension culture available, we tried to establish a robust protocol for isolating protoplasts from *Populus* leaves. Initially, we tried to isolate protoplasts from leaves of soil-grown *Populus* plants cultivated in a greenhouse. Newly emerged leaves from two different *Populus* clones (*Populus trichocarpa* clone Nisqually-1 and *P. tremula* X *alba* clone 717-1B4) were used for the study. No or very few protoplasts could be isolated from these plants (data not shown). Field or greenhouse-grown *Populus* under adequate irradiation conditions usually produce leaves with a thick cuticle and recalcitrant cell walls, which are likely to be the major factors limiting protoplast isolation. Because cuticle thickness is known to increase with high light intensity and low relative humidity [22], we then propagated *Populus* 717 on MS medium in a Magenta box through shoot propagation under relatively low light ($70 \mu\text{mol m}^{-2} \text{s}^{-1}$) conditions [23]. The propagated *Populus* plants were able to produce fully expanded and healthy-looking leaves (Figure 1A). We found that protoplasts with high yield and quality could be isolated from these leaves (Figure 1). Evans blue staining indicated that more than 90% of the protoplasts isolated were viable ($n > 500$) (Figure S1).

More importantly, when we followed the same protocol for isolating protoplasts from leaf mesophyll cells of *Populus* plants grown in a greenhouse or a Magenta box as that used in *Arabidopsis*, we found that the yields and quality of protoplasts were very poor. Therefore, we modified the protocol by adjusting enzyme concentrations and digestion time. We found that twice the enzyme concentration and a longer digestion time (4–5 h) are required for an optimal yield of protoplasts from *Populus* leaves. We could routinely obtain 1×10^7 protoplasts from one-gram (fresh weight) leaf tissue in 10 ml of enzyme solution. The high yield of protoplasts permits the application of this system for transfection studies [14].

Establishment of a *Populus* Protoplast Transient Expression System

We next examined whether *Populus* protoplasts can be efficiently transfected. The protoplasts were transfected with a 35S::GFP (green fluorescent protein) construct using a PEG (Polyethylene glycol)-mediated transfection approach [14]. In general, a greater than 50% transfection efficiency is required for obtaining reliable and reproducible results with the protoplast system [14]. After incubation overnight, a strong GFP signal could be observed in more than 70% ($n > 1,000$) of the protoplasts transfected with a 35S::GFP construct (Figure 1B). These results indicated that the transfection efficiency of *Populus* protoplasts was sufficiently high to be used for molecular assays.

Transiently Expressed Organelle Markers are Correctly Localized in the *Populus* Protoplasts

The next question we asked was whether the products of exogenously-introduced genes can be targeted to the proper subcellular locations. To test this, we expressed several well-known organelle-targeting proteins in *Populus* protoplasts [24] including a

plasma membrane marker (Figure 2A), a Golgi apparatus marker (Figure 2B), a nucleus marker (Figure 2C), a peroxisome marker (Figure 2G), an ER (endoplasmic reticulum) marker (Figure 2H), and an ubiquitously-localized protein (RACK1, Receptor for Activated C-protein Kinase 1) (Figure 2I) [25]. After overnight incubation for protein expression, all of the proteins were correctly targeted (Figure 2). These results indicate that the *Populus* protoplast system is suitable for studying protein subcellular localization as well as function.

Use the *Populus* Protoplast Transient Expression System to Study Protein-Protein Interaction

In order to examine whether the *Populus* protoplast transient expression system can be exploited to study protein-protein interaction, a previously identified interaction between *Arabidopsis* RACK1A and eIF6A (eukaryotic initiation factor 6A) [25] was tested in *Populus* protoplasts using a bi-molecular fluorescence complementation (BiFC) approach [26]. AtRACK1A was fused with the N-terminal half of YFP (Yellow Fluorescent Protein) and AteIF6A was fused with the C-terminal half of YFP. When both constructs were co-transfected into *Populus* protoplasts, a yellow fluorescent signal was detected (Figure 3), indicating a positive interaction between these two proteins. As a negative control, GUS (β -glucuronidase) was not found to interact with either RACK1A or eIF6A (Figure 3). These results demonstrated the usefulness of *Populus* protoplast system for studying protein-protein interactions.

Populus Protoplasts are Responsive to Exogenous Plant Hormone Treatments

It has been shown that plant hormones can trigger the expression of downstream genes of the respective signaling cascades [27,28,29,30] in *Arabidopsis* mesophyll protoplasts. Thus, the *Arabidopsis* mesophyll protoplasts have been used as an *in vivo* system for studying the dynamic interaction of molecules in many signaling pathways. We therefore examined the transcriptional response of *Populus* protoplasts to exogenous plant hormone application. We focused on the hormones that were known to play important roles in xylogenesis, including auxin, gibberellin (GA), cytokinin and ethylene [31]. The *Populus* homologues of *AthGH3.2* (*POPTR_01s30560*), a commonly used molecular marker for auxin response, was used as a reporter for auxin response [29]. Its expression was up-regulated 3.5 fold in response to 5 μM NAA (1-naphthaleneacetic acid, a synthetic auxin) treatment (Figure 4A). For gibberellin responses, *POPTR_14s08030* was chosen because it is a close homologue of *At2g45900*, a direct target of the DELLA protein that can be induced by GA treatment in *Arabidopsis* [32]. An approximately 4-fold increase of its expression was observed in response to 50 μM GA₃ (gibberellic acid) treatment (Figure 4B). *POPTR_10s00320*, also known as *PtRR13*, is a type B *Arabidopsis* Response Regulator (ARR) that was reported to respond to the treatment of BAP (6-benzylaminopurine, a synthetic cytokinin) in *Populus* leaves [33]. The expression of *PtRR13* increased 6 to 16 fold in response to 0.5 to 5 μM BAP treatments (Figure 4C). Finally, a *Populus* homologue of *Arabidopsis* type-B ARR, *POPTR_10s08300*, was chosen as a molecular marker for ethylene response [34]. About a 4-fold increase was observed in protoplasts treated with 10 μM ACC (1-aminocyclopropane-1-carboxylic acid, a precursor for ethylene) (Figure 4D). Auxin [35], ethylene [34], gibberellin [36], and cytokinin [37] have also been reported to promote secondary growth via stimulating cambial cell division in *Populus*. The validation of the presence of these hormone signaling cascades in




Figure 1. *Populus* leaf mesophyll protoplasts. (A) Optimal yield and quality of protoplasts can be isolated from one month-old *Populus* plants grown on MS medium in a magenta box. (B) High transfection efficiency is indicated with GFP signal.
doi:10.1371/journal.pone.0044908.g001

protoplasts enables us to study their potential roles in the regulation of secondary cell wall biosynthesis in the future.

In order to determine whether the *Populus* protoplasts respond to exogenous hormone treatment in a similar manner as that in intact leaves, we further examined the induction of these genes by plant hormones in intact *Populus* leaves. All these genes responded to the respective hormone treatments as they did in protoplasts (Figure 4E-H). Moreover, the level of transcriptional induction by plant hormones was largely similar between in protoplasts and in intact leaves.

Use the *Populus* Protoplast Transient Expression System to Study Signaling Events

In order to further examine the feasibility of utilizing *Populus* leaf mesophyll protoplasts to study signal transduction events, we explored an energy-signaling event that was discovered using *Arabidopsis* mesophyll protoplasts [38]. AthKIN10 (SNF1 kinase homolog 10) is recognized as an energy sensor that perceives the cellular energy status and reprograms the transcriptome for cellular energy homeostasis. One of the reporter genes in *Arabidopsis* that was up-regulated by AthKIN10 is *DIN6* (*At3g47340*, Dark Inducible 6). We searched the *Populus* genomic information in Phytozome (<http://www.phytozome.net/>) and found three sequence homologues of *DIN6*: *POPTR_09s07710.1* (*PtrDIN6-1*, with 91.1% similarity), *POPTR_01s28500.1* (*PtrDIN6-2*, with 91.3% similarity) and *POPTR_05s07720.1* (*PtrDIN6-3*, with

82% similarity). The transcript of *PtrDIN6-3* was not detected in protoplasts under our experimental conditions, and therefore was not studied further.

We first examined the response of *PtrDIN6-1* and *PtrDIN6-2* to dark and hypoxia, conditions that create cellular energy starvation. We found that both *DIN6* homologues showed high induction by these two conditions (Figure 5A). We then expressed AthKIN10 in *Populus* protoplasts to examine the effect of this energy sensor on the expression of *PtrDIN6* genes. We found that both *PtrDIN6* genes were strongly induced by ectopic expression of AthKIN10 (Figure 5B).

We further extended the experiments to test the effect of transient overexpression of *Populus* homologues of *AtKIN10*. Three *KIN10* homologues were identified via phytozome: *POPTR_04S11500*, *POPTR_17S12380* and *POPTR_13S09420*. While both *POPTR_04S11500* and *POPTR_17S12380* were expressed ubiquitously, *POPTR_13S09420* had extremely low expression (Figure 5C, Figure S2). We found that similar to *AtKIN10*, overexpression of both *POPTR_04S11500* and *POPTR_17S12380* induced the expression of *PtrDIN6* genes in *Populus* protoplasts (Figure 5D).

To further validate the induction of *PtrDIN6* by *PtrKIN10*, we constructed a reporter cassette in which the genomic DNA sequence that is 2 kb upstream of the start codon of *PtrDIN6-2* was fused with GUS reporter gene. When both *POPTR_0017s12380* and the *PtrDIN6-2* promoter fused with GUS reporter were co-




Figure 2. Subcellular localization of various organelle markers in *Populus* protoplasts. (A) Plasma membrane; (B) Golgi apparatus; (C) Nucleus; (G) Peroxisome; (H) endoplasmic reticulum (ER); (I) An ubiquitously-localized protein (RACK1, Receptor for Activated C-protein Kinase 1). Shown in (D), (E), (F), (J), (K) and (L) are bright field images for fluorescent images of (A), (B), (C), (G), (H) and (I), respectively. The organelle markers were fused with mCherry fluorescent protein, and RACK1 was fused with YFP fluorescent protein. The mCherry signal was separated from chloroplast autofluorescence using spectral imaging and linear unmixing. The mCherry and YFP signals are false-colored green and the chloroplast autofluorescence is shown in red. Scale bar, 1 μ m.
doi:10.1371/journal.pone.0044908.g002

transfected into the *Populus* protoplasts, we detected a 10-fold increase of GUS activity (Figure 5E), indicating that *Populus* KIN10 protein encoded by *POPTR_0017s12380* can be recruited to the promoter region of *PtDIN6-2* to activate the expression of *PtDIN6-2*.

Taken together, these results suggested the presence of a potentially conserved energy sensing signaling cascade mediated by KIN10 in *Populus*, and further demonstrated the suitability of using *Populus* protoplasts to study signal transduction events.




Figure 3. Using *Populus* protoplast system to test protein-protein interaction. Arabidopsis RACK1A (AtRACK1A) and eIF6A (AtelF6A) were fused with the N-terminal and C-terminal half of YFP, respectively. GUS-nYFP/cYFP fusions were used as negative controls. YFP, yellow fluorescent protein. BF, Bright field.
doi:10.1371/journal.pone.0044908.g003

Concluding Remarks

The Arabidopsis mesophyll protoplast transient expression system has proven to be an informative system to characterize genes and pathways. However, the use of *Populus* mesophyll protoplast transient expression system is still in its infancy. The optimal plant growth conditions associated with highly efficient protoplast isolation, the optimal enzyme concentration and digestion time, the yield of protoplasts, the estimate of transfection efficiency, and the use of the system for studying the effect of exogenously-introduced genes had not been described previously. All of these parameters are essential for the efficient application of this system. We have reported these parameters here.

We established a protocol for the isolation of high yield, physiologically active and transfection-ready protoplasts from *Populus* leaves. We validated the suitability of this system for studying gene activity *in vivo* by showing the correct subcellular localization of introduced proteins, protein-protein interaction, gene activation, as well as the molecular responses of protoplasts to plant hormones. As a further test of the system, we observed a potentially conserved cellular energy signaling pathway in *Populus* protoplasts. The establishment of an efficient *Populus* protoplast isolation and transfection system provides a useful approach to characterize *Populus* genes and pathways *in vivo*.

Because protoplasts were derived from single cell type and deprived of cell wall, hypothesis and conclusions generated from transient expression study with this system should be cautiously analyzed and supplemented with whole plant studies. Furthermore, the development of protein extraction protocol for *Populus* protoplasts would help make the *Populus* protoplasts a more robust and reliable system for studying protein-protein interactions and their involvement in signaling events. Because *Populus* is a promising feedstock for lignocellulosic biofuel production, it is worthwhile to further investigate whether one could draw reliable conclusions from observations in leaf mesophyll protoplasts for lignin-producing tissue. We have showed that *Populus* mesophyll protoplasts responded to exogenous plant hormone treatments in a manner similar to intact leaves (Figure 4). Earlier studies in *Zinnia elegans* showed that protoplasts isolated from mesophyll can be induced to mature tracheary elements [39,40], the units of xylem featured with various patterns of secondary cell wall deposition. In addition, ectopic expression of SND1, a transcription factor regulating the expression of secondary cell wall biosynthesis, induced ectopic deposition of secondary cell wall in mesophyll cells in *Arabidopsis* [41]. Taken together, these studies suggested that

Populus mesophyll protoplasts have the potential to be used for dissecting the functional aspects of secondary cell biosynthesis and deposition. This represents a promising area for future investigation.

Materials and Methods

Plant Growth Conditions

Experiments were performed with micropropagated *Populus* clone 717-1B4 (female, *Populus tremula* x *alba*). Fresh, young shoots from plants grown in a greenhouse were used as starting materials for plant propagation in media after surface-sterilization [5 min in 1% (v/v) Tween-20 solution, 1 min in 70% (v/v) ethanol, 15 min in 10% (v/v) bleach solution and then triple rinsed for 5 min in sterile water]. The plants were maintained in Magenta boxes (GA-7) on 100 ml of MS medium with micro and macronutrients, vitamins, and glycine (Murashige and Skoog, Sigma-Aldrich) supplemented with 1% (w/v) sucrose, 0.5 g MES (2-(N-Morpholino) ethanesulfonic acid sodium salt) hydrate (Sigma-Aldrich) and solidified with 0.8% (w/v) agar, and pH adjusted to 5.7. The plants were cultivated in a tissue culture growth room: 25°C with a 24-h photoperiod. The light is provided by fluorescence tubes (cool white, 95 W, F96T12/CW/HO/SS, Sylvania) at an intensity of 70 $\mu\text{mol m}^{-2} \text{s}^{-1}$ and plants were multiplied every 6 to 8 weeks [23].

Protoplast Isolation

Healthy, fully expanded *Populus* leaves were cut into 1–2 mm fine strips and digested in an enzyme solution [0.4 M mannitol, 20 mM KCl, 20 mM MES-KOH (pH 5.7), 3% (w/v) cellulase R10 (Yakult Pharmaceutical Ind. Co., Ltd., Japan), 0.8% (w/v) macerozyme R10 (Yakult Pharmaceutical Ind. Co., Ltd., Japan), 10 mM CaCl_2 , 5 mM β -mercaptoethanol and 0.1% (w/v) bovine serum albumin (BSA)] in the dark for 5 h with gentle shaking on a Boeker Rocker II (Boeker Inc.). The protoplasts were harvested by filtering through a 70 μm pore nylon cloth (Carolina Biological Supply Company) and then suspended in a transfection buffer (0.4 M mannitol, 15 mM MgCl_2 , 4 mM MES-KOH (pH 5.7)) to a concentration of 1×10^6 protoplasts ml^{-1} .

Protoplast Transfection

For each transfection assay used for qRT-PCR analysis, 50 μg plasmid [$1 \mu\text{g} \mu\text{l}^{-1}$, prepared with Qiagen Midi Prep Kit (Qiagen




Figure 4. The response of *Populus* protoplasts to various plant hormone treatments. Shown are the change *POPTR_01s30560* transcript in response to different concentrations of NAA in protoplasts (A) and intact leaves (E), the change of *POPTR_14s08030* transcript in response to different concentrations of GA₃ in protoplasts (B) and intact leaves (F), the change of *POPTR_10s00320* transcript in response to different concentrations of BAP in protoplasts (C) and intact leaves (G), and the change of *POPTR_10s08300* transcript in response to different concentrations of ACC in protoplasts (D) and intact leaves (H). The protoplasts or intact leaves were incubated with various concentrations of plant hormones for 3h before being harvested for qRT-PCR analysis. The experiments were repeated three times with similar results. The averages of three technical replicates \pm standard errors are shown. * indicates a significant difference (at $P \leq 0.01$, student's t-test) between each treatment and the untreated control. doi:10.1371/journal.pone.0044908.g004

Inc.) was added to 500 μ l of protoplast suspension, to which 550 μ l PEG solution [40% (w/v) polyethylene glycol (Sigma-Aldrich, MW 4000), 0.2 M mannitol, 0.1 M CaCl₂] was added. For transfection assays used for GUS reporter analysis, 10 μ g effector and 10 μ g reporter plasmids were transfected with 100 μ l protoplast, to which 120 μ l PEG solution was added. The mixture was incubated at room temperature for 5 min before 2.2 ml of W5 solution [154 mM NaCl, 125 mM CaCl₂, 5 mM KCl, 2 mM MES-KOH (pH 5.7)] were added to stop the reaction. The protoplasts were collected by centrifugation at 1,200 rpm (331 g) for 2 min on a swing-out rotor and then suspended overnight in 5 ml of incubation buffer [0.5 mM mannitol, 4 mM MES-KOH (pH 5.7), 20 mM KCl] in a 100 mm² petri-dish.

Imaging Methods

A set of vectors that transiently-expressed different organelle markers fused with mCherry [24] and an ubiquitously-expressed RACK1-YFP [25] were used for PEG-mediated transfection in *Populus* protoplasts. Images were captured using a Zeiss LSM 710 confocal laser scanning microscope equipped with a PlanAchromat 40x/1.4NA oil objective. Spectral imaging and linear unmixing were used to separate chloroplast autofluorescence from the mCherry-specific signal. Briefly, the cells were scanned with a 561 nm laser line and lambda-stack images were collected at a spectral resolution of 9.7 nm from 569–729 nm. Optical sections were collected at a z-scaling of 1 micron. After image acquisition, each set of lambda 3D confocal z-stacks was linearly unmixed with Zen 2009 software (Zeiss). The spectra (emission fingerprints) from unlabeled chloroplasts and the mCherry fusion proteins were unique in shape and were used to generate a 2-channel confocal stack representing chloroplasts (shown in red) and mCherry fusion protein (shown in green). Unmixed confocal stacks are displayed as maximum intensity projections using Zen 2009 software.

Other Methods

Total RNA was extracted from 5×10^5 protoplasts for each sample using Trizol® (Invitrogen Inc.). Two-hundred-fifty microliters of Trizol® was used for each RNA extraction, and 100 μ g Ambion® Glycogen was added in the RNA precipitation step as carrier. Five hundred nanograms of total RNA was used for reverse transcription using RevertAid™ Reverse Transcriptase (Fermentas Inc.) and oligo dT₁₆ was used as primer. The real-time PCR primers were designed using the NCBI Primer-BLAST tool [42]. The specificity of each pair of primers was determined by BLASTing the primers against Reference RNA sequence database for *Populus trichocarpa* (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>). The real-time PCR reactions were conducted on a StepOne Plus™ Realtime PCR system (Applied Biosystems) with the iTaq™-SYBR® Green Super Mix with ROX (Bio-RAD Inc.). The expression of an ubiquitin gene *POPTR_01s44440* was used to standardize the expression of each gene. A 35S::GFP (ABRC stock #: CD3-911) construct was co-transfected for each samples to monitor the transfection efficiency for each assay. Only assays with >60% transfection efficiency were used for qRT-PCR analysis.

For KIN10-DIN6 study, AthKIN10 and PtrKIN10 was each cloned into a plant cell transient expression vector pE3225 [26]. The GFP sequence of pE3225 was removed and replaced with double HA (Human influenza hemagglutinin) tag. The genomic sequence 2 kb upstream of the start codon of *PtrDIN6-2* was cloned and fused to a *GUS* reporter gene by replacing the *UBQ10* promoter of the HBT95-pUBQ10-GUS construct [43]. For co-transfection assay, a 35S::LUC (Luciferase) was included in each transfection and the LUC activity was used to normalize *GUS* activity to account for the potential variations in the transfection




Figure 5. Energy sensing signaling in *Populus* protoplasts. (A) Semi-quantitative RT-PCR analysis of *PtrDIN6* transcripts in response to dark and hypoxia treatments. L-Light; D-Dark; H-Hypoxia. The *PtrUBQ10* gene was used as a control. (B) The change of *PtrDIN6* transcripts in response to overexpression of AthKIN10 protein. After transfection, protoplasts were incubated overnight to allow the expression of AthKIN10 before samples were harvested for qRT-PCR and western blot analysis. Western blot was used to detect the presence of the introduced HA-tagged AthKIN10 protein. The experiments were repeated three times with similar results. The averages of three technical replicates \pm standard errors are presented in the graph. * indicates a significant difference (at $P \leq 0.01$, student's t-test) between protoplasts expressing AthKIN10 and the control (ctrl). (C) The expression of three *Populus KIN10* homologues in transfected protoplasts examined by semi-quantitative RT-PCR. The expression of *PtrUBQ10* was used as an internal control. (D) The response of *PtrDIN6* transcript to the overexpression of three *Populus KIN10* homologues. The experiments were repeated three times with similar results. The averages of three technical replicates \pm standard errors are shown. Protoplasts transfected with an empty vector was used as control (ctrl) for each comparison. (E) The activation of *PtrDIN6* by *PtrKIN10* in a GUS reporter assay. For each co-transfection, a 35S::LUC (Luciferase) was included and the LUC activity was used to normalize GUS activity to account for the potential variations in the transfection efficiency. The averages of three technical replicates \pm standard errors are shown. * indicates a significant difference (at $P \leq 0.01$, student's t-test) between each treatment and the control (ctrl). doi:10.1371/journal.pone.0044908.g005

efficiency [14]. The constructs used for BiFC were the same as those used previously [25].

For semi-quantitative RT-PCR, 1 μ l cDNA for each sample was used. The PCR was conducted on a vertiTM 96-well PCR machine (Applied Biosystems) for 30 cycles with the same primers used for qRT-PCR. All primers used in this study are listed in the Table S1.

For the detection of AthKIN10-HA, 2×10^5 protoplasts were lysed in 2xSDS PAGE protein loading buffer (Bio-Rad) and the protein bands were separated in a 10% SDS-PAGE gel. The proteins were then transferred onto a PVDF membrane using a semi-dry transferring cassette (Bio-Rad). For western blotting, the membrane was blocked with TBS containing 0.05% Tween-20 and 2% non-fat dry milk for 1 h. Anti-HA-HRP (Roche Inc) was diluted to 1:10,000 in TBS (25 mM Tris-HCl, 137 mM NaCl) and incubated with the blot for 1 h. The blot was washed three times (5 min each time) with TBS containing 0.05% Tween-20 and the signal was visualized by adding Pierce ECL western blotting substrate (Pierce Inc.) and exposure to X-ray film (Thermo Scientific).

For dark induction, the protoplasts were kept in the dark for 6 h whereas the control samples were kept in constant light with

$70 \mu\text{mol m}^{-2} \text{s}^{-1}$. For hypoxia induction, protoplasts were settled at the bottom of a 2 ml Eppendorf tube filled with incubation buffer solution for 6 h while the control samples were kept in a 100 mm (diameter) \times 15 (height) mm petri-dish with 0.2 mm depth of solution. For hormone treatments, plant hormones with different concentration were added in the incubation buffer together with protoplasts or whole leaves and incubated for 3 h before they were harvested for qRT-PCR analysis. The method for GUS activity assay was essentially the same as described previously [14].

The *Populus* homologues were identified using the Phytozome web tool (<http://www.phytozome.net>) [44]. Briefly, the locus numbers of Arabidopsis genes were used to locate the Arabidopsis gene. The closest *Populus* homologs identified under the "homolog peptide" tool were chosen as candidate homologues.

All of the primer sequences used in this paper are listed in the Table S1.

Supporting Information

Figure S1 Representative images of Evans Blue staining of *Populus* protoplasts.

(TIF)

Figure S2 The expression of three *Populus KLN10* homologues determined by the eFP tool (bar.toronto.ca).

(TIF)

Table S1 List of primers used in this study.

(XLSX)

Acknowledgments

We want to especially thank Dr. Jen Sheen (Department of Genetics, Harvard Medical School) for training J.G. on the protoplast system in her lab, Dr. Andreas Nebenfuhr (Department of Biochemistry and Cellular and Molecular Biology, University of Tennessee) for sharing the set of organelle vectors, Dr. Stanton Gelvin (Purdue University) for sharing the pE3225 construct and constructs for BiFC analysis, Dr. Olaf Czarnecki

References

- Carroll A, Somerville C (2009) Cellulosic biofuels. *Annu Rev Plant Biol* 60: 165–182.
- Dinus RJ (2001) Genetic improvement of poplar feedstock quality for ethanol production. *Appl Biochem Biotechnol* 91–93: 23–34.
- Ragauskas AJ, Williams CK, Davison BH, Britovsek G, Cairney J, et al. (2006) The path forward for biofuels and biomaterials. *Science* 311: 484–489.
- Somerville C (2006) The billion-ton biofuels vision. *Science* 312: 1277.
- Simmons BA, Loque D, Blanch HW (2008) Next-generation biomass feedstocks for biofuel production. *Genome Biol* 9: 242.
- Yuan JS, Tiller KH, Al-Ahmad H, Stewart NR, Stewart CN Jr (2008) Plants to power: bioenergy to fuel the future. *Trends Plant Sci* 13: 421–429.
- Tuskan GA, Difazio S, Jansson S, Bohlmann J, Grigoriev I, et al. (2006) The genome of black cottonwood, *Populus trichocarpa* (Torr. & Gray). *Science* 313: 1596–1604.
- Song J, Lu S, Chen ZZ, Lourenco R, Chiang VL (2006) Genetic transformation of *Populus trichocarpa* genotype Nisqually-1: a functional genomic tool for woody plants. *Plant Cell Physiol* 47: 1582–1589.
- Han KH, Meilan R, Ma C, Strauss SH (2000) An *Agrobacterium tumefaciens* transformation protocol effective on a variety of cottonwood hybrids (genus *Populus*). *Plant Cell Reports* 19: 315–320.
- Tzifira T, Jensen CS, Wang WX, Zuker A, Vinocur B, et al. (1997) Transgenic *Populus tremula*: a step-by-step protocol for its *Agrobacterium*-mediated transformation. *Plant Molecular Biology Reporter* 15: 219–235.
- Bradshaw HD, Ceulemans R, Davis J, Stettler R (2000) Emerging model systems in plant biology: Poplar (*Populus*) as a model forest tree. *Journal of Plant Growth Regulation* 19: 306–313.
- Busov VB, Strauss SH, Pilate G (2010) Transformation as a Tool for Genetic Analysis in *Populus*. In: *Genetics and Genomics of Populus* (Jansson S, Bhalerao R, Groover A, eds), Springer New York. 113–133.
- Faraco M, Di Sanebastiani GP, Spelt K, Koes RE, Quattrocchio FM (2011) One protoplast is not the other! *Plant Physiol* 156: 474–478.
- Yoo SD, Cho YH, Sheen J (2007) Arabidopsis mesophyll protoplasts: a versatile cell system for transient gene expression analysis. *Nat Protoc* 2: 1565–1572.
- Zhong R, Ye ZH (2010) The poplar PtrWINDs are transcriptional activators of secondary cell wall biosynthesis. *Plant Signal Behav* 5: 469–472.
- Zhong R, McCarthy RL, Lee C, Ye ZH (2011) Dissection of the transcriptional program regulating secondary wall biosynthesis during wood formation in Poplar. *Plant Physiol* 157: 1452–1468.
- Tiwari S, Wang S, Hagen G, Guilfoyle TJ (2006) Transfection assays with protoplasts containing integrated reporter genes. *Methods Mol Biol* 323: 237–244.
- Sheen J (2001) Signal transduction in maize and Arabidopsis mesophyll protoplasts. *Plant Physiol* 127: 1466–1475.
- Qiao J, Ishihara Y, Kuroda H, Sakai F, Sakai H, et al. (1997) Transient expression of goat growth hormone gene in poplar (*Populus alba* L.) protoplasts: a quick method for detection of foreign gene expression in mRNA level. *Biosci Biotechnol Biochem* 61: 1580–1581.
- Russell JA, Mccown BH (1986) Techniques for Enhanced Release of Leaf Protoplasts in Populus. *Plant Cell Reports* 5: 284–287.
- Tang RJ, Liu H, Bao Y, Lv QD, Yang L, et al. (2010) The woody plant poplar has a functionally conserved salt overly sensitive pathway in response to salinity stress. *Plant Mol Biol* 74: 367–380.
- Pallardy SG, Kozlowski TT (1980) Cuticle development in the stomatal region of *Populus* clones. *New Phytologist* 85: 363–368.
- Kang BG, Osburn L, Kopsell D, Tuskan GA, Cheng ZM (2009) Micro-propagation of *Populus trichocarpa* 'Nisqually-1': the genotype deriving the *Populus* reference genome. *Plant Cell Tissue and Organ Culture* 99: 251–257.

(Oak Ridge National Laboratory) for critical reading of the manuscript, and ABRC for distributing some of the plasmids used in this paper. We want to thank Lee Gunter, Sara Jawdy and Zackary Moore for their technical support. We also want to thank the anonymous reviewers for their constructive comments.

The submitted manuscript has been authored by a contractor of the U.S. Government under contract DE-AC05-00OR22725. Accordingly, the U.S. Government retains a nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or allow others to do so, for U.S. Government purposes.

Author Contributions

Conceived and designed the experiments: JG J-GC. Performed the experiments: JG JLM JLL. Analyzed the data: JG J-GC JLM JLL WM UCK GAT. Wrote the paper: JG J-GC JLM JLL WM UCK GAT.

- Nelson BK, Cai X, Nebenfuhr A (2007) A multicolored set of in vivo organelle markers for co-localization studies in Arabidopsis and other plants. *Plant J* 51: 1126–1136.
- Guo J, Wang S, Valerius O, Hall H, Zeng Q, et al. (2011) Involvement of Arabidopsis RACK1 in protein translation and its regulation by abscisic acid. *Plant Physiol* 155: 370–383.
- Citovsky V, Lee LY, Vyas S, Glick E, Chen MH, et al. (2006) Subcellular localization of interacting proteins by bimolecular fluorescence complementation in planta. *J Mol Biol* 362: 1120–1131.
- Fujii H, Chinnusamy V, Rodrigues A, Rubio S, Antoni R, et al. (2009) In vitro reconstitution of an abscisic acid signalling pathway. *Nature* 462: 660–664.
- Hwang I, Sheen J (2001) Two-component circuitry in Arabidopsis cytokinin signal transduction. *Nature* 413: 383–389.
- Wang S, Tiwari SB, Hagen G, Guilfoyle TJ (2005) AUXIN RESPONSE FACTOR7 restores the expression of auxin-responsive genes in mutant Arabidopsis leaf mesophyll protoplasts. *Plant Cell* 17: 1979–1993.
- Yanagisawa S, Yoo SD, Sheen J (2003) Differential regulation of EIN3 stability by glucose and ethylene signalling in plants. *Nature* 425: 521–525.
- Ohashi-Ito K, Fukuda H (2010) Transcriptional regulation of vascular cell fates. *Curr Opin Plant Biol* 13: 670–676.
- Zentella R, Zhang ZL, Park M, Thomas SG, Endo A, et al. (2007) Global analysis of DELLA direct targets in early gibberellin signaling in Arabidopsis. *Plant Cell* 19: 3037–3057.
- Ramirez-Carvajal GA, Morse AM, Davis JM (2008) Transcript profiles of the cytokinin response regulator gene family in *Populus* imply diverse roles in plant development. *New Phytologist* 177: 77–89.
- Love J, Bjorklund S, Vahala J, Hertzberg M, Kangasjarvi J, et al. (2009) Ethylene is an endogenous stimulator of cell division in the cambial meristem of *Populus*. *Proceedings of the National Academy of Sciences of the United States of America* 106: 5984–5989.
- Uggle C, Mellerowicz EJ, Sundberg B (1998) Indole-3-acetic acid controls cambial growth in Scots pine by positional signaling. *Plant Physiology* 117: 113–121.
- Bjorklund S, Antti H, Uddestrand I, Moritz T, Sundberg B (2007) Cross-talk between gibberellin and auxin in development of *Populus* wood: gibberellin stimulates polar auxin transport and has a common transcriptome with auxin. *Plant Journal* 52: 499–511.
- Nieminen K, Immanen J, Laxell M, Kauppinen L, Tarkowski P, et al. (2008) Cytokinin signaling regulates cambial development in poplar. *Proceedings of the National Academy of Sciences of the United States of America* 105: 20032–20037.
- Baena-Gonzalez E, Rolland F, Thevelein JM, Sheen J (2007) A central integrator of transcription networks in plant stress and energy signalling. *Nature* 448: 938–942.
- Turner S, Gallois P, Brown D (2007) Tracheary element differentiation. *Annual Review of Plant Biology* 58: 407–433.
- Fukuda H (1997) Tracheary element differentiation. *Plant Cell* 9: 1147–1156.
- Zhong R, Demura T, Ye ZH (2006) SND1, a NAC domain transcription factor, is a key regulator of secondary wall synthesis in fibers of Arabidopsis. *Plant Cell* 18: 3158–3170.
- Rozen S, Skaletsky H (2000) Primer3 for general users and for biologist programmers. *Methods Mol Biol* 132: 365–386.
- Norris SR, Meyer SE, Callis J (1993) The intron of Arabidopsis thaliana polyubiquitin genes is conserved in location and is a quantitative determinant of chimeric gene expression. *Plant Molecular Biology* 21: 895–906.
- Goodstein DM, Shu S, Howson R, Neupane R, Hayes RD, et al. (2012) Phytozome: a comparative platform for green plant genomics. *Nucleic Acids Res* 40: D1178–1186.