

**Highly sensitive and fast responsive ratiometric fluorescent probe for Cu<sup>2+</sup> based on a naphthalimide-rhodamine dyad and its application in living cell imaging**

Jun Tang<sup>a</sup>, Saige Ma<sup>a</sup>, Di Zhang<sup>b,\*</sup>, Yaqi Liu<sup>a</sup>, Yufen Zhao<sup>a,c</sup>, Yong Ye<sup>a,\*</sup>

<sup>a</sup> *Phosphorus Chemical Engineering Research Center of Henan Province, the College of Chemistry and Molecular Engineering, Zhengzhou University, Zhengzhou, China.*

<sup>b</sup> *Institute of Agricultural Quality Standards and Testing Technology, Henan Academy of Agricultural Sciences, Zhengzhou 450002, China.*

<sup>c</sup> *Department of Chemistry, The College of Chemistry and Chemical Engineering, The Key Laboratory for Chemical Biology of Fujian Province, Xiamen University, Xiamen, 361005, PR China*

---

\* Corresponding author; Email: [yeyong03@tsinghua.org.cn](mailto:yeyong03@tsinghua.org.cn) (Yong Ye), [pandy811@163.com](mailto:pandy811@163.com) (Di Zhang)

**Abstract:** Based on fluorescent resonance energy transfer (FRET), we developed a ratiometric fluorescent probe **1** for sensing Cu<sup>2+</sup>. As expected, probe **1** exhibited high selectivity and excellent sensitivity in both absorbance and fluorescence detection of Cu<sup>2+</sup> in aqueous solution. A significant color change from pale yellow to pink could be observed, enabling naked-eye detection of Cu<sup>2+</sup>. Furthermore, fluorescence imaging experiment of Cu<sup>2+</sup> in living MGC-803 cells demonstrated its value of practical applications in biological systems.

**Keywords:** Fluorescence, Ratiometric, FRET, Copper, Cell imaging, naphthalimide-rhodamine

## 1. Introduction

The determination of metal ions has received tremendous attention in the last two decades, because the metal ions play a significant role in the environment, biology and chemistry [1-3]. As the third-most abundant transition metal in human body and many living organisms, copper plays vital roles [4-5]. However, excess copper in the neuronal cytoplasm can lead many diseases, such as Parkinson's disease, Alzheimer's disease and Wilson disease [6-7]. Therefore, the detection of copper ion content in the physiological environment is very important.

Czarnik et al reported the rhodamine-B derivative and its ring-open reaction [8]. After then, many  $\text{Cu}^{2+}$  probes based on fluorescence intensity were synthesized [4, 9-16]. Although turn-on probes are more sensitive, a major limitation is that variations in the sample environment (pH, polarity, temperature, and so forth) might influence the fluorescence intensity measurements, due to the lack of background signal [17]. The ratiometric chemosensors can provide built-in correction for environmental factors by measuring the changes of emission intensities ratio at two different wavelengths followed by calculation of their intensity ratio [18-20]. From this point of view, fluorescence resonance energy transfer (FRET) based sensors were the most valuable [21-24]. To the best of our knowledge, only a few FRET probes for detecting  $\text{Cu}^{2+}$  based on rhodamine have been reported [25-30].

### Scheme 1 Three kinds of rhodamine-based FRET probe

According to the reported rhodamine-based FRET probe, we found they can be classified into three types (Scheme 1. Type A-C). Most of reported probe were Type A which the energy donor is linked to the interaction site of rhodamine [25-28]. This may induce cleavage of the FRET dyad and are not suitable for analytes. There are several Type B probes were reported. An energy donor is selectively incorporated into

the 4/5-position carboxylic acid group through an appropriate linker to form a Type B FRET platform. However, 4/5-position mixture isomers of FRET dyads are very difficult to separate by standard column chromatography [30]. For a FRET system, a substantial spectral overlap which is necessary between the donor emission band and the acceptor absorption band is closely related to the energy transfer efficiency. The naphthalimide derivatives have broad absorption spectra (450-650 nm) [31-33], which covers a part of absorption of rhodamine (500-560 nm) and fulfills a favorable condition for FRET (Fig. S1). In this paper, we construct a type C probe for sensing  $\text{Cu}^{2+}$  using piperazine to link the energy donor. Compared with Type A and Type B, this probe has the following advantages: 1) The interaction site is far away from the energy donor. 2) The strategy is suitable for target analytes that otherwise could induce the cleavage of the FRET dyad. 3) This approach only yields a single FRET dyad. The FRET energy transfer efficiency ( $E$ ) was calculated to be 82.1% as  $E=1-F_{\text{DA}}/F_{\text{D}}$ . Where,  $F_{\text{DA}}$  and  $F_{\text{D}}$  denote the donor fluorescence intensity with and without an acceptor, respectively (Fig. S2) [34-35]. The synthetic route of type C probe was shown in Scheme 2. Probe **1** exhibited excellent sensitivity and selectivity with a detection limit of 18.6 nM. It also has the shortest response time (< 1 min) for  $\text{Cu}^{2+}$  in ( $\text{CH}_3\text{CN}/\text{H}_2\text{O} = 1/9$ , v/v, 1 mM Tris-HCl buffer, pH = 7.40). Further, fluorescence imaging experiments of  $\text{Cu}^{2+}$  in living MGC-803 cells demonstrated its value of practical application in biological systems.

## 2. Experimental

### 2.1. Apparatus

Fluorescence spectra measurements were performed on a HITACHI F-7000 fluorescence spectrophotometer, and the excitation and emission wavelength band passes were both set at 10.0 nm. Absorption spectra were measured on a Lambda 35 UV/VIS spectrometer, Perkin Elmer precisely. NMR spectra were measured on a Bruker DTX-400 spectrometer in  $\text{CDCl}_3$  with TMS as internal standard. Mass spectral determination was carried on a HPLC Q-ToF HR-MS spectrometer (Waters Micromass) by using methanol as mobile phase.

## 2.2. Materials

All chemicals and reagents were used as received from commercial sources without further purification. Solvents for chemical synthesis and analysis were purified according to standard procedures. Double distilled water was used throughout the experiment. Chloride salts of metal ions ( $\text{Li}^+$ ,  $\text{K}^+$ ,  $\text{Na}^+$ ,  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ ,  $\text{Ba}^{2+}$ ,  $\text{Zn}^{2+}$ ,  $\text{Fe}^{2+}$ ,  $\text{Fe}^{3+}$ ,  $\text{Al}^{3+}$ ,  $\text{Mn}^{2+}$ ,  $\text{Cu}^{2+}$ ,  $\text{Co}^{2+}$ ,  $\text{Ni}^{2+}$ ,  $\text{Cd}^{2+}$ ,  $\text{Cr}^{3+}$ ) and the nitrate salt of  $\text{Pb}^{2+}$  and  $\text{Ag}^+$  were used to evaluate the metal ion binding properties by synthesized compounds. The metal ions were prepared as 10.00 mmol/L in water solution.

## 2.3. Synthesis

**Part 1** and **part 2** was synthesized by reported methods (the detailed synthetic routes were shown in Supporting information) [29, 36].

To a solution of **part 2** (0.2 g, 0.43 mmol) and triethylamine (5 mL) in anhydrous  $\text{CH}_2\text{Cl}_2$  (10 mL), **part 1** (0.15 g, 0.36 mmol) mixed with anhydrous  $\text{CH}_2\text{Cl}_2$  (10 mL) were added dropwise at 0 °C for 30 min. After then the resulting mixture was stirred at room temperature for 4 h. After removed of solvent, the residue was purified by silica gel column chromatography ( $\text{CH}_3\text{OH}:\text{CH}_2\text{Cl}_2 = 1:10$ , v/v) to afford the pure product 0.13 g (42 %) of **1** as a yellow foamy solid. Mp: 122-125 °C.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz, ppm): 1.18 (t, 6 H,  $J = 7.0$  Hz), 3.25 (s, 4 H), 3.36 (q, 4 H,  $J = 6.8$  Hz), 3.64-3.67 (m, 4 H), 3.70 (s, 4 H), 3.79 (s, 4 H), 5.5 (s, 2 H,  $\text{NH}_2$ ), 6.31-6.34 (m, 1 H), 6.56 (d, 2H,  $J = 1.2$  Hz), 6.66-6.69 (m, 4H), 7.09-7.11 (m, 1H), 7.26-7.32 (m, 3H), 7.47-7.49 (m, 2 H), 7.79 (t, 1 H,  $J = 7.8$  Hz), 7.95-7.97 (m, 1 H), 8.26 (d, 1 H,  $J = 7.6$  Hz), 8.64-8.66 (m, 1 H);  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ , 100 MHz, ppm): 12.58, 44.39, 48.74, 65.66, 66.79, 97.96, 102.97, 108.32, 109.68, 112.09, 114.21, 121.88, 123.11, 123.79, 124.50, 124.62, 127.06, 128.04, 128.12, 128.37, 129.40, 129.44, 129.86, 131.57, 131.74, 132.68, 134.51, 148.35, 148.41, 149.00, 151.27, 151.85, 153.51, 153.60, 160.98, 166.28, 170.95; HR-MS:  $\text{C}_{51}\text{H}_{47}\text{N}_7\text{O}_6$   $[\text{M}-\text{H}]^-$ , calcd for 852.2546, found 852.2543.

### Scheme 2. Synthetic route of **1**

## 3. Results and discussion

Fluorescence and UV-vis studies were performed using 10  $\mu\text{M}$  solution of **1** in  $\text{CH}_3\text{CN}/\text{H}_2\text{O}$  (1/9, v/v, 1 mM Tris-HCl buffer, pH = 7.40) with 10 equiv. amounts of metal ions. Solutions were stayed for 10 minutes before measuring the absorption and fluorescence in order to make the metal ions chelate with the sensors sufficiently.

Compound **1** was pale yellow in aqueous solution due to the naphthalimide donor moiety and found very stable in the above-mentioned solution system for more than one week. Upon addition of  $\text{Cu}^{2+}$ , a significant color change from pale yellow to pink could be observed, enabling colorimetric and naked-eye detection of  $\text{Cu}^{2+}$ .

Based on our work in rhodamine-based  $\text{Cu}^{2+}$  probe [11], we know that organic solvents have great effect on the coordination reaction. When the coordination reaction was performed in acetonitrile–water solution, high  $F/F_0$  and  $A$  values were obtained, indicating that acetonitrile–water media is favorable for fluorescent measurement. According to the reported [11, 37], it was found that the fluorescence signal can reach maximum value at 30% aqueous acetonitrile for rhodamine-based  $\text{Cu}^{2+}$  probe. Taking into account its application in cell imaging, we choose 10% aqueous acetonitrile in Tris-HCl buffer for fluorescent assay.

### 3.1. UV-vis spectra

An important feature of **1** was its high selectivity toward the  $\text{Cu}^{2+}$  over the other competitive species. Changes of UV-vis spectra of **1** caused by  $\text{Cu}^{2+}$  and other metal ions including  $\text{Li}^+$ ,  $\text{K}^+$ ,  $\text{Na}^+$ ,  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ ,  $\text{Ba}^{2+}$ ,  $\text{Zn}^{2+}$ ,  $\text{Fe}^{2+}$ ,  $\text{Fe}^{3+}$ ,  $\text{Al}^{3+}$ ,  $\text{Mn}^{2+}$ ,  $\text{Co}^{2+}$ ,  $\text{Ni}^{2+}$ ,  $\text{Cd}^{2+}$ ,  $\text{Pb}^{2+}$ ,  $\text{Ag}^+$  and  $\text{Cr}^{3+}$  in  $\text{CH}_3\text{CN}/\text{H}_2\text{O}$  solution (1/9, v/v, 1 mM Tris-HCl buffer, pH = 7.40) were recorded in Fig. 1. All the competitive cations did not lead to any significant absorption changes in the visible region.

**Fig. 1** UV-vis absorbance spectra of **1** (10  $\mu\text{M}$ ) in the absence and presence of 10 equiv. different metal ions in  $\text{CH}_3\text{CN}/\text{H}_2\text{O}$  (1/9, v/v, 1 mM Tris-HCl buffer, pH = 7.40) solution. Inset shows the photo **1** with different metal ions.

As shown in Fig. S14, an increase in the absorption intensity at 550 nm emerged soon after the addition of  $\text{Cu}^{2+}$ , which corresponded to the absorption of rhodamine. The absorption increased gradually after the addition of  $\text{Cu}^{2+}$  and a significant color change from pale yellow to pink could be observed easily by eyes, indicating that the addition of  $\text{Cu}^{2+}$  can promote the ring-opened reaction of the compound **1**.

To determine the stoichiometry of the  $\text{Cu}^{2+}$ -ligand complex, Job's method for absorbance measurement was applied. As shown in Fig. 2, the UV-vis titrations reached a maximum when the ratio of  $[\text{Cu}^{2+}]/\{[\text{Cu}^{2+}]+[\mathbf{1}]\}$  was 0.5, indicating a 1:1 binding stoichiometry between  $\text{Cu}^{2+}$  and **1**.

**Fig. 2** Job's plots of the complexation between **1** and  $\text{Cu}^{2+}$ .  $[\text{Cu}^{2+}]+[\mathbf{1}] = 20 \mu\text{M}$ .

**Fig. 3** Fluorescence spectra of **1** (10  $\mu\text{M}$ ) in  $\text{CH}_3\text{CN}/\text{H}_2\text{O}$  (1/9, v/v, 1 mM Tris-HCl buffer, pH = 7.40) solution with the presence of 10 equiv. of various species ( $\lambda_{\text{ex}} = 420 \text{ nm}$ , slit = 10 nm).

### 3.2. Fluorescence spectral

The selectivity of **1** for  $\text{Cu}^{2+}$  was further observed in the fluorescent spectra. As shown in Fig. 3, **1** alone exhibited a strongly emission of the naphthalimide centered about 540 nm when excited at 420 nm. When 10 eq metal ions such as  $\text{Li}^+$ ,  $\text{K}^+$ ,  $\text{Na}^+$ ,  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ ,  $\text{Ba}^{2+}$ ,  $\text{Zn}^{2+}$ ,  $\text{Fe}^{2+}$ ,  $\text{Fe}^{3+}$ ,  $\text{Al}^{3+}$ ,  $\text{Mn}^{2+}$ ,  $\text{Co}^{2+}$ ,  $\text{Ni}^{2+}$ ,  $\text{Cd}^{2+}$ ,  $\text{Pb}^{2+}$ ,  $\text{Ag}^+$ , and  $\text{Cr}^{3+}$  were added, no obvious changes of fluorescence intense could be observed. However, upon addition of  $\text{Cu}^{2+}$  under the same condition, a new enhancement of the fluorescence emission at 570 nm appeared quickly. These changes could be ascribed to the  $\text{Cu}^{2+}$  induced opening of the spirocyclic ring of the rhodamine moiety that facilitated the acceptance of fluorescence resonance energy transfer from the naphthalimide donor.

Moreover, the competitive metal ions binding studies shown in Fig. 4 confirmed that background metal ions showed very low interference with the detection of  $\text{Cu}^{2+}$  in  $\text{CH}_3\text{CN}/\text{H}_2\text{O}$  solution (1/9, v/v, 1 mM Tris-HCl buffer, pH = 7.40). These results also demonstrated that compound **1** was a selectivity sensor for  $\text{Cu}^{2+}$  over various other metal ions. Also, it was investigated that the fluorescence response of compound **1** toward  $\text{CuCl}_2$  in the presence of various coexistent anions such as  $\text{Br}^-$ ,  $\text{NO}_3^-$ ,  $\text{SO}_4^{2-}$ ,  $\text{OAc}^-$ ,  $\text{HCO}_3^-$  and  $\text{H}_2\text{PO}_4^-$ . It is gratifying to note that all the tested anions have no interference with the fluorescence response of compound **1** toward  $\text{Cu}^{2+}$  (Fig. 5).

**Fig. 4** Metal-ion selectivity of **1** in  $\text{CH}_3\text{CN}/\text{H}_2\text{O}$  (1/9, v/v, 1 mM Tris-HCl buffer, pH = 7.40) solution. The black bars represent the fluorescence emission ratio ( $F_{570}/F_{540}$ ) of a solution of **1** (10  $\mu\text{M}$ ) and 10 equiv. of other metal ions. The red bars show the  $F_{570}/F_{540}$  ratio after the addition of 10 equiv. of  $\text{Cu}^{2+}$  to the solution containing **1** (10  $\mu\text{M}$ ) and different metal ions (100  $\mu\text{M}$ ) ( $\lambda_{\text{ex}} = 420 \text{ nm}$ , slit = 10 nm).

**Fig. 5** The ratiometric fluorescence responses ( $F_{570}/F_{540}$ ) of **1** (10  $\mu\text{M}$ ) upon the addition of 100  $\mu\text{M}$   $\text{CuCl}_2$  in the presence of 100  $\mu\text{M}$  background anions in  $\text{CH}_3\text{CN}/\text{H}_2\text{O}$  (1/9, v/v, 1 mM Tris-HCl buffer, pH = 7.40) solution ( $\lambda_{\text{ex}} = 420 \text{ nm}$ , slit = 10 nm).

In order to investigate the influence of the different acid concentration on the spectra of **1** and find a suitable pH span in which **1** can selectively detect  $\text{Cu}^{2+}$  efficiently, the acid titration experiments were performed. As shown in Fig. 6, the fluorescent titration curve of free sensor **1** in  $\text{CH}_3\text{CN}/\text{H}_2\text{O}$  (1/9, v/v, 1 mM Tris-HCl buffer) solution did not show obvious changes of the fluorescence intensity ratio at ( $F_{570}/F_{540}$ ) between pH 1.0 and 10.0, suggesting that spiro lactam tautomer of **1** was insensitive to the pH changes in this range. However, the addition of  $\text{Cu}^{2+}$  led to the fluorescence intensity ratio enhancement over a comparatively wide pH range (4.0-10.0), which is attributed to opening of the rhodamine ring. Consequently, **1** might be used to detect  $\text{Cu}^{2+}$  in approximate physiological conditions.

**Fig. 6** The ratiometric fluorescence responses ( $F_{570}/F_{540}$ ) of free **1** (10  $\mu\text{M}$ ) and in the presence of 10 equiv.  $\text{Cu}^{2+}$  in  $\text{CH}_3\text{CN}/\text{Tris-HCl}$  (1/9, v/v, 1 mM) solution with different pH conditions ( $\lambda_{\text{ex}} = 420$  nm, slit = 10 nm).

Moreover, a time course of the fluorescence response of **1** upon addition of  $\text{Cu}^{2+}$  was shown in Fig. 7. The kinetics of fluorescence enhancement at 570 nm by the new developed ratiometric fluorescent probe was recorded, and results indicated that the recognizing event is very quickly (< 1 min) and could complete in 5 min. That means compound **1** can be used as a ratiometric fluorescence probe for the fast detection of  $\text{Cu}^{2+}$ .

**Fig. 7** Kinetics of the ratiometric fluorescence responses ( $F_{570}/F_{540}$ ) of free **1** (10  $\mu\text{M}$ ) and in the presence of 10 equiv.  $\text{Cu}^{2+}$  in  $\text{CH}_3\text{CN}/\text{H}_2\text{O}$  (1/9, v/v, 1 mM Tris-HCl buffer, pH = 7.40) solution ( $\lambda_{\text{ex}} = 420$  nm, slit = 10 nm).

We also have conducted the titration experiments with the addition of  $\text{Cu}^{2+}$  for probe **1** (Fig. S15). Upon addition gradual increase of  $\text{Cu}^{2+}$  to the solution of **1**, the emission at 570 nm was appeared and increased remarkably. The emission intensities ratio at 570 nm and 540 nm ( $F_{570}/F_{540}$ ) exhibited a change from 0.7 to 2, which showed that the FRET process occurred. One of the most important and useful application for a fluorescence sensor is the detection limit of metal ions. The



fluorescence intensity ratio was found to increase linearly with the  $\text{Cu}^{2+}$  concentration in the range of 0-16  $\mu\text{M}$  (Fig. 8) and the detection limit of  $\text{Cu}^{2+}$  was measured to be 18.6 nM [38, 39], which were far lower than the WHO recommended value for  $\text{Cu}^{2+}$  (2.0 mg/L) and the U.S. Environment Protection Agency (EPA) set the maximum allowable level of copper ( $\sim 20 \mu\text{M}$ ) in drinking water [40, 41].

**Fig. 8** Fluorescence intensity ratio changes ( $F_{570}/F_{540}$ ) of **1** (10  $\mu\text{M}$ ) upon gradual addition of  $\text{Cu}^{2+}$  in  $\text{CH}_3\text{CN}/\text{H}_2\text{O}$  (1/9, v/v, 1 mM Tris-HCl buffer, pH = 7.40) solution ( $\lambda_{\text{ex}} = 420 \text{ nm}$ , slit = 10 nm).

### 3.3. Mechanism

Further, it was of great interest to investigate the reversible binding nature of the sensor (Fig. 9). Upon addition of 10 equiv. EDTA to the solution of probe **1** (10  $\mu\text{M}$ ) with  $\text{Cu}^{2+}$  (10 equiv.), the fluorescence intensity at 570 nm was not quenched (blue line). These indicated that the coordination of probe **1** with  $\text{Cu}^{2+}$  is chemically irreversible.

**Fig. 9** Fluorescence intensity of **1** (10  $\mu\text{M}$ ) to  $\text{Cu}^{2+}$  in  $\text{CH}_3\text{CN}/\text{H}_2\text{O}$  (1/9, v/v, 1 mM Tris-HCl buffer, pH = 7.40) solution ( $\lambda_{\text{ex}} = 420 \text{ nm}$ , slit = 10 nm).

To further investigate the binding stoichiometry of **1** and  $\text{Cu}^{2+}$  ions, the form of **1**- $\text{Cu}^{2+}$  complex was also confirmed by ESI-MS analysis (Fig. S16). When  $\text{CuCl}_2$  was added to **1** in  $\text{CH}_3\text{CN}$ , the peak of  $[\mathbf{1}+\text{H}]^+$  (854.58) was disappeared, in the same time, a new peak 840.65 was appeared, which was suggested the compound  $[\mathbf{2}+\text{H}]^+$  (calcd = 840.34). Herein, according to our knowledge [29, 42-45], we broached a conceivable mechanism of  $\text{Cu}^{2+}$  complex with **1** (Scheme 3).

**Scheme 3.** Possible sensing mechanism of **1** with  $\text{Cu}^{2+}$ .

## 4. Bioimaging application of compound **1** in MGC-830 Cells [46, 47]

Bioimaging applications of compound **1** for monitoring of  $\text{Cu}^{2+}$  ions in living cells were then carried out. MGC-803 cells were used for this experiment. MGC-803 cells were incubated with **1** (10  $\mu\text{M}$ ) in RPMI1640 medium (containing 10% fetal

bovine serum) for about 30 min at 37 °C showed a strong green intracellular fluorescence (Fig. 10b) and a weak red one (Fig. 10c). After then, 20  $\mu\text{M}$  of  $\text{Cu}^{2+}$  were then supplemented to the cells, and all the reaction mixtures were incubated at 37 °C for another 30 min, a significant increase in the red fluorescence from the intracellular area was observed (Fig. 10f). A bright field transmission image of cells with  $\text{Cu}^{2+}$  and **1** confirmed that the cells were viable throughout the imaging experiments (Fig. 10a and d). These results clearly indicated that **1** might be used for ratiometric imaging of  $\text{Cu}^{2+}$  in biological samples.

**Fig. 10** Bright field and fluorescence microscopic images of MGC-803 cells. MGC-803 cells incubated by compound **1** (10  $\mu\text{M}$ ) observed under bright field (a), green channel (b), red channel (c), then further incubation with  $\text{Cu}^{2+}$  (20  $\mu\text{M}$ ) for 30 min at 37 °C under bright field (d), green channel (e), red channel (f).

## 5. Conclusion

In summary, we reported a novel FRET based fluorescent probe **1** for the ratiometric detection of  $\text{Cu}^{2+}$  in aqueous solution. It exhibited high sensitivity and selectivity toward  $\text{Cu}^{2+}$  and could be conveniently detected even by naked eye. Probe **1** was suitable in a broad pH range 4-10 and allowed the ratiometric detection of intracellular  $\text{Cu}^{2+}$  levels in live MGC-803 cells. Moreover, probe **1** exhibited a low detection limit (18.6 nM) and a rapid response time (< 1 min) for  $\text{Cu}^{2+}$ .

## Acknowledgements

This work was financially supported by the National Science Foundation of China (No. 21572209) and Science-Technology Foundation for Outstanding Young Scientists of Henan Academy of Agricultural Sciences (Grant no. 2016YQ22).

## References

- [1] J. S. Wu, I. C. Hwang, K. S. Kim, J. S. Kim, Rhodamine-based  $\text{Hg}^{2+}$ -selective chemodosimeter in aqueous solution: fluorescent off-on, *Org. Lett.* 9 (2007) 907-910.
- [2] S. Kabehie, M. Xue, A. Z. Stieg, M. Liong, K. L. Wang, J. I. Zink, Heteroleptic copper switches, *J. Am. Chem. Soc.* 132 (2010) 15987-15996.
- [3] J. F. Zhang, Y. Zhou, J. Yoon, J. S. Kim, Recent progress in fluorescent and colorimetric chemosensors for detection of precious metal ions (silver, gold and platinum ions), *Chem. Soc. Rev.* 40 (2011) 3416-3429.
- [4] E. L. Que, D. W. Domaille, C. J. Chang, Metals in neurobiology: probing their chemistry and biology with molecular imaging, *Chem. Rev.* 108 (2008) 1517-1549.
- [5] T. Francesco, M. Cristina, P. Marina, P. Maura, S. Carlo, Copper in diseases and treatments, and copper-based anticancer strategies, *Med. Res. Rev.* 30 (2010) 708-749.
- [6] J. C. Lee, H. B. Gray, J. R. Winkler, Copper(II) binding to  $\alpha$ -Synuclein, the Parkinson's protein, *J. Am. Chem. Soc.* 130 (2008) 6898-6899.
- [7] W. Chouyyok, Y. Shin, J. Davidson, W. D. Samuels, N. H. Lafemina, R.D. Rutledge, G.E. Fryxell, T. Sangvanich, W. Yantasee, Selective removal of copper(II) from natural waters by nanoporous sorbents functionalized with chelating diamines, *Environ. Sci. Technol.* 44 (2010) 6390-6395.
- [8] V. Dujols, F. Ford, A. W. Czarnik, A long-wavelength fluorescent chemodosimeter selective for Cu(II) ion in water, *J. Am. Chem. Soc.* 119 (1997) 7386-7387.
- [9] K. M. K. Swamy, S. K. Ko, S. K. Kwon, H. N. Lee, C. Mao, J. M. Kim, K. H. Lee, J. Kim, I. Shin, J. Yoon, Boronic acid-linked fluorescent and colorimetric probes for copper ions, *Chem. Commun.* 45 (2008) 5915-5917.
- [10] Y. B. Ruan, C. Li, J. Tang, J. Xie, Highly sensitive naked-eye and fluorescence "turn-on" detection of  $\text{Cu}^{2+}$  using Fenton reaction assisted signal amplification, *Chem. Commun.* 46 (2010) 9220-9222.
- [11] D. Zhang, M. Wang, M. M, Chai, X. P. Chen, Y. Ye, Y. F. Zhao, Three highly sensitive and selective colorimetric and off-on fluorescent chemosensors for  $\text{Cu}^{2+}$  in aqueous solution, *Sensor. Actuat. B-Chem.* 168 (2012) 200-206.
- [12] M. Wang, D. Zhang, M. Li, M. Fan, Y. Ye, Y. F. Zhao, A rhodamine-cyclen conjugate as chromogenic and fluorescent chemosensor for copper ion in aqueous

media, *J. Fluoresc.* 23 (2013) 417-423.

[13] W. Lin, L. Yuan, W. Tan, J. Feng, L. Long, Construction of fluorescent probes via protection/deprotection of functional groups: a ratiometric fluorescent probe for  $\text{Cu}^{2+}$ , *Chem. Eur. J.* 15 (2009) 1030-1035.

[14] L. Yuan, W. Lin, Z. Cao, L. Long and J. Song, Photocontrollable analyte-responsive fluorescent probes: a photocaged copper-responsive fluorescence turn-on probe. *Chem. Eur. J.* 17 (2011) 689-696.

[15] L. Yuan, W. Lin, B. Chen and Y. Xie, Development of FRET-based ratiometric fluorescent  $\text{Cu}^{2+}$  chemodosimeters and the applications for living cell imaging. *Org. Lett.* 14 (2012) 432-435.

[16] M. Ren, B. Deng, J. Y. Wang, Z. R. Liu and W. Lin. A dual-emission fluorescence-enhanced probe for imaging copper(II) ions in lysosomes. *J. Mater. Chem. B* 3 (2015) 6746-6752.

[17] J. L. Fan, P. Zhan, M. M. Hu, W. Sun, J. Z. Tang, J. Y. Wang, S. G. Sun, F.L. Song, X. J. Peng, A fluorescent ratiometric chemodosimeter for  $\text{Cu}^{2+}$  based on TBET and its application in living cells, *Org. Lett.* 15 (2013) 492-495.

[18] A. Coskun, E. U. Akkaya, Signal ratio amplification via modulation of resonance energy transfer: proof of principle in an emission ratiometric Hg(II) sensor, *J. Am. Chem. Soc.* 128 (2006) 14474-14475.

[19] W. Lin, L. Yuan, L. L. Long, C. C. Guo, J. B. Feng, A fluorescent cobalt probe with a large ratiometric fluorescence response via modulation of energy acceptor molar absorptivity on metal ion binding, *Adv. Funct. Mater.* 18 (2008) 2366-2372.

[20] C. C. Wang, D. Zhang, X. Y. Huang, P. G. Ding, Z. J. Wang, Y. F. Zhao, Y. Ye, A ratiometric fluorescent chemosensor for  $\text{Hg}^{2+}$  based on FRET and its application in living cells, *Sensor. Actuat. B-Chem.* 198 (2014) 33-40.

[21] L. Yuan, W. Lin, Y. Xie, B. Chen and J. Song. Development of a ratiometric fluorescent sensor for ratiometric imaging of endogenously produced nitric oxide in macrophage cells. *Chem. Commun.* 47 (2011) 9372-9374.

[22] L. Yuan, W. Lin, Y. Xie, B. Chen and S. Zhu, Single fluorescent probe responds to  $\text{H}_2\text{O}_2$ , NO, and  $\text{H}_2\text{O}_2/\text{NO}$  with three different sets of fluorescence signals. *J. Am. Chem. Soc.* 134 (2012) 1305-1315.

[23] L. Yuan, W. Lin, Z. Cao, J. Wang and B. Chen, Development of FRET-based dual-excitation ratiometric fluorescent pH probes and their photocaged derivatives. *Chem. Eur. J.* 18 (2012) 1247-1255.

- [24] L. Yuan, W. Lin, K. Zheng and S. Zhu, FRET-based small-molecule fluorescent probes: rational design and bioimaging applications. *Accounts of Chemical Research*. 46 (2013) 1462-1473.
- [25] D. Maity, D. Karthigeyan, T. K. Kundu, T. Govindaraju, FRET-based rational strategy for ratiometric detection of  $\text{Cu}^{2+}$  and live cell imaging, *Sensors and Actuators B* 176 (2013) 831-837.
- [26] C. Yu, Y. Wen, X. Qin, J. Zhang, A fluorescent ratiometric  $\text{Cu}^{2+}$  probe based on FRET by naphthalimide-appended rhodamine derivatives, *Anal. Methods* 6 (2014) 9825-9830.
- [27] Z. Hu, J. Hu, Y. Cui, G. Wang, X. Zhang, K. Uvdal, H.-W. Gao, A facile "click" reaction to fabricate a FRET-based ratiometric fluorescent  $\text{Cu}^{2+}$  probe, *J. Mater. Chem. B* 2 (2014) 4467-4472.
- [28] C. Kar, M. D. Adhikari, A. Ramesh, G. Das, NIR- and FRET-based sensing of  $\text{Cu}^{2+}$  and  $\text{S}^{2-}$  in physiological conditions and in live cells, *Inorg. Chem.* 52 (2013) 743-752.
- [29] X. Guan, W. Lin, W. Huang, Development of a new rhodamine-based FRET platform and its application as a  $\text{Cu}^{2+}$  probe, *Org. Biomol. Chem.* 12 (2014) 3944-3949.
- [30] S. Li, D. Zhang, M. Wang, S. Ma, J. Liu, Y. Zhao, Y. Ye, Synthesis and properties of a novel FRET-based ratiometric fluorescent sensor for  $\text{Cu}^{2+}$ , *J Fluoresc*, 2016, DOI: 10.1007/s10895-016-1768-5.
- [31] B. Dong, X. Song, C. Wang, X. Kong, Y. Tang and W. Lin, Dual site-controlled and Lysosome-targeted intramolecular charge transfer-photoinduced electron transfer-fluorescence resonance energy transfer fluorescent probe for monitoring pH changes in living cells, *Anal. Chem.* 88 (2016) 4085-4091.
- [32] Y. Tang, X. Kong, A. Xu, B. Dong and W. Lin, Development of a two-photon fluorescent probe for imaging of endogenous formaldehyde in living tissues, *Angew. Chem. Int. Ed.* 55 (2016) 3356-3359.
- [33] Q. Xu, L. He, H. Wei, W. Lin, An ICT-based hydrogen sulfide sensor with good water solubility for fluorescence imaging in living cells, *J. Fluoresc.* 2016, DOI 10.1007/s10895-015-1582-5.
- [34] Y. J. Gong, X. B. Zhang, C. C. Zhang, A. L. Luo, T. Fu, W. Tan, G. L. Shen and R. Q. Yu, Through bond energy transfer: a convenient and universal strategy toward

efficient ratiometric fluorescent probe for bioimaging applications, *Anal. Chem.* 84 (2012) 10777-10782.

[35] L. Yuan, W. Lin, Y. Xie, B. Chen and J. Song, Fluorescent detection of hypochlorous acid from turn-on to FRET-based ratiometry by a HOCl-mediated cyclization reaction, *Chem.–Eur. J.* 18 (2012) 2700-2705.

[36] C. Y. Wang, K. M. C Wong, Selective Hg<sup>2+</sup> sensing behaviors of rhodamine derivatives with extended conjugation based on two successive ring-opening processes, *Inorg. Chem.* 52 (2013) 13432-13441.

[37] M. Zhao, X. Yang, S. He, L. Wang, A rhodamine-based chromogenic and fluorescent chemosensor for copper ion in aqueous media, *Sensors and Actuators B* 135 (2009) 625-631

[38] Z. Li, X. H. Li, X. H. Gao, Y. Y. Zhang, W. Shi, H. M. Ma, Nitroreductase detection and hypoxic tumor cell imaging by a designed sensitive and selective fluorescent probe, 7-[(5-nitrofuran-2-yl)methoxy]-3H-phenoxanzin -3-one, *Anal. Chem.* 85 (2013) 3926-3932.

[39] D. Zhang, W. Chen, Z. R. Miao, Y. Ye, Y. F. Zhao, S. B. King, M. Xian, A reductive ligation based fluorescent probe for S-nitrosothiols, *Chem. Commun.* 50 (2014) 4806-4809.

[40] Y. R. Kim, H. J. Kim, J. S. Kim, H. Kim, Rhodamine-based “turn-on” fluorescent chemodosimeter for Cu(II) on Ultrathin Platinum films as molecular switches, *Adv. Mater.* 20 (2008) 4428-4432.

[41] M. Z. Tian, M. M. Hu, J. L. Fan, X. J. Peng, J. Y. Wang, S. G. Sun, R. Zhang, Rhodamine-based ‘turn-on’ fluorescent probe for Cu(II) and its fluorescence imaging in living cells, *Bioorg. Med. Chem. Lett.* 23 (2013) 2916-2919.

[42] Y. Xiang, A. J. Tong, P. Y. Jin, Y. Ju, New fluorescent rhodamine hydrazone chemosensor for Cu(II) with high selectivity and sensitivity, *Org. Lett.* 8 (2006) 2863-2866.

[43] F. J. Huo F, J. Su, Y. Q. Sun, C. X. Yin, H. B. Tong, Z. X. Nie, A rhodamine-based dual chemosensor for the visual detection of copper and the ratiometric fluorescent detection of vanadium, *Dyes Pigment.* 86 (2010) 50-55.

[44] Y. Xiang, Z. F. Li, X. T. Chen, A. J. Tong, Highly sensitive and selective optical chemosensor for determination of Cu<sup>2+</sup> in aqueous solution, *Talanta* 74 (2008) 1148-1153.

[45] X. T. Chen, Z. F. Li, Y. Xiang, A. J. Tong, Salicylaldehyde fluorescein hydrazone:


a colorimetric logic chemosensor for pH and Cu(II), *Tetrahedron Lett.* 49 (2008) 4697-4700.

[46] Y. Zhao, Y. Sun, X. Lv, Y. L. Liu, M. L. Chen, W. Guo, Rhodamine-based chemosensor for Hg<sup>2+</sup> in aqueous solution with a broad pH range and its application in live cell imaging, *Org. Biomol. Chem.* 8 (2010) 4143-4147.


[47] Y. J. Gong, X. B. Zhang, Z. Chen, Y. Yuan, Z. Jin, L. Mei, J. Zhang, W. H. Tan, G. L. Shen, R. Q. Yu, An efficient rhodamine thiospirolactam-based fluorescent probe for detection of Hg<sup>2+</sup> in aqueous samples, *Analyst* 137 (2012) 932-938.









**Fig. 2** Job's plots of the complexation between **1** and  $Cu^{2+}$ .  $[Cu^{2+}]+[1] = 20 \mu M$ .




**Fig. 3** Fluorescence spectra of **1** ( $10 \mu M$ ) in  $CH_3CN/H_2O$  (1/9, v/v, 1 mM Tris-HCl buffer, pH = 7.40) solution with the presence of 10 equiv. of various species ( $\lambda_{ex} = 420 \text{ nm}$ , slit = 10 nm).




**Fig. 4** Metal-ion selectivity of **1** in CH<sub>3</sub>CN/H<sub>2</sub>O (1/9, v/v, 1 mM Tris-HCl buffer, pH = 7.40) solution. The black bars represent the fluorescence emission ratio (F<sub>570</sub>/F<sub>540</sub>) of a solution of **1** (10 μM) and 10 equiv. of other metal ions. The red bars show the F<sub>570</sub>/F<sub>540</sub> ratio after the addition of 10 equiv. of Cu<sup>2+</sup> to the solution containing **1** (10 μM) and different metal ions (100 μM) (λ<sub>ex</sub> = 420 nm, slit = 10 nm).




**Fig. 5** The ratiometric fluorescence responses (F<sub>570</sub>/F<sub>540</sub>) of **1** (10 μM) upon the addition of 100 μM CuCl<sub>2</sub> in the presence of 100 μM background anions in CH<sub>3</sub>CN/H<sub>2</sub>O (1/9, v/v, 1 mM Tris-HCl buffer, pH = 7.40) solution (λ<sub>ex</sub> = 420 nm, slit = 10 nm).




**Fig. 6** The ratiometric fluorescence responses ( $F_{570}/F_{540}$ ) of free **1** ( $10\ \mu\text{M}$ ) and in the presence of 10 equiv.  $\text{Cu}^{2+}$  in  $\text{CH}_3\text{CN}/\text{Tris-HCl}$  (1/9, v/v, 1 mM) solution with different pH conditions ( $\lambda_{\text{ex}} = 420\ \text{nm}$ , slit = 10 nm).




**Fig. 7** Kinetics of the ratiometric fluorescence responses ( $F_{570}/F_{540}$ ) of free **1** ( $10\ \mu\text{M}$ ) and in the presence of 10 equiv.  $\text{Cu}^{2+}$  in  $\text{CH}_3\text{CN}/\text{H}_2\text{O}$  (1/9, v/v, 1 mM Tris-HCl buffer, pH = 7.40) solution ( $\lambda_{\text{ex}} = 420\ \text{nm}$ , slit = 10 nm).




**Fig. 8** Fluorescence intensity ratio changes ( $F_{570}/F_{540}$ ) of **1** ( $10\ \mu\text{M}$ ) upon gradual addition of  $\text{Cu}^{2+}$  in  $\text{CH}_3\text{CN}/\text{H}_2\text{O}$  ( $1/9$ , v/v,  $1\ \text{mM}$  Tris-HCl buffer,  $\text{pH} = 7.40$ ) solution ( $\lambda_{\text{ex}} = 420\ \text{nm}$ , slit =  $10\ \text{nm}$ ).



**Fig. 9** Fluorescence intensity of **1** ( $10\ \mu\text{M}$ ) to  $\text{Cu}^{2+}$  in  $\text{CH}_3\text{CN}/\text{H}_2\text{O}$  ( $1/9$ , v/v,  $1\ \text{mM}$  Tris-HCl buffer,  $\text{pH} = 7.40$ ) solution ( $\lambda_{\text{ex}} = 420\ \text{nm}$ , slit =  $10\ \text{nm}$ ).



**Scheme 3.** Possible sensing mechanism of **1** with  $\text{Cu}^{2+}$ .



**Fig. 10** Bright field and fluorescence microscopic images of MGC-803 cells. MGC-803 cells incubated by compound **1** ( $10\ \mu\text{M}$ ) observed under bright field (a), green channel (b), red channel (c), then further incubation with  $\text{Cu}^{2+}$  ( $20\ \mu\text{M}$ ) for 30 min at  $37\ ^\circ\text{C}$  under bright field (d), green channel (e), red channel (f).