

# **Highly Unsaturated Fatty Acid Synthesis in Atlantic Salmon: Characterisation of ELOVL5- and ELOVL2-like Elongases**

Sofia Morais\*, Oscar Monroig, Xiaozhong Zheng, Michael J. Leaver, Douglas R.

Tocher

*Institute of Aquaculture, University of Stirling, Stirling FK9 4LA, UK*

\*Corresponding author: Dr Sofia Morais, Institute of Aquaculture, University of Stirling, Stirling FK9 4LA, Scotland, United Kingdom. Tel: +44 1786 467993; Fax +44 1786 472133; E-mail: sofia.morais@stir.ac.uk

Keywords: Highly unsaturated fatty acids . Fatty acyl elongase . Genes . cDNA . Atlantic salmon

Running title: Characterisation of salmon ELOVL5- and ELOVL2 elongases

Abbreviations: aa, amino acid; ARA, arachidonic acid (20:4n-6); DHA, docosahexaenoic acid (22:6n-3); EPA, eicosapentaenoic acid (20:5n-3); ER, endoplasmic reticulum; FA, fatty acid; FO, fish oil; HUFA, highly unsaturated fatty acids (carbon chain length  $\geq C_{20}$  with  $\geq 3$  double bonds); LO, linseed oil; ORF, open reading frame; PUFA, polyunsaturated fatty acids; Q-PCR, quantitative (real-time) polymerase chain reaction; RACE, rapid amplification of cDNA ends; RO, rapeseed oil; SO, soybean oil; UTR, untranslated region; VO, vegetable oil.

**Abstract** Fish species vary in their capacity to biosynthesize the n-3 long-chain polyunsaturated fatty acids (LC-PUFA) eicosapentaenoic (EPA) and docosahexaenoic (DHA) acids that are crucial to the health of higher vertebrates. The synthesis of LC-PUFA involves enzyme-mediated fatty acyl desaturation and elongation. Previously, a cDNA for an elongase, now termed *elov15a*, had been cloned from Atlantic salmon. Here we report on the cloning of two new elongase cDNAs: a second *elov15b* elongase, corresponding to a 294 aa protein, and an *elov12*-like elongase, coding for a 287 aa protein, characterized for the first time in a non-mammalian vertebrate. Heterologous expression in yeast showed that the salmon Elov15b elongated C18 and C20 PUFA, with low activity towards C22, while Elov12 elongated C20 and C22 PUFA with lower activity towards C18 PUFA. All three transcripts showed predominant expression in the intestine and liver, followed by the brain. Elongase expression showed differential nutritional regulation. Levels of *elov15b* and particularly of *elov12*, but not of *elov15a*, transcripts were significantly increased in liver of salmon fed vegetable oils (VO) compared to fish fed fish oil (FO). Intestinal expression showed a similar pattern. Phylogenetic comparisons indicate that, in contrast to salmon and zebrafish, Acanthopterygian fish species lack *elov12* which is consistent with their negligible ability to biosynthesize LC-PUFA and to adapt to VO dietary inclusion, compared to predominantly freshwater salmonids. Thus the presence of *elov12* in salmon explains the ability of this species to biosynthesize LC-HUFA and may provide a biotechnological tool to produce enhanced levels of LC-PUFA, particularly DHA, in transgenic organisms.

## Introduction

In vertebrates, biosynthesis of long-chain polyunsaturated fatty acids (LC-PUFA) involves sequential desaturation and elongation of precursor essential PUFA, 18:2n-6 and 18:3n-3. Synthesis of arachidonic acid (ARA; 20:4n-6) is achieved by  $\Delta 6$  desaturation of 18:2n-6 to produce 18:3n-6 that is elongated to 20:3n-6 followed by  $\Delta 5$  desaturation (Cook, 1996). Synthesis of eicosapentaenoic acid (EPA; 20:5n-3) from 18:3n-3 uses the same enzymes and pathway as for ARA, but docosahexaenoic acid (DHA; 22:6n-3) synthesis requires two further elongation steps, a second  $\Delta 6$  desaturation and a chain shortening step (Sprecher, 2000). The extent to which any species can produce LC-PUFA varies and is dependent on their complement of fatty acyl desaturase and elongase enzymes. Freshwater fish and salmonids, including Atlantic salmon (*Salmo salar*), are capable of producing DHA from 18:3n-3 and so must express all the enzyme activities necessary for this biosynthetic pathway (Tocher, 2003).

Interest in LC-PUFA synthesis in fish results from the fact that they are the primary source in the human food basket of the omega-3 or n-3 LC-PUFA, that are crucial to the health of higher vertebrates. Aquaculture now supplies an increasing proportion of the fish for human consumption (Tidwell and Allan, 2002; FAO, 2006) and, until now, formulation of diets with fish oil (FO) has ensured that farmed fish are rich in n-3 LC-PUFA (Bell and Waagbø, 2008). As global FO supplies are at their sustainable limit, further expansion of aquaculture requires suitable alternatives, with vegetable oils (VO) being prime candidates. However, VOs are generally rich in C18 PUFA but lack the n-3 LC-PUFA abundant in FO (Sargent et al., 2002) and thus flesh of fish fed VO is characterised by increased levels of 18:2n-6 and 18:3n-3, and decreased levels of EPA and DHA, compromising their nutritional value to the human consumer (Izquierdo et al., 2003; Regost et al., 2003). Our hypothesis is that understanding the molecular basis of LC-PUFA biosynthesis and regulation in salmon

will allow us to optimise the activity of the pathway to enable efficient and effective use of dietary VOs while maintaining the nutritional quality of the fish. Alimuddin et al. (2008) take this further by suggesting genetic engineering as a strategy to create fish capable of being reared on VO diets while still maintaining their health-promoting composition and, to demonstrate this, were able to increase EPA and DHA contents in transgenic zebrafish by over-expressing a masu salmon *elovl5*-like gene.

In mammals, several fatty acyl elongase genes termed *ELOVL1* to *ELOVL7*, with differing fatty acid (FA) substrate specificities, have been described with *ELOVL2* and *ELOVL5* shown to participate in LC-PUFA biosynthesis (Leonard et al., 2000, 2002, 2004; Jakobsson et al., 2006). Mammalian *ELOVL5* is predominantly involved in the elongation of C18 and C20 PUFA, whereas *ELOVL2* has greatest activity in the elongation of C20 and C22 (Leonard et al., 2000, 2002, 2004). Previously, we cloned and characterised a cDNA for a fatty acyl elongase gene from Atlantic salmon (Hastings et al., 2005). Phylogenetic analysis grouped the salmon elongase cDNA into a cluster with greatest similarity to mammalian *ELOVL5* (Leaver et al., 2008a). Heterologous expression in yeast showed that the product of the salmon elongase cDNA had the ability to lengthen C18 and C20 PUFA with only low activity towards C22 (Hastings et al., 2005). Therefore, the salmon elongase appeared to have a similar substrate specificity to mammalian *ELOVL5*. More recently, a search of the Atlantic salmon EST database ([www.tigr.org](http://www.tigr.org)) showed it contained a second fatty acyl elongase transcript that appeared related to mammalian *ELOVL2* (Leaver et al., 2008a).

The aim of the present study was to isolate and characterise cDNAs for other fatty acyl elongases of Atlantic salmon, particularly for *elovl2*, which has not been previously reported in a non-mammalian vertebrate. Two new cDNAs identified as an *elovl2*-like elongase and a second *elovl5*-like fatty acyl elongase were cloned. Functional characterisation of their encoded polypeptides in yeast, *Saccharomyces cerevisiae*, showed that the salmon Elov12-like

protein elongated C20 and C22 PUFA with only low activity towards C18 PUFA. The second elongase, named *elovl5b*, elongated C18 and C20 PUFA, with residual activity towards C22. The tissue distribution of the three elongase transcripts was determined, and their nutritional regulation in response to changes in the FA composition of the diet was analysed in liver and intestine.

## **Materials and methods**

### cDNA cloning

The sequence of the previously cloned and characterised *elovl5a* cDNA (gb|AY170327|; Hastings et al., 2005) was used to query the GenBank dbEST ([www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov)) and Atlantic salmon Gene Index (<http://compbio.dfci.harvard.edu/tgi/>) databases. This led to the identification of salmon EST sequences (gb|DW546112| and ti|TC91192|) showing similarity to the *elovl5a* cDNA. Primers specific to these sequences were designed and the fragments obtained by polymerase chain reaction (PCR) using GoTaq<sup>®</sup> Colorless Master Mix (Promega, Southampton, U.K.), following manufacturer's instructions, were sequenced (CEQ-8800 Beckman Coulter Inc., Fullerton, U.S.A.) and further extended by 5' and 3' rapid amplification of cDNA ends (RACE) PCR (FirstChoice<sup>®</sup> RLM-RACE kit, Ambion, Applied Biosystems, Warrington, U.K.) to produce full-length cDNAs. These were deposited in the GenBank database under accession numbers gb|FJ237531| and gb|FJ237532| for the cDNAs produced from the gb|DW546112| and ti|TC91192| ESTs, respectively.

### Sequence and phylogenetic analysis

The deduced amino acid (aa) sequences of the two newly cloned elongase cDNAs were aligned with those of salmon (Ss) Elovl5a (NP\_001117039) and of human (Hs) ELOVL5 (NP\_068586) and ELOVL2 (NP\_060240) using ClustalW2 or, to compare sequences two by two, the EMBOSS Pairwise Alignment Algorithms tool (<http://www.ebi.ac.uk/Tools/emboss/align/>) was used. The aa sequences of *elovl* genes from other fish species were derived by searching the ENSEMBL genome database using BLASTX and the protein sequences of human and salmon *elovl5* and *elovl2* as keys. Genomic sequences with similarity were extracted and compared with the protein sequence of salmon *elovl2* and *elovl5* using Wise2 (<http://www.ebi.ac.uk/Tools/Wise2/index.html>) to generate deduced Elovl polypeptides from *Tetraodon nigroviridis*, *Takifugu rubripes*, *Danio rerio*, *Gasterosteus aculeatus* and *Oryzias latipes*. Deduced aa sequences of elongases from various species were aligned using ClustalX and sequence phylogenies were reconstructed using the Neighbour Joining method (Saitou and Nei, 1987). GenBank accession numbers for these sequences are: NP\_956747 and NP\_001035452 for *Danio rerio* (Dr), AAV67803 for *Oncorhynchus mykiss* (Om), AAL69984 for *Scophthalmus maximus* (Sm), AAT81404 for *Sparus aurata* (Sa), AAO13174 for *Oreochromis niloticus* (On), AAT81406 for *Gadus morhua* (Gm) and AAT81405 for *Clarias gariepinus* (Cg). Human and Atlantic salmon sequences were the same used in the aa alignment. Confidence in the resulting phylogenetic tree branch topology was measured by bootstrapping through 1000 iterations.

#### Functional characterisation

PCR fragments corresponding to the open reading frames (ORFs) of the putative elongases were amplified from salmon intestine cDNA using sense and antisense primers containing a digestion site (underlined) - Elo1BVF1

(CCCAAGCTTGAAATGGAGGCTTTTAATCATAAAC; *HindIII*) and Elo1BVR1 (CCGCTCGAGTCAGTCCACCCGCACTTT; *XhoI*), for gb|FJ237531|, and primers Elovl2VF2 (CCCGAGCTCAAGATGAACCATTTACAAAGTTTGG; *SacI*) and Elovl2VR1 (CCGCTCGAGCTACTTTCTTCTTGAAGCTG; *XhoI*) for gb|FJ237532|. PCR was performed using the high fidelity PfuTurbo<sup>®</sup> DNA Polymerase (Stratagene, Agilent Technologies, Cheshire, U.K.), with an initial denaturing step at 95°C for 2 min, followed by 32 cycles of denaturation at 95°C for 30 s, annealing at 60°C (for primers Elo1BVF1/Elo1BVR1) or 62°C (for primers Elovl2VF2/ Elovl2VR1) for 30 s, and extension at 72°C for 1 min, followed by a final extension at 72°C for 5 min. The ORF of the salmon *elov15a* cDNA was amplified using the primers described in Hastings et al. (2005). The DNA fragments were then digested with the corresponding restriction endonucleases (New England BioLabs, Herts, U.K.) and ligated into a similarly restricted pYES2 yeast expression vector (Invitrogen, Paisley, U.K.). Ligation products were used to transform Top10F' *Escherichia coli* competent cells (Invitrogen) which were screened for the presence of recombinants. The purified plasmids (illustra<sup>™</sup> GFX<sup>™</sup> Micro Plasmid Prep Kit, GE Healthcare Life Sciences, Buckinghamshire, U.K.) containing the three elongase ORFs or the pYES vector alone were then used to transform *Saccharomyces cerevisiae* competent cells (S.c. EasyComp Transformation Kit, Invitrogen). Transformation and selection of yeast with recombinant elongase-pYES2 plasmids, yeast culture and FA analysis was performed as described in detail previously (Hastings et al., 2001, 2005; Agaba et al., 2004; Zheng et al., 2005). Briefly, cultures of recombinant yeast were grown in *S. cerevisiae* minimal medium<sup>-uracil</sup> supplemented with one of the following FA substrates: stearidonic acid (18:4n-3),  $\gamma$ -linolenic acid (18:3n-6), EPA (20:5n-3), ARA (20:4n-6), docosapentaenoic acid (22:5n-3) or docosatetraenoic acid (22:4n-6). Docosapentaenoic and docosatetraenoic acids (>98-99% pure) were purchased from Cayman Chemical Co. (Ann Arbor, U.S.A.) and the remaining FA

substrates (>99% pure) and chemicals used to prepare the *S. cerevisiae* minimal medium<sup>-uracil</sup> were from Sigma Chemical Co. Ltd. (Dorset, U.K.). FA were added to the yeast cultures at final concentrations of 2.5 (C18), 5.0 (C20) and 10.0 (C22)  $\mu\text{M}$ . After 2-days, yeast were harvested and washed, and lipid extracted by homogenization in chloroform/methanol (2:1, v/v) containing 0.01% BHT as antioxidant. FA methyl esters were prepared, extracted, purified, and analyzed by GC in order to calculate the proportion of substrate FA converted to elongated FA product from as  $[\text{product area}/(\text{product area} + \text{substrate area})] \times 100$ . The identity of the FA peaks was based on GC retention time and confirmed by GC-MS as described previously (Hastings et al., 2001; Agaba et al., 2004).

#### Tissue distribution and nutritional regulation

For the tissue expression profile, tissues (intestine, liver, white muscle, red muscle, kidney, spleen, heart, brain, gill and adipose tissue) were dissected from three salmon, immediately frozen in liquid nitrogen and stored at  $-70\text{ }^{\circ}\text{C}$  pending RNA extraction. The fish were  $\sim 150\text{ g}$  post-smolts held in  $7\text{ m}^3$  seawater tanks at ambient temperature, salinity and photoperiod in the Marine Environment Research Laboratory, Machrihanish, Scotland, U.K., and fed a commercial salmon feed based on fish meal and fish oil. The effects of diet on elongase expression were investigated in samples from salmon post-smolts fed four diets with the same basal composition but formulated with different oils, FO, rapeseed oil (RO), linseed oil (LO) or soybean oil (SO). Full descriptions of the diets and the trial have been reported previously (Leaver et al. 2008b). At the end of the trial, 0.5 g of liver and small intestine of 5 fish per dietary treatment were dissected and rapidly disrupted in 5ml of TRI Reagent (Ambion, Applied Biosystems) using an Ultra-Turrax homogeniser (Fisher Scientific, Loughborough,



U.K.), and immediately frozen in liquid nitrogen and stored at  $-70\text{ }^{\circ}\text{C}$  prior to RNA extraction.

#### Tissue RNA extraction and quantitative real time PCR (qPCR)

Total RNA was extracted by organic solvent, according to manufacturer's instructions (Ambion, Applied Biosystems), and RNA quality and quantity assessed by electrophoresis (Bioanalyser 2100, Agilent Technologies, Santa Clara, U.S.A.) and spectrophotometry (NanoDrop ND-1000, Thermo Scientific, Wilmington, U.S.A.), respectively. One  $\mu\text{g}$  of total RNA per sample was reverse transcribed into cDNA using a Verso<sup>TM</sup> cDNA kit (ABgene, Surrey, U.K.), following manufacturer's instructions. Briefly, each 20  $\mu\text{l}$  reaction contained 1  $\mu\text{g}$  of total RNA, 300 ng of random hexamers and 125 ng of anchored oligo-dT, dNTP mix (500  $\mu\text{M}$  each), 1X cDNA synthesis buffer, RT enhancer and Verso enzyme mix. Following cDNA synthesis at  $42\text{ }^{\circ}\text{C}$  for 1 h, reactions were stopped by heating at  $95\text{ }^{\circ}\text{C}$  for 2 min and cDNA diluted 10-fold with water.

Expression of the three elongase transcripts was studied by quantitative real time PCR (qPCR). The qPCR primers were designed in regions corresponding to the 3'UTR region of *elovl5a* and of the two new elongase cDNAs. In addition, amplification of four reference genes (elongation factor-1 $\alpha$  (*elf-1 $\alpha$* ),  *$\beta$ -actin*, an EST previously shown to be unresponsive to VO dietary manipulation, and *18S rRNA*) was also performed, for normalization of the results. The unresponsive EST is an anonymous cDNA feature selected from a salmon cDNA microarray study and identified as a suitable reference gene on the basis of constant expression between different VO diets and time points (Taggart et al., 2008). Table 1 shows the sequence of the primers used, their specific annealing temperatures, the size of fragments produced and the reference sequences used for primer design, using the Primer3 software

([http://biotools.umassmed.edu/bioapps/primer3\\_www.cgi](http://biotools.umassmed.edu/bioapps/primer3_www.cgi)). For tissue distribution, qPCR amplicons corresponding to the three putative elongases were cloned into the pBluescript® II KS phagemid vector (Stratagene), while that of 18S rRNA was cloned into the pCR2.1-TOPO vector (TOPO TA cloning kit, Invitrogen). These recombinant vectors were then restricted with *SacI* or *XhoI* and the linearized plasmid DNA containing the target sequence for each gene quantified spectrophotometrically and serial-diluted to generate a standard curve of known copy numbers. For effects of diet, qPCR analysis used relative quantification with *elf-1 $\alpha$* ,  *$\beta$ -actin* and the unresponsive EST as reference genes, and amplification efficiency of the primer pairs assessed by serial dilutions of cDNA pooled from the samples being quantified. QPCR amplifications were carried out in duplicate using a Quantica machine (Techne, Cambridge, U.K.) in a final volume of 20  $\mu$ l containing 2  $\mu$ l (for the nutritional regulation trial and 18S rRNA) or 5  $\mu$ l (for tissue distribution) diluted (1/10) cDNA, 0.5  $\mu$ M of each primer and 10  $\mu$ l Absolute™ QPCR SYBR® Green mix (ABgene). Amplifications were carried out with a systematic negative control (NTC-non template control, containing no cDNA). The qPCR profiles contained an initial activation step at 95 °C for 15 min, followed by 30 to 40 cycles: 15 s at 95 °C, 15 s at the specific primer pair annealing  $T_m$  (Table I) and 30 s at 72 °C. After the amplification phase, a melt curve of 0.5 °C increments from 75 °C to 90 °C was performed, enabling confirmation of the amplification of a single product in each reaction. The qPCR product sizes were checked by agarose gel electrophoresis and their identity confirmed by sequencing. No primer-dimer formation occurred in the NTC.

### Statistical analysis

For tissue expression profiles, results were expressed as mean normalized values ( $\pm$  SD) corresponding to the ratio between the copy numbers of the putative elongase transcripts and

the copy numbers of the reference gene, 18S rRNA. For the tissues with highest expression level, intestine and liver, a one-way analysis of variance (ANOVA) followed by a Tukey HSD test, at a significance level of  $P < 0.05$ , was performed to compare the expression level of the three transcripts in both tissue samples, using Statistica 6 (StatSoft, Tulsa, U.S.A.). The effects of diet on elongase expression, expressed as the relative expression ratio of each gene in fish fed one of the VOs in relation to those fed FO (control) and normalized by three reference genes (*elf-1 $\alpha$* ,  *$\beta$ -actin* and the unresponsive EST), were analysed for statistical significance using the relative expression software tool (REST-MCS<sup>®</sup>, version 2, <http://www.gene-quantification.info/>), which employs a pair wise fixed reallocation randomisation test (10,000 randomisations) with efficiency correction (Pfaffl et al., 2002).

## Results

### Salmon fatty acyl elongase cDNAs sequences and phylogenetics

A 1650 bp full-length cDNA sequence was obtained by 5' and 3' RACE PCR from gb|DW546112| and was deposited in the GenBank database under the accession number gb|FJ237531|. It contains an ORF of 885 bp encoding a putative protein of 294 aa, sharing 91% aa sequence identity with the previously described Elov15-like salmon elongase (Elov15a) and 93% identity in nucleotide sequence, in the ORF, to *elov15a* (gb|AY170327|). This protein is 70% identical to several mammalian ELOVL5-like proteins (*Rattus norvegicus*, *Mus musculus* and *Homo sapiens*) and 74-81% identical to other fish Elov15-like proteins (*Danio rerio*, *Gadus morhua*, *Oreochromis niloticus*, *Clarias gariepinus*, *Scophthalmus maximus* and *Sparus aurata*). This new transcript was thus named *elov15b*.

For the sequence derived from ti|TC91192|, a full-length cDNA spanning 1792 bp, with a 864 bp ORF encoding a putative 287 aa protein, was obtained and determined to have 79% aa sequence identity to the zebrafish Elov12-like, and 68-71% aa identity to mammalian (rat, mouse and human) ELOVL2-like proteins. This sequence was deposited in the GenBank database under the accession number gb|FJ237532|.

Typically, these newly cloned Elov15-like and Elov12-like proteins both possess the diagnostic histidine box HXXHH motif (Fig. 1) conserved in all elongases and also characteristic of desaturase and hydrolase enzymes containing a di-iron-oxo cluster (Tvrdik et al., 2000; Jakobsson et al., 2006). In addition, they possess two lysine or arginine residuals at the carboxyl terminus, KXRXX in Elov15-like and KKXX in Elov12, which are proposed to function as endoplasmic reticulum (ER) retrieval signals (Jakobsson et al., 2006). By sequence comparison with a mouse ELOVL2 (Tvrdik et al., 2000), five putative transmembrane-spanning domains, containing hydrophobic aa stretches, can be predicted (Fig. 1). Noteworthy is the fact that, of the 17 aa residues found by Leonard et al. (2004) to be highly conserved across 22 members of the elongase family, one residue has been conservatively replaced in SsElov15b (Leu245 replaced by Phe245).

A phylogenetic tree was constructed on the basis of the aa sequence alignments between the putative two new salmon fatty acyl elongases, and other elongases of fish and mammals (Fig 2). The phylogenetic analysis showed that salmon Elov15a and Elov15b clustered with human ELOVL5 and several other fish proteins. In contrast, the putative salmon Elov12 clustered separately from all other Elov15 proteins and grouped with human ELOVL2 and an Elov1 from zebrafish. BLASTX searches of the ENSEMBL genomes of *T. nigroviridis*, *T. rubripes*, *G. aculeatus* and *O. latipes* demonstrated that these fish species do not contain *elov12* homologues.

## Functional characterisation

The elongases were functionally characterized by determining the FA profiles of *S. cerevisiae* transformed with pYES2 vector alone or containing *elovl5a*, *elovl5b* and *elovl2* cDNA ORF inserts grown in the presence of potential FA substrates. The FA composition of the wild yeast insert is characterised by having essentially only 16:0, 16:1n-7, 18:0 and 18:1n-9 (see Hastings et al., 2001). Yeast transformed with vector containing no insert contain these fatty acids together with whichever exogenous FA was added, consistent with the well established lack of PUFA elongase activity in *S. cerevisiae* (Agaba et al., 2004; Hastings et al., 2005). Heterologous expression of salmon *elovl5a* in yeast produced similar results to those previously observed (Hastings et al., 2005), with activity towards C18 and C20 substrates and only residual (1%) C22 to C24 activity (Table 2). The FA compositions of yeast transformed with the pYES2-*elovl5b* construct and grown in the presence of 18:4n-3, 20:5n-3 and 22:5n-3 are shown in Fig. 3A-C. The traces show the major endogenous FA (16:0, 16:1n-7, 18:0 and 18:1n-9, peaks 1-4) and additional peaks corresponding to the substrates and elongation products. Thus exogenously added 18:4n-3 (peak 6) was elongated to 20:4n-3 (peak 7) and 22:4n-3 (peak 8) (Fig. 3A), exogenously added 20:5n-3 (peak 9) was elongated to 22:5n-3 (peak 10) (Fig. 3B), and only a trace of exogenously added 22:5n-3 (peak 10) was elongated to 24:5n-3 (peak 11) (Fig. 3C). Other additional peaks including 18:1n-7 (peak 5), 20:1n-9 and 20:1n-7, indicated some capability of Elov15b to elongate endogenous monounsaturated FAs, as previously observed with Elov15a (Hastings et al., 2005). Elov15b elongated 62% of 18:4n-3 and 71% of 18:3n-6, mainly to the C20 products (58% and 65% converted to 20:4n-3 and 20:3n-6, respectively), and 69% of 20:5n-3 and 48% of 20:4n-6, but only 1-2% of 22:5n-3 and 22:4n-6 were converted into 24:5n-3 and 24:4n-6 (Table 2). These data confirmed *elovl5b* as an ELOVL5-like elongase and its functional similarity to *elovl5a*.

The FA compositions of yeast transformed with the pYES2-*elovl2* construct and grown in the presence of FAs show a distinct pattern (Fig. 3D-F). Exogenously added 18:4n-3 (peak 6) was only poorly elongated to 20:4n-3 (peak 7) (Fig. 3D), whereas exogenously added 20:5n-3 (peak 9) was elongated to 22:5n-3 (peak 10) and, especially, to 24:5n-3 (peak 11) (Fig. 3E), and exogenously added 22:5n-3 (peak 10) was substantially elongated to 24:5n-3 (peak 11) (Fig. 3F). Thus, only 5% and 12% of 18:4n-3 and 18:3n-6, respectively, were elongated, whereas 70% and 59% of 20:5n-3 and 20:4n-6, respectively, were elongated with 52% and 48% recovered as the C24 products (Table 2). Around 31% and 18% of exogenously added 22:5n-3 and 22:4n-6 were elongated to 24:5n-3 and 24:4n-6, respectively. Elongation of monounsaturated FAs was low, with conversion of 16:1n-7 to 18:1n-7 typically around 4%. No evidence for elongation of saturated FAs was observed with any of the salmon Elovls.

#### Tissue distribution of salmon fatty acyl elongase cDNAs

All three elongase genes in salmon were expressed mostly in intestine (pyloric caeca) and liver, followed by brain (Fig. 4). The previously characterized elongase gene, *elovl5a*, showed a more widespread expression than the two new elongase transcripts being also found, albeit at a lower level in terms of absolute copy numbers, in gill, kidney, spleen, heart, adipose tissue, red muscle and white muscle. In intestine, *elovl5a* and *elovl2* appear to have a similarly high expression level, with *elovl5b* showing significantly lower expression. In liver, no significant differences were found, but copy numbers of *elovl5b* were slightly higher than those of the other two transcripts.

#### Nutritional regulation of salmon fatty acyl elongase expression

The regulation of elongase genes, in response to dietary FA composition, was examined in liver and intestine of salmon that had been fed diets containing either FO rich in EPA and DHA, or VOs rich in C18 FA, RO (18:1n-9), SO (18:2n-6) or LO (18:3n-3). Previously, transcriptomic analysis of samples from this VO-feeding experiment revealed changes in the transcript levels of genes in response to the FA composition of the diets, but not of elongase (Leaver et al., 2008b). However, although the array contained *elovl5a*, both *elovl5b* and *elovl2* were absent, and since *elovl5a* and the newly discovered *elovl5b* will cross-hybridise, it is not surprising that changes in expression were not detected by microarray analysis. However, LC-HUFA biosynthesis was elevated in these VO-fed salmon (Leaver et al., 2008b), and thus we examined the expression of all known salmon *elovl* genes in the present work through qPCR using gene-specific primers. Compared to the FO-fed group, there was a significant increase of *elovl2* and *elovl5b* transcripts in the liver of VO-fed fish with normalized expression ratios for *elovl2* of 2.2, 2.6 and 1.9 for treatments RO, SO and LO, respectively, and a significant 1.7- and 1.5-fold increase of *elovl5b* transcript level in fish fed RO and SO diets, respectively (Fig. 5A). No significant differences were found between transcript levels for *elovl5a* in FO-fed fish and VO-fed fish, or between the different VO dietary treatments for any of the genes. For intestine, elongase mRNA showed higher variability within dietary treatments than observed in liver, and so no statistically significant differences were found in the relative expression of any of the elongase transcripts between fish fed the different dietary treatments (Fig. 5B). Nonetheless, the observed trends agreed with the results obtained in liver. Thus, the levels of *elovl2* transcript in fish fed RO, SO and LO diets were 3.0-, 4.2- and 4.4-fold higher, respectively, compared to salmon fed FO. As observed in liver, the level of *elovl5a* transcripts in the intestine was unaffected in fish fed the different VO diets, when compared to the FO group, while RO- and SO-fed fish showed a 1.7- and 1.8-fold increase, respectively, in *elovl5b* transcript level compared to the FO treatment.

## Discussion

Very long chain fatty acyl elongases (ELOVL) are ER membrane-bound proteins responsible for the first, regulatory step (condensation of activated FAs with malonyl-CoA) in the FA elongation pathway, elongating FA that are biosynthesised *de novo* or supplied by the diet. They belong to a gene family that consists of seven members in mice and humans, which differ in FA substrate specificity and have differing spatial and temporal expression patterns (Jakobsson et al., 2006). Of these seven ELOVLs enzymes, ELOVL2 and ELOVL5 have been demonstrated to have a substrate preference for PUFA. In addition, ELOVL4 is critical for normal human and mouse retinal function and there is evidence to indicate that it may also be involved specifically in DHA biosynthesis, which is a major membrane component in these tissues (Zhang et al., 2001; Jakobsson et al., 2006). In mammals, ELOVL2 has greatest activity in the elongation of C20 and C22, but low or, in the case of human, no activity towards C18 PUFA (Leonard et al., 2002). In contrast, mammalian ELOVL5 is very active towards C18 PUFA but does not appear to have the capacity to elongate beyond C22 (Leonard et al., 2000; Inagaki et al., 2002). Other species for which a PUFA elongase gene has been reported and functionally characterised are the nematode *Caenorhabditis elegans* (Beaudoin et al., 2000) and fungus *Mortierella alpina* (Parker-Barnes et al., 2000), with these enzymes being predominantly active on C18 PUFA with virtually no activity towards C20, and the marine microalgae *Pavlova*, which has a unique specificity towards C20 with no C18 or C22 activity (Pereira et al., 2004).

Fish *elovl* cDNAs have been cloned and functionally characterised from a number of species: the freshwater species zebrafish, common carp and tilapia, the salmonids, Atlantic salmon and rainbow trout, and the marine species cod, turbot and sea bream (Agaba et al.,



2004, 2005; Meyer et al., 2004; Hastings et al., 2005). More recently, a fatty acyl elongase from masu salmon has been cloned and over-expressed in zebrafish (Alimuddin et al., 2008). Phylogenetic analysis groups all these previously described *elovl* cDNAs into a cluster with greatest similarity to mammalian *ELOVL5* (Meyer et al., 2004; Agaba et al., 2005; Alimuddin et al., 2008). All the fish *elovl5* cDNAs tested lengthened monounsaturated FA and n-3 and n-6 PUFA with chain lengths from C18 to C22, with residual C22-C24 activity (Agaba et al., 2004, 2005; Hastings et al., 2005). A rainbow trout *Elov15* has also been described with C18 to C22, but no C22-24 activity (Meyer et al., 2004). Thus, fish *elovl5* appears to have wider PUFA specificity than its mammalian homologue, although activity towards C22 is minor, compared to C18 and C20 (Agaba et al., 2004, 2005; Hastings et al., 2005). No *ELOVL2*-like genes have been reported so far in a non-mammalian vertebrate. However, searches in the GenBank and Atlantic salmon EST databases, as well as in the zebrafish genome, demonstrated that a second PUFA *elovl* gene exists in Atlantic salmon and zebrafish and that this gene is clearly related to the mammalian *ELOVL2* (Fig. 2). In addition, a second Atlantic salmon *elovl5* gene, here named *elovl5b* to distinguish it from the previously cloned salmon gene (Hastings et al., 2005), now termed *elovl5a*, was identified. Preliminary work has revealed two distinct genomic sequences corresponding to these two transcripts, indicating that they are encoded by separate loci (unpublished results). Given the high degree of similarity of these two salmon *elovl5* genes (91% and 93% identical at the protein and ORF nucleotide sequence level, respectively), we can speculate that they may result from a duplicated locus, as the result of the recent salmonid tetraploidisation (Allendorf and Thorgaard, 1984). This conclusion is supported by searches of ENSEMBL fish genomes, which contain only single copies of *elovl5*. Analysis of the deduced aa sequences of *Elov15b* and *Elov12* showed that they possess characteristic features of microsomal membrane-bound enzymes, including a single histidine box redox centre motif, a canonical ER retention signal

(carboxyl-terminal dilysine/arginine targeting signal) and multiple transmembrane regions (Tvrdik et al., 2000; Jakobsson et al., 2006).

Functional characterisation of the new *elovl* cDNAs by heterologous expression in yeast confirmed the identities predicted by sequence homology and phylogenetic analysis. Elov15b had similar activity to the previously reported salmon Elov15a (Hastings et al., 2005) and to that of other fish Elov1's which clustered with mammalian ELOVL5 (Agaba et al., 2004, 2005). The only difference appears to be in terms of the relative conversion of n-3 and n-6 C18 (but not C20) FAs, as salmon Elov15a had a preference for n-3 FA substrates, which was not apparent in Elov15b. This preference seems variable as some fish Elov15 proteins showed similar activities with n-3 and n-6 FAs, and cod Elov15 was more active towards n-6 FAs (Agaba et al., 2005). The C18 and C20 PUFA specificity of both salmon Elov15 proteins was generally comparable to that of mammalian ELOVL5 (Leonard et al., 2000; Inagaki et al., 2002) but a residual activity towards C22 was also measured. In addition, as observed with human and rat ELOVL5 homologues (Leonard et al., 2000; Inagaki et al., 2002), some capacity to elongate monounsaturated FAs was found for both salmon Elov15 proteins, as in other fishes (Meyer et al., 2004; Agaba et al., 2005). More importantly, the other salmon cDNA described here is the first *elovl2*-like elongase to be cloned and functionally characterised in fish. In comparison to mouse ELOVL2 (Leonard et al., 2002), the salmon Elov12 protein showed similar low activity towards C18 and high activity towards C20 and C22 but had a low capacity to convert monounsaturated FAs (around 4% conversion of 16:1n-7), which was not observed in mammalian ELOVL2 (Leonard et al., 2002). Notably, the major difference in comparison to fish Elov15 and mammalian ELOVL5 proteins is the higher activity towards C22 LC-PUFA exhibited by Elov12.

Other fish EST's potentially corresponding to putative *elovl2* transcript are those of zebrafish (gb|NM\_001040362|), *Pimephales promelas* (gb|DT135746|; partial cDNA) and

*Ictalurus punctatus* (gb|BM438219.1]; partial cDNA), all Ostariophysian freshwater species. Although the analysis of fish ELOVL genes is not yet exhaustive, searches in the pufferfish, stickleback and medaka genomes have not indicated the presence of *elovl2*-like genes and thus it is possible to speculate that all fish of the order Acanthopterygii lack *ELOVL2* homologues. Therefore, the characterised *elovl5* cDNAs of sea bream, turbot and cod would be the sole PUFA *elovl* gene in these species. Elovl5 proteins have a very limited capacity towards C22 and because biosynthesis of DHA in vertebrates requires elongation from C22 to a C24 PUFA intermediate, followed by a peroxisomal  $\beta$ -oxidation chain-shortening step (Buzzi et al., 1997; Sprecher, 2000), fish which lack *elovl2* would be restricted in their ability to produce this essential FA. In fact where LC-PUFA biosynthesis has been directly measured in fish, it is clear that salmonids have substantially greater capacity than marine fish, such as sea bass, a member of the Acanthopterygii (Mourente et al., 2005). Previously, the ability of both salmon and zebrafish to produce LC-PUFA endogenously has been shown to be due to the presence of both  $\Delta 5$  and  $\Delta 6$  desaturase genes (Hastings et al., 2005; Zheng et al., 2005a), which contrasts with the single  $\Delta 6$  gene so far discovered in Acanthopterygii (Tocher et al., 2006). Here we further show that the varying competences of different fish to biosynthesise LC-PUFA, particularly DHA, might not only depend on their genome complement of desaturase genes but also of elongases. The phylogenetic distribution of the *elovl2* gene suggests that it must have been lost from the evolutionary line leading to the Acanthopterygii, but retained in Ostariophysi (e.g. zebrafish) and Salmoniformes. It is tempting to speculate that, since both Ostariophysi and Salmonid fish spend all, or at least a substantial period of time in freshwater, the retention of *elovl2* and the diversification of desaturase genes is an adaptation to the relative deficiency of preformed LC-PUFA in their freshwater habitats compared to marine ecosystems, which are fuelled by LC-PUFA producing phytoplankton (Sargent et al., 2002; Brett and Muller-Navarra, 1997). Although also speculative, it can be

observed that the vast majority of Acanthopterygian species inhabit marine environments. Those that inhabit freshwater tend to live in productive lacustrine or large riverine systems. Few, if any, exist in the nutrient-poor streams and lakes that salmonids inhabit.

The tissue distribution profile of salmon *elovl5a* had been reported previously and agreed broadly with that found here (Zheng et al., 2005a). The tissue distribution of *elovl5b* largely resembled that of *elovl5a*, apart from significantly lower transcript levels in intestine, slightly higher in liver and lower in brain. In general, *elovl5a*, *elovl5b* and *elovl2* are expressed predominately in intestine and liver. Since these tissues are the major sites of lipid synthesis and distribution, this expression profile is consistent with a role in FA biosynthesis. In the remaining tissues tested *elovl5a* was the most highly expressed. Rat and human *ELOVL5* was expressed in most tissues examined (Inagaki et al., 2002; Leonard et al., 2000; Wang et al., 2005) and *ELOVL2* transcripts, as in salmon, appeared to have a more restricted distribution in mammals (Leonard et al., 2002; Tvrdik et al., 2000; Wang et al., 2005). Salmon brain exhibited the third highest expression of the tissues analysed but expression levels of *elovl2* were very low, over one order of magnitude lower than *elovl5b* and even less than *elovl5a*. This might mean that DHA, which comprises a great proportion of the brain's cellular membranes, either has an exogenous source or that another gene, such as a homologue to the mammalian *ELOVL4*, for instance, is involved in DHA biosynthesis in the brain. This hypothesis should be investigated in the future.

This work has clearly shown that salmon *elovl5b* and *elovl2*, but not *elovl5a*, transcripts are increased in response to inclusion of VO in the diet. However, previous studies examining the effect of FO replacement by VO in the diets of Atlantic salmon on *elovl5a* expression have yielded inconsistent results (Zheng et al. 2004; Zheng et al., 2005a, 2005b). In mammals, *ELOVL5* and *ELOVL2* expression and regulation has been described. Compared to *Elov15*, levels of *Elov12* mRNA were low in rat liver (Wang et al., 2005), whilst in salmon

these genes are expressed at similar levels. If these transcript levels are reflected in protein concentrations then some differences in the relative importance and contribution of *elovl2* and *elovl5* to LC-PUFA synthesis in salmon and rats might be expected. In regard to nutritional regulation, Inagaki et al. (2002) reported a lack of dietary regulation, by cycles of fasting-refeeding, whilst Wang et al. (2005) observed that overnight starvation and feeding FO-enriched diets decreased *Elov15* mRNA levels in rat liver. No diet-induced changes were noted in *Elov12* expression (Wang et al., 2005). Thus, *Elov15* rather than *Elov12* appeared to play the major role (along with  $\Delta 5$  and  $\Delta 6$  desaturases) in response to changes in dietary lipid composition in rat (Wang et al., 2005), whilst *elovl2* may have a more prominent role in salmon. This might not be surprising, considering the differences in dietary regimes of mammals and carnivorous fish, particularly concerning the relative supply of C20 and C22 FAs.

Hepatic *Elov12* and *Elov15* are both regulated by sterol regulatory element binding protein (SREBP) transcription factors in mouse (Horton et al., 2003) and SREBP activation is increased by low cholesterol and decreased by increased PUFA (Espenshade, 2006), suggesting potential mechanisms for the observed changes in expression of these genes in salmon fed VO diets which are low in both PUFA and cholesterol (Leaver, 2008b). These increases in *elovl2* and *elovl5* are important in salmon aquaculture because they enable physiological adaptation to VO diets which are deficient in FA greater than 18 carbons, and enable LC-PUFA biosynthesis. Moreover, the observed difference in the nutritional regulation of *elovl5a* and *elovl5b* genes will provide an interesting opportunity to advance our knowledge on the regulation and transcriptional control of gene expression, by comparison of the presumably very similar upstream regulatory regions of these genes that have resulted from a recent gene duplication event.

In conclusion, two *elovl* genes, homologues of mammalian *ELOVL2* and *ELOVL5*, have been characterised in Atlantic salmon, which together with the previously characterised salmon  $\Delta 5$  and  $\Delta 6$  fatty acyl desaturases, explain the ability of these fish to biosynthesise LC-PUFA, including DHA, more efficiently than Acanthopterygian fish species which seem to lack an *elovl2* gene. This, and the ability of salmon to regulate these genes in response to dietary quality, might to some extent also explain the success of salmonids in colonising nutrient poor freshwater habitats and to better tolerate VO-based aquaculture diets than cultured Acanthopterygians. The results obtained are not only of high relevance in advancing our understanding of the molecular basis of LC-PUFA biosynthesis and regulation in fish, but also have biotechnological significance. The use of transgenic techniques to enhance EPA and DHA biosynthesis is likely to become routine in various organisms of commercial importance (Meyer et al., 2004). A previously cloned and characterised zebrafish desaturase (Hastings et al., 2001) has already been utilised to produce transgenic *Arabidopsis* capable of producing EPA and DHA at low levels (Robert et al., 2005). The identification of salmon *elovl2*, a fatty acyl elongase with high activity in the penultimate steps of LC-HUFA biosynthesis, might provide a route for increasing EPA and DHA production in transgenic organisms. Indeed, interest is also likely to become directed towards optimising HUFA biosynthesis pathways through transgenic techniques in farmed fish to enable efficient and effective use of dietary VO while maintaining the nutritional quality of the fish as a primary source in the human food basket of the omega-3 or n-3 LC-PUFA. Alimuddin et al. (2008) have already increased EPA and DHA biosynthesis in zebrafish by over-expressing an *elovl5*-like gene from masu salmon. Given the differences in substrate specificity of *elovl5* and *elovl2*, reported here for the first time in fish, and the necessity of C22 to C24 PUFA intermediate elongation for DHA biosynthesis (Sprecher, 2000), the potential benefits of over-expressing these two genes in parallel can easily be recognized.

**Acknowledgements** This work and XZ were supported by the Biotechnology and Biological Sciences Research Council (BBSRC) Responsive Mode Grant, BB/C51237X/1 (Transcriptional control of polyunsaturated fatty acid synthesis in fish). SM was supported by “Fundação para a Ciência e a Tecnologia”, Portugal (grant SFRH/BPD/34247/2006) and OM by the postdoctoral research programme of the Fundación Española para la Ciencia y la Tecnología (Ministerio de Educación y Ciencia).

## References

- Agaba, M., Tocher, D.R., Dickson, C., Dick, J.R., and Teale, A.J. (2004). Zebrafish cDNA encoding multifunctional fatty acid elongase involved in production of eicosapentaenoic (20:5n-3) and docosahexaenoic (22:6n-3) acids. *Mar Biotechnol* 6:251-261.
- Agaba, M.K., Tocher, D.R., Dickson, C.A., Zheng, X., Dick, J.R., and Teale, A.J. (2005). Cloning and functional characterisation of polyunsaturated fatty acid elongases from marine and freshwater teleost fish. *Comp Biochem Physiol* 142B:342-352.
- Alimuddin, Kiron, V., Satoh, S., Takeuchi, T., and Yoshizaki, G. (2008). Cloning and over-expression of a masu salmon (*Oncorhynchus masou*) fatty acid elongase-like gene in zebrafish. *Aquaculture* 282:13-18.
- Allendorf, F.W., and Thorgaard, G.H. (1984). Tetraploidy and the evolution of salmonid fishes. In: Turner, B.J. (ed) *Evolutionary genetics of salmonid fishes*. Plenum Press, New York, pp. 1-53.
- Beaudoin, F., Michaelson, L.V., Lewis, M.J., Shewry, P.R., Sayanova, O., and Napier, J.A. (2000). Production of C20 polyunsaturated fatty acids (PUFAs) by pathway engineering:

- identification of a PUFA elongase component from *Caenorhabditis elegans*. *Biochem Soc Trans* 28:661-663.
- Bell, J.G., and Waagbø, R. (2008). Safe and nutritious aquaculture produce: benefits and risks of alternative sustainable aquafeeds. In: Holmer, M., Black, K.D., Duarte, C.M., Marba, N., and Karakassis, I. (eds) *Aquaculture in the Ecosystem*. Springer Verlag BV, pp. 185-225.
- Brett, M.T., and Muller-Navarra, D.C. (1997). The role of highly unsaturated fatty acids in aquatic foodweb processes. *Freshwater Biol* 38:483-499.
- Buzzi, M., Henderson, R.J., and Sargent, J.R. (1997). Biosynthesis of docosahexaenoic acid in trout hepatocytes proceeds via 24-carbon intermediates. *Comp Biochem Physiol* 116:263-267.
- Cook, H.W. (1996). Fatty acid desaturation and chain elongation in eukaryotes. In: Vance, D.E., and Vance, J.E. (eds) *Biochemistry of Lipids, Lipoproteins and Membranes*. Elsevier, Amsterdam, pp. 129 -152.
- Espenshade, P.J. (2006). SREBPs: sterol-regulated transcription factors. *J Cell Sci* 119:973-976.
- Food and Agricultural Organisation, FAO (2006). *State of world aquaculture 2006*, FAO Fisheries Technical paper No. 500, FAO, Rome.
- Hastings, N., Agaba, M., Tocher, D.R., Leaver, M.J., Dick, J.R., Sargent, J.R., and Teale, A.J. (2001). A Vertebrate Fatty Acid Desaturase with  $\Delta 5$  and  $\Delta 6$  Activities. *Proc Natl Acad Sci USA* 98:14304-14309.
- Hastings, N., Agaba, M.K., Tocher, D.R., Zheng, X., Dickson, C.A., Dick, J.R., and Teale, A.J. (2005). Molecular cloning and functional characterization of fatty acyl desaturase and elongase cDNAs involved in the production of eicosapentaenoic and docosahexaenoic acids from  $\alpha$ -linolenic acid in Atlantic salmon (*Salmo salar*). *Mar Biotechnol* 6:463-474.



- Horton, J.D., Shah, N.A., Warrington, J.A., Anderson, N.N., Park, S.W., Brown, M.S., and Goldstein, J.L. (2003). Combined analysis of oligonucleotide microarray data from transgenic and knockout mice identifies direct SREBP target genes. *Proc Natl Acad Sci USA* 100: 12027-12032.
- Inagaki, K., Aki, T., Fukuda, Y., Kawamoto, S., Shigeta, S., Ono, K., and Suzuki, O. (2002). Identification and expression of a rat fatty acid elongase involved the biosynthesis of C18 fatty acids. *Biosci Biotechnol Biochem* 66:613-621.
- Izquierdo, M., Obach, A., Arantzamendi, L., Montero, D., Robaina, L., and Rosenlund, G. (2003). Dietary lipid sources for seabream and seabass: growth performance, tissue composition and flesh quality. *Aquacult Nutr* 9:397-407.
- Jakobsson, A., Westerberg, R., and Jacobsson, A. (2006). Fatty acid elongases in mammals: their regulation and roles in metabolism. *Prog Lipid Res* 45:237-249.
- Leaver, M.J., Bautista, J.M., Björnsson, B.T., Jönsson, E., Krey, G., Tocher, D.R., and Torstensen, B.E. (2008a). Towards fish lipid nutrigenomics: current state and prospects for fin-fish aquaculture. In: Sundell, K., and Power, D. (ed) *State-Of-The-Art Scientific and/or Technologic Reviews on the Use of Molecular Methods and Functional Genomics in Aquaculture*. *Rev Fish Sci* 16 Supplement 1:73-94.
- Leaver, M.J., Villeneuve, L.A.N., Obach, A., Jensen, L., Bron, J.E., Tocher, D.R., and Taggart, J.B. (2008b). Functional genomics reveals increases in cholesterol biosynthetic genes and highly unsaturated fatty acid biosynthesis after dietary substitution of fish oil with vegetable oils in Atlantic salmon (*Salmo salar*). *BMC Genomics* 9 299 doi:10.1186/1471-2164-9-299.
- Leonard, A.E., Bobik, E.G., Dorado, J., Kroeger, P.E., Chuang, L.-T., Thurmond, J.M., Parker-Barnes, J.M., Das, T., Huang, Y.-S., and Murkerji, P. (2000). Cloning of a human

- cDNA encoding a novel enzyme involved in the elongation of long-chain polyunsaturated fatty acids. *Biochem J* 350:765-770.
- Leonard, A.E., Kelder, B., Bobik, E.G., Chuang, L.-T., Lewis, C.J., Kopchick, J.J., Mukerji, P., and Huang, Y.-S. (2002). Identification and expression of mammalian long-chain PUFA elongation enzymes. *Lipids* 37:733-740.
- Leonard, A.E., Pereira, S.L., Sprecher, H., Huang, Y.-S. (2004). Elongation of long-chain fatty acids. *Prog Lipid Res* 43:36-54.
- Meyer, A., Kirsch, H., Domergue, F., Abbadi, A., Sperling, P., Bauer, J., Cirpus, P., Zank, T.K., Moreau, H., Roscoe, T.J., Zähringer, U., and Heinz, E. (2004). Novel fatty acid elongases and their use for the reconstitution of docosahexaenoic acid biosynthesis. *J Lipid Res* 45:1899-1909.
- Mourente, G., Dick, J.R., Bell, J.G., and Tocher, D.R. (2005). Effect of partial substitution of dietary fish oil by vegetable oils on desaturation and oxidation of [1-<sup>14</sup>C] 18:3n-3 and [1-<sup>14</sup>C]20:5n-3 in hepatocytes and enterocytes of European sea bass (*Dicentrarchus labrax* L.). *Aquaculture* 248:173-186.
- Parker-Barnes, J.M., Das, T., Bobik, E., Leonard, A.E., Thurmond, J.M., Chuang, L.-T., Huang, Y.-S., and Mukerji, P. (2000). Identification and characterization of an enzyme involved in the elongation of n-6 and n-3 polyunsaturated fatty acids. *Proc Natl Acad Sci USA* 97:8284-8289.
- Pereira, S.L., Leonard, A.E., Huang, Y.-S., Chuang, L.-T., and Mukerji, P. (2004). Identification of two novel microalgal enzymes involved in the conversion of the ω<sub>3</sub>-fatty acid, eicosapentaenoic acid, into docosahexaenoic acid. *Biochem J* 384:357-366.
- Pfaffl, M.W., Horgan, G.W., and Dempfle, L. (2004). Relative expression software tool (REST©) for group-wise comparison and statistical analysis of relative expression results in real-time PCR, *Nucleic Acids Res* 30 e36; doi:10.1093/nar/30.9.e36, 2002.

- Regost, C., Arzel, J., Robin, J., Rosenlund, G., and Kaushik, S.J. (2003). Total replacement of fish oil by soybean or linseed oil with a return to fish oil in turbot (*Psetta maxima*). 1. Growth performance, flesh fatty acid profile, and lipid metabolism. *Aquaculture* 217:465-482.
- Robert, S.S., Singh, S.P., Zhou, X.-R., Petrie, J.R., Blackburn, S.I., Mansour, P.M., Nichols, P.D., Liu, Q. and Green, A.G. (2005). Metabolic engineering of *Arabidopsis* to produce nutritionally important DHA in seed oil. *Functional Plant Biol* 32:473-479.
- Saitou, N., and Nei, M. (1987). The neighbor-joining method. A new method for reconstructing phylogenetic trees. *Mol Biol Evol* 4:406-425.
- Sargent, J.R., Tocher, D.R., and Bell, J.G. (2002). The Lipids. In: Halver, J.E., and Hardy, R.W. (eds) *Fish Nutrition*, 3<sup>rd</sup> ed. Academic Press, San Diego, pp. 181-257.
- Sprecher, H. (2000). Metabolism of highly unsaturated *n*-3 and *n*-6 fatty acids. *Biochim Biophys Acta* 1486:219-231.
- Taggart, J.B., Bron, J.E., Martin, S.A.M., Seear, P.J., Høyheim, B., Talbot, R., Carmichael, S.N., Villeneuve, L.A.N., Sweeney, G.E., Houlihan, D.F., Secombes, C.J., Tocher, D.R., and Teale, A.J. (2008). A description of the origins, designs and performance of the TRAIT-SGP Atlantic salmon *Salmo salar* L. cDNA microarray. *J Fish Biol* 72:2071-2094.
- Tidwell, J.H., and Allan, G.L. (2002). Fish as food: aquaculture's contribution. *World Aquac* 33:44-48.
- Tocher, D.R. (2003). Metabolism and functions of lipids and fatty acids in teleost fish. *Rev Fisheries Sci* 11:107-184.
- Tocher, D.R., Zheng, X., Schlechtriem, C., Hastings, N., Dick, J.D., and Teale, A.J. (2006). Highly unsaturated fatty acid synthesis in marine fish; cloning, functional characterization

- and nutritional regulation of fatty acid  $\Delta 6$  desaturase of Atlantic cod (*Gadus morhua* L.).  
Lipids 41:1003-1016.
- Tvrdik, P., Westerberg, R., Silve, S., Asadi, A., Jakobsson, A., Cannon, B., Loison, G., and  
Jacobsson, A. (2000). Role of a new mammalian gene family in the biosynthesis of very  
long chain fatty acids and sphingolipids. J Cell Biol 149:707-717.
- Wang, Y., Botolin, D., Christian, B., Busik, J., Xu, J., and Jump, D.B. (2005). Tissue-specific,  
nutritional, and developmental regulation of rat fatty acid elongases. J Lipid Res 46:706-  
715.
- Zhang, K., Kniazeva, M., Han, M., Li, W., Yu, Z., Yang, Z., Li, Y., Metzker, M.L.,  
Allikmets, R., Zack, D.J., Kakuk, L.E., Lagali, P.S., Wong, P.W., MacDonald, I.M.,  
Sieving, P.A., Figueroa, D.J., Austin, C.P., Gould, R.J., Ayyagari, R., and Petrukhin, K.  
(2001). A 5-bp deletion in *ELOVL4* is associated with two related forms of autosomal  
dominant macular dystrophy. Nat Gen 27:89-93.
- Zheng, X., Tocher, D.R., Dickson, C.A., Bell, J.G., and Teale, A.J. (2004). Effects of diets  
containing vegetable oil on expression of genes involved in highly unsaturated fatty acid  
biosynthesis in liver of Atlantic salmon (*Salmo salar*). Aquaculture 236:467-483.
- Zheng, X., Tocher, D.R., Dickson, C.A., Dick, J.R., Bell, J.G., and Teale, A.J. (2005a). Highly  
unsaturated fatty acid synthesis in vertebrates: new insights with the cloning and  
characterisation of a  $\Delta 6$  desaturase of Atlantic salmon. Lipids 40:13-24.
- Zheng, X., Torstensen, B.E., Tocher, D.R., Dick, J.R., Henderson, R.J., and Bell, J.G. (2005b).  
Environmental and dietary influences on highly unsaturated fatty acid biosynthesis and  
expression of fatty acyl desaturases and elongase genes in liver of Atlantic salmon (*Salmo  
salar*). Biochim Biophys Acta 1734:13-24.

Table 1

Sequence and annealing temperature (T<sub>m</sub>) of the primer pairs used, size of the fragment produced and accession number of the sequence used as reference for primer design, for real-time quantitative PCR (q-PCR) determinations of the transcript level of Atlantic salmon elongase genes.

Transcript	Primer name	Primer sequence	Fragment	T <sub>m</sub>	Accession No.
<i>elovl5a</i>	Elo1UTR-SM-1F	5'-ACAAGACAGGAATCTCTTTCAGATTAA-3'	137 bp	60°C	AY170327 <sup>1</sup>
	Elo1UTR-SM-1R	5'-TCTGGGGTACTGTGCTATAGTGAC-3'			
<i>elovl5b</i>	Elo2UTR-5F	5'-ACAAAAAGCCATGTTTATCTGAAAGA-3'	141 bp	60°C	DW546112 <sup>1</sup>
	Elo2UTR-5R	5'-CACAGCCCCAGAGACCCACTT-3'			
<i>elovl2</i>	Elo2-SM-1F	5'-CGGGTACAAAATGTGCTGGT-3'	145 bp	60°C	TC91192 <sup>2</sup>
	Elo2-SM-1R	5'-TCTGTTTGCCGATAGCCATT-3'			
<i>elf-1α</i>	ELF-1A jbt2	5'-CTGCCCCTCCAGGACGTTTACAA-3'	175 bp	60°C	AF321836 <sup>1</sup>
	ELF-1A jbt2	5'-CACCGGCATAGCCGATTCC-3'			
<i>β-actin</i>	BACT-F	5'-ACATCAAGGAGAAGCTGTGC-3'	141 bp	56°C	AF012125 <sup>1</sup>
	BACT-R	5'-GACAACGGAACCTCTCGTTA-3'			
unresp. EST	B2F	5'-AGCCTATGACCAACCCACTG-3'	224 bp	60°C	TC63899 <sup>2</sup>
	B2R	5'-TGTTACAGCTCGTTTACCG-3'			
<i>18S rRNA</i>	ss18S-2F	5'-GGCGCCCCCTCGATGCTCTTA-3'	189 bp	65°C	AJ427629 <sup>1</sup>
	ss18S-2R	5'-CCCCGGCCGTCCCTCTTAAT-3'			

<sup>1</sup> GenBank (<http://www.ncbi.nlm.nih.gov/>)

<sup>2</sup> Atlantic salmon Gene Index (<http://compbio.dfci.harvard.edu/tgi/>)

Table 2

Functional characterisation of three Atlantic salmon elongase genes, coding for Elovl5a, Elovl5b and Elovl2 proteins. Results are expressed as a percentage of total fatty acid (FA) substrate converted to elongated product. Percentage of stepwise conversion into intermediary products of the elongation pathway is also shown.

FA Substrate	Product	Conversion (%)			Activity
		Elovl5a	Elovl5b	Elovl2	
18:4n-3	20:4n-3	56	58	4	C18→20
	22:4n-3	7	4	2	C20→22
	24:4n-3	0	0	0	C22→24
		Total: 63	Total: 62	Total: 6	
18:3n-6	20:3n-6	43	65	7	C18→20
	22:3n-6	5	6	3	C20→22
	24:3n-6	0	0	2	C22→24
		Total: 48	Total: 71	Total: 12	
20:5n-3	22:5n-3	36	68	18	C20→22
	24:5n-3	1	1	52	C22→24
		Total: 37	Total: 69	Total: 70	
20:4n-6	22:4n-6	23	48	11	C20→22
	24:4n-6	1	1	48	C22→24
		Total: 24	Total: 49	Total: 59	
22:5n-3	24:5n-3	1	1	31	C22→24
22:4n-6	24:4n-6	1	2	18	C22→24

## Legends to Figures

**Fig. 1** ClustalW2 alignment of the deduced amino acid sequences of Atlantic salmon (*Salmo salar*; Ss) elongases, SsElov15a (Hastings et al., 2005; NP\_001117039) and the two newly cloned SsElov15b and SsElov12 (mRNA sequence deposited in the GenBank database under accession numbers gb|FJ237531| and gb| FJ237532|, respectively), together with human (*Homo sapiens*; Hs) elongases, HsELOVL5 (NP\_068586) and HsELOVL2 (NP\_060240). Identical residues are shaded black and similar residues (based on the Gonnet matrix, using ClustalW2 default parameters) are shaded grey. Indicated are the conserved histidine box motif, characteristic of desaturases and hydrolase enzymes containing a di-iron-oxo cluster (underlined), five (I-V) putative membrane-spanning domains predicted by Tvrdik et al. (2000) (underlined with dashed line) and, in bold, the lysine or arginine residuals proposed to function as ER retrieval signals (Jakobsson et al., 2006). An asterisk indicates the 17 aa residues found by Leonard et al. (2004) to be highly conserved across PUFA elongases.

**Fig. 2** Phylogenetic tree comparing putative amino acid sequences of Atlantic salmon, *Salmo salar* (Ss), Elov12 and Elov15b, cloned in this work, and that of Elov15a (NP\_001117039), with those of other fatty acyl elongases: Human, *Homo sapiens* (HsELOVL2 NP\_060240 and HsELOVL5 NP\_068586), zebrafish, *Danio rerio* (Dr-NP\_956747 and Dr-NP\_001035452), trout, *Oncorhynchus mykiss* (Om-AAV67803), turbot, *Scophthalmus maximus* (Sm-AAL69984), seabream, *Sparus aurata* (Sa-AAT81404), tilapia, *Oreochromis niloticus* (On-AAO13174), cod, *Gadus morhua* (Gm-AAT81406) and catfish, *Clarias gariepinus* (Cg-AAT81405). The tree was constructed using the Neighbour Joining method with ClustalX. Numbers represent the frequencies with which the tree topology presented was replicated after 1000 bootstrap iterations. The scale refers to the horizontal branch length and corresponds to the amino acid substitution rate per site.

**Fig. 3** Functional characterisation of Atlantic salmon putative fatty acyl elongases, *Elovl5b* (A-C) and *Elovl2* (D-F) in transgenic *Saccharomyces cerevisiae* grown in the presence of n-3 FAs; 18:4n-3 (A, D); 20:5n-3 (B, E); and 22:5n-3 (C, F). Fatty acids were extracted from yeast transformed with pYES2 vector containing the ORF of the putative fatty acyl elongase cDNA as an insert. Peaks 1-4 represent the main endogenous FAs of *S. cerevisiae*, namely 16:0 (1), 16:1n-7 (2), 18:0 (3) and 18:1n-9 (4). Peak 5 corresponds to 18:1n-7 resultant from the elongation of the yeast endogenous 16:1n-7 and the remaining main additional peaks (6-11) correspond to the exogenously added FAs and the products of their elongation – 18:4n-3 (6), 20:4n-3 (7), 22:4n-3 (8), 20:5n-3 (9), 22:5n-3 (10), and 24:5n-3 (11). Other minor peaks are 18:2n-6, 20:1n-9 and 20:1n-7, the latter two resulting from the elongation of 18:1n-9 and 18:1n-7 (panels A-C). Vertical axis, FID response; horizontal axis, retention time.

**Fig. 4** Tissue expression profile of *elovl5a*, *elovl5b* and *elovl2* in Atlantic salmon, determined by RT-qPCR. Absolute copy numbers were quantified for each transcript and were normalized by absolute levels of ribosomal 18s (values shown on top of each column and represented diagrammatically in logarithmic scale). I- Intestine, L- liver, WM- white muscle, RM- red muscle, K- kidney, SPL- spleen, H- heart, BR- brain. Results are means (n=3) ± S.D. Different letters in intestine and liver columns indicate significant differences (P<0.05) between transcripts in those tissues.

**Fig. 5** Nutritional regulation of elongase genes *elovl5a*, *elovl5b* and *elovl2*, in Atlantic salmon fed diets containing fish oil (FO), rapeseed oil (RO), soybean oil (SO) or linseed oil (LO), in the liver (A) and intestine (B) tissues, determined by RT-qPCR. The results shown are the normalised expression ratio (reference genes: *elf-1 $\alpha$* ,  *$\beta$ -actin* and a flatliner EST) of the target



transcripts in one of the VO treatments (RO, SO or LO), in relation to the FO control treatment. Values are means (n=5) with S.E. and asterisks represent significant differences (P<0.05) between the column's dietary treatment and the FO treatment, for the respective transcript (REST-MCS-2006).

```

SeElov15a  ---MEIFNYKLNMYIDSWMGRRERVQCGILLDNWPTFALTVMVTLIVWLGKRYMRHRQ 57
SeElov15b  ---MEAFNHKLNITYIDSWMGRRERVQCGILLDNWPTFALTVMVTLIVWLGKRYMRHRQ 57
HeELOVL5   ---MEHFDAISLTYFKALLGRRITRWGCFLLDNWPTFICSVVTLIVWLGKRYMRNRQ 57
HeELOVL2   MEHLKAFDDEINAFLNMEGRISRWGCFLLSGLDPTFFLTVVTLIVWLGKRYMRNRQ 60
SeElov12   MNHLQSLDERLNKLYFLFEERSEKRGKLLHSLLPTLSLTLIVVTSVWLGSKRYMRNRP 60
          .....
          I
SeElov15a  PVSCGCLLVVNLGTLILSFTMFTDMVSAVWHGDMNYCCDTHSAGEIDTKIINVLWVY 117
SeElov15b  PVSCGCLLVVNLGTLILSFTMFTDMVSAVWGGNNEYCCDTHSAGEIDTKIINVLWVY 117
HeELOVL5   PVSCGCLLVVNLGTLILSFTMFCDDVIGWVGKGNFYCCGTRIMGSDMKIIRVLWVY 117
HeELOVL2   ALSLGCLLVVNLGTLILSFTMLADLLSIWEGGNYCCDGLTSAGEIDIRVAKVLWVY 120
SeElov12   AYSLKAVLVVNSVGLSLVMDVPLVSATLSAGKPCGGLHEGSDLRVAKVWVY 120
          .....
          II
SeElov15a  ESHLIEEDTFFELRKNHDTFFLVYHHSVFNINWVFNWVPCGSRPCASINSEH 177
SeElov15b  ESHVIEEDTFFELRKNHDTFFLVYHHSVFNINWVFNWVPCGSRPCASINSEH 177
HeELOVL5   ESHLIEEDTFFELRKNHDTFFLVYHHSVFNINWVFNWVPCGSRPCASINSEH 177
HeELOVL2   ESHVIEEDTFFELRKNHDTFFLVYHHSVFNINWVFNWVPCGSRPCASINSEH 180
SeElov12   ESHVIEEDTFFELRKNHDTFFLVYHHSVFNINWVFNWVPCGSRPCASINSEH 180
          * * * * *
          III
SeElov15a  VLVSYVCLSAVCAIRPVLWKRKYITGCOLIORFLIMSCICAVWPCGSRPCAVFCIF 237
SeElov15b  VLVSYVCLSAVCAIRPVLWKRKYITGCOLIORFLIMSCICAVWPCGSRPCAVFCIF 237
HeELOVL5   VLVSYVCLSAVSMRPLWKRKYITGCOLIORVLTIIICSCGVWPCGSRPCAVFCIF 237
HeELOVL2   ILLSYVCLSVFSMRPLWKRKYITGCOLIORVLTIIIMSAVWPCGSRPCAVFCIF 240
SeElov12   VLVSYVCLSAVSMRPLWKRKYITGCOLIORVLTIIITLSAVWPCGSRPCAVFCIF 240
          * * * * *
          III
          IV
          V
SeElov15a  WLVVILALRSNFDLQTRKHLVSKK---ECHQNGSVASINGHVNGVTPTEI-IITHKVR 293
SeElov15b  WVASLALRSNFDLQTRKRVSKK---ETHQNGSVDSINGHANGVTPTEI-IITHKVR 292
HeELOVL5   WLVVILALRSNFDLQTRKSRKHLKQNGSMVAWNGHINSFPLENNVKKPLR 297
HeELOVL2   WLVVILALRSNFDLQTRKRMKKD---MCEPPACKEVNNGFSKAFFTAANGVAKR 296
SeElov12   WLVVILALRSNFDLQTRKRPDEE---IKSRP---NGHS---VSTNGTSRPAK 287
          .....
          V
SeElov15a  GD 295
SeElov15b  VD 294
HeELOVL5   KD 299
HeELOVL2   --
SeElov12   --

```

Fig 1




Fig 2




Fig 3




Fig 4




Fig 5