



Historical (1750–2014) anthropogenic emissions of reactive gases and aerosols from the Community Emissions Data System (CEDS)

Rachel M. Hoesly¹, Steven J. Smith^{1,2}, Leyang Feng¹, Zbigniew Klimont³, Greet Janssens-Maenhout⁴, Tyler Pitkanen¹, Jonathan J. Seibert¹, Linh Vu¹, Robert J. Andres⁵, Ryan M. Bolt¹, Tami C. Bond⁶, Laura Dawidowski⁷, Nazar Kholod¹, June-ichi Kurokawa⁸, Meng Li⁹, Liang Liu⁶, Zifeng Lu¹⁰, Maria Cecilia P. Moura¹, Patrick R. O'Rourke¹, and Qiang Zhang⁹

¹Joint Global Change Research Institute, Pacific Northwest National Lab, College Park, MD, USA

²Department of Atmospheric and Oceanic Science, University of Maryland, College Park, MD, USA

³International Institute for Applied Systems Analysis, Laxenburg, Austria

⁴European Commission, Joint Research Centre, Directorate Energy, Transport & Climate, Via Fermi 2749, 21027 Ispra, Italy

⁵Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, TN, USA

⁶Dept. of Civil & Environmental Engineering, University of Illinois at Urbana-Champaign, Urbana, IL, USA

⁷Comisión Nacional de Energía Atómica, Buenos Aires, Argentina

⁸Japan Environmental Sanitation Center, Asia Center for Air Pollution Research, Atmospheric Research Department, Niigata, Niigata, Japan

⁹Department of Earth System Science, Tsinghua University, Beijing, China

¹⁰Energy Systems Division, Argonne National Laboratory, Argonne, IL, USA

Correspondence: Rachel M. Hoesly (rachel.hoesly@pnnl.gov) and Steven J. Smith (ssmith@pnnl.gov)

Received: 20 February 2017 – Discussion started: 21 March 2017

Revised: 27 September 2017 – Accepted: 10 November 2017 – Published: 29 January 2018

Abstract. We present a new data set of annual historical (1750–2014) anthropogenic chemically reactive gases (CO, CH₄, NH₃, NO_x, SO₂, NMVOCs), carbonaceous aerosols (black carbon – BC, and organic carbon – OC), and CO₂ developed with the Community Emissions Data System (CEDS). We improve upon existing inventories with a more consistent and reproducible methodology applied to all emission species, updated emission factors, and recent estimates through 2014. The data system relies on existing energy consumption data sets and regional and country-specific inventories to produce trends over recent decades. All emission species are consistently estimated using the same activity data over all time periods. Emissions are provided on an annual basis at the level of country and sector and gridded with monthly seasonality. These estimates are comparable to, but generally slightly higher than, existing global inventories. Emissions over the most recent years are more uncertain, particularly in low- and middle-income regions where country-specific emission inventories are less available. Fu-

ture work will involve refining and updating these emission estimates, estimating emissions' uncertainty, and publication of the system as open-source software.

1 Introduction

Anthropogenic emissions of reactive gases, aerosols, and aerosol precursor compounds have substantially changed atmospheric composition and associated fluxes from land and ocean surfaces. As a result, increased particulate and tropospheric ozone concentrations since pre-industrial times have altered radiative balances of the atmosphere, increased human mortality and morbidity, and impacted terrestrial and aquatic ecosystems. Central to studying these effects are historical trends of emissions. Historical emission data and consistent emission time series are especially important for Earth systems models (ESMs) and atmospheric chemistry and transport models, which use emission time series as key

model inputs; integrated assessment models (IAMs), which use recent emission data as a starting point for future emission scenarios; and to inform management decisions.

Despite their wide use in research and policy communities, there are a number of limitations to current inventory data sets. Emission data from country- and region-specific inventories vary in methodology, level of detail, sectoral coverage, and consistency over time and space. Existing global inventories do not always provide comprehensive documentation for assumptions and methods, and few contain uncertainty estimates.

Several global emission inventories have been used in global research and modeling. The Emission Database for Global Atmospheric Research (EDGAR) is another widely used historical global emission data set. It provides an independent estimate of historical greenhouse gas (GHG) and pollutant emissions by country, sector, and spatial grid ($0.1 \times 0.1^\circ$) from 1970 to 2010 (Crippa et al., 2016; EC-JRC/PBL, 2016), with GHG emission estimates for more recent years. The most recent set of modeling exercises by the Task Force on Hemispheric Transport of Air Pollutants (TF HTAP) uses a gridded emission data set, HTAP v2 (Janssens-Maenhout et al., 2015), that merged EDGAR with regional and country-level gridded emission data for 2008 and 2010. The GAINS (Greenhouse Gas – Air Pollution Interactions and Synergies) model (Amann et al., 2011) has been used to produce regional and global emission estimates for several recent years (1990–2010; in 5-year intervals) together with projections to 2020 and beyond (Amann et al., 2013; Cofala et al., 2007; Klimont et al., 2009). These have been developed with substantial consultation with national experts, especially for Europe and Asia (Amann et al., 2008, 2015; Purohit et al., 2010; Sharma et al., 2015; Wang et al., 2014; Zhang et al., 2007; Zhao et al., 2013a). The newly developed ECLIPSE emission sets include several extensions and updates in the GAINS model and are also available in a gridded form (Klimont et al., 2017a) and have been used in a number of recent modeling exercises (Eckhardt et al., 2015; IEA, 2016b; Rao et al., 2016; Stohl et al., 2015).

Lamarque et al. (2010) developed a historical data set for the Coupled Model Intercomparison Project phase 5 (CMIP5), which includes global, gridded estimates of anthropogenic and open burning emissions from 1850 to 2000 at 10-year intervals. These data are also used as the historical starting point for the Representative Concentration Pathways (RCP) scenarios (van Vuuren et al., 2011) and in some research communities are referred to as the RCP historical data. In this article, these data are referred to as the CMIP5 data set. This was a compilation of “best available estimates” from many sources including EDGAR-HYDE (van Aardenne et al., 2001), which provides global anthropogenic emissions of carbon dioxide (CO_2), methane (CH_4), nitrous oxide (N_2O), nitrogen oxides (NO_x), non-methane volatile organic compounds (NMVOCs), sulfur dioxide (SO_2), and ammonia (NH_3) from 1890 to 1990 every 10 years at $1 \times 1^\circ$

grids; RETRO (Schultz and Sebastian, 2007), which estimated global emissions from 1960 to 2000; and emissions reported by, largely, the Organisation for Economic Cooperation and Development (OECD) countries over recent years. While this data set was an improvement upon the regional and country-specific inventories mentioned above, it lacks uncertainty estimates and reproducibility, has limited temporal resolution (10-year estimates to 2000), and does not have consistent methods across emission species. There are many existing inventories of various scope, coverage, and quality; however, no existing data set meets all the growing needs of the modeling community.

This paper describes the general methodology and results for an updated global historical emission data set that has been designed to meet the needs of the global atmospheric modeling community and other researchers for consistent long-term emission trends. The methodology was designed to produce annual estimates, be similar to country-level inventories where available, be complete and plausible, and use a consistent methodology over time with the same underlying driver data (e.g., fuel consumption). The data set described here provides a sectoral and gridded historical inventory of climate-relevant anthropogenic GHGs, reactive gases, and aerosols for use in the Coupled Model Intercomparison Project phase 6 (CMIP6). It does not include agricultural waste burning, which is included in van Marle et al. (van Marle et al., 2017). Gridded data were first released in summer 2016 through the Earth System Grid Federation (ESGF) system including SO_2 , NO_x , NH_3 , carbon monoxide (CO), black carbon (BC), organic carbon (OC), and NMVOCs, with a new release in May 2017 that corrected mistakes in the gridded data (links and details in Appendix Sects. A1 and A2). The May 2017 release also included CO_2 emissions (annual from 1750 to 2014) and CH_4 emissions (annual from 1970 to 2014 and a separate decadal historical extension from 1850 to 1970, also detailed in Appendix Sect. A2). This data set was created using the Community Emissions Data System (CEDs), which is being prepared for release as open-source software. Updated information on the system can be found at <http://www.globalchange.umd.edu/ceds/>.

An overview of the methodology and data sources is provided in Sect. 2, while further details on the methodology and data sources are included in the Supplement and outlined in Sect. 2.7. Section 3 compares this data set to existing inventories and Sect. 4 details future work involving this data set and system.

2 Data and methodology

2.1 Methodological overview

CEDs uses existing emission inventories, emission factors, and activity/driver data to estimate annual country-, sector-,

and fuel-specific emissions over time in several major phases (data system schematic shown in Fig. 1):

1. data are collected and processed into a consistent format and timescale (detailed in Sect. 2.2 and throughout the paper);
2. default emissions from 1960/1971 (1960 for most OECD countries and 1971 for all others) to 2014 are estimated using driver and emission factor data (emissions are equal to the driver multiplied by the emission factors) (Sect. 2.2);
3. default estimates are scaled to match existing emission inventories where available, complete, and plausible (Sect. 2.4);
4. scaled emission estimates are extended back to 1750 (Sect. 2.5) to produce final aggregate emissions by country, fuel, and sector;
5. emissions are checked and summarized to produce data for release and analysis; and
6. gridded emissions with monthly seasonality and volatile organic compound (VOC) speciation are produced from aggregate estimates using spatial proxy data (Sect. 2.6).

Rather than producing independent estimates, this methodology relies on matching default estimates to reliable, existing emission inventories (emission scaling) and extending those values to historical years (historical extension) to produce a consistent historical time series. While previous work (Lamarque et al., 2010) combined different data sets then smoothed over discontinuities, CEDS produces historical trends by extending the individual components (driver data and emission factors) separately to estimate emission trends. This method captures trends in fuel use, technology, and emission controls over time. Estimating emissions from drivers and emission factor components also allows the system to estimate emissions in recent years, using extrapolated emission factors and quickly released fuel use data, where detailed energy statistics and emission inventories are not yet available.

CEDS estimates emissions for 221 regions (and a global region for international shipping and aircraft), eight fuels, and 55 working sectors, summarized in Table 1. “Regions” refers to countries, regions, territories, or islands and are listed, along with mapping to summary regions and ISO codes in the Supplement files; they will henceforth be referred to as “countries”. CEDS working sectors (sectors 1A1–1A5) for combustion emissions follow the International Energy Agency (IEA) energy statistics sector definitions (Table A1). The IEA energy statistics are annually updated and the most comprehensive global energy statistics available, so this choice allows for maximal use of these data. Non-combustion emission sectors (sectors 1A1bc and 1B–7) are

drawn from EDGAR and generally follow EDGAR definitions (Table A2). Sector names were derived from Intergovernmental Panel on Climate Change (IPCC) reporting categories under the 1996 guidelines and Nomenclature for Reporting (NFR) 14 (Economic Commission for Europe, 2014) together with a short descriptive name¹. Note that CEDS data do not include open burning, e.g., forest and grassland fires, and agricultural waste burning on fields, which was developed by van Marle et al. (2017). Tables providing more detailed information on these mappings, which define the CEDS sectors and fuels, are provided in Sect. A3. We note that, while agriculture sectors include a large variety of activities, in practice, in the current CEDS system these sectors largely represent NH₃ and NO_x emissions from fertilizer application (under 3-D_Soil-emissions) and manure management, due to the focus in the current CEDS system on air-pollutant emissions.

In order to produce timely emission estimates for CMIP6, several CEDS emission sectors in this version of the system aggregate somewhat disparate processes to reduce the need for the development of detailed driver and emission factor information. For example, process emissions from the production of iron and steel, aluminum, and other non-ferrous metals are grouped together as an aggregate as 2C_Metal-production sector. Similarly, emissions from a variety of processes are reported in 2B_Chemical-industry. Also, the 1A1bc_Other-transformation sector includes emissions from combustion-related activities in energy transformation processes, including coal and coke production, charcoal production, and petroleum refining, but are combined in one working sector (see Sect. 2.3.2). Greater disaggregation for these sectors would improve these estimates but will require additional effort, described in Sect. 5.

The core outputs of the CEDS system are country-level emissions aggregated to the CEDS sector level. Emissions by fuel and by detailed CEDS sector are also documented within the system for analysis, although these are not released due to data confidentiality issues. Emissions are further aggregated and processed to provide gridded emission data with monthly seasonality, detailed in Sect. 2.6.

We note that the CEDS system does not reduce the need for more detailed inventory estimates. For example, CEDS does not include a representation of vehicle fleet turnover and emission control degradation (e.g., the effectiveness of catalytic converters over time) or multiple fuel combustion technologies that are included in more detailed inventories. The purpose of this system, as described further below, is to build on a combination of global emission estimation frameworks such as GAINS and EDGAR, combined with country-

¹Sector names were derived NFR14 nomenclature via a mapping table provided by the Centre on Emission Inventories and Projections (CEIP), available at http://www.ceip.at/ms/ceip_home1/ceip_home/reporting_instructions/




Figure 1. System summary. The key steps in calculation are to (1) collect and process activity, emission factors, and emission data; (2) develop default emission estimates; (3) calibrate default estimates to existing inventories; (4) extend present-day emission to historical time periods; (5) summarize emission outputs; and (6) produce data products including gridded emission and, in the future, uncertainty estimates.

level inventories, to produce reproducible, consistent emission trends over time, space, and emission species.

2.2 Activity data

Trends of energy consumption and other driver (activity) data are key inputs for estimating emissions. When choosing data to use in this system, priority was given to consistent trends over time rather than detailed data that might only be available for a limited set of countries or time span.

2.2.1 Energy data

Energy consumption data are used as drivers for emissions from fuel combustion. Core energy data for 1960–2013 are the International Energy Agency (IEA) energy statistics, which provide energy production and consumption estimates detailed by country, fuel, and sector from 1960 to 2013 for most OECD countries and 1971 to 2013 for non-OECD countries (IEA, 2015). While most data sources used in CEDS are open source, CEDS currently requires purchase of this proprietary data set. IEA data are provided at finer fuel and sector level so data are often aggregated to CEDS sectors and fuels. Mapping of IEA products to CEDS fuels is detailed in Sect. A4. Data for a number of small countries are provided by IEA only at an aggregate level, such as “Other Africa” and “Other Asia”, are disaggregated to CEDS countries using historical CO₂ emission data from the Carbon Dioxide Information Analysis Center (CDIAC) (Andres

et al., 2012; Boden et al., 1995). Sectoral splits for former Soviet Union (FSU) countries are smoothed over time to account for changes in reporting methodologies during the transition to independent countries (see the Supplement).

IEA energy statistics were extended to 2014 using the BP Statistical Review of World Energy (BP, 2015), which is freely available online and provides annual updates of country energy totals by aggregate fuel (oil, gas, and coal). BP trends for aggregate fuel consumption from 2013 to 2014 were applied to all CEDS sectors in the corresponding CEDS fuel estimates to extrapolate to 2014 energy estimates by sector and fuel from 2012 IEA values.

In a few cases, IEA energy data were adjusted to either smooth over discontinuities or to better match newer information. For international shipping, where a number of studies have concluded that IEA-reported consumption is incomplete (Corbett et al., 1999; Endresen et al., 2007; Eyring et al., 2010), we have added additional fuel consumption so that total consumption matches bottom-up estimates from the International Maritime Organization (IMO) (2014). For China, fuel consumption appears to be underestimated in national statistics (Guan et al., 2012; Liu et al., 2015b), so coal and petroleum consumption were adjusted to match the sum of provincial estimates as used in the MEIC inventory (Multi-resolution Emission Inventory for China) (Li et al., 2017) used to calibrate CEDS emission estimates. Several other changes were made, such as what appears to be spurious brown coal consumption over 1971–1984 in the IEA Other Asia region and a spike in agricultural diesel consumption

Table 1. CEDS working sectors and fuels (CEDS v2016-07-26). RCO indicates the “residential, commercial, other” sector.

CEDS working sectors		
Energy production	1A2g_Ind-Comb-other	RCO
1A1a_Electricity-public	2A1_Cement-production	1A4a_Commercial-institutional
1A1a_Electricity-autoproducer	2A2_Lime-production	1A4b_Residential
1A1a_Heat-production	2Ax_Other-minerals	1A4c_Agriculture-forestry-fishing
1A1bc_Other-transformation	2B_Chemical-industry	1A5_Other-unspecified
1B1_Fugitive-solid-fuels	2C_Metal-production	Agriculture
1B2_Fugitive-petr-and-gas	2-D_Other-product-use	3B_Manure-management
1B2d_Fugitive-other-energy	2-D_Paint-application	3-D_Soil-emissions
7A_Fossil-fuel-fires	2-D_Chemical-products-manufacture-processing	3I_Agriculture-other
Industry	2H_Pulp-and-paper-food-beverage-wood	3-D_Rice-Cultivation
1A2a_Ind-Comb-Iron-steel	2-D_Degreasing-Cleaning	3E_Enteric-fermentation
1A2b_Ind-Comb-Non-ferrous-metals	Transportation	Waste
1A2c_Ind-Comb-Chemicals	1A3ai_International-aviation	5A_Solid-waste-disposal
1A2d_Ind-Comb-Pulp-paper	1A3aii_Domestic-aviation	5E_Other-waste-handling
1A2e_Ind-Comb-Food-tobacco	1A3b_Road	5C_Waste-combustion
1A2f_Ind-Comb-Non-metallic-minerals	1A3c_Rail	5-D_Wastewater-handling
1A2g_Ind-Comb-Construction	1A3di_International-shipping	6A_Other-in-total
1A2g_Ind-Comb-transpequip	1A3di_Oil_tanker_loading	6B_Other-not-in-total
1A2g_Ind-Comb-machinery	1A3dii_Domestic-navigation	
1A2g_Ind-Comb-mining-quarrying	1A3eii_Other-transp	
1A2g_Ind-Comb-wood-products		
1A2g_Ind-Comb-textile-leather		
CEDS fuels		
Hard coal	Light oil	Natural gas
Brown coal	Diesel oil	Biomass
Coal coke	Heavy oil	

in Canada in 1984. All such changes are documented in the CEDS source code, input files, and the Supplement provided with this article.

Residential biomass was estimated by merging IEA energy statistics and Fernandes et al. (2007) to produce residential biomass estimates by country and fuel type over 1850–2013. Residential biomass data were reconstructed with the assumption that sudden drops in biomass consumption going back in time are due to data gaps, rather than sudden energy consumption changes. Both IEA and Fernandes et al. values were reconstructed to maintain smooth per capita (based on rural population) residential biomass use over time.

Details on methods and assumption for energy consumption estimates are available in the supplemental data and assumptions (see Sect. 3 of the Supplement).

2.2.2 Population and other data

Consistent historical time trends are prioritized for activity driver data. For non-combustion sectors, population is generally used as an activity driver. United Nations (UN) population data (UN, 2014, 2015) are used for 1950–2014, supplemented from 1960 to 2014 with World Bank population statistics (The World Bank, 2016). This series was merged with HYDE historical population data (Klein Gold-

ewijk et al., 2010). More details are available in Sect. 2.1 of the Supplement.

In this data version, population is used as the non-combustion emissions driver for all but three sectors. 5C_Waste-combustion, which includes industrial, municipal, and open waste burning, is driven by pulp and paper consumption, derived from Food and Agriculture Organization of the UN (FAO) Forestry Statistics (FAOSTAT, 2015). FAO statistics converted to per capita values were smoothed and linearly extrapolated backward in time. 1B2_Fugitive-petr-and-gas, which includes fugitive and flaring emissions from production of liquid and gaseous fuels together with oil refining, is driven by a composite variable that combines domestic oil and gas production with refinery inputs, derived from IEA energy statistics. This same driver is also used for 1B2d_Fugitive-other-energy. More details are available in Sect. 2.5 of the Supplement. While non-combustion emissions use population as an “activity driver” in calculations, emission trends are generally determined by a combination of EDGAR and country-level inventories. Final emission estimates, therefore, reflect recent emission inventories where these are available, rather than population trends.

2.3 Default estimates

Significant effort is devoted to creating reliable default emission estimates, including abatement measures, to serve as a starting point for scaling to match country-level inventories (Sect. 2.4) and historical extension back to 1750 (Sect. 2.5). While most default estimates do not explicitly appear in the final data set as they are altered to match inventories (Sect. 2.4), some are not altered because inventories are not available for all regions, sectors, and species. The method for calculating default emission factors varies by sectors and regions depending on available data.

Default emission estimates (box 2 in Fig. 1) are calculated using three types of data (box 1 in Fig. 1): activity data (usually energy consumption or population), emission inventories, and emission factors, according to Eq. (1).

$$E_{em}^{c, s, f, t} = A^{c, s, f, t} \times EF_{em}^{c, s, f, t}, \quad (1)$$

where E is total emissions, A is the activity or driver, EF is the emission factor, em is the emission species, c is the country, s is the sector, f is fuel (where applicable), and t is the year.

In general, default emissions for fuel combustion (sector 1A in Table 1) are estimated from emission factors and activity drivers (energy consumption), while estimates of non-combustion emissions (sectors 1B–7A and 1A1bc) are taken from a relevant inventory and the “implied emission factor” is inferred from total emissions and activity drivers.

2.3.1 Default fuel combustion emissions

Combustion sector emissions are estimated from energy consumption estimates (Sect. 2.2), and emission factors according to Eq. (1). Default emission factors for the combustion of fuels are derived from existing global data sets that detail emissions and energy consumption by sector and fuel, using Eq. (2):

$$EF_{em}^{c, s, f, t} = \frac{E_{em}^{c, s, f, t}}{A^{c, s, f, t}}, \quad (2)$$

where EF is the default emission factor, E is the total emissions as reported by other inventories, A is the activity data, measured in energy consumption as reported by inventories, em is the emission species, c is the country, s is the sector, f is fuel (where applicable), and t is the year.

The main data sets used to derive emission factors are shown in Table 2. Default emission factors for NO_x , NMVOCs, CO , and CH_4 are estimated from the global implementation of the GAINS model as released for the Energy Modeling Forum 30 project (<https://emf.stanford.edu/projects/emf-30-short-lived-climate-forcers-air-quality>) (Klimont et al., 2017a, b; Stohl et al., 2015). BC and OC emission factors from 1850 to 2000 are estimated from the latest version of the Speciated Pollutant Emission Wizard (SPEW) (Bond et al., 2007).

Emission factors for CO_2 emissions for coal and natural gas combustion are taken from CDIAC (Andres et al., 2012; Boden et al., 1995), with an additional coal mass balance check, as further described in Sect. 5.4 of the Supplement. For coal in China, a lower oxidation fraction of 0.96 was assumed; see discussion in the Supplement (Liu et al., 2015b). Because CEDS models liquid fuel emissions by fuel grade (light, medium, heavy), we use fuel-specific emission factors for liquid fuels also described in Sect. 5.4 of the Supplement.

Emission data are aggregated by sector and fuel to match CEDS sectors, while calculated emission factors from more aggregate data sets are applied to multiple CEDS sectors, fuels, or countries. When incomplete time series are available, emission factors are generally assumed constant back to 1970, linearly interpolated between data points, and extended forward to 2014 using trends from GAINS to produce a complete time series of default emission factors. Many of these interpolated and extended values are later scaled to match country inventories (Sect. 2.4).

Most of the default emission factors are derived from sources that account for technology efficiencies and mitigation controls over time, but some are estimated directly from fuel properties (e.g., fuel sulfur content for SO_2 emissions). A control percentage is used to adjust the emission factor in these cases. In the data reported here, the control percentage is primarily used in SO_2 calculations (see Sect. 5.1 of the Supplement) where the base emission factor is derived directly from fuel properties; however, this functionality is available when needed for other emission species. In most of these cases, emissions are later scaled to match inventory data.

2.3.2 Default non-combustion emissions

Default non-combustion emissions are generally taken from existing emission inventories, primarily EDGAR (EC-JRC/PBL, 2016) and some additional sources for specific sectors detailed in Table 2. Default emissions from sectors not specifically mentioned in Table 2 or the text below are taken from EDGAR (EC-JRC/PBL, 2016). Other data sources and detailed methods are explained in Sect. 6 of the Supplement. For detailed sector definitions, refer to Sect. A3.

When complete trends of emission estimates are not available, they are extended in a similar manner to combustion emissions: emission factors are inferred using Eq. (2) and (with few exceptions) using population as an activity driver; emission factors (e.g., per capita emissions) are linearly interpolated between data points and extended forward and back to 1970 and 2014 to create a complete trend of default emission factors; and default emission estimates are calculated using Eq. (1).

For this data set, all non-combustion sectors (except for 5C_Waste-combustion) use population as the activity driver since this provides a continuous historical time series to be used where interpolations were needed. In practice, since

Table 2. Data sources used to estimate default emission factors for fuel combustion and default emissions from non-combustion sectors.

Source sector	Emission species	Data source
Fuel combustion	(1A) NO _x , NMVOCs, CO, CH ₄ BC, OC SO ₂	GAINS energy use and emissions (Klimont et al., 2017a; Stohl et al., 2015) SPEW energy use and emissions (Bond et al., 2007) (Europe) GAINS sulfur content and ash retention (Amann et al., 2015; IASA., 2014a, b). Smith et al. (2011) and additional sources for other regions (Sect. 5.1 of the Supplement)
	NH ₃ CO ₂	US NEI energy use and emissions (US EPA, 2013) CDIAC (Boden et al., 2016) and additional data sources
Fugitive petroleum and gas (1B)	All	EDGAR emissions (EC-JRC/PBL, 2016), ECLIPSE V5a (Stohl et al., 2015)
Cement (2A1)	CO ₂	CDIAC (Boden et al., 2016)
Agriculture sectors (3)	CH ₄	For sectors 3B_Manure-management, 3B_Soil-emissions, and 3-D_Rice-Cultivation: FAOSTAT (FAO, 2016) All others: EDGAR emissions (EC-JRC/PBL, 2016)
	Other	EDGAR emissions (EC-JRC/PBL, 2016)
Waste combustion (5C)	All	(Akagi et al., 2011; Andreae and Merlet, 2001; Wiedinmyer et al., 2014) (Sect. 6.3 of the Supplement)
Waste water treatment (5-D)	NH ₃	CEDS estimate of NH ₃ from human waste (Sect. 6.4 of the Supplement)
Other non-combustion (2A–7A)	SO ₂	EDGAR (EC-JRC/PBL, 2016), Smith et al. (2011), and other sources (Sect. 6.5 of the Supplement)
	Other	EDGAR emissions (EC-JRC/PBL, 2016)

EDGAR is generally used for default non-combustion data sources, we are relying on EDGAR trends by country to extend emission data beyond years where additional inventory information does not exist (with exceptions as noted in Table 2). The pulp and paper sector uses pulp and paper consumption, detailed in Sect. 2.2; the waste combustion sector, which incorporates solid waste disposal (incineration) and residential waste combustion, which is the product of combustion, in this system, is methodologically treated as a non-combustion sector.

We note that, while emissions from sector 1A1bc_Other_transformation are also due to fuel combustion, due to the complexity of the processes included, this sector is treated as a non-combustion sector in CEDS in terms of methodology. This means that fuel is not used as an activity driver and that default emissions for this sector are taken from SPEW for BC and OC and EDGAR for other emissions. The major emission processes in this sector include coal coke production, oil refining, and charcoal production. A mass balance calculation for SO₂ and CO₂ focusing on coal transformation was also conducted to assure that these specific emissions were not underestimated, particularly for periods up to the mid-20th century (Sects. 5.4, 6.5.2, and 8.3.2 of the Supplement).

During the process of emission scaling, we found that default emissions were sometimes 1–2 orders of magnitude

different from emissions reported in national inventories. This is not surprising, since non-combustion emissions can be highly dependent on local conditions, technology performance, and there are also often issues of incompleteness of inventories. In these cases, we implemented a process whereby default non-combustion emissions were taken directly from national inventories, and gap-filled and trended over time using EDGAR estimates. These were largely fugitive and flaring emissions (1B) for SO₂; soil (3-D), manure (3B), and waste water (5-D) emissions for NH₃; and non-combustion emissions for NMVOCs, typically associated with solvent use.

2.4 Scaling emissions

CEDS uses a “mosaic” strategy to scale default emission estimates to authoritative country-level inventories when available. The goal of the scaling process is to match CEDS emission estimates to comparable inventories while retaining the fuel and sector detail of the CEDS estimates. The scaling process modifies CEDS default emissions and emission factors, but activity estimates remain the same.

A set of scaling sectors is defined for each inventory so that CEDS and inventory sectors overlap. These sectors are chosen to be broad, even when more inventory details are available, because it is often unclear if sector definitions and boundaries are comparable between data sets. For example, many inventories do not consistently break out industry auto-producer electricity from other industrial combustion, so they are combined together for scaling. Additionally, underlying driver data in inventories and CEDS may not match. Scaling detailed sectors that were calculated using different energy consumption estimates would yield unrealistic scaled emission factors at a detailed sector level. One example is off-road emissions; while often estimated in country inventories, energy consumption data at this level are not consistently available from the IEA energy statistics, so these emissions are combined into broader sector groupings, depending on the sector categories available in a specific inventory.

The first step in this process is to aggregate CEDS emissions and inventory emissions to common scaling sectors; then scaling factors are calculated with Eq. (3). Scaling factors represent the ratio between CEDS default estimates and scaling inventory estimates by scaling sector and provide a means for matching CEDS default estimates to scaling inventories.

$$SF_{em}^{c, ss, t} = \frac{Inv_{em}^{c, ss, t}}{CEDS_{em}^{c, ss, t}}, \quad (3)$$

where SF is the scaling factor, Inv is the inventory emissions estimate, CEDS is the CEDS emissions estimate, em is the emission species, c is the country, ss is the aggregate scaling sector (unique to inventory), and t is the year.

For each inventory, scaling factors are calculated for years when inventory data are available. Calculated scaling factors are limited to values between 1/100 and 100. Scaling factors outside this range may result from discontinuities or misreporting in inventory data; imperfect scaling maps between CEDS sectors, inventory sectors, and scaling sectors; or default CEDS emission estimates that are drastically different than reported inventories. Many of these cases were resolved by using the detailed inventory data as default emission data, as noted above in Sect. 2.3.2. Where inventory data are not available over the specified scaling time frame, remaining scaling factors are interpolated and extended to provide a continuous trend. Scaling factors are applied to corresponding CEDS default emission estimates and default emission factors to produce a set of scaled emission components (total emissions and emission factors, together with activity drivers, which are not changed), which are used in the historical extension (Sect. 2.5). Using scaling factors retains the sector and fuel level detail of CEDS default emission estimates, while matching total values to authoritative emission inventories.

We use a sequential methodology in which CEDS values are generally first scaled to EDGAR (EC-JRC/PBL, 2016) for most emission species, then national inventories, where

available. Final CEDS results, over the period these inventories were available, match the last inventory scaled. SO₂, CH₄, BC, and OC are not scaled to EDGAR values. For all pollutant species other than BC and OC, estimates are then scaled to match country-level emission estimates. These are available for most of Europe through the European Monitoring and Evaluation Programme (EMEP) for European countries after 1980 (EMEP, 2016); the United Nations Framework Convention on Climate Change (UNFCCC) GHG data for Belarus, Greece, and New Zealand (UNFCCC, 2015) after 1990; an updated version of the Regional Emissions Inventory in Asia (REAS) for Japan (Kurokawa et al., 2013a); MEIC for China (Li et al., 2017); and others detailed in Table 3. BC and OC emission estimates are entirely from default estimates calculated using predominantly SPEW data. While BC inventory estimates were available in a few cases, OC estimates were less available, so we have retained the consistent BC and OC estimates from SPEW for all countries. CH₄ emission estimates are scaled to match to the following inventories: EDGAR 4.2 (EC-JRC/PBL, 2012), UNFCCC submissions (UNFCCC, 2015) for most “Annex I” countries, and the US GHG inventory (US EPA, 2012b) for the United States.

The scaling process was designed to allow for exceptions when there are known discontinuities in inventory data or when the default scaling options resulted in large discontinuities. For example, former Soviet Union countries were only scaled to match EDGAR and other inventories after 1992 (where energy data become more consistent). Romania, for example, was only scaled to match EDGAR in 1992, 2000, and 2010 to avoid discontinuities. For the most part, these exceptions occur for countries with rather limited penetration of control measures or only low efficiency controls. Regions with more stringent emission standards requiring extensive application of high-efficiency controls have typically higher quality national inventories, e.g., the European Union, North America, and parts of Asia.

Description of the exceptions and assumptions for scaling inventories, as well as a detailed example of the scaling process, is available in Sect. 7 of the Supplement. Additionally, figures showing stacked area graphs of global emission, by final scaling inventory (or default estimate) are shown in Figs. S44–S55 in the supplement figures and tables. These show the percentage of final global emission estimates that are scaled to various inventories.

The scaling process operates on sectors where emissions are present in both the CEDS default data and the scaling inventories listed in Table 3. If the scaling inventory does not contain information for a particular sector, then the default data are used. This means that some gaps in the scaling inventories are automatically filled by this procedure and, as a result, the CEDS emission totals can be larger than those in the scaling inventory. For example, waste burning and fossil fuel fires are not included in some of the inventories, while these sectors are included in CEDS. In a few cases, specific

Table 3. Data sources for inventory scaling. All countries are scaled first to EDGAR and then to individual estimates.

Region/country	Species	Years	Data source
All, where available	NO _x , NMVOCs, CO, NH ₃ CH ₄	1970–2008	EDGAR v4.3 (EC-JRC/PBL, 2016)
		1970–2008	EDGAR v4.2 (EC-JRC/PBL, 2012)
Europe	SO ₂ , NO _x , NMVOCs, CO, NH ₃	1980–2012	(EMEP, 2016)
Greece, New Zealand, Belarus	SO ₂ , NO _x , NMVOCs, CO, CO ₂	1990–2012	(UNFCCC, 2015)
Other Asia	SO ₂ , NO _x , NMVOCs, CO, CH ₄	2000–2008	REAS 2.1 (Kurokawa et al., 2013a)
Argentina	SO ₂ , NO _x , NMVOCs, CO	1990–1999, 2001–2009, 2011	(Argentina UNFCCC Submission, 2016)
Australia	SO ₂ , NO _x , NMVOCs, CO	2000, 2006, 2012	(Australian Department of the Environment, 2016)
China	SO ₂ , NO _x , NMVOCs, CO, NH ₃	2008, 2010, 2012	MEIC (Li et al., 2017)
Canada	SO ₂ , NO _x , NMVOCs, CO	1985–2011	(Environment and Climate Change Canada, 2016; Environment Canada, 2013)
Japan	SO ₂ , NO _x , NMVOCs, CO, NH ₃	1960–2010	Preliminary update of Kurokawa et al. (2013b)
South Korea	SO ₂ , NO _x , NMVOCs, CO	1999–2012	(South Korea National Institute of Environmental Research, 2016)
Taiwan	SO ₂ , NO _x , NMVOCs, CO	2003, 2006, 2010	(TEPA, 2016)
USA	SO ₂ , NO _x , NMVOCs, CO NH ₃ CO ₂ CH ₄	1970, 1975, 1980, 1985, 1990–2014	EPA trends (US EPA, 2016b)
		1990–2014	
		1990–2014	US EPA (2016a)
		1990–2014	US GHG inventory (US EPA, 2012b)

additional data were added where gaps were known to be present. For example, the CEDS totals for China are slightly larger than the MEIC totals due to both the inclusion of waste burning and the addition of SO₂ emissions from metal smelting, which are not included in MEIC. Where necessary, discontinuities in inventory estimates were eliminated. For the USA, for example, discontinuities were present in the original EPA trend data due to methodological changes, particularly for transportation NO_x and agricultural NH₃.

2.5 Pre-1970 emissions extension

Historical emission and energy data before 1970 generally do not have the same details as more modern data. In general, we extend activity and emission factors back in time separately, with time- and sector-specific options to capture changes in technologies, fuel mixes, and activity. This allows for consistent methods across time and sectors, rather than piecing together different sources and smoothing over discontinuities, which was done in previous work (Lamarque et al., 2010). For most emission species and sectors, the assumed historical

trend in activity data has a large impact on emission trends. Activity for many sectors and fuels, such as fossil liquid and gas fuels, is small or zero by 1900. Some cases where emission factors are known to have changed over time have also been incorporated.

2.5.1 Pre-1970 activity drivers

IEA energy statistics, which are the foundation for energy estimates in this data set, go back to 1960 at the earliest. Fossil fuels are extended using CDIAC emissions, SPEW energy data, and assumptions about fuel type and sector splits in 1750, 1850, and 1900, detailed in Sect. 8.1 of the Supplement. First total fuel use for three aggregate fossil fuel types (coal, oil, and gas) is estimated over 1750–1960/70 for each country using historical national CO₂ estimates from the CDIAC (Andres et al., 1999; Boden et al., 2016).

For coal only, these extended trends were matched with SPEW estimates of total coal use, which are a composite of UN data (UN, 2016) and Andres et al. (1999). This resulted in a more accurate extension for a number of key countries. SPEW estimates for every 5 years were interpolated to annual values using CDIAC CO₂ time series, resulting in an annual time series. For coal and petroleum, aggregate fuel use was disaggregated into specific fuel types (e.g., brown coal, hard coal, and coal coke; light, medium, and heavy oil) by smoothly transitioning between fuel splits by aggregate sector from the IEA data to SPEW fuel type splits in earlier time periods. Finally, fuel use was disaggregated into sectors in a similar manner, smoothly transitioning between CEDS sectoral splits in either 1970 or 1960 to SPEW sectoral splits by 1850. A number of exogenous assumptions about fuel and sector splits over time were also needed in this process. More details on this method can be found in Sect. 8.1.1 of the Supplement.

While most biomass fuels are consumed in the residential sector, whose estimation was described above (Sect. 2.2.1), biomass consumed in other sectors is extended using SPEW energy data and population. The 1970 CEDS estimates of biomass used in industrial sectors are merged to SPEW values by 1920. Biomass estimates from 1750 to 1850 are estimated by assuming constant per capita values.

Activity drivers for non-combustion sectors in modern years are primarily population estimates. Most historical drivers for non-combustion sectors are also population, while some, shown in Table 4, are extended with other data. These are mostly sectors related to chemicals and solvents that are extended with CO₂ trends from liquid fuel use. Waste combustion is estimated by historical trends for pulp and paper consumption. The driver for sectors 1B2 and 1B2d, refinery and natural gas production, is extended using CDIAC CO₂ emissions for liquid and gas fuels.

2.5.2 Pre-1970 emission factors

In 1850, the only fuels are coal and biomass used in residential, industrial, rail, and international shipping sectors, and many non-combustion emissions are assumed to be zero. Emission factors are extended back in time by converging to a value in a specified year (often 0 in 1850 or 1900), remaining constant, or following a trend. For some non-combustion emissions, we use an emission trend instead of an emission factor trend. Ideally, sector-specific activity drivers would extend to zero, rather than emission factors; however, we often use population as the activity driver, because of the lack of complete, historical trends. Extending the emission factor (e.g., the per capita value) to zero approximates the decrease to zero in the actual activity.

BC and OC emission factors for combustion sectors were extended back to 1850 by sector and fuel using the SPEW database and held constant before 1850. Combustion emission factors for NO_x, NMVOCs, and CO in 1900 are drawn from a literature review, primarily Winijkul et al. (2016). These emission factors were held constant before 1900 and linearly interpolated between 1900 and 1970. Additional data sources and details are available in Sect. 8.2 of the Supplement.

Many non-combustion emissions were trended back with existing data from the literature. These include trends from SPEW (Bond et al., 2007), CDIAC (Boden et al., 2016), sector-specific sources such as SO₂ smelting and pig iron production, and others, detailed in Table 5. Emission factors for remaining sectors were linearly interpolated to zero in specified years based on a literature review (Bond et al., 2007; Davidson, 2009; Holland et al., 2005; Smith et al., 2011). Further methods and data sources are found in Sect. 8.3 of the Supplement.

NH₃ and NO_x emissions from minerals and manure (3B_Manure-management and 3-D_Soil-emissions) are grouped together. While CEDS total estimates should be reliable, there might be inconsistencies going back in time. We assume that the dominant trend from 1960 to 1970 is mineral fertilizer, then scaled back in time globally using Davidson et al. (2009).

2.6 Gridded emissions

Final emissions are gridded to facilitate use in Earth system, climate, and atmospheric chemistry models. Gridded outputs are generated as CF-compliant NetCDF files (<http://cfconventions.org/>). Aggregate emissions by country and CEDS sector are aggregated to 16 intermediate sectors (Table 6) and downscaled to a 0.5° × 0.5° grid. Country-aggregate emissions by intermediate gridding sector are spatially distributed using normalized spatial proxy distributions for each country, plus global spatial proxies for shipping and aircraft, then combined into global maps. For grid cells that contain more than one country, the proxy spatial distributions are ad-

Table 4. Historical driver extensions for non-combustion sectors.

Non-combustion sector	Modern activity driver	Historical extension trend
1B2_Fugitive-petr-and-gas	Refinery and natural gas production	CDIAC – liquid and gas fuels CO ₂
1B2d_Fugitive-other-energy	Refinery and natural gas production	CDIAC – liquid and gas fuels CO ₂
2B_Chemical-industry	Population	CDIAC – liquid fuels CO ₂
2-D_Degreasing-Cleaning	Population	CDIAC – liquid fuels CO ₂
2-D_Paint-application	Population	CDIAC – liquid fuels CO ₂
2-D3_Chemical-products-manufacture-processing	Population	CDIAC – liquid fuels CO ₂
2-D3_Other-product-use	Population	CDIAC – liquid fuels CO ₂
2L_Other-process-emissions	Population	CDIAC – liquid fuels CO ₂
5C_Waste-combustion	Pulp and paper consumption	Pulp and paper consumption
7A_Fossil-fuel-fires	Population	CDIAC – cumulative solid fuels CO ₂
All other process sectors	Population	Population

justed to be proportional to area fractions of each country occupying that cell. Gridded emissions are aggregated to nine sectors for final distribution: agriculture, energy, industrial, transportation, residential/commercial/other, solvents, waste, international shipping, and aircraft (shown in Table 6; more details can be found in Sect. 9.1 of the Supplement).

Proxy data used for gridding are primarily gridded emissions from EDGAR v4.2 (EC-JRC/PBL, 2012) and HYDE population (Goldewijk et al., 2011). Flaring emissions use a blend of grids from EDGAR and ECLIPSE (Klimont et al., 2017a). Road transportation uses the EDGAR v4.3 road transportation grid, which is significantly improved over previous versions (EC-JRC/PBL, 2016), but was only available for 2010, so this is used for all years. When the primary proxy for a specific country/region, sector, and year combination is not available, CEDS uses gridded population from Gridded Population of the World (GPW) (Doxsey-Whitfield et al., 2015) and HYDE as backup proxy. Whenever available, proxy data are from annual gridded data; however, proxy grids for sectors other than RCO (residential, commercial, other) and waste are held constant before 1970 and after 2008. Specific proxy data sources are detailed in Table 6. As noted above, these proxy data were used to distribute emissions spatially within each country such that country totals match the CEDS inventory estimates. More details on gridding can be found in Sect. 9 of the Supplement.

Emissions are aggregated to nine final gridding sectors (Table 6) and distributed over 12 months using spatially explicit, sector-specific monthly fractions, largely from the ECLIPSE project, except for international shipping (from EDGAR) and aircraft (from Lee et al. (2009), as used in Lamarque et al., 2010). Emissions are then converted to flux ($\text{kg m}^{-2} \text{s}^{-1}$). This process is further described in Sect. 9.4 of the Supplement.

2.7 Additional methodological details

The above sections discuss the general approach to the methodology used in producing this data set, but there are

a number of exceptions, details on additional processing and analysis, and data sources that are discussed in the Supplement files.

3 Results and discussion

3.1 Emission trends

Figures 2 and 3 show global emissions over time by aggregate sector and region, respectively, from 1750 to 2014. Definitions of aggregate sectors and regions are given in Sect. A of the supplemental figures and tables. Section B of the Supplement contains line graph versions of these figures, emissions by fuel, and regional versions of Figs. 2 and 3.

In 1850, the earliest year in which most existing data sets provide estimates, anthropogenic emissions are dominated by residential sector cooking and heating, and therefore products of incomplete combustion for BC, OC, CO, and NMVOCs. In 1850, anthropogenic emissions (sectors included in this inventory) made up approximately 20–30 % of total global emissions (which also include grassland and forest burning, estimated by Lamarque et al., 2010) for BC, OC, NMVOCs, and CO but only 3 % of global NO_x emissions.

In the late 1800s through the mid-20th century, global emissions transitioned to a mix of growing industrial, energy transformation and extraction (abbreviated as “Energy Trans/Ext”), and transportation emissions with a relatively steady global base of residential emissions (primarily biomass and later coal for cooking and heating). The 20th century brought a strong increase in emissions of pollutants associated with the industrial revolution and development of the transport sectors (SO₂, NO_x, CO₂, NMVOCs). BC and OC exhibit steadily growing emissions dominated by the residential sector over the century, while other sectors begin to contribute larger shares after 1950. The last few decades increasingly show, even at the global level, the impact of strong growth of Asian economies (Fig. 3). The Haber–Bosch process (ammonia synthesis) about 100 years ago allowed fast growth in agricultural production, stimulat-

Table 5. Historical extension method and data sources for emission factors.

Sector	Emission species	Extension method	Data source
All combustion sectors	NMVOCS, CO, NO _x	Interpolate to value in 1900	Detailed in the Supplement (Sect. 8.2.1)
All combustion sectors	BC, OC	EF trend	SPEW
2Ax_Other-minerals, 2-D_Degreasing-Cleaning, 2-D_Paint-application, 2-D3_Chemical-products- manufacture-processing, 2-D3_Other-product-use, 2H_Pulp-and-paper-food- beverage-wood, 2L_Other-process- emissions, 5A_Solid-waste-disposal, 5C_Waste-combustion, 5E_Other-waste-handling, 7A_Fossil-fuel-fires	All	Interpolate to zero in specified year (EFs are emissions per capita values)	Detailed in Sect. 8.3.1 of the Supplement
5-D_Wastewater-handling	NH ₃	Interpolate to value in specified year	
3B_Manure-management	NH ₃ , NO _x	EF trend Emissions trend	Manure nitrogen per capita (Holland et al., 2005) See Sect. 8.3.1 of the Supplement
3-D_Soil-emissions	NH ₃ , NO _x	EF trend Emissions trend	1961–1970: emissions trend using total nitrogen (N) fertilizer by country 1860–1960: per capita emissions scaled by global N fertilizer (Davidson, 2009) See Sect. 8.3.1 of the Supplement
1A1a_Electricity-public, 1A1a_Heat-production, 1A2g_Ind-Comb-other, 1A3c_Rail, 1A4a_Commercial- institutional, 1A4b_Residential	SO ₂	EF trend	(Gschwandtner et al., 1986)
1A1bc_Other- transformation	BC, OC	Emissions trend	Pig iron production (SPEW, USGS, other)
1A1bc_Other- transformation	Others	Emissions trend	Total fossil fuel CO ₂ (CDIAC)
2A1_Cement-production, 2A2_Lime-production	All	Emissions trend	CDIAC cement CO ₂
2C_Metal-production	SO ₂	Emissions trend	Smith et al. (2011) emissions
2C_Metal-production	CO	Emissions trend	Pig iron production
2C_Metal-production	Others	Emissions trend	CDIAC solid fuel CO ₂

ing population growth and a consequent explosion of NH₃ emissions (Erisman et al., 2008). Before 1920, global emis-

sions for all species were less than 10 % of the year 2000 global values.

Table 6. Proxy data used for gridding.

CEDS final gridding sector	CEDS intermediate gridding sector definition	Proxy data source	Years
Residential, commercial, other (RCO)	Residential, commercial, other (residential and commercial)	HYDE population (decadal values, interpolated annually)	1750–1899
		EDGAR v4.2 (1970) blended with HYDE population	1900–1969
		EDGAR v4.2 RCORC	1970–2008
	Residential, commercial, other (other)	HYDE population (decadal values, interpolated annually)	1750–1899
		EDGAR v4.2 (1970) blended with HYDE Population	1900–1969
		EDGAR v4.2 RCOO	1970–2008
Agriculture (AGR)	Agriculture	EDGAR v4.2 AGR	1970–2008
Energy sector (ENE)	Electricity and heat production	EDGAR v4.2 ELEC	1970–2008
	Fossil fuel fires	EDGAR v4.2 FFFI	1970–2008
	Fuel production and transformation	EDGAR v4.2 ETRN	1970–2008
	Oil and gas fugitive/flaring	ECLIPSE FLR 1990, 2000, 2010 EDGAR v4.2 ETRN (1970–2008)	1970–2010
Industrial sector (IND)	Industrial combustion	EDGAR v4.2 INDC	1970–2008
	Industrial process and product use	EDGAR v4.2 INPU	1970–2008
Transportation section (TRA)	Road transportation	EDGAR v4.3 ROAD (2010)	1750–2014
	Non-road transportation	EDGAR v4.2 NRTR	1970–2008
International shipping (SHP)	International shipping	ECLIPSE and additional data (1990–2015)	1990–2010
	International shipping (tanker loading)	ECLIPSE and additional data (1990–2015)	1990–2010
Solvent production and application (SLV)	Solvent production and application	EDGAR v4.2 SLV	1970–2008
Waste (WST)	Waste	HYDE population, GPW v3 (modified rural population)	1750–2014
Aircraft (AIR)	Aircraft	CMIP5 (Lamarque et al., 2010; Lee et al., 2009)	1850–2008

* Spatial proxy data within each country are held constant before and after the years shown. See the Supplement for further details on the gridding proxy data including definitions for the EDGAR gridding codes in this table.

For several decades after 1950, global emissions grew quickly for all species. SO₂ continued to be dominated by industry and energy transformation and extraction sectors. In the later parts of the century, while Europe and North American SO₂ emissions declined as a result of emission control policies, SO₂ emissions in Asia continued to grow. NH₃ was dominated by the agriculture sectors and NMVOCs by industry and energy transformation and extraction sectors. Trans-

portation emissions have grown steadily and became an important contribution to NO_x, NMVOCs, and CO emissions. Growth in CO emissions over the century is due to transportation emissions globally until the 1980s and 1990s when North America and Europe introduced catalytic converters. Other regions followed more recently, resulting in a declining transport contribution; however, CO emissions in Asia and Africa have continued to rise due to population-driven res-




Figure 2. CEDEs emission estimates by aggregate sector compared to Lamarque et al. (2010) (dots) and CDIAC (line) for CO₂. For a like-with-like comparison, these figures do not include aviation or agricultural waste burning on fields. “RCO” stands for residential, commercial, and other.

identical biomass burning. Similarly, while NO_x from transportation sectors has decreased in recent years, total global NO_x emissions have increased quickly since 2005 due to industry and energy sectors in all parts of Asia. BC and OC increases since 1950 have been dominated by residential emissions from Africa and Asia, but growing fleets of diesel vehicles in the last decades added to the burden of BC emissions.

BC emissions from residential biomass are shown in Fig. 4 alongside rural population by region. Other Asia, Africa, and China dominate residential biomass BC emissions, which are regions with the largest rural populations. While residential biomass in most regions follow rural population trends, emissions in Latin America stay flat as its rural population has steadily increased since 1960. Emissions in China flatten




Figure 3. Emission estimates by region compared to Lamarque et al. (2010) (dots) and CDIAC (line) for CO₂. For a like-with-like comparison, these figures do not include aviation or agricultural waste burning on fields. The “International” region shows international shipping emissions.

more dramatically after 1990 than rural population, presumably reflecting the spread of modern energy sources as rural residential per capita biomass use decreases in this data set.

Of the emission species estimated, SO₂ is the most responsive to global events such as war and depressions. SO₂ emissions are primarily from non-residential fuel burning and industrial processes which vary with economic activity, where

other species have a base of residential biomass burning or agriculture and waste emissions. In this data set, these emissions remain steady within the backdrop of variable economic conditions, while events such as World Wars or the collapse of the Soviet Union can be seen most clearly in annual SO₂ emissions. We note that the relative constancy of residential and agricultural emissions is, to some extent, a re-

sult of a lack of detailed time series data for the drivers of these emissions in earlier periods. Variability for these sectors in earlier years, therefore, might be underestimated.

3.2 Emission trends in recent years (2000–2014)

After 2000, many species' emissions follow similar trends as the late 20th century, as shown in Fig. 5, with further details in the supplemental figures (Sects. C, E, and G).

BC and OC steadily grow in Africa and Other Asia from residential biomass emissions, which are driven by continued growth of rural populations. While most BC emission growth in China is due to energy transformation, primarily coke production, the residential, transportation, industry, and waste sectors all contribute smaller but similar growth over 2000–2014 (Fig. S19). See Sect. 3.4 for a discussion of uncertainty.

NH₃ continues its steady increase mostly due to agriculture in Asia and Africa. Global CO₂ emissions increase due to steadily rising emissions across most sectors in China and Asia and moderately rising emissions in Africa and Latin America, while emissions in North America and Europe flatten or decline after 2007 (largely due to the energy transformation and extraction sectors).

Global CO emissions flatten, despite increasing CO emissions in China and Other Asia, and Africa, which is offset by a continuing decrease of transportation CO emissions in North America and Europe, shown in Fig. 2 and in more detail in the supplemental figures. CO emissions in China increase then flatten after 2007, despite continually decreasing transportation CO emissions, which are offset by an increase in industrial emissions (Fig. S19). Similarly, after an increase from 2000–2005, global SO₂ emissions flatten despite increasing emissions in China and Other Asia due to steadily decreasing emissions in Europe, North America, and the former Soviet Union (Figs. 2 and S3). SO₂ emissions from energy transformation in China have declined since 2005 with the onset of emission controls in power plants; however, industrial emissions remained largely uncontrolled and became the dominant sector in China (Fig. S19).

Global NO_x emissions rise and then flatten around 2008. The growth in industrial emissions after 2000 is offset in 2007 by the decrease in international shipping emissions, while global emissions in other sectors stay flat. NO_x emissions in North America and Europe decline due to transportation and energy transformation (Simon et al., 2015), while emissions in China and Other Asia continue to grow, also in the transportation and energy transformation. Growth of NO_x emissions in Other Asia almost completely offset reductions in NO_x emissions in North America from 2000–2014. In China, industry has continually grown since 2003, transportation began to flatten around 2007, and the energy transformation and extraction sectors began declining in 2011 (Fig. S19) following the introduction of more stringent emission standards for power plants (Liu et al., 2016).

Globally, NMVOC emissions increase over the period, due to varying developments across the regions but in large part due to increases in energy emissions. NMVOC emissions increase in China from solvents (Fig. S19), Other Asia from transportation (Fig. S24), and Africa from energy transformation (Fig. S18); they decline in Europe and North America due to transportation and solvents (Figs. S20 and S23), and stay flat in other regions.

As discussed in Sect. 3.5, trends in recent years are more uncertain as they rely on sometimes preliminary activity data and emission factors extended outside inventory scaling years. Some of the notable trends in CEDS emission estimates in recent years are also from particularly uncertain sources. OC and BC emission estimates have some of the highest degrees of uncertainty in global inventories, and waste sectors in particular are highly uncertain. Additionally, a lot of global growth can be attributed to sectors that, in the CEDS system, follow population trends over the most recent few years (e.g., waste, agriculture, and residential biomass); are from inherently uncertain sectors (e.g., waste); or are located in China where emissions remain uncertain because the accounting of emission factors, fuel properties, and energy use data have been subject to corrections and subsequent debate (Hong et al., 2017; Korsbakken et al., 2016; Liu et al., 2015b; Olivier et al., 2015).

3.3 Gridded emissions

Figure 6 shows gridded CEDS estimates of total emissions in 2010 for all emission species. CEDS maps are similar to existing maps such as EDGAR (EC-JRC/PBL, 2012) and CMIP5 (Lamarque et al., 2010) as these data sets are used in the gridding process. Emissions for most species are concentrated in high-population areas such as parts of China, India, and the eastern US. BC and OC, whose emissions are dominated by heating and cooking fueled by biomass are also more concentrated in Africa. Shipping emissions are concentrated along ocean shipping lanes for NO_x, SO₂, and CO₂. Discussion of how gridded data differ from CMIP5 (Lamarque et al., 2010) gridded data is included in Sect 3.4.1.

3.4 Comparison with other inventories

Differences between CEDS emissions and other inventory estimates are described below. The reasons depend on emission species but are largely due to updated emission factors, increased detail in fuel and sector data, and a new estimate of waste emissions (however, see Sect. 3.5).

3.4.1 CMIP5 (Lamarque et al., 2010)

The emission data used for CMIP5 (Lamarque et al., 2010) also used a “mosaic” methodology, combining emission estimates from different sources. The CEDS methodology provides a more consistent estimate over time since driver data are used to produce consistent trends. Emissions in earlier




Figure 4. (a) BC residential biomass emissions by region and (b) rural population by region.

years, particularly before 1900, also differ because CEDS differentiates between biomass and coal combustion, which have a large impact on CO and NO_x emissions. The Lamarque et al. (2010) estimates for early years were drawn from the EDGAR-HYDE estimates (van Aardenne et al., 2001), which did not distinguish between these fuels. Figures showing comparisons between CMIP5 and CEDS globally by sector and for the top five emitting CMIP5 regions are shown in Sect. H of the supplemental figures and tables.

CEDS global SO₂ estimates are similar to CMIP5 estimates, although slightly lower (~ 10 %) in the mid-20th century and slightly higher (~ 5 %) near the end of the 20th century. Similar methods and data were used to develop both estimates (Smith et al., 2011). FSU SO₂ emissions are larger in CEDS (see Smith et al., 2011) from 1970 to 2000 but smaller in Europe from 1930 to 1980. Shipping SO₂ emissions are lower in the early 20th century due to updated methodologies (Smith et al., 2011) and slightly lower in recent years due to updated parameter estimates (see the Supplement and Fig. S43).

CEDS NO_x emissions are smaller than the CMIP5 estimates until the mid-20th century. This is largely because of explicit representation of the lower NO_x emissions from biomass fuels in early periods, which combusts at lower temperatures as compared to coal. In 1970, CEDS NO_x emissions began to diverge from CMIP5 estimates, generally becoming larger due to waste, transportation, and energy sectors. CEDS emissions remain about 10 % larger than those of CMIP5 in 1980 and 1990. Both global estimates increase and start to flatten around 1990. However, CEDS values flatten until 2000 and then increase again, while CMIP5 values decrease from 1990 to 2000.

CEDS CO estimates before 1960 are increasingly larger than CMIP5 estimates going back in time, reaching a factor of 2 by 1850 due to the explicit representation of biomass. In 1900, CEDS estimates were 70 % larger than those of CMIP5, 98 % of which is due to the RCO sector. CEDS estimates are slightly larger than those of CMIP5 after 1960 (8 % in 1960 and 1970 and less than 5 % from 1980 to 2000).

CEDS OC estimates are within 10 % but smaller than CMIP5 estimates through 1970, when CEDS estimates quickly increase and become larger (at most 25 % larger) than CMIP5 estimates. BC emissions are similar, although CEDS estimates are smaller (sometimes by 25 %) than those of CMIP5 until 1960 when CEDS estimates increase quickly, up to 25 % larger than CMIP5 estimates, in part due to larger waste sector emissions (see Sect. 3.5). Differences in BC in the early 20th century are mostly from residential fuel use in the US. In 1910, 98 % of the difference between the two inventories was from residential energy use, with 77 % of that difference in the USA. US residential biomass consumption in 1949 is estimated using the Energy Information Administration (EIA) data and propagated back in time to merge with Fernandes et al. (2007) used by SPEW in 1920. This US biomass estimate may be lower than that used in CMIP5.

NH₃ and NMVOC emissions are similar to CMIP5 estimates until 1950 when CEDS emissions began to grow at a faster rate than CMIP5 emissions through 1990 when they were about 20–30 % larger. Between 1990 and 2000, CMIP5 estimates show a decrease in emissions while CEDS estimates show flattening emissions, then a steep increase. Differences in NH₃ emissions are largely due to steadily increasing agricultural emissions and a larger estimate from wastewater/human waste, which makes up 14 % of CEDS NH₃ estimates in recent decades but was largely missing in the RCP estimates. CEDS NMVOC emissions are much larger for global waste, while they are much smaller for global transportation.

Global CEDS CH₄ emissions range from 93 % of CMIP5 values in 1970 to 109 % of CMIP5 values in 2000. CEDS estimates change more smoothly over time, without a dip in 2000. CEDS energy estimates are consistently larger than CMIP5 emissions, by 22–58 %, while CEDS agriculture emissions are consistently 10–15 % smaller than CMIP5 estimates, except in 2000 (6 % smaller) when CMIP5 estimates dip and CEDS emissions flatten due to our inclusion of FAO agriculture data.




Figure 5. Recent emission estimates (1990–2014) by region compared to Lamarque et al. (2010) (dots) and CDIAC (line) for CO_2 . This shows the same data as Fig. 3 over a shorter timescale. For like-with-like comparison, these figures do not include aviation or agricultural waste burning on fields. The “International” region shows international shipping emissions.

Figure 7 shows differences between total gridded emissions for CEDS and CMIP5 for BC and SO_2 in 1900 and 2000. In 1900, CEDS BC emissions were lower over the US and Europe (especially cities in the UK), and larger over parts of India and China. Larger differences are concentrated in high-population areas. In 2000, emissions followed a similar pattern. CEDS BC emissions are smaller over Europe and the

eastern US, but larger over populated areas of India, China, and western Africa (particularly Nigeria), reflecting, in part, higher country totals (e.g., Fig. S41).

Additional text and similar difference maps for NO_x , CO, OC, NH_3 , and NMVOCs, as well as high-resolution figures for SO_2 , are included in the supplemental figures and tables (Sect. K). The magnitude of most differences in 1850 is




Figure 6. Total gridded CEDS emissions by emission species for 2010.




Figure 7. Difference between CEDS and CMIP5 total gridded emissions for BC (top) and SO₂ (bottom) in 1900 (left) and 2000 (right) at 10° grid cells. Values shown are CEDS – CMIP5 estimates. For like-with-like comparison, these figures do not include aviation or agricultural waste burning on fields.

small, as total global emissions are small and tend to be more concentrated in populated areas, with larger differences by 1900. Differences in 2000 are a bit larger and tend to be consistent across countries. For example, total CEDS CO emission in India in 2000 are smaller than CMIP5 values, so most grid cells in India have smaller values.

However, differences in gridded SO₂ emissions in 2000 are not as consistent across countries or regions and tend to be highly concentrated into small groups of grid cells. Globally, CEDS SO₂ emissions are very similar to CMIP5 emissions, and emissions are dominated by large point sources, so these differences are likely due to updated proxy data for power plants and metal smelters. The distribution of SO₂ emissions over the US also differs from CMIP5 grids, shown in Fig. 7 and Sect. K of the supplemental figures and tables, and detailed in Section A2.1.

BC, OC, and NH₃ CEDS emissions in 2000 are larger over India, China, and parts of Africa than CMIP5 estimates, similar to BC emissions in Fig. 7. CEDS NO_x emissions in 2000 are also larger over China and India, while they are smaller over the Middle East and eastern Europe. NMVOC estimates are smaller over China and the Middle East.

As discussed further in the Supplement (Sect. K), these differences are due to a combination of differences in aggregate country-level emission estimates, spatial proxy data, and methodologies for mapping aggregate emissions to spatial grids. We note that the spatial proxy that is most important will also depend on emission species: for SO₂, power plants will generally be a key sector, while for NO₂, mobile sources are an important sector over recent decades.

3.4.2 GAINS and EDGAR v4.3

CEDS estimates are compared to GAINS and EDGAR v4.3 emission estimates in Fig. S40, shown in the supplemental figures and tables.

Comparing GAINS with CEDS for BC, OC, NO_x, and SO₂ CEDS estimates is within $\pm 20\%$ of global GAINS values in 2000, 2005, and 2010. OC and SO₂ CEDS emissions are smaller than GAINS values in 2000 but become larger than GAINS global values by 2010. CEDS NO_x, CO₂, and BC emissions are consistently smaller than GAINS estimates and CEDS CO estimates are consistently larger than GAINS but within 6%, while CEDS NMVOCs are 26–43% larger than GAINS estimates from 2000 to 2010.

BC emissions increase by about 10% from 2000 to 2010 in GAINS while the increase is 33% in CEDS. Two particularly large differences are due to coke production in China, which is particularly uncertain, and residential emissions from biofuel use (see Fig. 4), both of which increase significantly over this period in CEDS.

Between 2000 and 2010, global CEDS emissions for all species (except CO₂) increase more than the GAINS estimates, with CEDS estimates higher than GAINS by 2010 for a number of species (Fig. S40). GAINS emissions ex-

hibit slower growth than CEDS emissions in recent years, indicating that GAINS includes more emission controls or other changes over this period than CEDS (and the inventories to which CEDS is calibrated). The divergence in recent years is particularly present in SO₂ and NO_x emissions for power generation in China and India, and SO₂ globally from refineries. This divergence continues to 2015 (IEA, 2016b, based on an updated version of GAINS), in which global SO₂ emissions decline by $\sim 25\,000$ Gg from 2005 to 2015, while CEDS emissions decline by only $\sim 10\,000$ Gg over 2005 to 2014.

CEDS estimates are consistently larger than EDGAR v4.3 global estimates for most emission species. CEDS emissions follow the similar trends as EDGAR from 1970 to 2000 or all species but OC. CEDS emissions for OC grow somewhat linearly over the period, while EDGAR estimates stay relatively flat. Sectors driving the differences between CEDS and EDGAR estimates vary by emission species. However, these differences are largely due to waste burning and aggregate sector 1A4, which is dominated by residential emissions but also includes commercial/institutional emissions and agriculture/forestry/fishing. A key difference is associated with estimates for waste (trash) burning which are much higher in CEDS (based on Wiedinmyer et al., 2014) and have a strong influence on totals, particularly OC, with smaller relative impacts on NMVOCs and BC (see Sect. 3.5).

Global CEDS CH₄ emission estimates are slightly smaller than, but similar to, EDGAR v4.2 estimates, ranging from 94 to 98% of the EDGAR estimates. The similarity is because much of our methane emissions are either from EDGAR or FAO (which uses similar methodologies). The largest differences can be found in 1B2 (fugitive petroleum and gas emissions) in Central and South America, Africa, and the former Soviet Union, as these default emissions also incorporate data from ECLIPSE V5a (Stohl et al., 2015), and rice cultivation in China (FAO, 2016).

3.5 Uncertainty

Emission uncertainty estimates in inventories are a critical need; however, this is difficult to quantify and most inventories do not include uncertainty estimates. All the components and assumptions used in this analysis are uncertain to varying degrees, which means that uncertainty will vary with time, space, and emission species making quantification of uncertainties challenging.

There are some consistent trends in uncertainty estimates by emission species. Uncertainty is generally lowest for CO₂ and SO₂ emissions, which depend primarily on quality of fossil fuel statistical data and fuel properties, e.g., carbon and sulfur content, with straightforward stoichiometric relationships. Global CO₂ and SO₂ uncertainty has been estimated to be on the order of 8% for CO₂ (Andres et al., 2012) and 8–14% for SO₂ (Smith et al., 2011), for a roughly 5–95% confidence interval. Global uncertainties for these species tend to

be relatively low also because fuel properties are not thought to be highly correlated between major emitting regions.

Uncertainty in specific countries can be much higher, however. China is a major emitter of both CO₂ and SO₂, and uncertainties regarding the level of coal consumption (Guan et al., 2012; Liu et al., 2015b) will directly impact emission estimates as well as actual implementation and efficiency of control equipment (Xu et al., 2009; Zhang et al., 2012). Since China energy consumption uncertainties appear to be largest in sectors with limited emission controls, they can have a large impact on SO₂ emissions in particular (Hong et al., 2017). There is also uncertainty regarding the appropriate CO₂ emission factor for coal in China (Liu et al., 2015b; Olivier et al., 2015) as discussed further in Sect. 5.4 of the Supplement.

Emission factors for CO, NO_x, NMVOCs, BC, and OC, tend to be dependent on details of the emitting process and therefore have higher uncertainties (Blanco et al., 2014). This is particularly true for carbonaceous aerosol emissions, where emission factors can range over several orders of magnitude depending on the conditions under which combustion occurs. Uncertainties in global BC emissions have been estimated to be a factor of 2 (Bond et al., 2004). Uncertainty in country-level BC emissions in China was estimated to be –43 to +93 % by Lu et al. (2011), –50 to +164 % by Qin and Xie (2012), ±176 % by Kurokawa et al. (2013a), and –28 to +126 % by Zhao et al. (2013b). Uncertainty in activity levels also contributes. Solid biomass consumption is difficult to track, and both absolute values and trends are generally much more uncertain than fossil fuel consumption data, which will contribute to BC and OC emission uncertainty.

Emission uncertainties for CO, NO_x, and NMVOCs typically lie between those of carbonaceous aerosols and those of CO₂ and SO₂. In part, this is because, particularly in industrialized economies, a number of sectors contribute to emissions, and sectoral uncertainties will largely be independent of each other. Substantial uncertainty can still be present for specific sectors, even in countries with well-developed emission inventory processes (Parrish, 2006). For example, studies combining observations and modeling suggest that the recent US national emission inventory overestimates on road vehicle NO_x emissions by about a factor of 2 (Anderson et al., 2014; Hassler et al., 2016; Travis et al., 2016), while recent updates of Canadian NMVOC emissions (Environment and Climate Change Canada, 2016) are, for some sectors, a factor of 2 larger than previous estimates (Environment Canada, 2013).

There are specific sectors with particularly uncertain emissions. The level of fugitive emissions often depends on procedures and practices, leading to large uncertainty. Emissions that result from biological processes, such as NO_x from fertilized soils or NH₃ from wastewater and agriculture, also generally depend on environmental conditions and would, in principle, require detailed modeling to improve estimates. Our NH₃ emissions from human waste, for example, adapt

the methodologies used in REAS (Kurokawa et al., 2013a) and use a single global default emission factor (modified to account for wastewater treatment as described in the Supplement). Not only is this emission factor uncertain, but there will certainly be regional variations due to differing environmental conditions that we were unable to take into account. For agricultural emissions, the actual practices of managing livestock manures will affect true emissions; such practices vary significantly across the world but are not always well understood or reflected in the emission factors used in global inventories (Paulot et al., 2014). We note that in the CEDS historical extrapolation before either 1960 or 1970, depending on the sector, global trends were used for agricultural emissions, which means that country-specific trends were not taken into account, leading to additional uncertainties at the country level.

Residential waste burning emissions depend on the amount of waste combusted, composition of the waste, and combustion conditions. This sector globally contributes a substantial fraction of OC emissions in particular but substantial amounts of BC and other species. The CEDS estimate for this sector, except where scaled to country emission estimates (available only in a few OECD countries), is based on 2010 estimates from Wiedinmyer et al. (2014). Wiedinmyer et al. followed IPCC guidelines and assumed that 60 % of all waste that is not reported as collected is burnt. This could be an overestimate in countries where there is informal waste collection and recycling. Klimont et al. (2017a) recently estimated BC and OC emissions from this sector, estimating that from 115 to 160 Tg of waste was openly burned, while Wiedinmyer et al. (2014) derived a value of 970 Tg. It is possible that the CEDS values, therefore, are overestimates of emissions from this source. Note, however, that the Wiedinmyer et al. (2014) estimate only includes residential waste burning. In the USA, for example, a large portion of CO₂ from waste burning is from industrial waste, particularly from tires (US EPA, 2015), which implies there may also be additional air pollutant emissions from industrial waste combustion. Outside of the specific OECD countries where country-specific inventories include this sector, industrial waste estimates were not explicitly included in the CEDS estimates. Overall, there is substantial uncertainty for emissions from this sector.

All other factors being equal, uncertainty will tend to increase backwards in time, as driver data become more uncertain and older technologies are used, for which emission factors are not well quantified. We generally expect that uncertainty in this data set will be smaller for those years and countries where robust inventory development mechanisms are in place. However, as noted above for NO_x in the USA, this does not eliminate uncertainty. Official country inventories can sometimes be developed with outdated methodologies or can be incomplete. Many countries have regular evaluation activities, which indicate deficiencies and potential areas for improvement. However, assessments of completeness

and plausibly are always useful, and inventories developed for scientific use, including CEDS, can help contribute in this area.

Our data system also allows us to examine the emission factors implied by scaling to country inventories. This can reveal potential inconsistencies or regional differences. One example is shown in Fig. 8, which shows the implied emission factor for CO emissions from gasoline road vehicles. Even where there is a mix of fuels in the road sector, the much higher CO emission factor for gasoline tends to lead to gasoline dominating emissions, making this comparison a fairly unambiguous reflection of underlying inventory assumptions. There is over a factor of 2 difference in implied emission factors before 1990, with some inventories indicating steadily increasing emission factors going back in time while others flatten out. It is unclear if these differences are due to local variations in vehicle types, operation, or environmental conditions, or if differences reflect inventory assumptions, which implies some inventories might be biased high or low.

There are specific issues with uncertainty over the most recent few years in most emission data sets. We have, in this data set, provided emissions up to 2014. Emissions estimated for the most recent several years are likely to have larger uncertainty due to the use of incomplete or preliminary data. Uncertainty in recent years comes from three main sources: activity data, emission inventories that are used in our estimate, and the treatment of emission factors. Uncertainty from activity data comes from both uncertainty in country totals and their sector split. While activity data are often updated annually, recent estimates sometimes change for a few years after their initial release. For example, the BP estimate of Russian coal use in 2012 may be different in the 2013, 2014, and 2015 data releases. The BP estimates we use to extrapolate fuel use for the most recent 2 years (Sect. 2.2.1) also lack sectoral detail, which adds to uncertainty. Values in the inventory estimates we use in this data set for the most recent year are often preliminary and are later revised, which is an additional source of uncertainty.

Finally, we use emission factor trends from GAINS to project emission factors for combustion sectors for recent years beyond where inventory data are available. The last inventory year varies: 2010 for EDGAR, which is our default inventory for most species, 2008 for REAS, 2012 for China, 2013 for most of Europe, and 2014 for the USA. Using emission factor trends that are not from detailed country-specific inventories is an additional source of uncertainty.

In future versions of CEDS, a quantitative uncertainty analysis will be included for all time periods, as further discussed in Sect. 5.

4 Comparisons with observations

It is challenging to evaluate emissions against observations since, other than facility-specific emission monitors, concentrations of emitted species are observed rather than emission fluxes into the atmosphere. Satellite data (Jacob et al., 2016; Streets et al., 2013), road-side measurements (Pant and Harrison, 2013), and inversion of surface observations (Bruhwiler et al., 2014; Houweling et al., 2017) can all be used to estimate emissions using observational data. These techniques can be used to gain insights into the accuracy of emission inventories, although each has associated uncertainties. Emission ratios are a particularly valuable technique, and we compare in this section CEDS data with observations for two cases.

Hassler et al. (2016) compare observed ambient NO_x/CO enhancement ratios (measurements taken during morning rush hour) with NO_x/CO road emission trends for London, Paris, and several US cities. Hassler et al. compare to the MACCity inventory (Granier et al., 2011), which is based on CMIP5 (Lamarque et al., 2010) inventory estimates and RCP projections. They find that log linear trends in observed ratios in US cities, London, and Paris are steeper than MACCity ratios by a factor of 2.8–5.5. CEDS country-level NO_x/CO emissions ratios match observed trends much closer than MACCity, where observed trends are only 2–18% steeper than CEDS trends, as shown in Table 7. Further, CEDS gridded road emissions match even better with the observed trends for London and Paris.

Kanaya et al. (2016) present observations of BC/CO ratios over 6 years (2009–2015) at Fukue Island, Japan, which, depending on wind conditions, gives region-specific emission ratios under dry conditions for Japan, South Korea, and four regions in China, shown in Table 8 compared to CEDS and REAS BC/CO emissions ratios, both of which do not include open biomass burning. Both CEDS and REAS emissions ratios are similar to observed ratios for Japan, 1.64 and 1.1 times larger than observed ratios, respectively, but near the observational uncertainty. The 2008–2015 average CEDS emission ratio is 2.1–2.7 times larger than observed ratios from China regions.

CEDS emissions ratios are substantially larger than both observed and REAS ratios for South Korea. Kanaya et al. attribute the difference between REAS and observations in South Korea to the overestimation of industry and transportation BC/CO ratios in inventories. CEDS South Korean sector-specific BC/CO emission ratios are high compared to observations: 370 and 41 $\text{ng m}^{-3} \text{ppb}^{-1}$ for industry and transportation sectors, respectively, compared to 42 and 27 $\text{ng m}^{-3} \text{ppb}^{-1}$ in REAS. CEDS CO estimates, which are scaled to the South Korean national inventory from 1999 to 2012, are 5–47% lower than REAS2.1 estimates over 2000–2008. CEDS CO emission estimates are dominated by energy transformation (20%) and transportation (68%). CEDS BC estimates use SPEW assumptions. CEDS BC emission




Figure 8. Implied CO emission factor for gasoline road vehicles obtained by the CEDS system after scaling to match country inventories. Data points are only shown where an inventory value was available in units of g CO g^{-1} fuel.

Table 7. Trends in observed and inventory NO_x/CO emission ratios.

City/country	Years	Observed*	MACCity*	CEDS (gridded – road)	CEDS aggregate (road)	CEDS (total)
USA (various cities)	1989–2013	4.1	1.45		3.86	2.37
UK (London)	1989–2015	7.2	1.88	6.92	6.90	5.90
France (Paris)	1995–2014	8.8	1.59	8.09	7.47	3.39

Values are shown in log linear trends in units of $\% \text{ yr}^{-1}$
* (Hassler et al., 2016)

estimates for South Korea are 5–8 times larger than REAS estimates. While CEDS estimates are larger over all sectors, the other transformation (e.g., coal coke production) and road sectors are the primary sources. Emissions from the CEDS other transformation sector, which are zero in REAS estimates, make up 35 % of CEDS South Korean estimates. CEDS road BC emissions over 2000–2008 are 2–3 times larger than REAS estimates and 34 % of the CEDS total.

These comparisons are approximate, given that the CEDS data represent entire countries, and the air trajectories sampled at Fukue Island will preferentially sample only portions of each country. In future versions of CEDS, we plan to produce emissions for large countries such as China at the province level which will aid in such comparisons. In general, differences in these ratios could be attributed to the overestimation of BC, underestimation of CO emissions, or both. Overall, CEDS emissions appear consistent for Japan but perhaps slightly too high for China. CEDS BC estimates for South Korea are quite high compared to other inventories and the observations, and suggest that the SPEW emission factors for South Korea may not have incorporated the impact of transportation emission controls and new technologies for coal coke production.

These examples illustrate that further comparisons would be of substantial value in better resolving emissions. The use of multiple observations and methodologies would add confidence to conclusions regarding the accuracy of emission inventory data.

5 Limitations and future work

While this data set includes many improvements upon existing comprehensive, long-term inventories, there are some specific limitations of the current methodology, and plans for improvement, that we discuss here.

Disaggregation of key non-combustion sectors, particularly 1A1bc_Other-transformation and 2C_Metal-production, should allow a more accurate estimation of emission trends. This will require collection of additional activity data and default emission factors. At the current level of aggregation, emission trends for these sectors will be less accurate, particularly for years where country-level emission data sets are not available.

Table 8. Observed and inventory BC/CO emission ratios.

Country	Observed* 2009–2015 ($\Delta\text{BC}/\Delta\text{CO}$)	CEDS 2009–2014 (BC/CO)	CEDS 2008 (BC/CO)	REAS2.1 2008 (BC/CO)
Japan	5.9 ± 3.4	9.7	9.5	6.5
South Korea	6.7 ± 3.7	89.8	82.3	23
China (north east)	6.0 ± 2.8	14.3	12.8	
China (north central east)	5.3 ± 2.1			8.3
China (south central east)	6.4 ± 2.2			9.9
China (south)	6.9 ± 1.2			

Values shown in $\text{ngm}^{-3} \text{ppb}^{-1}$
* (Kanaya et al., 2016)

Emission trends could be further improved for the mid-20th century. Emission factors here are often the result of scaling at later inventory years (e.g., Fig. 4), and further work to better constrain emission factors over this period is needed. The sectoral split for fuel use is also approximated over this period; incorporation of regional activity data would improve this as well. Non-combustion emissions are particularly uncertain in the era before modern inventory data sets, which is generally before 1970/1980, since these emissions can depend on process details.

We plan to incorporate more detailed data from the US National Emissions Inventory, although as with the current estimate, discontinuities due to methodological changes will need to be addressed. Use of these data to estimate emissions at the US state level is underway, which will also be used to improve the spatial gridding of emissions over time.

Currently, a number of gridding proxies are static over time. Residential (and related) emissions are distributed using population distribution, which does change over time. Because residential emissions are dominant in earlier years, much of the major shifts in spatial distribution within countries are being captured. Other sectors have a mix of spatial proxies, few of which are newer than 2010, and many were kept static over time. Shipping emission patterns have changed over time; however, we lack consistently constructed spatial proxies over time. The shipping spatial data used here (from ECLIPSE) have a higher fraction of emissions in the North Atlantic than the spatial distribution used in CMIP5. It is not clear if this difference is due to different methodologies or an actual change in spatial distribution over the last decade. Consistent data sets over time for spatial proxy information would be a useful addition.

A major next step in this project will be estimation of uncertainty. Our first step will be quantification of the additional uncertainty that stems from producing estimates out to the most recent full year, followed by comprehensive un-

certainty estimates that will be used to produce ensembles of emissions to more fully reflect the uncertainty in these data.

In addition to updates, refinements, and uncertainty analyses, the CEDS system will be released as open-source software, along with associated input data. Where previous work has only released final emission estimates, this entire data system will be released to facilitate evaluation of trends in and the relationships between emissions, emission factors, and their drivers across time, countries, sectors, and fuels; foster transparency in assumption and methods; and allow community input and participation. While the current data system requires purchase of the IEA energy statistics, we will explore options to facilitate use with publicly available data as well.

6 Summary

This paper described the methodology and results for a new annual data set of historical anthropogenic GHGs, reactive gases, aerosols, and aerosol and ozone precursor compounds from 1750 to 2014 for use in CMIP6. This data set relies heavily on IEA energy statistics, EDGAR, and other inventory data sets to produce consistent trends over time. Key steps in estimating emissions include collecting existing activity, emission factors, and emission data; developing default emission estimates; calibrating default estimates to existing inventories; extending present-day emissions to historical time periods; and gridding emissions.

Emissions before 1850 are dominated by residential biomass burning and agricultural emissions. As the industrial revolution expanded, energy-, industry-, and transportation-related emissions then began to grow and then quickly increased in the mid-20th century. Emissions of some species began to slow or see global reductions in the late 20th century with the introduction of emission control policies, but emissions of many of those species increased again in recent years due to increased economic activity in rapidly

industrializing regions. While comparable to existing data sets, such as CMIP5 (Lamarque et al., 2010), EDGAR (EC-JRC/PBL, 2016) and GAINS (Amann et al., 2011; Klimont et al., 2017a), CEDS estimates are generally slightly higher than those inventories in recent years.

Future work on this data system will involve refining and updating these emission estimates, adding details, and the release of the CEDS as an open-source data system. In order to be able to release the current data set in time for use in CMIP6, the focus was on the development and use of a consistent methodology, relying largely on IEA energy statistics and existing inventory data over recent years. As described above and in the Supplement, a number of additions were made where inconsistencies or incompleteness in these core data sets were known and improved data were readily available. There are many further corrections that would likely be useful to implement. For example, the inventories used here for calibration may already be known to contain deficiencies, for example, through regular validation activities. There are likely also country-level energy and other driver data that can be used to improve the data used here. Finally, further detailed comparisons with observations may help to indicate additional areas where changes to emission factor or other assumptions are warranted.

With the release of this data set, and soon the entire data system, it is our intention that further improvements will be made through feedback from the global emission inventory community. The CEDS data system, including R code and all input data other than the IEA energy statistics, is being prepared for public release in fall 2017 through the GitHub collaboration website. This will facilitate community comments and direct contributions to improving these emission data. The next data release is planned for fall/winter 2017, which will extend the time series to 2016 and correct, to the extent possible, any known issues with the data set. We aim to continue annual updates in subsequent years. We welcome comments, including notes on any potential inconsistencies or relevant new data sources, so that these data can be improved in future releases.

Data availability. Gridded versions of these data are available through the Earth System Grid Federation (ESGF) (<https://esgf-node.llnl.gov/search/input4mips/>) under the activity_id = “input4MIPs” and institution = “PNNL-JGCRI”. More information on the CEDS project, system release, and updates can be found at <http://www.globalchange.umd.edu/ceds/>. Note that known issues with the data are listed at <https://github.com/JGCRI/CEDS>, and users can also submit issues via the GitHub site.

Appendix A

Table A1. Sector definitions of combustion emissions (IEA and NFR14 codes).

IEA energy statistics	IEA name	NFR14 code	CEDS working sector name	Aggregate sector (gridding)
MAINELEC	Main-Activity-Producer-Electricity-Plants	1A1a	1A1a_Electricity-public	Power_and_Heat
AUTOELEC	Autoproducer-Electricity-Plants	1A1a	1A1a_Electricity-autoproducer	Industrial_Combustion
MAINCHP	Main-Activity-Producer-CHP-Plants	1A1a	1A1a_Electricity-public	Power_and_Heat
AUTOCHP	Autoproducer-CHP-Plants	1A1a	1A1a_Electricity-autoproducer	Industrial_Combustion
MAINHEAT	Main-Activity-Producer-Heat-Plants	1A1a	1A1a_Heat-production	Power_and_Heat
AUTOHEAT	Autoproducer-Heat-Plants	1A1a	1A1a_Heat-production	Power_and_Heat
IRONSTL	Iron-and-Steel	1A2a	1A2a_Ind-Comb-Iron-steel	Industrial_Combustion
NONFERR	Non-Ferrous-Metals	1A2b	1A2b_Ind-Comb-Non-ferrous-metals	Industrial_Combustion
CHEMICAL	Chemical-and-Petrochemical	1A2c	1A2c_Ind-Comb-Chemicals	Industrial_Combustion
PAPERPRO	Paper,-Pulp-and-Print	1A2d	1A2d_Ind-Comb-Pulp-paper	Industrial_Combustion
FOODPRO	Food-and-Tobacco	1A2e	1A2e_Ind-Comb-Food-tobacco	Industrial_Combustion
NONMET	Non-Metallic-Minerals	1A2f	1A2f_Ind-Comb-Non-metallic-minerals	Industrial_Combustion
CONSTRUC	Construction	1A2g	1A2g_Ind-Comb-Construction	Industrial_Combustion
TRANSEQ	Transport-Equipment	1A2g	1A2g_Ind-Comb-transpequip	Industrial_Combustion
MACHINE	Machinery	1A2g	1A2g_Ind-Comb-machinery	Industrial_Combustion
MINING	Mining-and-Quarrying	1A2g	1A2g_Ind-Comb-mining-quarrying	Industrial_Combustion
WOODPRO	Wood-and-Wood-Products	1A2g	1A2g_Ind-Comb-wood-products	Industrial_Combustion
TEXTILES	Textile-and-Leather	1A2g	1A2g_Ind-Comb-textile-leather	Industrial_Combustion
INONSPEC	Non-specified-(Industry)	1A2g	1A2g_Ind-Comb-other	Industrial_Combustion
WORLDNAV	World-Aviation-Bunkers	1A3ai	1A3ai_International-aviation	Aviation
DOMESAIR	Domestic-Aviation	1A3aii	1A3aii_Domestic-aviation	Aviation
ROAD	Road	1A3b	1A3b_Road	Road
NA*	Evaporative emissions from road transport and tanker loading do not have corresponding IEA sectors, denoted with "NA"	1A3b	1A3b_Road	Road
RAIL	Rail	1A3c	1A3c_Rail	Other_Surface_Transport
WORLDMAR	World-Marine-Bunkers	1A3di	1A3di_International-shipping	International-Shipping
NA*	Evaporative emissions from tanker loading	1A3di	1A3di_Oil_tanker_loading	International-Shipping
DOMESNAV	Domestic-Navigation	1A3dii	1A3dii_Domestic-navigation (shipping)	Other_Surface_Transport
PIPELINE	Pipeline-Transport	1A3ei	1A3eii_Other-transp	Other_Surface_Transport
TRNONSPE	Non-specified-(Transport)	1A3eii	1A3eii_Other-transp	Other_Surface_Transport
COMMPUB	Commercial-and-Public-Services	1A4a	1A4a_Commercial-institutional	Residential_Commercial_Other
RESIDENT	Residential	1A4b	1A4b_Residential	Residential_Commercial_Other
AGRICULT	Agriculture/Forestry	1A4c	1A4c_Agriculture-forestry-fishing	Residential_Commercial_Other
FISHING	Fishing	1A4c	1A4c_Agriculture-forestry-fishing	Residential_Commercial_Other
ONONSPEC	Non-specified-(Other)	1A5	1A5_Other-unspecified	Residential_Commercial_Other

A1 Data release and known issues

A1.1 Known issues

This section lists known issues with the data released as of this writing (August 2017). Readers should refer to the project website for general updates (globalchange.umd.edu/CEDS) and the project's GitHub site for an updated list of issues (<https://github.com/JGCRI/CEDS>).

- Combustion emissions become zero in earlier years for several countries that have inconsistent temporal coverage in the IEA energy data. These include Sint Maarten, Suriname, Cambodia, Mongolia, Palau, Botswana, Namibia, and Niger. Some of these instances, where alternative data sources are available, will be corrected in the next release.
- Some of the countries in the IEA “other” aggregations (e.g., “Other Asia”, “Other Africa”, and “Other Non-OECD Americas”) have spurious sector splits due to the simple methods used to assign fuel use to these countries (e.g., there is fuel use in the Afghanistan international shipping sector).
- There are a few spurious small-magnitude process emissions (particularly in 2C_Metal-production) for smaller countries before 1900 that are artifacts of the extension process. These have negligible impacts on emission totals.
- There are some spurious emission results for early years at the sectoral level in the current database due to the sectoral resolution of the data used to extend emissions back in time. For example, aircraft emissions are present back to 1851, even though actual aircraft emissions did not begin until the early 20th century. The magnitudes of these emissions are small and, while these emissions should be zero in early years, these small magnitudes will not materially impact climate model results. (NO_x in the CEDS aviation sector in 1920 is 0.2 % of estimated NO_x from lighting (Schumann and Huntrieser, 2007), for example, and very much smaller in earlier years.) The historical energy code is being revised to be more flexible to improve our ability to incorporate additional energy data sets including, for example, historical estimates of aircraft fuel consumption.
- Due to an error, SO₂ emissions in the US are overestimated from about 1961 to 1969. The overestimate averages 22 % over this period. This has been corrected for inclusion in the next data release. The previous and corrected time series is shown in the Supplement.
- SO₂ emissions in the gridded data are overestimated in the western United States relative to the eastern United States. This spatial allocation is present in the EDGAR

emission grids used for spatial mapping within each country.

A1.2 Gridded data release history

There have been several releases of the CEDS gridded data. The underlying emissions by country, sector and fuel have been identical in all of these releases, as have total emissions by country and gridding sector (with the exception of small changes in 1850 emissions noted below).

v2016-05-20: Pre-industrial 1750–1850 data release

v2016-06-18: 1851–2014 data

v2016-06-18-sectorDim: Re-release of both pre-industrial and 1851–2014 in a new NetCDF format with sectors as an additional dimension in the data variable. This reformatting was necessary due to a limitation that was discovered within the ESGF system in summer 2016. The reformatted data were released in early fall 2016.

v2017-05-18: Re-release of entire data set in order to correct two gridding errors discovered by users. The first error was inconsistent emission allocation to spatial grids within countries that resulted in incorrect spatial allocations and some large discontinuities in the gridded data. These issues were particularly apparent in spatially large countries such as the USA and China. The second error was minor inconsistencies in seasonal allocation, resulting largely in emissions that were too high in February. Total annual emissions within each country were not impacted by either of these issues.

Emissions are also fully consistent across 1850 in this release. There were small discontinuities in 1850 between the CEDS CMIP6 pre-industrial release (v2016-06-18) and later the full CEDS release (v2016-07-26) due to updates in the data system. These differences are 0.5 % for all species (except NMVOCs which reach 1.5 %). In absolute terms, these differences are very small (relative to, for example, open biomass burning emissions) and will not have a significant impact on simulation results.

A link to further examination of these issues, including comparison maps and time series comparisons, can be found at the project website (globalchange.umd.edu/CEDS).

A1.3 Methane historical extension

As several modeling groups participating in CMIP6 requested CH₄ emissions from 1850, we were not able to extend the consistent CH₄ time series before 1970 due to the additional data that would need to be collected and processed. We have, however, produced a “rough cut” supplementary extension of CH₄ emissions from 1850–1970 by scaling with CMIP5 historical CH₄ estimates (Lamarque et al., 2010). These estimates were generated by scaling the CEDS 1970 estimates with the CMIP5 trends (i.e., shifting CMIP5 trends to match CEDS values in 1970) by aggregate sector and the 26 subregion levels of the CMIP5 data. While these emission estimates are not fully consistent with the other CEDS emis-

sions, they provide a longer time series, albeit with some additional uncertainty, for groups that would like to have these trends. These data are available as supplementary gridded information for CEDS version 2017-05-18 data through ESGF (see data availability section).

Biases in this extended data set have already been identified. The waste sector is 30 % of total anthropogenic CH₄ emissions by 1850. This is likely because earlier CMIP5 data are scaled back in time with population data. This is an overestimate of anthropogenic CH₄ emissions from this source at that time since landfills and wastewater treatment plants, which create the anaerobic conditions conducive to CH₄ emissions, did not start to come into widespread use until around 1930. However, as noted in the main paper, earlier CMIP5 emission estimates did not distinguish between biomass and coal combustion. CH₄ emissions from biomass combustion are much larger than those from coal combustion, which means methane emissions from the residential sector are underestimated in this extrapolation. A rough estimate indicates that these two effects are of similar (and offsetting) magnitude. Further work is necessary to better refine historical CH₄ emissions.

Table A2. Sector definitions of non-combustion emissions (drawn from EDGAR processes).

EDGAR process description	CEDS working sector name	Aggregate sector (gridding)
Fuel combustion petroleum refineries	1A1bc_Other-transformation	Industrial_Combustion
Coal mines	1A1bc_Other-transformation	Industrial_Combustion
Fuel combustion BKB (brown coal briquette) plants	1A1bc_Other-transformation	Industrial_Combustion
Fuel combustion blast furnaces	1A1bc_Other-transformation	Industrial_Combustion
Fuel combustion charcoal production plants	1A1bc_Other-transformation	Industrial_Combustion
Fuel combustion coal liquefaction plants	1A1bc_Other-transformation	Industrial_Combustion
Fuel combustion coke ovens	1A1bc_Other-transformation	Industrial_Combustion
Fuel combustion gasification plants for biogas	1A1bc_Other-transformation	Industrial_Combustion
Fuel combustion liquefaction/regasification plants	1A1bc_Other-transformation	Industrial_Combustion
Fuel combustion non-specified transformation active	1A1bc_Other-transformation	Industrial_Combustion
Fuel combustion oil and gas extraction	1A1bc_Other-transformation	Industrial_Combustion
Fuel combustion patent fuel plants	1A1bc_Other-transformation	Industrial_Combustion
Gas works	1A1bc_Other-transformation	Industrial_Combustion
BKB plants	1B1_Fugitive-solid-fuels	Fugitive_Energy_Emissions
Fuel transformation coal liquefaction plants	1B1_Fugitive-solid-fuels	Fugitive_Energy_Emissions
Fuel transformation patent fuel plants	1B1_Fugitive-solid-fuels	Fugitive_Energy_Emissions
Production of brown coal	1B1_Fugitive-solid-fuels	Fugitive_Energy_Emissions
Production of hard coal	1B1_Fugitive-solid-fuels	Fugitive_Energy_Emissions
Production of peat	1B1_Fugitive-solid-fuels	Fugitive_Energy_Emissions
Fuel transformation charcoal production plants	1B1_Fugitive-solid-fuels	Fugitive_Energy_Emissions
Fuel transformation coke ovens	1B1_Fugitive-solid-fuels	Fugitive_Energy_Emissions
Fuel transformation in gas works	1B1_Fugitive-solid-fuels	Fugitive_Energy_Emissions
Chemical heat for electricity production	1B2_Fugitive-petr-and-gas	Fugitive_Energy_Emissions
For blended natural gas	1B2_Fugitive-petr-and-gas	Fugitive_Energy_Emissions
Fuel transformation gasification plants for biogas	1B2_Fugitive-petr-and-gas	Fugitive_Energy_Emissions
Fuel transformation liquefaction/regasification plants	1B2_Fugitive-petr-and-gas	Fugitive_Energy_Emissions
Gas-to-liquid (GTL) plants	1B2_Fugitive-petr-and-gas	Fugitive_Energy_Emissions
Non-specified transformation activity	1B2_Fugitive-petr-and-gas	Fugitive_Energy_Emissions
Petrochemical industry	1B2_Fugitive-petr-and-gas	Fugitive_Energy_Emissions
Transformation in GTL plants	1B2_Fugitive-petr-and-gas	Fugitive_Energy_Emissions
Fuel transformation petroleum refineries	1B2_Fugitive-petr-and-gas	Fugitive_Energy_Emissions
Production of oil	1B2_Fugitive-petr-and-gas	Fugitive_Energy_Emissions
Production of gas	1B2_Fugitive-petr-and-gas	Fugitive_Energy_Emissions
Production of oil	1B2_Fugitive-petr-and-gas	Fugitive_Energy_Emissions
(None)	1B2d_Fugitive-other-energy	Fugitive_Energy_Emissions
Cement production	2A1_Cement-production	Minerals
Lime production	2A2_Lime-production	Minerals
Lime production	2A2_Lime-production	Minerals
Soda ash production and use	2Ax_Other-minerals	Minerals
Brick production	2Ax_Other-minerals	Minerals
Glass bottles	2Ax_Other-minerals	Minerals
Glass production	2Ax_Other-minerals	Minerals
Other non-metallic minerals	2Ax_Other-minerals	Minerals
Other uses of carbonate	2Ax_Other-minerals	Minerals
Ammonia production	2B_Chemical-industry	Chemical-industry
Bulk chemicals production	2B_Chemical-industry	Chemical-industry
Nitric acid production	2B_Chemical-industry	Chemical-industry
Adipic acid production	2B_Chemical-industry	Chemical-industry
Silicon carbide production	2B_Chemical-industry	Chemical-industry
Calcium carbide production	2B_Chemical-industry	Chemical-industry
Bulk chemicals production	2B_Chemical-industry	Chemical-industry
Caprolactam production	2B_Chemical-industry	Chemical-industry
Bulk chemicals production	2B_Chemical-industry	Chemical-industry
N-fertilizer production	2B_Chemical-industry	Chemical-industry
Specialities production	2B_Chemical-industry	Chemical-industry
Sulfuric acid production	2B_Chemical-industry	Chemical-industry
Titanium oxide production	2B_Chemical-industry	Chemical-industry
Bulk chemicals production	2B_Chemical-industry	Chemical-industry
Glyoxal production	2B_Chemical-industry	Chemical-industry
Glyoxylic acid production	2B_Chemical-industry	Chemical-industry
Crude steel production	2C_Metal-production	Metals-industry
Blast furnaces	2C_Metal-production	Metals-industry
Pig iron production	2C_Metal-production	Metals-industry
Sinter production	2C_Metal-production	Metals-industry
Pellet production	2C_Metal-production	Metals-industry

* This sector is currently equal to zero in all years and countries, and not included in data files.

Table A2. Continued.

EDGAR process description	CEDS working sector name	Aggregate sector (gridding)
Steel casting	2C_Metal-production	Metals-industry
Ferrous alloy production	2C_Metal-production	Metals-industry
Aluminium production	2C_Metal-production	Metals-industry
Magnesium production	2C_Metal-production	Metals-industry
Aluminium production	2C_Metal-production	Metals-industry
Other non-ferrous production	2C_Metal-production	Metals-industry
Gold production	2C_Metal-production	Metals-industry
Copper production	2C_Metal-production	Metals-industry
Mercury production	2C_Metal-production	Metals-industry
Other non-ferrous production	2C_Metal-production	Metals-industry
Lead production	2C_Metal-production	Metals-industry
Other non-ferrous production	2C_Metal-production	Metals-industry
Magnesium production	2C_Metal-production	Metals-industry
Zinc production	2C_Metal-production	Metals-industry
Paper production	2H_Pulp-and-paper-food-beverage-wood	Pulp-and-paper-food-beverage-wood
Wood pulp production	2H_Pulp-and-paper-food-beverage-wood	Pulp-and-paper-food-beverage-wood
Beer production	2H_Pulp-and-paper-food-beverage-wood	Pulp-and-paper-food-beverage-wood
Bread production	2H_Pulp-and-paper-food-beverage-wood	Pulp-and-paper-food-beverage-wood
Other food production	2H_Pulp-and-paper-food-beverage-wood	Pulp-and-paper-food-beverage-wood
Wine production	2H_Pulp-and-paper-food-beverage-wood	Pulp-and-paper-food-beverage-wood
Non-energy use in petrochemical industry	2L_Other-process-emissions*	Other_Non-Combustion
Non-energy use in industry, transformation industry	2L_Other-process-emissions*	Other_Non-Combustion
Non-energy use in transport sector	2L_Other-process-emissions*	Other_Non-Combustion
Other non-energy use	2L_Other-process-emissions*	Other_Non-Combustion
Non-energy use in petrochemical industry	2L_Other-process-emissions*	Other_Non-Combustion
Non-energy use in industry, transformation industry	2L_Other-process-emissions*	Other_Non-Combustion
Non-energy use in transport sector	2L_Other-process-emissions*	Other_Non-Combustion
Other non-energy use	2L_Other-process-emissions*	Other_Non-Combustion
Other non-combustion not elsewhere (not in EDGAR)	2L_Other-process-emissions*	Other_Non-Combustion
Solvents in glues and adhesives	2-D_Paint-application	Solvents
Solvents in graphic arts	2-D_Paint-application	Solvents
Solvents in paint	2-D_Paint-application	Solvents
Solvents in dry cleaning	2-D_Degreasing-Cleaning	Solvents
Solvents in household products	2-D_Degreasing-Cleaning	Solvents
Solvents in industrial degreasing	2-D_Degreasing-Cleaning	Solvents
Solvents in chemical industry	2-D_Chemical-products-manufacture-processing	Solvents
Other solvent use	2-D_Other-product-use	Solvents
Production and use of other products	2-D_Other-product-use	Solvents
Use of N ₂ O as anesthesia	2-D_Other-product-use	Solvents
Solvents in leather production	2-D_Other-product-use	Solvents
Solvents in pesticides	2-D_Other-product-use	Solvents
Solvents in rubber and plastic industry	2-D_Other-product-use	Solvents
Solvents in vegetative oil extraction	2-D_Other-product-use	Solvents
Enteric fermentation by cattle	3E_Enteric-fermentation	Agriculture_non-combustion
Enteric fermentation by buffalo	3E_Enteric-fermentation	Agriculture_non-combustion
Enteric fermentation by sheep	3E_Enteric-fermentation	Agriculture_non-combustion
Enteric fermentation by goats	3E_Enteric-fermentation	Agriculture_non-combustion
Enteric fermentation by camels	3E_Enteric-fermentation	Agriculture_non-combustion
Enteric fermentation by horses	3E_Enteric-fermentation	Agriculture_non-combustion
Enteric fermentation by asses	3E_Enteric-fermentation	Agriculture_non-combustion
Enteric fermentation by swine	3E_Enteric-fermentation	Agriculture_non-combustion
Manure management of cattle	3B_Manure-management	Agriculture_non-combustion
Manure management of buffalo	3B_Manure-management	Agriculture_non-combustion
Manure management of sheep	3B_Manure-management	Agriculture_non-combustion
Manure management of geese	3B_Manure-management	Agriculture_non-combustion
Manure management of goats	3B_Manure-management	Agriculture_non-combustion
Manure management of camels	3B_Manure-management	Agriculture_non-combustion
Manure management of horses	3B_Manure-management	Agriculture_non-combustion
Manure management of asses	3B_Manure-management	Agriculture_non-combustion
Manure management of swine	3B_Manure-management	Agriculture_non-combustion
Manure management of chickens	3B_Manure-management	Agriculture_non-combustion
Manure management of ducks	3B_Manure-management	Agriculture_non-combustion
Manure management of turkeys	3B_Manure-management	Agriculture_non-combustion

* This sector is currently equal to zero in all years and countries, and not included in data files.

Table A2. Continued.

EDGAR process description	CEDS working sector name	Aggregate sector (gridding)
Separate category for rice CH ₄ emissions (not in EDGAR)	3-D_Rice-Cultivation	Agriculture_non-combustion
Agricultural soils, rice cultivation	3-D_Soil-emissions	Agriculture_non-combustion
Agricultural soils, nitrogen fertilizers	3-D_Soil-emissions	Agriculture_non-combustion
Agricultural soils, animal waste as fertilizer	3-D_Soil-emissions	Agriculture_non-combustion
Agricultural soils, N-fixing crops	3-D_Soil-emissions	Agriculture_non-combustion
Agricultural soils, crop residues	3-D_Soil-emissions	Agriculture_non-combustion
Agricultural soils, histosols	3-D_Soil-emissions	Agriculture_non-combustion
Agricultural soils, buffalo in pasture	3-D_Soil-emissions	Agriculture_non-combustion
Agricultural soils, camels in pasture	3-D_Soil-emissions	Agriculture_non-combustion
Agricultural soils, cattle in pasture	3-D_Soil-emissions	Agriculture_non-combustion
Agricultural soils, chickens in pasture	3-D_Soil-emissions	Agriculture_non-combustion
Agricultural soils, ducks in pasture	3-D_Soil-emissions	Agriculture_non-combustion
Agricultural soils, goats in pasture	3-D_Soil-emissions	Agriculture_non-combustion
Agricultural soils, horses in pasture	3-D_Soil-emissions	Agriculture_non-combustion
Agricultural soils, mules and asses in pasture	3-D_Soil-emissions	Agriculture_non-combustion
Agricultural soils, pigs in pasture	3-D_Soil-emissions	Agriculture_non-combustion
Agricultural soils, sheep in pasture	3-D_Soil-emissions	Agriculture_non-combustion
Agricultural soils, turkeys in pasture	3-D_Soil-emissions	Agriculture_non-combustion
Indirect N ₂ O emissions	3-D_Soil-emissions	Agriculture_non-combustion
Indirect N ₂ O emissions – deposition, other	3-D_Soil-emissions	Agriculture_non-combustion
Indirect N ₂ O emissions – deposition, agriculture	3-D_Soil-emissions	Agriculture_non-combustion
Indirect N ₂ O emissions – leaching and runoff	3-D_Soil-emissions	Agriculture_non-combustion
Agricultural soils, CO ₂ from urea fertilization	3-D_Soil-emissions	Agriculture_non-combustion
Agricultural soils, liming	3-D_Soil-emissions	Agriculture_non-combustion
Solid waste disposal (landfills)	5A_Solid-waste-disposal	Waste
Industrial waste water	5-D_Wastewater-handling	Waste
Domestic waste water	5-D_Wastewater-handling	Waste
Human waste (not in EDGAR)	5-D_Wastewater-handling	Waste
Solid waste disposal (incineration)	5C_Waste-combustion	Waste
Residential waste combustion (not in EDGAR)	5C_Waste-combustion	Waste
Other waste handling	5E_Other-waste-handling	Waste
Coal fires underground	7A_Fossil-fuel-fires	Fossil_Fuel_Files
Oil fires	7A_Fossil-fuel-fires	Fossil_Fuel_Files
Gas fires	7A_Fossil-fuel-fires	Fossil_Fuel_Files

* This sector is currently equal to zero in all years and countries, and not included in data files.

A2 Sector definitions

A2.1 Combustion emissions

Fuel combustion emission sectors in CEDS are defined in reference to corresponding IEA energy statistics energy flows as given in this table. One exception is evaporative emissions from road transport, which are mapped to the 1A3b road transport sector, following general air pollutant inventory practice, even though this is a non-combustion emissions source. Also, NMVOC evaporative emissions from oil tanker loading are not combustion emissions but are categorized together with international shipping emissions.

Note that the current calibration (e.g., scaling) to country emission inventories is generally not performed at this level of detail, which means that sectoral emission values are more reliable at the aggregate sector level.

A2.2 Non-combustion emissions

Non-combustion emission sectors (also generally referred to as process emissions in CEDS documentation) are defined in reference to corresponding EDGAR categories as given in this table. Note that the 1A1bc sector is actually combustion-related emissions; however, this sector is processed the same as non-combustion emissions in CEDS (see Sect. 2.3.2).

Table A3. Fuel mapping to IEA products.

CEDS fuel	IEA product	
biomass	Industrial waste (terajoule (TJ) net)	Primary solid biofuels (TJ net)
	Municipal waste (renewable) (TJ net)	Non-specified primary biofuels/waste (TJ net)
	Municipal waste (non-renewable) (TJ net)	Charcoal (kt)
brown_coal	Brown coal (if no details) (kt)	Peat (kt)
	Lignite (kt)	Peat products (kt)
coal_coke	Coke oven coke (kt)	
hard_coal	Hard coal (if no details) (kt)	Patent fuel (kt)
	Anthracite (kt)	Gas coke (kt)
	Coking coal (kt)	Coal tar (kt)
	Other bituminous coal (kt)	BKB (kt)
	Sub-bituminous coal (kt)	
light_oil	Refinery feedstocks (kt)	Kerosene-type jet fuel excl. biofuels (kt)
	Additives/blending components (kt)	Other kerosene (kt)
	Other hydrocarbons (kt)	Other Kerosene (kt)
	Ethane (kt)	Naphtha (kt)
	Liquefied petroleum gases (LPGs) (kt)	White spirit & SBP (kt)
	Motor gasoline excl. biofuels (kt)	Biogasoline (kt)
	Aviation gasoline (kt)	Other liquid biofuels (kt)
	Gasoline type jet fuel (kt)	
diesel_oil	Natural gas liquids (kt)	Lubricants (kt)
	Gas/diesel oil excl. biofuels (kt)	Biodiesels (kt)
heavy_oil	Oil shale and oil sands (kt)	Bitumen (kt)
	Crude/NGL/feedstocks (if no details) (kt)	Paraffin waxes (kt)
	Crude oil (kt)	Petroleum coke (kt)
	Fuel oil (kt)	Other oil products (kt)
natural_gas	Gas works gas (TJ gross)	Natural gas (TJ gross)
	Coke oven gas (TJ gross)	Natural gas (TJ gross)
	Blast furnace gas (TJ gross)	Refinery gas (kt)
	Other recovered gases (TJ gross)	Biogases (TJ net)
NOT MAPPED	Electric/heat output from non-specified manufactured gases	Wind
	Heat output from non-specified combustible fuels	Other sources
	Nuclear	Electricity (GWh)
	Hydro	Heat (TJ)
	Geothermal (direct use in TJ net)	Total
	Solar photovoltaics	Total of all energy sources
	Solar thermal (direct use in TJ net)	Memo: renewables
	Tide, wave, and ocean	Heat from chemical sources
	Electric boilers	
	Heat pumps	

Information about the Supplement

Supplement files related to this article include:

- Supplemental data and assumptions text (pdf)
- Supplemental figures and tables (pdf)
- Data files (zipped set of csv files)
- Emissions by country and sector (all species)
- Global emissions by sector (all species)
- Total emissions by country (all species)
- Country mapping and ISO codes

The Supplement for this article describes a number of additional data sources used in this work, including the following:

Bartoňová, 2015; Blumberg et al., 2003; Denier van der Gon et al., 2015; EIA, 2013; Endresen et al., 2007; Environment Canada, 2016; Eyring et al., 2005; Fletcher, 1997; Foell et al., 1995; Fouquet and Pearson, 1998; Gschwandtner et al., 1986; Huo et al., 2012; IEA, 2016a; IMO, 2014; Kaur et al., 2012; Kholod and Evans, 2015; Liu et al., 2015a; Ludek and Holub, 2009; McLinden et al., 2016; Mester, 2000; Mitchell, 2003, 2007, 1983; Mylona, 1996; OECD, 2016; Pretorius et al., 2015; Rowe and Morrison, 1999; Ryaboshapko et al., 1996; Sanger, 1997; Simachaya, 2015; Smith et al., 2014; Tushingham, 1996; UK DEFRA, 2015; US EPA, 2012a; Wu et al., 2012; Zhou et al., 2011.

The Supplement related to this article is available online at <https://doi.org/10.5194/gmd-11-369-2018-supplement>.

Author contributions. RMH and SJS prepared the paper with contributions from LF, ZK, GJM, LV, RA, MCPM, LL, ZL, and QZ. The CEDS system was developed by RMH, SJS, LF, TP, JJS, LV, and RB. Analysis was performed by RMH, SJS, LF, LV, MCPM, NK, and PO. Data were contributed by ZK, GJM, RJA, TCB, LD, JK, ML, LL, ZL, and QZ.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. This research was based on work supported by the US Department of Energy (DOE), Office of Science, Biological and Environmental Research as part of the Earth System Modeling program. Additional support for the development of the gridded data algorithm was from the National Atmospheric and Space Administration's Atmospheric Composition: Modeling and Analysis Program (ACMAP), award NNH15AZ64I. The Pacific Northwest National Laboratory is operated for DOE by the Battelle Memorial Institute under contract DE-AC05-76RLO1830. RJA was sponsored

by US Department of Energy, Office of Science, Biological and Environmental Research (BER) programs and performed at Oak Ridge National Laboratory (ORNL) under US Department of Energy contract DE-AC05-00OR22725.

The authors would like to acknowledge Grace Duke and Han Chen for data collection and processing; Alison Delgado, Minji Joeng, and Bo Liu for data collection and translation; and Benjamin Bond-Lamberty and Robert Link for code review. We thank Kostas Tsigaridis for pointing out a discontinuity due to a data anomaly (spurious brown coal consumption over 1971–1984 in the IEA Other Asia region) in a review version of the inventory data.

Edited by: Fiona O'Connor

Reviewed by: Alistair Sellar and one anonymous referee

References

- Akagi, S. K., Yokelson, R. J., Wiedinmyer, C., Alvarado, M. J., Reid, J. S., Karl, T., Crounse, J. D., and Wennberg, P. O.: Emission factors for open and domestic biomass burning for use in atmospheric models, *Atmos. Chem. Phys.*, 11, 4039–4072, <https://doi.org/10.5194/acp-11-4039-2011>, 2011.
- Amann, M., Bertok, I., Borcen, J., Chambers, A., Cofala, J., Dentener, F., Heyes, C., Hoglund, L., Klimont, Z., Purohit, P., Rafaj, P., Schöpp, W., Teixeira, E., Toth, G., Wagner, F., and Winiwarter, W.: GAINS-Asia: a tool to combat air pollution and climate change simultaneously, *Methodology*, available at: <http://pure.iiasa.ac.at/8669/1/XO-08-044.pdf> (last access: 15 January 2018), 2008.
- Amann, M., Bertok, I., Borcen-Kleefeld, J., Cofala, J., Heyes, C., Höglund-Isaksson, L., Klimont, Z., Nguyen, B., Posch, M., Rafaj, P., Sandler, R., Schöpp, W., Wagner, F., and Winiwarter, W.: Cost-effective control of air quality and greenhouse gases in Europe: modeling and policy applications, *Environ. Modell. Softw.*, 26, 1489–1501, <https://doi.org/10.1016/j.envsoft.2011.07.012>, 2011.
- Amann, M., Klimont, Z., and Wagner, F.: Regional and global emissions of air pollutants: recent trends and future scenarios, *Annu. Rev. Env. Resour.*, 38, 31–55, <https://doi.org/10.1146/annurev-environ-052912-173303>, 2013.
- Amann, M., Bertok, I., Borcen-Kleefeld, J., Cofala, J., Heyes, C., Hoglund-Isaksson, L., Kiesewetter, G., Klimont, Z., Schöpp, W., Vellinga, N., and Winiwarter, W.: Adjusted historic emission data, projections, and optimized emission reduction targets for 2030 – a comparison with COM data 2013, IASIA, Laxenburg, Austria, available at: http://ec.europa.eu/environment/air/pdf/review/TSAP_16a.pdf (last access: 15 January 2018), 2015.
- Anderson, D. C., Loughner, C. P., Diskin, G., Weinheimer, A., Canty, T. P., Salawitch, R. J., Worden, H. M., Fried, A., Mikoviny, T., Wisthaler, A., and Dickerson, R. R.: Measured and modeled CO and NO_y in DISCOVER-AQ: an evaluation of emissions and chemistry over the eastern US, *Atmos. Environ.*, 96, 78–87, <https://doi.org/10.1016/j.atmosenv.2014.07.004>, 2014.
- Andreae, M. O. and Merlet, P.: Emission of trace gases and aerosols from biomass burning, *Global Biogeochem. Cy.*, 15, 955–966, <https://doi.org/10.1029/2000GB001382>, 2001.

- Andres, R. J., Marland, G., Boden, T., and Bischof, S.: Carbon dioxide emissions from fossil fuel consumption and cement manufacture, 1751–1991; and an estimate of their isotopic composition and latitudinal distribution, Oak Ridge National Lab., TN (United States); Oak Ridge Inst. for Science and Education, TN (United States), available at: <http://www.osti.gov/scitech/biblio/10185357>, last access: 17 October 2016, 1994.
- Andres, R. J., Fielding, D. J., Marland, G., Boden, T. A., Kumar, N., and Kearney, A. T.: Carbon dioxide emissions from fossil-fuel use, 1751–1950, *Tellus B*, 51, 759–765, 1999.
- Andres, R. J., Boden, T. A., Bréon, F.-M., Ciais, P., Davis, S., Erickson, D., Gregg, J. S., Jacobson, A., Marland, G., Miller, J., Oda, T., Olivier, J. G. J., Raupach, M. R., Rayner, P., and Treanton, K.: A synthesis of carbon dioxide emissions from fossil-fuel combustion, *Biogeosciences*, 9, 1845–1871, <https://doi.org/10.5194/bg-9-1845-2012>, 2012.
- Argentina UNFCCC Submission: Argentinian Inventory 1990–2012, Submitted to UNFCCC, 2016.
- Australian Department of the Environment: National Pollutant Inventory (NPI), Canberra, Australia, <http://www.npi.gov.au/substances> (last access: 15 January 2018), 2016.
- Bartoňová, L.: Unburned carbon from coal combustion ash: an overview, *Fuel Process. Technol.*, 134, 136–158, <https://doi.org/10.1016/j.fuproc.2015.01.028>, 2015.
- Blanco, G., Gerlagh, R., Suh, S., Barrett, J., de Coninck, H. C., Diaz Morejon, C. F., Mathur, R., Nakicenovic, N., Ofori Ahenkora, A., Pan, J., Pathak, H., J. Rice, Richels, R., Smith, S. J., Stern, D. I., Toth, F. L., and Zhou, P.: Drivers, trends and mitigation, in: *Climate Change 2014: Mitigation of Climate Change. Contribution of Working Group III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Edenhofer, O., Pichs-Madruga, R., Sokona, Y., Farahani, E., Kadner, S., Seyboth, K., Adler, A., Baum, I., Brunner, S., Eickemeier, P., Kriemann, B., Savolainen, J., Schlömer, S., von Stechow, C., Zwickel, T., and Minx, J. C., Cambridge University Press, Cambridge, UK and New York, NY, USA, 2014.
- Blumberg, K. O., Walsh, M. P., and Pera, C.: Low-sulfur gasoline & diesel: the key to lower vehicle emissions, *The International Council on Clean Transportation (ICCT)*, Napa, CA, USA, 2003.
- Boden, T., Marland, G., and Andres, R. J.: Global, Regional, and National Fossil-Fuel CO₂ Emissions, Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, US Department of Energy, Oak Ridge, TN, USA, 2016.
- Boden, T. A., Marland, G., and Andres, R. J.: Estimates of Global, Regional, and National Annual CO₂ Emissions from Fossil-Fuel Burning, Hydraulic Cement Production, and Gas Flaring: 1950–1992, Oak Ridge National Laboratory, available at: <http://cdiac.ornl.gov/epubs/ndp/ndp030/ndp0301.htm>, 1995.
- Bond, T. C., Streets, D. G., Yarber, K. F., Nelson, S. M., Woo, J.-H., and Klimont, Z.: A technology-based global inventory of black and organic carbon emissions from combustion, *J. Geophys. Res.-Atmos.*, 109, D14203, <https://doi.org/10.1029/2003JD003697>, 2004.
- Bond, T. C., Bhardwaj, E., Dong, R., Jogani, R., Jung, S., Roden, C., Streets, D. G., and Trautmann, N. M.: Historical emissions of black and organic carbon aerosol from energy-related combustion, 1850–2000: historical BC/OC emissions, *Global Biogeochem. Cy.*, 21, GB002840, <https://doi.org/10.1029/2006GB002840>, 2007.
- BP: BP Statistical Review of World Energy, bp.com/statisticalreview (last access: 15 January 2018), 2015.
- Bruhwyler, L., Dlugokencky, E., Masarie, K., Ishizawa, M., Andrews, A., Miller, J., Sweeney, C., Tans, P., and Worthy, D.: CarbonTracker-CH₄: an assimilation system for estimating emissions of atmospheric methane, *Atmos. Chem. Phys.*, 14, 8269–8293, <https://doi.org/10.5194/acp-14-8269-2014>, 2014.
- Cofala, J., Amann, M., Klimont, Z., Kupiainen, K., and Höglund-Isaksson, L.: Scenarios of global anthropogenic emissions of air pollutants and methane until 2030, *Atmos. Environ.*, 41, 8486–8499, <https://doi.org/10.1016/j.atmosenv.2007.07.010>, 2007.
- Corbett, J. J., Fischbeck, P. S., and Pandis, S. N.: Global nitrogen and sulfur inventories for oceangoing ships, *J. Geophys. Res.-Atmos.*, 104, 3457–3470, <https://doi.org/10.1029/1998jd100040>, 1999.
- Crippa, M., Janssens-Maenhout, G., Dentener, F., Guizzardi, D., Sindelarova, K., Muntean, M., Van Dingenen, R., and Granier, C.: Forty years of improvements in European air quality: regional policy-industry interactions with global impacts, *Atmos. Chem. Phys.*, 16, 3825–3841, <https://doi.org/10.5194/acp-16-3825-2016>, 2016.
- Davidson, E. A.: The contribution of manure and fertilizer nitrogen to atmospheric nitrous oxide since 1860, *Nat. Geosci.*, 2, 659–662, <https://doi.org/10.1038/ngeo608>, 2009.
- Denier van der Gon, H. A. C., Bergström, R., Fountoukis, C., Johansson, C., Pandis, S. N., Simpson, D., and Visschedijk, A. J. H.: Particulate emissions from residential wood combustion in Europe – revised estimates and an evaluation, *Atmos. Chem. Phys.*, 15, 6503–6519, <https://doi.org/10.5194/acp-15-6503-2015>, 2015.
- Doxsey-Whitfield, E., MacManus, K., Adamo, S. B., Pistolesi, L., Squires, J., Borkovska, O., and Baptista, S. R.: Taking advantage of the improved availability of census data: a first look at the gridded population of the world, version 4, *Pap. Appl. Geogr.*, 1, 226–234, <https://doi.org/10.1080/23754931.2015.1014272>, 2015.
- EC-JRC/PBL: Emission Database for Global Atmospheric Research (EDGAR), release EDGARv4.2 FT2012, <http://edgar.jrc.ec.europa.eu> (last access: 15 January 2018), 2012.
- EC-JRC/PBL: Emission Database for Global Atmospheric Research (EDGAR), release version 4.3.1, <http://edgar.jrc.ec.europa.eu/overview.php?v=431> (last access: 15 January 2018), 2016.
- Eckhardt, S., Quennehen, B., Olivieri, D. J. L., Berntsen, T. K., Cherian, R., Christensen, J. H., Collins, W., Crepinsek, S., Daskalakis, N., Flanner, M., Herber, A., Heyes, C., Hodnebrog, Ø., Huang, L., Kanakidou, M., Klimont, Z., Langner, J., Law, K. S., Lund, M. T., Mahmood, R., Massling, A., Myriokefalitakis, S., Nielsen, I. E., Nøjgaard, J. K., Quaas, J., Quinn, P. K., Raut, J.-C., Rumbold, S. T., Schulz, M., Sharma, S., Skeie, R. B., Skov, H., Uttal, T., von Salzen, K., and Stohl, A.: Current model capabilities for simulating black carbon and sulfate concentrations in the Arctic atmosphere: a multi-model evaluation using a comprehensive measurement data set, *Atmos. Chem. Phys.*, 15, 9413–9433, <https://doi.org/10.5194/acp-15-9413-2015>, 2015.
- Economic Commission for Europe: Guidelines for reporting emissions and projections data under the conven-

- tion on long-range transboundary air pollution, http://www.ceip.at/fileadmin/inhalte/emep/2014_Guidelines/ece.eb.air.125_ADVANCE_VERSION_reporting_guidelines_2013.pdf (last access: 15 January 2018), 2014.
- EIA: US total adjusted sales of distillat fuel oil by end use, http://www.eia.gov/dnav/pet/pet_cons_821dsta_dcu_nus_a.htm (last access: 15 January 2018), 2013.
- EMEP: Officially reported emission data, Level 1, <http://www.ceip.at/> (last access: 15 January 2018), 2016.
- Endresen, O., Sorgard, E., Behrens, H. L., Brett, P. O., and Isakson, I. S. A.: A historical reconstruction of ships' fuel consumption and emissions, *J. Geophys. Res.-Atmos.*, 112, D12301, <https://doi.org/10.1029/2006jd007630>, 2007.
- Environment and Climate Change Canada: Air pollutant emissions inventory report: 1990–2014, environment and climate change Canada, Gatineau, Canada, http://www.ec.gc.ca/Pollution/A17452DA-CFC0-4222-985F-36B12FCBAA0B/APEI2017-E-Feb_15_2017.pdf, 2016.
- Environment Canada: National emission trends for key air pollutants, 1985–2011, <http://www.ec.gc.ca/inrp-npri/default.asp?lang=en&n=0EC58C98-#sommaries> (last access: 15 January 2018), 2013.
- Environment Canada: Sulfur in liquid fuels 2005, http://publications.gc.ca/site/archivee-archived.html?url=http://publications.gc.ca/collections/collection_2014/ec/En11-6-2005-eng.pdf, 2016.
- Erismann, J. W., Sutton, M. A., Galloway, J., Klimont, Z., and Winiwarter, W.: How a century of ammonia synthesis changed the world, *Nat. Geosci.*, 1, 636–639, <https://doi.org/10.1038/ngeo325>, 2008.
- Eyring, V., Köhler, H. W., van Aardenne, J., and Lauer, A.: Emissions from international shipping: 1. the last 50 years, *J. Geophys. Res.-Atmos.*, 110, D17305, <https://doi.org/10.1029/2004JD005619>, 2005.
- Eyring, V., Isakson, I. S. A., Berntsen, T., Collins, W. J., Corbett, J. J., Endresen, O., Grainger, R. G., Moldanova, J., Schlager, H., and Stevenson, D. S.: Transport impacts on atmosphere and climate: shipping, *Transp. Impacts Atmosphere Clim. ATTICA Assess. Rep.*, 44, 4735–4771, <https://doi.org/10.1016/j.atmosenv.2009.04.059>, 2010.
- FAOSTAT: FAOSTAT-Forestry database, <http://www.fao.org/forestry/statistics/84922/en/> (last access: 15 January 2018), 2015.
- FAO: FAOSTAT database, <http://www.fao.org/faostat/en/#data> (last access: 15 January 2018), 2016.
- Fernandes, S. D., Trautmann, N. M., Streets, D. G., Roden, C. A., and Bond, T. C.: Global biofuel use, 1850–2000, *Global Biogeochem. Cy.*, 21, GB2019, <https://doi.org/10.1029/2006GB002836>, 2007.
- Fletcher, M. E.: From coal to oil in British shipping, in: *The World of Shipping*, edited by: Williams, D. M., Ashgate Publishing, Brookfield VT., 1997.
- Foell, W., Amann, G., Carmichael, M., Hettelingh, J., Hordijk, L., and Dianwu, Z.: RAINS – Asia: An Assessment Model for Air Pollution in Asia, IIASA, 1995.
- Fouquet, R. and Pearson, P. J. G.: A thousand years of energy use in the UK, *Energ. J.*, 19, 1–41, 1998.
- Goldewijk, K. K., Beusen, A., van Drecht, G., and de Vos, M.: The HYDE 3.1 spatially explicit database of human-induced global land-use change over the past 12 000 years, *Global Ecol. Biogeogr.*, 20, 73–86, <https://doi.org/10.1111/J.1466-8238.2010.00587.X>, 2011.
- Granier, C., Bessagnet, B., Bond, T., D'Angiola, A., Gon, H. D. van der, Frost, G. J., Heil, A., Kaiser, J. W., Kinne, S., Klimont, Z., Kloster, S., Lamarque, J.-F., Liousse, C., Masui, T., Meleux, F., Mieville, A., Ohara, T., Raut, J.-C., Riahi, K., Schultz, M. G., Smith, S. J., Thompson, A., Aardenne, J. van, Werf, G. R. van der and Vuuren, D. P. van: Evolution of anthropogenic and biomass burning emissions of air pollutants at global and regional scales during the 1980–2010 period, *Climatic Change*, 109, 163, <https://doi.org/10.1007/s10584-011-0154-1>, 2011.
- Gschwandtner, G., Gschwandtner, K., Eldridge, K., Mann, C., and Mobley, D.: Historic emissions of sulfur and nitrogen oxides in the United States from 1900 to 1980, *JAPCA J. Air Waste Ma.*, 36, 139–149, <https://doi.org/10.1080/00022470.1986.10466052>, 1986.
- Guan, D. B., Liu, Z., Geng, Y., Lindner, S., and Hubacek, K.: The gigatonne gap in China's carbon dioxide inventories, *Nat. Clim. Change*, 2, 672–675, <https://doi.org/10.1038/Nclimate1560>, 2012.
- Hassler, B., McDonald, B. C., Frost, G. J., Borbon, A., Carslaw, D. C., Civerolo, K., Granier, C., Monks, P. S., Monks, S., Parrish, D. D., Pollack, I. B., Rosenlof, K. H., Ryonson, T. B., von Schneidmesser, E., and Trainer, M.: Analysis of long-term observations of NO_x and CO in megacities and application to constraining emissions inventories, *Geophys. Res. Lett.*, 43, 2016GL069894, <https://doi.org/10.1002/2016GL069894>, 2016.
- Holland, E. A., Lee-Taylor, J., Nevison, C., and Sulzman, J. M.: Global N Cycle: fluxes and N₂O mixing ratios originating from human activity, http://daac.ornl.gov/cgi-bin/dsvviewer.pl?ds_id=797, last access: 12 January 2017, 2005.
- Hong, C., Zhang, Q., He, K., Guan, D., Li, M., Liu, F., and Zheng, B.: Variations of China's emission estimates: response to uncertainties in energy statistics, *Atmos. Chem. Phys.*, 17, 1227–1239, <https://doi.org/10.5194/acp-17-1227-2017>, 2017.
- Houweling, S., Bergamaschi, P., Chevallier, F., Heimann, M., Kaminski, T., Krol, M., Michalak, A. M., and Patra, P.: Global inverse modeling of CH₄ sources and sinks: an overview of methods, *Atmos. Chem. Phys.*, 17, 235–256, <https://doi.org/10.5194/acp-17-235-2017>, 2017.
- Huo, H., Lei, Y., Zhang, Q., Zhao, L., and He, K.: China's coke industry: recent policies, technology shift, and implication for energy and the environment, *Energ. Policy*, 51, 397–404, <https://doi.org/10.1016/j.enpol.2012.08.041>, 2012.
- IEA: IEA Energy Statistics, <http://www.iea.org/statistics/> (last access: 15 January 2018), 2015.
- IEA: CO₂ Emissions from Fuel Combustion: 2016 Edition, IEA, Paris, France, 2016a.
- IEA: World energy outlook special report 2016: energy and air pollution, IEA, <https://www.iea.org/publications/freepublications/publication/WorldEnergyOutlookSpecialReport2016EnergyandAirPollution.pdf> (last access: 15 January 2018), 2016b.
- IIASA: GAINS – Sulfur Content of Fuels, doi:Scenario: TSAP_Consultation_2014, available at: <http://gains.iiasa.ac.at/models/index.html> (last access: 15 January 2018), 2014a.

- IIASA: GAINS – Sulfur Retention in Ash, doi:Scenario: TSAP_Consultation_2014, available at: <http://gains.iiasa.ac.at/models/index.html> (last access: 15 January 2018), 2014b.
- IMO: Reduction of GHG Emissions From Ships: Third IMO GHG Study 2014, MEPC-67-6-INF3-2014, 2014.
- Jacob, D. J., Turner, A. J., Maasackers, J. D., Sheng, J., Sun, K., Liu, X., Chance, K., Aben, I., McKeever, J., and Frankenberg, C.: Satellite observations of atmospheric methane and their value for quantifying methane emissions, *Atmos. Chem. Phys.*, 16, 14371–14396, <https://doi.org/10.5194/acp-16-14371-2016>, 2016.
- Janssens-Maenhout, G., Crippa, M., Guizzardi, D., Dentener, F., Muntean, M., Pouliot, G., Keating, T., Zhang, Q., Kurokawa, J., Wankmüller, R., Denier van der Gon, H., Kuenen, J. J. P., Klimont, Z., Frost, G., Darras, S., Koffi, B., and Li, M.: HTAP_v2.2: a mosaic of regional and global emission grid maps for 2008 and 2010 to study hemispheric transport of air pollution, *Atmos. Chem. Phys.*, 15, 11411–11432, <https://doi.org/10.5194/acp-15-11411-2015>, 2015.
- Kanaya, Y., Pan, X., Miyakawa, T., Komazaki, Y., Taketani, F., Uno, I., and Kondo, Y.: Long-term observations of black carbon mass concentrations at Fukue Island, western Japan, during 2009–2015: constraining wet removal rates and emission strengths from East Asia, *Atmos. Chem. Phys.*, 16, 10689–10705, <https://doi.org/10.5194/acp-16-10689-2016>, 2016.
- Kaur, R., Wani, S. P., Singh, A. K., and Lal, K.: Wastewater production, treatment and use in India, in: National Report presented at the 2nd regional workshop on Safe Use of Wastewater in Agriculture, available at: http://www.ais.unwater.org/ais/pluginfile.php/356/mod_page/content/111/CountryReport_India.pdf, 2012.
- Kholod, N. and Evans, M.: Circumpolar best practices: policy and financing options for black carbon emission reductions from diesel sources, circumpolar best practices, <https://oaarchive.arctic-council.org/handle/11374/1953> (last access: 15 January 2018), 2015.
- Klein Goldewijk, K., Beusen, A., and Janssen, P.: Long-term dynamic modeling of global population and built-up area in a spatially explicit way: HYDE 3.1, Holocene, 20, 565–573, <https://doi.org/10.1177/0959683609356587>, 2010.
- Klimont, Z., Cofala, J., Xing, J., Wei, W., Zhang, C., Wang, S., Kejun, J., Bhandari, P., Mathur, R., Purohit, P., Rafaj, P., Chambers, A., Amann, M., and Hao, J.: Projections of SO₂, NO_x and carbonaceous aerosols emissions in Asia, *Tellus B*, 61, 602–617, <https://doi.org/10.1111/j.1600-0889.2009.00428.x>, 2009.
- Klimont, Z., Kupiainen, K., Heyes, C., Purohit, P., Cofala, J., Rafaj, P., Borken-Kleefeld, J., and Schöpp, W.: Global anthropogenic emissions of particulate matter including black carbon, *Atmos. Chem. Phys.*, 17, 8681–8723, <https://doi.org/10.5194/acp-17-8681-2017>, 2017a.
- Klimont, Z., Hoglund-Isaksson, L., Heyes, C., Rafaj, P., Schopp, W., Cofala, J., Purohit, P., Borken-Kleefeld, J., Kupiainen, K., Kiesewetter, G., Winiwarter, W., Amann, M., Zhao, B., Wang, S., Bertok, I., and Sander, R.: Global scenarios of air pollutants and methane: 1990–2050, ACP, in preparation, 2017b.
- Korsbakken, J. I., Peters, G. P., and Andrew, R. M.: Uncertainties around reductions in China's coal use and CO₂ emissions, *Nat. Clim. Change*, 6, 687–690, <https://doi.org/10.1038/nclimate2963>, 2016.
- Kurokawa, J., Ohara, T., Morikawa, T., Hanayama, S., Janssens-Maenhout, G., Fukui, T., Kawashima, K., and Akimoto, H.: Emissions of air pollutants and greenhouse gases over Asian regions during 2000–2008: Regional Emission inventory in ASia (REAS) version 2, *Atmos. Chem. Phys.*, 13, 11019–11058, <https://doi.org/10.5194/acp-13-11019-2013>, 2013a.
- Kurokawa, J., Ohara, T., Morikawa, T., Hanayama, S., Janssens-Maenhout, G., Fukui, T., Kawashima, K., and Akimoto, H.: Emissions of air pollutants and greenhouse gases over Asian regions during 2000–2008: Regional Emission inventory in ASia (REAS) version 2, *Atmos. Chem. Phys.*, 13, 11019–11058, <https://doi.org/10.5194/acp-13-11019-2013>, 2013b.
- Lamarque, J.-F., Bond, T. C., Eyring, V., Granier, C., Heil, A., Klimont, Z., Lee, D., Liousse, C., Mieville, A., Owen, B., Schultz, M. G., Shindell, D., Smith, S. J., Stehfest, E., Van Aardenne, J., Cooper, O. R., Kainuma, M., Mahowald, N., McConnell, J. R., Naik, V., Riahi, K., and van Vuuren, D. P.: Historical (1850–2000) gridded anthropogenic and biomass burning emissions of reactive gases and aerosols: methodology and application, *Atmos. Chem. Phys.*, 10, 7017–7039, <https://doi.org/10.5194/acp-10-7017-2010>, 2010.
- Lee, D. S., Pitari, G., Grewe, V., Gierens, K., Penner, J. E., Petzold, A., Prather, M. J., Schumann, U., Bais, A., Bernsten, T., Iachetti, D., Lim, L. L., and Sausen, R.: Transport impacts on atmosphere and climate: aviation, *Atmos. Environ.*, 44, 4678–4734, <https://doi.org/10.1016/j.atmosenv.2009.06.005>, 2009.
- Li, M., Zhang, Q., Kurokawa, J.-I., Woo, J.-H., He, K., Lu, Z., Ohara, T., Song, Y., Streets, D. G., Carmichael, G. R., Cheng, Y., Hong, C., Huo, H., Jiang, X., Kang, S., Liu, F., Su, H., and Zheng, B.: MIX: a mosaic Asian anthropogenic emission inventory under the international collaboration framework of the MICS-Asia and HTAP, *Atmos. Chem. Phys.*, 17, 935–963, <https://doi.org/10.5194/acp-17-935-2017>, 2017.
- Liu, F., Zhang, Q., Tong, D., Zheng, B., Li, M., Huo, H., and He, K. B.: High-resolution inventory of technologies, activities, and emissions of coal-fired power plants in China from 1990 to 2010, *Atmos. Chem. Phys.*, 15, 13299–13317, <https://doi.org/10.5194/acp-15-13299-2015>, 2015a.
- Liu, Z., Guan, D., Wei, W., Davis, S. J., Ciais, P., Bai, J., Peng, S., Zhang, Q., Hubacek, K., Marland, G., Andres, R. J., Crawford-Brown, D., Lin, J., Zhao, H., Hong, C., Boden, T. A., Feng, K., Peters, G. P., Xi, F., Liu, J., Li, Y., Zhao, Y., Zeng, N., and He, K.: Reduced carbon emission estimates from fossil fuel combustion and cement production in China, *Nature*, 524, 335–338, <https://doi.org/10.1038/nature14677>, 2015b.
- Liu, F., Zhang, Q., A. R. J. van der, Zheng, B., Tong, D., Yan, L., Zheng, Y., and He, K.: Recent reduction in NO_x emissions over China: synthesis of satellite observations and emission inventories, *Environ. Res. Lett.*, 11, 114002, <https://doi.org/10.1088/1748-9326/11/11/114002>, 2016.
- Lu, Z., Zhang, Q., and Streets, D. G.: Sulfur dioxide and primary carbonaceous aerosol emissions in China and India, 1996–2010, *Atmos. Chem. Phys.*, 11, 9839–9864, <https://doi.org/10.5194/acp-11-9839-2011>, 2011.
- Ludek and Holub: The Century of Petrol – The History of the Refining Industry, in: The Czech Lands, Edited by: Jaroslav Pánek and Oldrich Tuma, Karolinum Press, Charles University, 2009.
- McLinden, C. A., Fioletov, V., Shephard, M. W., Krotkov, N., Li, C., Martin, R. V., Moran, M. D., and Joiner, J.: Space-based detec-

- tion of missing sulfur dioxide sources of global air pollution, *Nat. Geosci.*, 9, 496–500, <https://doi.org/10.1038/ngeo2724>, 2016.
- Mester, Z. C.: Meeting sulfur specifications for 2000 and beyond, San Francisco, available at: <http://www.environmentalconsultation.com/Meeting%20Sulfur%20Specifications%20for%202000%20and%20Beyond.pdf> (last access: 15 January 2018), 2000.
- Mitchell, B. R.: International historical statistics: The Americas, 1750–2000, Palgrave Macmillan, Houndmills, Basingstoke, Hampshire, New York, 2003.
- Mitchell, B. R.: International historical statistics: Europe, 1750–2005, Sixth Edition, Palgrave Macmillan, Basingstoke, Hampshire, New York, N. Y., 2007.
- Mitchell, B. R. and Brian, R.: International historical statistics: The Americas and Australasia, Macmillan, London, 1983.
- Mylona, S.: Sulphur dioxide emissions in Europe 1880–1991 and their effect on sulphur concentrations and depositions, *Tellus B*, 48, 662–689, <https://doi.org/10.1034/j.1600-0889.1996.t01-2-00005.x>, 1996.
- OECD: OECD Waste water treatment, <https://data.oecd.org/water/waste-water-treatment.htm> (last access: 15 January 2018), 2016.
- Olivier, J., Janssens-Maenhout, G., Muntean, M., and Peters, J.: Trends in global CO₂ emissions: 2015 Report, The Hague: PBL Netherlands Environmental Assessment Agency, Ispra: European Commission, Joint Research Centre, 2015.
- Pant, P. and Harrison, R. M.: Estimation of the contribution of road traffic emissions to particulate matter concentrations from field measurements: a review, *Atmos. Environ.*, 77, 78–97, <https://doi.org/10.1016/j.atmosenv.2013.04.028>, 2013.
- Parrish, D. D.: Critical evaluation of US on-road vehicle emission inventories, *Atmos. Environ.*, 40, 2288–2300, <https://doi.org/10.1016/j.atmosenv.2005.11.033>, 2006.
- Paulot, F., Jacob, D. J., Pinder, R. W., Bash, J. O., Travis, K., and Henze, D. K.: Ammonia emissions in the United States, European Union, and China derived by high-resolution inversion of ammonium wet deposition data: interpretation with a new agricultural emissions inventory (MASAGE_NH3), *J. Geophys. Res.-Atmos.*, 119, 4343–4364, <https://doi.org/10.1002/2013JD021130>, 2014.
- Pretorius, I., Piketh, S., Burger, R., and Neomagus, H.: A perspective on South African coal fired power station emissions, *J. Energy South Afr.*, 26, 27–40, 2015.
- Purohit, P., Amann, M., and Mathur, R.: GAINS-Asia: Scenarios for Cost-Effective Control of Air Pollution and Greenhouse Gases in India, <http://pure.iiasa.ac.at/9379/1/XO-10-061.pdf> (last access: 15 January 2018), 2010.
- Qin, Y. and Xie, S. D.: Spatial and temporal variation of anthropogenic black carbon emissions in China for the period 1980–2009, *Atmos. Chem. Phys.*, 12, 4825–4841, <https://doi.org/10.5194/acp-12-4825-2012>, 2012.
- Rao, S., Klimont, Z., Smith, S. J., Van Dingenen, R., Dentener, F., Bouwman, L., Riahi, K., Amann, M., Bodirsky, B. L., van Vuuren, D. P., Aleluia Reis, L., Calvin, K., Drouet, L., Fricko, O., Fujimori, S., Gernaat, D., Havlik, P., Harmsen, M., Hasegawa, T., Heyes, C., Hilaire, J., Luderer, G., Masui, T., Stehfest, E., Strefler, J., van der Sluis, S., and Tavoni, M.: Future air pollution in the Shared Socio-economic Pathways, *Global Environ. Chang.*, 42, 346–358, <https://doi.org/10.1016/j.gloenvcha.2016.05.012>, 2016.
- Rowe, J. and Morrison, J.: Heart of a mountain, soul of a town: the story of Algoma Ore and the town of Wawa, Privately Published, Wawas, Ontario, available at: <http://www.ourroots.ca/page.aspx?id=910392>, 1999.
- Ryaboshapko, A. G., Brukhanov, P. A., Gromov, S. A., Proshina, Y. V., and Afinogenova, O. G.: Anthropogenic Emissions of Oxidized Sulfur and Nitrogen into the Atmosphere of the Former Soviet Union in 1985 and 1990, Department of Meteorology, Stockholm University, 1996.
- Sanger, R. P.: Motor vehicle emission regulations and fuel specifications – Part 2: Detailed information and historic review (1970–1996), <https://www.concawe.eu/publication/report-no-697/> (last access: 15 January 2018), 1997.
- Schultz, M. and Sebastian, R.: Emission data sets and methodologies for estimating emissions, http://retro-archive.iek.fz-juelich.de/data/documents/reports/D1-6_final.pdf (last access: 15 January 2018), 2007.
- Schumann, U. and Huntrieser, H.: The global lightning-induced nitrogen oxides source, *Atmos. Chem. Phys.*, 7, 3823–3907, <https://doi.org/10.5194/acp-7-3823-2007>, 2007.
- Sharma, S., Goel, A., Gupta, D., Kumar, A., Mishra, A., Kundu, S., Chatani, S., and Klimont, Z.: Emission inventory of non-methane volatile organic compounds from anthropogenic sources in India, *Atmos. Environ.*, 102, 209–219, <https://doi.org/10.1016/j.atmosenv.2014.11.070>, 2015.
- Simachaya, W.: Thailand’s coal-fired power plant pollution control, <https://www.concawe.eu/publication/report-no-697/> (last access: 15 January 2018), 2015.
- Simon, H., Reff, A., Wells, B., Xing, J., and Frank, N.: Ozone trends across the United States over a period of decreasing NO_x and VOC emissions, *Environ. Sci. Technol.*, 49, 186–195, <https://doi.org/10.1021/es504514z>, 2015.
- Smith, S. J., van Aardenne, J., Klimont, Z., Andres, R. J., Volke, A., and Delgado Arias, S.: Anthropogenic sulfur dioxide emissions: 1850–2005, *Atmos. Chem. Phys.*, 11, 1101–1116, <https://doi.org/10.5194/acp-11-1101-2011>, 2011.
- Smith, T. W. P., Jalkanen, J. P., Anderson, B. A., Corbett, J. J., Faber, J., Hanayama, S., O’Keefe, E., Parker, S., Johansson, L., Aldous, L., Raucci, C., Traut, M., Ettinger, S., Nelissen, D., Lee, D. S., Ng, S., Agrawal, A., Winebrake, J. J., Hoen, M., Chesworth, S., and Pandey, A.: Reduction of GHG Emissions from Ships: Third IMO GHG Study 2014 – Final Report, IMO, available at: http://www.imo.org/en/OurWork/Environment/PollutionPrevention/AirPollution/Documents/Third_Greenhouse_Gas_Study/GHG3_Executive_Summary_and_Report.pdf (last access: 15 January 2018), 2014.
- South Korea National Institute of Environmental Research: National air pollutants emission service, <http://airemiss.nier.go.kr/> (last access: 15 January 2018), 2016.
- Stohl, A., Aamaas, B., Amann, M., Baker, L. H., Bellouin, N., Berntsen, T. K., Boucher, O., Cherian, R., Collins, W., Daskalakis, N., Dusinska, M., Eckhardt, S., Fuglestedt, J. S., Harju, M., Heyes, C., Hodnebrog, Ø., Hao, J., Im, U., Kanakidou, M., Klimont, Z., Kupiainen, K., Law, K. S., Lund, M. T., Maas, R., MacIntosh, C. R., Myhre, G., Myriokefalitakis, S., Olivie, D., Quaas, J., Quennehen, B., Raut, J.-C., Rumbold, S. T., Samset, B. H., Schulz, M., Seland, Ø., Shine, K. P., Skeie, R. B., Wang, S., Yttri, K. E., and Zhu, T.: Evaluating the climate

- and air quality impacts of short-lived pollutants, *Atmos. Chem. Phys.*, 15, 10529–10566, <https://doi.org/10.5194/acp-15-10529-2015>, 2015.
- Streets, D. G., Canty, T., Carmichael, G. R., de Foy, B., Dickerson, R. R., Duncan, B. N., Edwards, D. P., Haynes, J. A., Henze, D. K., Houyoux, M. R., Jacob, D. J., Krotkov, N. A., Lamsal, L. N., Liu, Y., Lu, Z., Martin, R. V., Pfister, G. G., Pinder, R. W., Salawitch, R. J., and Wecht, K. J.: Emissions estimation from satellite retrievals: a review of current capability, *Atmos. Environ.*, 77, 1011–1042, <https://doi.org/10.1016/j.atmosenv.2013.05.051>, 2013.
- TEPA: Taiwan Emission Data System, https://erdb.epa.gov.tw/eng/DataRepository/EnvMonitor/ReportInspectAirTEDS.aspx?topic1=Environmental_and_Biological_Monitoring&topic2=Air&subject=Air_Quality (last access: 15 January 2018), 2016.
- The World Bank: World Development Indicators, databank.worldbank.org/data/download/WDI_excel.zip (last access: 15 January 2018), 2016.
- Travis, K. R., Jacob, D. J., Fisher, J. A., Kim, P. S., Marais, E. A., Zhu, L., Yu, K., Miller, C. C., Yantosca, R. M., Sulprizio, M. P., Thompson, A. M., Wennberg, P. O., Crouse, J. D., St. Clair, J. M., Cohen, R. C., Laughner, J. L., Dibb, J. E., Hall, S. R., Ullmann, K., Wolfe, G. M., Pollack, I. B., Peischl, J., Neuman, J. A., and Zhou, X.: Why do models overestimate surface ozone in the Southeast United States?, *Atmos. Chem. Phys.*, 16, 13561–13577, <https://doi.org/10.5194/acp-16-13561-2016>, 2016.
- Tushingham, M.: Regional Composition of Gasoline and Diesel, Environment Canada, 1996.
- UK DEFRA: National Atmospheric Emissions Inventory (NAEI): emissions of air quality pollutants 1990–2013, http://naei.defra.gov.uk/reports/reports?report_id=853 (last access: 15 January 2018), 2015.
- UN: World Urbanization Prospects: The 2014 Revision, https://esa.un.org/unpd/wup/CD-ROM/WUP2014_XLS_CD_FILES/WUP2014-F01-Total_Urban_Rural.xls (last access: 15 January 2018), 2014.
- UN: World Population Prospects: The 2015 Revision, <http://esa.un.org/unpd/wpp/DVD/> (last access: 15 January 2018), 2015.
- UN: UN Energy Statistics Database, UN, <http://unstats.un.org/unsd/energy/> (last access: 15 January 2018), 2016.
- UNFCCC: National Inventory Submissions of Annex I Parties to the UNFCCC, <http://unfccc.int/> (last access: 15 January 2018), 2015.
- US EPA: Report to Congress On Black Carbon, EPA report 450/R-12-001, US Environmental Protection Agency, Washington, DC, available at: <https://www3.epa.gov/airquality/blackcarbon/2012report/fullreport.pdf>, 2012a.
- US EPA: Inventory of US Greenhouse Gas Emissions and Sinks 1990–2014, EPA report 430-R-16-002, US Environmental Protection Agency, Washington DC, available at: <https://www.epa.gov/sites/production/files/2016-04/documents/us-ghg-inventory-2016-main-text.pdf>, 2012b.
- US EPA: 2011 National Emissions Inventory (NEI) v2, <https://www.epa.gov/air-emissions-inventories/2011-national-emissions-inventories> (last access: 15 January 2018), 2013.
- US EPA: Inventory of US Greenhouse Gas Emissions and Sinks: 1990–2013, EPA report 430-R-15-004: US Environmental Protection Agency, Washington, DC, available at: <https://www.epa.gov/sites/production/files/2016-03/documents/us-ghg-inventory-2015-main-text.pdf>, 2015.
- US EPA: Inventory of US Greenhouse Gas Emissions and Sinks: 1990–2014, US EPA, Table 3-25, <https://www.epa.gov/ghgemissions/us-greenhouse-gas-inventory-report-1990-2014> (last access: 15 January 2018), 2016a.
- US EPA: Air Pollutant Emissions Trends Data: Average Annual Emissions, https://www.epa.gov/sites/production/files/2015-07/national_tier1_caps.xlsx (last access: 15 January 2018), 2016b.
- van Aardenne, J. A., Dentener, F. J., Olivier, J. G. J., Goldewijk, C. G. M. K., and Lelieveld, J.: A $1^\circ \times 1^\circ$ resolution data set of historical anthropogenic trace gas emissions for the period 1890–1990, *Global Biogeochem. Cy.*, 15, 909–928, <https://doi.org/10.1029/2000GB001265>, 2001.
- van Marle, M. J. E., Kloster, S., Magi, B. I., Marlon, J. R., Daniu, A.-L., Field, R. D., Arneth, A., Forrest, M., Hantson, S., Kehrwald, N. M., Knorr, W., Lasslop, G., Li, F., Mangeon, S., Yue, C., Kaiser, J. W., and van der Werf, G. R.: Historic global biomass burning emissions for CMIP6 (BB4CMIP) based on merging satellite observations with proxies and fire models (1750–2015), *Geosci. Model Dev.*, 10, 3329–3357, <https://doi.org/10.5194/gmd-10-3329-2017>, 2017.
- van Vuuren, D. P., Edmonds, J., Kainuma, M., Riahi, K., Thomson, A., Hibbard, K., Hurtt, G. C., Kram, T., Krey, V., Lamarque, J.-F., Masui, T., Meinshausen, M., Nakicenovic, N., Smith, S. J., and Rose, S. K.: The representative concentration pathways: an overview, *Climatic Change*, 109, 5, <https://doi.org/10.1007/s10584-011-0148-z>, 2011.
- Wang, S. X., Zhao, B., Cai, S. Y., Klimont, Z., Nielsen, C. P., Morikawa, T., Woo, J. H., Kim, Y., Fu, X., Xu, J. Y., Hao, J. M., and He, K. B.: Emission trends and mitigation options for air pollutants in East Asia, *Atmos. Chem. Phys.*, 14, 6571–6603, <https://doi.org/10.5194/acp-14-6571-2014>, 2014.
- Wiedinmyer, C., Yokelson, R. J., and Gullett, B. K.: Global emissions of trace gases, particulate matter, and hazardous air pollutants from open burning of domestic waste, *Environ. Sci. Technol.*, 48, 9523–9530, <https://doi.org/10.1021/es502250z>, 2014.
- Winijkul, E., Fierce, L., and Bond, T. C.: Emissions from residential combustion considering end-uses and spatial constraints: Part I, methods and spatial distribution, *Atmos. Environ.*, 125, 126–139, <https://doi.org/10.1016/j.atmosenv.2015.10.013>, 2016.
- Wu, Q. R., Wang, S. X., Zhang, L., Song, J. X., Yang, H., and Meng, Y.: Update of mercury emissions from China's primary zinc, lead and copper smelters, 2000–2010, *Atmos. Chem. Phys.*, 12, 11153–11163, <https://doi.org/10.5194/acp-12-11153-2012>, 2012.
- Xu, Y., Williams, R. H., and Socolow, R. H.: China's rapid deployment of SO₂ scrubbers, *Energ. Environ. Sci.*, 2, 459–465, <https://doi.org/10.1039/B901357C>, 2009.
- Zhang, Q., Streets, D. G., He, K., and Klimont, Z.: Major components of China's anthropogenic primary particulate emissions, *Environ. Res. Lett.*, 2, 045027, <https://doi.org/10.1088/1748-9326/2/4/045027>, 2007.
- Zhang, Q., He, K., and Huo, H.: Policy: Cleaning China's air, *Nature*, 484, 161–162, <https://doi.org/10.1038/484161a>, 2012.
- Zhao, B., Wang, S. X., Liu, H., Xu, J. Y., Fu, K., Klimont, Z., Hao, J. M., He, K. B., Cofala, J., and Amann, M.: NO_x emissions in China: historical trends and future perspectives, *At-*

- mos. Chem. Phys., 13, 9869–9897, <https://doi.org/10.5194/acp-13-9869-2013>, 2013a.
- Zhao, Y., Zhang, J., and Nielsen, C. P.: The effects of recent control policies on trends in emissions of anthropogenic atmospheric pollutants and CO₂ in China, Atmos. Chem. Phys., 13, 487–508, <https://doi.org/10.5194/acp-13-487-2013>, 2013b.
- Zhou, Y., Li, J., Chang, C.-C., Wei, S., Ni, Y., and Zhang, Y.: Status and development for municipal wastewater reuse in China, IEEE, 3183–3186, 2011.