

History of Neural Stem Cell Research and Its Clinical Application

Yasushi TAKAGI¹

¹Department of Neurosurgery, Kyoto University Graduate School of Medicine, Sakyo, Kyoto

Abstract

“Once development was ended...in the adult centers, the nerve paths are something fixed and immutable. Everything may die, nothing may be regenerated,” wrote Santiago Ramón y Cajal, a Spanish neuro-anatomist and Nobel Prize winner and the father of modern neuroscience. This statement was the central dogma in neuroscience for a long time. However, in the 1960s, neural stem cells (NSCs) were discovered. Since then, our knowledge about NSCs has continued to grow. This review focuses on our current knowledge about NSCs and their surrounding microenvironment. In addition, the clinical application of NSCs for the treatment of various central nervous system diseases is also summarized.

Key words: neural stem cell, neurogenesis, clinical application

Introduction

“Once development was ended...in the adult centers, the nerve paths are something fixed and immutable. Everything may die, nothing may be regenerated,” wrote Santiago Ramón y Cajal, a Spanish neuro-anatomist and Nobel Prize winner and the father of modern neuroscience. This has been the central dogma in neuroscience for a long time. However, in the 1960s, neural stem cells (NSCs) were discovered. Since then, a great amount of research has focused on increasing our understanding of NSCs.

The method of neural induction has also been clarified, and in 2006, Takahashi and Yamanaka first reported on induced pluripotent stem (iPS) cells.¹⁾ Numerous studies have provided new hope for the development of cell therapy to treat central nervous system (CNS) diseases. Recently, clinical trials using stem cells to treat CNS diseases have been undertaken. In this review, the history and characteristics of NSCs are described, and the clinical applications of stem cells, mainly focusing on NSCs, are introduced.

Adult NSC and Neurogenesis

I. The discovery of adult NSCs and neurogenesis

The discovery of adult NSCs and neurogenesis occurred in the 1960s. Gage and Temple have summarized the events in a recent review.²⁾ The

observations of cell division and differentiation in the adult brain emerged from studies of brain development and were greatly advanced by the application of tritiated thymidine, which incorporates into the DNA of dividing cells and can be detected by autoradiography. Using this labeling technique, Smart and Leblond observed and concluded that glial cells were likely dividing throughout the parenchyma.³⁾ Following these pioneering studies, Altman, using the same techniques, observed dividing cells in the subventricular zone (SVZ) and speculated that neurogenesis occurred in the adult rat and cat dentate gyrus (DG).^{4,5)} Then, in 1965, Altman and Das provided the first strong evidence for neurogenesis in the adult brain,⁶⁾ reporting on the migration of cells born postnatally in the SVZ, which matured into neurons in the olfactory bulb. In 1969, Altman was the first to describe the rostral migratory stream, which is located between the SVZ and olfactory bulb and serves as the migratory path for the NSCs born in the SVZ.⁷⁾ Surprisingly, there was little follow-up to these discoveries for about 10 years.

The next phase in the early history of adult neurogenesis focused on the avian brain, where Goldman and Nottebohm first detected what they reported as neurogenesis in adult birds.⁸⁾ Then, in the 1990s, there was the first evidence that proliferation levels of the early progenitor cells and subsequent numbers of newborn neurons are regulated; Gould et al. demonstrated that stress levels negatively affect the number of proliferating cells in the DG.⁹⁾ This

Received December 18, 2015; Accepted January 15, 2016

finding was followed by a series of observations demonstrating that neurogenesis could be substantially increased by running,¹⁰⁾ that housing animals even for short periods in complex enriched environments robustly increased the number of surviving newborn neurons,¹¹⁾ that learning itself could influence adult neurogenesis,^{12,13)} and that antidepressant drugs, as well as alcohol,¹⁴⁾ could influence components of the adult neurogenesis process.¹⁵⁾ Around this same time, neurogenesis was shown to decrease with age, but persist throughout life.¹⁶⁾

The next development was the advancement in immunohistological techniques combined with the application of confocal microscopy, which allowed researchers to visualize adult neurogenesis and to follow individually labeled cells. Importantly, the application of stereological techniques for quantifying dividing cells [in particular cells labeled with bromodeoxyuridine (BrdU)] and neuron-specific antibodies (initially NeuN), has allowed researchers to determine the absolute numbers of dividing and surviving cells in various brain regions. These techniques convincingly demonstrated that dividing cells from the DG indeed differentiated into neurons.^{17,18)} Subsequently, through the use of BrdU labeling in cancer patients, these techniques were also applied to determine that adult neurogenesis also takes place in humans.¹⁹⁾

Another important development was the establishment of methods to isolate, propagate, and differentiate neuronal progenitors from the adult CNS in special culture conditions. This breakthrough was first achieved by dissecting the lateral wall of the lateral ventricle to obtain cells from the

SVZ; these cells were then expanded in culture in proliferating populations that are now referred to as “neurospheres.”^{20,21)} The ability to isolate, maintain, expand, and differentiate these precursor cells *in vitro* allowed for researchers to explore in greater detail the cellular and molecular nature of these cells, as well as the mechanisms that regulate their behavior.

II. NSCs in the adult mammalian brain

A neurogenic niche is a region where neurogenesis takes place.²²⁾ There are two major neurogenic niches that exist in the adult mammalian brain where endogenous NSCs reside—the SVZ lining the lateral ventricles and the subgranular zone (SGZ) within the DG of the hippocampus (Fig. 1). The SVZ resides next to the ependymal cell layer, which is directly exposed to the cerebral spinal fluid and separates the ventricular space from the SVZ (Fig. 2A). Adult NSCs from the SVZ (also named “type B cells”) extend a basal process that terminates on blood vessels, as well as an apical process with a primary cilium that pokes through the ependymal cell layer to contact the cerebrospinal fluid (CSF) in the ventricle.²³⁾ Type B NSCs give rise to transient amplifying progenitors (C cells),²⁴⁾ which undergo multiple divisions before becoming neuroblasts (A cells). Neuroblasts then form a chain and migrate into the olfactory bulb where they radially migrate and differentiate into different subtypes of interneurons. Radial glia-like NSCs (named RGLs or type 1 cells) in the SGZ, which reside at the border between the inner granule cell (GC) layer and hilus, give rise to intermediate progenitor cells (IPCs),²⁵⁾ which exhibit




Fig. 1 Behavior of neural stem cells (NSCs) in the adult rodent brain. A sagittal view of the adult rodent brain, focusing on two major niches where adult NSCs reside: the subventricular zone (SVZ) and the subgranular zone (SGZ). The SVZ is located along the lateral ventricle in the forebrain and the SGZ is located in the hippocampus along the dentate granule cell layer, where it abuts the hilus.²²⁾




Fig. 2 Adult neural stem cell (NSC) niches. **A:** A schematic diagram depicting cellular and molecular components of the SVZ niche. Ependymal cells line the lateral ventricle and border the SVZ. Radial glia-like neural stem cells (B cells) reside along the ependymal zone in the SVZ and extend a radial process to contact blood vessels. They also extend a single cilium through the ependymal rosettes to contact the cerebrospinal fluid in the ventricular space. Radial glia-like NSCs generate transit-amplifying cells (C cells), which generate neuroblasts (A cells). Neuroblasts migrate along the rostral migratory stream to the olfactory bulb, where they differentiate into olfactory bulb interneurons. In addition to the aforementioned cell types, astrocytes and microglia contribute to the cellular architecture of the niche.²²⁾ **B:** A schematic diagram depicting cellular and molecular components of the SGZ niche. Radial glia-like NSCs (type I cells) reside in the SGZ and extend a radial process through the granule cell layer of the dentate gyrus into the molecular layer. Radial glia-like NSCs generate intermediate progenitor cells (IPCs), which generate neuroblasts, and these progenitor cells are closely associated with the vasculature. Neuroblasts differentiate into dentate granule cells, which migrate into the granule cell layer of the dentate gyrus. In addition to the aforementioned cell types, astrocytes, microglia, and interneurons contribute to the cellular architecture of the niche. CSF: cerebrospinal fluid.²²⁾

limited rounds of proliferation before generating neuroblasts (Fig. 2B).²⁶⁾ Neuroblasts tangentially migrate along the SGZ and develop into immature neurons, which radially migrate into the GC layer to differentiate into dentate granule neurons.²⁷⁾

Stem cells are characterized by two fundamental properties: the ability to self-renew and the ability to rise to differentiated progeny (Fig. 3A). It had long been postulated that adult neurogenesis originates from tri-potent NSCs with the capacity to generate neurons, astrocytes, and oligodendrocytes. The existence of self-renewing, multipotent adult NSCs was originally suggested by the long-term expansion and differentiation of neurospheres (nonadherent, spherical cultures of clonally derived precursors), or monolayer cultures that differentiated into three neural lineages.^{21,28)} However, recent genetic fate-mapping and clonal lineage-tracing of NSCs in the adult hippocampus *in vivo* have determined that these cells generate neurons and astrocytes, but not oligodendrocytes as originally thought.²⁹⁾ In the adult SVZ, results from population fate-mapping studies had suggested the generation of both neurons and oligodendrocytes, but recent *in vivo* clonal analysis has determined only neuronal lineages from individual NSCs.³⁰⁾ Nevertheless, *in vitro* time-lapse analysis revealed the generation

of either neurons or oligodendrocytes from acutely isolated individual precursor cells, but never both (Fig. 3B).³¹⁾

Studies using an anti-mitotic drug to eliminate dividing precursors showed that adult NSCs are largely quiescent *in vivo*.³²⁾ The quiescent state has long been viewed as dormant and passive, and quiescence is thought to allow adult stem cells to withstand metabolic stress and to preserve genome integrity over a lifetime. Emerging evidence suggests just the opposite. Single-cell transcriptome analysis of quiescent adult SGZ NSCs revealed active expression of various receptors for niche signals and downstream signaling components, the majority of which are downregulated once NSCs become activated.³³⁾ In most adult somatic stem cell systems, quiescent and active stem cell populations co-exist.³⁴⁾ The most obvious example of heterogeneity among adult NSCs is the difference in their progeny between two niches: SVZ NSCs generate olfactory bulb interneurons and corpus callosum oligodendrocytes, whereas SGZ NSCs generate dentate granule neurons and astrocytes. Notably, NSCs derived from both niches generate all three neural lineages once propagated in culture with high concentrations of growth factors,^{21,28)} suggesting that the *in vivo* niche may limit adult NSC potential.




Fig. 3 Behavior of neural stem cells (NSCs) within adult niches. A schematic diagram illustrating the potential behavior of an adult stem cell (A) and, more specifically, of an adult NSC (B) over its life cycle. Adult NSCs can transition between quiescent and active states by exiting and entering the cell cycle, respectively. Once activated, NSCs choose between different modes of division. Asymmetric division is self-renewing and yields an NSC and a progenitor, whereas symmetric division yields either two NSCs (self-renewing) or two progenitors (not self-renewing). Progenitors may be fate-restricted, meaning that they can only differentiate into a particular cell type, or they may be multipotent and must make a fate choice before differentiation. It is also possible that NSCs may directly differentiate into mature glial cell types.²²⁾

III. Functions of adult neurogenesis

Once the evidence for the existence of adult neurogenesis was generally accepted, the question of its functional relevance emerged. A series of correlational studies clearly revealed that upregulated neurogenesis in the DG also increased behavioral performance in a variety of hippocampus-related tasks and, conversely, downregulated neurogenesis resulted in behavioral impairments. Experiments designed to decrease neurogenesis by irradiation, virus manipulation, antimetabolic agents, or the engineering of transgenic animals that allowed for genetic or pharmacological regulation of adult neurogenesis, confirmed a functional role for adult neurogenesis in the DG.³⁵⁾

To more completely understand the functional importance of adult neurogenesis, it is important to consider adult neurogenesis in the context of the hippocampus and its theoretical function as a whole. Individual GCs in the DG receive inputs from thousands of entorhinal cortex neurons, suggesting that they are capable of simultaneously representing a highly complex combination of spatial and object features. Several studies^{36–38)} have suggested that the DG's encoding role can be thought of as a computational perspective that has attracted considerable attention in recent years, where the DG is critical for "pattern separation." Pattern separation, as related to the DG, can be described as recoding cortical input information into a sparse, essentially orthogonal representation.^{39,40)} By manipulating the rate of adult neurogenesis, several studies have

used ablation techniques or the overexpression of adult neurogenesis in tests that reflect the computational characteristics of pattern separation to show that newborn neurons are critical for making fine discriminations between neighboring spatial locations or highly similar environments.^{41–45)} Together, these studies support the concept that a DG network dominated by young GCs is biased toward interpreting similar, but not identical inputs, as distinct, whereas older GCs are biased toward interpreting similar inputs as equivalent.

Although adult neurogenesis in the DG is now generally accepted to occur in all adult mammals, further studies are needed to determine the mechanisms involved in the functional contribution of hippocampus-mediated behaviors. Additionally, a greater understanding of how hippocampal circuits mediate behaviors is needed. Future studies focused on adult neurogenesis and its contribution to hippocampal function will greatly advance this field of study.

Clinical Application of NSCs for the Treatment of CNS Diseases

I. Parkinson disease

Parkinson disease (PD), which is characterized by an extensive loss of dopaminergic (DA) neurons in the substantia nigra pars compacta and their striatal terminals, affects more than 500,000 people in the United States, with 50,000 new cases reported annually.^{46,47)} While the etiology of idiopathic PD is not

known, several predisposing factors for dopamine depletion associated with the disease have been suggested, including programmed cell death, viral infection, and environmental toxins. As an effective treatment for PD, patients have been administered L-dihydroxyphenyl alanine (L-DOPA), a precursor of dopamine. However, long-term L-DOPA administration results in grave side effects.^{48,49)} More recently, surgical deep-brain stimulation has been adopted as a more successful treatment for PD patients.⁵⁰⁾

Since the late 1980s, the transplantation of human fetal ventral mesencephalic tissues into the striatum of PD patients has been used as a successful therapy for patients with advanced disease.^{51–54)} However, fetal tissue transplantation has been associated with ethical and religious questions, as well as problematic logistics of acquiring fetal tissues. Additionally, recent reports have indicated that the survival of transplanted fetal mesencephalic cells in the patient brain was very low, and it was difficult to obtain sufficient fetal tissues needed for transplantation.⁵⁵⁾ To overcome these difficulties, neurons with a DA phenotype have been generated from ESCs, iPSCs, MSCs, or NSCs and could serve as a practical and an effective alternative for the transplantation of fetal brain tissues. DA neurons have been generated from mouse ESCs following treatment with fibroblast growth factor 8 (FGF8) and sonic hedgehog,^{56,57)} overexpression of *Nurr1*^{58,59)} or *Bcl-XL*,⁶⁰⁾ or co-culture with a mouse bone marrow stromal cell line.⁶¹⁾ Neurons with a DA phenotype have also been generated from monkey ESCs by co-culturing with mouse bone marrow stromal cells; behavioral improvement was observed in MPTP-lesioned monkeys following intra-striatal transplantation of these cells.⁶²⁾

DA neurons have also been generated from neural progenitor cells derived from fetal brain and were shown to induce functional recovery following transplantation into the brains of parkinsonian monkeys.⁶³⁾ Transplantation of NSCs in the brain attenuates anatomical or functional deficits associated with injury or disease in the CNS via cell replacement, the release of specific neurotransmitters, and/or the production of neurotrophic factors that provide neuroprotection and promote neuronal growth.

Previous studies have reported that mouse or human ESC-derived DA neurons are efficacious in PD animal models; however, considerable safety concerns exist, such as risk of tumor formation and neural overgrowth. More recent studies have indicated that functional human DA neurons could be efficiently generated from human ES cells, and upon transplantation in rat PD models, the ES cell-derived DA neurons induced behavioral recovery.^{64–66)}

Human DA neurons derived from iPS cells may provide an ideal cellular source for PD transplantation therapy because they can be generated from the patient's own fibroblasts and do not cause immune rejection. However, the development of an effective cellular therapy for PD using iPS cells relies on the optimization of *in vitro* production of iPS cell-derived DA neurons, as well as the prevention of teratoma formation *in vivo*. A recent study reported on the generation of DA neurons from iPS cells derived from fibroblasts and improved behavior following transplantation of these DA neurons in PD model rats.⁶⁷⁾ When multiple human iPSC lines derived by virus- and protein-based reprogramming were compared, DA neurons derived from protein-based iPSCs were best suited for transplantation, because they induce gene expression and exhibit physiological and electrophysiological properties similar to human midbrain DA neurons.⁶⁸⁾ DA neurons have also been generated from iPS cells from PD patients; these DA neurons were transplanted without signs of neurodegeneration into a PD animal model. Additionally, the neurons survived at high numbers and mediated functional effects in PD animals.⁶⁹⁾ These PD iPS cell-derived DA neurons could be used for screening novel drugs in the development of PD therapeutic strategies.

More recently, human fibroblasts were directly converted into DA neuron-like cells through the use of a combination of five transcriptional factors: *Mash1*, *Ngn2*, *Sox2*, *Nurr1*, and *Pitx3*. Results showed that the reprogrammed cells expressed various markers for DA neurons. Although further research is still required, cell therapy strategies based on iPS-derived DA neurons⁷⁰⁾ or DA neurons directly converted from fibroblasts may become a promising treatment strategy for PD patients in the coming years.

II. Huntington disease

Huntington disease (HD) is an autosomal dominant neurodegenerative disorder characterized by involuntary choreic movements, cognitive impairment, and emotional disturbances.^{71,72)} Despite the identification of the HD gene and associated protein, the mechanisms involved in HD pathogenesis remain largely unknown, thereby hampering effective therapeutic interventions. The transplantation of fetal human brain tissue may serve as a useful strategy in reducing neuronal damage in the HD brain. A recent study documented improvements in motor and cognition performance in HD patients following fetal cell transplantation.⁷³⁾ This trial follows previous reports in HD experimental animals describing the positive effects of fetal striatal cell transplantation in ameliorating neuronal dysfunction⁷⁴⁾ and that striatal

graft tissue could integrate and survive within the progressively degenerated striatum in a transgenic HD mouse model.⁷⁵⁾ This latter study is consistent with results obtained from HD patients indicating survival and differentiation of implanted human fetal tissue in the affected regions.⁷⁶⁾ Cell replacement therapy using human fetal striatal grafts has shown clinical success in HD patients.

It is, however, important to note that a recent study reported neural overgrowth of grafted tissue in a HD patient who survived 5 years post-transplantation.⁷⁷⁾ Overgrown grafts were composed of neurons and glia embedded in a disorganized neuropil. This report highlights the safety concerns related to fetal cell grafts and the potential risk of neural overgrowth following transplantation in the brain of HD patients. The use of NSC transplantation to replace degenerated neurons or genetically modified NSCs producing neurotrophic factors has been employed to protect striatal neurons against excitotoxic insults.⁷⁸⁾ At present, little is known regarding whether NSC implantation prior to neuropathological damage could alter the progressive degeneration of striatal neurons and motor deficits that occur in HD. This question is important, because genetic studies of HD gene mutations⁷⁹⁾ and neuroimaging have provided details about the factors involved in HD progression,^{80,81)} suggesting that early intervention using brain transplantation could be effective in “pre-clinical” HD patients that express the mutant HD gene.

III. Amyotrophic lateral sclerosis

Amyotrophic lateral sclerosis (ALS), also known as Lou Gehrig's disease, is a relentlessly progressive, adult-onset, neurodegenerative disorder characterized by the degeneration and loss of motor neurons in the cerebral cortex, brain stem, and spinal cord, leading to muscle wasting and weakness, and eventually to death within 5 years after onset of clinical symptoms.^{82,83)} The proposed pathogenetic mechanisms of ALS, which have not been fully determined, include oxidative stress, protein aggregation, mitochondrial dysfunction, impaired axonal transport, glutamate-mediated excitotoxicity, and insufficient production of neurotrophic factors.⁸⁴⁾ To date, there is no effective treatment for patients suffering from ALS. Recent studies have indicated that it is possible to generate motor neurons in culture from stem cells, such as ESCs and NSCs.^{85–88)} A phase I trial of intraspinal injections of fetal-derived NSCs in ALS patients was conducted in the United States. Ten total injections were made into the lumbar spinal cord at a dose of 100,000 cells per injection in 12 ALS patients. Clinical assessments, ranging

from 6 months to 18 months after transplantation, demonstrated no evidence of acceleration of disease progression due to the intervention.⁸⁹⁾ A previous study reported that iPSCs isolated from an ALS patient were differentiated into motor neurons,⁹⁰⁾ suggesting that these patient-derived neurons could serve as an ideal cellular source for screening new drug candidates. Neurons and glia induced from patient-derived iPSCs are autologous and easily accessible, and also lack the problems associated with immune rejection and ethical concerns.

It is unrealistic to expect transplantation of stem cells or stem cell-derived motor neurons to replace lost neurons in ALS patients or to integrate into existing neural circuitry and restore motor function. Rather, preventing cell death in host motor neurons via provision of neurotrophic factors by transplanted stem cells or stem cell-derived motor neurons is more realistic and an achievable approach.⁹¹⁾

IV. Alzheimer disease

Alzheimer disease (AD) is characterized by the degeneration and loss of neurons and synapses throughout the brain, particularly in the basal forebrain, amygdala, hippocampus, and cortical area. Symptoms include progressive decline in memory and cognitive function, as well as dementia and premature death.^{92–94)} To date, there is no effective treatment. Acetylcholinesterase inhibitors have been shown to augment cholinergic function, but this is not curative and is only a temporary measure.

With regard to AD pathogenesis, the amyloid cascade hypothesis postulates that memory deficits are caused by increased levels of both soluble and insoluble amyloid beta (A β) peptides, which are derived from the larger amyloid precursor protein (APP) sequential proteolytic processing.^{92–94)}

Previously, a phase 1 clinical trial utilizing *ex vivo* NGF gene delivery was performed in eight mild AD patients; autologous fibroblasts genetically modified to express human NGF were implanted into the forebrain. After a mean follow-up of 22 months in six subjects, long-term adverse effects were not observed. Evaluation by Mini-Mental State Examination and Alzheimer's Disease Assessment Scale-Cognitive Subcomponent (ADAS-Cog) suggested improvement in the rate of cognitive decline, and serial positron emission tomography (PET) scans showed significant increases in cortical fluorodeoxyglucose after treatment.⁹⁵⁾ In AD patients, a dysfunctional presynaptic cholinergic system is one of the causes for the cognitive disorders involved in patients where decreased activity of choline acetyltransferase (ChAT), which is responsible for acetylcholine (ACh) synthesis, is observed.⁹⁶⁾ To date, AD therapy has been largely

based on small molecules designed to increase ACh concentration by inhibiting acetylcholinesterase.⁹⁷ Because therapies with these drugs is only palliative and does not provide protection against progressive tissue destruction, there is a continued need for effective therapies for AD patients. Stem cell-based therapeutic approaches targeting AD provide hope for fulfilling this requirement.

A recent review article indicated that stem cell transplant therapy is an extension of NSC use in other neurological treatments, such as PD and stroke, and could serve as a highly effective therapeutic approach for AD.⁹⁸

VI. Cell-based therapies against stroke

In a recent review by George and Steinberg, the use of stem cell therapy to treat stroke patients was discussed.⁹⁹ Stem cell therapy is an exciting area of research that has entered the clinical arena with multiple ongoing trials. Stem cells are pluripotent or multipotent cells that have the ability to transform into multiple cell types and are self-perpetuating. Endogenous therapeutic strategies have focused on increasing mobilization, longevity, and production of NSCs in the SVZ and dentate gyrus. Exogenous stem cell treatments refer to transplanted cells from another source into a patient. Exogenous stem cells have been delivered to the brain via the blood stream, or direct transplantation and have shown great promise in stroke animal models to enhance recovery.

VI. Endogenous stem cells

NPCs from the SVZ, which normally migrate along the rostral migratory system to the olfactory lobe, have also been shown to traverse to injured brain areas following neurological insult.¹⁰⁰ Brain ischemia results in the upregulation of endogenous NPCs and occasional differentiation into the predominant cell type of the injured region.^{101,102} Therapeutic approaches have focused on augmenting the brain's normal endogenous reaction to injury. Multiple pathways have been used to induce neurogenesis, including those triggered by numerous neurotrophic and growth factors such as GDNF, BDNF, granulocyte colony-stimulating factor (G-CSF), and insulin growth factor (IGF-1).^{103,104} Alternative mechanisms of increasing endogenous NPC proliferation include anti-inflammatory drugs like indomethacin, non-coding RNA, and hormones such as erythropoietin.¹⁰⁵⁻¹⁰⁷ The delivery of G-CSF and IGF-1 to alter key survival pathways, such as the phosphoinositide 3-kinase-Akt pathway, has been shown to reduce NPC death.¹⁰⁸ Current clinical trials are investigating the ability of G-CSF to mobilize endogenous bone marrow

cells, as well as utilizing the neuroprotective effects of G-CSF for stroke recovery.^{109,110} Increasing the number of migrating endogenous stem cells can be achieved using various chemokine receptors, such as stromal-derived factor 1 and integrin beta-1.^{111,112} However, this research is currently limited to the preclinical arena.

VII. Exogenous stem cells

Exogenous stem cells are typically divided into three categories: (1) immortalized cell lines, (2) NPCs or NSCs, and (3) bone marrow-derived hematopoietic/endothelial progenitors and stromal cells.¹¹³ Immortalized cell lines have been developed from tumor cells or from oncogene manipulation (such as *myc* in the human fetal neural cell line ReN001 of ReNeuron). NT2N cells, which are derived from teratocarcinomas, differentiate into post-mitotic neuron-like cells with the addition of retinoic acid and mitotic inhibitors,^{114,115} and have been shown to improve outcome in several ischemic models.¹¹⁶ ReNeuron's cells have shown to exhibit dose-dependent recovery in stroke rodent models,¹¹⁷ and have been engineered to be immortal only in the presence of tamoxifen to reduce the risk of tumor formation.¹¹⁸

Human NPCs are derived from embryonic and fetal tissue and have the ability to produce astrocytes, neurons, and oligodendrocytes.¹¹⁹ In stroke models, NPCs are able to migrate to the injured regions and improve recovery.¹²⁰⁻¹²² NPCs sometimes integrate into the host tissue, differentiate, and exhibit neuronal characteristics, including expression of synaptic proteins, synapse formation, and electrophysiological properties.¹²³⁻¹²⁵ Progenitor cells derived from bone marrow, umbilical cord blood, and adipose tissue have all been shown to improve recovery in stroke models.¹²⁶ Many of these sources are already used for the clinical treatment of other disorders, such as malignancy, and can be obtained from autologous harvesting.

Although many cell types are included in each of these sources and it appears that the mononuclear or marrow stromal cell component mediates recovery, it is not clear which subtype is responsible for improving functional outcomes. Multiple trials using exogenous stem cells have been performed or are ongoing.

The discovery of iPS cells created a paradigm shift in cell therapy. The ability to transform host somatic cells, such as fibroblasts, into pluripotent stem cells bypassed many of the concerns of traditional stem cell therapy, such as ethical discussions, supply limitations, and the possible requirement of immunosuppression.^{1,127,128} Further development has led to vector- and transgene-free techniques to

derive iPSCs that improve functional outcome after brain ischemia.¹²⁹⁾ A recent study was able to generate neural cells directly from mouse or human fibroblasts using transcription factors, without passing through a pluripotent phase, which may ultimately have clinical relevance.¹³⁰⁾

Apart from efficacy, safety is an important consideration to further develop cell transplantation therapy. Careful biological classification and understanding will be critical for reducing any possibility of tumor formation or adverse effects.¹³¹⁾ Immortalized cell lines (NT2N) were the first human cells used in a clinical phase I stroke trial and were implanted into the infarcted region of 12 patients, 6 months to 6 years after a basal ganglia stroke.¹³²⁾ There were no significant adverse events and functional improvement was observed in this small patient group. A subsequent phase II trial with NT2N cells implanted into the peri-infarct or peri-hemorrhagic cavity also showed no increase in adverse events.¹³³⁾ An

open-label, single-blinded, randomized trial using mesenchymal stem cells (MSCs) showed significant improvement in functional outcome based on the modified Rankin scale (a functional outcome scale with 0–3 representing the ability to walk with varying degrees of disability) in the treatment group without a difference in adverse events, and multiple other trials have indicated safety and feasibility.^{134–136)} In addition, trials using bone marrow mononuclear cells (BMMNCs) have shown safety and feasibility in the acute and chronic phases of recovery.^{137–139)} A phase 1/2A study that transplanted human-modified bone marrow-derived stromal cells were safe and feasible for direct intracerebral transplantation 6 months to 5 years post-stroke, with improvement in neurological outcomes.¹⁴⁰⁾ Many questions have been raised about the translation of cell therapy to clinical applications, and numerous clinical trials are underway (Table 1) to determine whether cell-based therapy will become the next modality for restorative stroke therapeutics.⁹⁹⁾

Table 1 List of completed or ongoing trials using exogenous stem or progenitor cells

Study type	Sponsor	Cell type	Planned enrollment	Timing of delivery	Delivery route	Status/Results
Ph1-NR-OL	Universidade Federal do Rio de Janeiro (Rio de Janeiro, Brazil)	BMMNC	15	3–90 days	IA or i.v.	Complete, no reported results
Ph1-NR-OL	Neurogen Brain and Spine Institute (Mumbai, India)	BMMNC	30	chronic	IT	No reported results
Ph2-R-OL	Manipal Acunova Ltd. (Manipal, India)	BMMNC	11	7–30 days	i.v.	Safe, feasible
Ph1/2-R-DB	University of California (Irvine, USA)	BMMNC	40	1–3 days	i.v.	Not currently recruiting
Ph1-NR-OL	The University of Texas Health Science Center (Houston, USA)	BMMNC	10	1–3 days	i.v.	Safe, feasible
Ph2-R-SB	All India Institute of Medical Sciences (New Delhi, India)	BMMNC	120	7–30 days	i.v.	Safe, feasible, no efficacy benefit
Ph1/2-NR-OL	Chaitanya Hospital (Pune, India)	BMMNC	50	chronic	IT	Recruiting
Ph1-NR-OL	Neurogen Brain and Spine Institute (Mumbai, India)	BMMNC	200	subacute/chronic	IT	Recruiting
Ph1/2-NR-OL	India	BMMNC	11	3–12 months	i.v.	Safe, feasible, improved neurologic outcomes
Ph2-R-OL	Andalusian Initiative for Advanced Therapies (Andalusia, Spain)	BMMNC	76	1–7 days	IA	Recruiting
Ph2-R-DB	Athersys, Inc. (Cleveland, USA)	multistem	126	1–2 days	i.v.	Safe, feasible, no efficacy benefit
Ph2-NR-OL	ReNeuron Limited (Guildford, UK)	CTX0E03, NSC	41	2–3 months	IC	Safe, improved neurologic outcomes

Continued

Table 1 Continued

Study type	Sponsor	Cell type	Planned enrollment	Timing of delivery	Delivery route	Status/Results
Ph1-NR-OL	ReNeuron Limited (Guildford, UK)	CTX0E03, NSC	12	6–60 months	IC	Not currently recruiting
Ph1/2- NR-OL	Ageless Regenerative Institute (FL, USA)	ASC	10	subacute	IA	Not currently recruiting
Ph1/2-R-DB	Stempeutics Research Pvt Ltd (Whitefield, India)	MSC	78	< 10 days	i.v.	Not currently recruiting
Ph1/2-R-OL-SB	South Korea	MSC	85	5–7 weeks	i.v.	Safe, feasible, improved neurologic outcomes
Ph2-R-OL	Stempeutics Research Pvt Ltd (Whitefield, India)	MSC	30	< 6 weeks	i.v.	Not currently recruiting
Ph1/2-NR-OL	Stemmedica Cell Technologies, Inc. (San Diego, USA)	MSC	35	> 6 months	i.v.	Recruiting
Ph1/2-R-DB	Instituto de Investigación Hospital Universitario La Paz (Madrid, Spain)	MSC	40	< 14 days	i.v.	Not currently recruiting
Ph1-NR-OL	Japan	MSC	12	1–4 months	i.v.	Safe, feasible, decreased infarct
Ph3-R-OL	Samsung Medical Center (Seoul, South Korea)	MSC	60	< 90 days	i.v.	recruiting
Ph1/2-R-DB	Sean Savitz (Houston, USA)	MSC	48	3–10 days	i.v.	Not currently recruiting
Ph1/2-R-DB	Southern Medical University (GuangZhou, China)	MSC, EPC	90	5 weeks	i.v.	Recruiting
Ph1/2-NR-SB	Hospital Universitario Central de Asturias (Asturias, Spain)	CD34+	20	5–9 days	IA	Safe, feasible, increased b-NGF
Ph2-R-OL	China Medical University Hospital (Taichung City, Taiwan)	CD34+	30	6–60 months	IC	Complete, no reported results
Ph1/2-NR-OL	Imperial College London (London, UK)	CD34+	5	7 days	IA	Safe, feasible, reduced infarct volume
Ph1-R-OL	Zhejiang Hospital (Zhejiang, China)	CD34+	40	< 12 months	IA	Recruiting
Ph1-NR-OL	China Medical University Hospital (Taichung City, Taiwan)	CD34+	6	6–60 months	IC	Not currently recruiting
Ph2-R-DB	Celgene Corporation (Summit, USA)	PDC	44	acute	i.v.	Stopped by sponsor
Ph1/2-NR-OL	SanBio, Inc. (Mountain View, USA)	SB623	18	6–36 months	IC	Safe, improved neurologic outcomes
Ph2-R-OL-SB	University of Pittsburgh (Pittsburgh, USA)	NT2	18	1–5 years	IC	Safe, feasible, improved neurologic outcomes in secondary endpoints
Ph1-NR-OL	University of Pittsburgh (Pittsburgh, USA)	NT2	12	6–72 months	IC	Safe, improved neurologic outcomes

ASC: adipose-derived stromal cell, DB: double blind, EPC: endothelial progenitor cell, IA: intra-arterial, IC: intracranial, i.v.: intravenous, NR: nonrandomized, NSC: neural stem cell, NT2: tetracarcoma cell-derived neurons, OEC: olfactory ensheathing cell, OL: open label, P1: Phase 1 trial, P2: Phase 2 trial, PDC: placenta-derived stem cell, R: randomized, SB: single blind, SB623: human mesenchymal stromal cells.

Conclusion

Recent scientific advancements have suggested the novel use of NSC for therapy, with numerous clinical trials underway. Results from these studies will provide valuable information for future strategies. However, the demonstration of efficacy in randomized, double-blinded trials is needed to prove the efficacy of cell therapy for the treatment of CNS diseases.

Conflicts of Interest Disclosure

None declared. The author is a member of the Japan Neurosurgical Society and has registered online self-reported COI disclosure statement forms.

References

- 1) Takahashi K, Yamanaka S: Induction of pluripotent stem cells from mouse embryonic and adult fibroblast cultures by defined factors. *Cell* 126: 663–676, 2006
- 2) Gage FH, Temple S: Neural stem cells: generating and regenerating the brain. *Neuron* 80: 588–601, 2013
- 3) Smart I, Leblond C: Evidence for division and transformations of neuroglia cells in the mouse brain, as derived from radioautography after injection of thymidine- H^3 . *J Comp Neurol* 116: 349–367, 1961
- 4) Altman J: Are new neurons formed in the brains of adult mammals? *Science* 135: 1127–1128, 1962
- 5) Altman J: Autoradiographic investigation of cell proliferation in the brains of rats and cats. *Anat Rec* 145: 573–591, 1963
- 6) Altman J, Das GD: Autoradiographic and histological evidence of postnatal hippocampal neurogenesis in rats. *J Comp Neurol* 124: 319–335, 1965
- 7) Altman J: Autoradiographic and histological studies of postnatal neurogenesis. IV. Cell proliferation and migration in the anterior forebrain, with special reference to persisting neurogenesis in the olfactory bulb. *J Comp Neurol* 137: 433–457, 1969
- 8) Goldman SA, Nottebohm F: Neuronal production, migration, and differentiation in a vocal control nucleus of the adult female canary brain. *Proc Natl Acad Sci USA* 80: 2390–2394, 1983
- 9) Gould E, Cameron HA, Daniels DC, Woolley CS, McEwen BS: Adrenal hormones suppress cell division in the adult rat dentate gyrus. *J Neurosci* 12: 3642–3650, 1992
- 10) van Praag H, Kempermann G, Gage FH: Running increases cell proliferation and neurogenesis in the adult mouse dentate gyrus. *Nat Neurosci* 2: 266–270, 1999
- 11) Kempermann G, Kuhn HG, Gage FH: More hippocampal neurons in adult mice living in an enriched environment. *Nature* 386: 493–495, 1997
- 12) Döbrössy MD, Drapeau E, Arousseau C, Le Moal M, Piazza PV, Abrous DN: Differential effects of learning on neurogenesis: learning increases or decreases the number of newly born cells depending on their birth date. *Mol Psychiatry* 8: 974–982, 2003
- 13) Gould E, McEwen BS, Tanapat P, Galea LA, Fuchs E: Neurogenesis in the dentate gyrus of the adult tree shrew is regulated by psychosocial stress and NMDA receptor activation. *J Neurosci* 17: 2492–2498, 1997
- 14) Nixon K, Crews FT: Binge ethanol exposure decreases neurogenesis in adult rat hippocampus. *J Neurochem* 83: 1087–1093, 2002
- 15) Malberg JE, Eisch AJ, Nestler EJ, Duman RS: Chronic antidepressant treatment increases neurogenesis in adult rat hippocampus. *J Neurosci* 20: 9104–9110, 2000
- 16) Kuhn HG, Dickinson-Anson H, Gage FH: Neurogenesis in the dentate gyrus of the adult rat: age-related decrease of neuronal progenitor proliferation. *J Neurosci* 16: 2027–2033, 1996
- 17) Kempermann G, Kuhn HG, Gage FH: More hippocampal neurons in adult mice living in an enriched environment. *Nature* 386: 493–495, 1997
- 18) Kuhn HG, Winkler J, Kempermann G, Thal LJ, Gage FH: Epidermal growth factor and fibroblast growth factor-2 have different effects on neural progenitors in the adult rat brain. *J Neurosci* 17: 5820–5829, 1997
- 19) Eriksson PS, Perfilieva E, Björk-Eriksson T, Alborn AM, Nordborg C, Peterson DA, Gage FH: Neurogenesis in the adult human hippocampus. *Nat Med* 4: 1313–1317, 1998
- 20) Kilpatrick TJ, Bartlett PF: Cloning and growth of multipotential neural precursors: requirements for proliferation and differentiation. *Neuron* 10: 255–265, 1993
- 21) Reynolds BA, Weiss S: Generation of neurons and astrocytes from isolated cells of the adult mammalian central nervous system. *Science* 255: 1707–1710, 1992
- 22) Bond AM, Ming GL, Song H: Adult mammalian neural stem cells and neurogenesis: five decades Later. *Cell Stem Cell* 17: 385–395, 2015
- 23) Mirzadeh Z, Merkle FT, Soriano-Navarro M, Garcia-Verdugo JM, Alvarez-Buylla A: Neural stem cells confer unique pinwheel architecture to the ventricular surface in neurogenic regions of the adult brain. *Cell Stem Cell* 3: 265–278, 2008
- 24) Doetsch F, Caillé I, Lim DA, García-Verdugo JM, Alvarez-Buylla A: Subventricular zone astrocytes are neural stem cells in the adult mammalian brain. *Cell* 97: 703–716, 1999
- 25) Seri B, García-Verdugo JM, McEwen BS, Alvarez-Buylla A: Astrocytes give rise to new neurons in the adult mammalian hippocampus. *J Neurosci* 21: 7153–7160, 2001
- 26) Berg DK, Yoon KJ, Will B, Xiao AY, Kim NS, Christian KM, Song H, Ming G: Tbr2-expressing intermediate progenitor cells in the adult mouse hippocampus are unipotent neuronal precursors with limited amplification capacity under homeostasis. *Front Biol* 10: 262–271, 2015

- 27) Sun GJ, Zhou Y, Stadel RP, Moss J, Yong JH, Ito S, Kawasaki NK, Phan AT, Oh JH, Modak N, Reed RR, Toni N, Song H, Ming GL: Tangential migration of neuronal precursors of glutamatergic neurons in the adult mammalian brain. *Proc Natl Acad Sci USA* 112: 9484–9489, 2015
- 28) Palmer TD, Takahashi J, Gage FH: The adult rat hippocampus contains primordial neural stem cells. *Mol Cell Neurosci* 8: 389–404, 1997
- 29) Bonaguidi MA, Wheeler MA, Shapiro JS, Stadel RP, Sun GJ, Ming GL, Song H: In vivo clonal analysis reveals self-renewing and multipotent adult neural stem cell characteristics. *Cell* 145: 1142–1155, 2011
- 30) Calzolari F, Michel J, Baumgart EV, Theis F, Götz M, Ninkovic J: Fast clonal expansion and limited neural stem cell self-renewal in the adult subependymal zone. *Nat Neurosci* 18: 490–492, 2015
- 31) Ortega F, Gascón S, Masserdotti G, Deshpande A, Simon C, Fischer J, Dimou L, Chichung Lie D, Schroeder T, Berninger B: Oligodendroglial and neurogenic adult subependymal zone neural stem cells constitute distinct lineages and exhibit differential responsiveness to Wnt signalling. *Nat Cell Biol* 15: 602–613, 2013
- 32) Doetsch F, Caillé I, Lim DA, García-Verdugo JM, Alvarez-Buylla A: Subventricular zone astrocytes are neural stem cells in the adult mammalian brain. *Cell* 97: 703–716, 1999
- 33) Shin J, Berg DA, Zhu Y, Shin JY, Song J, Bonaguidi MA, Enikolopov G, Nauen DW, Christian KM, Ming GL, Song H: Single-cell RNA-Seq with waterfall reveals molecular cascades underlying adult neurogenesis. *Cell Stem Cell* 17: 360–372, 2015
- 34) Li L, Clevers H: Coexistence of quiescent and active adult stem cells in mammals. *Science* 327: 542–545, 2010
- 35) Deng W, Aimone JB, Gage FH: New neurons and new memories: how does adult hippocampal neurogenesis affect learning and memory? *Nat Rev Neurosci* 11: 339–350, 2010
- 36) Morris AM, Weeden CS, Churchwell JC, Kesner RP: The role of the dentate gyrus in the formation of contextual representations. *Hippocampus* 23: 162–168, 2013
- 37) O'Reilly RC, McClelland JL: Hippocampal conjunctive encoding, storage, and recall: avoiding a trade-off. *Hippocampus* 4: 661–682, 1994
- 38) Rolls ET, Kesner RP: A computational theory of hippocampal function, and empirical tests of the theory. *Prog Neurobiol* 79: 1–48, 2006
- 39) McNaughton BL, Morris RGM: Hippocampal synaptic enhancement and information storage within a distributed memory system. *Trends Neurosci* 10: 408–415, 1987
- 40) Treves A, Rolls ET: Computational constraints suggest the need for two distinct input systems to the hippocampal CA3 network. *Hippocampus* 2: 189–199, 1992
- 41) Clelland CD, Choi M, Romberg C, Clemenson GD Jr, Fagniere A, Tyers P, Jessberger S, Saksida LM, Barker RA, Gage FH, Bussey TJ: A functional role for adult hippocampal neurogenesis in spatial pattern separation. *Science* 325: 210–213, 2009
- 42) Creer DJ, Romberg C, Saksida LM, van Praag H, Bussey TJ: Running enhances spatial pattern separation in mice. *Proc Natl Acad Sci USA* 107: 2367–2372, 2010
- 43) Gu Y, Arruda-Carvalho M, Wang J, Janoschka SR, Josselyn SA, Frankland PW, Ge S: Optical controlling reveals time-dependent roles for adult-born dentate granule cells. *Nat Neurosci* 15: 1700–1706, 2012
- 44) Nakashiba T, Cushman JD, Pelkey KA, Renaudineau S, Buhl DL, McHugh TJ, Rodriguez Barrera V, Chittajallu R, Iwamoto KS, McBain CJ, Fanselow MS, Tonegawa S: Young dentate granule cells mediate pattern separation, whereas old granule cells facilitate pattern completion. *Cell* 149: 188–201, 2012
- 45) Sahay A, Scobie KN, Hill AS, O'Carroll CM, Kheirbek MA, Burghardt NS, Fenton AA, Dranovsky A, Hen R: Increasing adult hippocampal neurogenesis is sufficient to improve pattern separation. *Nature* 472: 466–470, 2011
- 46) Kish SJ, Shannak K, Hornykiewicz O: Uneven pattern of dopamine loss in the striatum of patients with idiopathic Parkinson's disease. Pathophysiologic and clinical implications. *N Engl J Med* 318: 876–880, 1988
- 47) Agid Y: Parkinson's disease: pathophysiology. *Lancet* 337: 1321–1324, 1991
- 48) Qureshi F, Pestian J, Davis P, Zaritsky A: Effect of nebulized ipratropium on the hospitalization rates of children with asthma. *N Engl J Med* 339: 1030–1035, 1998
- 49) Lang AE, Lozano AM: Parkinson's disease. First of two parts. *N Engl J Med* 339: 1044–1053, 1998
- 50) Lyons MK: Deep brain stimulation: current and future clinical applications. *Mayo Clin Proc* 86: 662–672, 2011
- 51) Lindvall O, Brundin P, Widner H, Rehnström S, Gustavii B, Frackowiak R, Leenders KL, Sawle G, Rothwell JC, Marsden CD: Grafts of fetal dopamine neurons survive and improve motor function in Parkinson's disease. *Science* 247: 574–577, 1990
- 52) Kordower JH, Goetz CG, Freeman TB, Olanow CW: Dopaminergic transplants in patients with Parkinson's disease: neuroanatomical correlates of clinical recovery. *Exp Neurol* 144: 41–46, 1997
- 53) Dunnett SB, Björklund A: Prospects for new restorative and neuroprotective treatments in Parkinson's disease. *Nature* 399: A32–A39, 1999
- 54) Lindvall O, Björklund A: Cell therapy in Parkinson's disease. *NeuroRx* 1: 382–393, 2004
- 55) Hagell P, Brundin P: Cell survival and clinical outcome following intrastriatal transplantation in Parkinson disease. *J Neuropathol Exp Neurol* 60: 741–752, 2001
- 56) Lee SH, Lumelsky N, Studer L, Auerbach JM, McKay RD: Efficient generation of midbrain and hindbrain neurons from mouse embryonic stem cells. *Nat Biotechnol* 18: 675–679, 2000

- 57) Kim JH, Auerbach JM, Rodríguez-Gómez JA, Velasco I, Gavin D, Lumelsky N, Lee SH, Nguyen J, Sánchez-Pernaute R, Bankiewicz K, McKay R: Dopamine neurons derived from embryonic stem cells function in an animal model of Parkinson's disease. *Nature* 418: 50–56, 2002
- 58) Wagner J, Akerud P, Castro DS, Holm PC, Canals JM, Snyder EY, Perlmann T, Arenas E: Induction of a midbrain dopaminergic phenotype in Nurr1-overexpressing neural stem cells by type 1 astrocytes. *Nat Biotechnol* 17: 653–659, 1999
- 59) Chung S, Sonntag KC, Andersson T, Bjorklund LM, Park JJ, Kim DW, Kang UJ, Isacson O, Kim KS: Genetic engineering of mouse embryonic stem cells by Nurr1 enhances differentiation and maturation into dopaminergic neurons. *Eur J Neurosci* 16: 1829–1838, 2002
- 60) Shim JW, Koh HC, Chang MY, Roh E, Choi CY, Oh YJ, Son H, Lee YS, Studer L, Lee SH: Enhanced in vitro midbrain dopamine neuron differentiation, dopaminergic function, neurite outgrowth, and 1-methyl-4-phenylpyridium resistance in mouse embryonic stem cells overexpressing Bcl-XL. *J Neurosci* 24: 843–852, 2004
- 61) Kawasaki H, Mizuseki K, Nishikawa S, Kaneko S, Kuwana Y, Nakanishi S, Nishikawa SI, Sasai Y: Induction of midbrain dopaminergic neurons from ES cells by stromal cell-derived inducing activity. *Neuron* 28: 31–40, 2000
- 62) Takagi Y, Takahashi J, Saiki H, Morizane A, Hayashi T, Kishi Y, Fukuda H, Okamoto Y, Koyanagi M, Ideguchi M, Hayashi H, Imazato T, Kawasaki H, Suemori H, Omachi S, Iida H, Itoh N, Nakatsuji N, Sasai Y, Hashimoto N: Dopaminergic neurons generated from monkey embryonic stem cells function in a Parkinson primate model. *J Clin Invest* 115: 102–109, 2005
- 63) Redmond DE, Bjugstad KB, Teng YD, Ourednik V, Ourednik J, Wakeman DR, Parsons XH, Gonzalez R, Blanchard BC, Kim SU, Gu Z, Lipton SA, Markakis EA, Roth RH, Elsworth JD, Sladek JR, Sidman RL, Snyder EY: Behavioral improvement in a primate Parkinson's model is associated with multiple homeostatic effects of human neural stem cells. *Proc Natl Acad Sci USA* 104: 12175–12180, 2007
- 64) Cho MS, Lee YE, Kim JY, Chung S, Cho YH, Kim DS, Kang SM, Lee H, Kim MH, Kim JH, Leem JW, Oh SK, Choi YM, Hwang DY, Chang JW, Kim DW: Highly efficient and large-scale generation of functional dopamine neurons from human embryonic stem cells. *Proc Natl Acad Sci USA* 105: 3392–3397, 2008
- 65) Chung S, Moon JI, Leung A, Aldrich D, Lukianov S, Kitayama Y, Park S, Li Y, Bolshakov VY, Lamonerie T, Kim KS: ES cell-derived renewable and functional midbrain dopaminergic progenitors. *Proc Natl Acad Sci USA* 108: 9703–9708, 2011
- 66) Kriks S, Shim JW, Piao J, Ganat YM, Wakeman DR, Xie Z, Carrillo-Reid L, Auyeung G, Antonacci C, Buch A, Yang L, Beal MF, Surmeier DJ, Kordower JH, Tabar V, Studer L: Dopamine neurons derived from human ES cells efficiently engraft in animal models of Parkinson's disease. *Nature* 480: 547–551, 2011
- 67) Wernig M, Zhao JP, Pruszak J, Hedlund E, Fu D, Soldner F, Broccoli V, Constantine-Paton M, Isacson O, Jaenisch R: Neurons derived from reprogrammed fibroblasts functionally integrate into the fetal brain and improve symptoms of rats with Parkinson's disease. *Proc Natl Acad Sci USA* 105: 5856–5861, 2008
- 68) Rhee YH, Ko JY, Chang MY, Yi SH, Kim D, Kim CH, Shim JW, Jo AY, Kim BW, Lee H, Lee SH, Suh W, Park CH, Koh HC, Lee YS, Lanza R, Kim KS, Lee SH: Protein-based human iPS cells efficiently generate functional dopamine neurons and can treat a rat model of Parkinson disease. *J Clin Invest* 121: 2326–2335, 2011
- 69) Hargus G, Cooper O, Deleidi M, Levy A, Lee K, Marlow E, Yow A, Soldner F, Hockemeyer D, Hallett PJ, Osborn T, Jaenisch R, Isacson O: Differentiated Parkinson patient-derived induced pluripotent stem cells grow in the adult rodent brain and reduce motor asymmetry in Parkinsonian rats. *Proc Natl Acad Sci USA* 107: 15921–15926, 2010
- 70) Studer L: Derivation of dopaminergic neurons from pluripotent stem cells. *Prog Brain Res* 200: 243–263, 2012
- 71) Greenamyre JT, Shoulson I: Huntington disease, in Calne D (ed): *Neurodegenerative Disease*. Philadelphia, WB Saunders, 1994, pp 657–704
- 72) Harper PS: *Huntington's Disease, ed 2*. Philadelphia, WB Saunders, 1996
- 73) Bachoud-Lévi AC, Rémy P, Nguyen JP, Brugières P, Lefaucheur JP, Bourdet C, Baudic S, Gaura V, Maison P, Haddad B, Boissé MF, Grandmougin T, Jény R, Bartolomeo P, Dalla Barba G, Degos JD, Lisovsky F, Ergis AM, Pailhous E, Cesaro P, Hantraye P, Peschanski M: Motor and cognitive improvements in patients with Huntington's disease after neural transplantation. *Lancet* 356: 1975–1979, 2000
- 74) Nakao N, Itakura T: Fetal tissue transplants in animal models of Huntington's disease: the effects on damaged neuronal circuitry and behavioral deficits. *Prog Neurobiol* 61: 313–338, 2000
- 75) Dunnett SB, Carter RJ, Watts C, Torres EM, Mahal A, Mangiarini L, Bates G, Morton AJ: Striatal transplantation in a transgenic mouse model of Huntington's disease. *Exp Neurol* 154: 31–40, 1998
- 76) Freeman TB, Cicchetti F, Hauser RA, Deacon TW, Li XJ, Hersch SM, Nauert GM, Sanberg PR, Kordower JH, Saporta S, Isacson O: Transplanted fetal striatum in Huntington's disease: phenotypic development and lack of pathology. *Proc Natl Acad Sci USA* 97: 13877–13882, 2000
- 77) Keene CD, Chang RC, Leverenz JB, Kopyov O, Perlman S, Hevner RF, Born DE, Bird TD, Montine TJ: A patient with Huntington's disease and long-surviving fetal neural transplants that developed mass lesions. *Acta Neuropathol* 117: 329–338, 2009

- 78) Björklund A, Lindvall O: Cell replacement therapies for central nervous system disorders. *Nat Neurosci* 3: 537–544, 2000
- 79) A novel gene containing a trinucleotide repeat that is expanded and unstable on Huntington's disease chromosomes. The Huntington's Disease Collaborative Research Group. *Cell* 72: 971–983, 1993
- 80) Harris GJ, Codori AM, Lewis RF, Schmidt E, Bedi A, Brandt J: Reduced basal ganglia blood flow and volume in pre-symptomatic, gene-tested persons at-risk for Huntington's disease. *Brain* 122 (Pt 9): 1667–1678, 1999
- 81) Thieben MJ, Duggins AJ, Good CD, Gomes L, Mahant N, Richards F, McCusker E, Frackowiak RS: The distribution of structural neuropathology in pre-clinical Huntington's disease. *Brain* 125(Pt 8): 1815–1828, 2002
- 82) Hudson AJ. *Amyotrophic Lateral Sclerosis: Concepts in Pathogenesis and Etiology*. Toronto, University of Toronto Press, 1990
- 83) Rowland LP, Shneider NA: Amyotrophic lateral sclerosis. *N Engl J Med* 344: 1688–1700, 2001
- 84) Boillée S, Vande Velde C, Cleveland DW: ALS: a disease of motor neurons and their nonneuronal neighbors. *Neuron* 52: 39–59, 2006
- 85) Wichterle H, Lieberam I, Porter JA, Jessell TM: Directed differentiation of embryonic stem cells into motor neurons. *Cell* 110: 385–397, 2002
- 86) Harper JM, Krishnan C, Darman JS, Deshpande DM, Peck S, Shats I, Backovic S, Rothstein JD, Kerr DA: Axonal growth of embryonic stem cell-derived motoneurons in vitro and in motoneuron-injured adult rats. *Proc Natl Acad Sci USA* 101: 7123–7128, 2004
- 87) Miles GB, Yohn DC, Wichterle H, Jessell TM, Rafuse VF, Brownstone RM: Functional properties of motoneurons derived from mouse embryonic stem cells. *J Neurosci* 24: 7848–7858, 2004
- 88) Li P, Tessler A, Han SS, Fischer I, Rao MS, Selzer ME: Fate of immortalized human neuronal progenitor cells transplanted in rat spinal cord. *Arch Neurol* 62: 223–229, 2005
- 89) Glass JD, Boulis NM, Johe K, Rutkove SB, Federici T, Polak M, Kelly C, Feldman EL: Lumbar intraspinal injection of neural stem cells in patients with amyotrophic lateral sclerosis: results of a phase I trial in 12 patients. *Stem Cells* 30: 1144–1151, 2012
- 90) Dimos JT, Rodolfa KT, Niakan KK, Weisenthal LM, Mitsumoto H, Chung W, Croft GF, Saphier G, Leibel R, Goland R, Wichterle H, Henderson CE, Eggan K: Induced pluripotent stem cells generated from patients with ALS can be differentiated into motor neurons. *Science* 321: 1218–1221, 2008
- 91) Lindvall O, Kokaia Z: Stem cells for the treatment of neurological disorders. *Nature* 441: 1094–1096, 2008
- 92) Whitehouse PJ, Price DL, Clark AW, Coyle JT, DeLong MR: Alzheimer disease: evidence for selective loss of cholinergic neurons in the nucleus basalis. *Ann Neurol* 10: 122–126, 1981
- 93) Bartus RT, Dean RL, Beer B, Lippa AS: The cholinergic hypothesis of geriatric memory dysfunction. *Science* 217: 408–414, 1982
- 94) Coyle JT, Price DL, DeLong MR: Alzheimer's disease: a disorder of cortical cholinergic innervation. *Science* 219: 1184–1190, 1983
- 95) Tuszynski MH, Thal L, Pay M, Salmon DP, U HS, Bakay R, Patel P, Blesch A, Vahlsing HL, Ho G, Tong G, Potkin SG, Fallon J, Hansen L, Mufson EJ, Kordower JH, Gall C, Conner J: A phase 1 clinical trial of nerve growth factor gene therapy for Alzheimer disease. *Nat Med* 11: 551–555, 2005
- 96) Terry AV, Buccafusco JJ: The cholinergic hypothesis of age and Alzheimer's disease-related cognitive deficits: recent challenges and their implications for novel drug development. *J Pharmacol Exp Ther* 306: 821–827, 2003
- 97) Musiał A, Bajda M, Malawska B: Recent developments in cholinesterases inhibitors for Alzheimer's disease treatment. *Curr Med Chem* 14: 2654–2679, 2007
- 98) Borlongan CV: Recent preclinical evidence advancing cell therapy for Alzheimer's disease. *Exp Neurol* 237: 142–146, 2012
- 99) George PM, Steinberg GK: Novel stroke therapeutics: unraveling stroke pathophysiology and its impact on clinical treatments. *Neuron* 87: 297–309, 2015
- 100) Goings GE, Sahni V, Szele FG: Migration patterns of subventricular zone cells in adult mice change after cerebral cortex injury. *Brain Res* 996: 213–226, 2004
- 101) Arvidsson A, Collin T, Kirik D, Kokaia Z, Lindvall O: Neuronal replacement from endogenous precursors in the adult brain after stroke. *Nat Med* 8: 963–970, 2002
- 102) Parent JM, Vexler ZS, Gong C, Derugin N, Ferriero DM: Rat forebrain neurogenesis and striatal neuron replacement after focal stroke. *Ann Neurol* 52: 802–813, 2002
- 103) Kobayashi T, Ahlenius H, Thored P, Kobayashi R, Kokaia Z, Lindvall O: Intracerebral infusion of glial cell line-derived neurotrophic factor promotes striatal neurogenesis after stroke in adult rats. *Stroke* 37: 2361–2367, 2006
- 104) Dempsey RJ, Sailor KA, Bowen KK, Türeyen K, Vemuganti R: Stroke-induced progenitor cell proliferation in adult spontaneously hypertensive rat brain: effect of exogenous IGF-1 and GDNF. *J Neurochem* 87: 586–597, 2003
- 105) Hoehn BD, Palmer TD, Steinberg GK: Neurogenesis in rats after focal cerebral ischemia is enhanced by indomethacin. *Stroke* 36: 2718–2724, 2005
- 106) Schouten M, Buijink MR, Lucassen PJ, Fitzsimons CP: New neurons in aging brains: molecular control by small non-coding RNAs. *Front Neurosci* 6: 25, 2012
- 107) Wang L, Zhang Z, Wang Y, Zhang R, Chopp M: Treatment of stroke with erythropoietin enhances neurogenesis and angiogenesis and improves neurological function in rats. *Stroke* 35: 1732–1737, 2004

- 108) Lee SR, Kim HY, Rogowska J, Zhao BQ, Bhide P, Parent JM, Lo EH: Involvement of matrix metalloproteinase in neuroblast cell migration from the subventricular zone after stroke. *J Neurosci* 26: 3491–3495, 2006
- 109) Dunac A, Frelin C, Popolo-Blondeau M, Chatel M, Mahagne MH, Philip PJ: Neurological and functional recovery in human stroke are associated with peripheral blood CD34+ cell mobilization. *J Neurol* 254: 327–332, 2007
- 110) Kawada H, Takizawa S, Takanashi T, Morita Y, Fujita J, Fukuda K, Takagi S, Okano H, Ando K, Hotta T: Administration of hematopoietic cytokines in the subacute phase after cerebral infarction is effective for functional recovery facilitating proliferation of intrinsic neural stem/progenitor cells and transition of bone marrow-derived neuronal cells. *Circulation* 113: 701–710, 2006
- 111) Ohab JJ, Fleming S, Blesch A, Carmichael ST: A neurovascular niche for neurogenesis after stroke. *J Neurosci* 26: 13007–13016, 2006
- 112) Yan YP, Sailor KA, Lang BT, Park SW, Vemuganti R, Dempsey RJ: Monocyte chemoattractant protein-1 plays a critical role in neuroblast migration after focal cerebral ischemia. *J Cereb Blood Flow Metab* 27: 1213–1224, 2007
- 113) Bliss TM, Andres RH, Steinberg GK: Optimizing the success of cell transplantation therapy for stroke. *Neurobiol Dis* 37: 275–283, 2010
- 114) Andrews PW, Damjanov I, Simon D, Banting GS, Carlin C, Dracopoli NC, Føgh J: Pluripotent embryonal carcinoma clones derived from the human teratocarcinoma cell line Tera-2. Differentiation in vivo and in vitro. *Lab Invest* 50: 147–162, 1984
- 115) Pleasure SJ, Lee VM: NTera 2 cells: a human cell line which displays characteristics expected of a human committed neuronal progenitor cell. *J Neurosci Res* 35: 585–602, 1993
- 116) Saporta S, Borlongan CV, Sanberg PR: Neural transplantation of human neuroteratocarcinoma (hNT) neurons into ischemic rats. A quantitative dose-response analysis of cell survival and behavioral recovery. *Neuroscience* 91: 519–525, 1999
- 117) Stroemer P, Patel S, Hope A, Oliveira C, Pollock K, Sinden J: The neural stem cell line CTX0E03 promotes behavioral recovery and endogenous neurogenesis after experimental stroke in a dose-dependent fashion. *Neurorehabil Neural Repair* 23: 895–909, 2009
- 118) Stroemer P, Hope A, Patel S, Pollock K, Sinden J: Development of a human neural stem cell line for use in recovery from disability after stroke. *Front Biosci* 13: 2290–2292, 2008
- 119) Gage FH: Mammalian neural stem cells. *Science* 287: 1433–1438, 2000
- 120) Zhang SC, Wernig M, Duncan ID, Brüstle O, Thomson JA: In vitro differentiation of transplantable neural precursors from human embryonic stem cells. *Nat Biotechnol* 19: 1129–1133, 2001
- 121) Reubinoff BE, Itsykson P, Turetsky T, Pera MF, Reinhartz E, Itzik A, Ben-Hur T: Neural progenitors from human embryonic stem cells. *Nat Biotechnol* 19: 1134–1140, 2001
- 122) Kelly S, Bliss TM, Shah AK, Sun GH, Ma M, Foo WC, Masel J, Yenari MA, Weissman IL, Uchida N, Palmer T, Steinberg GK: Transplanted human fetal neural stem cells survive, migrate, and differentiate in ischemic rat cerebral cortex. *Proc Natl Acad Sci USA* 101: 11839–11844, 2004
- 123) Bühnenmann C, Scholz A, Bernreuther C, Malik CY, Braun H, Schachner M, Reymann KG, Dihné M: Neuronal differentiation of transplanted embryonic stem cell-derived precursors in stroke lesions of adult rats. *Brain* 129: 3238–3248, 2006
- 124) Daadi MM, Lee SH, Arac A, Grueter BA, Bhatnagar R, Maag AL, Schaar B, Malenka RC, Palmer TD, Steinberg GK: Functional engraftment of the medial ganglionic eminence cells in experimental stroke model. *Cell Transplant* 18: 815–826, 2009
- 125) Daadi MM, Li Z, Arac A, Grueter BA, Sofilos M, Malenka RC, Wu JC, Steinberg GK: Molecular and magnetic resonance imaging of human embryonic stem cell-derived neural stem cell grafts in ischemic rat brain. *Mol Ther* 17: 1282–1291, 2009
- 126) Shen LH, Li Y, Chen J, Zacharek A, Gao Q, Kapke A, Lu M, Raginski K, Vanguri P, Smith A, Chopp M: Therapeutic benefit of bone marrow stromal cells administered 1 month after stroke. *J Cereb Blood Flow Metab* 27: 6–13, 2007
- 127) Meissner A, Wernig M, Jaenisch R: Direct reprogramming of genetically unmodified fibroblasts into pluripotent stem cells. *Nat Biotechnol* 25: 1177–1181, 2007
- 128) Wernig M, Meissner A, Foreman R, Brambrink T, Ku M, Hochedlinger K, Bernstein BE, Jaenisch R: In vitro reprogramming of fibroblasts into a pluripotent ES-cell-like state. *Nature* 448: 318–324, 2007
- 129) Mohamad O, Drury-Stewart D, Song M, Faulkner B, Chen D, Yu SP, Wei L: Vector-free and transgene-free human iPS cells differentiate into functional neurons and enhance functional recovery after ischemic stroke in mice. *PLoS ONE* 8: e64160, 2013
- 130) Pang ZP, Yang N, Vierbuchen T, Ostermeier A, Fuentes DR, Yang TQ, Citri A, Sebastiano V, Marro S, Südhof TC, Wernig M: Induction of human neuronal cells by defined transcription factors. *Nature* 476: 220–223, 2011
- 131) Jandial R, Snyder EY: A safer stem cell: on guard against cancer. *Nat Med* 15: 999–1001, 2009
- 132) Kondziolka D, Wechsler L, Goldstein S, Meltzer C, Thulborn KR, Gebel J, Jannetta P, DeCesare S, Elder EM, McGrogan M, Reitman MA, Bynum L: Transplantation of cultured human neuronal cells for patients with stroke. *Neurology* 55: 565–569, 2000
- 133) Kondziolka D, Steinberg GK, Wechsler L, Meltzer CC, Elder E, Gebel J, Decesare S, Jovin T, Zafonte

- R, Lebowitz J, Flickinger JC, Tong D, Marks MP, Jamieson C, Luu D, Bell-Stephens T, Teraoka J: Neurotransplantation for patients with subcortical motor stroke: a phase 2 randomized trial. *J Neurosurg* 103: 38–45, 2005
- 134) Lee JS, Hong JM, Moon GJ, Lee PH, Ahn YH, Bang OY; STARTING collaborators: A long-term follow-up study of intravenous autologous mesenchymal stem cell transplantation in patients with ischemic stroke. *Stem Cells* 28: 1099–1106, 2010
- 135) Bhasin A, Srivastava MV, Kumaran SS, Mohanty S, Bhatia R, Bose S, Gaikwad S, Garg A, Airan B: Autologous mesenchymal stem cells in chronic stroke. *Cerebrovasc Dis Extra* 1: 93–104, 2011
- 136) Bhasin A, Srivastava MV, Mohanty S, Bhatia R, Kumaran SS, Bose S: Stem cell therapy: a clinical trial of stroke. *Clin Neurol Neurosurg* 115: 1003–1008, 2013
- 137) Friedrich MA, Martins MP, Araújo MD, Klamt C, Vedolin L, Garicochea B, Raupp EF, Sartori El Ammar J, Machado DC, Costa JC, Nogueira RG, Rosado-de-Castro PH, Mendez-Otero R, Freitas GR: Intra-arterial infusion of autologous bone marrow mononuclear cells in patients with moderate to severe middle cerebral artery acute ischemic stroke. *Cell Transplant* 21(Suppl 1): S13–S21, 2012
- 138) Savitz SI, Misra V, Kasam M, Juneja H, Cox CS Jr, Alderman S, Aisiku I, Kar S, Gee A, Grotta JC: Intravenous autologous bone marrow mononuclear cells for ischemic stroke. *Ann Neurol* 70: 59–69, 2011
- 139) Moniche F, Gonzalez A, Gonzalez-Marcos JR, Carmona M, Piñero P, Espigado I, Garcia-Solis D, Cayuela A, Montaner J, Boada C, Rosell A, Jimenez MD, Mayol A, Gil-Peralta A: Intra-arterial bone marrow mononuclear cells in ischemic stroke: a pilot clinical trial. *Stroke* 43: 2242–2244, 2012
- 140) Steinberg GK, Kondziolka D, Schwartz NE, Wechsler L, Lunsford LD, Coburn ML, Billigen JB, Keren-Gill H, McGrogan M, Case C, Mori K, Yankee EW: A novel phase 1/2a study of intraparenchymal transplantation of human modified bone marrow derived cells in patients with stable ischemic stroke. *Stroke* 45, A149, 2014

Address reprint requests to: Yasushi Takagi, MD, PhD, Department of Neurosurgery, Kyoto University, Graduate School of Medicine, 54 Kawahara-cho, Shogoin, Sakyo, Kyoto 606-8507, Japan.
e-mail: ytakagi@kuhp.kyoto-u.ac.jp