

## HOLOMORPHIC MAPPING INTO ALGEBRAIC VARIETIES OF GENERAL TYPE

PEICHU HU

### §1. Introduction

We will study holomorphic mappings

$$f: M \longrightarrow N$$

from a connected complex manifold  $M$  of dimension  $m$  to a projective algebraic manifold  $N$  of dimension  $n$ . Assume first that  $N$  is of general type, i.e.

$$\varliminf_{k \rightarrow \infty} \frac{\dim H^0(N, K_N^k)}{k^n} > 0,$$

where  $K_N \rightarrow N$  is the canonical bundle of  $N$ . If  $K_N$  is positive, then  $N$  is of general type.

In 1971, Kodaira [6] obtained that

**THEOREM A.** *Any holomorphic mapping  $f: C^m \rightarrow N$  has everywhere rank less than  $n$ .*

P. Griffiths & J. King [2], [3] furthermore proved that

**THEOREM B.** *If  $M$  is a smooth affine algebraic variety, then any holomorphic mapping  $f: M \rightarrow N$  whose image contains an open set is necessarily rational.*

In 1977, W. Stoll [6] extended Theorems A, B to parabolic manifolds  $M$ . To state it, we let  $M$  possess a parabolic exhaustion  $\tau$  and denote

$$(1) \quad \nu = dd^c \tau, \quad \sigma = d^c \log \tau \wedge (dd^c \log \tau)^{m-1}.$$

For a form  $\varphi$  of bidegree  $(1, 1)$  on  $M$ , write

$$(2) \quad A(t, \varphi) = t^{2-2m} \int_{M[t]} \varphi \wedge \nu^{m-1}, \quad T(r, s; \varphi) = \int_s^r \frac{A(t, \varphi)}{t} dt$$

---

Received July 10, 1989.

if the integrals exist, where  $M[t] = \{x \in M: \tau(x) \leq t^2\}$ . Suppose throughout that  $L$  is a positive holomorphic line bundle over  $N$  with a hermitian metric  $\rho$  along the fibers of  $L$  such that the Chern form  $c(L, \rho) > 0$ . The characteristic function of  $f$  for  $L$  is defined by

$$(3) \quad T(r, s) = T(r, s; f^*c(L, \rho)).$$

THEOREM C. *If  $M$  is a parabolic manifold and if  $F$  is an effective Jacobian section such that*

(i)  *$F$  is dominated by  $\tau$  with  $Y$  as dominator, there exist positive constants  $c_1, c_2, c_3$  such that for  $\varepsilon > 0$*

$$(4) \quad T(r, s) \leq c_1 \log Y(r) + c_2 \text{Ric}_\tau(r, s) + c_3 \varepsilon \log r$$

*with the exception of a set of values  $(r)$  of finite measure.*

The condition (i) implies  $m \geq n = \text{rank } f$  ([8], Lemma 18.1). We remove this restriction (see [4]). To state the generalization of the Theorem C which we shall prove, we take a positive form  $\psi$  of class  $C^\infty$  and bidegree  $(1, 1)$  on  $N$  and set

$$(5) \quad \psi_f = \begin{cases} f^*(\psi^m) & \text{if } m \leq n \\ f^*(\psi^n) \wedge \chi & \text{if } m > n \end{cases}$$

where  $\chi$  be a positive  $(m - n, m - n)$ -form of class  $C^\infty$  on  $M$ . Then the form

$$(6) \quad \chi_f = f^*(\text{Ric } \psi^n) - \frac{n}{b} \text{Ric } \psi_f \quad \text{where } b = \min(m, n),$$

is well-defined. Take a holomorphic form  $B$  of bidegree  $(m - 1, 0)$  on  $M$ . Define

$$\begin{aligned} \check{\psi}_f &= \check{\psi}_f(B) = m i_{m-1} f^*(\psi) \wedge B \wedge \bar{B}, \\ e_f &= e_f(\psi) = f^*(\text{Ric } \psi^n) - n \text{Ric } \check{\psi}_f, \end{aligned}$$

where  $i_{m-1}$  is defined in Section 3. Then  $\chi_f(h\psi) = \chi_f(\psi)$ ,  $e_f(h\psi) = e_f(\psi)$  for positive functions  $h$  of class  $C^2$  on  $N$ . Define  $\eta$  by  $\check{\psi}_f = \eta f^*(\psi) \wedge \nu^{m-1}$  and denote

$$(7) \quad B(r, \eta) = \frac{1}{2} \int_{\partial M[r]} \log \eta \sigma,$$

$$(8) \quad E_f(r, s) = T(r, s; e_f) + nB(t, \eta)|_s^r,$$

where  $B(t)|_r^s$  means  $B(r) - B(s)$ . For  $\psi = c(L, \rho)$ , we obtain that

**THEOREM 1.** *If there exists an effective Jacobian section of  $f$  and if  $\text{rank } f = b = \min(m, n)$ , then exist positive constants  $c_1$  and  $c_2$  such that for  $\varepsilon > 0$*

$$(9) \quad c_1 T(r, s) \leq n \text{ Ric}_f(r, s) + E_f(r, s) + c_2 \varepsilon \log r$$

with the exception of a set of values  $(r)$  of finite measure.

**COROLLARY 2.** *If  $M$  is smooth affine algebraic variety, any non-degenerate holomorphic mapping  $f: M \rightarrow N$  with*

$$(ii) \quad \overline{\lim}_{r \rightarrow \infty} \frac{E_f(r, s)}{\log r} < \infty$$

is necessarily rational.

To draw geometrical consequences, here assume that  $M$  and  $N$  are hermitian manifolds. Relative to the local coordinates  $z^i$  let

$$(10) \quad ds_M^2 = \sum_{i,j} h_{ij} dz^i d\bar{z}^j \quad 1 \leq i, j \leq m$$

be a positive definite hermitian metric on  $M$  with the associated 2-form

$$(11) \quad \varphi = \frac{\sqrt{-1}}{2\pi} \sum_{i,j} h_{ij} dz^i \wedge d\bar{z}^j.$$

Similarly, let

$$(12) \quad ds_N^2 = \sum_{k,l} \tilde{h}_{kl} dw^k d\bar{w}^l \quad 1 \leq k, l \leq n$$

be a positive definite hermitian metric on  $N$ , with the local coordinates  $w^k$ , and

$$(13) \quad \psi = \frac{\sqrt{-1}}{2\pi} \sum_{k,l} \tilde{h}_{kl} dw^k \wedge d\bar{w}^l$$

be the associated 2-form. Define the function  $u$  on  $M$  by

$$(14) \quad \psi_f = u\varphi^m.$$

Then we have

$$(15) \quad \partial\bar{\partial} \log u = \text{Ric}_M - \frac{b}{n} f^*(\text{Ric}_N) + \frac{2\pi b \sqrt{-1}}{n} \chi_f.$$

When  $m \leq n$ ,

$$(16) \quad u = \frac{\det(\hat{h}_{ij})}{\det(h_{ij})}$$

is geometrically the ratio of the volume elements, where

$$\hat{h}_{ij} = \sum_{k,l} \tilde{h}_{kl} \frac{\partial w^k}{\partial z^i} \frac{\partial \bar{w}^l}{\partial \bar{z}^j}$$

under the mapping  $f$ . If  $m = n$ , (15) implies the Chern formula [1]

$$(17) \quad \frac{1}{2} \Delta \log u = R - \text{Tr}(f^*(\text{Ric}_N)),$$

where  $\Delta$  is the Laplacian in  $M$  and  $R$  denotes the scalar curvature of  $M$ .

Let  $D_f$  be the zero divisor of  $\psi_f$ , which independent of the choices of  $\psi$  and  $\chi$ . Then  $\chi_f$  determines an element  $[\chi_f] \in H_{DR}^2(M - D_f, \mathbf{R})$ , the de Rham cohomology group of closed  $C^\infty$  differential forms modulo exact ones. We extend the Chern Theorems [1] on holomorphic mappings of hermitian manifolds of the same dimension to non-equidimensional cases. This includes a non-equidimensional version of the Schwarz lemma, which says that if  $M$  is the unit  $m$ -ball and  $N$  is almost einsteinian with  $\sqrt{-1} \text{Tr}(\chi_f) \geq 0$ , the mapping  $f$  does not increase volume.

The author learned about value distribution theory from Mo Ye and Yum-Tong Siu, whom he wishes to thank for sharing their insights with him. Also he would like to thank the referee for his suggestions to correct errors in this paper.

## §2. The Ricci form and proof of the formula (15)

As usual, we let

$$d = \partial + \bar{\partial} \quad \text{and} \quad d^c = \frac{\sqrt{-1}}{4\pi} (\bar{\partial} - \partial).$$

Then

$$dd^c = \frac{\sqrt{-1}}{2\pi} \partial\bar{\partial}.$$

The Chern form of the line bundle  $L$  for the hermitian metric  $\rho$  is defined by

$$c(L, \rho) = -dd^c \log |s|_\rho^2 \quad \text{on } U$$

for all open subsets  $U$  in  $N$  and all  $s \in H^0(U, L)$ . Let  $\Psi$  be a volume form

on  $N$ . This is the same as a metric on the canonical line bundle  $K_N$ , which is denoted by  $\rho_\psi$ . In terms of complex coordinates  $w^1, \dots, w^n$ , such a form is one which can be written

$$\Psi(w) = \rho(w)\Phi(w) \quad \text{where } \Phi(w) = \prod_{j=1}^n \frac{\sqrt{-1}}{2\pi} dw^j \wedge d\bar{w}^j$$

and  $\rho$  is real  $>0$ . In practice one often has

$$\rho(w) = \lambda(w)|g(w)|^{2q},$$

where  $g$  is holomorphic not identically zero,  $q$  is some fixed rational number  $>0$  and  $\lambda$  is  $C^\infty$  and  $>0$ . We define the Ricci form of  $\Psi$  to be the Chern form of this metric  $\rho_\Psi$  on  $K_N$ , so

$$\text{Ric } \Psi = c(K_N, \rho_\Psi) = dd^c \log \rho = dd^c \log \lambda,$$

which is independent of the choice of complex coordinates, and defines a real (1, 1)-form.

Now we prove the formula (15). It is well known that the Ricci form of  $M$  for the metric  $ds_M^2$  is of

$$(18) \quad \text{Ric}_M = -\partial\bar{\partial} \log \det(h_{i,j}).$$

Then we have

$$(19) \quad \text{Ric } \varphi^m = dd^c \log \det(h_{i,j}) = \frac{1}{2\pi\sqrt{-1}} \text{Ric}_M.$$

It follows that

$$\begin{aligned} \chi_f &= f^*(\text{Ric } \psi^n) - \frac{n}{b} \text{Ric } \psi_f \\ &= f^*\left(\frac{1}{2\pi\sqrt{-1}} \text{Ric}_N\right) - \frac{n}{b} (dd^c \log u + \text{Ric } \varphi^m), \end{aligned}$$

which implies (15) by (19).

For convenience, we let  $\chi = 1$  if  $m \leq n$ , so that

$$\psi_f = f^*(\psi^b) \wedge \chi.$$

Hence when  $m \leq n$ ,  $u$  is independent of the choice of  $\chi$  and of the expression (16). Thus

$$u = \frac{\det(\tilde{h}_{ki})}{\det(h_{i,j})} \left| \det\left(\frac{\partial w^k}{\partial z^t}\right) \right|^2$$

if  $m = n$ . When  $m > n$ ,  $u = u_\chi$  depends on the choice of  $\chi$  with

$$u_{h\chi} = hu_\chi,$$

where  $h$  is a function on  $M$ . Locally we may choose an orthonormal co-frame  $\theta_1, \dots, \theta_m$  for  $M$  such that

$$ds_M^2 = \sum_{j=1}^m \theta_j \bar{\theta}_j.$$

It is well-known that  $ds_M^2$  induces an intrinsic connection on  $M$  and we let

$$\Omega_{ij} = \frac{1}{2} \sum_{k,l} R_{ijkl} \theta_k \wedge \bar{\theta}_l$$

be the curvature. Then

$$\text{Ric}_M = \sum_{i=1}^m \Omega_{ii} = \frac{1}{2} \sum_{k,l} R_{kl} \theta_k \wedge \bar{\theta}_l,$$

where

$$R_{kl} = \sum_{i=1}^m R_{iikl}.$$

From them we form the scalar curvature

$$R = \sum_{k=1}^m R_{kk}.$$

Similarly, let  $\omega_1, \dots, \omega_n$  be an orthonormal co-frame for  $N$  such that

$$ds_N^2 = \sum_{k=1}^n \omega_k \bar{\omega}_k$$

and let  $S_{ijkl}$ ,  $S_{ij}$  and  $S$  be the curvature tensor, the Ricci tensor and scalar curvature of  $N$  respectively. We put

$$\begin{aligned} du &= \sum_i (u_i \theta_i + \bar{u}_i \bar{\theta}_i), \\ \delta \bar{\partial} u &= -d\bar{\partial} u = \sum_{i,j} u_{ij} \theta_i \wedge \bar{\theta}_j. \end{aligned}$$

Then the Laplacian of  $u$  is defined by

$$\Delta u = 4 \sum_i u_{ii}.$$

If  $u > 0$ , we find

$$(20) \quad \Delta \log u = \frac{1}{u} \Delta u - \frac{4}{u^2} \sum_i u_i \bar{u}_i.$$

Under the mapping  $f$  let us set

$$(21) \quad \omega_i = \sum_{j=1}^m a_{ij} \theta_j \quad 1 \leq i \leq n.$$

If  $u > 0$ , it follows from (15) that

$$(22) \quad \frac{1}{2} \Delta \log u = R - \frac{b}{n} \sum_{k,l,i} S_{k\bar{l}} a_{ki} \bar{a}_{li} + \frac{2b}{n} \lambda_f,$$

where

$$(23) \quad \lambda_f = 2\pi\sqrt{-1} \operatorname{Tr}(\chi_f).$$

When  $m = n$ , (22) implies (17).

To draw geometrical conclusions we start with some definitions:  $f$  is said to be degenerate at  $p \in M$ , if  $u$  vanishes at  $p$ , totally degenerate if  $u$  vanishes identically, volume decreasing or volume increasing according as  $u \leq 1$  or  $u \geq 1$  for a  $\lambda$ . Proceeding in similar manner as Chern [1], we have

PROPOSITION 3. *Let  $f: M \rightarrow N$  be a holomorphic mapping, where  $M, N$  are hermitian manifolds of dimension  $m$  and  $n$  respectively, with  $M$  compact and  $N$  einsteinian. Let  $R$  and  $S$  be their scalar curvature respectively. Then we have*

- (1) *If  $R > 0, S \leq 0, \lambda_f \geq 0$ , then  $f$  is totally degenerate.*
- (2) *If  $R < 0, S \geq 0, \lambda_f \leq 0$ , then there is a point of  $M$  at which  $f$  is degenerate.*

To obtain an upper bound for the scalar function  $u$ , Chern impose some conditions on the domain manifold  $M$  and the image manifold  $N$ . The first property is:

( $DO_K$ ).  $M$  is exhausted by a sequence of open submanifolds

$$M_1 \subset M_2 \subset M_3 \subset \dots \subset M$$

whose closures  $\bar{M}_\alpha$  are compact, such that: (1) to each  $\alpha = 1, 2, \dots$  there is a smooth function  $\nu_\alpha \geq 0$  defined in  $M_\alpha$ , which satisfies the inequality

$$(24) \quad \frac{1}{2} \Delta \nu_\alpha \leq R + K \exp(\nu_\alpha/m),$$

where  $K$  is a given positive constant; (2)  $\nu_\alpha(p_\beta) \rightarrow \infty$ , if  $p_\beta$  is a divergent sequence of points in  $M_\alpha$ .

For example, the unit ball  $M = D_1$  defined by

$$r^2 = z_1\bar{z}_1 + \dots + z_m\bar{z}_m < 1$$

in the  $m$ -dimensional number space  $C^m$  with coordinates  $(z_1, \dots, z_m)$  has the property  $(DO_K)$ , with

$$(25) \quad \nu_\rho = \log \left( \frac{1 - r^2}{\rho^2 - r^2} \right)^{2m}$$

in the exhaustion submanifolds  $D_\rho$  of  $D_1$ , where  $D_\rho$  be defined by  $r < \rho$  ( $< 1$ ), and  $K = 2m(m + 1)$ . The unit ball is einsteinian with its scalar curvature  $R = -2m(m + 1)$  under the kählerian metric

$$(26) \quad ds_M^2 = \frac{1}{1 - r^2} \sum_k dz_k d\bar{z}_k + \frac{4r^2}{(1 - r^2)^2} \partial r \bar{\partial} r.$$

$(IM_K)$ .  $N$  is said to have the property  $(IM_K)$  (or almost einsteinian), if

$$(27) \quad \sum_{i,k} S_{ik} \zeta_i \bar{\zeta}_k \leq -\frac{K}{n} \sum_i \zeta_i \bar{\zeta}_i, \quad \text{for all } \zeta_i.$$

For the rest of this section we let  $m \leq n$ . Define

$$A_{jk} = \sum_{i=1}^n a_{ij} \bar{a}_{ik}.$$

Then we have

$$(28) \quad u = \det(A_{jk}).$$

By Hadamard's well-known determinant inequality we have

$$\frac{1}{m} \sum_{j,k} |A_{jk}|^2 \geq |\det(A_{jk})|^{2/m} = u^{2/m}.$$

Hence Cauchy-Hölder's inequality implies

$$(29) \quad (m^{1/2}/n)u^{1/m} \leq \frac{1}{n} (\sum_{j,k} |A_{jk}|^2)^{1/2} \leq \frac{1}{n} \sum_{i,j} |a_{ij}|^2.$$

It follows from (22) that if  $N$  have the property  $(IM_K)$  and  $u > 0$  we have

$$(30) \quad \frac{1}{2} \Delta \log u \geq R + (m^{3/2}/n^2)Ku^{1/m} + \frac{2m}{n} \lambda_j.$$

Now proceeding in similar manner as Chern [1], we have

**PROPOSITION 4.** *Let  $f: M \rightarrow N$  be a holomorphic mapping, where  $M$  and  $N$  are hermitian manifolds of dimension  $m$  and  $n$  having the properties*



$(DO_K)$  and  $(IM_{K_0})$  respectively, with  $K_0 = (n^2/m^{3/2})K$  and  $m \leq n$ . If  $\lambda_f \geq 0$ , then  $u \leq \exp(\nu_a)$ .

PROPOSITION 5. Let  $f: D_1 \rightarrow N$  be a holomorphic mapping, where  $D_1$  is the unit  $m$ -ball with the standard kähler metric and where  $N$  is an  $n$ -dimensional hermitian einsteinian manifold with scalar curvature  $\leq -2n^2(m+1)/m^{1/2}$  and  $n \geq m$ . If  $\lambda_f \geq 0$ , then  $f$  is volume-decreasing.

§ 3. Notes on parabolic manifolds

From now on, we will study value distribution on the holomorphic mapping  $f: M \rightarrow N$ . Let  $L_f \rightarrow M$  be the pull-back of  $L \rightarrow N$  and  $s_f$  the pull-back of  $s \in H^0(N, L)$ . Then  $K_M \otimes (K_N^*)_{s_f}$  is called the Jacobian bundle, its holomorphic sections over  $M$  are called Jacobian sections. A Jacobian section  $F$  is called effective if the set  $F^{-1}(0)$  of zeroes is thin, its zero divisor  $D_F$  is called the ramification divisor of  $f$  for  $F$ . Let  $A_k^p(U)$  be the vector space of forms of class  $C^k$  and degree  $p$  on  $U \subset N$ . Define

$$i_p = \left(\frac{\sqrt{-1}}{2\pi}\right)^p (-1)^{p(p-1)/2} p!.$$

Then a Jacobian section  $F$  operates on forms of degree  $2n$  as follows: Take  $\Psi \in A_k^{2n}(U)$  with  $\tilde{U} = f^{-1}(U) \neq \emptyset$ . Relative to the local coordinates  $z^i$  and  $w^k$  of  $M$  and  $N$  respectively, write

$$\begin{aligned} F &= g dz^1 \wedge \dots \wedge dz^m \otimes \left(\frac{\partial}{\partial w^1} \wedge \dots \wedge \frac{\partial}{\partial w^n}\right)_f, \quad g \in \text{Hol}(\tilde{U}), \\ \Psi &= i_n h dw^1 \wedge \dots \wedge dw^n \wedge d\bar{w}^1 \wedge \dots \wedge d\bar{w}^n. \end{aligned}$$

Then

$$F[\Psi] = i_m (h \circ f) |g|^2 dz^1 \wedge \dots \wedge dz^m \wedge d\bar{z}^1 \wedge \dots \wedge d\bar{z}^m.$$

If  $M$  is Stein and if  $f$  has strict rank  $\min(m, n)$ , effective Jacobian sections exist (see [8]).

Assume that  $\tau$  is a parabolic exhaustion of  $M$ , i.e., a proper map  $\tau: M \rightarrow \mathbf{R}^+$  of class  $C^\infty$  which satisfies

$$\begin{cases} dd^c \log \tau \geq 0, \\ (dd^c \tau)^m \neq 0 \text{ but } (dd^c \log \tau)^m \equiv 0, \\ M[0] \text{ has measure zero.} \end{cases}$$

For any regular value  $r$  of  $\tau$ , then

$$c = \int_{\partial M[r]} \sigma$$

is a constant. Take a positive form  $\Omega$  of degree  $2m$  and class  $C^2$  on  $M$ . Define  $\nu$  by  $\nu^m = \nu\Omega$ . The Ricci function of  $\tau$  is defined by

$$(31) \quad \text{Ric}_\tau(r, s) = T(r, s; \text{Ric } \Omega) + B(t, \nu)|_s^r,$$

which does not depend on the choice of  $\Omega$ . Let  $D$  be a divisor on  $M$  and set  $D[r] = D \cap M[r]$ . We define

$$n(t, D) = t^{2-2m} \int_{D[t]} \nu^{m-1},$$

$$N(r, s; D) = \int_s^r n(t, D) \frac{dt}{t}.$$

If we define  $\nu$  by  $\nu^m = \nu F[\Psi]$  for an effective Jacobian section  $F$  and a positive volume form  $\Psi$  of class  $C^\infty$  and degree  $2n$  on  $N$ , then

$$(32) \quad \text{Ric}_\tau(r, s) = T(r, s; f^*(\text{Ric } \Psi)) + B(t, \nu)|_s^r + N(r, s; D_F)$$

(For a detailed proof see [8] Theorem 15.5).

Take an effective Jacobian section  $F$  and a positive form  $\psi$  of class  $C^\infty$  and bidegree  $(1, 1)$  on  $N$ . Define  $u_0$  and  $u_1$  by

$$(33) \quad \nu^m = u_0 \check{\psi}_f, \quad \nu^m = u_1 F[\psi^n].$$

By the definitions of  $\eta$  and  $\check{\psi}_f$ , we have

$$\nu^m = u_0 \eta f^*(\psi) \wedge \nu^{m-1}.$$

Let  $D_f$  be the zero divisor of  $\check{\psi}_f$ . Then

$$(34) \quad S_f(r, s) = N(r, s; D_F) - nN(r, s; D_f) + B\left(t, \frac{u_1}{u_0}\right)|_s^r$$

is defined such that

$$(35) \quad E_f(r, s) + S_f(r, s) = (1 - n) \text{Ric}_\tau(r, s) + nB(t, \eta)|_s^r.$$

In fact, the form  $\check{\psi}_f$  determines a section  $s_f$  of  $K_M$  such that  $\check{\psi}_f = |s_f|_\rho^2 \Omega$  for a volume form  $\Omega$  and a hermitian metric  $\rho$  along the fibers of  $K_M$ . Then by Green Residue Theorem [9]

$$(36) \quad T(r, s; dd^c \log |s_f|_\rho^2) + N(r, s; D_f) = B(t, |s_f|_\rho^2)|_s^r$$

for all regular values  $s$  and  $r$  of  $\tau$  with  $0 < s < r$ . Since

$$\text{Ric } \check{\psi}_f = dd^c \log |s_f|_\rho^2 + \text{Ric } \Omega ,$$

we have

$$\begin{aligned} (37) \quad \text{Ric}_r(r, s) &= T(r, s; \text{Ric } \Omega) + B(t, u_0 \cdot |s_f|_\rho^2)|_s^r && \text{(by (31)) ,} \\ &= T(r, s; \text{Ric } \check{\psi}_f) + N(r, s; D_f) + B(t, u_0)|_s^r && \text{(by (36)) .} \end{aligned}$$

It follows from (32) that

$$(38) \quad \text{Ric}_r(r, s) = T(r, s; f^*(\text{Ric } \psi^n)) + B(t, u_1)|_s^r + N(r, s; D_F) .$$

Multiply (37) by  $n$  and minus (38) to obtain (35).

Let  $D$  be a divisor given by the zeroes of a holomorphic section  $\alpha \in H^0(N, L)$ . Since  $\alpha$  and  $\lambda\alpha$  ( $\lambda \neq 0$ ) define the same divisor and  $N$  is compact, we shall assume that  $|\alpha(x)|_\rho \leq 1$  for  $x \in N$ , i.e., the metric  $\rho$  is distinguished. Assume that  $\alpha_f \neq 0$ . The proximity form is defined by

$$m(r, D) = B(r, |\alpha_f|^{-2}) \geq 0 .$$

Then we have F. M. T. for any effective divisor (see [3], [8])

$$(39) \quad N(r, s; D_f^c) + m(t, D)|_s^r = T(r, s) ,$$

where  $D_f^c$  be the divisor of  $\alpha_f \in H^0(M, L_f)$ .

The following Lemma is well-known (see Nevanlinna [7]):

LEMMA 6. *Let  $h(r) \geq 0$ ,  $g(r) \geq 0$  and  $\alpha(r) > 0$  be increasing continuous functions of  $r$  where  $g'(r)$  is continuous and  $h'(r)$  is piecewise continuous. Suppose moreover that  $\int^\infty (dr/\alpha(r)) < \infty$ . Then*

$$h'(r) \leq g'(r)\alpha(h(r))$$

except for a union of intervals  $I \subset \mathbf{R}^+$  such that  $\int_I dg < \infty$ .

We use the notation

$$\|_\varepsilon a(r) \leq b(r)$$

to mean that the stated inequality holds except on an open set  $I \subset \mathbf{R}^+$  such that  $\int_I r^\varepsilon dr < \infty$  for  $\varepsilon > 0$ .

LEMMA 7. *Let  $\varphi \geq 0$  be a form of bidegree (1, 1) on  $M$  such that  $T(r, s; \varphi)$  exists. Let  $u \geq 0$  be a function on  $M$  such that*

$$u\omega^m \leq \varphi \wedge \nu^{m-1} .$$

Then

$$\|_{\varepsilon} B(r, u) \leq \frac{c}{2} \{(1 + 2\varepsilon) \log T(r, s; \varphi) + 4\varepsilon \log r\}.$$

*Proof.* Define

$$\hat{B}(r, u) = \frac{1}{c} \int_{\partial M[r]} u \sigma.$$

Since

$$\begin{aligned} 0 \leq r^{2m-2} A(r, u\nu) &= m \int_{M[r]} u \tau^{m-1} d\tau \wedge \sigma = 2m \int_0^r \left\{ \int_{\partial M[t]} u \sigma \right\} t^{2m-1} dt \\ &= 2mc \int_0^r \hat{B}(t, u) t^{2m-1} dt \leq r^{2m-2} A(r, \varphi), \end{aligned}$$

$\hat{B}(t, u)$  exists for almost all  $t > 0$ . Now

$$\frac{2}{c} B(r, u) = \frac{1}{c} \int_{\partial M[r]} \log u \sigma \leq \log \hat{B}(r, u)$$

implies

$$\begin{aligned} H(r) &= \int_s^r t^{1-2m} dt \int_0^t r^{2m-1} \exp\left(\frac{2}{c} B(r, u)\right) dr \\ &\leq \int_s^r t^{1-2m} dt \int_0^t r^{2m-1} \hat{B}(r, u) dr \\ &= \frac{1}{2mc} \int_s^r A(t, u\nu) \frac{dt}{t} = \frac{1}{2mc} T(r, s; u\nu) \leq \frac{1}{2mc} T(r, s; \varphi). \end{aligned}$$

Taking  $h(r) = H(r)$ ,  $g(r) = r^{1+\varepsilon}/(1+\varepsilon)$ ,  $\alpha(r) = r^\lambda$  with  $\varepsilon > 0$  and  $\lambda > 1$ , we obtain from Lemma 6 that

$$\begin{aligned} \|_{\varepsilon} H'(r) &= r^{1-2m} \int_0^r r^{2m-1} \exp\left(\frac{1}{c} B(r, u)\right) dr \leq r^\varepsilon (h(r))^\lambda \\ &\leq r^\varepsilon (T(r, s; \varphi)/(2mc))^\lambda. \end{aligned}$$

Keeping the same  $\alpha$  and  $g$  and taking  $h(r) = r^{2m-1} H'(r)$ , we find

$$\begin{aligned} \|_{\varepsilon} r^{2m-1} \exp\left(\frac{2}{c} B(r, u)\right) &= \frac{d}{dr} \left( r^{2m-1} \frac{dH}{dr} \right) \leq r^\varepsilon \left( r^{2m-1} \frac{dH}{dr} \right)^\lambda \\ &\leq r^\varepsilon \{ r^{\varepsilon+2m-1} (T(r, s; \varphi)/(2mc))^\lambda \}^\lambda, \end{aligned}$$

which implies

$$\begin{aligned} (40) \quad \|_{\varepsilon} B(r, u) &\leq \frac{c}{2} \{ \lambda^2 \log T(r, s; \varphi) + (\lambda(\varepsilon + 2m - 1) + (\varepsilon + 1 - 2m)) \log r \\ &\quad - \lambda^2 \log(2mc) \}. \end{aligned}$$

Take  $0 < \delta < \min(1, \epsilon)$  such that  $\epsilon(4 + \delta) + \delta(2m - 1) < 6\epsilon$ . Let  $\lambda = 1 + \delta/2$ . Then  $\lambda^2 < 1 + 2\epsilon$  and

$$\lambda(\epsilon + 2m - 1) + \epsilon + 1 - 2m = \frac{1}{2}\{\epsilon(4 + \delta) + \delta(2m - 1)\} < 3\epsilon.$$

Hence Lemma 7 follows if  $r$  is large enough. q.e.d.

**§ 4. Holomorphic maps into algebraic varieties of general type**

*Proof of Theorem 1.* By Kobayashi-Ochiai [5] and Kodaira [6], an integer  $p \in \mathbb{N}$  exists such that  $L^p$  is ample and  $k \in \mathbb{N}$  exists such that  $H^0(N, I)$  has positive dimension with  $I = K_N^k \otimes (L^p)^*$ . Take  $\alpha \in H^0(N, I)$ . Let  $D_f$  be the divisor of  $\alpha_f \in H^0(M, I_f)$  and let  $\hat{\rho}$  be a distinguished hermitian metric along the fibers of  $I$ . Then (39) implies

$$T(r, s; f^*c(I, \hat{\rho})) = N(r, s; D_f) + m(t, D)|_s^r.$$

A form  $\Psi > 0$  of class  $C^\infty$  and degree  $2n$  exists such that  $\text{Ric } \Psi = c(K_N, \rho_\Psi)$  and  $\hat{\rho} = (\rho_\Psi)^k \otimes (\rho^*)^p$ . Hence

$$c(I, \hat{\rho}) = k \text{ Ric } \Psi - pc(L, \rho),$$

which implies

$$kT(r, s; f^*(\text{Ric } \Psi)) - m(t, D)|_s^r = pT(r, s) + N(r, s; D_f).$$

A function  $v \geq 0$  of class  $C^\infty$  exists on  $M - F^{-1}(0)$  such that  $\nu^m = vF[\Psi]$  and such that

$$\text{Ric}_t(r, s) = N(r, s; D_F) + B(t, v)|_s^r + T(r, s; f^*(\text{Ric } \Psi))$$

from (32), where  $F$  is an effective Jacobian section of  $f$ . Define  $\tilde{\zeta} = |\alpha_f|_s^{2/k} v^{-1}$ . Then

$$\begin{aligned} \text{Ric}_t(r, s) + B(t, \tilde{\zeta})|_s^r &= N(r, s; D_F) + T(r, s; f^*(\text{Ric } \Psi)) \\ &- \frac{1}{k} m(t, D)|_s^r = N(r, s; D_F) + \frac{1}{k} N(r, s; D_f) + \frac{p}{k} T(r, s). \end{aligned}$$

Therefore

$$(41) \quad nN(r, s; D_f) + \frac{p}{k} T(r, s) \leq \text{Ric}_t(r, s) - S_f(r, s) + B(t, \zeta)|_s^r,$$

where  $\zeta = u_1 u_0^{-n} \tilde{\zeta}$  and

$$\psi = c(L, \rho).$$

Define  $\hat{\psi} = |\alpha|_{\beta}^{2/k} \Psi$ . Then

$$F[\hat{\psi}] = |\alpha_f|_{\beta}^{2/k} F[\Psi] = \tilde{\zeta} \nu^m.$$

Since  $\hat{\psi}$  is continuous and  $c(L, \rho) > 0$ , a constant  $\gamma_1 > 0$  exists such that  $(\gamma_1 c(L, \rho))^n \geq \hat{\psi}$ , which implies

$$u_1 \tilde{\zeta} = u_1 \frac{F[\hat{\psi}]}{\nu^m} \leq u_1 \frac{F[(\gamma_1 c(L, \rho))^n]}{\nu^m} \leq \gamma_1^n.$$

Hence

$$\zeta^{1/n} \nu^m \leq \frac{\gamma_1}{u_0} \nu^m = \eta \gamma_1 f^*(c(L, \rho)) \wedge \nu^{m-1}.$$

It follows from Lemma 7 that

$$\begin{aligned} \left\|_{\varepsilon} B\left(t, \frac{\zeta}{\eta^n}\right) \Big|_s^r \right. &= nB(r, \zeta^{1/n}(\eta\gamma)^{-1}) + \frac{c}{2} \log \gamma_1^n - B\left(s, \frac{\zeta}{\eta^n}\right) \\ &\leq \frac{nc}{2} \{(1 + 2\varepsilon) \log T(r, s) + 5\varepsilon \log r\} \leq \frac{P}{2k} T(r, s) + 3nc\varepsilon \log r \end{aligned}$$

if  $r$  is large enough. Therefore

$$(42) \quad \left\|_{\varepsilon} nN(r, s; D_f) + \frac{P}{2k} T(r, s) \leq \text{Ric}_\varepsilon(r, s) - S_f(r, s) + nB(t, \eta) \Big|_s^r \right. \\ \left. + 3nc\varepsilon \log r . \right.$$

Now (35) and (42) yield (9).

q.e.d.

*Remark.* If  $F$  be dominated by  $\tau$  with  $Y$  as dominator, i.e.

$$n\left(\frac{F[\psi^n]}{\nu^m}\right)^{1/n} \nu^m \leq Y(r) f^*(\psi) \wedge \nu^{m-1} \quad \text{on } M[r]$$

holds for all continuous form  $\psi \geq 0$  of bidegree  $(1, 1)$  on  $M$ , which implies

$$n\left(\frac{u_0^n}{u_1}\right)^{1/n} \eta \leq Y(r).$$

Then

$$(43) \quad S_f(r, s) \geq -nN(r, s; D_f) - \frac{nc}{2} \log \frac{Y(r)}{n} + nB(t, \eta) \Big|_s^r.$$

Hence (42) and (43) yield

$$\left\|_{\varepsilon} \frac{P}{2k} T(r, s) \leq \text{Ric}_\varepsilon(r, s) + \frac{nc}{2} \log \frac{Y(r)}{n} + 3nc\varepsilon \log r , \right.$$

which is the (4) in Theorem C.

*Proof of Corollary 2.* By Stoll [8], there exist effective Jacobian sections of  $f$  and holds the following

$$0 \leq \lim_{r \rightarrow \infty} \frac{\text{Ric}_\tau(r, s)}{\log r} < \infty .$$

Then the condition (ii) and Theorem 1 imply

$$A(\infty) = \lim_{r \rightarrow \infty} A(r) = \lim_{r \rightarrow \infty} \frac{T(r, s)}{\log r} < \infty ,$$

where  $A(r) = A(r, f^*c(L, \rho))$ . Hence  $f$  is rational (see [8]). q.e.d.

*Remark.* The condition (ii) can be replaced by

$$(ii)' \quad E_f = \overline{\lim}_{r \rightarrow \infty} \frac{E_f(r, s) - nN(r, s; D_f)}{\log r} < \infty .$$

If  $M$  is smooth affine algebraic variety with  $m \geq n$ , then there exists an effective Jacobian section of  $f$  and dominated by  $\tau$  with a constant dominator  $Y = m$ . It follows from (35) and (43) that

$$\overline{\lim}_{r \rightarrow \infty} \frac{E_f(r, s) - nN(r, s; D_f)}{\log r} \leq \overline{\lim}_{r \rightarrow \infty} \frac{(1 - n) \text{Ric}_\tau(r, s)}{\log r} \leq 0 .$$

Hence (ii)' holds for this case and Theorem B follows from Corollary 2.

*Remark.* If  $M = C^m$ , then  $\text{Ric}_\tau(r, s) = 0$  where  $\tau$  is defined by  $\tau(z) = |z|^2$ . Now (9) yields

$$E_f \geq c_1 A(\infty) > 0 ,$$

because the line bundle  $L$  is positive and  $\text{rank } f = b$ . Hence we have

**COROLLARY 8.** *Let  $N$  be a connected,  $n$ -dimensional projective algebraic manifold of general type. Then any holomorphic mappings  $f: C^m \rightarrow N$  with  $E_f \leq 0$  has everywhere rank less than  $\min(m, n)$ .*

Theorem A follows from Corollary 8 and Remark above.

*Remark.* If  $\psi$  satisfies

$$\overline{\lim}_{r \rightarrow \infty} \log T(r, s; f^*(\psi)) / T(r, s) = 0$$

by the proof of Theorem 1, Theorem 1 holds for such  $\psi$ .

## REFERENCES

- [ 1 ] S. S. Chern, Shiing-Shen Chern selected papers, Springer-Verlag New York Heidelberg Berlin, 347-360.
- [ 2 ] Ph. Griffiths, Holomorphic mapping into canonical algebraic varieties, *Ann. of Math.*, (2) **93** (1971), 439-458.
- [ 3 ] Ph. Griffiths and J. King, Nevanlinna theory and holomorphic mappings between algebraic varieties, *Acta. Math.*, **130** (1973), 145-220.
- [ 4 ] P. C. Hu, On Griffiths' Conjecture in value distribution of holomorphic maps, to appear.
- [ 5 ] S. Kobayashi and T. Ochiai, Mappings into compact complex manifolds with negative Chern class, *J. Math. Soc. Japan*, **23** (1971), 137-148.
- [ 6 ] K. Kodaira, On holomorphic mappings of polydiscs into compact complex manifolds, *J. Diff. Geom.*, **6** (1971), 33-46.
- [ 7 ] R. Nevanlinna, *Eindeutige analytische Funktionen*. Die Grundle., d. Math. Wiss. XLVC Springer Verlag. Berlin-Göttingen-Heidelberg 2 ed. 1953.
- [ 8 ] W. Stoll, *Value distribution on parabolic spaces*. Lecture Notes in Mathematics 600 Springer-Verlag, Berlin-Heidelberg-New York. 1977.
- [ 9 ] —, *Value distribution theory for meromorphic maps*. Aspects of Mathematics, Vieweg. 1985.

*Department of Mathematics*  
*Shandong University*  
*Jinan, Shandong,*  
*China*