

DARWIN REVIEW

# Homogalacturonan-modifying enzymes: structure, expression, and roles in plants

Fabien Sénéchal\*, Christopher Wattier\*, Christine Rustérucci and Jérôme Pelloux†

EA3900 BIOPI Biologie des Plantes et Innovation, Université de Picardie Jules Verne, 33 Rue St Leu, F-80039 Amiens, France

\* These authors contributed equally to this work.

† To whom correspondence should be addressed. E-mail: [jerome.pelloux@u-picardie.fr](mailto:jerome.pelloux@u-picardie.fr)

Received 13 March 2014; Revised 20 May 2014; Accepted 22 May 2014

## Abstract

Understanding the changes affecting the plant cell wall is a key element in addressing its functional role in plant growth and in the response to stress. Pectins, which are the main constituents of the primary cell wall in dicot species, play a central role in the control of cellular adhesion and thereby of the rheological properties of the wall. This is likely to be a major determinant of plant growth. How the discrete changes in pectin structure are mediated is thus a key issue in our understanding of plant development and plant responses to changes in the environment. In particular, understanding the remodelling of homogalacturonan (HG), the most abundant pectic polymer, by specific enzymes is a current challenge in addressing its fundamental role. HG, a polymer that can be methylesterified or acetylated, can be modified by HGMEs (HG-modifying enzymes) which all belong to large multigenic families in all species sequenced to date. In particular, both the degrees of substitution (methylesterification and/or acetylation) and polymerization can be controlled by specific enzymes such as pectin methylesterases (PMEs), pectin acetylerases (PAEs), polygalacturonases (PGs), or pectate lyases-like (PLLs). Major advances in the biochemical and functional characterization of these enzymes have been made over the last 10 years. This review aims to provide a comprehensive, up to date summary of the recent data concerning the structure, regulation, and function of these fascinating enzymes in plant development and in response to biotic stresses.

**Key words:** Biotic stress, development, homogalacturonans, pectate lyase-like, pectin methylesterase, pectins, polygalacturonase.

## Introduction

The growth of plant organs involves cell expansion and cell division. Although the regulation of cell division is relatively well understood, very little is known about the control of cell expansion. It is driven by turgor pressure and notably depends on changes in the extensibility of the primary cell wall. In dicotyledonous species, such as the model plant *Arabidopsis thaliana*, the primary cell wall consists of a hydrogen-bonded network of cellulose microfibrils and xyloglucans (XyGs) embedded in a complex pectic and protein matrix (Carpita and Gibeaut, 1993). Cell growth requires creep between cellulose and XyG, which is facilitated by the presence of specific proteins such as expansins (Rose *et al.*, 2002; Cosgrove, 2005). In addition, in dicots and gymnosperms, secondary growth and biomass

production involve the synthesis of large amounts of lignified secondary cell wall.

Pectins, which control cell wall porosity and hydration level as well as cellular adhesion, are known to show structural variation during growth and in response to stresses (Willats *et al.*, 2001). Recent findings underline the importance of pectins in the control of cell expansion and differentiation, presumably through changes in the rheological properties of the cell wall. Pectins, which are complex polysaccharides rich in galacturonic acid (Gal-A), contain distinct domains—homogalacturonans (HGs), rhamnogalacturonans I (RGIs), rhamnogalacturonans II (RGIIs), and xylogalacturonans (XGs)—based on two molecular backbones and differing in the diversity of their side chains (Vincken *et al.*, 2003;

Mohnen, 2008; Burton *et al.*, 2010). HG, one of the main pectic constituents, is a linear homopolymer of  $\alpha$ -(1–4)-linked D-Gal-A, which can either be methylesterified at the C-6 carboxyl (typically 80%; Wolf *et al.*, 2009b) or carry acetyl groups at O2 or O3 (up to 10%; Gou *et al.*, 2012). HG is synthesized from nucleotide sugars in the Golgi apparatus, and then secreted in a fully methylesterified form into the cell wall where its structure can be modified by the activity of cell wall enzymes. These will subsequently be referred to as HGMEs (HG-modifying enzymes), all of which, in *A. thaliana*, belong to large multigenic families. In particular, HGs can be modified by pectin methylesterases (PMEs; EC 3.1.1.11), whose activity is regulated by endogenous PMEIs (pectin methylesterase inhibitors) and which control the degree of methylesterification (DM) of HGs (Pelloux *et al.*, 2007; Wolf *et al.*, 2009b). Pectin acetyltransferases (PAEs; EC 3.1.1.6) play a similar role by hydrolysing the *O*-acetylated bonds. Overall, the partially demethylesterified HGs can either form Ca<sup>2+</sup> bonds, which promote the development of the so-called ‘egg box’ structures that underlie the formation of pectin gels, or become a target for pectin-degrading enzymes such as polygalacturonases (endo-PGs, EC 3.2.1.15; and exo-PGs, EC 3.2.1.67) and pectate lyases-like (PLLs), including pectate lyases (endo-PLs, EC 4.2.2.2; and exo-PLs, EC 4.2.2.9) and pectin lyases (endo-PNLs; EC 4.2.2.10). In either case, the demethylesterification of HGs has dramatic consequences on the rheological properties of the cell wall (Peaucelle *et al.*, 2011a, b), and is predicted to affect cell growth. In addition, the hydrolysis of partially demethylesterified HGs can release oligogalacturonides (OGs) with a signalling function, for instance during plant–pathogen interactions (Lionetti

*et al.*, 2007; Osorio *et al.*, 2008) or in modulating growth by inhibiting auxin-induced processes such as stem elongation, rhizogenesis, and flower and stomata formation (Ridley *et al.*, 2001; Falasca *et al.*, 2008; Camejo *et al.*, 2011). However, to date, the presence of endogenous OGs *in planta* has remained elusive.

This review describes the most recent findings concerning the biochemical characterization of HGMEs and highlights the diversity of their roles during plant development and in responses to biotic stresses.

## Inventory and structure of HGMEs

*All HGMEs belong to large multigenic families*

An analysis of the data generated by sequencing projects shows that all HGMEs belong to rather large multigenic families in several plant species [CAZy, <http://www.cazy.org/> (Cantarel *et al.*, 2009); Cell Wall Genomics, <http://cellwall.genomics.purdue.edu/>; TAIR, <http://www.arabidopsis.org/>]. For instance, in dicotyledonous species such as *A. thaliana* and poplar (*Populus trichocarpa*), 66 and 88 open reading frames (ORFs) have been annotated, respectively, as putative full-length PMEs (Geisler-Lee *et al.*, 2006; Pelloux *et al.*, 2007). In contrast, in grass species such as *Brachypodium distachyon* and rice (*Oryza sativa* japonica group), only 41 putative PME-encoding genes have been identified for both species (Table 1). This lower number of *PME* genes in grass species is likely to be related to the differences in the structure of cell wall polysaccharides between poales and dicots. In particular, HGs are known to be much less abundant and

**Table 1.** Inventory of the *PME*, *PAE*, *PG*, and *PLL* isoforms in four sequenced plant species

Species	Pectin methylesterase (EC 3.1.1.11) CE8	Pectin acetyltransferase (EC 3.1.1.6) CE13	Endo-polygalacturonase (EC 3.2.1.15), exo-polygalacturonase (EC 3.2.1.67), exo-polygalacturonosidase (EC 3.2.1.82), rhamnogalacturonase (EC 3.2.1.171), endo-xylogalacturonan hydrolase (EC 3.2.1.–), rhamnogalacturonan $\alpha$ -L-rhamnopyranohydrolase (EC 3.2.1.40) GH28	Endo-pectate lyase (EC 4.2.2.2), exo-pectate lyase (EC 4.2.2.9), endo-pectin lyase (EC 4.2.2.10) PL1
<i>Arabidopsis thaliana</i> (dicots) <sup>a</sup>	66	12	68	26
<i>Populus trichocarpa</i> (dicots) <sup>b</sup>	88	10	89	28
<i>Brachypodium distachyon</i> (monocots) <sup>c</sup>	41	7	41	7
<i>Oryza sativa</i> Japonica group (monocots) <sup>d</sup>	41	10	45	12

<sup>a</sup> Full-length *Arabidopsis* protein data were retrieved from Cell Wall Genomics (<http://cellwall.genomics.purdue.edu/>), CAZy (<http://www.cazy.org/>), TAIR (<http://www.arabidopsis.org/>), the PFAM database (<http://pfam.sanger.ac.uk/>), Tian *et al.* (2006), Gonzalez-Carranza *et al.* (2012), Sun and Van Nocker (2010), and Cao (2012).

<sup>b</sup> *Populus* protein information were retrieved from Geisler-Lee *et al.* (2006).

<sup>c</sup> *Brachypodium* protein data were retrieved from the International Brachypodium Initiative (2010), Tyler *et al.* (2010), and Davidson *et al.* (2012).

<sup>d</sup> Full length *Oryza* protein sequence information were retrieved from Cell Wall Genomics (<http://cellwall.genomics.purdue.edu/>), CAZy (<http://www.cazy.org/>), and Davidson *et al.* (2012).

The CAZy code for each family protein was retrieved from CAZy (<http://www.cazy.org/>).

less methylesterified in grass species (Carpita and Gibeaut, 1993; Vogel, 2008; Burton *et al.*, 2010). The need for HGMEs would therefore be more limited and could explain the differences in the figures observed between type I and type II cell wall species when considering other HGMEs such as PGs and PLLs (Table 1). Surprisingly, regarding the PAE gene family, dicotyledonous and grass species have largely the same number of isoforms, with 12, 10, 7, and 11 ORFs annotated as a putative PAE for *A. thaliana*, *P. trichocarpa*, *B. distachyon*, and *O. sativa* japonica group, respectively (Table 1). The large difference between the number of PME and PAE isoforms in dicots could be related to the occurrence of specific substrates. HGs are known to be highly methylesterified (up to 80%; Wolf *et al.*, 2009b) while their degree of acetylation (DA) falls within the range of 0.25–10% (Gou *et al.*, 2012) depending on the species. In addition to HG, a number of plant cell wall polysaccharides, such as RGIs, XyGs, and glucuronoxarabinoxylans (GAXs), can be *O*-acetylated (Kiefer *et al.*, 1989; Ishii, 1997; Kabel *et al.*, 2003; Fry, 2004; Gibeaut *et al.*, 2005; Gille and Pauly, 2012). The large amount of GAX and the presence of *O*-acetyl-substituents observed in grass species (Carpita and Gibeaut, 1993; Vogel, 2008; Burton *et al.*, 2010) could thus constitute targets for PAE activity. The diversity of putative targets for PAEs would therefore explain the relatively similar number of isoforms between type I and type II cell wall species.

In order to act on HGs *in muro*, HGMEs, which are synthesized in the endoplasmic reticulum and post-translationally modified in the Golgi apparatus (glycosylation and other modifications), must be secreted into the cell wall by exocytosis (Wolf *et al.*, 2009a, b; Worden *et al.*, 2012). Several isoforms have been shown to be present in the cell wall proteome (Borderies *et al.*, 2003; Boudart *et al.*, 2005; Charmont *et al.*, 2005; Irshad *et al.*, 2008; Jamet *et al.*, 2009). A signal peptide and/or a transmembrane domain at the N-terminus enables their secretion. Analysis of HGME sequences in *A. thaliana* and *O. sativa* japonica group reveals that the majority of sequences (65% and 57%, respectively) show a signal peptide (SP). Many sequences also possess a transmembrane domain (TM; 24% and 22%, respectively) and some possess both (Supplementary Table S1 available at *JXB* online). Surprisingly, for each family, a number of putative soluble isoforms can be identified. The proportion of soluble protein is higher in grass species (17%) than in dicot species (8%). The targets and the functional role of these soluble isoforms remain to be determined. The biochemical characterization of a soluble AtPME suggested that it could act in defence mechanisms against pathogens (Dedeurwaerder *et al.*, 2009).


### Structure of HGMEs

To date, among plant HGMEs, 3D crystallographic structures have only been resolved for carrot and tomato PMEs (Johansson *et al.*, 2002; D'Avino *et al.*, 2003; Di Matteo *et al.*, 2005). The first crystallization of bacterial PME from *Erwinia chrysanthemi* was previously performed (Jenkins *et al.*, 2001), and its co-crystallization with HGs helped in determining the key amino acids involved at the catalytic site in

enzyme–substrate interactions (Fries *et al.*, 2007). This study not only highlighted the hydrolysis mechanism, but also the processive action of *Erwinia* PME (Fries *et al.*, 2007). So far, no PGs and PLs from plants have been resolved at their structural level. However, 3D crystallographic structures of PLs (Pickersgill *et al.*, 1994; Yoder and Jurnak, 1995; Lietzke *et al.*, 1996; Jenkins and Pickersgill, 2001; Jenkins *et al.*, 2004; Creze *et al.*, 2008; Seyedarabi *et al.*, 2010; Zheng *et al.*, 2012) and PGs (Pickersgill *et al.*, 1998; Van Santen *et al.*, 1999; Cho *et al.*, 2001; Jenkins and Pickersgill, 2001; Van Pouderoyen *et al.*, 2003; Bonivento *et al.*, 2007) from plant pathogens are available (PDB Protein Data Bank <http://www.rcsb.org/pdb/home/home.do>; Punta *et al.*, 2012). Identification of structural motifs was inferred from these structures, enabling models of plant PGs and PLLs to be generated (PFAM database <http://pfam.sanger.ac.uk/>). The structure of plant PAEs has not yet been resolved.

In *A. thaliana*, 66 PME isoforms are divided into two groups (Pelloux *et al.*, 2007; Dedeurwaerder *et al.*, 2009). The first, called group 1 (or type II), is composed of 21 isoforms that contain a mature active part (PME catalytic domain, Pfam01095), either preceded by different targeting motifs (SP for 14 sequences, SP/TM for 1 sequence, TM for 3 sequences) or without a motif for three soluble isoforms (Fig. 1A). In previous publications (Tian *et al.*, 2006; Pelloux *et al.*, 2007), two group 2 PMEs (At3g10720 and At4g33220) were considered to be group 1 PMEs as the protein sequences used to determine domains were truncated. The second group, so-called group 2 (or type I), is composed of 45 isoforms which have, in addition to the mature part, an N-terminal extension (PRO region) showing similarities with the PME1 domain (Pfam04043). As for group 1 PMEs, different targeting motifs can be identified for the 44 sequences, one being predicted to be soluble (Fig. 1A). Figures for other species are shown in Supplementary Table S1 at *JXB* online. Among group 2 PMEs, 43 isoforms show conserved dibasic amino acid sequences (such as RLL, RKLL, KKDL, RKLM, RRLM, RKLA, RCLK, RCLR, and RRML) upstream of the mature active part, which could constitute one or more proteolytic cleavage sites. This domain, the so-called processing motif (PM), could be targeted by subtilisin-like proteases (SBTs; 56 isoforms in *Arabidopsis*) during PME *trans*-Golgi trafficking (Wolf *et al.*, 2009a; Schaller *et al.*, 2012). A large number of group 2 PMEs and SBTs are co-expressed during development and in response to stresses. Processing of group 2 PMEs was shown to be a prerequisite for the secretion of active enzymes in the cell wall (Wolf *et al.*, 2009a) and it was suggested that the PRO region could prevent group 2 PME activity during their transport through the secretory pathway (Bosch and Hepler, 2005; Bosch *et al.*, 2005; Dorokhov *et al.*, 2006). These results are notably backed up by the fact that, in the cell wall proteome, identified PMEs often lack this domain (E. Jamet, personal communication).


When considering AtPGs, the majority of the sequences show a single glycosyl hydrolase family 28 domain (GH28; Pfam00295). This is preceded by a targeting motif at the N-terminus for 59 isoforms, while five isoforms are predicted as soluble (Fig. 1B). Among PG isoforms, there are also four



**Fig. 1.** *Arabidopsis* PME (A) and PG (B) structural motifs. The domains, average size of proteins in amino acid (AA), isoelectric point (pI), and molecular weight (MW) ranges of representative members are indicated according to the PFAM database (<http://pfam.sanger.ac.uk/>), SignalP4.0 (<http://www.cbs.dtu.dk/services/SignalP/>), and ExpASyProtParam Tool (<http://web.expasy.org/protparam/>).

atypical protein sequences, three of which harbour two GH28 domains and one with a reverse transcriptase-like 3 domain (RVT3; Pfam13456) downstream of the GH28 domain. Two asparagine and histidine residues have been identified in the GH28 domain for 53 sequences. One ORF, previously annotated as putative PG QRT3 (At4g20050; Rhee et al., 2003), lacks the GH28 domain but has a pectate lyase 3 domain (Pfam12708). The number of putative PGs would therefore be 67, in line with recent published data (Cao, 2012).

For AtPLLs, all the sequences annotated as PL (26 sequences) show a Pec\_lyase\_C domain (Pfam00544), and only three isoforms are predicted as soluble proteins (Fig. 2A). This multigenic family also has specific features, with the presence of an N-terminal domain, called Pec\_lyase\_N (Pfam04431), in four sequences. Lastly, one protein (At3g54920 or PMR6), which plays a role in powdery mildew resistance, shows an additional C-terminal glycosylphosphatidylinositol (GPI) anchor site, which may be responsible



**Fig. 2.** *Arabidopsis* PLL (A) and PAE (B) structural motifs. The domains, average size of proteins in amino acid (AA), isoelectric point (pI), and molecular weight (MW) ranges of representative members are indicated according to the PFAM database (<http://pfam.sanger.ac.uk/>), SignalP4.0 (<http://www.cbs.dtu.dk/services/SignalP/>), and ExpASyProtParam Tool (<http://web.expasy.org/protparam/>).

for this resistance (Vogel *et al.*, 2002; Sun and Van Nocker, 2010).


Concerning AtPAE, all 12 annotated sequences have a PAE domain (Pfam03283) and N-terminal targeting motifs (SP, TM, or SP/TM) (Fig. 2B). No isoform is predicted to be soluble.

## HGMEs display distinct patterns of expression

### *A number of isoforms are co-expressed*

Analysis of public microarray data sets for all genes encoding HGMEs has led to the identification of several non-exhaustive co-expression clusters in *A. thaliana* (Genevestigator, <http://www.genevestigator.com/>; Hruz *et al.*, 2008). For instance, for the six clusters reported in this review, a number of genes encoding PME, PAE, PG, and PLL are co-expressed (Fig. 3). In cluster 1, six *PME* genes, five *PAE* genes, three *PG* genes, and 11 *PLL* genes are mainly co-expressed in seedlings,

leaf, and root tissues. In contrast, the majority of these genes are not expressed in seed-related tissues. In this cluster, ~42% of *PAE* and *PLL* genes are present. Cluster 2 contains only *PME* and *PG* genes specifically expressed in root tissue, while cluster 3 shows *PME*, *PAE*, *PG*, and *PLL* genes expressed in all tissues selected for the analysis apart from roots. Cluster 4 comprises *PME*, *PG*, and *PLL* genes specifically co-expressed in pollen. The expression and roles of *At1g69940*, *At2g47030*, *At2g47040*, and *At3g62170* in pollen germination and pollen tube growth have been highlighted (Jiang *et al.*, 2005; Röckel *et al.*, 2008; Wolf *et al.*, 2009b; Mollet *et al.*, 2013). In addition, several genes encoding PGs, PMEs, and PLLs are expressed in anthers (Gonzalez-Carranza *et al.*, 2002, 2007; Rhee *et al.*, 2003; Jiang *et al.*, 2005; Francis *et al.*, 2006; Wolf *et al.*, 2009b; Sun and Van Nocker, 2010) with the encoded proteins, namely AtPME At5g55590 and putative AtPG At4g20050, playing a role in pollen grain formation and tetrad separation (Rhee *et al.*, 2003; Francis *et al.*, 2006). Furthermore, transcriptomic analysis in the stamen abscission zone of *Arabidopsis* revealed a co-expression between



**Fig. 3.** Clustering analysis of *PME*, *PAE*, *PG*, and *PLL* mRNA expression during development in *A. thaliana*. *PME* (red circles), *PAE* (green circles), *PG* (orange circles), and *PLL* (blue circles) are shown together. Microarray data and cluster analysis was carried out using Genevestigator (<https://www.genevestigator.com/gv/>; Hruz et al., 2008). Only some specific tissues were selected. Probes with a single gene and genes showing a minimal expression were used for the cluster analysis. In all clusters, for each HGME family, the proportion (%) of genes expressed among all family members is indicated.

*PAE*, *PME*, *PG*, and *PLL* genes (Lashbrook and Cai, 2008). In cluster 5, a few *PME*, *PAE*, *PG*, and *PLL* genes exhibit a ubiquitous expression. Interestingly, a small cluster composed of three *PME* genes only expressed in general and chalazal seed coat was also identified. It could relate to the emerging roles of *PME*–*PMEI*-mediated control of the DM of HG in mucilage structure and extrusion (Rautengarten et al., 2008; Arsovski et al., 2010; Saez-Aguayo et al., 2013; Voiniciuc et al., 2013). From this analysis, it appears that neither *PAE* nor *PLL* genes are specifically expressed in root and seed coat. In addition, no *PAE* genes appear to be solely expressed in pollen. In contrast, *PME* genes are identified in all of these clusters, potentially highlighting the major role of the control of the DM of HG in a large range of vegetative and reproductive developmental processes. Overall, this analysis shows that the control of HG structure and chemistry is likely to be a highly integrated process involving, in some developmental processes, several different HGMEs.

#### Regulation of HGME gene expression through hormone signalling

The gene expression of a number of HGMEs has been shown to be modulated, either directly or indirectly, through hormone signalling. A number of publications related to the analysis of phytohormone signalling, including mutants in hormone synthesis pathways and/or hormone application, report changes in the expression of several genes encoding HGMEs (Goda, 2004; Vanneste et al., 2005; Che et al., 2006; Laskowski et al., 2006; Derbyshire et al., 2007; Palusa et al., 2007; Swarup et al., 2008; Quesada et al., 2009; Curvers et al., 2010; Kanai et al., 2010; Di Matteo et al., 2010; Sun et al., 2010; Osorio et al., 2011a; Savatin et al., 2011; Ribeiro et al., 2012; Braybrook and Peaucelle, 2013). This has consequences on the structure of HGs affecting plant development. It has also been shown

that local auxin accumulation at the shoot apex of *A. thaliana* leads to local demethylesterification of HG, suggesting a role for auxin in the control of *PME* activity (Braybrook and Peaucelle, 2013). Several expression data sets reveal that auxin specifically regulates the expression of *PME*, *PAE*, *PG*, and *PLL* genes during various developmental events, such as lateral root emergence (Vanneste et al., 2005; Laskowski et al., 2006; Swarup et al., 2008), adventitious root formation (Savatin et al., 2011), fruit development (Quesada et al., 2009; Osorio et al., 2011a), and seedling development (Goda et al., 2004; Palusa et al., 2007). In addition, in the latter study, *PME*, *PAE*, and *PG* genes appeared to be up-regulated by brassinosteroids (BRs). Potential regulation of *PME* expression by BRs was also observed in the transcriptome analysis of mutant lines for the *Arabidopsis* transcription factor *AtBZR1*, responsible for regulating the expression of specific target genes involved in developmental processes as diverse as cell elongation and root development. Based on these results, *AtPME2* and *AtPME3* were identified as putative targets of *AtBZR1* (Sun et al., 2010). The regulation of *AtPME41* gene expression during chilling stress in *atbzr1-1D* and *atbri1*, two mutants of the BR signalling pathway, together with the recent data obtained using a *PMEI* overexpressor, reinforce the hypothesis of the control of *PME* by BRs (Qu et al., 2011; Wolf et al., 2012b). When considering other HGMEs, it has been shown that auxin, together with cytokinins, has an effect on the expression of *PLL* and *PGIP* (*PG* inhibitor protein) genes in shoot, root, and callus development in *Arabidopsis* tissue culture (Che et al., 2006). In general, it is likely that all phytohormones directly or indirectly control HGME gene expression. For instance, gibberellic acids (GAs) have an effect on the expression of genes encoding *PLLs* and *PMEs* during rosette expansion in *A. thaliana* (Ribeiro et al., 2012). This is consistent with results showing a change in *PME* activity and in the DM of pectins in GA-deficient mutants over the course

of hypocotyl growth (Derbyshire *et al.*, 2007). Abscisic acid (ABA) can also regulate development, through the modulation of changes in pectin. For example, ABA-deficient tomato mutants cause changes in pectin composition, including levels of Gal-A (Curvers *et al.*, 2010). This could be related to changes in the expression/activity of pectin-degrading enzymes such as PLLs, known to be regulated by this phytohormone (Palusa *et al.*, 2007), or in the expression of the gene encoding PGIP, which was suggested to control PG activity during germination (Kanai *et al.*, 2010). Identification of the major involvement of HGMEs in changes in fruit texture during maturation has prompted research into the role of ethylene in regulating their gene expression. Early work on tomato (Gaffe *et al.*, 1994) and banana (*Musa acuminata*) showed that PME and PL gene expression is ethylene dependent (Dominguez-Puigjaner *et al.*, 1997) during fruit maturation. More recently, several genes encoding PME, PME1, PG, and PLL were shown to be strongly expressed in the early stages of banana ripening, a period corresponding to the burst in ethylene production (Mbéguié-A-Mbéguié *et al.*, 2009; Srivastava *et al.*, 2012). In tomato, the expression of one PME and one PG gene was associated with inhibition of ripening, when the ethylene receptor *Never-ripe* was mutated (Osorio *et al.*, 2011b). Similarly, the overexpression of the *FaPEI* gene, encoding a PME from strawberry (*Fragaria vesca*), led to up-regulation of the expression of one gene involved in the ethylene response (Osorio *et al.*, 2011a). In distinct species, several PG genes showed specific changes in gene expression, which were ethylene dependent in oil palm (Roongsattham *et al.*, 2012). The role of ethylene in the control of HGME gene expression might go beyond the specific case of fleshy fruit maturation as the *Arabidopsis* transcription factor AtRAP2.6L (a member of the ethylene response factor family), which has a function in shoot regeneration, targets the *AtPG* gene *At3g15720*, which was down-regulated in the *atrap2.6l* T-DNA mutant (Che *et al.*, 2006). These regulation pathways, in particular that of auxin, may be controlled through OG-mediated negative feedback (Ferrari *et al.*, 2013). The antagonism between auxin and OGs has previously been shown during root formation in tobacco (Bellincampi *et al.*, 1993) and by using exogenous OGs in maize seedlings (Peña-Urbe *et al.*, 2012). Exogenous OGs inhibit the expression of auxin-induced genes (*IAA5*, *SAUR16*, etc.), leading to inhibition of adventitious root formation (Savatin *et al.*, 2011). Inhibition by OGs targets late rather than early auxin-responsive genes (Mauro *et al.*, 2002). In *A. thaliana*, a few PME and PG genes showed significant changes in expression in response to treatment with OGs, which could be associated with calcium signalling pathways (Moscatello *et al.*, 2006).


## Regulation of biochemical activity

### *Mode of action and regulation of plant PMEs*

As shown previously, PMEs are commonly present in plants and microorganisms such as fungi and bacteria (Giovane *et al.*, 2004; Pelloux *et al.*, 2007). In plants, several neutral and basic isoforms (Micheli, 2001; Giovane *et al.*, 2004;

Verlent *et al.*, 2007; Jolie *et al.*, 2010; Dixit *et al.*, 2013), as well as a few acidic isoforms (Lin *et al.*, 1989; Bordenave and Goldberg, 1994; Mareck *et al.*, 1995; Micheli *et al.*, 2000; Ding *et al.*, 2000, 2002; Thonar *et al.*, 2006), have been identified. Although all isoforms catalyse the specific hydrolysis of methylester bonds at C-6 from Gal-A residues, and the subsequent formation of free carboxyl groups, release of methanol (MeOH), and acidification of the cell wall (Micheli, 2001; Giovane *et al.*, 2004; Pelloux *et al.*, 2007; Jolie *et al.*, 2010), PME activity is dependent upon a rather large range of factors. For instance, the presence of free carboxyl groups near the active site appears to be required for enzyme action, which might explain the affinity of PME for partially demethylesterified HG (Rexova-Benkova and Markovic, 1976; Bordenave, 1996; Grasdalen *et al.*, 1996; Van Alebeek *et al.*, 2003; Fries *et al.*, 2007). The reaction mechanism of PMEs, which act as monomers according to resolved crystallographic structures (Johansson *et al.*, 2002; Di Matteo *et al.*, 2005; Fries *et al.*, 2007), consists of a nucleophilic attack and an acid/base catalysis by conserved aspartate catalytic residues on the carbonyl carbon of the C-6 methylester of Gal-A (Fries *et al.*, 2007). Co-crystallization between *Erwinia* PME and various substrates highlighted a preference towards substrates with an alternation of methylesterified and non-methylesterified Gal-A residues, corresponding to partially methylesterified HGs. Consequently, the presence of methylesterified Gal-A residues upstream and non-methylesterified Gal-A residues downstream of the catalytic site could determine the processive action of *Erwinia* PME (Fries *et al.*, 2007). Moreover, recent results using molecular dynamics approaches on the *Erwinia* PME suggest that the rotation of the substrate is necessary for access to the subsequent site and the processive demethylesterification of HG by the enzyme (Mercadante *et al.*, 2013). A similar mechanism could be hypothesized for plant PMEs that have a similar mode of action.

Several factors can affect plant PME activity, which appears to be very sensitive to changes in pH and in the concentrations of cations (Catoire *et al.*, 1998; Denès *et al.*, 2000; Ly-Nguyen *et al.*, 2004; Do Amaral *et al.*, 2005; Verlent *et al.*, 2007; Jolie *et al.*, 2009; Dixit *et al.*, 2013). The effect of pH on PME activity may be related to the pI of the isoforms, which is neutral to alkaline for most plant and bacterial PMEs and acidic to neutral for fungal PMEs. In an acidic environment, alkaline plant PMEs may be positively charged and poorly active due to a strong interaction between PMEs and the negatively charged free carboxyl groups of HG. This has consequent effects on growth (Fig. 4). In contrast, at slightly alkaline pH, basic isoforms are less positively charged and can be released from the substrate due to electrostatic repulsion between the enzyme and the free carboxyl groups (Bordenave, 1996; Jolie *et al.*, 2010). However, as some acidic plant PMEs have been identified in various species such as flax (Mareck *et al.*, 1995), mung bean (Bordenave and Goldberg, 1994), jelly fig (Lin *et al.*, 1989; Ding *et al.*, 2000, 2002), chicory (Thonar *et al.*, 2006), and aspen (Micheli *et al.*, 2000), distinct scenarios are likely to occur in the cell wall (Fig. 4). Thus, the control of the DM of pectins probably depends on the recruitment of specific PME pools, according to the DM of the substrate,



**Fig. 4.** Schematic diagram showing the regulation of HGME pools in distinct cell wall microenvironments and consequent effects on growth. In the presence of auxin, the auxin-binding protein 1 (ABP1) receptor, localized at the plasma membrane (PM), activates  $\text{Ca}^{2+}$  and  $\text{K}^{+}$  influx into the cytoplasm (Philippart et al., 1999, 2004; Shishova and Lindberg, 2010; Perrot-Rechenmann, 2010; Tomas et al., 2010; Sauer and Kleine-Vehn, 2011) as well as  $\text{H}^{+}$ -ATPase, leading to  $\text{H}^{+}$  efflux into the cell wall and acidification of the apoplast (Perrot-Rechenmann, 2010; Tomas et al., 2010; Sauer and Kleine-Vehn, 2011). In parallel, auxin imported into the cytoplasm through AUX1/LAX penetrates the nucleus, where it acts on the expression of auxin-regulated genes via the action of TIR1, AUX/IAA, and ARF proteins (Robert and Friml, 2009; Shishova and Lindberg, 2010; Hayashi, 2012). ABP1 can induce downstream responses by regulating gene expression (Rayle and Cleland, 1992; Rober-Kleber et al., 2003; Perrot-Rechenmann, 2010; Monshausen et al., 2011), including that of HGME genes (Vanneste et al., 2005; Laskowski et al., 2006; Swarup et al., 2008). PMEs (red, green), PAEs (yellow), PNLs (dark purple), PGs (orange), and PMEIs (blue) are synthesized in the endoplasmic reticulum (ER) and matured in the *trans*-Golgi network (TGN), before secretion into the cell wall (Wolf et al., 2009b; De Caroli et al., 2011a, b). In parallel, HGs are synthesized in the *cis*-Golgi by galacturonosyltransferases (GAUTs), methylated and acetylated by pectin methyltransferases (PMTs) and pectin acetyltransferases (PATs), respectively, in the medial-Golgi, and secreted into the cell wall as highly methylesterified (HM) and slightly acetylated (SA) forms (Atmodjo et al., 2013). In an acidic cell wall context (left), basic PMEs (in green), which are the major PME isoforms, are strongly positively charged (represented by positive symbols) and can be trapped by free carboxyl groups (represented by negative symbols) (Jolie et al., 2010) (1). In parallel, acidic pH stimulates inhibition of basic PMEs by several PMEIs (Raiola et al., 2004). This leads to a decrease in HG demethylesterification. HGs can become targets of PG and PNL, which depolymerize HGs, leading to the release of OGs, cell wall loosening, and rapid growth. In the same conditions, acidic PMEs (in red) are neutrally charged (represented by positive and negative symbols) and can act on HGs by multiple chain demethylesterification and/or be inhibited by PMEIs (2) (Catoire et al., 1998; Denès et al., 2000). Following PME action, PAEs can deacetylate HGs (Williamson, 1991; Bordenave et al., 1995). Randomly demethylesterified and deacetylated HGs are depolymerized by PGs and PNLs (Pressey and Avants, 1973; Themmen et al., 1982; Mayans et al., 1997; Protsenko et al., 2008), leading to the release of OGs, cell wall loosening, and rapid growth. OGs can increase the apoplasmic pH and control gene expression through binding to membrane receptors such as PBD-EGF-EGF-WAK1 (Wolf et al., 2012a; Ferrari et al., 2013). In the absence of auxin (right),  $\text{H}^{+}$  is not imported into the cell wall, and  $\text{Ca}^{2+}$  and  $\text{K}^{+}$  are not exported into the cytoplasm. Consequently, the cell wall is slightly alkaline with a high ionic concentration. Auxin-independent HGME genes can be expressed, generating distinct pools. In these conditions, acidic PMEs (in red) are strongly negatively charged (represented by negative symbols). By electrostatic repulsion with the free carboxyl group, acidic PMEs cannot bind to their substrate. In parallel, basic PMEs (in green) are less positively or neutrally charged (represented by positive and negative symbols) and can bind to HG HM/SA (Jolie et al., 2010). Slightly alkaline pH can decrease PME–PMEI complex formation and stability, but other PMEIs showing various pH sensitivities can be stimulated and inhibit acidic and basic PMEs (Raiola et al., 2004). Basic PMEs perform the demethylesterification of the HG HM/SA, by single chain or multiple attacks, leading to processively demethylesterified HG (Catoire et al., 1998; Denès et al., 2000; Ngouémazong et al., 2012). PAEs could subsequently act on processively demethylesterified HGs (Williamson, 1991; Bordenave et al., 1995). Depending on  $\text{Ca}^{2+}$  content and PG/PL presence, processively demethylesterified and deacetylated HGs can form so-called egg box structures through binding with  $\text{Ca}^{2+}$ , leading to cell wall strengthening and growth retardation (3). Several basic PGs and PLs, stimulated by  $\text{Ca}^{2+}$  (Pressey and Avants, 1973; Themmen et al., 1982; Protsenko et al., 2008; Chotigeat et al., 2009; Wang et al., 2010), can depolymerize HG to form OGs, which could increase apoplasmic pH and lead to residual growth. The egg box and OGs can regulate gene expression



the pI of the isoforms, and the pH microenvironment at the cell wall. Therefore, depending on the patterns of demethylesterification produced, the properties of pectin are likely to be modified and could differentially affect cell wall rheology and cell growth (Fig. 4).

In plants, cations are often essential for PME activity (Jolie *et al.*, 2010). Although some salt-independent plant PME isoforms have been described, PME activity normally increases up to an optimal cation concentration, above which it decreases (Do Amaral *et al.*, 2005; Verlent *et al.*, 2007; Jolie *et al.*, 2009; Videcoq *et al.*, 2011; Dixit *et al.*, 2013). This optimal concentration is highly dependent upon the type of cation. For example, some salts, such as NaCl or KCl, have a positive effect on the activity up to a certain range of concentration, while others, such as LiCl, are strong inhibitors regardless of the concentration (Do Amaral *et al.*, 2005; Verlent *et al.*, 2007; Dixit *et al.*, 2013). This latter effect is related to competition between cations and positively charged PMEs for interaction with the free carboxyl groups of HG (Fig. 4). In acidic or slightly alkaline cell wall contexts, basic PMEs, positively charged, could be more sensitive to the ionic environment than acidic PMEs.

Different modes of action of plant PMEs have been reported. Three mechanisms have been described: (i) a single-chain mechanism, where PMEs remove all contiguous methyl-esters from a single chain of HG before dissociating from the substrate; (ii) a multiple-attack mechanism, in which several PMEs catalyse the release of a limited number of methyl-esters on several chains of HG; these two modes of action produce similar processively demethylesterified HG and are often attributed to basic PMEs, in plants and in bacteria (Johansson *et al.*, 2002; Willats *et al.*, 2006; Jolie *et al.*, 2010); (iii) a multiple-chain mechanism, where PMEs remove only one methylester before dissociating from Gal-A. This mechanism is likely to be that of acidic PMEs in plants and in fungi (*Aspergillus* sp.) and allows a random demethylesterification (Micheli, 2001; Johansson *et al.*, 2002; Willats *et al.*, 2006; Jolie *et al.*, 2010). However, fungal PME from *Trichoderma reesei* shows a processive demethylesterification. Thus these different modes of action are probably dependent upon several factors including the enzyme properties characterized by the pH and ionic environment as well as the substrate specificity, rather than the origin of PMEs (Denès *et al.*, 2000; Johansson *et al.*, 2002; Videcoq *et al.*, 2011). In fact, the action of apple PME has been shown to be pH dependent, with a possible shift between a blockwise and non-blockwise mode of action (Catoire *et al.*, 1998). In addition, the activity of isoforms can vary for the same given substrate (Ly-Nguyen *et al.*, 2004; Do Amaral *et al.*, 2005; Jolie *et al.*, 2009).

Plant PMEs can also be regulated by inhibition of their activities by proteinaceous or non-proteinaceous compounds. Protein inhibitors of plant PMEs (PMEIs) can form

a stoichiometric 1:1 complex with PMEs (Di Matteo *et al.*, 2005), leading to an inhibition of PME activity which is dependent on the pH and ionic environment of the cell wall (Bellincampi *et al.*, 2004; Giovane *et al.*, 2004; Raiola *et al.*, 2004). Like PMEs, PMEIs belong to large multigenic families in plants, which further questions the occurrence of specific PME–PMEI pairs in the cell wall. Non-proteinaceous compounds include a wide range of substances including OGs, iodine, detergents, tannins, phenolic acids, glycerol, some sugars, and epigallocatechingallate (Rexova-Benkova and Markovic, 1976; Lewis *et al.*, 2008).

#### Regulation of biochemical activity of plant PAEs

In plant cell walls, several polysaccharides such as pectins, XyGs, xylans, and mannans can be acetylated (Ishii, 1997; Ralet *et al.*, 2005; Scheller and Ulvskov, 2010; Orfila *et al.*, 2012). However, acetyl groups are not distributed homogeneously in the cell wall, and for pectins they are mainly clustered in specific regions of HG and RG-I, as shown in sugar beet (Ralet *et al.*, 2005; Orfila *et al.*, 2012). Gal-A residues from RG-I can be *O*-acetylated at the O2 and/or O3 positions (Ishii, 1997), while Gal-A from the HG backbone can be *O*-acetylated at the O2 or O3 position, but not di-acetylated (Ralet *et al.*, 2005). In the HG backbone, the simultaneous presence of acetyl and methyl groups on the same Gal-A residues is observed infrequently, but might influence either PME or PAE activities (Ralet *et al.*, 2005). Several investigations have indicated an effect of acetylation on pectin properties. The presence of acetyl groups on Gal-A of HG has an effect on cell wall viscosity (Gou *et al.*, 2012; Orfila *et al.*, 2012) and impairs the formation of Ca<sup>2+</sup> bonds between HG chains (Renard and Jarvis, 1999; Turquois *et al.*, 1999). Acetylation can also hinder the degradation of HG by some endo-PGs (Bonnin *et al.*, 2003). The deacetylation of HG by enzymatic activity is thus likely to be necessary to trigger changes in the cell wall mediating plant growth.

In this context, PAEs can act by specific hydrolysis of acetylene ester bonds at O2 and/or O3 from Gal-A residues, making up the linear HG and RG-I of pectins (Gou *et al.*, 2012). Indeed, partial deacetylation of HG improved the gelation properties of sugar beet (Williamson, 1991; Ralet *et al.*, 2003) and increased the enzymatic degradation of pectins (Biely *et al.*, 1986; Schols and Voragen, 1994; Chen and Mort, 1996; Benen *et al.*, 1999; Bonnin *et al.*, 2003). PAE enzymes have been identified in plants (Williamson, 1991; Bordenave *et al.*, 1995; Christensen *et al.*, 1996), bacteria (Shevchik and Hugouvieux-Cotte-Pattat, 1997; Bolvig *et al.*, 2003; Shevchik and Hugouvieux-Cotte-Pattat, 2003), and fungi (Kauppinen *et al.*, 1995; Bonnin *et al.*, 2008). Generally, they act specifically on HG and RG-I polymers and seem to be inactive against acetyl groups present in XyGs, xylans, and

---

through their binding to membrane receptor such as PBD-EGF-EGF-WAK1 (Wolf *et al.*, 2012a; Ferrari *et al.*, 2013). Otherwise, in acidic or slightly alkaline cell wall contexts, basic PMEs, positively or neutrally charged, could be more sensitive to the ionic environment than acidic PMEs. Particularly in a slightly alkaline cell wall context, basic PMEs could be in competition with cations such as K<sup>+</sup> (pink circle) and Ca<sup>2+</sup> (light blue circle) for interaction with free carboxyl groups of HG, avoiding basic PME trapping by processively demethylesterified HG. The solid arrows correspond to the actions taking place in the presence of auxin (purple circle), whereas dotted arrows represent the lack of action when there is no auxin signal.

mannans (Bolvig *et al.*, 2003; Bonnin *et al.*, 2008). This differentiates them from rhamnogalacturonan acetyltransferase from *Aspergillus aculeatus*, which is highly specific for RG-I and does not deacetylate HG (Kauppinen *et al.*, 1995). In *Arabidopsis*, the majority of PAE isoforms have a neutral to basic pI. Purified plant PAEs characterized from mung bean and orange, which are highly active against synthetic substrates such as triacetin and *p*-nitrophenyl acetate and sugar beet pectins, have an optimal activity at pH ranging from 5 to 6.5, depending on the substrates used in the assay (Williamson, 1991; Bordenave *et al.*, 1995; Christensen *et al.*, 1996). Moreover, PAE activity is increased when the substrate has previously been demethylated (Williamson, 1991; Bordenave *et al.*, 1995; Oosterveld *et al.*, 2000). Therefore, a synergistic effect between PME and PAE is likely to occur at the cell wall to mediate either egg box formation or degradation of HG by PGs and PLLs.

#### Mode of action and regulation of plant PGs

PGs belong to an enzyme family identified in plants (Pressey *et al.*, 1973, 1977; Themmen *et al.*, 1982; Nogota *et al.*, 1993; Chun and Huber, 1998; Hadfield and Bennett, 1998; Pathak and Sanwal, 1998; Pathak *et al.*, 2000; Verlent *et al.*, 2004, 2005), herbivorous insects (Shen *et al.*, 2003), and microorganisms such as bacteria, fungi, and nematodes (Pickersgill *et al.*, 1998; Jaubert *et al.*, 2002; André-Leroux *et al.*, 2005; Jayani *et al.*, 2005; Kars *et al.*, 2005; Mertens and Bowman, 2011). Whatever their origin, PGs cleave by hydrolysing the  $\alpha$ -(1–4) bonds linking D-Gal-A residues, mainly from the HG linear homopolymer (Brummell and Harpster, 2001; Verlent *et al.*, 2005; Protsenko *et al.*, 2008). In both plants and microorganisms, PGs are constituted of two types, the endo-PGs (EC 3.2.1.15) and the exo-PGs (EC 3.2.1.67) (Brummell and Harpster, 2001; Verlent *et al.*, 2005; Protsenko *et al.*, 2008). Generally, their activities increase with the decrease in the DM of HG (Thibault and Mercier, 1979; Bonnin *et al.*, 2002). Consequently, and in agreement with previous comments, prior action of PME seems to be necessary to enable the degradation of HGs by plant PGs and, for the same DM, the pattern of demethylation could affect the action of plant PGs (Verlent *et al.*, 2005). For example, PG isolated from tomato prefers HG previously demethylated blockwise by the endogenous PME of tomato, rather than randomly demethylated by the exogenous fungal PME of *A. aculeatus* (Verlent *et al.*, 2005). More precisely, endo-PGs catalyse the hydrolytic cleavage of the  $\alpha$ -(1–4) linkage between at least two demethylated Gal-A residues of the HG backbone with a random action pattern, leading to the formation of OGs of various degrees of polymerization (DPs; Verlent *et al.*, 2005; Ferrari *et al.*, 2013). In plants, endo-PGs particularly prefer HG with a chain region containing more than four demethylated Gal-A residues between methylated Gal-A (Clausen *et al.*, 2003; Protsenko *et al.*, 2008). Consequently, endo-PGs could act against HG demethylated blockwise, and the variation of the demethylation pattern may regulate the action of endo-PGs. In contrast, exo-PGs act only at the terminal position of the

HG backbone to hydrolyse the  $\alpha$ -(1–4) linkage, producing monogalacturonides (Verlent *et al.*, 2005; Protsenko *et al.*, 2008). In this case, they could act together against HG, which is partially, randomly, or blockwise demethylated at the terminal position.

In plants such as *Arabidopsis*, all isoforms have a pI varying from 4.8 to 9.8 (Cao, 2012), which could regulate their optimum activity depending on the pH and the ionic microenvironment. With regard to pH, several plant PGs have been partially purified from plants and biochemically characterized. Overall, like PGs from microorganisms such as *Botrytis cinerea*, that show optimal activities at approximately pH 4.5 (Kars *et al.*, 2005), plant PGs prefer an acidic pH from 3.3 to 6 (Pressey *et al.*, 1973, 1977; Themmen *et al.*, 1982; Nogota *et al.*, 1993; Chun and Huber, 1998; Pathak and Sanwal, 1998; Pathak *et al.*, 2000; Verlent *et al.*, 2004). Furthermore, the pH of optimal activity of plant PGs seems to be dependent on the ionic microenvironment. For example, PG purified from tomato shows an optimal activity at pH 4–4.5 with NaCl, compared with pH 5–6 in the presence of KCl (Chun and Huber, 1998). Other ions such as Cd<sup>2+</sup> and Ca<sup>2+</sup> could affect PG activity in *Avena sativa* (Pressey *et al.*, 1977). Regarding Ca<sup>2+</sup>, an exo-PG from peach fruit is Ca<sup>2+</sup> dependent (Pressey *et al.*, 1973) as is an endo-PG from strawberry fruit because with EDTA, which chelates Ca<sup>2+</sup> ions, PG activity is inhibited (Nogota *et al.*, 1993). Nevertheless, not all plant PGs are Ca<sup>2+</sup> dependent. For instance, among three PG isoforms isolated from banana, where two endo-PGs (PG1 and PG3) and one exo-PG (PG2) are characterized, only PG1 was shown to be activated by Ca<sup>2+</sup> (Pathak *et al.*, 1998). Consequently, ion dependence does not seem to be related to the mode of action of PGs (endo- or exo-PGs).

PG activity can also be inhibited by proteinaceous compounds (Juge, 2006; Protsenko *et al.*, 2008). It has been widely demonstrated in plants that there are PGIPs directed against secreted pathogen PGs (Federici *et al.*, 2001; Di Matteo *et al.*, 2003; Ferrari *et al.*, 2006; Oelofse *et al.*, 2006; Sathiyaraj *et al.*, 2010; Benedetti *et al.*, 2011). Plant PGIPs are bound to HG substrate as a guardian, thus inhibiting pathogen PG activity by formation of a stoichiometric complex, which prevents access to the substrate (Spadoni *et al.*, 2006). More precisely, when PGIPs interact with pathogen PG, the complex formed can move the substrate away, thus preventing PG action against the HG substrate (Spadoni *et al.*, 2006). Moreover, it appears that PGIPs act mainly against endo-PGs from pathogens (Protsenko *et al.*, 2008). This is confirmed by the mode of action of PGIPs, which bind strongly to HG demethylated blockwise, in contrast to HG randomly demethylated, and prevent the action of endo-PGs (Spadoni *et al.*, 2006). Currently, limited data indicate a role for plant PGIPs in the regulation of plant PG activity and development. However, a recent publication shows changes in radicle protrusion in *pgip1* mutants and PGIP1 overexpressors, probably caused by an alteration of *Arabidopsis* PG activity (Kanai *et al.*, 2010).

#### Mode of action and regulation of plant PLLs

PLLs are mainly found in plants (Domingo *et al.*, 1998; Chourasia *et al.*, 2006; Chotigeat *et al.*, 2009; Wang *et al.*,

2010) and microorganisms such as bacteria, fungi, and nematodes (Popejusz *et al.*, 2000; Jayani *et al.*, 2005). They cleave, by  $\beta$ -elimination, the  $\alpha$ -(1–4) bond linking methylesterified or non-methylesterified D-Gal-A units mainly from the HG backbone, giving rise to an unsaturated C4–C5 bond at the non-reducing end of the newly formed OG (Mayans *et al.*, 1997; Pilnik and Rombouts, 1981). PLLs comprise the PL and PNL family of enzymes. PLs are more specific for non-methylesterified or slightly methylesterified HG and require  $\text{Ca}^{2+}$  for their activity whose optimal pH is near 8.5. In contrast, PNLs degrade highly methylesterified HG with an optimal pH of 5.5 for their activity and do not require  $\text{Ca}^{2+}$  (Mayans *et al.*, 1997; Herron *et al.*, 2000). PLs have both endo (EC 4.2.2.2) and exo (EC 4.2.2.9) activities, while only endo (EC 4.2.2.10) activity has been discovered for PNLs (Sinitsyna *et al.*, 2007). PLs and PNLs are commonly found in microorganisms; fungi usually secrete PNLs (Sinitsyna *et al.*, 2007) while bacteria produce predominantly PLs (Payasi and Sanwal, 2003).

To date, of PNLs and PLs, only PLs have been discovered and biochemically characterized in plants. For instance, the plant PL from *Zinnia elegans* shares homology with the microbial PLs from *Bacillus subtilis* and *Pectobacterium carotovorum* (Domingo *et al.*, 1998). Moreover, the presence of PL isoforms has been demonstrated in plants such as *Hevea brasiliensis*, *Z. elegans*, and *Gossypium hirsutum*, and in the ripening fruits of banana. More precisely, coding sequences isolated from these latter and expressed in a heterologous system led to the production of *Escherichia coli* or yeast expressed recombinant enzymes showing PL activities (Domingo *et al.*, 1998; Pua *et al.*, 2001; Chotigeat *et al.*, 2009; Wang *et al.*, 2010). PLs from pathogens are rather active at alkaline pH (Jayani *et al.*, 2005); recombinant PLs from *H. brasiliensis*, *Z. elegans*, and mango fruit also show optimal activities at basic pH, namely pH 7, 10, and 8, respectively (Domingo *et al.*, 1998; Chourasia *et al.*, 2006; Chotigeat *et al.*, 2009). Moreover, as in microorganisms, PL activity from *H. brasiliensis* and mango fruit is inhibited by EDTA and stimulated by  $\text{CaCl}_2$  (Mayans *et al.*, 1997; Chourasia *et al.*, 2006; Chotigeat *et al.*, 2009). A multiple sequence alignment suggests that the calcium-binding site of plant PLs could be conserved and may involve three aspartate residues (Chotigeat *et al.*, 2009). Lastly, plant PLs, like those from bacteria (*E. chrysanthemi*), seem to be more active against non-methylesterified HG because their optimal activities have often been quantified with non-methylesterified substrates (Tardy *et al.*, 1997; Chotigeat *et al.*, 2009; Wang *et al.*, 2010). To date, a PNL inhibitor protein has been found (Bugbee, 1993), but no proteic inhibitor of PL enzyme.

## Roles of HG modifications in vegetative and reproductive development

Early reports showed the multiple roles of pectin modifications in the control of vegetative development (Hasunuma *et al.*, 2004; Pilling *et al.*, 2004). The current review focuses on the role(s) of HGME-mediated HG modification during

cell elongation and differentiation in specific developmental processes.

### Roles of HGMEs in organ growth

Understanding the role of pectin modifications in the control of growth requires the use of simple models in which developmental and cell biology, genomics, biochemistry, and biophysics can be integrated at a cellular level. The pollen tube, such as dark-grown hypocotyl, is a powerful system to analyse the roles of the cell wall in modulating cell elongation.

PMEs have been reported to play a role in pollen grain formation. In *A. thaliana*, the *atqrt1* mutant, which does not express the *AtQRT1* gene encoding a PME (At5g55590), does not show cell wall degradation and separation of the haploid spores during microsporogenesis (Francis *et al.*, 2006). Consequently, the spores remain fused and pollen grains are released as tetrads. The modifications of HGs by PMEs could play a central role in the first step of wall degradation, by creating substrates for pectin-degrading enzymes. A similar phenotype was shown for mutants in the *AtQRT3* gene encoding a putative PG (At4g20050; Rhee *et al.*, 2003). A model was proposed in which the action of PME would create specific substrates for downstream enzymes. The structure of the pollen grain might be a determinant of the subsequent capacity for pollen tube emergence. In *A. thaliana*, the *AtVGD1* gene (At2g47040), which encodes a PME, is involved in pollen tube elongation. The *atvgd1* knockout (KO) mutant showed a slight reduction in pollen PME activity together with a retarded pollen tube growth within the style and transmitting tract. In addition, the pollen tube had an abnormal shape with frequent tip explosions (Jiang *et al.*, 2005). The mutant had lower than wild-type levels of pollen fertility, and hence smaller siliques with fewer seeds. A similar phenotype, albeit less drastic, was observed for the *AtPPME1* KO mutant (Tian *et al.*, 2006). In other species, such as *Nicotiana tabacum*, silencing of *NtPPME1*, encoding the main tobacco PME isoforms, led to a decrease in pollen tube growth (Bosch and Hepler, 2006). Recent advances in the understanding of the role of HG modifications in pollen tube elongation include the tight spatial and temporal regulation of PMEs by PMEIs. For instance, it was shown that *AtPMEI2* inactivates *AtPPME1* *in vitro* and that both proteins are located in the pollen tube where they physically interact (Röckel *et al.*, 2008). More recently, *BoPMEI1*, a novel *Brassica oleracea* gene, was characterized (Zhang *et al.*, 2010). Heterologously expressed *BoPMEI1* showed PME activity while a transgenic *Arabidopsis* plant, expressing antisense *BoPMEI1*, suppressed the expression of the orthologous gene *At1g10770* altering pollen tube growth. The fine control of PME activity modulates HG structure on the lateral sides of the pollen tube, with consequences on cell wall rheology, enabling apical growth. In contrast to PMEs, the functional characterization of the roles of PAEs, PGs, and PLLs in pollen tube growth has remained elusive. However, a recent report showed a role for the control of the DA of pectins in pollen tube growth (Gou *et al.*, 2012). In *Brassica campestris*, a putative BcPG encoded by *BcMF2* is specifically expressed in tapetum and

pollen after the tetrad stage of anther development. In a transgenic plant with reduced levels of *BcMF2* expression, mature pollen presents a distorted morphology with abnormal intine development, leading to abnormal pollen tube growth and a consequent reduction in male fertility (Huang *et al.*, 2009a). In the same species, *BcMF9*, encoding a distinct BcPG, was shown to play a role in intine and exine formation (Huang *et al.*, 2009b). It is therefore likely that PGs are involved in the changing intine cell wall structure affecting subsequent pollen tube development. Similarly, the activity of PL Cry j I, expressed in pollen of *Cryptomeria japonica*, could cause cell wall loosening during pollen development, thus improving pollen tube emergence and growth as well as its penetration within the style (Taniguchi *et al.*, 1995). Overall, recent data generated about the control of HG structure, including the role of HG acetylation (Gou *et al.*, 2012) during pollen development, have enabled new models to be established linking pollen tube growth to the spatial distribution of polysaccharides (Zonia and Munnik, 2011; Chebli *et al.*, 2012; Mollet *et al.*, 2013).

On the vegetative side, PME play a role in the early stages of radicle emergence, as shown by the faster germination rate in *A. thaliana* plants overexpressing *AtPMEI5* (Müller *et al.*, 2013). PME activity would therefore modulate the mechanical properties of the cell wall, between opposing forces of radicle elongation and resistance of the endosperm. PG activity can also be involved in radicle protrusion. For example, when the *AtPGIPI* gene, which is regulated by the transcription factor ABA insensitive 5 (ABI5), is mutated or overexpressed, PG activity is modified, leading to changes in the seed coat mucilage released and the timing of radicle protrusion (Kanai *et al.*, 2010). Seed coat mucilage production has been shown to be affected by PME (Rautengarten *et al.*, 2008; Voiniciuc *et al.*, 2013; Saez-Aguayo *et al.*, 2013).

Once germinated, dark-grown hypocotyl has a simple anatomy that elongates, in the absence of cell division, from 10 µm up to 1 mm during post-embryonic development (Gendreau *et al.*, 1997). Previous work has shown that the initial elongation rate of hypocotyl cells is developmentally controlled. At first, all cells elongate uniformly and slowly up to 48 h after germination, after which abrupt growth acceleration takes place (Refregier *et al.*, 2004). An important relationship has been demonstrated between the DM of HGs in primary cell wall and hypocotyl elongation in *A. thaliana*, as well as other species (Al-Qsous *et al.*, 2004; Paynel *et al.*, 2009). First, two mutants deficient in GA biosynthesis (*gal-3* and *gai*) showed changes in the DM of HGs, with consequences on hypocotyl length (Derbyshire *et al.*, 2007). More recently, microarray analysis in *A. thaliana* showed that several genes, encoding PMEs, PMEIs, PGs, and PLLs, were up- or down-regulated at the growth phase transition (Pelletier *et al.*, 2010). The down-regulation of PME expression, using the *atpme3* KO mutant, had consequences on hypocotyl length (Guénin *et al.*, 2011). The overexpression of one of these genes, *AtPMEI4* (*At4g25250*), increased carboxylic ester bonds in the primary cell wall of dark-grown hypocotyls and delayed growth (Pelletier *et al.*, 2010). A similar approach showed that the overexpression of *AtPMEI5* (*At2g31430*) had

dramatic effects on plant growth, including a drastic reduction in dark-grown hypocotyl length (Wolf *et al.*, 2012b). It was further shown that feedback signalling from the cell wall is integrated by the BR signalling module to ensure homeostasis of cell wall biosynthesis and remodelling. This adds a new component to understanding the regulation and roles of pectin modifications in the control of growth rate. BRs have been shown to modulate PME activity and the expression of specific PME genes (Qu *et al.*, 2011). Surprisingly, the overexpression of another *PMEI*, *AtPMEI-2*, did not show similar results (Lionetti *et al.*, 2010). This suggests that, depending on the *PMEI* used, specific PME targets could be present or absent, with consequences on the cell wall and growth phenotype. This introduces a fascinating perspective in understanding the role of the sensing of cell wall integrity and its consequences on modulating cell growth (Kohorn and Kohorn, 2012).

When considering root growth, a study showed that the *AtPME3* gene encoding one of the major PME isoforms in *A. thaliana* plays a role in controlling root elongation. The *atpme3* KO mutant, which showed decreased PME activity, had a 20% reduction in root length compared with the wild type, while *AtPME3* overexpressors showed the opposite phenotype (Hewezi *et al.*, 2008). By using a distinct KO allele, it was shown that the reduction in PME activity in the *atpme3* mutant correlated with an increased DM of HGs (Guénin *et al.*, 2011). The control of PME activity by specific PMEIs is likely to play a key role in the regulation of root growth. Transgenic plants overexpressing *AtPMEI-1* and *AtPMEI-2* showed a 50% decrease in PME activity and an increased root length compared with the wild type; this was notably related to changes in cell size in the root expansion zone (Lionetti *et al.*, 2007). These changes were associated with modifications in leaf shape/size. Recent data demonstrated a link between expression of specific PME genes and the AI-induced inhibition of root elongation in rice (Yang *et al.*, 2012). Interestingly, the number of adventitious roots was modified in PME mutants, suggesting that these enzymes influence both root elongation and root differentiation (Guénin *et al.*, 2011).

### Roles of HGMEs in organ formation

As mentioned previously, the changes in the pectic network affect root emergence as shown by the modifications of the number of adventitious roots in specific *atpme* KO mutant lines (Guénin *et al.*, 2011). This could be related to OG-mediated signalling (Savatin *et al.*, 2011) and/or to phytohormone-related signalling. In this respect, auxin homeostasis is likely to be a major signal regulating the expression of HG-modifying genes, such as *PG* and *PLL* genes, during root emergence (Swarup *et al.*, 2008). Although the functional role of PLLs in root differentiation has not yet been elucidated, the increase in transcript accumulation in auxin-treated roots (Laskowski *et al.*, 2006) as well as the large number of isoforms expressed in this organ zone suggest a major involvement of these enzymes (Sun and Van Nocker, 2010). Recent data showing a role for a *Lotus japonicus* PL in the

modification of roots required for rhizobia infection support this hypothesis (Xie *et al.*, 2012).

The phyllotaxis of plant organs, which is the precise emergence of lateral organs, is controlled by a gradient of the plant hormone auxin (Rybel *et al.*, 2010; Vernoux *et al.*, 2010; Besnard *et al.*, 2011; Santuari *et al.*, 2011; Sassi *et al.*, 2012), but the chemical and mechanical status of the cell wall is also important in the formation of new organs. More particularly, the DM of HGs can affect cell wall rheology, modifying phyllotaxis. A first study has shown that the formation of flower primordia in *A. thaliana* shoot apical meristem is accompanied by a demethylesterification of HGs. In fact, the overexpression of *AtPME3* (*At5g20740*) and *AtPME5* (*At5g47500*), which are co-expressed in the shoot apical meristem area, alters the methylesterification status of HGs. This can lead to inhibition of either primordia formation when *AtPME3* is expressed or ectopic primordia formation with *AtPME5* expression (Peaucelle *et al.*, 2008). A second study reported the role of the homeodomain transcription factor BELLRINGER (*AtBLR*) in the establishment and maintenance of the phyllotaxis pattern in *A. thaliana* by the control of *AtPME5* expression. The study of the KO mutant *atblr-6* showed that *AtBLR* is required to establish normal phyllotaxis through the exclusion of *AtPME5* expression from the meristem; in contrast, phyllotaxis is maintained by the activation of *AtPME5* in the elongating stem (Peaucelle *et al.*, 2011b). Local accumulation of auxin in the shoot apex of *Arabidopsis* leads to local demethylesterification of HGs, suggesting a role for auxin in the control of PME activity, necessary for the decrease in tissue rigidity promoting organ formation. In an *AtPME3*-overexpressing line, which shows decreased demethylesterification of HGs, local accumulation of auxin did not induce organ formation, confirming that the control of the DM of HGs occurs downstream of auxin accumulation during organ formation (Braybrook and Peaucelle, 2013). The changes in HG structure underlie changes in cell wall rheology that are key elements of primordia emergence at the shoot apical meristem (Hamant *et al.*, 2011; Peaucelle *et al.*, 2011a). Pectins thus play a major role in controlling plant morphogenesis during development (Palin and Geitmann, 2012).

Although the parts played by other HGMEs in flower development have been less documented, some results suggest putative roles in this developmental process. For instance, a cDNA encoding a *PGIP* gene has been isolated from cotton flower. This gene, designated *GhPSI*, is specifically expressed in cotton petals and is gradually up-regulated over the course of petal development. However, its expression level declines rapidly in senesced petals after flowering, suggesting that the *GhPSI* gene may be involved in cotton petal development and senescence (Shi *et al.*, 2009).

#### *Roles of HGMEs in the modulation of the physical properties of the end-product*

Among other post-fecundation processes, the roles of pectin modifications during fleshy fruit development and maturation have been extensively studied (Brummell *et al.*, 2004;

Ericksson *et al.*, 2004; Louvet *et al.*, 2011; Lunn *et al.*, 2013; Terao *et al.*, 2013). Early reports showed that, during tomato ripening, PME activity regulates MeOH and ethanol (EtOH) accumulation in the pericarp (Frenkel, 1998). Recent results showed an additional role for PME in cellular calcium distribution and blossom-end rot development in tomato fruit (De Freitas *et al.*, 2012). The link between PME levels and fruit susceptibility to pathogens was reported in the strawberry–*Botrytis* interaction (Osorio *et al.*, 2008), and the consequences of changes in PME gene expression on metabolic and signalling pathways were described (Osorio *et al.*, 2011a, b). Overall, the fine control of PME activity could be related to the interaction with specific inhibitors during fruit development, as recently shown (Reca *et al.*, 2012). The role of PME-mediated changes in the pectic network is likely to be highly conserved during fruit development, as shown by the changes in PME activity and/or transcript accumulation in various plant species (Barnavon *et al.*, 2001; Deytieu-Belleau *et al.*, 2008; Draye and Cutsem, 2008; Mbéguié-A-Mbéguié *et al.*, 2009; Cação *et al.*, 2012; Roongsatham *et al.*, 2012; Wen *et al.*, 2013). For instance, in *Musa acuminata*, PMEs are involved in cell wall modifications, responsible for softening the pedicel abscission area after induction of ripening. However, the observed effects could also be caused by cooperation of PME with other cell wall-modifying genes. In tomato, PG activity appears necessary for HG modifications, following the first stage of ripening; in transgenic plants underexpressing a PG gene, the fruit does not show any degradation in the last stage of fruit ripening (Hadfield and Bennett, 1998). The close relationship between PG activities and fruit firmness has recently been shown in apple (Atkinson *et al.*, 2012) and in strawberry (García *et al.*, 2009; Quesada *et al.*, 2009). In addition, PGs have been reported to play a central role in the modifications of the HG network associated with fruit abscission (Swain *et al.*, 2011). In banana, as in other climacteric fruits, ripening is accompanied by a high production of ethylene, which suggests a strong regulation of HGMEs by this phytohormone (Dominguez-Puigjaner *et al.*, 1997; Mbéguié-A-Mbéguié *et al.*, 2009; Srivastava *et al.*, 2012). For instance, in *M. acuminata*, the *BAN17* gene encoding PL is expressed under the control of ethylene. The role of PL in fruit softening was functionally demonstrated in strawberry (Jimenez-Bermudez *et al.*, 2002; Santiago-Doménech *et al.*, 2008). The changes in the HG network and its methylesterification status, mediated by HGMEs, are thus key elements of fruit texture that could be used for quantitative and genetic association studies (Chapman *et al.*, 2012; Lahaye *et al.*, 2012). Other post-fecundation processes appear to be under the control of HG structure. For instance, HG modification was shown to play a role in fibre elongation in cotton. In *Gossypium hirsutum*, the *GhPEL* gene, whose product was biochemically characterized as a PL, is preferentially expressed in fibres at 10 d post-anthesis. In anti-sense *GhPEL* transgenic cotton plants, where the expression is significantly suppressed, a reduction in PL activity was observed. This reduction led to a decreased degradation of demethylesterified HG epitopes in the primary cell wall with

consequent effects on cell wall loosening; ultimately, the elongation of the fibre was repressed (Wang *et al.*, 2010).

Recent findings also highlight emerging roles for PME in wood development and wood mechanical properties. In *A. thaliana*, five different PME genes are expressed in the xylem, one of which is more highly expressed in this tissue than in any other examined. Similarly, transcripts of a dozen PME genes have been found in poplar wood-forming tissues (Geisler-Lee *et al.*, 2006) and shown to be tightly regulated within the cambial meristem and during xylogenesis. In addition, several genes encoding HGMEs have been found to be regulated during wood formation in *Pinus radiata* (Li *et al.*, 2011, 2012) and *Eucalyptus* (Carvalho *et al.*, 2008; Goulao *et al.*, 2011). A role for PMEs has been demonstrated in the regulation of fibre length in poplar (Siedlecka *et al.*, 2007) as well as in the modulation of stem mechanical properties in *Arabidopsis* (Hongo *et al.*, 2012). For the latter, AtPME35 is involved in the demethylesterification of the primary cell wall, which directly regulates the mechanical strength of the supporting tissue. Mutants affected in the expression of the AtPME35 gene showed a striking bending phenotype. Furthermore, the presence of HG and its de-esterification are likely to be essential for xylem lignification. For example, Ca<sup>2+</sup>-bridged de-esterified HG is known to bind class III peroxidases that might initiate lignin polymerization (Jenkins *et al.*, 2001). In support of this hypothesis, it was shown that PME, de-esterified HGs, peroxidase, and the start of the lignification process co-localize at cell junctions in woody tissues (Wi *et al.*, 2005) and that pectins interact with lignin monomers and affect lignin polymerization *in vitro* (Lairez *et al.*, 2005; Habrant *et al.*, 2009). The control of the methylesterification status of HGs has dramatic consequences on the chemical and rheological properties of the cell wall (Wolf *et al.*, 2009b), and thus is likely to affect the properties of biomass-derived by-products. For instance, the PME-mediated changes in pectin structure have been shown to influence the texture of cooked potato tubers (Ross *et al.*, 2011a, b), the saccharification of plant tissues for biomass bioconversion (Lionetti *et al.*, 2010), and the solid wood properties of *Eucalyptus pilularis* (Sexton *et al.*, 2012). The differences in pectins between juvenile and adult *Eucalyptus globulus* wood, revealed by various nuclear magnetic resonance (NMR) analyses, support a major role for these enzymes during wood formation (Rencoret *et al.*, 2011).

## Role of HG modifications in plant responses to biotic stress

### *Biotic stresses modify gene expression of HG-modifying enzymes*

Plant HGMEs are involved in various biotic interactions: necrotrophic, hemibiotrophic, or biotrophic pathogens (fungi, oomycetes, bacteria, and viruses); phytophagous organisms (piercing–sucking insects, chewing insects, and nematodes); endosymbiotic microorganisms (arbuscular mycorrhizal fungi and bacteria); or plant parasites that feed and then develop on host plants. Piercing–sucking insects

wound the plant by inserting their stylets (mouthpart) into tissue to suck the cell contents or sieves. This wounding stress may be mimicked using needle punctures on leaves. Plant HGME gene expression is modified by all of these stresses, but the observed patterns of expression are dependent upon the bioaggressor (Table 2). For instance, no induction of PG gene expression has yet been reported after chewing insect and plant parasite infestations. Similarly, no induction of PAE or PL gene expression was shown following infections by viruses, endosymbiotic microorganisms, and plant parasites. For each bioaggressor, the pattern of HGME expression differs depending on the plant species, ecotype (or cultivar), and plant phenology. While *A. thaliana* AtPME3 (At3g14310) was overexpressed in the C24 ecotype (Hewezi *et al.*, 2008) after 3 d of infestation with 250 J<sub>2</sub> nematodes (*Heterodera schachtii*), no such effect was reported in Col-0 for a similar infestation level. In contrast, in this ecotype, the expression of two other PMEs (At2g45220 and At1g53830) was down-regulated following infestation (Puthoff *et al.*, 2003). This difference could also be related to plant phenology as, although of similar age (14 d old for C24 and 12 d old for Col-0), the plants were grown under distinct photoperiods (16/8 h and 12/12 h day/night, respectively).

The analysis of available microarray data sets from *A. thaliana* Col-0 ecotype (www.genevestigator.com; Hruz *et al.*, 2008) for distinct pathogens shows that the expression of a number of HGME genes is mostly down-regulated whatever the type of biotic stress: necrotrophic (*Botrytis cinerea* and *Fusarium oxysporum*), hemibiotrophic (*Phytophthora infestans*), or biotrophic (*Pseudomonas syringae*) pathogens, phytophagous organisms (*Bemisia tabaci* and *Meloidogyne incognita*), virus (*Cabbage leaf curl virus* CalCuV), or wounding (needles, Fig. 5). In general, within the smallest multigenic families of HGMEs such as PAEs and PLs, up to 67% and 42% of genes, respectively, show distinct expression patterns in response to biotic stresses. In contrast, figures are in the range of 29% and 24% for PMEs and PGs, respectively. In more detail, when considering the number of regulated HGME genes and their expression levels (up- or down-regulation), two clusters can be distinguished. The first includes responses to *P. infestans*, *B. tabaci*, *B. cinerea*, CalCuV, and *P. syringae*, and shows an overall down-regulation of most of the genes. The second cluster, which has far fewer modifications of gene expression, comprises the responses to *M. incognita*, *F. oxysporum*, or wounding. Some genes are specifically expressed in each cluster (PL, At5g63180; PGs, At4g23820, At3g06770, and At3g16850; or PME, At1g53830; and PL, At5g55720 in cluster one and two, respectively). On this basis, it would be possible to distinguish between rather ubiquitous biotic stress-related HGMEs (PME, At2g45220; and PL, At3g07010) and specific ones (PMEs At5g04960–At3g10710 with *M. incognita*; PME At2g47280 and PGs At3g48950–At5g27530 with CalCuV, or PGs At1g70500–At3g57510 with *P. syringae*). As both clusters include a necrotrophic fungus (*B. cinerea* and *F. oxysporum*), the changes in the expression of HGMEs do not appear to be dependent on the bioaggressor lifestyle *per se*. The expression of HGME genes is more likely to depend on the bioaggressor species (Fig. 5). For instance, the number

**Table 2.** Gene expression variations of homogalacturonan-modifying enzymes after biotic stresses

Pectin methyl esterases, pectin acetyl esterases, polygalacturonases, and pectin lyase-like involved in plant–bioaggressor interactions.

Gene name	AGI or accession	Stress	Species name(s)	Induction	References
Pectin methyl esterases (PMEs)					
		Aphid	<i>Brevicoryne brassicae</i>	32 aphids/plant; 48 hpi	Kusnierczyk <i>et al.</i> (2008)
<i>AtPME17</i>	At2g45220			Up-regulated 48 hpi (×4.53)	
<i>AtPMEPCR1 (61)</i>	At5g53370			Up-regulated 6, 12, 24, 48 hpi (×1.35, 1.44, 1.83, 1.85)	
<i>Apium graveolens</i> cv. Dulce (celery)		Aphid	<i>Myzus persicae</i>	20 (3 dpi) or 100 aphids (7 dpi)/plant	Divol <i>et al.</i> (2005)
<i>AgPME</i>	CN254944			Up-regulated (×3.41, 2.41)	
<i>AgPME</i>	CN254453			Up-regulated (×3.48, 2.19)	
		Whitefly	<i>Bemisia tabaci</i>	100 whiteflies/plant; 21 dpi	Kempema <i>et al.</i> (2007)
<i>AtPME17</i>	At2g45220			Up-regulated (×8.06)	
<i>AtPMEPCR1 (18)</i>	At1g11580			Down-regulated (×−1.70)	
		Chewing insect	<i>Spodoptera littoralis</i> OS	1 mm holes punctured and 1 µl of insect OS applied; 6, 24 hpi	Consales <i>et al.</i> (2011)
<i>AtPME32</i>	At3g43270		(oral secretion)	Up-regulated (×2.17, 1.58)	
<i>AtPMEPCR1 (18)</i>	At1g11580			Up-regulated (×3.33, 1.77)	Little <i>et al.</i> (2007)
		Chewing insect	<i>Pieris brassicae</i>	10–40 eggs/plant; 24, 48, 72 hpi	
<i>AtPME44</i>	At4g33220			Down-regulated (× −1.54, −1.65, −1.39)	
<i>Nicotiana attenuata</i> (tobacco)		Chewing insect	<i>Manduca sexta</i> OS	Leaf wounded with a pattern wheel + 20 µl of diluted insect OS; 9 hpi	Von <i>et al.</i> (2006)
<i>NaPME</i>	DQ115979			Up-regulated (×2.59)	
		Nematode	<i>Meloidogyne javanica</i>	10–12 J <sub>2</sub> nematodes/root tip; 3 dpi	Barcala <i>et al.</i> (2010)
<i>AtPME</i>	At5g20860			Up-regulated (×3.05)	
<i>AtPME</i>	At1g11580			Up-regulated (×3.56)	
<i>AtPME17</i>	At2g45220			Down-regulated (× −3.54)	
		Nematode	<i>Heterodera schachtii</i>	250 J <sub>2</sub> nematodes/plant; 3, 8, 13 dpi	Hewezi <i>et al.</i> (2008)
<i>AtPME3</i>	At3g14310		or <i>Meloidogyne incognita</i>	Up-regulated (×2.0, 3.5, 3.0)	
<i>Solanum lycopersicum</i> (tomato)		Nematode	<i>Globodera rostochiensis</i>	10 000 J <sub>2</sub> nematodes/plant; 14 dpi	Uehara <i>et al.</i> (2007)
<i>LePME</i>	SNG-U213346			Up-regulated (×7.0)	
		Nematode	<i>Heterodera schachtii</i>	250 J <sub>2</sub> nematodes/plant; 3 dpi	Puthoff <i>et al.</i> (2003)
<i>AtPME2</i>	At1g53830			Down-regulated (× −3.8)	
<i>AtPME17</i>	At2g45220			Down-regulated (× −3.9)	
		Bacterium	<i>Pectobacterium carotovorum</i>	5 × 10 <sup>7</sup> cfu ml <sup>−1</sup> ; 14 hpi	Raiola <i>et al.</i> (2011)
<i>AtPME3</i>	At3g14310			Up-regulated (×2.5)	
		Fungus	<i>Botrytis cinerea</i>	5 × 10 <sup>5</sup> conidia ml <sup>−1</sup> ; 72 hpi	Raiola <i>et al.</i> (2011)
<i>AtPME3</i>	At3g14310			Up-regulated (×7.0)	
		Fungus	<i>Alternaria brassicola</i>	10, 24, 48 hpi	Narusaka <i>et al.</i> (2005)
<i>AtPME3</i>	At3g14310			Up-regulated (×7.2, 2.9, 4.1)	
		Fungus	<i>Alternaria alternata</i>	10, 24, 48 hpi	Narusaka <i>et al.</i> (2005)
<i>AtPME3</i>	At3g14310			Up-regulated (×11.2, 11.5, 6.8)	
<i>Linum usitatissimum</i> (flax)		Fungus	<i>Fusarium oxysporum</i> or <i>F. culmorum</i>	2 dpi (RT–PCR)	Wojtasik <i>et al.</i> (2011)
<i>LuPME3</i>	AF188895			Down-regulated	
<i>LuPME5</i>	AF355057			Down-regulated	
<i>Medicago truncatula</i> (barrel medic)		AM fungus	<i>Glomus mosseae</i>	28 dpi	Hohnjec <i>et al.</i> (2005)
<i>MtPME</i>	TC78420		or <i>Glomus intraradices</i>	Up-regulated (×2.56 GM, ×2.66 GI)	

Table 2. Continued

Gene name	AGI or accession	Stress	Species name(s)	Induction	References
<i>MtPME</i>	TC82059			Up-regulated ( $\times 2.33$ GM, $\times 4.11$ GI)	
<i>Sesbania rostrata</i> ( <i>Sesbania</i> )		Bacterium	<i>Azorhizobium caulinodans</i>	48 hpi	Lievens et al. (2001)
<i>SrPME1</i>	Srdd18			Up-regulated (RT-PCR)	Lievens et al. (2002)
<i>Solanum tuberosum</i> cv Igor (potato)		Virus	PVY <sup>NTN</sup>	0.5 hpi	Baebler et al. (2009)
<i>StPME</i>	STMHY50			Down-regulated ( $\times -1.27$ )	
		Virus	TuMV	5 dpi	Yang et al. (2007)
<i>AtPME3</i>	At3g14310			Down-regulated ( $\times -2.64$ )	
		Virus	CaLCuV	12 dpi	Ascencio-Ibanez et al. (2008)
<i>AtPMEPCRA (18)</i>	At1g11580			Down-regulated ( $\times -1.14$ )	
		Virus		4 dpi	Whitham et al. (2003)
<i>AtPMEPCRA (18)</i>	At1g11580		CMV, ORMV, TuMV, PVX, TVCV	Up-regulated ( $\times 2.2-3.8$ )	
<i>Vigna unguiculata</i> (cowpea)		Parasitic plant	<i>Striga gesnerioides</i>	6 dpi or 13 dpi	Huang et al. (2012)
<i>VuPME</i>	33686210			Up-regulated 6 dpi ( $\times 3.24$ ) and 13 dpi ( $\times 3.29$ )	
Pectin acetylsterases (PAEs)					
		Whitefly	<i>Bemisia tabaci</i>	100 silverfly/plant, 21 dpi	Kempema et al. (2007)
<i>AtPAE</i>	At4g19420			Up-regulated ( $\times 2.89$ )	
<i>AtPAE</i>	At5g45280			Down-regulated ( $\times -1.78$ )	
		Aphid	<i>Myzus persicae</i>	20 (3 dpi) or 100 aphids (7 dpi) /plant	Divol et al. (2005)
<i>AtPAE</i>	CN254169			Up-regulated ( $\times 3.78$ , 2.16)	
<i>Malus domestica</i> (apple tree)		Aphid	<i>Dysaphis plantaginea</i>	20 aphids/leaf, 72 hpi	Qubbaj et al. (2005)
<i>MdPAE</i>	CB035291			Up-regulated	
		Chewing insect	<i>Spodoptera littoralis</i> OS	1 mm holes punctured and 1 $\mu$ l of insect OS applied; 6, 24 hpi	Consales et al. (2011)
<i>AtPAE</i>	At2g46930			Up-regulated ( $\times 2.83$ , 2.30)	
		Nematode	<i>Meloidogyne incognita</i> or <i>Heterodera schachtii</i>	1, 2, 3, 5, 7 dpi	Vercauteren et al. (2002)
<i>AtPAE</i>	AY050847			Up-regulated	
Polygalacturonases (PGs)					
		Aphid	<i>Brevicoryne brassicae</i>	32 aphids/plant	Kusnierczyk et al. (2008)
<i>AtPG</i>	At5g49215			Down-regulated 6, 12, 24, 48 hpi ( $\times -1.21$ , $-1.46$ , $-1.38$ , $-1.52$ )	
<i>AtPG</i>	At3g62110			Down-regulated 6, 12, 24, 48 hpi ( $\times -1.53$ , $-1.43$ , $-1.56$ , $-1.55$ )	
<i>AtPG</i>	At1g60590			Down-regulated 6, 12, 24, 48 hpi ( $\times -1.33$ , $-1.45$ , $-1.39$ , $-1.51$ )	
<i>AtPG</i>	At4g23820			Down-regulated 6, 12, 24, 48 hpi ( $\times -1.14$ , $-1.27$ , $-1.51$ , $-1.62$ )	
<i>AtPG</i>	At3g06770			Down-regulated 6, 12, 24, 48 hpi ( $\times -1.48$ , $-1.83$ , $-1.92$ , $-1.83$ )	
<i>AtPG</i>	At1g10640			Down-regulated 6, 24, 48 hpi ( $\times -1.78$ , $-1.41$ , $-1.88$ )	
		Aphid	<i>Myzus persicae</i> saliva infiltration	50 aphids/plant, 24 (OS), 48, 72 (OS+feeding) hpi	De Vos and Jander (2009)
<i>AtPG</i>	At1g60590			Down-regulated ( $\times -2.69$ , $-22.7$ )	
		Nematode	<i>Heterodera schachtii</i>	250 J <sub>2</sub> nematodes/plant, 3 dpi	Puthoff et al. (2003)




Table 2. Continued

Gene name	AGI or accession	Stress	Species name(s)	Induction	References
<i>AtPG</i>	At2g41850			Up-regulated (×3.7)	
<i>AtPG</i>	At1g05660	Nematode	<i>Meloidogyne javanica</i>	Down-regulated (×3.1) 10–12 J <sub>2</sub> nematodes/root tip; 3 dpi	Barcala <i>et al.</i> (2010)
<i>AtPG</i>	At4g23820	AM fungus	<i>Glomus mosseae</i>	Up-regulated (×3.03) 28 dpi	Hohnjec <i>et al.</i> (2005)
<i>Medicago truncatula</i> (barrel medic)			or <i>Glomus intraradices</i>		
<i>MtPG</i>	TC88957			Up-regulated (×2.53 GM, ×4.50 GI)	
<i>Medicago truncatula</i> (barrel medic)		Bacterium	<i>Sinorhizobium meliloti</i>	24, 48 dpi	Lohar <i>et al.</i> (2006)
<i>MtPG</i>	TC78631			Up-regulated (×1.14, 1.41)	
<i>MtPG</i>	TC89800			Up-regulated (×1.45, 1.25)	
<i>MtPG</i>	TC91368			Up-regulated (×1.14, 1.12)	
<i>MtPG</i>	TC87651			Up-regulated (×2.88, 2.23)	
<i>Medicago sativa</i> (alfafa)		Bacterium	<i>Sinorhizobium meliloti</i>	24 dpi	Munoz <i>et al.</i> (1998)
<i>MsPG</i>	MsPG3			Up-regulated (RT-PCR)	
<i>Solanum tuberosum</i> cv Igor (potato)		Virus	PVY <sup>NTN</sup>	0.5 hpi	Baebler <i>et al.</i> (2009)
<i>StPG</i>	STMIS28			Down-regulated (× -1.43)	
Pectate lyase-like proteins (PLLs)		Aphid	<i>Brevicoryne brassicae</i>	32 aphids/plant, 6, 12, 24, 48 hpi	Kusnierczyk <i>et al.</i> (2008)
<i>AtPL</i>	At1g67750			Down-regulated (× -2.00, -1.62, -1.64, -1.85)	
<i>Glycine max</i> (soybean)		Aphid	<i>Aphis glycines</i>	40 aphids/plant, 6 hpi	Li <i>et al.</i> (2008)
<i>GmPL</i>	At748551 (At1g67750)			Down-regulated (× -1.52)	
<i>GmPL</i>	AW099533			Down-regulated (× -1.54)	
<i>GmPL</i>	AW309146 (At1g11920)			Down-regulated (× -1.62)	
		Chewing insect	<i>Spodoptera littoralis</i> OS	1 mm holes punctured and 1 µl of insect OS applied; 6, 24 hpi	Consales <i>et al.</i> (2011)
<i>AtPNL</i>	At3g15720			Up-regulated (×5.28, 2.0)	
<i>AtPNL</i>	At3g61490	Chewing insect	<i>Pieris brassicae</i>	Up-regulated (×12.13, 2.0) Down-regulated (× -1.48, -1.45, -1.23)	Little <i>et al.</i> (2007)
<i>AtPNL</i>	At5g48900			10–40 eggs/plant, 24, 48, 72 hpi	
		Nematode	<i>Meloidogyne javanica</i>	10–12 J <sub>2</sub> nematodes/root tip; 3 dpi	Barcala <i>et al.</i> (2010)
<i>AtPL</i>	At3g53190	AM fungus	<i>Glomus mosseae</i>	Up-regulated (×4.88) 28 dpi	Hohnjec <i>et al.</i> (2005)
<i>Medicago truncatula</i> (barrel medic)			or <i>Glomus intraradices</i>		
<i>MtPL</i>	TC88957			Up-regulated (×2.03 GM, ×3.00 GI)	
<i>Medicago truncatula</i> (barrel medic)		Bacterium	<i>Rhizobium meliloti</i>	6, 12, 24, 48 dpi	Lohar <i>et al.</i> (2006)
<i>MtPL</i>	TC89125			Up-regulated (×5.12, 2.37, 2.73, 1.18)	
<i>Vigna unguiculata</i> (cowpea)		Parasitic plant	<i>Striga gesnerioides</i>	13 dpi	Huang <i>et al.</i> (2012)
<i>VuPL</i>	33691436			Down-regulated (× -8.86)	

Piercing-sucking insects (yellow box), chewing insects (turquoise), nematodes (blue), bacteria (black), fungi (light grey), arbuscular mycorrhizal fungus (dark grey), viruses (red), and parasitic flowering plants (green).

Species names in bold indicate necrotrophic pathogens and underlined gene names refer to characterized enzymes.



**Fig. 5.** Expression pattern of PME, PAE, PG and PLL mRNA in *A. thaliana* Col-0 in response to necrotrophic fungi (48h, *Botrytis cinerea*, *Fusarium oxysporum*), hemibiotrophic oomycete (24h, *Phytophthora infestans*), biotrophic bacterium (24h, *Pseudomonas syringae*), herbivore insects (21 d, *Bemisia tabaci*), nematode (28 d, *Meloidogyne incognita*), virus (24h, *Cabbage leaf curl virus* CalCuV), and leaf wounding with needles (24h). PMEs, PAEs, PGs, and PLLs (red, green, orange, and blue circles) are shown together. Data were analysed using the Genevestigator

of HGME genes regulated following a 48 h infestation by *B. cinerea* was 3-fold higher than for an *F. oxysporum* infestation. This suggests distinct response pathways at the cell wall level. Interestingly, whatever the HGME family considered, *F. oxysporum* induces the fewest changes in gene expression levels. While it has been suggested that necrotrophs and hemibiotrophs can modulate the expression of a higher number of PMEs than biotrophs (Lionetti et al., 2012), it seems important to consider the pathogen species and probably the duration of infection. For example, after 48 h of infestation, HGME modifications induced by the two necrotrophic fungi, *F. oxysporum* and *B. cinerea*, belong to the two different clusters defined before, while, within the same cluster, most HGME genes are more extensively modified after 24 h of infection by the biotrophic bacterium *P. syringae* than by the necrotrophic fungus *B. cinerea* (29, 24, and 42% against 23, 16, and 35% of PME, PG, and PL, respectively). Only PAEs are recruited at the same level in both infections. Changes in the expression of HGME genes were observed for the duration of an interaction that can reach several weeks in the case of piercing-sucking insects and nematodes; the time needed to complete development steps or even a whole reproduction cycle. After 21 d of continuous feeding in a single phloem sieve element by the whitefly nymph *B. tabaci*, or after 28 d of infestation by the nematode *M. incognita*, the number of genes with modified expression was almost the same as after *B. cinerea* infection. Comparing the two piercing-sucking phytophagous insects, each belonging to the two different clusters, the overall number of PMEs and PLs differentially expressed is relatively similar, but differences lie in the identity of the isoforms. The changes in gene expression are likely to concern isoforms that are organ specific; nematodes feeding on roots and whiteflies on leaves. Interestingly, 24 h after leaf wounding, the HGME pattern belongs to the same cluster as that of nematode infestation. Current challenges include the identification of the specific roles of HGME isoforms in biotic interactions (host species, bioaggressor lifestyle, plant phenology, attacked organ, and duration of infestation/infection). This might help to build general and specific models of the response of the plant at the cell wall (pectin) level.

#### Role of HGMEs in the establishment of feeding structures during biotic interactions

Except for chewing insects and necrotrophic pathogens, in most cases a feeding connection is established during plant-bioaggressor interactions. Most hemibiotrophic and obligate biotrophic fungi, or oomycetes use an appressorium followed by a penetration peg that goes through the plant cell wall to develop a haustorium for host cell nutrient absorption. Nutrient trafficking passes through the plant extrahaustorial membrane and matrix to the hyphal wall and membrane

Meta-Analyzer Tools ([www.genevestigator.com/gv/](http://www.genevestigator.com/gv/); Hruz et al., (2008).

Only probes with a single gene and genes showing a minimal expression were used for the analysis. For each HGME family, the percentage (%) of genes expressed among all family members is indicated.

(Szabo and Bushnell, 2001; Underwood, 2012), while the whole haustorium is encased. Haustoria are also described as special organs of most parasitic plants and arbuscular mycorrhizal fungi for penetration and development of vascular connections within the host plant for feeding (Leake, 1994; Lee, 2007). Cell wall remodelling, and thus cell wall-degrading enzymes (CWDEs), including HGMEs, are necessary to establish a conductive system (Deising *et al.*, 1995). CWDEs are secreted by pathogens, plant parasites, or mycorrhizal fungi to develop the haustorium and by the plant to restrict its development, especially through the cell wall apposition (papilla). Although callose and lignin have been described as the main embedded compounds for cell wall strengthening in the vicinity of the haustorium, pectate gel generated following PME activity could also play a role (Micheli, 2001) (Fig. 4). In fact, two PME genes are overexpressed in *Medicago truncatula* roots infected by the endosymbiotic fungus *Glomus* sp. (Hohnjec *et al.*, 2005). In cowpea (*Vigna unguiculata*) infected by the holoparasite *Striga gesnerioides*, in parallel with the overexpression of one PME gene, one PLL is underexpressed in roots (Huang *et al.*, 2012). Although to date no specific role has been demonstrated for plant HGMEs during haustorium differentiation, in wheat leaves infected by the pathogen *Blumeria graminis*, papilla size correlated with peroxidase activity and H<sub>2</sub>O<sub>2</sub> accumulation, depending on the DA of plant pectin fragments used as elicitors (Randoux *et al.*, 2010).

HGMEs are also good candidates for modulating cell wall structure during intrusive cell growth that occurs when plants interact with biotrophic, hemibiotrophic, and arbuscular mycorrhizal fungi (Perfect and Green, 2001) and with parasitic plants (Leake, 1994). HGMEs could also modulate the interaction with aphids and nematodes (Wyss and Zunke, 1986; Tjallingii and Esch, 1993). For instance, the trophic behaviour of the aphid *Myzus persicae* was modified within 8 h of infestation in *A. thaliana* plants knocked out for the expression of the *AtPME17* gene (C. Wattier *et al.*, unpublished). Interestingly, the expression of *PME17* is markedly increased following interactions with other aphids (*Brevicoryne brassicae*) or whitefly (*B. tabaci*) (Fig. 5) (Kempema *et al.*, 2007; Kuśnierczyk *et al.*, 2008). As the salivation phases were longer in the *pme17* mutant, *PME17* could have a role in facilitating the progression of the stylets in wild-type plants. Similarly, very early in root knot and cyst nematode infections (*M. incognita* and *Heterodera schachtii*, respectively), an up-regulation of one *AtPAE* was shown using *in situ* localization (Vercauteren *et al.*, 2002). The activity of this HGME could be involved in softening the plant cell wall to establish a feeding site, which is an induced multinucleate and physiologically active aggregation of fused root cells that exclusively provides nutrients during the nematode sedentary life (Szabo and Bushnell, 2001; Vercauteren *et al.*, 2002). Similarly, *AtPME3* seems to be crucial in the early phase of the establishment of the syncytium, the feeding site of the cyst nematode, as it appeared to be a virulent target in the *A. thaliana*–*H. schachtii* interaction (Hewezi *et al.*, 2008). Using a yeast two-hybrid screen, *PME3*, whose gene expression was increased during infestation, was identified as

interacting with a cellulose-binding protein (CBP) secreted by *H. schachtii* (Hewezi *et al.*, 2008). Results obtained using transgenic lines suggest that *PME3* could be recruited by CBP to modify the plant cell wall, thus helping cyst nematode parasitism. However, this is unlikely to be the whole mechanism for parasitism success. Similarly, a so-called ‘welcoming programme’, characterized by huge cell reorganizations and by the formation of root hairs, has been related to the facilitation of the intercellular progression of infection threads or hyphae during the initiation of endosymbiotic interactions (van Brussel *et al.*, 1992; Bonfante, 2001; Hause and Fester, 2004). For instance, during legume–*Rhizobium* or arbuscular mycorrhizal fungus symbioses, several plant HGMEs were involved in the early stages of the interaction (Lohar *et al.*, 2006; Oldroyd *et al.*, 2011). Four PG genes and one PLL gene were overexpressed in *M. truncatula*–*Rhizobium meliloti* interactions, while in *Medicago sativa* one PG was specifically localized in the cell wall of nodule primordia and invasion zones from 1 h to 2 d post-inoculation (Muñoz *et al.*, 1998; Lohar *et al.*, 2006). Using mutant plants, a PL from *L. japonicus* appears essential for the proper initiation of *Rhizobium trifolii* infection (Xie *et al.*, 2012). Moreover, *in situ* localization showed that PMEs could contribute to the development of new vascular tissues during rhizobial infection (Lievens *et al.*, 2002). This is consistent with the analysis of microarray data sets, showing that *M. truncatula* PGs, PLLs, and PMEs are overexpressed 28 d after inoculation with *Glomus* sp. (Hohnjec *et al.*, 2005) and that PG activity is localized in the cell wall of lateral root primordia (Peretto *et al.*, 1995). The specific case of parasitic plants is of interest as, like host plants, they secrete their own HGMEs to facilitate their penetration into the host plant root cortex. For instance, *Orobanche* sp. secretes PMEs and their activity is correlated with the localization of low esterified HG in the cell wall and the middle lamella at the site of contact between host and parasitic cells (Ben-Hod *et al.*, 1993; Losner-Goshen *et al.*, 1998). This plant–plant interaction raises fascinating questions concerning the roles and specificity of host and parasite HGMEs. In particular, the study of the potential inhibition of parasitic plant PMEs by host plant PMEIs would be of great interest.

Host plant HGMEs are thus largely involved during the establishment of the feeding structures of bioaggressors, both to facilitate their intrusion and to restrict their excessive spreading. Their role in the fine-tuning of cell wall remodelling in favour of parasitic success at an early stage suggests that HGME activity has been diverted during a co-evolution process. Virulence factors of bioaggressors that target plant protein involved in defence responses, and the binding of exogenous protein to some plant PMEs, appear as examples. Reported for plant–nematode interactions, this targeting has previously been highlighted for plant–virus interactions. The movement protein (MP) of *Tobacco mosaic virus* (TMV), which can be transmitted by piercing–sucking insects, interacts with a PME purified from tobacco leaves and this interaction is required for TMV cell to cell movement in the host plant (Dorokhov *et al.*, 1999; Chen *et al.*, 2000). The reduction of total PME activity, using antisense suppression of the

expression of one PME or the overexpression of a characterized PME1, led to the delayed systemic movement of TMV (Chen and Citovsky, 2003). In the meantime, using transgenic plants overexpressing PME, an inverse correlation between PME activity and TMV lesion sizes has been demonstrated (Gasanova *et al.*, 2008). In this respect, the common hypothesis that plant HGMEs play a role in plant resistance by mediating cell wall strengthening or producing endogenous elicitors is probably too simplistic.

#### *Roles of HGMEs in structural resistance*

The cell wall represents an impenetrable physical barrier with constitutive rigidity to fend off bioaggressor attacks (Vallarino and Osorio, 2012). A high DM of HG has been correlated to genotype resistance to the aphid *Schizaphis graminum*, the biotrophic fungus *Colletotrichum lindemuthianum*, and the necrotrophic bacterium *Ralstonia solanacearum* (Table 3) (Dreyer and Campbell, 1984; Boudart *et al.*, 1998; Wydra and Beri, 2006). This could be related to changes in cell wall elasticity and mechanics. As shown during organ initiation (Peaucelle *et al.*, 2011a), the lowest wall elasticity is correlated to the highest DM. Furthermore, a high constitutive DM of pectin was structurally related to the borate-RGII cross-link in regulating cell wall stiffness (Ishii and Matsunaga, 2001). As such, plant resistance to the necrotrophic bacterium *Pectobacterium carotovorum* is associated with a high DM of pectin in wild potato plants (McMillan *et al.*, 1993; Marty *et al.*, 1997) and in the *pme3* mutant of *A. thaliana* (Raiola *et al.*, 2011). The enhanced DM level obtained by mutation of *PME* genes or by overexpression of *PMEI* genes *in planta* increased the resistance of dicotyledonous and monocotyledonous species to biotrophic or necrotrophic pathogens, but not to the full range of pathogens for each plant (Table 3) (Lionetti *et al.*, 2007; An *et al.*, 2008; Raiola *et al.*, 2011; Volpi *et al.*, 2011). Furthermore, *in vitro*, the hydrolysis of plant pectin by endo-PGs from several necrotrophic pathogens is decreased when pectins of high DM are used as carbon sources (Bonnin *et al.*, 2002). The resistance observed could therefore be related to a decrease in the number of specific substrates for endogenous PGs, as well as the physical or chemical properties of pectins, such as the isoform pattern of HGME activity or the amount of branched pectins. Using antisense tomato plants, the decrease in plant PG activity reduced, as expected, fruit softening and ripening, but also increased tomato resistance against biotrophic or necrotrophic pathogens (Damasceno *et al.*, 2011). Nevertheless, enhancing resistance through the modulation of the DM of HG is unlikely to be an easy strategy as it appears to be pathogen specific. For instance, increased PME activity and the associated lower DM of pectins, in transgenic strawberry overexpressing one PME, enhanced fruit resistance to the necrotrophic fungus *B. cinerea* (Osorio *et al.*, 2008). Understanding this opposite effect is complex as PME activity on HG can have two distinct consequences. On one hand, the linear activity of PME can give rise to blocks of free carboxyl groups that non-covalently interact with  $\text{Ca}^{2+}$  ions, conferring a gel-like structure and cell wall strengthening

(Morris *et al.*, 1982; Micheli, 2001). On the other hand, random PME activity promotes the action of pectin depolymerases (endo-PG or lyase activities) increasing both cell wall loosening and porosity and producing OGs that elicit plant defence responses (Baron-Epel *et al.*, 1988; Ehwald *et al.*, 1992). Thus, in transgenic plants with modified HGME activity, either the direct or the indirect effect on HG structure could play a role in plant resistance. For example, constitutive gene expression of antimicrobial proteins (*PR5*) in transgenic strawberry overexpressing one PME enhanced basal resistance to *B. cinerea*, while a decrease in their level (*PR1* and *PR10*) in *PMEII*-silenced pepper conferred decreased basal resistance to the biotrophic bacterium *Xanthomonas campestris* (An *et al.*, 2008; Osorio *et al.*, 2008). Among PMEs, some (At1g11580) that have a ribosome-inactivating protein (RIP) activity might be considered antimicrobial proteins themselves. RIPs are known to be involved in plant defence against viruses (De-la-Peña *et al.*, 2008). Resistance to the fungus *Puccinia graminis* was associated with a random distribution of the methylesters of HGs in the near-isogenic resistant line as compared with a more blockwise distribution in the susceptible cultivar (Wiethölter *et al.*, 2003). The DA of pectins is also likely to play an important role in plant resistance. While the *Arabidopsis* mutant *reduced cell wall acetylation rwa2* was more resistant against the necrotrophic fungus *B. cinerea*, it was susceptible to the biotrophic fungus *Golovinomyces cichoracearum*. The ‘antagonistic responses’ to these pathogens are consistent with the two distinct plant defence pathways induced [jasmonic acid (JA)/ethylene (ET), versus salicylic acid (SA)]. All these results suggest an indirect link between cell wall-related basal structural resistance and inducible plant defences. The mutation of the plant putative pectate lyase *PMR6* (*POWDERY MILDEW RESISTANT 6*), required for the virulence of *Erysiphe* sp., increased the content and DM of pectin as well as plant resistance, but surprisingly did not change either fungus penetration success or SA- and JA/ET-dependent defence responses. Similar results were obtained using the *pmr5* mutant; *PMR5* encodes a protein of unknown function required for pectin production and is likely to be targeted to the endoplasmic reticulum/secretory pathway. These results suggest non-elicitor OG production, highlighting the dual role of HGMEs in the control of cell wall stiffness during plant bioaggressor interactions. Finally, since HGME activities result in direct (strengthening) or indirect (OG elicitor production) resistance, they appear good candidates for virulent factor targeting.

#### *HGMEs are involved in induced resistance against biotic stresses*

The role of HGMEs in the production of endogenous elicitors during plant–bioaggressor interactions has been indirectly shown through the induction of all the main known plant defence responses following application of purified OGs, end-products of HGMEs. Indeed, within early signalling events, OGs induce  $\text{H}^+$  and  $\text{Ca}^{2+}$  influx,  $\text{K}^+$  efflux, membrane depolarization, extracellular medium alkalization (Mathieu *et al.*, 1991), protein phosphorylation/dephosphorylation,

**Table 3.** Biochemical implication of HG-modifying enzymes (PMEs, PAEs, and PLLs) and their inhibitor proteins (PMEIs, PGIPs, and PNLIP) in plant resistance against bioaggressors

Gene name	Utilization	Stress	Induction	References
Pectin methylation and PME activity <i>Sorghum bicolor</i> (sorgho) DM of pectin		Aphid <i>Schzaphis graminum</i>	Resistant variety has higher methylated pectins than the susceptible	Dreyer and Campbell (1984)
<i>Phaseolus vulgaris</i> (bean)		Fungus <i>Collectotrichum lindemuthianum</i>	Resistant line has higher methylated pectins than the susceptible	Boudart <i>et al.</i> (1998)
DM of pectin <i>Solanum tuberosum</i> cv Bintje or ADG (potato) DM of pectin	Near-isogenic lines	Bacterium <b><i>Pectobacterium carotovorum</i></b>	Resistant genotype (ADG) has higher methylated pectins than the susceptible genotype (Bintje)	Marty <i>et al.</i> (1997)
Potato DM of pectin	Somatic hybrid of 3 cv (Record, Estima, Katahdin)	Bacterium <b><i>Pectobacterium carotovorum</i></b>	Resistant genotype has higher methylated pectins than the susceptible genotype	McMillan <i>et al.</i> (1993)
<i>Solanum lycopersicum</i> (tomato) DM of pectin		Bacterium <b><i>Ralstonia solanacearum</i></b>	Resistant genotype (Hawaii7996) has higher methylated pectins than the susceptible genotype (Wva700)	Wydra and Beri (2006)
<i>Nicotiana attenuata</i> (tobacco) PME activity	(Hawaii7996, Wva700)	Chewing insect <i>Manduca sexta</i> OS	Leaf wounded with a pattern wheel; 20 µl of diluted insect OS applied; PME activity increased (29%) 30 min after OS applied	Von Dahl <i>et al.</i> (2006)
<i>Nicotiana tabacum</i> (tobacco) PME	Wild type	Virus TMV	PME specifically recognized the TMV MP (movement protein)	Dorokhov <i>et al.</i> (1999)
<i>Nicotiana tabacum</i> cv. Turk (tobacco) PME		Virus TMV	Mutant TMV without MP proteins cannot link to tobacco PME (no lesions on leaves after TMV mutant infection)	Chen <i>et al.</i> (2000)
<i>Nicotiana tabacum</i> cv. Samsun (tobacco)		Virus TMV	PME activity increased >resistance increased (size of leaf necrosis and short- and long-distance transport decreased)	Gasanova <i>et al.</i> (2008)
PME activity	ProPME			
<i>Nicotiana tabacum</i> cv. Turk (tobacco)		Virus TMV	PME activity decreased >symptome appearance delayed (5–12 times slower in the antisense line than in the wild type)	Chen and Citovsky (2003)
PME activity	Antisense suppression <i>pme3</i> KO	Fungus <b><i>Botrytis cinerea</i></b>	PME activity decreased >DM decreased >resistance decreased	Raiola <i>et al.</i> (2011)
<i>AtPME3</i> <i>AtPME3</i>	At3g14310 <i>pme3</i> KO At3g14310	Bacterium <b><i>Pectobacterium carotovorum</i></b>	PME activity decreased >DM decreased >resistance decreased	Raiola <i>et al.</i> (2011)
<i>Fragaria vesca</i> (wild strawberry)		Fungus <b><i>Botrytis cinerea</i></b>	OGA with low DM >resistance increased	Osorio <i>et al.</i> (2008)

Table 3. Continued

Gene name	Utilization	Stress	Induction	References
PME activity	Overexpression line (FaPE1)			
Pectin acetylation	Mutant <i>Atrwa2</i>	Fungus <i>Botrytis cinerea</i>	Pectin acetylation decreased >resistance increased	Manabe et al. (2011)
DA of pectin	(with 20% decreased acetyléster content)			
<i>Triticum aestivum</i> (wheat)	Chemical acetylation	Fungus <i>Blumeria graminis</i>	OGA with high DA >resistance increased	Randoux et al. (2010)
DA of pectin				
Pectate lyase-like (PLL)		Fungus <i>Erysiphe cichoracearum</i>	10 <sup>8</sup> cfu ml <sup>-1</sup> ; 1, 2, 4 dpi confers resistance to <i>E. cichoracearum</i>	Vogel et al. (2002)
<i>AtPME6</i>	At3g54920			
Pectin methyl esterase inhibitors (PMEIs)	overexpression lines	Fungus <i>Botrytis cinerea</i>	ATPMEI increased >PME activity decreased >DM increased >resistance increased	Lionetti et al. (2007)
<i>AtPME1</i>	At1g48020			
<i>ATPME2</i>	At3g17220			
<i>Capsicum annuum</i> (pepper)	Transgenic pepper silences CaPME1	Bacterium <i>Xanthomonas campestris</i> pv. <i>vesicatoria</i>	CaPMEI inhibited >susceptibility increased	An et al. (2008)
<i>CaPME1</i>				
<i>Capsicum annuum</i> (pepper)	Transgenic <i>A. thaliana</i> overexpresses CaPME1	Bacterium <i>Pseudomonas syringae</i> pv. <i>tomato</i>	CaPMEI overexpressed >resistance increased, but no resistance to the biotrophic fungus <i>Hyaloperonospora parasitica</i>	An et al. (2008)
<i>CaPME1</i>				
<i>Actinidia chinensis</i> (kiwi)	Transgenic wheat expresses AcPMEI	Fungus <i>Fusarium graminearum</i>	AcPMEI expression >PME activity decreased >DM increased >resistance increased	Volpi et al. (2011)
<i>AcPMEI</i>				
<i>Actinidia chinensis</i> (Kiwi)	Transgenic wheat expresses AcPMEI	Fungus <i>Bipolaris sorokiniana</i>	AcPMEI expression >PME activity decreased >DM increased >resistance increased	Volpi et al. (2011)
<i>AcPMEI</i>				
Polygalacturonase inhibitor proteins (PGIPs)		Fungus <i>Botrytis cinerea</i>	AtPGIP1 increased >resistance increased	Ferrari et al. (2006)
AtPGIP1	Antisense suppression			
<i>Brassica rapa</i> (Chinese cabbage)		Bacterium <i>Pectobacterium carotovorum</i>	BrPGIP2 overexpressed >resistance increased	Hwang et al. 2010
PGIP	Overexpression lines	Fungus <i>Botrytis cinerea</i>	pPGIP increased >resistance increased ( <i>B. cinerea</i> endo-PGs inhibited)	Powell et al. (2000)
<i>Solanum lycopersicum</i> (tomato)				
pPGIP	Expression of a pear PGIP	Fungus <i>Botrytis cinerea</i>	VvPGIP1 increased >resistance increased (BcPG1 inhibited)	Joubert et al. (2006)
<i>Vitis vinifera</i> (grape)				


Table 3. Continued

Gene name	Utilization	Stress	Induction	References
VvPGIP1 <i>Vitis vinifera</i> (grape)	Overexpression lines	Fungus <b><i>Botrytis cinerea</i></b>	pPGIP increased >resistance increased	Agüero <i>et al.</i> (2005)
pPGIP	Expression of a pear PGIP	Fungus <b><i>Botrytis cinerea</i></b>	AtPGIPs increased >resistance increased	Ferrari <i>et al.</i> (2003)
AtPGIP1, AtPGIP2 <i>Phaseolus vulgaris</i> (bean)	Overexpression lines	Fungus <b><i>Botrytis cinerea</i></b>	PvPGIP2 increased >resistance increased (BcPG1 inhibited)	Manfredini <i>et al.</i> (2005)
PvPGIP2 <i>Phaseolus vulgaris</i> (bean)	Overexpression lines	Fungus <b><i>Fusarium moniliforme</i></b>	PvPGIP2 inhibits FmPG	Federeci <i>et al.</i> (2001)
PvPGIP2 <i>Phaseolus vulgaris</i> (bean)		Fungus <i>Aspergillus niger</i>	PvPGIP inhibits AnPG (endoPGII)	King <i>et al.</i> (2002)
PvPGIP <i>Phaseolus vulgaris</i> (bean)		Fungus <b><i>Botrytis cinerea</i></b>	PvPGIP2 increased >resistance increased (BcPG1 inhibited)	Sicilia <i>et al.</i> (2005)
PvPGIP2 Pectin lyase inhibitor protein (PNLIP) <i>Beta vulgaris</i> (sugar beet)		Fungus <b><i>Rhizoctonia solani</i></b>	Barley-grain inoculum applied for 2 weeks at 25 °C	Bugbee (1993)
PNLIP	PNLIP		PNLIP activity higher in rotted tissues than in healthy	

Piercing–sucking insects (yellow box), chewing insects (turquoise), bacteria (black), fungi (light grey), and viruses (red). Species names in bold indicate necrotrophic pathogens.

mitogen-activated protein kinase (MAPK) cascades, GTP-binding protein, and reactive oxygen species (ROS) production ( $H_2O_2$ ,  $O_2^-$ ) (Shibuya and Minami, 2001; Vallarino and Osorio, 2012). They also induce the expression of defence genes encoding proteins involved in (i) defence protein accumulation such as protease inhibitors, pathogenesis-related proteins (PRs), or PGIPs; (ii) SA and JA/ET biosynthesis and signalling; (iii) biosynthesis of defensive secondary metabolites such as phytoalexins; or (iv) plant cell wall reinforcement (Shibuya and Minami, 2001) (Fig. 6; Supplementary Table S2 at JXB online). Some cell wall proteins were reported to be involved in cell wall strengthening and papilla formation [peroxidase and hydroxyproline-rich glycoprotein (HRGP)]. Among HGMEs, only PG and PL were identified following elicitation by OGs. Wound-inducible PG activity was correlated to  $H_2O_2$  production in most plants belonging to different families (Orozco-Cardenas and Ryan, 1999). Both responses were induced by OG treatment in tomato leaves (Orozco-Cardenas and Ryan, 1999), where PG gene expression and its corresponding activity were transiently increased (Bergey *et al.*, 1999). In contrast, in *Arabidopsis* seedlings, a mixture of OGs (DP 9–16) did not change overall PG gene expression but repressed PL-PMR6 expression (Denoux *et al.*, 2008). These differences are probably related to the plant species as well as the type and concentrations of OGs used. In fact, several studies have shown that the ability of OGs to act as elicitors is dependent on their chemical structure (DP, DM,

pattern of methylesterification, and DA) (Ochoa-Villarreal *et al.*, 2012). OGs with DP ranging from 1 to 20 are efficient elicitors (Côté and Hahn, 1994) (Fig. 6). The treatment by flagellin, a microbial-associated molecular pattern (MAMP) from bacteria, is mimicked by OGs in a range of DP from 9 to 16 to induce plant defences including cell wall reinforcement (Denoux *et al.*, 2008). The critical DP triggering plant elicitation is dependent on both the type of defence responses measured for a given plant and the type of plant species (Côté and Hahn, 1994). OGs with a high DA led to wheat resistance against the necrotrophic fungus *B. graminis* (Randoux *et al.*, 2010), while fully methylated OGs failed to induce defence signalling in soybean (Navazio *et al.*, 2002). This variability highlights the difficulty in unravelling the relationship between the structure and function of OGs. Furthermore, in adequate ionic conditions, the  $Ca^{2+}$ -egg box conformation of OGs improves their biological activity (Fig. 6). For example, dimeric and trimeric association of OGs induced a higher level of early and late defence responses in carrot (Messiaen *et al.*, 1993). The ability of multimeric forms of OGs to elicit plant responses appears to depend on their maturation, which is likely to be necessary for their fixation on OG receptors (Cabrera *et al.*, 2008). To date, wall-associated kinase 1 (WAK1), belonging to the WAK gene family (five in *Arabidopsis*), is the only receptor characterized for OG recognition inducing the defence signalling cascade (Wolf *et al.*, 2012a) (Fig. 6). Using chimeric proteins, the binding



**Fig. 6.** Involvement of HGMEs in plant defence responses to biotic stresses. Homogalacturonans (HGs) embedded in the cell wall after their synthesis in the Golgi apparatus with a high degree of polymerization (DP, minimum 72–100 galacturonic acid residues; Thibault *et al.*, 1993) are O-acetylated [with a degree of acetylation (DA) of ~10%; Ralet *et al.* (2005)], and highly methylated [with a degree of methylesterification (DM) of ~80%; O’Neill and Albersheim (1990)]. Depending on the stress and growth states, these HGs will be demethylesterified by pectin methylesterases (PMEs) and/or deacetylated by pectin acylesterases (PAEs), both enzyme activities being associated with the release of the volatile compounds methanol and ethanol, respectively. Depending on the cell wall properties, PMEs can act linearly, giving rise to blocks of free carboxyl groups that interact with bivalent ions (Ca<sup>2+</sup>) and contributing to cell wall strengthening (green frame), or can act randomly, promoting the action of downstream cell wall hydrolases [polygalacturonases (PGs) and pectate lyases (PLs)] and then contributing to cell wall loosening (Micheli, 2001). Cell wall strengthening restricts bioaggressor progression,



of OGs on its ectodomain was shown to activate specific plant defences following recognition by a leucine-rich repeat (LRR)-receptor kinase (EFR). Conversely, its intracellular kinase domain induced OG-specific plant defences after treatment by the EFR-specific elicitor (Brutus *et al.*, 2010). Interestingly, OG monomers, dimers, and trimers have been reported to be inhibitors of disease resistance reactions independently of the way they are produced (plant cell wall autolysis or pathogen CWDE digestion) (Messiaen *et al.*, 1993; Moerschbacher *et al.*, 1999) (Fig. 6). As monomeric OGs inhibit phytoalexin accumulation [i.e. phenylalanine ammonia lyase (PAL) activity] in carrot in contrast to the dimeric OG (Messiaen *et al.*, 1993), HGME activity associated with ionic cell status may be a way to regulate plant responses by modifying OG structure. In non-challenged plants, the difference in the rate of activity between intrinsic plant exo-PG and endo-PG that give rise to different OG structures (elicitors or not) supports a regulation of the balance between active/non-active OG forms by plant HGMEs during stress. Nevertheless, in plant–bioaggressor interactions, although the endogenous OG production is mostly described as being in favour of the plant, it may also benefit the bioaggressor. The combination of the action of both virulent microbes and plant HGMEs might suppress OGs with an eliciting function. PG, PL, and PME are found in plants and in secretions of most bioaggressors (microbes, nematodes, and insects) (Shen *et al.*, 2003, 2005; Harmel *et al.*, 2010; Sharma *et al.*, 2013) (Fig. 6). The types and distribution of HGMEs vary depending on the bioaggressor considered. For instance, among phytophagous insects, no HGME activity has been measured in chewing insects. In aphididae, PME activity has been detected in the saliva of all tested aphids except for *Sitobion avenae*, while they all possess PG activity (Harmel *et al.*, 2010). The differences in HGME content of bioaggressors, together with the activity of plant enzymes, may have an effect on the quantity and structure of OGs released, thus enhancing or inhibiting specific MAMPs or HAMPs (herbivore-associated molecular patterns) plant defence responses (Felton and Tumlinson, 2008). While an oxidative burst can be induced by OGs, ROS themselves (H<sub>2</sub>O<sub>2</sub>) can give rise to a non-enzymatic OG production by oxidative breakdown of pectins (Miller, 1986; Fry, 1998) (Fig. 6).

The aphid *Diuraphis noxia* induced massive H<sub>2</sub>O<sub>2</sub> accumulation in resistance of wheat, while this effect was not detectable in the *Brevicoryne brassicae*–*Arabidopsis* interaction (Moloi and van der Westhuizen, 2006; Kuśnierczyk *et al.*, 2008). The observation of H<sub>2</sub>O<sub>2</sub> accumulation in both compatible and incompatible interactions between the aphid

*Macrosiphum euphorbiae* and tomatoes suggested that they might not have a preponderant role as enhancers of endogenous elicitor production (De Ilarduya *et al.*, 2003). If the production of effective OGs is considered to be related to the balance between endogenous and exogenous HGME activities, specific regulators of HGMEs (PMEI, PGIP, and PNLIP) have to be taken into account (Table 3; Fig. 6). Among these protein inhibitors, at least one PGIP appears specifically directed to exogenous HGMEs and is used as a marker of the plant defence response. Plant PGIPs interact with PGs from various bioaggressors such as bacteria, fungi, and phytophagous insects (Albersheim and Anderson, 1971; D’Ovidio *et al.*, 2004; Schacht *et al.*, 2011) but are unlikely to target plant PGs (Cervone *et al.*, 1990; Federici *et al.*, 2001). PGIPs from different plant species can reduce PG activity from distinct pathogen species (Table 3), and one single PGIP can inhibit different fungal PGs; the PGIP2 of common bean inhibits *Fusarium moniliforme*, *Aspergillus niger*, and *B. cinerea* PGs (Federici *et al.*, 2001; King *et al.*, 2002; Sicilia *et al.*, 2005). The expression of a pea PGIP gene was induced during pea defence against the cyst nematode *Heterodera goettingiana*; none of the PGs from the nematode was shown to interact with the plant PGIP (Veronico *et al.*, 2011). Results concerning potential pectin lyase inhibitor protein (PNLIP) are rather scarce. A PNLIP from sugar beet (*Beta vulgaris*) was shown to inhibit fungal pectin lyases from *Rhizoctonia solani*, *Phoma betae*, and *Aspergillus japonicus*, but information about the inhibitor structure and regulation in plants is so far lacking (Table 3) (Bugbee, 1993; Juge, 2006). Plant PMEIs, which only target plant PMEs, could, however, play a role in resistance to pathogens by targeting specific plant PME isoforms, thus modulating the structure of pectins (Lionetti *et al.*, 2007).

Plant HGMEs are involved in the emission of two volatile organic compounds (VOCs), MeOH and EtOH (Yadav *et al.*, 2009), released by PME and PAE activity, respectively (Fig. 6). PME-mediated pectin remodelling appears to be the main MeOH producer, while EtOH is mostly attributed to fermentive reactions of glucose (Fall, 1999; Seco *et al.*, 2007). Among the huge range of VOCs produced during plant defence responses, an increase in MeOH emissions has been measured after wounding (Dorokhov *et al.*, 2012a) and feeding by herbivore caterpillars (Peñuelas *et al.*, 2005; Körner *et al.*, 2009). In tobacco leaves, rapid and sustained emission of MeOH was observed after *Manduca sexta* wounding, and was enhanced in the presence of caterpillar oral secretions, due to both up-regulation of gene expression and activity of plant PMEs, and a decrease in the DM of pectins (Von Dahl

while cell wall loosening may facilitate their penetration. The various bioaggressors (blue frames) that attempt to breach the plant cell wall release their own HGMEs (exogenous), including pectin lyases (PNLs), that could amplify or compete with the hydrolysis by plant HGMEs (endogenous). Consequently, following the activity of plant and bioaggressor hydrolases, small HG fragments called oligogalacturonides (OGs) are released. Most of these enzymes (plant PMEs or bioaggressor PGs, PNLs) could be modulated by PME, PG, or PNL inhibitors (PMEIs, PGIPs, and PNLIPs). Depending on their DP and the amount of free Ca<sup>2+</sup>, the OG monomers obtained can form multimers with an egg box conformation (Cabrera *et al.*, 2008) and/or can act as endogenous elicitors. After OG recognition by a specific receptor [wall-associated kinase 1 (WAK1); Brutus *et al.* (2010)], various defence responses (red frame) are induced against the bioaggressor (Ridley *et al.*, 2001) and could interact with growth modifications. Among them, ROS production (H<sub>2</sub>O<sub>2</sub>) can contribute to cell wall loosening with a non-enzymatic degradation of cell wall polysaccharides including HG (OG production; Miller, 1986; Fry, 1998). Plant PMEs also appear to act more directly during an interaction with some bioaggressors as they (bottom blue frame) specifically bind to a virus movement protein (MP; Chen *et al.*, 2000) or a nematode cellulose-binding protein (CBP; Hewezi *et al.*, 2008).

*et al.*, 2006). The role of MeOH in induced plant resistance appears dual. It may either diffuse as a signal or be catabolized into compounds that might be used in plant defence. First converted by a putative methanol oxidase into formaldehyde, a lethal compound, it is quickly bound to a nucleophile such as glutathione (*S*-formylglutathione), turned into formate, and used as a carbon source incorporated into plant one-carbon metabolism (C1 folate pool) or the Calvin–Benson cycle (CO<sub>2</sub>) (Gout *et al.*, 2000; Achkor *et al.*, 2003; Wojtasik *et al.*, 2011). Ethanol may also be oxidized into acetyl-coenzyme A via alcohol dehydrogenases involved in primary metabolism (Leblová *et al.*, 1977). In *Arabidopsis*, glutathione-dependent formaldehyde dehydrogenase, known as *S*-nitrosoglutathione reductase (GSNOR), plays a key role in regulating nitric oxide and *S*-nitrosoglutathione levels as well as being a signal in systemic resistance against pathogens (Martínez *et al.*, 1996; Rustérucchi *et al.*, 2007). As nitric oxide and *S*-nitrosothiols are signalling molecules that regulate immunity, MeOH release by PMEIs seems to act as a quantitative signal during plant–herbivore interactions. Silencing the endogenous PME gene suppressed MeOH release and led to reduced accumulation of PGIP involved in tobacco leaf resistance against *M. sexta* (Körner *et al.*, 2009). While wounding regulated the GSNOR gene, the direct effect of MeOH on its regulation is still not clear (Downie *et al.*, 2004). For instance, several MeOH-inducible genes were identified, some of which are known to encode proteins involved in plant resistance, especially in antibacterial resistance, virus spreading (Dorokhov *et al.*, 2012a), or anthocyanin and flavonoid content (Downie *et al.*, 2004). PME-mediated MeOH production was recently shown to act as a cross-kingdom signal (Dorokhov *et al.*, 2012b). Indeed, in mice, some MeOH-inducible genes are involved in their preference for MeOH sources such as wounded leaves. As caterpillar oral secretions increased VOC emission, which is known to attract predators or parasitoids against insects or nematodes (Kahl *et al.*, 2000; Heil, 2008), as well as MeOH (Von Dahl *et al.*, 2006), PME appears to play a role in both indirect (VOCs) and direct plant defences (signal/elicitor producer).

## Concluding remarks and perspectives

The contribution of the changes in the pectic network to the changes in cell wall rheology, enabling anisotropic growth or response to biotic stress, has been well documented over recent years. However, how the changes in HG-type pectins are spatially and temporally mediated, through the specific action of HGMEs, remains a central issue in our understanding of plant development. Until now, major advances in understanding the contribution of these enzymes to changes in development have mainly concerned PMEIs. This notably includes their roles in mediating discrete changes in HG structure during the interaction with pathogens, the regulation of primordia emergence, pollen tube and hypocotyl elongation, as well as the identification of novel post-translational control of their activity through the processing of the proteins by serine proteases and/or their interaction with

specific inhibitors (PMEIs). Although much progress has been made when considering PMEIs, many challenges remain; for instance, the identification of specific PME–SBT and PME–PMEI pairs in the cell wall, the understanding of the potential polarity of the trafficking of PMEIs to the cell wall, and its role in generating specific localized demethylesterification patterns through interaction of the enzymes with pH and ion microdomains. In addition, given the recent discovery of the interplay between hormonal signalling and PMEIs, further research could include the determination of the upstream regulators of PME transcription, including transcription factors and hormone levels, and possible feedback loops.

Other classes of plant HGMEs, including PGs, PLLs, and PAEs, have received surprisingly little attention over the last few years, which probably does not reflect the importance of these enzymes in mediating changes in HG structure. As for PMEIs, this could be related to the difficulties in determining strong phenotypes in KO mutants, with the occurrence of compensation mechanisms. When considering these multigenic families, the identification of compensation isoforms using dedicated tools, at both the transcript and protein levels, will help to provide a comprehensive overview of the underlying changes in cell wall structure. In particular, the roles of pectic fragments in generating plant responses to stress and in mediating changes in development will need further investigation. This will involve the identification of potential receptors of cell wall fragments, and of their specificity with regard to DP, DM, or DA. In parallel, the biochemical characterization of PMEIs, PGs, PLLs, and PAEs will enable their substrate specificity and pH preference to be determined. In particular, how the PME-mediated changes in HG structure can influence the activity of PGs, PLLs, and PAEs will be a key issue in our understanding of the possible interplay of these enzymes *in muro*. This could be used to implement the current models illustrating the interaction between HGs and HGMEs in the cell wall environment (Figs 4, 6).

## Supplementary data

Supplementary data are available at *JXB* online.

**Table S1.** Comparative inventory of the structural motifs of PME, PAE, PG, and PLL isoforms between dicot and monocot species.

**Table S2.** Gene expression variations of HG-modifying enzyme inhibitor proteins (PMEIs and PGIPs) after biotic stresses.

## Acknowledgements

The authors thank Ministry of Education support for the PhD scholarship CW. This work was supported by a grant from the Agence Nationale de la Recherche (ANR-09-BLANC-0007-01, GROWPEC project), by the EC INTERREG IVA No. 4166 ‘Trans channel Wallnet’ program, and by the Conseil Régional de Picardie through a PhD studentship awarded to FS. Financial support from the Institut Universitaire de France (IUF) to JP is gratefully acknowledged.

## References

Achkor H, Díaz M, Fernández MR, Biosca JA, Parés X, Martínez MC. 2003. Enhanced formaldehyde detoxification by overexpression of

- glutathione-dependent formaldehyde dehydrogenase from *Arabidopsis*. *Plant Physiology* **132**, 2248–2255.
- Agüero CB, Uratsu SL, Greve C, Powell AL, Labavitch JM, Meredith CP, Dandekar AM.** 2005. Evaluation of tolerance to Pierce's disease and *Botrytis* in transgenic plants of *Vitis vinifera* L. expressing the pear PGIP gene. *Molecular Plant Pathology* **6**, 43–51.
- Albersheim P, Anderson AJ.** 1971. Proteins from plant cell walls inhibit polygalacturonases secreted by plant pathogens. *Proceedings of the National Academy of Sciences, USA* **68**, 1–5.
- Al-Qsous S, Carpentier E, Klein-Eude D, Burel C, Mareck A, Dauchel H, Gomord V, Balangé A.** 2004. Identification and isolation of a pectin methyl-esterase isoform that could be involved in flax cell wall stiffening. *Planta* **219**, 369–378.
- An SH, Sohn KH, Choi HW, Hwang IS, Lee SC, Hwang BK.** 2008. Pepper pectin methyl-esterase inhibitor protein CaPME1 is required for antifungal activity, basal disease resistance and abiotic stress tolerance. *Planta* **228**, 61–78.
- André-Leroux G, Tessier D, Bonnin E.** 2005. Action pattern of *Fusarium moniliforme* endopolygalacturonase towards pectin fragments: comprehension and prediction. *Biochimica et Biophysica Acta* **1749**, 53–64.
- Arsovski AA, Haughn GW, Western TL.** 2010. Seed coat mucilage cells of *Arabidopsis thaliana* as a model for plant cell wall research. *Plant Signaling and Behavior* **5**, 796–801.
- Ascencio-Ibáñez J, Sozzani R, Lee T, Chu T, Wolfinger R, Cella R, Hanley-Bowdoin L.** 2008. Global analysis of *Arabidopsis* gene expression uncovers a complex array of changes impacting pathogen response and cell cycle during geminivirus infection. *Plant Physiology* **148**, 436–454.
- Atkinson RG, Sutherl PW, Johnston SL, Gunaseelan K, Hallett IC, Mitra D, Brummell DA, Schroder R, Johnston JW, Schaffer RJ.** 2012. Down-regulation of POLYGALACTURONASE1 alters firmness, tensile strength and water loss in apple (*Malus domestica*) fruit. *BMC Plant Biology* **12**, 1–13.
- Atmodjo MA, Hao Z, Mohnen D.** 2013. Evolving views of pectin biosynthesis. *Annual Review of Plant Biology* **64**, 747–779.
- Baebler Š, Krecic Stres H, Rotter A, Kogovsek P, Cankar K, Kok EJ, Gruden K, Kovac M, ŽEL J, Pompe Novak M.** 2009. PVYNTN elicits a diverse gene expression response in different potato genotypes in the first 12 h after inoculation. *Molecular Plant Pathology* **10**, 263–275.
- Barcala M, García A, Cabrera J, Casson S, Lindsey K, Favery B, García-Casado G, Solano R, Fenoll C, Escobar C.** 2010. Early transcriptomic events in microdissected *Arabidopsis* nematode-induced giant cells. *The Plant Journal* **61**, 698–712.
- Barnavon L, Doco T, Terrier N, Ageorges A, Romieu C, Pellerin P.** 2001. Involvement of pectin methyl-esterase during the ripening of grape berries: partial cDNA isolation, transcript expression and changes in the degree of methyl-esterification of cell wall pectins. *Phytochemistry* **58**, 693–701.
- Baron-Epel O, Gharyal PK, Schindler M.** 1988. Pectins as mediators of wall porosity in soybean cells. *Planta* **175**, 389–395.
- Bellincampi D, Camardella L, Delcour JA, et al.** 2004. Potential physiological role of plant glycosidase inhibitors. *Biochimica et Biophysica Acta* **1696**, 265–274.
- Bellincampi D, Salvi G, Lorenzo G, Cervone F, Marfa V, Eberhard S, Darvill A, Albersheim P.** 1993. Oligogalacturonides inhibit the formation of roots on tobacco explants. *The Plant Journal* **4**, 207–213.
- Benedetti M, Leggio C, Federici L, de Lorenzo G, Pavel NV, Cervone F.** 2011. Structural resolution of the complex between a fungal polygalacturonase and a plant polygalacturonase-inhibiting protein by small-angle X-ray scattering. *Plant Physiology* **157**, 599–607.
- Benen J, Kester H, Visser J.** 1999. Kinetic characterization of *Aspergillus niger* N400 endopolygalacturonases I, II and C. *European Journal of Biochemistry* **259**, 577–585.
- Ben-Hod G, Losner D, Joel DM, Mayer AM.** 1993. Pectin methyl-esterase in calli and germinating seeds of *Orobanche aegyptiaca*. *Phytochemistry* **32**, 1399–1402.
- Bergey DR, Orozco-Cardenas M, de Moura DS, Ryan CA.** 1999. A wound- and systemin-inducible polygalacturonase in tomato leaves. *Proceedings of the National Academy of Sciences, USA* **96**, 1756–1760.
- Besnard F, Vernoux T, Hamant O.** 2011. Organogenesis from stem cells in planta: multiple feedback loops integrating molecular and mechanical signals. *Cellular and Molecular Life Sciences* **68**, 2885–2906.
- Biely P, MacKenzie C, Puls J, Schneider H.** 1986. Cooperativity of esterases and xylanases in the enzymatic degradation of acetyl xylan. *Nature Biotechnology* **4**, 731–733.
- Bolvig P, Pauly M, Orfila C, Scheller H, Schnorr K.** 2003. Sequence analysis and characterisation of a novel pectin acetyl esterase from *Bacillus subtilis*. In: Voragen F, Schols H, Visser R. eds. *Advances in pectins and pectinase research*. Berlin: Springer, 315–330.
- Bonfante P.** 2001. At the interface between mycorrhizal fungi and plants: the structural organization of cell wall, plasma membrane and cytoskeleton. In: Esser K, ed. *The mycota, fungal associations*. Berlin: Springer, 45–61.
- Bonivento D, Pontiggia D, Matteo A, Fernandez Recio J, Salvi G, Tsernoglou D, Cervone F, Lorenzo G, Federici L.** 2007. Crystal structure of the endopolygalacturonase from the phytopathogenic fungus *Colletotrichum lupini* and its interaction with polygalacturonase-inhibiting proteins. *Proteins* **70**, 294–299.
- Bonnin E, Clavurier K, Daniel S, Kauppinen S, Mikkelsen JD, Thibault JF.** 2008. Pectin acetyl-esterases from *Aspergillus* are able to deacetylate homogalacturonan as well as rhamnogalacturonan. *Carbohydrate Polymers* **74**, 411–418.
- Bonnin E, Dolo E, Le Goff A, Thibault JF.** 2002. Characterisation of pectin subunits released by an optimised combination of enzymes. *Carbohydrate Research* **337**, 1687–1696.
- Bonnin E, Le Goff A, Alebeek G, Voragen A, Thibault JF.** 2003. Mode of action of *Fusarium moniliforme* endopolygalacturonase towards acetylated pectin. *Carbohydrate Polymers* **52**, 381–388.
- Bonnin E, Le Goff A, Körner R, Vigouroux J, Roepstorff P, Thibault JF.** 2002. Hydrolysis of pectins with different degrees and patterns of methylation by the endopolygalacturonase of *Fusarium moniliforme*. *Biochimica et Biophysica Acta* **1596**, 83–94.
- Bordenave M.** 1996. Analysis of pectin methyl esterases. *Plant Cell Wall Analysis* **17**, 165–180.
- Bordenave M, Goldberg R.** 1994. Immobilized and free apoplastic pectin methyl-esterases in mung bean hypocotyl. *Plant Physiology* **106**, 1151–1156.
- Bordenave M, Goldberg R, Huet JC, Pernollet JC.** 1995. A novel protein from mung bean hypocotyl cell walls with acetyl esterase activity. *Phytochemistry* **38**, 315–319.
- Borderies G, Jamet E, Lafitte C, Rossignol M, Jauneau A, Boudart G, Monsarrat B, Esquerré Tugayé MT, Boudet A, Pont-Lezica R.** 2003. Proteomics of loosely bound cell wall proteins of *Arabidopsis thaliana* cell suspension cultures: a critical analysis. *Electrophoresis* **24**, 3421–3432.
- Bosch M, Hepler PK.** 2005. Pectin methyl-esterases and pectin dynamics in pollen tubes. *The Plant Cell* **17**, 3219–3226.
- Bosch M, Cheung AY, Hepler PK.** 2005. Pectin methyl-esterase, a regulator of pollen tube growth. *Plant Physiology* **138**, 1334–1346.
- Bosch M, Hepler PK.** 2006. Silencing of the tobacco pollen pectin methyl-esterase NtPPME1 results in retarded *in vivo* pollen tube growth. *Planta* **223**, 736–745.
- Boudart G, Jamet E, Rossignol M, Lafitte C, Borderies G, Pont-Lezica R.** 2005. Cell wall proteins in apoplastic fluids of *Arabidopsis thaliana* rosettes: identification by mass spectrometry and bioinformatics. *Proteomics* **5**, 212–221.
- Boudart G, Latte C, Barthe JP, Frasez D, Esquerré-Tugayé MT.** 1998. Differential elicitation of defense responses by pectic fragments in bean seedlings. *Planta* **206**, 86–94.
- Braybrook SA, Peaucelle A.** 2013. Mechano-chemical aspects of organ formation in *Arabidopsis thaliana*: the relationship between auxin and pectin. *PLoS One* **8**, e57813.
- Brummell DA, Cin VD, Crisosto C, Labavitch J.** 2004. Cell wall metabolism during maturation, ripening and senescence of peach fruit. *Journal of Experimental Botany* **55**, 2029–2039.
- Brummell DA, Harpster MH.** 2001. Cell wall metabolism in fruit softening and quality and its manipulation in transgenic plants. *Plant Molecular Biology* **47**, 311–340.

- Brutus A, Sicilia F, Maccone A, Cervone F, De Lorenzo G.** 2010. A domain swap approach reveals a role of the plant wall-associated kinase 1 (WAK1) as a receptor of oligogalacturonides. *Proceedings of the National Academy of Sciences, USA* **107**, 9452–9457.
- Bugbee WM.** 1993. A pectin lyase inhibitor protein from cell walls of sugar beet. *Physiologia Plantarum* **83**, 63–67.
- Burton RA, Gidley MJ, Fincher GB.** 2010. Heterogeneity in the chemistry, structure and function of plant cell walls. *Nature Chemical Biology* **6**, 724–732.
- Cabrera JC, Boland A, Messiaen J, Cambier P, van Cutsem P.** 2008. Egg box conformation of oligogalacturonides: the time-dependent stabilization of the elicitor-active conformation increases its biological activity. *Glycobiology* **18**, 473–482.
- Cação SM, Leite TF, Budzinski IG, Santos TB, Scholz MB, Carpentieri-Pipolo V, Domingues DS, Vieira LG, Pereira LF.** 2012. Gene expression and enzymatic activity of pectin methylesterase during fruit development and ripening in *Coffea arabica* L. *Genetics and Molecular Research* **11**, 3186–3197.
- Cantarel BL, Coutinho PM, Rancurel C, Bernard T, Lombard V, Henrissat B.** 2009. The Carbohydrate-Active EnZymes database (CAZY): an expert resource for glycogenomics. *Nucleic Acids Research* **37**, 233–238.
- Camejo D, Martí MC, Jiménez A, Cabrera JC, Olmos E, Sevilla F.** 2011. Effect of oligogalacturonides on root length, extracellular alkalization and O<sub>2</sub>-accumulation in alfalfa. *Journal of Plant Physiology* **168**, 566–575.
- Cao J.** 2012. The pectin lyases in *Arabidopsis thaliana*: evolution, selection and expression profiles. *PLoS One* **7**, e46944.
- Carpita NC, Gibeault DM.** 1993. Structural models of primary cell walls in flowering plants: consistency of molecular structure with the physical properties of the walls during growth. *The Plant Journal* **3**, 1–30.
- Carvalho MC, Caldas DG, Carneiro RT, Moon DH, Salvatierra GR, Franceschini LM, Andrade A, Celedon PA, Oda S, Labate CA.** 2008. SAGE transcript profiling of the juvenile cambial region of *Eucalyptus grandis*. *Tree Physiology* **28**, 905–919.
- Catoire L, Pierron M, Morvan C, Penhoat du CH, Goldberg R.** 1998. Investigation of the action patterns of pectinmethylesterase isoforms through kinetic analyses and NMR spectroscopy. Implications in cell wall expansion. *Journal of Biological Chemistry* **273**, 33150–33156.
- Cervone F, De Lorenzo G, Pressey R, Darvill AG, Albersheim P.** 1990. Can Phaseolus PGIP inhibit pectic enzymes from microbes and plants? *Phytochemistry* **29**, 447–449.
- Chapman NH, Bonnet J, Grivet L, et al.** 2012. High-resolution mapping of a fruit firmness-related quantitative trait locus in tomato reveals epistatic interactions associated with a complex combinatorial locus. *Plant Physiology* **159**, 1644–1657.
- Charmont S, Jamet E, Pont-Lezica R, Canut H.** 2005. Proteomic analysis of secreted proteins from *Arabidopsis thaliana* seedlings: improved recovery following removal of phenolic compounds. *Phytochemistry* **66**, 453–461.
- Che P, Lall S, Nettleton D, Howell S.** 2006. Gene expression programs during shoot, root, and callus development in *Arabidopsis* tissue culture. *Plant Physiology* **141**, 620–637.
- Chebli Y, Kaneda M, Zerzour R, Geitmann A.** 2012. The cell wall of the *Arabidopsis* pollen tube—spatial distribution, recycling, and network formation of polysaccharides. *Plant Physiology* **160**, 1940–1955.
- Chen M, Citovsky V.** 2003. Systemic movement of a tobamovirus requires host cell pectin methylesterase. *The Plant Journal* **35**, 386–392.
- Chen E, Mort A.** 1996. Nature of sites hydrolyzable by endopolygalacturonase in partially-esterified homogalacturonans. *Carbohydrate Polymers* **29**, 129–136.
- Chen MH, Sheng J, Hind G, Handa AK, Citovsky V.** 2000. Interaction between the tobacco mosaic virus movement protein and host cell pectin methylesterases is required for viral cell-to-cell movement. *EMBO Journal* **19**, 913–920.
- Cho S, Lee S, Shin W.** 2001. The X-ray structure of *Aspergillus aculeatus* polygalacturonase and a modeled structure of the polygalacturonase–octagalacturonate complex. *Journal of Molecular Biology* **314**, 863–878.
- Chotigeat W, Duangchu S, Wititsuwannakun R, Phongdara A.** 2009. Cloning and characterization of pectate lyase from *Hevea brasiliensis*. *Plant Physiology and Biochemistry* **47**, 243–247.
- Chourasia A, Sane VA, Nath P.** 2006. Differential expression of pectate lyase during ethylene-induced postharvest softening of mango (*Mangifera indica* var. Dashehari). *Physiologia Plantarum* **128**, 546–555.
- Christensen T, Nielsen J, Mikkelsen J.** 1996. Isolation, characterization and immunolocalization of orange fruit acetyl esterase. *Progress in Biotechnology* **14**, 723–730.
- Chun J, Huber D.** 1998. Polygalacturonase-mediated solubilization and depolymerization of pectic polymers in tomato fruit cell walls. Regulation by pH and ionic conditions. *Plant Physiology* **117**, 1293–1299.
- Clausen MH, Willats WG, Knox JP.** 2003. Synthetic methyl hexagalacturonate hapten inhibitors of anti-homogalacturonan monoclonal antibodies LM7, JIM5 and JIM7. *Carbohydrate Research* **338**, 1797–1800.
- Consales F, Schweizer F, Erb M, Gouhier-Darimont C, Bodenhausen N, Bruessow F, Sobhy I, Reymond P.** 2011. Insect oral secretions suppress wound-induced responses in *Arabidopsis*. *Journal of Experimental Botany* **63**, 727–737.
- Cosgrove DJ.** 2005. Growth of the plant cell wall. *Nature Reviews Molecular Cell Biology* **6**, 850–861.
- Côté F, Hahn MG.** 1994. Oligosaccharins: structures and signal transduction. *Plant Molecular Biology* **26**, 1379–1411.
- Creze C, Castang S, Derivery E, Haser R, Hugouvieux-Cotte-Pattat N, Shevchik EV, Gouet P.** 2008. The crystal structure of pectate lyase Pell from soft rot pathogen *Erwinia chrysanthemi* in complex with its substrate. *Journal of Biological Chemistry* **283**, 18260–18268.
- Curvers K, Seifi H, Mouille G, et al.** 2010. Abscisic acid deficiency causes changes in cuticle permeability and pectin composition that influence tomato resistance to *Botrytis cinerea*. *Plant Physiology* **154**, 847–860.
- Damasceno CMB, Lohse M, Usadel B, Rose JKC, Fernie AR.** 2011. Systems biology of tomato fruit development: combined transcript, protein, and metabolite analysis of tomato transcription factor (*nor*, *rin*) and ethylene receptor (*Nr*) mutants reveals novel regulatory interactions. *Plant Physiology* **157**, 405–425.
- Davidson RM, Gowda M, Moghe G, Lin H, Vaillancourt B, Shiu S, Jiang N, Buell CR.** 2012. Comparative transcriptomics of three Poaceae species reveals patterns of gene expression evolution. *The Plant Journal* **71**, 492–202.
- D’Avino R, Camardella L, Christensen TMIE, Giovane A, Servillo L.** 2003. Tomato pectin methylesterase: modeling, fluorescence, and inhibitor interaction studies—comparison with the bacterial (*Erwinia chrysanthemi*) enzyme. *Proteins* **53**, 830–839.
- De Caroli M, Lenucci MS, Di Sansebastiano G, Dalessandro G, de Lorenzo G, Piro G.** 2011a. Protein trafficking to the cell wall occurs through mechanisms distinguishable from default sorting in tobacco. *The Plant Journal* **65**, 295–308.
- De Caroli M, Lenucci MS, Di Sansebastiano GP, Dalessandro G, de Lorenzo G, Piro G.** 2011b. Dynamic protein trafficking to the cell wall. *Plant Signaling and Behavior* **6**, 1012–1015.
- Dedeurwaerder S, Menu-Bouaouiche L, Mareck A, Lerouge P, Guerinéau F.** 2009. Activity of an atypical *Arabidopsis thaliana* pectin methylesterase. *Planta* **229**, 311–321.
- De Freitas ST, Handa AK, Wu Q, Park S, Mitcham EJ.** 2012. Role of pectin methylesterases in cellular calcium distribution and blossom-end rot development in tomato fruit. *The Plant Journal* **71**, 824–835.
- Deising H, Fritrang AK, Kunz S, Mendgen K.** 1995. Regulation of pectin methylesterase and polygalacturonate lyase activity during differentiation of infection structures in *Uromyces viciae-fabae*. *Microbiology* **141**, 561–571.
- De Ilarduya OM, Xie Q, Kaloshian I.** 2003. Aphid-induced defense responses in Mi-1-mediated compatible and incompatible tomato interactions. *Molecular Plant-Microbe Interactions* **16**, 699–708.
- De-la-Peña C, Badri DV, Vivanco JM.** 2008. Novel role for pectin methylesterase in *Arabidopsis*: a new function showing ribosome-inactivating protein (RIP) activity. *Biochimica et Biophysica Acta* **1780**, 773–783.

- Denès JM, Baron A, Renard CM, Péan C, Drilleau JF.** 2000. Different action patterns for apple pectin methylesterase at pH 7.0 and 4.5. *Carbohydrate Research* **327**, 385–393.
- Denoux C, Galletti R, Mammarella N, Gopalan S, Werck D, De Lorenzo G, Ferrari S, Ausubel FM, Dewdney J.** 2008. Activation of defense response pathways by OGs and Flg22 elicitors in *Arabidopsis* seedlings. *Molecular Plant* **1**, 423–445.
- Derbyshire P, McCann MC, Roberts K.** 2007. Restricted cell elongation in *Arabidopsis* hypocotyls is associated with a reduced average pectin esterification level. *BMC Plant Biology* **7**, 1–12.
- De Vos M, Jander G.** 2009. *Myzus persicae* (green peach aphid) salivary components induce defence responses in *Arabidopsis thaliana*. *Plant, Cell and Environment* **32**, 1548–1560.
- Deytieu-Belleau C, Vallet A, Donèche B, Geny L.** 2008. Pectin methylesterase and polygalacturonase in the developing grape skin. *Plant Physiology and Biochemistry* **46**, 638–646.
- Di Matteo A, Federici L, Mattei B, Salvi G, Johnson KA, Savino C, de Lorenzo G, Tsernoglou D, Cervone F.** 2003. The crystal structure of polygalacturonase-inhibiting protein (PGIP), a leucine-rich repeat protein involved in plant defense. *Proceedings of the National Academy of Sciences, USA* **100**, 10124–10128.
- Di Matteo A, Giovane A, Raiola A, Camardella L, Bonivento D, de Lorenzo G, Cervone F, Bellicampi D, Tsernoglou D.** 2005. Structural basis for the interaction between pectin methylesterase and a specific inhibitor protein. *The Plant Cell* **17**, 849–858.
- Di Matteo A, Sacco A, Anacleria M, Pezzotti M, Delledonne M, Ferrarini A, Frusciante L, Barone A.** 2010. The ascorbic acid content of tomato fruits is associated with the expression of genes involved in pectin degradation. *BMC Plant Biology* **10**, 1–11.
- Ding JL, Hsu JS, Wang MM, Tzen JT.** 2002. Purification and glycosylation analysis of an acidic pectin methylesterase in jelly fig (*Ficus awkeotsang*) achenes. *Journal of Agricultural and Food Chemistry* **50**, 2920–2925.
- Ding JL, Lee TT, Wang MM, Tai SS, Tzen JT.** 2000. Cloning and expression of an acidic pectin methylesterase from Jelly Fig (*Ficus awkeotsang*). *Journal of Agricultural and Food Chemistry* **4**, 3052–3057.
- Divol F, Vilaine F, Thibivilliers S, Amselem J, Palauqui J-C, Kusiak C, Dinant S.** 2005. Systemic response to aphid infestation by *Myzus persicae* in the phloem of *Apium graveolens*. *Plant Molecular Biology* **57**, 517–540.
- Dixit S, Upadhyay S, Singh H, Pandey B, Chandrashekar K, Verma P.** 2013. Pectin methylesterase of *Datura* species, purification, and characterization from *Datura stramonium* and its application. *Plant Signaling and Behavior* **8**, e25681.
- Do Amaral S, De Assis S, De Oliveira OF.** 2005. Partial purification and characterization of pectin methylesterase from orange (*Citrus sinensis*) cv. Pera-rio. *Journal of Food Biochemistry* **29**, 367–380.
- Domingo C, Roberts K, Stacey NJ, Connerton I, Ruiz-Teran F, McCann MC.** 1998. A pectate lyase from *Zinnia elegans* is auxin inducible. *The Plant Journal* **13**, 17–28.
- Dominguez-Puigjaner E, Llop I, Vendrell M, Prat S.** 1997. A cDNA clone highly expressed in ripe banana fruit shows homology to pectate lyases. *Plant Physiology* **114**, 1071–1076.
- Dorokhov YL, Komarova TV, Petrunia I.** 2012a. Airborne signals from a wounded leaf facilitate viral spreading and induce antibacterial resistance in neighboring plants. *PLoS Pathogens* **8**, e1002640.
- Dorokhov YL, Komarova TV, Petrunia IV, Kosorukov VS, Zinovkin RA, Shindyapina AV, Frolova OY, Gleba YY.** 2012b. Methanol may function as a cross-kingdom signal. *PLoS One* **7**, e36122.
- Dorokhov YL, Mäkinen K, Frolova O, Merits A, Saarinen J, Kalkkinen N, Atabekov J, Saarma M.** 1999. A novel function for a ubiquitous plant enzyme pectin methylesterase: the host-cell receptor for the tobacco mosaic virus movement protein. *FEBS Letters* **461**, 223–228.
- Dorokhov YL, Skurat EV, Frolova OY, et al.** 2006. Role of the leader sequence in tobacco pectin methylesterase secretion. *FEBS Letters* **580**, 3329–3334.
- D'Ovidio R, Raiola A, Capodicasa C, Devoto A, Pontiggia D, Roberti S, Galletti R, Conti E, O'Sullivan D, De Lorenzo G.** 2004. Characterization of the complex locus of bean encoding polygalacturonase-inhibiting proteins reveals subfunctionalization for defense against fungi and insects. *Plant Physiology* **135**, 2424–2435.
- Downie A, Miyazaki S, Bohnert H, John P, Coleman J, Parry M, Haslam R.** 2004. Expression profiling of the response of *Arabidopsis thaliana* to methanol stimulation. *Phytochemistry* **65**, 2305–2316.
- Draye M, Cutsem P.** 2008. Pectin methylesterases induce an abrupt increase of acidic pectin during strawberry fruit ripening. *Journal of Plant Physiology* **165**, 1152–1160.
- Dreyer D, Campbell B.** 1984. Association of the degree of methylation of intercellular pectin with plant resistance to aphids and with induction of aphid biotypes. *Experientia* **40**, 224–226.
- Ehwald R, Woehlecke H, Titel C.** 1992. Cell wall microcapsules with different porosity from suspension cultured *Chenopodium album*. *Phytochemistry* **31**, 3033–3038.
- Ericksson E, Bovy A, Manning K, Harrison L, Andrews J, Silva J, Tucker G, Seymour G.** 2004. Effect of the colorless non-ripening mutation on cell wall biochemistry and gene expression during tomato fruit development and ripening. *Plant Physiology* **136**, 4184–4197.
- Falasca G, Capitani F, Rovere F, Zaghi D, Franchin C, Biondi S, Altamura M.** 2008. Oligogalacturonides enhance cytokinin-induced vegetative shoot formation in tobacco explants, inhibit polyamine biosynthetic gene expression, and promote long-term remobilisation of cell calcium. *Planta* **227**, 835–852.
- Fall R.** 1999. Biogenic emissions of volatile organic compounds from higher plants. In: Hewitt CN, ed. *Reactive hydrocarbons in the atmosphere*. New York: Academic Press, 43–96.
- Federici L, Caprari C, Mattei B, Savino C, Di Matteo A, De Lorenzo G, Cervone F, Tsernoglou D.** 2001. Structural requirements of endopolygalacturonase for the interaction with PGIP (polygalacturonase-inhibiting protein). *Proceedings of the National Academy of Sciences, USA* **98**, 13425–13430.
- Felton GW, Tumlinson JH.** 2008. Plant–insect dialogs: complex interactions at the plant–insect interface. *Current Opinion in Plant Biology* **11**, 457–463.
- Ferrari S, Galletti R, Vairo D, Cervone F, De Lorenzo G.** 2006. Antisense expression of the *Arabidopsis thaliana* *AtPGIP1* gene reduces polygalacturonase-inhibiting protein accumulation and enhances susceptibility to *Botrytis cinerea*. *Molecular Plant-Microbe Interactions* **19**, 931–936.
- Ferrari S, Savatin D, Sicilia F, Gramegna G, Cervone F, de Lorenzo G.** 2013. Oligogalacturonides: plant damage-associated molecular patterns and regulators of growth and development. *Frontiers in Plant Science* **4**, 1–9.
- Ferrari S, Vairo D, Ausubel FM, Cervone F, De Lorenzo G.** 2003. Tandemly duplicated *Arabidopsis* genes that encode polygalacturonase-inhibiting proteins are regulated coordinately by different signal transduction pathways in response to fungal infection. *The Plant Cell* **15**, 93–106.
- Francis KE, Lam SY, Copenhaver GP.** 2006. Separation of *Arabidopsis* pollen tetrads is regulated by *QUARTET1*, a pectin methylesterase gene. *Plant Physiology* **142**, 1004–1013.
- Frenkel C.** 1998. Pectin methylesterase regulates methanol and ethanol accumulation in ripening tomato (*Lycopersicon esculentum*) fruit. *Journal of Biological Chemistry* **273**, 4293–4295.
- Fries M, Ihrig J, Brocklehurst K, Shevchik V, Pickersgill R.** 2007. Molecular basis of the activity of the phytopathogen pectin methylesterase. *EMBO Journal* **26**, 3879–3887.
- Fry SC.** 1998. Oxidative scission of plant cell wall polysaccharides by ascorbate-induced hydroxyl radicals. *Biochemical Journal* **332**, 507–515.
- Fry SC.** 2004. Primary cell wall metabolism: tracking the careers of wall polymers in living plant cells. *New Phytologist* **161**, 641–675.
- Gaffe J, Tieman D, Handa A.** 1994. Pectin methylesterase isoforms in tomato (*Lycopersicon esculentum*) tissues (effects of expression of a pectin methylesterase antisense gene). *Plant Physiology* **105**, 199–203.
- García JM, Posé S, Muñoz-Blanco J, Quesada MA, Mercado JA.** 2009. The polygalacturonase FaPG1 gene plays a key role in strawberry fruit softening. *Plant Signaling and Behavior* **4**, 766–768.

- Gasanova T, Skurat E, Frolova O, Semashko M, Dorokhov Y.** 2008. Pectin methylesterase as a factor of plant transcriptome stability. *Molecular Biology* **42**, 421–429.
- Geisler-Lee J, Geisler M, Coutinho PM, et al.** 2006. Poplar carbohydrate-active enzymes. Gene identification and expression analyses. *Plant Physiology* **140**, 946–962.
- Gendreau E, Traas J, Desnos T, Grandjean O, Caboche M, Höfte H.** 1997. Cellular basis of hypocotyl growth in *Arabidopsis thaliana*. *Plant Physiology* **114**, 295–305.
- Gibeaut DM, Pauly M, Bacic A.** 2005. Changes in cell wall polysaccharides in developing barley (*Hordeum vulgare*) coleoptiles. *Planta* **221**, 729–738.
- Gille S, Pauly M.** 2012. O-Acetylation of plant cell wall polysaccharides. *Frontiers in Plant Science* **3**, 1–7.
- Giovane A, Servillo L, Balestrieri C, Raiola A, D'Avino R, Tamburrini M, Ciardiello MA, Camardella L.** 2004. Pectin methylesterase inhibitor. *Biochimica et Biophysica Acta* **1696**, 245–252.
- Goda H, Sawa S, Asami T, Fujioka S, Shimada Y, Yoshida S.** 2004. Comprehensive comparison of auxin-regulated and brassinosteroid-regulated genes in *Arabidopsis*. *Plant Physiology* **134**, 1555–1573.
- Gonzalez-Carranza ZH, Elliott KA, Roberts JA.** 2007. Expression of polygalacturonases and evidence to support their role during cell separation processes in *Arabidopsis thaliana*. *Journal of Experimental Botany* **58**, 3719–3730.
- Gonzalez-Carranza ZH, Shahid AA, Zhang L, Liu Y, Ninsuwan U, Roberts JA.** 2012. A novel approach to dissect the abscission process in *Arabidopsis*. *Plant Physiology* **160**, 1342–1356.
- Gonzalez-Carranza ZH, Whitelaw CA, Swarup R, Roberts JA.** 2002. Temporal and spatial expression of a polygalacturonase during leaf and flower abscission in oilseed rape and *Arabidopsis*. *Plant Physiology* **128**, 534–543.
- Gou J, Miller LM, Hou G, Yu X, Chen X, Liu C.** 2012. Acetylase-mediated deacetylation of pectin impairs cell elongation, pollen germination, and plant reproduction. *The Plant Cell* **24**, 50–65.
- Gout E, Aubert S, Bligny R, Rébeillé F, Nonomura AR, Benson AA, Douce R.** 2000. Metabolism of methanol in plant cells. Carbon-13 nuclear magnetic resonance studies. *Plant Physiology* **123**, 287–296.
- Goulao LF, Vieira-Silva S, Jackson PA.** 2011. Association of hemicellulose- and pectin-modifying gene expression with *Eucalyptus globulus* secondary growth. *Plant Physiology and Biochemistry* **49**, 873–881.
- Grasdalen H, Andersen A, Larsen B.** 1996. NMR spectroscopy studies of the action pattern of tomato pectinesterase: generation of block structure in pectin by a multiple-attack mechanism. *Carbohydrate Research* **289**, 105–114.
- Guénin S, Mareck A, Rayon C, et al.** 2011. Identification of pectin methylesterase 3 as a basic pectin methylesterase isoform involved in adventitious rooting in *Arabidopsis thaliana*. *New Phytologist* **192**, 114–126.
- Habrant A, Gaillard C, Ralet M, Lairez D, Cathala B.** 2009. Relation between chemical structure and supramolecular organization of synthetic lignin-pectin particles. *Biomacromolecules* **10**, 3151–3156.
- Hadfield K, Bennett A.** 1998. Polygalacturonases: many genes in search of a function. *Plant Physiology* **117**, 337–343.
- Hamant O, Meyerowitz EM, Traas J.** 2011. Is cell polarity under mechanical control in plants? *Plant Signaling and Behavior* **6**, 137–139.
- Harmel N, Haubruge É, Francis F.** 2010. Étude des salives de pucerons: un préalable au développement de nouveaux bio-insecticides. *Biotechnologie, Agronomie, Société et Environnement* **14**, 369–378.
- Hasunuma T, Fukusaki E, Kobayashi A.** 2004. Expression of fungal pectin methylesterase in transgenic tobacco leads to alteration in cell wall metabolism and a dwarf phenotype. *Journal of Biotechnology* **111**, 241–251.
- Hause B, Fester T.** 2004. Molecular and cell biology of arbuscular mycorrhizal symbiosis. *Planta* **221**, 184–196.
- Hayashi K.** 2012. The interaction and integration of auxin signaling components. *Plant and Cell Physiology* **53**, 965–975.
- Heil M.** 2008. Indirect defence via tritrophic interactions. *New Phytologist* **178**, 41–61.
- Herron SR, Benen JA, Scavetta RD, Visser J, Jurnak F.** 2000. Structure and function of pectic enzymes: virulence factors of plant pathogens. *Proceedings of the National Academy of Sciences, USA* **97**, 8762–8769.
- Hewezi T, Howe P, Maier TR, Hussey RS, Mitchum MG, Davis EL, Baum TJ.** 2008. Cellulose binding protein from the parasitic nematode *Heterodera schachtii* interacts with *Arabidopsis* pectin methylesterase: cooperative cell wall modification during parasitism. *The Plant Cell* **20**, 3080–3093.
- Hohnjec N, Vieweg MF, Pühler A, Becker A, Küster H.** 2005. Overlaps in the transcriptional profiles of *Medicago truncatula* roots inoculated with two different Glomus fungi provide insights into the genetic program activated during arbuscular mycorrhiza. *Plant Physiology* **137**, 1283–1301.
- Hongo S, Sato K, Yokoyama R, Nishitani K.** 2012. Demethylesterification of the primary wall by PECTIN METHYLESTERASE35 provides mechanical support to the *Arabidopsis* stem. *The Plant Cell* **24**, 2624–2634.
- Hruz T, Laule O, Szabo G, Wessendorp F, Bleuler S, Oertle L, Widmayer P, Gruissem W, Zimmermann P.** 2008. Genevestigator V3: a reference expression database for the meta-analysis of transcriptomes. *Advances in Bioinformatics* **2008**, 1–5.
- Huang L, Cao J, Zhang A, Ye Y, Zhang Y, Liu T.** 2009a. The polygalacturonase gene *BcMF2* from *Brassica campestris* is associated with intine development. *Journal of Experimental Botany* **60**, 301–313.
- Huang K, Mellor KE, Paul SN, Lawson MJ, Mackey AJ, Timko MP.** 2012. Global changes in gene expression during compatible and incompatible interactions of cowpea (*Vigna unguiculata* L.) with the root parasitic angiosperm *Striga gesnerioides*. *BMC Genomics* **13**, 1–15.
- Huang L, Ye Y, Zhang Y, Zhang A, Liu T, Cao J.** 2009b. *BcMF9*, a novel polygalacturonase gene, is required for both *Brassica campestris* intine and exine formation. *Annals of Botany* **104**, 1339–1351.
- Hwang BH, Bae H, Lim H-S, Kim KB, Kim SJ, Im M-H, Park B-S, Kim DS, Kim J.** 2010. Overexpression of polygalacturonase-inhibiting protein 2 (*PGIP2*) of Chinese cabbage (*Brassica rapa* ssp. *pekinensis*) increased resistance to the bacterial pathogen *Pectobacterium carotovorum* ssp. *carotovorum*. *Plant Cell, Tissue and Organ Culture* **103**, 293–305.
- International Brachypodium Initiative.** 2010. Genome sequencing and analysis of the model grass *Brachypodium distachyon*. *Nature* **463**, 763–768.
- Irshad M, Canut H, Borderies G, Pont-Lezica R, Jamet E.** 2008. A new picture of cell wall protein dynamics in elongating cells of *Arabidopsis thaliana*: confirmed actors and newcomers. *BMC Plant Biology* **8**, 1–16.
- Ishii T.** 1997. O-Acetylated oligosaccharides from pectins of potato tuber cell walls. *Plant Physiology* **113**, 1265–1272.
- Ishii T, Matsunaga T.** 2001. Pectic polysaccharide rhamnogalacturonan II is covalently linked to homogalacturonan. *Phytochemistry* **57**, 969–974.
- Jamet E, Roujol D, San-Clemente H, Irshad M, Soubigou-Taconnat L, Renou J, Pont-Lezica R.** 2009. Cell wall biogenesis of *Arabidopsis thaliana* elongating cells: transcriptomics complements proteomics. *BMC Genomics* **10**, 1–12.
- Jaubert S, Laffaire J, Abad P, Rosso M.** 2002. A polygalacturonase of animal origin isolated from the root-knot nematode *Meloidogyne incognita*. *FEBS Letters* **522**, 109–112.
- Jayani R, Saxena S, Gupta R.** 2005. Microbial pectinolytic enzymes: a review. *Process Biochemistry* **40**, 2931–2944.
- Jenkins J, Mayans O, Smith D, Worboys K, Pickersgill RW.** 2001. Three-dimensional structure of *Erwinia chrysanthemi* pectin methylesterase reveals a novel esterase active site. *Journal of Molecular Biology* **305**, 951–960.
- Jenkins J, Pickersgill R.** 2001. The architecture of parallel  $\beta$ -helices and related folds. *Biophysics and Molecular Biology* **77**, 111–175.
- Jenkins J, Shevchik V, Hugouvieux-Cotte-Pattat N.** 2004. The crystal structure of pectate lyase Pel9A from *Erwinia chrysanthemi*. *Journal of Biological Chemistry* **279**, 9139–9145.
- Jiang S, Yang SL, Xie LF, San Pua CH, Zhang XQ, Yang WC, Sundaresan V, Ye D.** 2005. VANGUARD1 encodes a pectin methylesterase that enhances pollen tube growth in the *Arabidopsis* style and transmitting tract. *The Plant Cell* **17**, 584–596.

- Jimenez-Bermudez S, Redondo-Nevado J, Munoz-Blanco J, Caballero JL, Lopez-Aranda JM, Valpuesta V, Pliego-Alfaro F, Quesada MA, Mercado JA.** 2002. Manipulation of strawberry fruit softening by antisense expression of a pectate lyase gene. *Plant Physiology* **128**, 751–759.
- Johansson K, El-Ahmad M, Friemann R, Jörnvall H, Markovic O, Eklund H.** 2002. Crystal structure of plant pectin methyltransferase. *FEBS Letters* **514**, 243–249.
- Jolie RP, Duvetter T, Houben K, Clynen E, Sila DN, van Loey AMV, van Loey AM, Hendrickx ME.** 2009. Carrot pectin methyltransferase and its inhibitor from kiwi fruit: study of activity, stability and inhibition. *Innovative Food Science and Emerging Technologies* **10**, 601–609.
- Jolie RP, Duvetter T, van Loey AMV, van Loey AM, Hendrickx ME.** 2010. Pectin methyltransferase and its proteinaceous inhibitor: a review. *Carbohydrate Research* **345**, 2583–2595.
- Joubert DA, Slaughter AR, Kemp G, Becker JW, Krooshof GH, Bergmann C, Benen J, Pretorius IS, Vivier MA.** 2006. The grapevine polygalacturonase-inhibiting protein (VvPGIP1) reduces *Botrytis cinerea* susceptibility in transgenic tobacco and differentially inhibits fungal polygalacturonases. *Transgenic Research* **15**, 687–702.
- Juge N.** 2006. Plant protein inhibitors of cell wall degrading enzymes. *Trends in Plant Science* **11**, 359–367.
- Kabel MA, Waard P, Schols HA, Voragen AG.** 2003. Location of O-acetyl substituents in xylo-oligosaccharides obtained from hydrothermally treated Eucalyptus wood. *Carbohydrate Research* **338**, 69–77.
- Kahl J, Siemens DH, Aerts RJ, Gäbler R, Kühnemann F, Preston CA, Baldwin IT.** 2000. Herbivore-induced ethylene suppresses a direct defense but not a putative indirect defense against an adapted herbivore. *Planta* **210**, 336–342.
- Kanai M, Nishimura M, Hayashi M.** 2010. A peroxisomal ABC transporter promotes seed germination by inducing pectin degradation under the control of ABI5. *The Plant Journal* **62**, 936–947.
- Kars I, Krooshof GH, Wagemakers L, Joosten R, Benen JA, van Kan JA.** 2005. Necrotizing activity of five *Botrytis cinerea* endopolygalacturonases produced in *Pichia pastoris*. *The Plant Journal* **43**, 213–225.
- Kauppinen S, Christgau S, Kofod L, Halkier T, Dorreich K, Daldoge H.** 1995. Molecular cloning and characterization of a rhamnogalacturonan acetyltransferase from *Aspergillus aculeatus*. *Journal of Biological Chemistry* **270**, 27172–27178.
- Kempema LA, Cui X, Holzer FM, Walling LL.** 2007. *Arabidopsis* transcriptome changes in response to phloem-feeding silverleaf whitefly nymphs. Similarities and distinctions in responses to aphids. *Plant Physiology* **143**, 849–865.
- Kiefer L, York W, Darvill A, Albersheim P.** 1989. Xyloglucan isolated from suspension-cultured sycamore cell walls is O-acetylated. *Phytochemistry* **28**, 2105–2107.
- King D, Bergmann C, Orlando R, Benen JA, Kester HC, Visser J.** 2002. Use of amide exchange mass spectrometry to study conformational changes within the endopolygalacturonase II-homogalacturonan-polygalacturonase inhibiting protein system. *Biochemistry* **41**, 10225–10233.
- Kohorn BD, Kohorn SL.** 2012. The cell wall-associated kinases, WAKs, as pectin receptors. *Frontier in Plant Science* **3**, 1–5.
- Körner E, von Dahl CC, Bonaventure G, Baldwin IT.** 2009. Pectin methyltransferase NaPME1 contributes to the emission of methanol during insect herbivory and to the elicitation of defence responses in *Nicotiana attenuata*. *Journal of Experimental Botany* **60**, 2631–2640.
- Kuśnierczyk A, Winge P, Jørstad TS, TrocZYńska J, Rossiter JT, Bones AM.** 2008. Towards global understanding of plant defence against aphids—timing and dynamics of early *Arabidopsis* defence responses to cabbage aphid (*Brevicoryne brassicae*) attack. *Plant, Cell and Environment* **31**, 1097–1115.
- Lahaye M, Falourd X, Quemener B, Ralet M, Howad W, Dirlwanger E, Arús P.** 2012. Cell wall polysaccharide chemistry of peach genotypes with contrasted textures and other fruit traits. *Journal of Agricultural and Food Chemistry* **60**, 6594–6605.
- Lairez D, Cathala B, Monties B, Bedos-Belval F, Duran H, Gorrichon L.** 2005. Aggregation during coniferyl alcohol polymerization in pectin solution: a biomimetic approach of the first steps of lignification. *Biomacromolecules* **6**, 763–774.
- Lashbrook CC, Cai S.** 2008. Cell wall remodeling in Arabidopsis stamen abscission zones. *Plant Signaling and Behavior* **3**, 733–736.
- Laskowski M, Biller S, Stanley K, Kajstura T, Prusty R.** 2006. Expression profiling of auxin-treated Arabidopsis roots: toward a molecular analysis of lateral root emergence. *Plant and Cell Physiology* **47**, 788–792.
- Leake JR.** 1994. The biology of myco-heterotrophic ('saprophytic') plants. *New Phytologist* **127**, 171–216.
- Leblová S, Perglerová E, Hlochová J.** 1977. Comparative study of plant alcohol dehydrogenases. *Biologia Plantarum* **19**, 88–95.
- Lee KB.** 2007. Structure and development of the upper haustorium in the parasitic flowering plant *Cuscuta japonica* (Convolvulaceae). *American Journal of Botany* **94**, 737–745.
- Lewis KC, Selzer T, Shahar C, Udi Y, Tworowski D, Sagi I.** 2008. Inhibition of pectin methyl esterase activity by green tea catechins. *Phytochemistry* **69**, 2586–2592.
- Li X, Wu HX, Southerton SG.** 2011. Transcriptome profiling of *Pinus radiata* juvenile wood with contrasting stiffness identifies putative candidate genes involved in microfibril orientation and cell wall mechanics. *BMC Genomics* **12**, 1–16.
- Li X, Wu HX, Southerton SG.** 2012. Identification of putative candidate genes for juvenile wood density in *Pinus radiata*. *Tree Physiology* **32**, 1046–1057.
- Li Y, Zou J, Li M, Bilgin DD, Vodkin LO, Hartman GL, Clough SJ.** 2008. Soybean defense responses to the soybean aphid. *New Phytologist* **179**, 185–195.
- Lietzke S, Scavetta R, Yoder M, Jurnak F.** 1996. The refined three-dimensional structure of pectate lyase E from *Erwinia chrysanthemi* at 2.2 Å resolution. *Plant Physiology* **111**, 73–92.
- Lievens S, Goormachtig S, Holsters M.** 2001. A critical evaluation of differential display as a tool to identify genes involved in legume nodulation: looking back and looking forward. *Nucleic Acids Research* **29**, 3459–3468.
- Lievens S, Goormachtig S, Herman S, Holsters M.** 2002. Patterns of pectin methyltransferase transcripts in developing stem nodules of *Sesbania rostrata*. *Molecular Plant-Microbe Interactions* **15**, 164–168.
- Lin TP, Liu CC, Chen SW, Wang WY.** 1989. Purification and characterization of pectin methyltransferase from *Ficus awkeotsang* makino achenes. *Plant Physiology* **91**, 1445–1453.
- Lionetti V, Francocci F, Ferrari S, Volpi C, Bellincampi D, Galletti R, D'Ovidio R, De Lorenzo G, Cervone F.** 2010. Engineering the cell wall by reducing de-methyl-esterified homogalacturonan improves saccharification of plant tissues for bioconversion. *Proceedings of the National Academy of Sciences, USA* **107**, 616–621.
- Lionetti V, Cervone F, Bellincampi D.** 2012. Methyl esterification of pectin plays a role during plant-pathogen interactions and affects plant resistance to diseases. *Journal of Plant Physiology* **169**, 1623–1630.
- Lionetti V, Raiola A, Camardella L, Giovane A, Obel N, Pauly M, Favaron F, Cervone F, Bellincampi D.** 2007. Overexpression of pectin methyltransferase inhibitors in Arabidopsis restricts fungal infection by *Botrytis cinerea*. *Plant Physiology* **143**, 1871–1880.
- Little D, Gouhier-Darimont C, Bruessow F, Reymond P.** 2007. Oviposition by pierid butterflies triggers defense responses in *Arabidopsis*. *Plant Physiology* **143**, 784–800.
- Lohar DP, Sharopova N, Endre G, Peñuela S, Samac D, Town C, Silverstein KAT, Vandenbosch KA.** 2006. Transcript analysis of early nodulation events in *Medicago truncatula*. *Plant Physiology* **140**, 221–234.
- Losner-Goshen D, Portnoy VH, Mayer AM, Joel DM.** 1998. Pectolytic activity by the haustorium of the parasitic plant *Orobancha* L. (Orobanchaceae) in host roots. *Annals of Botany* **81**, 319–326.
- Louvet R, Rayon C, Doman J, et al.** 2011. Major changes in the cell wall during silique development in *Arabidopsis thaliana*. *Phytochemistry* **72**, 59–67.
- Lunn D, Phan TD, Tucker GA, Lycett GW.** 2013. Cell wall composition of tomato fruit changes during development and inhibition of vesicle trafficking is associated with reduced pectin levels and reduced softening. *Plant Physiology and Biochemistry* **66**, 91–97.
- Ly-Nguyen B, van Loey AM, Smout C, Verlent I, Duvetter T, Hendrickx ME.** 2004. Effect of intrinsic and extrinsic factors on the

- interaction of plant pectin methylesterase and its proteinaceous inhibitor from kiwi fruit. *Journal of Agricultural and Food Chemistry* **52**, 8144–8150.
- Manabe Y, Nafisi M, Verherbruggen Y, et al.** 2011. Loss-of-function mutation of *REDUCED WALL ACETYLATION2* in *Arabidopsis* leads to reduced cell wall acetylation and increased resistance to *Botrytis cinerea*. *Plant Physiology* **155**, 1068–1078.
- Manfredini C, Sicilia F, Ferrari S, Pontiggia D, Salvi G, Caprari C, Lorito M, Lorenzo GD.** 2005. Polygalacturonase-inhibiting protein 2 of *Phaseolus vulgaris* inhibits BcPG1, a polygalacturonase of *Botrytis cinerea* important for pathogenicity, and protects transgenic plants from infection. *Physiological and Molecular Plant Pathology* **67**, 108–115.
- Mareck A, Gaffé J, Morvan O, Alexandre C, Morvan C.** 1995. Characterization of isoforms of pectin methylesterase of *Linum usitatissimum* using polyclonal antibodies. *Plant and Cell Physiology* **36**, 409–417.
- Martínez MC, Achkor H, Persson B, Fernández MR, Shafqat J, Farrés J, Jörnvall H, Parés X.** 1996. *Arabidopsis* formaldehyde dehydrogenase. *European Journal of Biochemistry* **241**, 849–857.
- Marty P, Jouan B, Bertheau Y, Vian B, Goldberg R.** 1997. Charge density in stem cell walls of *Solanum tuberosum* genotypes and susceptibility to blackleg. *Phytochemistry* **44**, 1435–1441.
- Mathieu Y, Armen K, Xia H, Guern J, Koller A.** 1991. Membrane responses induced by oligogalacturonides in suspension-cultured tobacco cells. *The Plant Journal* **1**, 333–343.
- Mauro M, Lorenzo G, Costantino P, Bellincampi D.** 2002. Oligogalacturonides inhibit the induction of late but not of early auxin-responsive genes in tobacco. *Planta* **215**, 494–501.
- Mayans O, Scott M, Connerton I, Gravesen T, Benen J, Visser J, Pickersgill R, Jenkins J.** 1997. Two crystal structures of pectin lyase A from *Aspergillus* reveal a pH driven conformational change and striking divergence in the substrate-binding clefts of pectin and pectate lyases. *Current Biology* **5**, 677–689.
- Mbégué-A-Mbégué D, Hubert O, Baurens F, Matsumoto T, Chillet M, Fils-Lycaon B, Sidibé-Bocs S.** 2009. Expression patterns of cell wall-modifying genes from banana during fruit ripening and in relationship with finger drop. *Journal of Experimental Botany* **60**, 2021–2034.
- McMillan GP, Hedley D, Fyffe L, Pérombelon MCM.** 1993. Potato resistance to soft-rot erwinias is related to cell wall pectin esterification. *Physiological and Molecular Plant Pathology* **42**, 279–289.
- Mercadante D, Melton L, Jameson G, Williams MK, De Simone A.** 2013. Substrate dynamics in enzyme action: rotations of monosaccharide subunits in the binding groove are essential for pectin methylesterase processivity. *Biophysical Journal* **104**, 1731–1739.
- Mertens JA, Bowman MJ.** 2011. Expression and characterization of fifteen *Rhizopus oryzae* 99–880 polygalacturonase enzymes in *Pichia pastoris*. *Current Microbiology* **62**, 1173–1178.
- Messiaen J, Read N, Cutsem P, Trewavas A.** 1993. Cell wall oligogalacturonides increase cytosolic free calcium in carrot protoplasts. *Journal of Cell Science* **104**, 365.
- Micheli F.** 2001. Pectin methylesterases: cell wall enzymes with important roles in plant physiology. *Trends in Plant Science* **6**, 414–419.
- Micheli F, Sundberg B, Goldberg R, Richard L.** 2000. Radial distribution pattern of pectin methylesterases across the cambial region of hybrid aspen at activity and dormancy. *Plant Physiology* **124**, 191–199.
- Miller AR.** 1986. Oxidation of cell wall polysaccharides by hydrogen peroxide: a potential mechanism for cell wall breakdown in plants. *Biochemical and Biophysical Research Communications* **141**, 238–244.
- Moerschbacher BM, Mierau M, Graebner B, Noll U, Mort AJ.** 1999. Small oligomers of galacturonic acid are endogenous suppressors of disease resistance reactions in wheat leaves. *Journal of Experimental Botany* **50**, 605–612.
- Mohnen D.** 2008. Pectin structure and biosynthesis. *Current Opinion in Plant Biology* **11**, 266–277.
- Mollet J, Leroux C, Dardelle F, Lehner A.** 2013. Cell wall composition, biosynthesis and remodeling during pollen tube growth. *Plants* **2**, 107–147.
- Moloi M, van der Westhuizen A.** 2006. The reactive oxygen species are involved in resistance responses of wheat to the Russian wheat aphid. *Journal of Plant Physiology* **163**, 1118–1125.
- Monshausen GB, Miller ND, Murphy AS, Gilroy S.** 2011. Dynamics of auxin-dependent  $Ca^{2+}$  and pH signaling in root growth revealed by integrating high-resolution imaging with automated computer vision-based analysis. *The Plant Journal* **65**, 309–318.
- Morris ER, Powell DA, Rees DA.** 1982. Conformations and interactions of pectins. I. Polymorphism between gel and solid states of calcium polygalacturonate. *Journal of Molecular Biology* **155**, 507–516.
- Moscatiello R, Mariani P, Sanders D, Maathuis FJM.** 2006. Transcriptional analysis of calcium-dependent and calcium-independent signalling pathways induced by oligogalacturonides. *Journal of Experimental Botany* **57**, 2847–2865.
- Müller K, Levesque-Tremblay G, Bartels S, Weitbrecht K, Wormit A, Usadel B, Haughn G, Kermodé AR.** 2013. Demethylesterification of cell wall pectins in *Arabidopsis* plays a role in seed germination. *Plant Physiology* **161**, 305–316.
- Muñoz JA, Coronado C, Pérez-Hormaeche J, Kondorosi A, Ratet P, Palomares AJ.** 1998. *MsPG3*, a *Medicago sativa* polygalacturonase gene expressed during the alfalfa–*Rhizobium meliloti* interaction. *Proceedings of the National Academy of Sciences, USA* **95**, 9687–9692.
- Narusaka Y, Narusaka M, Motoaki S, Ishida J, Shinozaki K, Nan Y, Park P, Shiraiishi T, Kobayashi M.** 2005. Cytological and molecular analyses of non-host resistance of *Arabidopsis thaliana* to *Alternaria alternata*. *Molecular Plant Pathology* **6**, 615–627.
- Navazio L, Moscatiello R, Bellincampi D, Baldan B, Meggio F, Brini M, Bowler C, Mariani P.** 2002. The role of calcium in oligogalacturonide-activated signalling in soybean cells. *Planta* **215**, 596–605.
- Ngouémazong DE, Jolie RP, Cardinaels R, Fraeye I, van Loey AV, van Loey A, Moldenaers P, Hendrickx M.** 2012. Stiffness of  $Ca^{2+}$ -pectin gels: combined effects of degree and pattern of methylesterification for various  $Ca^{2+}$  concentrations. *Carbohydrate Research* **348**, 69–76.
- Nogota Y, Ohta H, Voragen A.** 1993. Polygalacturonase in strawberry fruit. *Phytochemistry* **34**, 617–620.
- Ochoa-Villarreal M, Aispuro-Hernández E, Vargas-Arispuro I, Martínez-Téllez MÁ.** 2012. Plant cell wall polymers: function, structure and biological activity of their derivatives. In: De Souza Gomes A, ed. *Polymerization*. InTech, 64–89.
- Oelofse D, Dubery IA, Meyer R, Arendse MS, Gazendam I, Berger DK.** 2006. Apple polygalacturonase inhibiting protein1 expressed in transgenic tobacco inhibits polygalacturonases from fungal pathogens of apple and the anthracnose pathogen of lupins. *Phytochemistry* **67**, 255–263.
- Oldroyd GE, Murray JD, Poole PS, Downie JA.** 2011. The rules of engagement in the legume–rhizobial symbiosis. *Annual Review of Genetics* **45**, 119–144.
- Oosterveld A, Beldman G, Searle-van Leeuwen, Voragen AGJ.** 2000. Effect of enzymatic deacetylation on gelation of sugar beet pectin in the presence of calcium. *Carbohydrate Polymers* **43**, 249–256.
- Orfila C, Degan FD, Jørgensen B, Scheller HV, Ray PM, Ulvskov P.** 2012. Expression of mung bean pectin acetyl esterase in potato tubers: effect on acetylation of cell wall polymers and tuber mechanical properties. *Planta* **236**, 185–196.
- Orozco-Cardenas M, Ryan C.** 1999. Hydrogen peroxide is generated systemically in plant leaves by wounding and systemin via the octadecanoid pathway. *Proceedings of the National Academy of Sciences, USA* **96**, 6553–6557.
- Osorio S, Alba R, Damasceno CMB, et al.** 2011b. Systems biology of tomato fruit development: combined transcript, protein, and metabolite analysis of tomato transcription factor (nor, rin) and ethylene receptor (Nr) mutants reveals novel regulatory interactions. *Plant Physiology* **157**, 405–425.
- Osorio S, Bombarely A, Giavalisco P, Usadel B, Stephens C, Aragüez I, Medina-Escobar N, Botella MA, Fernie AR, Valpuesta V.** 2011a. Demethylation of oligogalacturonides by FaPE1 in the fruits of the wild strawberry *Fragaria vesca* triggers metabolic and transcriptional changes associated with defence and development of the fruit. *Journal of Experimental Botany* **62**, 2855–2873.
- Osorio S, Castillejo C, Quesada MA, Medina-Escobar N, Brownsey GJ, Suau R, Heredia A, Botella MA, Valpuesta V.** 2008. Partial demethylation of oligogalacturonides by pectin methyl esterase 1 is required for eliciting defence responses in wild strawberry (*Fragaria vesca*). *The Plant Journal* **54**, 43–55.



- Palin R, Geitmann A.** 2012. The role of pectin in plant morphogenesis. *Biosystem* **109**, 397–402.
- Palusa SG, Golovkin M, Shin SB, Richardson DN, Reddy AS.** 2007. Organ-specific, developmental, hormonal and stress regulation of expression of putative pectate lyase genes in Arabidopsis. *New Phytologist* **174**, 537–550.
- Pathak N, Mishra S, Sanwal GG.** 2000. Purification and characterization of polygalacturonase from banana fruit. *Phytochemistry* **54**, 147–152.
- Pathak N, Sanwal GG.** 1998. Multiple forms of polygalacturonase from banana fruits. *Phytochemistry* **48**, 249–255.
- Payasi A, Sanwal GG.** 2003. Pectate lyase activity during ripening of banana fruit. *Phytochemistry* **63**, 243–248.
- Paynel F, Schaumann A, Arkoun M, Douchiche O, Morvan C.** 2009. Temporal regulation of cell-wall pectin methyltransferase and peroxidase isoforms in cadmium-treated flax hypocotyl. *Annals of Botany* **104**, 1363–1372.
- Peaucelle A, Braybrook SA, Le Guillou L, Bron E, Kuhlemeier C, Höfte H.** 2011a. Pectin-induced changes in cell wall mechanics underlie organ initiation in Arabidopsis. *Current Biology* **21**, 1720–1726.
- Peaucelle A, Louvet R, Johansen JN, Höfte H, Laufs P, Pelloux J, Mouille G.** 2008. Arabidopsis phyllotaxis is controlled by the methyl-esterification status of cell-wall pectins. *Current Biology* **18**, 1943–1948.
- Peaucelle A, Louvet R, Johansen JN, et al.** 2011b. The transcription factor BELLRINGER modulates phyllotaxis by regulating the expression of a pectin methyltransferase in Arabidopsis. *Development* **138**, 4733–4741.
- Pelletier S, Van Orden J, Wolf S, et al.** 2010. A role for pectin de-methyltransferase in a developmentally regulated growth acceleration in dark-grown Arabidopsis hypocotyls. *New Phytologist* **188**, 726–739.
- Pelloux J, Rustérucci C, Mellerowicz EJ.** 2007. New insights into pectin methyltransferase structure and function. *Trends in Plant Science* **12**, 267–277.
- Peña-Urbe C, García-Pineda E, Beltrán-Peña E, Cruz H.** 2012. Oligogalacturonides inhibit growth and induce changes in S6K phosphorylation in maize (*Zea mays* L. var. Chalqueño). *Plant Growth Regulation* **67**, 151–159.
- Peñuelas J, Filella I, Stefanescu C, Llusia J.** 2005. Caterpillars of *Euphydryas aurinia* (Lepidoptera: Nymphalidae) feeding on *Succisa pratensis* leaves induce large foliar emissions of methanol. *New Phytologist* **167**, 851–857.
- Peretto R, Bonfante P, Bettini V, Favaron F, Alghisi P.** 1995. Polygalacturonase activity and location in arbuscular mycorrhizal roots of *Allium porrum* L. *Mycorrhiza* **5**, 157–163.
- Perfect SE, Green JR.** 2001. Infection structures of biotrophic and hemibiotrophic fungal plant pathogens. *Molecular Plant Pathology* **2**, 101–108.
- Perrot-Rechenmann C.** 2010. Cellular responses to auxin: division versus expansion. *Cold Spring Harbor Perspectives in Biology* **2**, 1–15.
- Philipp K, Fuchs I, Luthen H, et al.** 1999. Auxin-induced K<sup>+</sup> channel expression represents an essential step in coleoptile growth and gravitropism. *Proceedings of the National Academy of Sciences, USA* **96**, 12186–12191.
- Philipp K, Ivashikina N, Ache P, Christian M, Luthen H, Palme K, Hedrich R.** 2004. Auxin activates KAT1 and KAT2, two K<sup>+</sup>-channel genes expressed in seedlings of *Arabidopsis thaliana*. *The Plant Journal* **37**, 815–827.
- Pickersgill R, Jenkins J, Harris G, Nasser W, Robert-Baudouy.** 1994. The structure of *Bacillus subtilis* pectate lyase in complex with calcium. *Nature Structural Biology* **1**, 717–723.
- Pickersgill R, Smith D, Worboys K, Jenkins J.** 1998. Crystal structure of polygalacturonase from *Erwinia carotovora* ssp. *carotovora*. *Journal of Biological Chemistry* **273**, 24660–24664.
- Pilling J, Willmitzer L, Bucking H, Fisahn J.** 2004. Inhibition of a ubiquitously expressed pectin methyl transferase in *Solanum tuberosum* L. affects plant growth, leaf growth polarity, and ion partitioning. *Planta* **219**, 32–40.
- Pilnik W, Rombouts FM.** 1981. Pectic enzymes. In: Birch GG, Blakebrough N, Parker KJ. *Enzymes and food processing*. Dordrecht, The Netherlands: Springer, 105–128.
- Popeijus H, Overmars H, Jones J, Blok V, Goverse A, Helder J, Schots A, Bakker J, Smant G.** 2000. Enzymology: degradation of plant cell walls by a nematode. *Nature* **406**, 36–37.
- Powell AL, van Kan J, Have ten A, Visser J, Greve LC, Bennett AB, Labavitch JM.** 2000. Transgenic expression of pear PGIP in tomato limits fungal colonization. *Molecular Plant-Microbe Interactions* **13**, 942–950.
- Pressey R, Avants JK.** 1973. Separation and characterization of endopolygalacturonase and exopolygalacturonase from peaches. *Plant Physiology* **52**, 252–256.
- Pressey R, Avants JK.** 1977. Occurrence and properties of polygalacturonase in *Avena* and other plants. *Plant Physiology* **60**, 548–553.
- Protsenko MA, Buza NL, Krinitsyna AA, Bulantseva EA, Korableva NP.** 2008. Polygalacturonase-inhibiting protein is a structural component of plant cell wall. *Biochemistry (Moscow)* **73**, 1053–1062.
- Pua EC, Ong CK, Liu P, Liu JZ.** 2001. Isolation and expression of two pectate lyase genes during fruit ripening of banana (*Musa acuminata*). *Physiologia Plantarum* **113**, 92–99.
- Punta M, Coghill PC, Eberhardt RY, et al.** 2012. The Pfam protein families database. *Nucleic Acids Research* **40**, 290–301.
- Puthoff D, Nettleton D, Rodermeil S, Baum T.** 2003. *Arabidopsis* gene expression changes during cyst nematode parasitism revealed by statistical analyses of microarray expression profiles. *The Plant Journal* **33**, 911–921.
- Qu T, Liu R, Wang W, An L, Chen T, Liu G, Zhao Z.** 2011. Brassinosteroids regulate pectin methyltransferase activity and AtPME41 expression in Arabidopsis under chilling stress. *Cryobiology* **63**, 111–117.
- Qubbaj T, Reineke A, Zebitz C.** 2005. Molecular interactions between rosy apple aphids, *Dysaphis plantaginea*, and resistant and susceptible cultivars of its primary host *Malus domestica*. *Entomologia Experimentalis et Applicata* **115**, 145–152.
- Quesada MA, Blanco-Portales R, Pose S, Garcia-Gago JA, Jimenez-Bermudez S, Munoz-Serrano A, Caballero JL, Piiego-Alfaro F, Mercado JA, Munoz-Blanco J.** 2009. Antisense down-regulation of the FaPG1 gene reveals an unexpected central role for polygalacturonase in strawberry fruit softening. *Plant Physiology* **150**, 1022–1032.
- Raiola A, Camardella L, Giovane A, Mattei B, de Lorenzo G, Cervone F, Bellincampi D.** 2004. Two *Arabidopsis thaliana* genes encode functional pectin methyltransferase inhibitors. *FEBS Letters* **557**, 199–203.
- Raiola A, Lionetti V, Elmaghraby I, Immerzeel P, Mellerowicz E, Salvi G, Cervone F, Bellincampi D.** 2011. Pectin methyltransferase is induced in *Arabidopsis* upon infection and is necessary for a successful colonization by necrotrophic pathogens. *Molecular Plant-Microbe Interactions* **24**, 432–440.
- Ralet M, Cabrera JC, Bonnin E, Quémener B, Hellin P, Thibault J.** 2005. Mapping sugar beet pectin acetylation pattern. *Phytochemistry* **66**, 1832–1843.
- Ralet M, Crépeau M, Buchholt H, Thibault J.** 2003. Polyelectrolyte behaviour and calcium binding properties of sugar beet pectins differing in their degrees of methylation and acetylation. *Biochemical Engineering Journal* **16**, 191–201.
- Randoux B, Renard-Merlier D, Mulard G, Rossard S, Duyme F, Sanssené J, Courtois J, Durand R, Reignault P.** 2010. Distinct defenses induced in wheat against powdery mildew by acetylated and nonacetylated oligogalacturonides. *Phytopathology* **100**, 1352–1363.
- Rautengarten C, Usadel B, Neumetzler L, Hartmann J, Büssis D, Altmann T.** 2008. A subtilisin-like serine protease essential for mucilage release from Arabidopsis seed coats. *The Plant Journal* **54**, 466–480.
- Rayle DL, Cleland RE.** 1992. The acid growth theory of auxin-induced cell elongation is alive and well. *Plant Physiology* **99**, 1271–1274.
- Reca IB, Lionetti V, Camardella L, D'Avino R, Giardina T, Cervone F, Bellincampi D.** 2012. A functional pectin methyltransferase inhibitor protein (SolyPMEI) is expressed during tomato fruit ripening and interacts with PME-1. *Plant Molecular Biology* **79**, 429–442.
- Refregier G, Pelletier S, Jaillard D, Höfte H.** 2004. Interaction between wall deposition and cell elongation in dark-grown hypocotyl cells in Arabidopsis. *Plant Physiology* **135**, 959–968.

- Renard C, Jarvis M.** 1999. Acetylation and methylation of homogalacturonans 2: effect on ion-binding properties and conformations. *Carbohydrate Polymers* **39**, 209–216.
- Rencoret J, Gutierrez A, Nieto L, Jimenez-Barbero J, Faulds CB, Kim H, Ralph J, Martinez AT, Rio JC.** 2011. Lignin composition and structure in young versus adult *Eucalyptus globulus* plants. *Plant Physiology* **155**, 667–682.
- Rexova-Benkova L, Markovic O.** 1976. Pectic enzymes. *Advances in Carbohydrate Chemistry and Biochemistry* **33**, 323–385.
- Rhee SY, Osborne E, Poindexter P, Somerville C.** 2003. Microspore separation in the quartet 3 mutants of *Arabidopsis* is impaired by a defect in a developmentally regulated polygalacturonase required for pollen mother cell wall degradation. *Plant Physiology* **133**, 1170–1180.
- Ribeiro DM, Araujo WL, Fernie AR, Schippers JH, Mueller-Roeber B.** 2012. Translatome and metabolome effects triggered by gibberellins during rosette growth in *Arabidopsis*. *Journal of Experimental Botany* **63**, 2769–2786.
- Ridley BL, O'Neill MA, Mohnen D.** 2001. Pectins: structure, biosynthesis, and oligogalacturonide-related signaling. *Phytochemistry* **57**, 929–967.
- Rober-Kleber N, Albrechtová JT, Fleig S, Huck N, Michalke W, Wagner E, Speth V, Neuhaus G, Fischer-Iglesias C.** 2003. Plasma membrane H<sup>+</sup>-ATPase is involved in auxin-mediated cell elongation during wheat embryo development. *Plant Physiology* **131**, 1302–1312.
- Robert H, Friml J.** 2009. Auxin and other signals on the move in plants. *Nature Chemical Biology* **5**, 325–332.
- Roongsattham P, Morcillo F, Jantasuriyarat C, et al.** 2012. Temporal and spatial expression of polygalacturonase gene family members reveals divergent regulation during fleshy fruit ripening and abscission in the monocot species oil palm. *BMC Plant Biology* **12**, 1–15.
- Rose J, Braam J, Fry S.** 2002. The XTH family of enzymes involved in xyloglucan endotransglucosylation and endohydrolysis: current perspectives and a new unifying nomenclature. *Plant and Cell Physiology* **43**, 1421–1435.
- Ross HA, Morris WL, Ducreux LJM, et al.** 2011a. Pectin engineering to modify product quality in potato. *Plant Biotechnology Journal* **9**, 848–856.
- Ross HA, Wright KM, McDougall GJ, et al.** 2011b. Potato tuber pectin structure is influenced by pectin methyl esterase activity and impacts on cooked potato texture. *Journal of Experimental Botany* **62**, 371–381.
- Rustérucci C, Espunya MC, Díaz M, Chabannes M, Martínez MC.** 2007. S-nitrosoglutathione reductase affords protection against pathogens in *Arabidopsis*, both locally and systemically. *Plant Physiology* **143**, 1282–1292.
- Rybel BD, Rybel B, Vassileva V, et al.** 2010. A novel Aux/IAA28 signaling cascade activates GATA23-dependent specification of lateral root founder cell identity. *Current Biology* **20**, 1697–1706.
- Röckel N, Wolf S, Kost B, Rausch T, Greiner S.** 2008. Elaborate spatial patterning of cell-wall PME and PME1 at the pollen tube tip involves PME1 endocytosis, and reflects the distribution of esterified and de-esterified pectins. *The Plant Journal* **53**, 133–143.
- Saez-Aguayo S, Ralet MC, Berger A, Botran L, Ropartz D, Marion-Poll A, North HM.** 2013. PECTIN METHYLESTERASE INHIBITOR6 promotes *Arabidopsis* mucilage release by limiting methylesterification of homogalacturonan in seed coat epidermal cells. *The Plant Cell* **25**, 308–323.
- Santiago-Doménech N, Jimenez-Bermudez S, Matas A, Rose J, Munoz-Blanco J, Mercado J, Quesada M.** 2008. Antisense inhibition of a pectate lyase gene supports a role for pectin depolymerization in strawberry fruit softening. *Journal of Experimental Botany* **59**, 2769–2779.
- Santuari L, Scacchi E, Rodriguez-Villalon A, Salinas P, Dohmann EN, Brunoud G, Vernoux T, Smith R, Hardtke C.** 2011. Positional information by differential endocytosis splits auxin response to drive *Arabidopsis* root meristem growth. *Current Biology* **21**, 1918–1923.
- Sassi M, Lu Y, Zhang Y, et al.** 2012. COP1 mediates the coordination of root and shoot growth by light through modulation of PIN1- and PIN2-dependent auxin transport in *Arabidopsis*. *Development* **139**, 3402–3412.
- Sathiyaraj G, Srinivasan S, Subramaniam S, Kim Y, Kim Y, Kwon W, Yang D.** 2010. Polygalacturonase inhibiting protein: isolation, developmental regulation and pathogen related expression in *Panax ginseng* C.A. Meyer. *Molecular Biology Reports* **37**, 3445–3454.
- Sauer M, Kleine-Vehn J.** 2011. AUXIN BINDING PROTEIN1: the outsider. *The Plant Cell* **23**, 2033–2043.
- Savatini DV, Ferrari S, Sicilia F, Lorenzo G.** 2011. Oligogalacturonide–auxin antagonism does not require post-transcriptional gene silencing or stabilization of auxin response repressors in *Arabidopsis*. *Plant Physiology* **157**, 1163–1174.
- Schacht T, Unger C, Pich A, Wydra K.** 2011. Endo- and exopolygalacturonases of *Ralstonia solanacearum* are inhibited by polygalacturonase-inhibiting protein (PGIP) activity in tomato stem extracts. *Plant Physiology and Biochemistry* **49**, 377–387.
- Schaller A, Stintzi A, Graff L.** 2012. Subtilases—versatile tools for protein turnover, plant development, and interactions with the environment. *Physiologia Plantarum* **145**, 52–66.
- Scheller HV, Ulvskov P.** 2010. Hemicelluloses. *Annual Review of Plant Biology* **61**, 263–289.
- Schols H, Voragen A.** 1994. Occurrence of pectic hairy regions in various plant cell wall materials and their degradability by rhamnogalacturonase. *Carbohydrate Research* **256**, 83–95.
- Seco R, Penuelas J, Filella I.** 2007. Short-chain oxygenated VOCs: emission and uptake by plants and atmospheric sources, sinks, and concentrations. *Atmospheric Environment* **41**, 2477–2499.
- Sexton TR, Henry RJ, Harwood CE, Thomas DS, McManus LJ, Raymond C, Henson M, Shepherd M.** 2012. Pectin methyl esterase genes influence solid wood properties of *Eucalyptus pilularis*. *Plant Physiology* **158**, 531–541.
- Seyedarabi A, To TT, Ali S, Hussain S, Fries M, Madsen R, Clausen MH, Teixeira S, Brocklehurst K, Pickersgill RW.** 2010. Structural insights into substrate specificity and the anti beta-elimination mechanism of pectate lyase. *Biochemistry* **49**, 539–546.
- Sharma N, Rathore M, Sharma M.** 2013. Microbial pectinase: sources, characterization and applications. *Reviews in Environmental Science and Biotechnology* **12**, 45–60.
- Shen Z, Denton M, Mutti N, Pappan K, Kanost M, Reese J, Reeck G.** 2003. Polygalacturonase from *Sitophilus oryzae*: possible horizontal transfer of a pectinase gene from fungi to weevils. *Journal of Insect Science* **3**, 1–9.
- Shen Z, Pappan K, Mutti NS, He Q-J, Denton M, Zhang Y, Kanost MR, Reese JC, Reeck GR.** 2005. Pectinmethylesterase from the rice weevil, *Sitophilus oryzae*: cDNA isolation and sequencing, genetic origin, and expression of the recombinant enzyme. *Journal of Insect Science* **5**, 1–9.
- Shevchik V, Hugouvieux-Cotte-Pattat N.** 1997. Identification of a bacterial pectin acetyl esterase in *Erwinia chrysanthemi* 3937. *Molecular Microbiology* **24**, 1285–1301.
- Shevchik EV, Hugouvieux-Cotte-Pattat N.** 2003. PaeX, a second pectin acetyl esterase of *Erwinia chrysanthemi* 3937. *Journal of Bacteriology* **185**, 3091–3100.
- Shi H, Zhu L, Zhou Y, Li G, Chen L, Li X.** 2009. A cotton gene encoding a polygalacturonase inhibitor-like protein is specifically expressed in petals. *Acta Biochimica et Biophysica Sinica* **41**, 316–324.
- Shibuya N, Minami E.** 2001. Oligosaccharide signalling for defence responses in plant. *Physiological and Molecular Plant Pathology* **59**, 223–233.
- Shishova M, Lindberg S.** 2010. A new perspective on auxin perception. *Journal of Plant Physiology* **167**, 417–422.
- Sicilia F, Fernandez-Recio J, Caprari C, De Lorenzo G, Tsernoglou D, Cervone F, Federici L.** 2005. The polygalacturonase-inhibiting protein PGIP2 of *Phaseolus vulgaris* has evolved a mixed mode of inhibition of endopolygalacturonase PG1 of *Botrytis cinerea*. *Plant Physiology* **139**, 1380–1388.
- Siedlecka A, Wiklund S, Peronne M, Micheli F, Lesniewska J, Sethson I, Edlund U, Richard L, Sundberg B, Mellerowicz EJ.** 2007. Pectin methyl esterase inhibits intrusive and symplastic cell growth in developing wood cells of *Populus*. *Plant Physiology* **146**, 554–565.
- Sinitsyna OA, Fedorova EA, Semenova MV, Gusakov AV, Sokolova LM, Bubnova TM, Okunev ON, Chulkin AM, Vavilova EA, Vinetsky YP.** 2007. Isolation and characterization of extracellular pectin lyase from *Penicillium canescens*. *Biochemistry* **72**, 565–571.

- Spadoni S, Zobotina O, Di Matteo A, Mikkelsen JD, Cervone F, De Lorenzo G, Mattei B, Bellincampi D.** 2006. Polygalacturonase-inhibiting protein interacts with pectin through a binding site formed by four clustered residues of arginine and lysine. *Plant Physiology* **141**, 557–564.
- Srivastava S, Gupta SM, Sane AP, Nath P.** 2012. Isolation and characterization of ripening related pectin methyltransferase inhibitor gene from banana fruit. *Physiology and Molecular Biology of Plants* **18**, 191–195.
- Sun L, Van Nocker S.** 2010. Analysis of promoter activity of members of the PECTATE LYASE-LIKE (PLL) gene family in cell separation in *Arabidopsis*. *BMC Plant Biology* **10**, 1–13.
- Sun Y, Fan X, Cao D, et al.** 2010. Integration of brassinosteroid signal transduction with the transcription network for plant growth regulation in *Arabidopsis*. *Developmental Cell* **19**, 765–777.
- Swain S, Kay P, Ogawa M.** 2011. Preventing unwanted breakups: using polygalacturonases to regulate cell separation. *Plant Signaling and Behavior* **6**, 93–97.
- Swarup K, Benková E, Swarup R, et al.** 2008. The auxin influx carrier LAX3 promotes lateral root emergence. *Nature Cell Biology* **10**, 946–954.
- Szabo LJ, Bushnell WR.** 2001. Hidden robbers: the role of fungal haustoria in parasitism of plants. *Proceedings of the National Academy of Sciences, USA* **98**, 7654–7655.
- Taniguchi Y, Ono A, Sawatani M, Nanba M, Kohno K, Usui M, Kurimoto M, Matuhasi T.** 1995. *Cry j 1*, a major allergen of Japanese cedar pollen, has pectate lyase enzyme activity. *Allergy* **50**, 90–93.
- Tardy F, Nasser W, Robert-Baudouy J, Hugouvieux-Cotte-Pattat N.** 1997. Comparative analysis of the five major *Erwinia chrysanthemi* pectate lyases: enzyme characteristics and potential inhibitors. *Journal of Bacteriology* **179**, 2503–2511.
- Terao A, Hyodo H, Satoh S, Iwai H.** 2013. Changes in the distribution of cell wall polysaccharides in early fruit pericarp and ovule, from fruit set to early fruit development, in tomato (*Solanum lycopersicum*). *Journal of Plant Research* **126**, 719–728.
- Themmen AP, Tucker GA, Grierson D.** 1982. Degradation of isolated tomato cell walls by purified polygalacturonase *in vitro*. *Plant Physiology* **69**, 122–124.
- Thibault JJ, Mercier C.** 1979. *Aspergillus niger* endopolygalacturonase: 2. Characterization and some properties. *Journal of Food Biochemistry* **2**, 379–393.
- Thonar C, Liners F, Van Cutsem P.** 2006. Polymorphism and modulation of cell wall esterase activities in the chicory root during the growing season. *Journal of Experimental Botany* **57**, 81–89.
- Tian G, Chen M, Zaltsman A, Citovsky V.** 2006. Pollen-specific pectin methyltransferase involved in pollen tube growth. *Developmental Biology* **294**, 83–91.
- Tjallingii WF, Esch TH.** 1993. Fine structure of aphid stylet routes in plant tissues in correlation with EPG signals. *Physiological Entomology* **18**, 317–328.
- Tromas A, Tromas R, Paponov I, Perrot-Rechenmann C.** 2010. AUXIN BINDING PROTEIN 1: functional and evolutionary aspects. *Trends in Plant Science* **15**, 436–446.
- Turquois T, Rinaudo M, Taravel F, Heyraud A.** 1999. Extraction of highly gelling pectic substances from sugar beet pulp and potato pulp: influence of extrinsic parameters on their gelling properties. *Food Hydrocolloids* **13**, 255–262.
- Tyler L, Bragg JN, Wu J, Yang X, Tuskan GA, Vogel JP.** 2010. Annotation and comparative analysis of the glycoside hydrolase genes in *Brachypodium distachyon*. *BMC Genomics* **11**, 1–21.
- Uehara T, Sugiyama S, Masuta C.** 2007. Comparative serial analysis of gene expression of transcript profiles of tomato roots infected with cyst nematode. *Plant Molecular Biology* **63**, 185–194.
- Underwood W.** 2012. The plant cell wall: a dynamic barrier against pathogen invasion. *Frontiers in Plant Science* **3**, 85.
- Vallarino JG, Osorio S.** 2012. Signaling role of oligogalacturonides derived during cell wall degradation. *Plant Signaling and Behavior* **7**, 1447–1449.
- Vanneste S, De Rybel B, Beemster GT, et al.** 2005. Cell cycle progression in the pericycle is not sufficient for SOLITARY ROOT/IAA14-mediated lateral root initiation in *Arabidopsis thaliana*. *The Plant Cell* **17**, 3035–3050.
- Van Alebeek G, van Scherpenzeel K, Beldman G, Schols H, Voragen A.** 2003. Partially esterified oligogalacturonides are the preferred substrates for pectin methyltransferase of *Aspergillus niger*. *Biochemical Journal* **372**, 211–218.
- Van Brussel AAN, Bakhuizen R, van Spronsen PC, Spaik HP, Tak T, Lugtenberg BJJ, Kijne JW.** 1992. Induction of pre-infection thread structures in the leguminous host plant by mitogenic lipo-oligosaccharides of *Rhizobium*. *Science Signaling* **257**, 70–72.
- Van Pouderooyen G, Snijder H, Benen J, Dijkstra B.** 2003. Structural insights into the processivity of endopolygalacturonase I from *Aspergillus niger*. *FEBS Letters* **554**, 462–466.
- Van Santen Y, Benen J, Schröter K, Kalk K, Armand S, Visser J, Dijkstra B.** 1999. 1.68-Å Crystal structure of endopolygalacturonase II from *Aspergillus niger* and identification of active site residues by site-directed mutagenesis. *Journal of Biological Chemistry* **274**, 30474–30480.
- Vercauteren I, de Almeida Engler J, De Groot R, Gheysen G.** 2002. An *Arabidopsis thaliana* pectin acetyltransferase gene is upregulated in nematode feeding sites induced by root-knot and cyst nematodes. *Molecular Plant-Microbe Interactions* **15**, 404–407.
- Verlent I, Hendrickx M, Verbeyst L, van Loey A.** 2007. Effect of temperature and pressure on the combined action of purified tomato pectinmethyltransferase and polygalacturonase in presence of pectin. *Enzyme and Microbial Technology* **40**, 1141–1146.
- Verlent I, Smout C, Duvetter T, Hendrickx ME, Van Loey A.** 2005. Effect of temperature and pressure on the activity of purified tomato polygalacturonase in the presence of pectins with different patterns of methyl esterification. *Innovative Food Science and Emerging Technologies* **6**, 293–303.
- Verlent I, Van Loey A, Smout C, Duvetter T, Hendrickx ME.** 2004. Purified tomato polygalacturonase activity during thermal and high-pressure treatment. *Biotechnology and Bioengineering* **86**, 63–71.
- Vernoux T, Besnard F, Traas J.** 2010. Auxin at the shoot apical meristem. *Cold Spring Harbor Perspectives in Biology* **2**, 1–14.
- Veronico P, Melillo MT, Saponaro C, Leonetti P, Picardi E, Jones JT.** 2011. A polygalacturonase-inhibiting protein with a role in pea defence against the cyst nematode *Heterodera goettingiana*. *Molecular Plant Pathology* **12**, 275–287.
- Videcoq P, Garnier C, Robert P, Bonnin E.** 2011. Influence of calcium on pectin methyltransferase behaviour in the presence of medium methylated pectins. *Carbohydrate Polymers* **86**, 1657–1664.
- Vincken JP, Schols HA, Oomen RJFJ, McCann MC, Ulvskov P, Voragen AGJ, Visser RGF.** 2003. If homogalacturonan were a side chain of rhamnogalacturonan I. Implications for cell wall architecture. *Plant Physiology* **132**, 1781–1789.
- Vogel J.** 2008. Unique aspects of the grass cell wall. *Current Opinion in Plant Biology* **11**, 301–307.
- Vogel J, Raab T, Schiff C, Somerville S.** 2002. *PMR6*, a pectate lyase-like gene required for powdery mildew susceptibility in *Arabidopsis*. *The Plant Cell* **14**, 2095–2106.
- Voiniciuc C, Dean GH, Griffiths JS, Kirchsteiger K, Hwang YT, Gillett A, Dow G, Western TL, Estelle M, Haughn GW.** 2013. FLYING SAUCER1 is a transmembrane RING E3 ubiquitin ligase that regulates the degree of pectin methylesterification in *Arabidopsis* seed mucilage. *The Plant Cell* **25**, 944–959.
- Volpi C, Janni M, Lionetti V, Bellincampi D, Favaron F, D'Ovidio R.** 2011. The ectopic expression of a pectin methyl transferase inhibitor increases pectin methyl esterification and limits fungal diseases in wheat. *Molecular Plant-Microbe Interactions* **24**, 1012–1019.
- Von Dahl CC, Hävecker M, Schlögl R, Baldwin IT.** 2006. Caterpillar-elicited methanol emission: a new signal in plant-herbivore interactions? *The Plant Journal* **46**, 948–960.
- Wang H, Guo Y, Lv F, Zhu H, Wu S, Jiang Y, Li F, Zhou B, Guo W, Zhang T.** 2010. The essential role of *GhPEL* gene, encoding a pectate lyase, in cell wall loosening by depolymerization of the de-esterified pectin during fiber elongation in cotton. *Plant Molecular Biology* **72**, 397–406.
- Wen B, Ström A, Tasker A, West G, Tucker GA.** 2013. Effect of silencing the two major tomato fruit pectin methyltransferase isoforms on cell wall pectin metabolism. *Plant Biology* **15**, 1025–1032.
- Whitham S, Quan S, Chang H, Cooper B, Estes B, Zhu T, Wang X, Hou Y.** 2003. Diverse RNA viruses elicit the expression of common sets

of genes in susceptible *Arabidopsis thaliana* plants. *The Plant Journal* **33**, 271–283.

**Wi SG, Singh AP, Lee KH, Kim YS.** 2005. The pattern of distribution of pectin, peroxidase and lignin in the middle lamella of secondary xylem fibres in alfalfa (*Medicago sativa*). *Annals of Botany* **95**, 863–868.

**Wiethölter N, Graebner B, Mierau M, Mort AJ, Moerschbacher BM.** 2003. Differences in the methyl ester distribution of homogalacturonans from near-isogenic wheat lines resistant and susceptible to the wheat stem rust fungus. *Molecular Plant-Microbe Interactions* **16**, 945–952.

**Willats WG, Knox JP, Mikkelsen JD.** 2006. Pectin: new insights into an old polymer are starting to gel. *Trends in Food Science and Technology* **17**, 97–104.

**Willats WGT, McCartney L, Mackie W, Knox JP.** 2001. Pectin: cell biology and prospects for functional analysis. *Plant Molecular Biology* **47**, 9–27.

**Williamson G.** 1991. Purification and characterization of pectin acetyltransferase from orange peel. *Phytochemistry* **30**, 445–449.

**Wojtasik W, Kulma A, Kostyn K, Szopa J.** 2011. The changes in pectin metabolism in flax infected with *Fusarium*. *Plant Physiology and Biochemistry* **49**, 862–872.

**Wolf S, Hématy K, Höfte H.** 2012a. Growth control and cell wall signaling in plants. *Annual Review of Plant Biology* **63**, 381–407.

**Wolf S, Mouille G, Pelloux J.** 2009b. Homogalacturonan methylesterification and plant development. *Molecular Plant* **2**, 851–860.

**Wolf S, Mravec J, Greiner S, Mouille G, Höfte H.** 2012b. Plant cell wall homeostasis is mediated by brassinosteroid feedback signaling. *Current Biology* **22**, 1732–1737.

**Wolf S, Rausch T, Greiner S.** 2009a. The N-terminal pro region mediates retention of unprocessed type-I PME in the Golgi apparatus. *The Plant Journal* **58**, 361–375.

**Worden N, Park E, Drakakaki G.** 2012. Trans-golgi network—an intersection of trafficking cell wall components. *Journal of Integrative Plant Biology* **54**, 875–886.

**Wydra K, Beri H.** 2006. Structural changes of homogalacturonan, rhamnogalacturonan I and arabinogalactan protein in xylem cell walls of

tomato genotypes in reaction to *Ralstonia solanacearum*. *Physiological and Molecular Plant Pathology* **68**, 41–50.

**Wyss U, Zunke U.** 1986. Observations on the behaviour of second stage juveniles of *Heterodera schachtii* inside host roots. *Revue de Nématologie* **9**, 153–165.

**Xie F, Murray JD, Kim J, Heckmann AB, Edwards A, Oldroyd GED, Downie JA.** 2012. Legume pectate lyase required for root infection by rhizobia. *Proceedings of the National Academy of Sciences, USA* **109**, 633–638.

**Yadav S, Yadav P, Yadav D, Yadav K.** 2009. Pectin lyase: a review. *Process Biochemistry* **44**, 1–10.

**Yang C, Guo R, Jie F, Nettleton D, Peng J, Carr T, Yeakley J, Fan J, Whitham S.** 2007. Spatial analysis of *Arabidopsis thaliana* gene expression in response to *Turnip mosaic virus* infection. *Molecular Plant-Microbe Interactions* **20**, 358–370.

**Yang XY, Zeng ZH, Yan JY, Fan W, Bian HW, Zhu MY, Yang JL, Zheng SJ.** 2012. Association of specific pectin methylesterases with AI-induced root elongation inhibition in rice. *Physiologia Plantarum* **148**, 502–511.

**Yoder M, Jurnak F.** 1995. The refined three-dimensional structure of pectate lyase C from *Erwinia chrysanthemi* at 2.2 angstrom resolution (implications for an enzymatic mechanism). *Plant Physiology* **107**, 349–364.

**Zhang GY, Feng J, Wu J, Wang XW.** 2010. BoPME11, a pollen-specific pectin methylesterase inhibitor, has an essential role in pollen tube growth. *Planta* **231**, 1323–1334.

**Zheng Y, Huang C, Liu W, Ko T, Xue Y, Zhou C, Guo R, Ma Y.** 2012. Crystal structure and substrate-binding mode of a novel pectate lyase from alkaliphilic *Bacillus* sp. N16-5. *Biochemical and Biophysical Research Communications* **420**, 269–274.

**Zonia L, Munnik T.** 2011. Understanding pollen tube growth: the hydrodynamic model versus the cell wall model. *Trends in Plant Science* **16**, 347–352.