

## HOMOTOPY IDEMPOTENTS ON FINITE- DIMENSIONAL COMPLEXES SPLIT<sup>1</sup>

HAROLD M. HASTINGS<sup>2</sup> AND ALEX HELLER

**ABSTRACT.** We prove that (unpointed) homotopy idempotents on finite-dimensional complexes split, and describe some geometric consequences.

**1. Introduction.** A *homotopy idempotent* on a space  $X$  is a map  $f: X \rightarrow X$  such that  $f^2 \simeq f$ . It is said to *split* if there are maps  $W \xrightarrow{u} X \xrightarrow{v} W$  with  $vu \simeq 1_W$ , and  $uv \simeq f$ . It is well known (see E. M. Brown [2], D. A. Edwards and R. Geoghegan [8], P. Freyd [10]) that *pointed* homotopy idempotents on *pointed connected CW complexes* split. Questions about splitting unpointed homotopy idempotents have arisen in several areas. In homotopy theory (Freyd and Heller [11], Heller [13]) this question is closely linked with Brown's representation theorem for half-exact functors. This question is also closely related with the study of FANR's (fundamental absolute neighborhood retracts). A compact metric space  $C$  is a FANR if it is a shape (fundamental) retract of a (compact) ANR (absolute neighborhood retract)  $A$ , that is, if there is a map  $j: C \rightarrow A$  and a shape map  $r: A \rightarrow C$  with  $rj$  shape equivalent to  $1_C$ . By J. West [16],  $A$  is homotopy equivalent to a finite complex  $X$ . Then, cf. [8], the composite mapping  $jr$  induces a homotopy idempotent on  $X$ . In pro-homotopy (D. A. Edwards and Hastings [9]), the question of splitting homotopy idempotents is a special case of the more general question of whether weak pro-homotopy equivalences are strong pro-homotopy equivalences. Similar questions arise in shape theory (J. Dydak and Hastings [6], Dydak and J. Segal [7], Edwards and R. Geoghegan [8]: Is every shape equivalence a strong shape equivalence?) and proper homotopy theory (T. A. Chapman [3], Chapman and L. Siebenmann [4, Appendix II], and Edwards and Geoghegan [8]: Is every weak proper homotopy equivalence a proper homotopy equivalence?).

Recently, Dydak and P. Minc [5], and Freyd and Heller [11] independently found an unpointed homotopy idempotent on an infinite-dimensional complex which does not split. See §2. This answered the pro-homotopy question in the negative. However, the shape and proper homotopy questions remained open, because they involve implicit finiteness restrictions.

We shall prove the following.

---

Received by the editors August 18, 1980 and, in revised form, April 20, 1981.

1980 *Mathematics Subject Classification.* Primary 55P65; Secondary 55P55, 55U35.

<sup>1</sup>These results were announced at the conference "Shape theory and pro-homotopy", Dubrovnik, Yugoslavia, January 1981 [12].

<sup>2</sup>Partially supported by NSF Grant MCS 81-02053.

©1982 American Mathematical Society  
0002-9939/82/0000-0173/\$02.00

**THEOREM 1.** *(Unpointed) homotopy idempotents on finite-dimensional complexes split.*

**COROLLARY.** *Every FANR is a pointed FANR.*

We need consider, without loss of generality, pointed, connected CW complexes. All maps, with such obvious exceptions as deck-transformations of covering spaces, preserve basepoints. We shall consider both pointed ( $\sim$ ) and unpointed ( $\simeq$ ) homotopies.

We thank J. Dydak and R. Geoghegan for helpful conversations.

**2. Proof of Theorem 1.** We begin by recalling an earlier result [5, 11]. Let  $f: X \rightarrow X$  be a homotopy idempotent, and let  $H: f^2 \simeq f$  be a homotopy. Let  $F$  be the group with presentation

$$\langle x_0, x_1, \dots \mid x_i x_j x_i^{-1} = x_{j+1}, \text{ all } i < j \rangle.$$

Then  $F$  admits an endomorphism  $\phi$  with  $\phi x_i = x_{i+1}$  for all  $i$ . Under the homotopy  $H$ , the basepoint of  $X$  traces out an element  $\xi_0$  in  $\pi_1 X$ . Then there is a unique homomorphism  $\Phi: F \rightarrow \pi_1 X$  such that  $\Phi x_0 = \xi_0$  and  $\Phi \phi = (\pi_1 f)\Phi$ .

**THEOREM 2** [5, 11]. *The homotopy idempotent  $f: X \rightarrow X$  fails to split if and only if  $\Phi$  is injective. In particular, the Eilenberg-Mac Lane space  $K(F, 1)$  has an unsplit homotopy idempotent, the induced map  $K(\phi, 1)$ .*

If a homotopy idempotent  $f$  splits, then  $f$  has an image. (Of course, images do not exist, in general in homotopy theory.) Here is a candidate for homotopy “image” of  $f$ . Let  $Y$  be the homotopy of the diagram colimit of the diagram

$$(2.1) \quad X \xrightarrow{f} X \xrightarrow{f} X \xrightarrow{f} \dots$$

For an explicit construction, let  $Y$  be the Milnor telescope [13], of (2.1), that is,

$$(2.2) \quad Y = \text{Tel} \left( X \xrightarrow{f} X \xrightarrow{f} X \rightarrow \dots \right) = M_f \cup_X M_f \cup_X M_f \cup_X \dots$$

Let  $i_k: X \rightarrow Y$  include  $X$  as the base of the  $k$ th mapping cylinder in  $Y$ . Then

$$(2.3) \quad \begin{aligned} & \text{(a) } i_{k+1} f \sim i_k, \\ & \text{(b) } g i_{k+1} \sim i_k, \text{ and} \\ & \text{(c) } \pi_* Y = \text{colim} \left( \pi_* X \xrightarrow{f_*} \pi_* X \xrightarrow{f_*} \dots \right), \\ & \text{(d) } H_* Y = \text{colim} \left( H_* X \xrightarrow{f_*} H_* X \xrightarrow{f_*} \dots \right). \end{aligned}$$

Thus for all  $q$ ,  $H_q Y = \bigcup_k \text{Im}(H_q i_k)$ . By (2.3)(b), all of the images  $\text{Im}(H_q i_k)$  are isomorphic. Thus unless  $H_q Y = 0$ , all  $\text{Im}(H_q i_k) \neq 0$ .

Similarly form

$$(2.4) \quad \tilde{Y} = \text{Tel} \left( \tilde{X} \xrightarrow{\tilde{f}} \tilde{X} \xrightarrow{\tilde{f}} \tilde{X} \rightarrow \dots \right).$$

Then, up to homotopy,  $\tilde{Y}$  is the universal cover of  $Y$  (compare  $\pi_* Y$  and  $\pi_* \tilde{Y}$ ) and analogues of (2.3) hold for  $\tilde{Y}$ . We consider the left action of fundamental groups on universal covering spaces and their homology. Then for any element  $\xi$  of  $\pi_1 X$ , the following equivariance conditions hold:

$$(2.5) \quad \tilde{f}\xi \simeq ((\pi_1 f)\xi)\tilde{f}; \quad \tilde{i}_k \xi \simeq ((\pi_1 \tilde{i}_k)\xi)\tilde{i}_k, \quad k = 0, 1, \dots$$

We may now proceed to the proof of the theorem. Let  $f: X \rightarrow X$  be a homotopy idempotent on a finite-dimensional complex. Assume that  $f$  does not split. First observe that by taking a suitable covering space we may assume that  $\pi_1 X = F$ ,  $\Phi = 1_F$ , and  $\phi = \pi_1 f$ . We may then compute, the homotopy “image” of  $f$  in (2.2) and (2.4):

$$(2.6) \quad \pi_1 Y = G = \langle \dots, x_{-1}, x_0, x_1, \dots \mid x_i x_j x_i^{-1} = x_{j+i}, \text{ all } i < j \rangle \quad \text{and} \\ (\pi_1 \tilde{i}_k)x_j = x_{j-k}, \quad j = 0, 1, \dots$$

For the deck-transformations on the universal coverings, (2.6) yields the formula  $\tilde{i}_k x_j \simeq x_{j-k} \tilde{i}_k$ .

The homotopy  $\tilde{f}^2 \simeq \tilde{f}$  lifts to a homotopy  $\tilde{f}^2 \simeq x_0 \tilde{f}$ . Thus for all  $k > 0$ ,  $\tilde{f}^{k+1} \simeq x_0 \tilde{f}^k$  and, composing with  $i_{k+1}$ ,  $\tilde{i}_0 \simeq x_{-k-1} \tilde{i}_1$ . Thus all  $x_l$ ,  $l \leq -2$ , operate in the same way on the subgroup  $\text{Im}(H_q \tilde{i}_1) \subset H_q Y$ . For each integer  $n > 0$ , let  $T_n$  be the subgroup of  $G$  generated by the  $n$  elements

$$(2.7) \quad x_{-3}^{-1} x_{-2}, x_{-6}^{-1} x_{-5}, \dots, x_{-3n}^{-1} x_{-3n+1}.$$

It is shown in [11, cf. also 6, 7] that  $T_n$  is free abelian on these  $n$  generators; we have just seen that it operates trivially on  $\text{Im}(H_q \tilde{i}_1)$ .

Because  $X$  is finite-dimensional, there is a largest degree  $r$  such that  $H_r \tilde{Y} \neq 0$ . Also, for each  $n$ , let  $Y_n$  be the covering space of  $Y$  with fundamental group  $T_n$ , and compute  $H^* Y_n$  by the spectral sequence of the covering  $\tilde{Y} \rightarrow Y_n$ ; cf. [14] for a detailed description. In this spectral sequence  $E_{pq}^2 = H_p(T_n, H_q Y)$ . Thus  $E_{pq}^2 \neq 0$  if  $p > n$  or  $q > r$ . Accordingly

$$(2.8) \quad E_{nr}^2 = E_{nr} = H_n(T_n, H_r Y) = (H_r Y)^{T_n},$$

where  $(H_r Y)^{T_n}$  denotes the subgroup fixed under  $T_n$ . The last equality may be obtained directly (cf. [14]) or using Poincaré duality in local coefficients (following a suggestion of the referee):

$$(2.9) \quad H_n(K(T_n, 1), H_r Y) \cong H^0(K(T_n, 1), H_r Y).$$

Thus

$$(2.10) \quad H_{n+r} Y_n = (H_r Y)^{T_n} = \text{Im}(H_r \tilde{i}_1) \neq 0, \quad n = 1, 2, \dots$$

Hence the dimension of  $Y$ , and hence also of  $X$ , is infinite, a contradiction. This completes the proof.  $\square$

**3. A geometric application.** We sketch a proof of the following, as an example of the use of Theorem 1 in geometric topology.

**THEOREM 3.** *Every (compact) FANR is a pointed FANR.*

PROOF. Let  $C$  be a (compact) FANR. Following [8], as explained in our Introduction,  $C$  is a shape retract a finite complex  $X$  via shape maps  $j: C \rightarrow X$  and  $r: X \rightarrow C$ . These yield a homotopy idempotent  $f: X \rightarrow X$ . By [8],  $f$  splits in weak pro-homotopy through the inverse sequence

$$(3.1) \quad X \xleftarrow{f} X \xleftarrow{f} X \xleftarrow{f} \dots,$$

which defines the shape of  $C$ . Also,  $C$  has the shape of a complex if and only if  $f$  splits in the homotopy category of complexes. (In this case,  $f$  splits through the homotopy limit,  $\text{holim}$ , of (3.1); see A. K. Bousfield and D. M. Kan [1], also [8], [9].) We provide a splitting in the unpointed case; Edwards and Geoghegan [8] had only been able to consider pointed shape and pointed FANR's. Our Theorem 1 extends the Edwards-Geoghegan argument to unpointed shape, thus  $C$  has the shape of a complex  $Y$ . Hence  $C$  has the strong shape of  $Y$ , and the pointed shape of  $Y$  [9, §5]. The conclusion now follows from [8].  $\square$

#### REFERENCES

1. A. K. Bousfield and D. M. Kan, *Homotopy limits, completions, and localizations*, Lecture Notes in Math., vol. 304, Springer-Verlag, Berlin and New York, 1973.
2. E. M. Brown, *Cohomology theories*, Ann. of Math. (2) **75** (1962), 467–484.
3. T. A. Chapman, *On some applications of infinite-dimensional manifolds to the theory of shape*, Fund. Math. **76** (1972), 181–193.
4. T. A. Chapman and L. Siebenmann, *Finding a boundary for a Hilbert cube manifold*, Acta Math. **139** (1976), 171–208.
5. J. Dydak, *A simple proof that pointed, connected FANR-spaces are regular fundamental retracts of ANR's*, Bull. Acad. Polon. Sci. Sér. Sci. Math. Astronom. Phys. **25** (1977), 55–62.
6. J. Dydak and H. M. Hastings, *Homotopy idempotents on two-dimensional complexes split*, Proc. Internat. Conf. on Geometric Topology (Warsaw, 1978), (ed. K. Borsuk and A. Kirkor), PWN, Warsaw, 1980, pp. 127–133.
7. J. Dydak and J. Segal, *Shape theory*, Lecture Notes in Math., vol. 688, Springer-Verlag, Berlin and New York, 1978.
8. D. A. Edwards and R. Geoghegan, *Shapes of complexes, ends of manifolds, homotopy limits, and the Wall obstruction*, Ann. of Math. (2) **101** (1975), 521–535; Correction **104** (1976), 379.
9. D. A. Edwards and H. M. Hastings, *Čech and Steenrod homotopy theory, with applications to geometric topology*, Lecture Notes in Math., vol. 542, Springer-Verlag, Berlin and New York, 1976.
10. P. Freyd, *Splitting homotopy idempotents*, Proc. Conf. on Categorical Algebra (La Jolla 1965), (ed. S. Eilenberg and G. M. Kelley), Springer, Berlin and New York, 1966, pp. 173–176.
11. P. Freyd and A. Heller, *Splitting homotopy idempotents* (to appear).
12. H. M. Hastings and A. Heller, *Splitting homotopy idempotents*, Conf. on Shape and Pro-Homotopy (Dubrovnik, 1981), (ed. S. Mardešić and J. Segal), Lecture Notes in Math., vol. 870, Springer-Verlag, Berlin and New York, 1981, pp. 23–36.
13. A. Heller, *On the representability of homotopy functors* (to appear).
14. P. J. Hilton and S. Wiley, *Homology theory: an introduction to algebraic topology*, Cambridge, 1962.
15. J. Milnor, *On axiomatic homology theory*, Pacific J. Math. **12** (1962), 337–341.
16. J. West, *Mapping Hilbert cube manifolds to ANR's: A solution to a conjecture of Borsuk*, Ann. of Math. (2) **106** (1977), 1–18.

DEPARTMENT OF MATHEMATICS, HOFSTRA UNIVERSITY, HEMPSTEAD, NEW YORK 11550

DEPARTMENT OF MATHEMATICS, GRADUATE SCHOOL & UNIVERSITY CENTER, CITY UNIVERSITY OF NEW YORK, 33 WEST 42 STREET, NEW YORK, NEW YORK 10036