

# Homozygous and compound heterozygous mutations at the Werner syndrome locus

Junko Oshima\*, Chang-En Yu<sup>1</sup>, Charles Piussan<sup>2</sup>, Georg Klein<sup>3</sup>, Jörg Jabkowski<sup>3</sup>, Sevim Balci<sup>4</sup>, Tetsuro Miki<sup>5</sup>, Jun Nakura<sup>5</sup>, Toshio Ogihara<sup>5</sup>, James Ells<sup>6</sup>, Marilia de A. C. Smith<sup>7</sup>, Maria I. Melaragno<sup>7</sup>, Marco Fraccaro<sup>8</sup>, Susi Scappaticci<sup>8</sup>, John Matthews<sup>9</sup>, Samir Ouais<sup>10</sup>, Amy Jarzebowicz, Gerard D. Schellenberg<sup>1</sup> and George M. Martin

Box 357470, Department of Pathology, University of Washington, Seattle, WA 98195, USA, <sup>1</sup>Geriatrics Research Education and Clinical Center, Veterans Affairs Puget Sound Health Care System, Seattle Division, Seattle, WA, USA, <sup>2</sup>Pediatric Genetics, University of Amiens, Amiens, 80054, France, <sup>3</sup>Dermatologische Abteilung, Allgem. öffentliche Krankenhaus der Elisabethinen, Linz, 239 Austria, <sup>4</sup>Department of Clinical Genetics, Hacettepe University Children's Hospital, Ankara, Turkey, <sup>5</sup>Department of Geriatric Medicine, Osaka University, Osaka, 565, Japan, <sup>6</sup>Internal Medicine Clinic, Nellis Air Force Base Hospital, Las Vegas, NV 89101, USA, <sup>7</sup>Division of Genetics, Escola Paulista de Medicina, São Paulo, 04023, Brazil, <sup>8</sup>Medical Genetics, University of Pavia, Pavia, Italy, <sup>9</sup>Group Health, Fairfax, VA 22039, USA and <sup>10</sup>Section of Endocrinology, Damascus City Hospital, Damascus, Syria

Received June 22, 1996; Revised and Accepted September 23, 1996

**The Werner syndrome (WS) is a rare autosomal recessive progeroid disorder. The Werner syndrome gene (*WRN*) has recently been identified as a member of the helicase family. Four distinct mutations were previously reported in three Japanese and one Syrian WS pedigrees. The latter mutation was originally described as a 4 bp deletion spanning a spliced junction. It is now shown that this mutation results in a 4 bp deletion at the beginning of an exon. Nine new *WRN* mutations in 10 additional WS patients, both Japanese and Caucasian, are described. These include three compound heterozygotes (one Japanese and two Caucasian). The new mutations are located all across the coding region.**

## INTRODUCTION

Werner syndrome (WS) is a rare autosomal recessive segmental progeroid syndrome (2). Patients exhibit not only an appearance of accelerated aging (premature graying, thinning of hair, skin atrophy and atrophy of subcutaneous fat), but also several disorders commonly associated with aging. These include bilateral cataracts, diabetes mellitus, osteoporosis, several forms of arteriosclerosis and a variety of benign and malignant neoplasms (3,4).

WS fibroblasts have very limited proliferative capacities as compared with age-matched controls (5–7). A prolongation of the S phase has been demonstrated both in WS fibroblasts and lymphoblastoid cell lines (8). Cultured cells exhibit a propensity


for chromosomal and intragenic mutations (9–12). The rate of repair of X-ray- or UV-damaged DNA appears to be normal in WS fibroblasts (13).

*WRN* was initially mapped to chromosome 8p (14,15). Physical and genetic maps of the region were constructed (16–19). *WRN* has recently been identified (GenBank accession number L76937) and four distinct *WRN* mutations were described (1). The *WRN* gene encodes a 1432 amino acid protein partially homologous to RecQ helicases (20). The *WRN* protein contains seven helicase motifs; two of them have been identified in all ATP-binding proteins (21).

DNA helicases have been implicated in a number of molecular processes. One of the most important functions of DNA helicases is the unwinding of DNA during DNA replication as a component in a replication complex (22–24). Another function of helicase involves DNA repair. It has been hypothesized that some forms of nucleotide excision repair are coupled with transcription; mutant helicases responsible for the DNA instability syndromes may impair lesion recognition and/or lesion removal of the damaged nucleotides during transcription (25–27). Examples include: *ERCC2* helicase, which complements xeroderma pigmentosum B and its yeast homologue *RAD3* (28,29); *ERCC3*, which complements xeroderma pigmentosum D and its yeast homologue *RAD25* (30–34); *ERCC3* and *ERCC6*, which complement a Cockayne syndrome mutation (35,36). In *Escherichia coli*, the *RecQ* helicase is involved in the initial step of DNA repair by recombination (37).

Helicases are required for accurate chromosomal segregation. In yeast, precise chromosome segregation requires *Sgs1*, a eukaryotic homologue of *RecQ* (38).

\*To whom correspondence should be addressed



**Figure 1.** Locations of the *WRN* mutations. The rectangular box indicates the predicted *WRN* protein. The light shadowed segment indicates the highly acidic repeat region; dark shadows indicate the locations of the helicase consensus motifs. The locations of the *WRN* mutations are grouped based upon the type of mutation and are shown underneath the *WRN* protein along with Registry codes. Parentheses indicate the heterozygous mutations. ZM and MH mutations were previously described (1).

Given the several potential roles of the *WRN* protein, a careful delineation of spontaneous mutations at this locus could facilitate the characterizations of its functions. We report nine *WRN* gene mutations from 10 WS patients, three of which were compound heterozygotes. Two other mutations found in three patients have been previously reported (1). These various mutations involve sequences throughout the coding region.

## RESULTS

Four Japanese and eight non-Japanese WS patients were selected from our International Registry. Six of them (AUS, KO, MIM3, SEP, TUR, UH) were classified as 'definite WS' and three (LGS, OW, SUG) as 'probable WS'. Clinical and laboratory data for members of BLS, KUN and SYR remain incomplete, but the affected subjects had been diagnosed as WS by the submitting physicians.

Three new mutations were found in regions N-terminal with respect to the helicase consensus motifs. The point mutation at nt 1336, CGA (Arg) to TGA (Stp), was found as a homozygous mutation in one Caucasian (LGS) and two consanguineous Japanese (OW, KO) WS subjects and as a heterozygous mutation in one Japanese WS subject (KUN). LGS denied consanguinity; non-consanguinity was supported by haplotype data (19). A single nucleotide deletion at 1194–1196, AAA to AA, was seen as a heterozygous mutation in AUS. This mutation would create a frameshift which ends at 1406–1408 TGA (Stp). A four nucleotide insertion (ATCT) between 1509 and 1520 was homozygous in MIM3. This frameshift mutation would terminate at 1535–1537 TGA (Stp).

Three mutations were found within or just 3' to the helicase motifs in two Caucasian patients. One (SEP) mutation was a 105 bp insertion between 2319 and 2320. The insertion results in a termination codon, creating a truncated protein that excludes helicase domains III and the subsequent C terminus of the *WRN* protein. A second mutation was a deletion of nucleotide 2320–3056 seen in SUG as a heterozygous mutation, terminating at nt 3081–3083 TGA (Stp). The third mutation was a heterozygous termination mutation found in SUG, located 30 amino acids after the last helicase motif.

Three new mutations were found in regions C-terminal to the helicase motifs. A Japanese patient, IB, was homozygous for an A deletion at nt 3677. The mutated protein stops at nt 3713–3715 TAG (Stp). BLS (French) and TUR (Turkish) patients shared the same mutation at nt 3724, CGA (Gln) to TGA (Stp), which was

previously found in the Japanese SY family (1). A 74 bp deletion of nt 3541–3614 was seen as a heterozygous mutation in a Japanese WS, KUN. This deletion results in a termination at 3720–3722 TAG (Stp). A 113 bp deletion of nt 3691–3803, which would result in a termination at nt 3816–3818 TGA (Stp), was found as a heterozygous deletion in the Caucasian WS, AUS.

These mutations were confirmed by sequencing of genomic PCR products, using the primers from the intron sequences of *WRN* (39). A summary of the newly discovered mutations is given in Figure 1.

The mutation in the SYR pedigree was previously reported as a 4 bp deletion at the intron–exon boundary, 2 bp from the putative intron and 2 bp from the contiguous exon (gtagACA-GACC at the DNA level). This was expected to cause an in-frame deletion of the exon. Our RT–PCR protocol, however, showed a deletion of 4 bp, ACAG, from the beginning of this exon. The ACAG deletion would result in a termination at nt 3971–3973 TAG (Stp).

## DISCUSSION

In our original report of the positional cloning of the *WRN* locus, four distinct homozygous mutations in the 3' region of the *WRN* gene were described (1). Using the present RT–PCR strategy mutations were readily found in various locations within the gene. The biochemical consequences of these mutations are not known.

All of the *WRN* mutations we have found to date either create a stop codon mutation or cause frameshifts that lead to premature terminations. We have not yet found an amino acid substitution in *WRN* that seems to be responsible for the pathogenesis of WS. It is quite possible that the various truncated *WRN* proteins may be rapidly degraded, resulting in comparable null mutations and comparable phenotypes. Such altered mRNAs are thought to be degraded via a specific pathway (40). In preliminary experiments, we do observe evidence for reduced levels of *WRN* mRNA expression in WS LCLs with four different mutations.

Identical mutations were found across a variety of ethnic groups, raising the question of potential mutationally susceptible sequences. Although the total number of mutations so far found in the *WRN* protein is not extensive, candidate sequences for such susceptibility would include nt 3677–3920, nt 1336–1395 and nt 2319–2320.

Three instances of compound heterozygous mutations were found: KUN (Japanese), AUS (Caucasian) and SUG (Caucasian). There have been numerous reports of compound heterozygous

mutations in 'disease genes' (41,42). However, comparatively few compound heterozygotes have been reported in the genomic instability syndromes. Given the comparatively low prevalence of consanguinity in the USA, clinicians should therefore be alert to the diagnosis of WS in the absence of a history of consanguinity. Our experience suggests that WS is underdiagnosed in the USA.

## MATERIALS AND METHODS

### Samples

WS patients were from an International Registry of Werner Syndrome (George M. Martin, MD, Junko Oshima, MD, PhD, Amy Jarzebowicz, BS). Diagnostic criteria were previously described (18). This study was approved by the University of Washington Institutional Review Board.

### RT-PCR

Five µg of poly(A) RNA, isolated from total RNA, using Oligotex (Qiagen Inc.) was reverse-transcribed with random hexamers in 100 µl reaction volume with GeneAmp RNA PCR kit (Perkin Elmer Cetus). Two µl of the RT product were amplified in a 50 µl PCR reaction buffer containing 5 units *Taq* DNA polymerase, 10 mM Tris-HCl (pH 8.3), 50 mM KCl, 1.5 mM MgCl<sub>2</sub>, 50 µM each of dGTP, dATP, dTTP and dCTP. The cycle program was typically: 94°C for 5 min, then 94°C for 45 s, 55°C for 45 s, 72°C for 3.5 min with 2 s increase per cycle for 35 cycles, followed by 72°C for 10 min. Five µl aliquots of the first amplification products were subjected to a nested second amplification in 100 µl reaction volumes. The primer sequences for RT-PCR are listed in Table 1. The secondary PCR products were separated on 1% agarose/1×TBE (100 mM Tris-HCl pH 8.0, 90 mM boric acid and 1 mM ethylenediaminetetraacetic acid) to estimate the concentrations of DNA before sequencing.

**Table 1.** Primer sequences for the RT-PCR sequencing template

Region of the amplification	1st amplification primers (5' to 3')	2nd amplification primers (5' to 3')	Size of PCR product
5' end	GTGGTGGCGCTCCACAGTCATCC CTTTATGAAGCCAATTTCTACCC	AAGACCTGTTGGACTGGATCTTCTC TACTCCAAAATCTCTAAATTTCCGG	838
Translation start site to helicase region	GTGGTGGCGCTCCACAGTCATCC CTTTATGAAGCCAATTTCTACCC	TAGGACTTTCAAAGATGAGTG CGTATACAATCCGGTATTACC	1936
Helicase region	GTGGTGGCGCTCCACAGTCATCC CTTTATGAAGCCAATTTCTACCC	AGATGTACTTTGGCCATTCCAG GCAATGATCCAATCTGGACC	1218
3' region	GCATTAATAAAGCTGACATTCGCC CGGAAGGCTGATTTAAGATGCC	CATTACGGTGCTCCTAAGGACATG CGGAAGGCTGATTTAAGATGCC	1946

**Table 2.** *WRN* mutations in Japanese and Caucasian WS patients

Registry no.	Country	Ethnicity	M/F	Location	Mutation	Predicted protein
LGS90610	USA	Caucasian	F	1336	CGA-TGA Arg Stp	368
OW90650	Japan	Japanese	M	1336	CGA-TGA Arg Stp	368
KO90375	Japan	Japanese	M	1336	CGA-TGA Arg Stp	368
KUN9001	Japan	Japanese	M	1336	CGA-TGA Arg Stp	368
AUS40025	Austria	Caucasian	M	3541-3614 1395	Deletion A deletion	1138 391
MIM37100	Brazil	Caucasian	F	3691-3803 1509	Deletion ATCT insertion	1157 429
SEP9000	Sardinia	Caucasian	F	2319-2320	105 bp insertion	708
SUG17802	USA	Caucasian	M	2320-3056 2896	Deletion CGA-TGA Arg Stp	704 888
IB90550	Japan	Japanese	F	3677	A deletion	1160
BLS60010	France	Caucasian	M	3724	CAG-TAG Gln Stp	1164
TUR90010	Turkey	Caucasian	M	3724	CAG-TAG Gln Stp	1164
SYR10006	Syria	Syrian	M	3919-3922	ACAG deletion	1245

## Direct sequencing of PCR products

RT-PCR products were sequenced using a T7 sequence PCR product sequencing kit (UBS, Amersham Life Science, Inc.). Seven  $\mu$ l of PCR product was pretreated with 15 U of exonuclease I and 1.5 U of shrimp alkaline phosphatase at 37°C for 15 min followed by inactivation of the enzymes at 80°C for 15 min, then mixed with 100 ng of sequencing primers. The sequencing reaction followed the manufacturer's instructions.

The sequencing gel contained 6.6% LongRanger polyacrylamide (J. T. Baker Inc.), 6 M urea and 1.2 $\times$  TBE. The running buffer contained 0.6 $\times$  TBE. The gel was run at 55 W, dried and exposed overnight to Biomax MR film (Eastman Kodak Co.).

## ACKNOWLEDGEMENTS

We thank Dr Goberdhan P. Dimri for a human cDNA library, an important contribution leading to the original positional cloning of *WRN*. We also thank Annette Smith, Charles E. Ogburn, Thao Dang, Susan Fredell, Ellen Nemens and Deanne Sparlin for their technical support. This work was supported by National Institute on Aging grants P1 AG08303 (GMM), R37 AG08303 (GMM), T32 AG00057 (GMM), RO1 AG12019 (GDS) and CNPq and FAPESP, Brazil (MIM).

## ABBREVIATIONS

WS, Werner syndrome; *WRN*, Werner syndrome gene; UV, ultraviolet; ERCC, excision repair-cross-complementing; LCL, lymphoblastoid cell line; RT-PCR, reverse transcription-polymerase chain reaction; PCR, polymerase chain reaction.

## REFERENCES

1. Yu, C.E., Oshima, J., Fu, Y.W., Hisama, F., Wijsman, E.M., Alisch, R., Matthews, S., Nakura, J., Miki, T., Ouais, S., Martin, G.M., Mulligan, J. and Schellenberg, G.D. (1996) Positional cloning of the Werner's syndrome gene. *Science*, **272**, 258–262.
2. Martin, G.M. (1978) Genetic syndromes in man with potential relevance to the pathology of aging. *Birth Defects*, **14**, 5–39.
3. Epstein, C.J., Martin, G.M., Schultz, A.L. and Motulsky, A.G. (1966) Werner's syndrome: a review of its symptomatology, natural history, pathologic features, genetics and relationships to the natural aging process. *Medicine*, **45**, 172–221.
4. Tollefsbol, T.O. and Cohen, H.J. (1984) Werner's syndrome: an underdiagnosed disorder resembling premature aging. *Age*, **7**, 75–88.
5. Martin, G.M., Sprague, C.A. and Epstein, C.J. (1970) Replicative life-span of cultivated human cells. Effects of donor's age, tissue and genotype. *Lab. Invest.*, **23**, 86–92.
6. Salk, D., Bryant, E., Hoehn, H., Johnston, P. and Martin, G.M. (1985) Growth characteristics of Werner syndrome cells *in vitro*. *Adv. Exp. Med. Biol.*, **190**, 305–311.
7. Kill, I.R., Faragher, R.G., Lawrence, K. and Shall, S. (1994) The expression of proliferation-dependent antigens during the lifespan of normal and progeroid human fibroblasts in culture. *J. Cell Sci.*, **107**, 571–579.
8. Poot, M., Hoehn, H., Runger, T.M. and Martin, G.M. (1992) Impaired S-phase transit of Werner syndrome cells expressed in lymphoblastoid cell lines. *Exp. Cell Res.*, **202**, 267–273.
9. Hoehn, H., Bryant, E.M., Au, K., Norwood, T.H., Boman, H. and Martin, G.M. (1975) Variegated translocation mosaicism in human skin fibroblast culture. *Cytogenet. Cell. Genet.*, **15**, 282–298.
10. Fukuchi, K., Martin, G.M. and Monnet, R.J. (1989) Mutator phenotype of Werner syndrome is characterized by extensive deletion. *Proc. Natl Acad. Sci. USA*, **86**, 5893–5897.
11. Fukuchi, K., Tayama, K., Kumahara, Y., Murano, K., Pride, A.B., Martin, G.M., Monnet, R.J. (1990) Increased frequency of 6-thioguanine-resistant peripheral blood lymphocytes in Werner syndrome patients. *Hum. Genet.*, **84**, 249–252.
12. Runger, T.M., Bauer, C., Dekant, B., Moller, K., Sobotta, P., Czerny, C., Poot, M. and Martin, G.M. (1994) Hypermutable ligation of plasmid DNA ends in cells from patients with Werner syndrome. *J. Invest. Dermatol.*, **102**, 45–48.
13. Fujiwara, Y., Higashikawa, T. and Tatsumi, M. (1977) A retarded rate of DNA replication and normal level of DNA repair in Werner's syndrome fibroblasts in culture. *J. Cell. Physiol.*, **92**, 365–374.
14. Goto, M., Rubenstein, M., Weber, J., Woods, K. and Drayna, D. (1992) Genetic linkage of Werner's syndrome to five markers on chromosome 8. *Nature*, **355**, 735–758.
15. Schellenberg, G.D., Martin, G.M., Wijsman, E.M., Nakura, J., Miki, T. and Ogihara, T. (1992) Homozygosity mapping and Werner syndrome. *Lancet*, **339**, 1002.
16. Yu, C.E., Oshima, J., Goddard, K.A.B., Miki, T., Nakura, J., Ogihara, T., Fraccaro, M., Puissan, C., Martin, G.M., Schellenberg, G.D. and Wijsman, E.M. (1994) Linkage disequilibrium and haplotype analysis of chromosome 8p11.1–21.1 markers and Werner syndrome. *Am. J. Hum. Genet.*, **55**, 356–364.
17. Oshima, J., Yu, C.E., Boehnke, M., Weber, J.L., Edelfhoff, S., Wagner, M.J., Wells, D.E., Wood, S., Disteshe, D.M., Martin, G.M. and Schellenberg, G.D. (1994) Integrated mapping analysis of the Werner syndrome region of chromosome 8. *Genomics*, **23**, 100–113.
18. Nakura, J., Wijsman, E.J., Miki, T., Kamino, K., Yu, C.E., Oshima, J., Fukuchi, K.I., Weber, J.L., Puissan, C., Melaragno, M.I., Epstein, C.J., Scappaticci, S., Fraccaro, M., Matusmura, T., Murano, S., Yoshida, S., Fujiwara, Y., Saida, T., Ogihara, T., Martin, G.M. and Schellenberg, G.D. (1994) Homozygosity mapping of the Werner syndrome locus (*WRN*). *Genomics*, **23**, 600–608.
19. Goddard, K.A.B., Yu, C.E., Oshima, J., Martin, G.M., Schellenberg, G.D. and Wijsman, E.M. (1996) Towards localization of the Werner syndrome gene by linkage disequilibrium and ancestral haplotyping: lessons learned from analysis of 35 chromosome 8p11.1–21.1 markers. *Am. J. Hum. Genet.*, **58**, 1286–1302.
20. Schmid, S.R. and Linder, P. (1992) D-E-A-D protein family of putative RNA helicases. *Mol. Microbiol.*, **6**, 283–292.
21. Gorbalenya, A.E., Koonin, E.V., Donchenko, A.P. and Blinov, V.M. (1989) Two related superfamilies of putative helicases involved in replication, recombination, repair and expression of DNA and RNA genome. *Nucleic Acids Res.*, **17**, 4713–4730.
22. Lohman, T.M. (1993) Helicase-catalyzed DNA unwinding. *J. Biol. Chem.*, **268**, 2269–2272.
23. Conaway, R.C. and Lehman, L.R. (1982) A DNA primase activity associated with DNA polymerase alpha from *Drosophila melanogaster* embryos. *Proc. Natl Acad. Sci. USA*, **79**, 2523–2527.
24. Dodson, M.S. and Lehman, I.R. (1991) Association of DNA helicase and primase activities with subassembly of the herpes simplex virus 1 helicase-primase composed of the UL5 and UL52 gene products. *Proc. Natl Acad. Sci. USA*, **88**, 1105–1109.
25. Hanawalt, P.C. (1994) Transcription-coupled repair and human disease. *Science*, **266**, 1957–1958.
26. Drapkin, R. and Reinberg, D. (1992) The essential twist. *Nature*, **369**, 523–533.
27. Bootsma, D., Weeda, G., Vermeulen, W., van Vuuren, H., Troelstra, C., van der Spek, P. and Hoeijmakers, J. (1995) Nucleotide excision repair syndrome: molecular basis and clinical symptoms. *Phil. Trans. R. Soc. London B. Biol. Sci.*, **347**, 75–81.
28. Weber, C.A., Salazer, E.P., Stewart, S.A. and Thompson, L.H. (1990) ERCC2: cDNA cloning and molecular characterization of a human nucleotide excision repair gene with high homology to yeast RAD3. *EMBO J.*, **9**, 1437–1447.
29. Guzder, S.N., Qiu, H., Sommers, C.H., Sung, P., Prakash, L. and Prakash, S. (1994a) DNA repair gene RAD3 of *S. cerevisiae* is essential for transcription by RNA polymerase II. *Nature*, **367**, 91–94.
30. Weeda, G., van Ham, R.C.A., Masurel, R., Westerveld, A., Odijk, H., de Wit, J., Bootsma, D., van del Eb, A.J. and Hoeijmakers, J.H.J. (1990a) Molecular cloning and biological characterization of the human excision repair gene ERCC-3. *Mol. Cell. Biol.*, **10**, 2570–2581.
31. Flejter, W.L., McDaniel, L.D., Johns, D., Friedberg, E.C. and Schults, R.A. (1992) Correction of xeroderma pigmentosum complementation group D mutant cell phenotypes by chromosome and gene transfer: involvement of human ERCC2 DNA repair gene. *Proc. Natl Acad. Sci. USA*, **89**, 261–265.

32. Sung, P., Bailly, V., Weber, C., Thompson, L.H., Prakash, L. and Prakash, S. (1993) Human xeroderma pigmentosum group D gene encodes a DNA helicase. *Nature*, **365**, 852–855.
33. Guzder, S.N., Sung, P., Bailly, V., Prakash, L. and Prakash, S. (1994) RAD25 is a DNA helicase required for DNA repair and RNA polymerase II transcription. *Nature*, **369**, 578–581.
34. Ma, L., Westbroek, A., Jochemsen, A.G., Weeda, G., Bosch, A., Bootsma, D., Hoeijmakers, J.H. and van der Eb, A.J. (1994) Mutational analysis of ECRR3, which is involved in DNA repair and transcription initiation: identification of domains essential for the DNA repair function. *Mol. Cell. Biol.*, **14**, 4126–4136.
35. Weeda, G., van Ham, R.C.A., Vermeulen, W., Bootsma, D., van der Eb, A.J. and Hoeijmakers, J.H.J. (1990) A presumed DNA helicase encoded by ERRC-3 is involved in the human repair disorders xeroderma pigmentosum and Cockayne's syndrome. *Cell*, **62**, 777–791.
36. Troelstra, C., van Gool, A., de Wit, J., Vermeulen, W., Bootsma, D. and Hoeijmaker, J.H.J. (1992) ECCR6, a member of a subfamily of putative helicases, is involved in Cockayne's syndrome and preferential repair of active gene. *Cell*, **71**, 939–953.
37. West, S.C. (1994) The processing of recombination intermediates: mechanistic insights from studies of bacterial protein. *Cell*, **76**, 9–15.
38. Watt, P.M., Louis, E.J., Borts, R.H., Hickson, I.D. (1995) Sgs1: A eukaryotic homolog of *E. coli* RecQ that interacts with topoisomerase II *in vivo* and is required for faithful chromosome segregation. *Cell*, **81**, 253–260.
39. Yu, C.E., Oshima, J., Wijsman, E. M., Nakura, J., Miki, T., Puissan, C., Matthews, S., Martin, G.M., Schellenberg, G.D. and the Werner's Syndrome Collaborative Group. Mutations in the consensus helicase domains of the Werner's syndrome gene (submitted).
40. Jacobson, A. and Peltz, S.W. (1996) Interrelationships of the pathways of mRNA decay and translation in eukaryotic cells. *Annu. Rev. Biochem.*, **65**, 693–739.
41. Satokata, I., Tanaka, K. and Okada, Y. (1992) Molecular basis of group A xeroderma pigmentosum: a missense mutation and two deletions located in a zinc finger consensus sequence of the XPAC gene. *Hum. Genet.*, **88**, 603–607.
42. Sato, M., Nishigori, C., Yagi, T. and Tanabe, H. (1996) Aberrant splicing and truncated-protein expression due to a newly identified XPA gene mutation. *Mutat. Res.*, **362**, 199–208.