

 Open access • Journal Article • DOI:10.1088/0951-7715/4/4/003

## Hopf bifurcation on a square lattice — [Source link](#)

Mary Silber, Edgar Knobloch

**Institutions:** University of Minnesota

**Published on:** 01 Nov 1991 - Nonlinearity (IOP Publishing)

**Topics:** Saddle-node bifurcation, Bifurcation diagram, Transcritical bifurcation, Bifurcation theory and Hopf bifurcation

Related papers:

- [Hopf bifurcation with the symmetry of the square](#)
- [Singularities and groups in bifurcation theory](#)
- [Pattern selection in three-dimensional magnetoconvection](#)
- [Symmetry-breaking Hopf bifurcation in anisotropic systems](#)
- [Symmetries and pattern selection in Rayleigh-Bénard convection](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/hopf-bifurcation-on-a-square-lattice-2o1llwbto5>

**HOPF BIFURCATION  
ON A SQUARE LATTICE**

By

**Mary Silber  
and  
Edgar Knobloch**

**IMA Preprint Series # 675**

July 1990

# HOPF BIFURCATION ON A SQUARE LATTICE

MARY SILBER\* and EDGAR KNOBLOCH†

**Abstract.** A complete classification of the generic  $D_4 \times_s T^2$ -equivariant Hopf bifurcation problems is presented. This bifurcation arises naturally in the study of extended systems, invariant under the Euclidean group  $E(2)$ , when a spatially uniform quiescent state loses stability to waves of wavenumber  $k \neq 0$  and frequency  $\omega \neq 0$ . The  $D_4 \times_s T^2$  symmetry group applies when periodic boundary conditions are imposed in two orthogonal horizontal directions. The center manifold theorem allows a reduction of the infinite dimensional problem to a bifurcation problem on  $C^4$ . In normal form, the vector field on  $C^4$  commutes with an  $S^1$  symmetry, which is interpreted as a time translation symmetry. The spatial and spatio-temporal symmetries of all possible solutions are classified in terms of isotropy subgroups of  $D_4 \times_s T^2 \times S^1$ . On a neighborhood of the Hopf bifurcation point, all small amplitude oscillatory solutions are found and their stabilities calculated relative to perturbations that preserve the spatial periodicity of the square lattice. There are five oscillatory solutions with maximal isotropy that bifurcate from the trivial solution; these are interpreted in terms of standing and travelling wave convection patterns. An unstable submaximal solution is also found to exist in an open region of parameter space. All possible bifurcation diagrams are given. The possibility of a primary bifurcation to a structurally stable heteroclinic cycle is explored.

## 1. Introduction.

Many pattern-forming systems produce approximately spatially periodic patterns. This observation has motivated a number of studies of pattern selection in systems with periodic boundary conditions. For doubly periodic patterns in a plane the resulting spatial domain may be identified with a two-torus. There are three doubly periodic lattices that tile a plane and for which the wavelength is the same in the two directions of periodicity. These are the rhombic, square and hexagonal lattices. The formulation of the problem on such a lattice introduces important symmetries into the problem. These are the symmetries of the unit cell  $D_n$ ,  $n = 2, 4, 6$  respectively, in addition to the two-torus of translations. The presence of these symmetries is responsible for the multiple solution branches that bifurcate from the trivial (homogeneous) solution, and can be exploited in analyzing their relative stability, *i.e.*, their stability properties with respect to perturbations preserving the lattice. This approach has met with considerable success in describing pattern formation arising from a steady state bifurcation [1–3]. Pattern formation arising from a Hopf bifurcation is less well studied. In this paper we focus attention on patterns, periodic on a square lattice, that form at a Hopf bifurcation. The spatial symmetry of the problem is described by the semi-direct product group  $D_4 \times_s T^2$ . Certain aspects of this problem have been considered previously. Swift [4] derived the form of the general equivariant vector field and determined the solutions of

---

\*Institute for Mathematics and its Applications, University of Minnesota, Minneapolis, MN 55455.

†Department of Physics, University of California, Berkeley, CA 94720.

the third order truncation. However, he did not consider the stability properties of these solutions. Pismen [5] determined the stabilities of the standing wave solutions in the case of the third order truncation, but did not complete the stability analysis for the travelling wave solutions. Neither paper addresses the symmetry properties of the solutions. Additional work, complementary to this paper, has been done on Hopf bifurcation with  $O(2)$  symmetry [6–9],  $D_4$  symmetry [10,11], and  $D_6 \times_s T^2$  symmetry [12].

In this paper we present an essentially complete description of the generic Hopf bifurcation on a square lattice. We employ the techniques of equivariant bifurcation theory relying heavily on the symmetry properties of the possible solutions. In the next section we point out that the symmetry is responsible for the quadrupling of the eigenvalues at the bifurcation and state explicitly the assumptions that allow a formulation of the bifurcation problem on an eight–dimensional center manifold. We discuss the symmetries of this problem and observe that in Poincaré–Birkhoff normal form it will have an additional  $S^1$  phase shift symmetry which we interpret as a time translation symmetry. In section 3 we determine the isotropy subgroup structure of the resulting symmetry group  $D_4 \times_s T^2 \times S^1$ . This analysis determines not only the possible solutions but also classifies them by their spatial and spatio–temporal symmetries. Five nontrivial solutions, guaranteed by the equivariant Hopf theorem [6], are found. In section 4, the general form of the equivariant vector field on the center manifold is derived. The oscillatory solutions that bifurcate from the trivial solution at the Hopf bifurcation are determined from the third order truncation of this vector field in section 5. It is found that in addition to the five solutions guaranteed by abstract theory a sixth solution with submaximal isotropy bifurcates in a primary bifurcation in an open region of parameter space. The stabilities of all six solutions are calculated in section 6, and all possible bifurcation diagrams depicted in section 7. In section 8 we describe the possible heteroclinic orbits connecting the various solutions and remark on the possible dynamics when all primary branches bifurcate supercritically but none are stable. In particular, we determine the region of the coefficient space where we expect a primary bifurcation to a structurally stable heteroclinic cycle. Finally, in section 9 we summarize our results and discuss possible applications.

## 2. Problem Formulation.

We consider the oscillatory instability of the trivial state of an isotropic homogeneous spatially extended system invariant under  $E(2)$ , the Euclidean group of symmetries of the plane, *i.e.*, translations, rotations and reflections. Here a “trivial state” is any time–independent solution which is uniform in the horizontal and hence does not break the  $E(2)$  symmetry. It is assumed that the instability sets in with a unique *nonzero* wavenumber  $k_c$  and frequency  $\omega_c$  leading to solutions of the linear problem that take the form of *standing* and *travelling* waves. Specifically, let  $\mathcal{S}(\mu, k; R) = 0$  be the dispersion relation for linear

modes of the form  $e^{\mu t + i\mathbf{k} \cdot \mathbf{x}} g(y)$ , where  $g(y)$  is an appropriate vertical eigenfunction,  $\mathbf{x} \in \mathbb{R}^2$  is a horizontal vector and  $S, \mu \in \mathbb{C}$ ,  $k, R \in \mathbb{R}$ ,  $k \equiv |\mathbf{k}|$ . Here  $R$  is a control parameter of the system. The neutral stability curve  $R(k)$  is determined from the real and imaginary parts of  $S(\mu = i\omega, k; R) = 0$ . Its minimum defines  $k_c$ ,  $R_c = R(k_c)$  (see Figure 1). The critical frequency satisfies  $S(i\omega_c, k_c; R_c) = 0$ . For the remainder of the paper, we let  $\lambda \propto (R - R_c)$  be the bifurcation parameter and further assume

$$\left. \frac{\partial \operatorname{Re}(\mu)}{\partial \lambda} \right|_{k=k_c, \lambda=0} > 0. \quad (2.1)$$




Figure 1: Hopf bifurcation curve  $R(k)$ . The trivial solution is unstable above the curve and linearly stable below it.

At  $\lambda = 0$  there is an entire circle of wavevectors  $\mathbf{k}$  with magnitude  $k_c$  and hence an infinite number of neutrally stable modes at the Hopf bifurcation point. In the presence of periodic boundary conditions in the horizontal this orientational degeneracy is absent. Specifically, we restrict the space of solutions to those that are periodic on a square lattice  $\mathcal{L}_s$ , where

$$\mathcal{L}_s = \{n_1 \mathbf{a}_1 + n_2 \mathbf{a}_2 : n_1, n_2 \in \mathbb{Z}, \mathbf{a}_1 = \frac{2\pi}{k_c} \hat{\mathbf{x}}_1, \mathbf{a}_2 = \frac{2\pi}{k_c} \hat{\mathbf{x}}_2\}. \quad (2.2)$$

Here,  $\hat{\mathbf{x}}_1, \hat{\mathbf{x}}_2$  are orthonormal vectors that span a horizontal plane. The symmetry of the problem thereby reduces to that of a compact subgroup of the Euclidean group; translational symmetry is identified with the two-torus  $\mathbb{T}^2 = \mathbb{R}^2/\mathcal{L}_s$  since translations by vectors in  $\mathcal{L}_s$  act trivially on the space of periodic solutions. The reflectional and rotational symmetries

are those of the unit square cell, corresponding to the discrete group  $D_4$ . The symmetry group is thus the semi-direct product of these groups:  $\Gamma_s = D_4 \times_s T^2$ .

The periodic boundary conditions are essential to the analysis which follows. Specifically, at  $\lambda = 0$  only a finite number of the linear modes have eigenvalues on the imaginary axis while the rest of the spectrum is bounded away from the imaginary axis. This allows us to invoke the center manifold theorem [13] and rigorously formulate the bifurcation problem in terms of ordinary differential equations for the amplitudes of the critical modes. The  $\Gamma_s$ -invariant subspace of neutrally stable modes at  $\lambda = 0$  is

$$\mathcal{F}_s = \left\{ (v_1(t)e^{ik_c x_1} + v_2(t)e^{ik_c x_2} + w_1(t)e^{-ik_c x_1} + w_2(t)e^{-ik_c x_2})g(y) : \right. \\ \left. \mathbf{z} \equiv (v_1, v_2, w_1, w_2) \in \mathbb{C}^4 \right\}. \quad (2.3)$$

The linear part of the ordinary differential equation for the critical amplitudes is  $\dot{\mathbf{z}} = \mu(\lambda)\mathbf{z}$ , where  $\mu(0) = i\omega_c$  and  $\text{Re}(\mu'(0)) > 0$ . Hence, we interpret  $|w_j|$  as the amplitude of a travelling wave in the positive  $\hat{x}_j$ -direction and  $|v_j|$  as the amplitude of a travelling wave in the negative  $\hat{x}_j$ -direction,  $j = 1, 2$ .

The action of the symmetry group  $\Gamma_s$  on the amplitudes of the critical modes is inherited from its natural action on  $\mathbb{R}^2$ . From (2.3) it follows that a translation  $(x_1, x_2) \rightarrow (x_1 + l_1, x_2 + l_2)$  is equivalent to a phase shift of the amplitudes

$$(\theta_1, \theta_2) : \begin{pmatrix} v_1 \\ v_2 \\ w_1 \\ w_2 \end{pmatrix} \rightarrow \begin{pmatrix} e^{i\theta_1} v_1 \\ e^{i\theta_2} v_2 \\ e^{-i\theta_1} w_1 \\ e^{-i\theta_2} w_2 \end{pmatrix}, \quad (\theta_1, \theta_2) \in T^2, \quad (2.4a)$$

where  $\theta_j = k_c l_j$ ,  $j = 1, 2$ . The discrete group  $D_4$  is generated by a reflection  $\sigma_v : x_2 \rightarrow -x_2$  and a  $\pi/2$  rotation  $r_{\pi/2} : (x_1, x_2) \rightarrow (x_2, -x_1)$ . As indicated in Figure 2 the corresponding action on  $\mathbb{C}^4$  is

$$\sigma_v : \begin{pmatrix} v_1 \\ v_2 \\ w_1 \\ w_2 \end{pmatrix} \rightarrow \begin{pmatrix} v_1 \\ w_2 \\ w_1 \\ v_2 \end{pmatrix}, \quad r_{\pi/2} : \begin{pmatrix} v_1 \\ v_2 \\ w_1 \\ w_2 \end{pmatrix} \rightarrow \begin{pmatrix} w_2 \\ v_1 \\ v_2 \\ w_1 \end{pmatrix}. \quad (2.4b)$$

This representation of  $D_4 \times_s T^2$  is not irreducible; the group representations relevant to Hopf bifurcation problems are (generically) either (a) irreducible, but not absolutely irreducible, or (b) isomorphic to the direct sum of two copies of an absolutely irreducible representation

[6]. The group representation of  $\Gamma_s$ , generated by (2.4), is of type (b): it readily decomposes into two copies of an absolutely irreducible representation acting diagonally on

$$\begin{pmatrix} v_1 + \bar{w}_1 \\ v_2 + \bar{w}_2 \\ i(v_1 - \bar{w}_1) \\ i(v_2 - \bar{w}_2) \end{pmatrix}. \quad (2.5)$$

Here, absolutely irreducible means that the only  $4 \times 4$  real matrices that commute with the group action on  $\mathbb{R}^4$  are scalar multiples of the identity matrix. The 4-dimensional representation is relevant when the trivial solution loses stability via a steady state bifurcation. In this case, the action of  $D_4 \times_s T^2$  is that of (2.4) on, say, the first two components of (2.5). This equivariant bifurcation problem was analyzed by Swift [4].




Figure 2: Action of  $D_4$  on the amplitudes  $(v_1, v_2, w_1, w_2)$ .

The vector field that commutes with the spatial symmetries (2.4) can be put into Poincaré–Birkhoff normal form. As with other Hopf bifurcation problems this procedure introduces an extra phase shift symmetry in time into the vector field. We denote this symmetry by  $S^1$ , and henceforth assume that the full spatio-temporal symmetry of the problem is specified by the group  $\Gamma_s \times S^1$ , where

$$\varphi : z \rightarrow e^{i\varphi} z, \quad \varphi \in S^1. \quad (2.6)$$

In the next section the isotropy subgroups of  $\Gamma_s \times S^1$ , together with their fixed point subspaces are determined. The isotropy subgroups determine the symmetries of all possible solutions, and the equivariant Hopf theorem [6] guarantees the existence of primary branches with

two-dimensional fixed point subspaces. This abstract analysis provides much important information without explicit computation.

### 3. Isotropy Subgroups.

The symmetry of a solution  $\mathbf{z}$  is characterized by the isotropy subgroup  $\Sigma_{\mathbf{z}} \subset \Gamma_s \times S^1$ , which leaves it fixed, *i.e.*,

$$\Sigma_{\mathbf{z}} = \{\sigma \in \Gamma_s \times S^1 : \sigma \mathbf{z} = \mathbf{z}\}. \quad (3.1)$$

Two solutions on the same group orbit have conjugate isotropy subgroups: if  $\mathbf{z}$  is a solution with symmetry  $\Sigma_{\mathbf{z}}$ , then  $\gamma \mathbf{z}$  has isotropy subgroup

$$\Sigma_{\gamma \mathbf{z}} = \gamma \Sigma_{\mathbf{z}} \gamma^{-1}. \quad (3.2)$$

We consider two solutions on the same group orbit to be equivalent. In this section, we determine all of the isotropy subgroups of  $\Gamma_s \times S^1$  up to conjugacy. In addition, we associate a fixed point subspace  $\text{Fix}(\Sigma)$  with every isotropy subgroup  $\Sigma$ , where

$$\text{Fix}(\Sigma) = \{\mathbf{z} \in \mathbb{C}^4 : \sigma \mathbf{z} = \mathbf{z} \text{ for all } \sigma \in \Sigma\}. \quad (3.3)$$

In section 5, we will use the observation that  $\text{Fix}(\Sigma)$  is invariant under the dynamics [14] to determine the periodic solutions with symmetry  $\Sigma$ .

We denote the group elements of  $\Gamma_s \times S^1$  by  $[\gamma, \varphi]$ , where  $\gamma \in \Gamma_s$  and  $\varphi \in S^1$ . The elements of  $\Gamma_s$  may be denoted by  $r_{\pi/2}^n \sigma_v^m(\theta_1, \theta_2)$ , with  $n \in \{0, 1, 2, 3\}$ ,  $m \in \{0, 1\}$ , and  $(\theta_1, \theta_2) \in \mathbb{T}^2$ . Hence, the action of a general element of  $\Gamma_s \times S^1$  on  $\mathbb{C}^4$  is

$$[r_{\pi/2}^n \sigma_v^m(\theta_1, \theta_2), \varphi] : \begin{pmatrix} v_1 \\ v_2 \\ w_1 \\ w_2 \end{pmatrix} \rightarrow r_{\pi/2}^n \sigma_v^m \begin{pmatrix} e^{i(\varphi+\theta_1)} v_1 \\ e^{i(\varphi+\theta_2)} v_2 \\ e^{i(\varphi-\theta_1)} w_1 \\ e^{i(\varphi-\theta_2)} w_2 \end{pmatrix}, \quad (3.4)$$

where the actions of  $\sigma_v$  and  $r_{\pi/2}$  are given by (2.4b). The subgroup of  $\Gamma_s \times S^1$  generated by  $\sigma_1, \dots, \sigma_k \in \Gamma_s \times S^1$  is denoted  $\langle \sigma_1, \dots, \sigma_k \rangle$ . Note that  $[(\pi, \pi), \pi]$  acts trivially on  $\mathbb{C}^4$  and hence is contained in every isotropy subgroup.

The remainder of this section is devoted to compiling Table 1 which contains a representative isotropy subgroup for each conjugacy class, together with its fixed point subspace. We begin by noting that every isotropy subgroup is conjugate to one whose fixed point subspace is of the form

$$(v_1, v_2, w_1, w_2) = (r_1, r_2 e^{i\psi}, r_3, r_4 e^{i\psi}), \quad r_j \geq 0, \quad 0 \leq \psi < \pi. \quad (3.5a)$$

That three of the phases of a solution may be chosen is a consequence of the three continuous symmetries (*i.e.*,  $\mathbb{T}^2 \times S^1$ ) which act on  $\mathbb{C}^4$  by phase shifts. If any amplitude  $r_j$  is zero then we



may, without loss of generality, further assume  $\psi = 0$ . The action of the discrete symmetries of  $D_4$  allow us to order the amplitudes  $r_j$  such that

$$r_1 \geq r_2 \geq r_4 \geq 0, \quad r_1 \geq r_3 \geq 0. \quad (3.5b)$$

The trivial solution has the symmetry of the full group, whereas if all four amplitudes  $r_j$  are unequal, the isotropy subgroup is the trivial one:  $Z_2^c = \langle [(\pi, \pi), \pi] \rangle$ . We now consider the cases where  $r_1 \neq 0$  and  $r_i = r_j$  for some  $i \neq j$ ; these fall into the following categories:

$$\begin{aligned} (a) \quad & r_2 = r_3 = r_4 = 0 \\ (b) \quad & r_3 = r_4 = 0, \quad r_2 \neq 0 \\ (c) \quad & r_2 = r_4 = 0, \quad r_3 \neq 0 \\ (d) \quad & r_1 = r_2 \neq 0, \quad r_3 \neq 0 \\ (e) \quad & r_1 = r_3 \neq 0, \quad r_2 \neq 0. \end{aligned} \quad (3.6)$$

For cases (d) and (e) we do not assume the ordering of (3.5b).

The entries in Table 1 are verified by considering in turn each of the cases in (3.6):

- (a)  $\mathbf{z} = (r_1, 0, 0, 0)$  corresponds to travelling rolls (TR) and has isotropy subgroup  $O(2) \times S^1 = \langle \sigma_v, (0, \theta_2) \rangle \times [(-\varphi, 0), \varphi]$ . Here  $S^1$  is a spatio-temporal symmetry; a time translation of the oscillation by approximately  $\varphi/\omega_c$  is followed by a translation in the  $\hat{x}_1$  direction by  $-\varphi/k_c$ .
- (b)  $\mathbf{z} = (r_1, r_2, 0, 0)$  has isotropy subgroup  $S^1 = [(-\varphi, -\varphi), \varphi]$  if  $r_1 \neq r_2$  (VIII in Table 1), and isotropy subgroup  $Z_2 \times S^1$  if  $r_1 = r_2$ , where  $Z_2 = \langle r_{\pi/2} \sigma_v \rangle$  (travelling squares (TS)).
- (c)  $\mathbf{z} = (r_1, 0, r_3, 0)$  has isotropy subgroup  $Z_2^c \times O(2)$  if  $r_1 \neq r_3$  (IX in Table 1). If  $r_1 = r_3$  then the isotropy subgroup is enlarged by  $Z_2 = \langle r_{\pi/2}^2 \rangle$  (standing rolls (SR)).
- (d)  $\mathbf{z} = (r_1, r_1 e^{i\psi}, r_3, r_4 e^{i\psi})$  must have  $r_3 = r_4$  for the isotropy subgroup to be nontrivial. If  $r_3 = r_4 \neq r_1$ , then the isotropy subgroup is nontrivial for those values of  $\psi$  satisfying

$$[r_{\pi/2} \sigma_v(\theta_1, \theta_2), \varphi] \begin{pmatrix} r_1 \\ r_1 e^{i\psi} \\ r_3 \\ r_3 e^{i\psi} \end{pmatrix} = \begin{pmatrix} r_1 e^{i(\varphi + \theta_2 + \psi)} \\ r_1 e^{i(\varphi + \theta_1)} \\ r_3 e^{i(\varphi - \theta_2 + \psi)} \\ r_3 e^{i(\varphi - \theta_1)} \end{pmatrix} = \begin{pmatrix} r_1 \\ r_1 e^{i\psi} \\ r_3 \\ r_3 e^{i\psi} \end{pmatrix}. \quad (3.7)$$

This equation has a solution  $[(\theta_1, \theta_2), \varphi] \in T^2 \times S^1$  provided  $\psi = 0, \pi/2$ . For  $\mathbf{z} = (r_1, r_1, r_3, r_3)$  the isotropy subgroup is  $Z_2^c \times Z_2$ , where  $Z_2 = \langle r_{\pi/2} \sigma_v \rangle$  (X in Table 1). For  $\mathbf{z} = (r_1, ir_1, r_3, ir_3)$  the isotropy subgroup is  $Z_2^c \times \tilde{Z}_2$ , where  $\tilde{Z}_2 = \langle [r_{\pi/2} \sigma_v(0, \pi), \pi/2] \rangle$  (XI in Table 1). Here the tilde on  $\tilde{Z}_2$  indicates that it is a spatio-temporal symmetry of the solution.

In the case where  $\mathbf{z} = (r_1, r_1 e^{i\psi}, r_1, r_1 e^{i\psi})$ , the isotropy subgroup is  $Z_2^c \times Z_2^2 = Z_2^c \times$

Name	Fixed Point Subspace	Isotropy Subgroup
I. Trivial Solution (T)	$v_1 = v_2 = w_1 = w_2 = 0$	$\Gamma_s \times S^1$
II. Travelling Rolls (TR)	$v_1 \neq 0, v_2 = w_1 = w_2 = 0$	$O(2) \times S^1 : O(2) = \langle \sigma_v, (0, \theta_2) \rangle,$ $S^1 = [(-\varphi, 0), \varphi]$
III. Travelling Squares (TS)	$v_1 = v_2 \neq 0, w_1 = w_2 = 0$	$Z_2 \times S^1 : Z_2 = \langle r_{\pi/2} \sigma_v \rangle,$ $S^1 = [(-\varphi, -\varphi), \varphi]$
IV. Standing Rolls (SR)	$v_1 = w_1 \neq 0, v_2 = w_2 = 0$	$Z_2^c \times Z_2 \times O(2) : Z_2 = \langle r_{\pi/2}^2 \rangle$
V. Standing Squares (SS)	$v_1 = v_2 = w_1 = w_2 \neq 0$	$Z_2^c \times D_4$
VI. Alternating Rolls (AR)	$v_1 = -iv_2 = w_1 = -iw_2 \neq 0$	$Z_2^c \times \tilde{D}_4 :$ $\tilde{D}_4 = \langle \sigma_v, [r_{\pi/2}(0, \pi), \pi/2] \rangle$
VII. Standing Cross-Rolls (SCR)	$v_1 = w_1 \neq 0, v_2 = w_2 \neq 0,$ $ v_1  \neq  v_2 $	$Z_2^c \times Z_2^2 : Z_2^2 = \langle \sigma_v, r_{\pi/2}^2 \rangle$
VIII.	$v_1 \neq 0, v_2 \neq 0, w_1 = w_2 = 0,$ $ v_1  \neq  v_2 $	$S^1 : S^1 = [(-\varphi, -\varphi), \varphi]$
IX.	$v_1 \neq 0, w_1 \neq 0, v_2 = w_2 = 0,$ $ v_1  \neq  w_1 $	$Z_2^c \times O(2)$
X.	$v_1 = v_2 \neq 0, w_1 = w_2 \neq 0,$ $ v_1  \neq  w_1 $	$Z_2^c \times Z_2 : Z_2 = \langle r_{\pi/2} \sigma_v \rangle$
XI.	$v_1 = -iv_2 \neq 0, w_1 = -iw_2 \neq 0,$ $ v_1  \neq  w_1 $	$Z_2^c \times \tilde{Z}_2 :$ $\tilde{Z}_2 = \langle [r_{\pi/2} \sigma_v(0, \pi), \pi/2] \rangle$
XII.	$v_1 = w_1 \neq 0, v_2 \neq 0,$ $ v_2  \neq  w_2 $	$Z_2^c \times Z_2 : Z_2 = \langle r_{\pi/2}^2 \sigma_v \rangle$
XIII.	$v_1 \neq 0, w_1 \neq 0, v_2 \neq 0,$ $ v_2  \neq  w_2  \neq  v_1  \neq  w_1  \neq  w_2 $	$Z_2^c \equiv [(\pi, \pi), \pi]$

Table 1: All isotropy subgroups of  $\Gamma_s \times S^1$ , up to conjugacy, listed with their associated fixed point subspaces.

$\langle \sigma_v, r_{\pi/2}^2 \rangle$  for  $\psi \neq 0, \pi/2$ . If  $\psi = 0$  the isotropy subgroup is  $Z_2^c \times D_4$  (standing squares (SS)), whereas if  $\psi = \pi/2$  the isotropy subgroup is  $Z_2^c \times \tilde{D}_4$ , where  $\tilde{D}_4 = \langle \sigma_v, [r_{\pi/2}(0, \pi), \pi/2] \rangle$  (alternating rolls (AR)).

(e)  $\mathbf{z} = (r_1, r_2 e^{i\psi}, r_1, r_4 e^{i\psi})$  has isotropy subgroup  $Z_2^c \times Z_2 = Z_2^c \times \langle r_{\pi/2}^2 \sigma_v \rangle$  if  $r_2 \neq r_4$  (XII in Table 1). If  $r_2 = r_4 \neq r_1$  then the isotropy subgroup is  $Z_2^c \times Z_2^2$  where  $Z_2^2 = \langle \sigma_v, r_{\pi/2}^2 \rangle$  as in case (d) above (standing cross-rolls (SCR) in Table 1). The case  $r_1 = r_2 = r_4$  is also examined in (d).






Figure 3: Lattice of isotropy subgroups of  $\Gamma_s \times S^1$ . Each entry represents the entire conjugacy class of the isotropy subgroup. Inclusion is indicated by an arrow.

The “lattice” of isotropy subgroups in Figure 3 summarizes the ways in which symmetry may be broken in this problem. The five maximal isotropy subgroups describe the symmetries of solutions II–VI in Table 1; each has a two-dimensional fixed point subspace. The equivariant Hopf theorem [6] guarantees that these five solutions bifurcate from the trivial solution at  $\lambda = 0$ . The planforms associated with these solutions are determined from

## TRAVELLING ROLLS




## TRAVELLING SQUARES



(Figure 4 continued on the next page)

## STANDING ROLLS




$t = 0$

$t = 1/8$

$t = 1/4$

$t = 3/8$




$t = 1/2$

$t = 5/8$

$t = 3/4$

$t = 7/8$

## STANDING SQUARES




$t = 0$

$t = 1/8$

$t = 1/4$

$t = 3/8$



$t = 1/2$

$t = 5/8$

$t = 3/4$

$t = 7/8$

(Figure 4 continued on the next page)

# ALTERNATING ROLLS




Figure 4: Planforms associated with the solutions guaranteed by the equivariant Hopf theorem. The square lattice periodicity is indicated by depicting four of the periodic cells. Note the symmetry between the solutions at  $t = 0$  and half a cycle later at  $t = 1/2$  (i.e., the trivial symmetry  $Z_2^c$ ).

the real part of the linear eigenfunctions in (2.3). The time evolution of each planform is depicted in Figure 4 as it would appear in a shadowgraph image of a convecting fluid layer. In this case dark regions correspond to hot rising fluid and light regions to cold descending fluid. Note that the solutions TR, TS and AR have nontrivial spatio-temporal symmetries, whereas the symmetries of SR and SS are purely spatial (excluding the trivial symmetry  $Z_2^c$ ).

Finally, it is of interest to determine the normalizer  $\mathcal{N}(\Sigma)$  of each isotropy subgroup  $\Sigma \in \Gamma_s \times S^1$ , where

$$\mathcal{N}(\Sigma) = \{\gamma \in \Gamma_s \times S^1 : \gamma^{-1}\Sigma\gamma = \Sigma\}. \quad (3.8)$$

The elements of  $\mathcal{N}(\Sigma)$  map the fixed point subspace  $\text{Fix}(\Sigma)$  to itself, with the elements of  $\mathcal{N}(\Sigma)/\Sigma$  acting nontrivially. Hence, a  $\Gamma_s \times S^1$ -equivariant vector field restricted to the invariant subspace  $\text{Fix}(\Sigma)$  is  $\mathcal{N}(\Sigma)$ -equivariant. For example, the maximal isotropy subgroups have  $\mathcal{N}(\Sigma) = S^1$ . In Table 2 we specify  $\mathcal{N}(\Sigma)/\Sigma$  for the isotropy subgroups with four-dimensional fixed point subspaces (*i.e.*, entries VII–XI of Table 1). Note that  $\mathcal{N}(\Sigma)/\Sigma \approx O(2) \times S^1$  for isotropy subgroups VIII–XI and hence that the Hopf bifurcation problem with  $O(2) \times S^1$  symmetry applies in these invariant subspaces. This bifurcation problem has been extensively analyzed using singularity theory methods [8,9]. In particular, universal unfoldings of the normal forms for all degenerate bifurcation problems through codimension-two have been determined. The  $D_4 \times S^1$ -equivariant Hopf bifurcation problem, which applies in the SCR subspace, has been analyzed by Swift [11]. It should be noted that the results obtained in this subspace apply not only to the Hopf bifurcation problem with periodic boundary conditions (as posed in the previous section), but also to the same problem posed on a square domain with Neumann boundary conditions. Specifically, it is well known that, suitably extended, solutions obtained with Neumann boundary conditions also solve the corresponding problem with periodic boundary conditions. This extension can introduce additional symmetries into the problem (see, for example, Crawford *et al.* [15]). In the case of the square domain, these symmetries are precisely those listed for the SCR subspace in Table 2. Note that the “reflection symmetry”  $(v_1, v_2) \rightarrow (v_1, -v_2)$  actually corresponds to a translation  $(0, \pi) \in T^2$ .

In the next section, we show by direct calculation that the form of the vector field restricted to any of the four-dimensional subspaces  $\text{Fix}(\Sigma)$  is exactly what we expect from considerations of the normalizer  $\mathcal{N}(\Sigma)$  (*i.e.*, there are no additional restrictions placed on a  $\mathcal{N}(\Sigma)$ -equivariant vector field obtained in this way). This calculation is necessary in order to exclude the possibility that a degenerate  $\mathcal{N}(\Sigma)$ -equivariant bifurcation problem arises naturally when the equivariant vector field is restricted to  $\text{Fix}(\Sigma)$ .

Representative Fixed Point Subspace	$\mathcal{N}(\Sigma)/\Sigma$
VII. (SCR) $(v_1, v_2, w_1, w_2) = (z_1, z_2, z_1, z_2) \in \mathbb{C}^2$	$D_4 \times S^1$ $r_{\pi/2} : (z_1, z_2) \rightarrow (z_2, z_1)$ $(0, \pi) : (z_1, z_2) \rightarrow (z_1, -z_2)$ $\varphi : (z_1, z_2) \rightarrow e^{i\varphi}(z_1, z_2)$ $\{r_{\pi/2}, (0, \pi)\} \in D_4; \quad \varphi \in S^1$
VIII. $(v_1, v_2, w_1, w_2) = (z_1, z_2, 0, 0) \in \mathbb{C}^2$	$O(2) \times S^1$ $(r_{\pi/2}\sigma_v) : (z_1, z_2) \rightarrow (z_2, z_1)$ $(\theta, -\theta) : (z_1, z_2) \rightarrow (e^{i\theta}z_1, e^{-i\theta}z_2)$ $\varphi : (z_1, z_2) \rightarrow e^{i\varphi}(z_1, z_2)$ $\{(r_{\pi/2}\sigma_v), (\theta, -\theta)\} \in O(2); \quad \varphi \in S^1$
IX. $(v_1, v_2, w_1, w_2) = (z_1, 0, z_2, 0) \in \mathbb{C}^2$	$O(2) \times S^1$ $r_{\pi/2}^2 : (z_1, z_2) \rightarrow (z_2, z_1)$ $(\theta, 0) : (z_1, z_2) \rightarrow (e^{i\theta}z_1, e^{-i\theta}z_2)$ $\varphi : (z_1, z_2) \rightarrow e^{i\varphi}(z_1, z_2)$ $\{r_{\pi/2}^2, (\theta, 0)\} \in O(2); \quad \varphi \in S^1$
X. $(v_1, v_2, w_1, w_2) = (z_1, z_1, z_2, z_2) \in \mathbb{C}^2$	$O(2) \times S^1$ $r_{\pi/2}^2 : (z_1, z_2) \rightarrow (z_2, z_1)$ $(\theta, \theta) : (z_1, z_2) \rightarrow (e^{i\theta}z_1, e^{-i\theta}z_2)$ $\varphi : (z_1, z_2) \rightarrow e^{i\varphi}(z_1, z_2)$ $\{r_{\pi/2}^2, (\theta, \theta)\} \in O(2); \quad \varphi \in S^1$
XI. $(v_1, v_2, w_1, w_2) = (z_1, iz_1, z_2, iz_2) \in \mathbb{C}^2$	$O(2) \times S^1$ $r_{\pi/2}^2 : (z_1, z_2) \rightarrow (z_2, z_1)$ $(\theta, \theta) : (z_1, z_2) \rightarrow (e^{i\theta}z_1, e^{-i\theta}z_2)$ $\varphi : (z_1, z_2) \rightarrow e^{i\varphi}(z_1, z_2)$ $\{r_{\pi/2}^2, (\theta, \theta)\} \in O(2); \quad \varphi \in S^1$

Table 2: Action of  $\mathcal{N}(\Sigma)$  on  $\text{Fix}(\Sigma)$  for all four-dimensional fixed point subspaces.



#### 4. Equivariant Vector Field.

The symmetry  $\Gamma_s \times S^1$  severely restricts the form of the dynamical equation on the center manifold

$$\dot{\mathbf{z}} = \mathbf{f}(\mathbf{z}, \lambda); \quad \mathbf{f} : \mathbb{C}^4 \times \mathbb{R} \rightarrow \mathbb{C}^4. \quad (4.1a)$$

Specifically,  $\mathbf{f}$  must be  $\Gamma_s \times S^1$ -equivariant:

$$\gamma \mathbf{f}(\mathbf{z}, \lambda) = \mathbf{f}(\gamma \mathbf{z}, \lambda) \quad \text{for all } \gamma \in \Gamma_s \times S^1. \quad (4.1b)$$

We further assume that  $\mathbf{f}$  is smooth ( $C^\infty$ ). The general form of such a vector field is

$$\begin{aligned} \dot{v}_1 &= v_1 g(N_1, \delta_1, N_2, \Delta_2, Q) + \bar{w}_1 v_2 w_2 h(N_1, \delta_1, N_2, \Delta_2, \bar{Q}) \\ \dot{v}_2 &= v_2 g(N_2, \delta_2, N_1, \Delta_1, \bar{Q}) + \bar{w}_2 v_1 w_1 h(N_2, \delta_2, N_1, \Delta_1, Q) \\ \dot{w}_1 &= w_1 g(N_1, -\delta_1, N_2, \Delta_2, Q) + \bar{v}_1 v_2 w_2 h(N_1, -\delta_1, N_2, \Delta_2, \bar{Q}) \\ \dot{w}_2 &= w_2 g(N_2, -\delta_2, N_1, \Delta_1, \bar{Q}) + \bar{v}_2 v_1 w_1 h(N_2, -\delta_2, N_1, \Delta_1, Q), \end{aligned} \quad (4.2)$$

where  $g$  and  $h$  are  $C^\infty$  complex-valued functions of  $N_j \equiv |v_j|^2 + |w_j|^2$ ,  $\delta_j \equiv |w_j|^2 - |v_j|^2$ ,  $\Delta_j \equiv \delta_j^2$ , and  $Q \equiv v_1 \bar{v}_2 w_1 \bar{w}_2$ . This section is devoted to the derivation of (4.2).

Consider a general term in the Taylor expansion of the first component of (4.1a), *i.e.*, of  $\dot{v}_1 = f_1(v_1, v_2, w_1, w_2)$ :

$$A_{\alpha\beta\gamma\delta\sigma\tau\mu\nu} v_1^\alpha v_2^\beta w_1^\gamma w_2^\delta \bar{v}_1^\sigma \bar{v}_2^\tau \bar{w}_1^\mu \bar{w}_2^\nu. \quad (4.3)$$

The coefficient  $A_{\alpha\dots\nu}$  is nonzero for those terms that commute with the action of  $T^2 \times S^1$ . For the  $T^2$  action (2.4a) this implies  $A_{\alpha\dots\nu} \neq 0$  for

$$\begin{aligned} (\alpha - \sigma) - (\gamma - \mu) &= 1 \\ (\beta - \tau) - (\delta - \nu) &= 0. \end{aligned} \quad (4.4a)$$

Thus  $(\alpha - \sigma) = 1 + (\gamma - \mu) \equiv m$  and  $(\tau - \beta) = (\nu - \delta) \equiv n$ ,  $m, n \in \mathbb{Z}$ . The term (4.3) must also commute with the  $S^1$  action (2.6), so that  $A_{\alpha\dots\nu} \neq 0$  for

$$\alpha + \beta + \gamma + \delta - \sigma - \tau - \mu - \nu = 1. \quad (4.4b)$$

It follows that  $m = n + 1$ . Thus the terms in the Taylor expansion of  $f_1$  are

$$B_{n\alpha\beta\gamma\delta} v_1^{\alpha+n+1} v_2^\beta w_1^{\gamma+n} w_2^\delta \bar{v}_1^\alpha \bar{v}_2^{\beta+n} \bar{w}_1^\gamma \bar{w}_2^{\delta+n}. \quad (4.5)$$

Those terms with  $n \geq 0$  may be expressed as

$$B_{n\alpha\beta\gamma\delta} |v_1|^{2\alpha} |v_2|^{2\beta} |w_1|^{2\gamma} |w_2|^{2\delta} Q^n v_1, \quad (4.6)$$

where  $Q \equiv v_1 \bar{v}_2 w_1 \bar{w}_2$  is an invariant of the  $T^2 \times S^1$  action. The terms with  $n \leq -1$  have the form

$$B_{n\alpha\beta\gamma\delta} |v_1|^{2(\alpha+n+1)} |v_2|^{2(\beta+n)} |w_1|^{2(\gamma+n)} |w_2|^{2(\delta+n)} \bar{Q}^{-(n+1)} v_2 w_2 \bar{w}_1. \quad (4.7)$$

Thus the form of  $f_1(v_1, v_2, w_1, w_2)$ , which commutes with  $T^2 \times S^1$ , is

$$f_1 = v_1 G + v_2 w_2 \bar{w}_1 H, \quad (4.8)$$

where  $G \equiv G(N_1, \delta_1, N_2, \delta_2, Q)$  and  $H \equiv H(N_1, \delta_1, N_2, \delta_2, \bar{Q})$  are general invariant functions of  $T^2 \times S^1$ . In addition,  $G$  and  $H$  must be invariant under the action of  $\sigma_v : (v_2, w_2) \rightarrow (w_2, v_2)$ . Any smooth function  $F(x, y)$  with the property  $F(x, y) = F(y, x)$  may be expressed in terms of the invariants  $N = x + y$  and  $\Delta = (y - x)^2$  (i.e.,  $F = \widehat{F}(N, \Delta)$ ). Hence we write  $G = g(N_1, \delta_1, N_2, \Delta_2, Q)$  and  $H = h(N_1, \delta_1, N_2, \Delta_2, \bar{Q})$ . The remaining components of (4.2) are generated from (4.8) by the action of  $D_4$  on a  $\Gamma_s \times S^1$ -equivariant vector field. Therefore

$$\begin{aligned} \dot{v}_2 &= f_1(v_2, w_1, w_2, v_1) \\ \dot{w}_1 &= f_1(w_1, w_2, v_1, v_2) \\ \dot{w}_2 &= f_1(w_2, v_1, v_2, w_1). \end{aligned} \quad (4.9)$$

As noted in the previous section, the  $\Gamma_s \times S^1$ -equivariant vector field (4.2) restricted to the four-dimensional fixed point subspaces of Table 1 commutes with either the action of  $D_4 \times S^1$  or  $O(2) \times S^1$ . The general  $D_4 \times S^1$ -equivariant vector field is

$$\begin{aligned} \dot{v}_1 &= v_1 g_{D_4}(|v_1|^2, |v_2|^2, v_1^2 \bar{v}_2^2) + \bar{v}_1 v_2^2 h_{D_4}(|v_1|^2, |v_2|^2, \bar{v}_1^2 v_2^2) \\ \dot{v}_2 &= v_2 g_{D_4}(|v_2|^2, |v_1|^2, v_2^2 \bar{v}_1^2) + \bar{v}_2 v_1^2 h_{D_4}(|v_2|^2, |v_1|^2, \bar{v}_2^2 v_1^2). \end{aligned} \quad (4.10)$$

This vector field pertains to the standing cross-rolls subspace where  $v_1 = w_1$ ,  $v_2 = w_2$ . The third order truncation of (4.10) was analyzed by Swift [11]. The general  $O(2) \times S^1$ -equivariant vector field is

$$\begin{aligned} \dot{z}_1 &= z_1 g_{O_2}(|z_1|^2, |z_2|^2) \\ \dot{z}_2 &= z_2 g_{O_2}(|z_2|^2, |z_1|^2). \end{aligned} \quad (4.11)$$

These equations pertain to the four-dimensional fixed point subspaces VIII–XI in Table 1 and have been extensively studied by a number of authors [6–9]. Note that the equations for the magnitudes of the complex amplitudes decouple from the phases so that we can study the two-dimensional  $D_4$ -equivariant system [4]:

$$\begin{aligned} \dot{r}_1 &= r_1 g_r(r_1^2, r_2^2) \\ \dot{r}_2 &= r_2 g_r(r_2^2, r_1^2), \end{aligned} \quad (4.12)$$

where  $z_j = r_j e^{i\phi_j}$ ,  $j = 1, 2$ , and  $g_r : \mathbf{R}^2 \rightarrow \mathbf{R}$ .

## 5. Periodic Solutions.

In this section we let the functions  $g$  and  $h$  in (4.2) depend on a bifurcation parameter  $\lambda$  as discussed in section 2. We assume that the trivial solution is stable subcritically, and that the bifurcation parameter is scaled such that

$$g(0, \lambda) = \lambda + i\omega(\lambda) + \mathcal{O}(\lambda^2), \quad \omega(0) = \omega_c. \quad (5.1)$$

We use a result of Golubitsky and Stewart [6] to determine the small amplitude periodic solutions of (4.2) that bifurcate from the trivial solution at  $\lambda = 0$ . Specifically, these authors show that the small amplitude periodic solutions with period  $2\pi/(\omega_c + \tau)$  are in one-to-one correspondence with the solutions of

$$\phi(\mathbf{z}, \lambda, \tau) \equiv \mathbf{f}(\mathbf{z}, \lambda) - i(\omega_c + \tau)\mathbf{z} = \mathbf{0} \quad (5.2)$$

on a neighborhood of  $(\mathbf{z}, \lambda, \tau) = (\mathbf{0}, 0, 0)$ .

Consider the Taylor expansion of (5.2) about  $(\mathbf{z}, \lambda, \tau) = (\mathbf{0}, 0, 0)$  through third order in  $\mathbf{z}$ :

$$\Phi \equiv \begin{pmatrix} \nu v_1 + (a|w_1|^2 + bN_1 + cN_2)v_1 + d\bar{w}_1 v_2 w_2 \\ \nu v_2 + (a|w_2|^2 + bN_2 + cN_1)v_2 + d\bar{w}_2 v_1 w_1 \\ \nu w_1 + (a|v_1|^2 + bN_1 + cN_2)w_1 + d\bar{v}_1 v_2 w_2 \\ \nu w_2 + (a|v_2|^2 + bN_2 + cN_1)w_2 + d\bar{v}_2 v_1 w_1 \end{pmatrix} = \mathbf{0}, \quad (5.3)$$

where  $\nu \equiv \lambda + i\sigma$ ,  $\sigma \equiv \omega - (\omega_c + \tau)$ . The cubic coefficients in (5.3) are complex and may be identified with the coefficients of the cubic terms in the Taylor expansion of (4.2) as follows

$$\begin{aligned} a &= 2g_2(\mathbf{0}), & b &= g_1(\mathbf{0}) - g_2(\mathbf{0}), \\ c &= g_3(\mathbf{0}), & d &= h(\mathbf{0}). \end{aligned} \quad (5.4)$$

Here  $g_j(\mathbf{0})$  is the partial derivative of  $g$  with respect to its  $j^{\text{th}}$  argument evaluated at  $(\mathbf{z}, \lambda) = (\mathbf{0}, 0)$ . Solutions of (5.3) with  $Q \equiv v_1 \bar{v}_2 w_1 \bar{w}_2 \neq 0$  satisfy

$$\hat{\Phi} \equiv \begin{pmatrix} \nu r_1 + (ar_3^2 + bN_1 + cN_2)r_1 + de^{2i\psi}r_2 r_3 r_4 \\ \nu r_2 + (ar_4^2 + bN_2 + cN_1)r_2 + de^{-2i\psi}r_3 r_4 r_1 \\ \nu r_3 + (ar_1^2 + bN_1 + cN_2)r_3 + de^{2i\psi}r_4 r_1 r_2 \\ \nu r_4 + (ar_2^2 + bN_2 + cN_1)r_4 + de^{-2i\psi}r_1 r_2 r_3 \end{pmatrix} = \mathbf{0}, \quad (5.5)$$

where  $r_1 = |v_1|$ ,  $r_2 = |v_2|$ ,  $r_3 = |w_1|$ ,  $r_4 = |w_2|$  and  $\psi = -\frac{1}{2}\arg(Q)$ . The remainder of this section is devoted to compiling Table 3 which contains all nontrivial solutions of (5.3).

We begin by noting that the nontrivial solutions have either one, two, or four nonzero amplitudes. This follows from the fourth component of (5.3) which shows that  $w_2 =$

Name	Branching Equations	Solution
Travelling Rolls (TR)	$\nu + b v_1 ^2 = 0$ $v_1 \neq 0, v_2 = w_1 = w_2 = 0$	$ v_1 ^2 = -\lambda/b_r$ $\sigma = -b_i v_1 ^2$
Travelling Squares (TS)	$\nu + (b + c) v_1 ^2 = 0$ $v_1 = v_2 \neq 0, w_1 = w_2 = 0$	$ v_1 ^2 = -\lambda/(b_r + c_r)$ $\sigma = -(b_i + c_i) v_1 ^2$
Standing Rolls (SR)	$\nu + (a + 2b) v_1 ^2 = 0$ $v_1 = w_1 \neq 0, v_2 = w_2 = 0$	$ v_1 ^2 = -\lambda/(a_r + 2b_r)$ $\sigma = -(a_i + 2b_i) v_1 ^2$
Standing Squares (SS)	$\nu + (a + 2b + 2c + d) v_1 ^2 = 0$ $v_1 = v_2 = w_1 = w_2 \neq 0$	$ v_1 ^2 = -\lambda/(a_r + 2b_r + 2c_r + d_r)$ $\sigma = -(a_i + 2b_i + 2c_i + d_i) v_1 ^2$
Alternating Rolls (AR)	$\nu + (a + 2b + 2c - d) v_1 ^2 = 0$ $v_1 = -iv_2 = w_1 = -iw_2 \neq 0$	$ v_1 ^2 = -\lambda/(a_r + 2b_r + 2c_r - d_r)$ $\sigma = -(a_i + 2b_i + 2c_i - d_i) v_1 ^2$
Standing Cross-Rolls (SCR)	$\nu + (a + 2b)r_1^2 + (2c + de^{2i\psi})r_2^2 = 0$ $\nu + (a + 2b)r_2^2 + (2c + de^{-2i\psi})r_1^2 = 0$ $r_1 =  v_1 , r_2 =  v_2 , 2\psi = \arg(\bar{v}_1^2 v_2^2)$ $v_1 = w_1 \neq 0, v_2 = w_2 \neq 0, r_1 \neq r_2$	$r_1^2 + r_2^2 = -2\lambda \operatorname{Im}(\bar{d}f)/[d_i( f ^2 -  d ^2) + 2(a_r + 2b_r) \operatorname{Im}(\bar{d}f)]$ $\sigma = -[d_r( d ^2 -  f ^2) + 2(a_i + 2b_i) \operatorname{Im}(\bar{d}f)](\tau_1^2 + r_2^2)/2 \operatorname{Im}(\bar{d}f)$ $\cos(2\psi) = \operatorname{Re}(\bar{d}f)/ d ^2, \sin(2\psi) = \sqrt{1 - \cos^2(2\psi)}$ $(r_1/r_2)^2 = [\operatorname{Im}(\bar{d}f) -  d ^2 \sin(2\psi)]/[\operatorname{Im}(\bar{d}f) +  d ^2 \sin(2\psi)]$

Table 3: Solutions of (5.3). Subscripts  $r$  and  $i$  on the coefficients  $a, b, c, d$  refer to real and imaginary parts,  $\nu \equiv \lambda + i\sigma$ , and  $f \equiv a + 2b - 2c$ .

0,  $v_1 v_2 w_1 \neq 0$ , implies  $d = 0$ . Thus generically there are no solutions with exactly three nonzero amplitudes. To verify Table 3, consider the four cases

$$\begin{aligned}
(1) \quad & v_1 \neq 0, \quad v_2 = w_1 = w_2 = 0 \\
(2) \quad & v_1 \neq 0, \quad v_2 \neq 0, \quad w_1 = w_2 = 0 \\
(3) \quad & v_1 \neq 0, \quad w_1 \neq 0, \quad v_2 = w_2 = 0 \\
(4) \quad & Q \neq 0.
\end{aligned} \tag{5.6}$$

By restricting (5.3) to the appropriate invariant fixed point subspace it is straightforward to check that these equations admit only the solutions TR, TS, and SR for cases (1), (2), and (3), respectively. For case (4), we consider equation (5.5). Let  $\hat{\Phi}_j$  denote the  $j^{\text{th}}$  component of  $\hat{\Phi}$ ; then

$$\begin{aligned}
r_1 \hat{\Phi}_1 - r_3 \hat{\Phi}_3 &= (\nu + bN_1 + cN_2)(r_1^2 - r_3^2) = 0 \\
r_2 \hat{\Phi}_2 - r_4 \hat{\Phi}_4 &= (\nu + bN_2 + cN_1)(r_2^2 - r_4^2) = 0.
\end{aligned} \tag{5.7}$$

Thus case (4) divides into three distinct subcases:

$$\begin{aligned}
(4a) \quad & r_1^2 = r_3^2, \quad r_2^2 = r_4^2 \\
(4b) \quad & \nu + bN_1 + cN_2 = \nu + bN_2 + cN_1 = 0 \\
(4c) \quad & \nu + bN_1 + cN_2 = 0, \quad r_2^2 = r_4^2.
\end{aligned} \tag{5.8}$$

The solutions SS, AR and SCR all satisfy the conditions of (4a). These are determined as follows. Consider the following equation evaluated at  $r_1 = r_3$ ,  $r_2 = r_4$ ,

$$\frac{\hat{\Phi}_1}{r_1} - \frac{\hat{\Phi}_2}{r_2} = (f - d \cos(2\psi))(r_1^2 - r_2^2) + id(r_1^2 + r_2^2) \sin(2\psi) = 0, \tag{5.9}$$

where  $f \equiv a + 2b - 2c$ . Solutions with  $r_1^2 = r_2^2$  have  $\sin(2\psi) = 0$ . These correspond to SS and AR for  $\psi = 0, \pi/2$ , respectively. If, on the other hand,  $r_1^2 \neq r_2^2$ , then the real and imaginary parts of (5.9) may be solved for  $\cos(2\psi)$  and  $(r_1/r_2)^2$  to give

$$\begin{aligned}
\cos(2\psi) &= \frac{\text{Re}(\bar{d}f)}{|d|^2} \\
\left(\frac{r_1}{r_2}\right)^2 &= \frac{\text{Im}(\bar{d}f) - |d|^2 \sin(2\psi)}{\text{Im}(\bar{d}f) + |d|^2 \sin(2\psi)}.
\end{aligned} \tag{5.10}$$

This is the solution SCR in Table 3. Without loss of generality we may assume that  $\sin(2\psi) > 0$  since letting  $\sin(2\psi) \rightarrow -\sin(2\psi)$  in (5.10) has the same effect as interchanging  $r_1$  and  $r_2$ . Thus a representative solution on the group orbit of SCR has  $\sin(2\psi) = \sqrt{1 - \cos^2(2\psi)}$ . From the conditions  $|\cos(2\psi)| < 1$  and  $(r_1/r_2)^2 > 0$ , we find that the solution (5.10) exists provided

$$|f|^2 > |d|^2 > |\text{Re}(\bar{d}f)|. \tag{5.11}$$

It remains to show that cases (4b) and (4c) do not contain any new solutions to (5.5).

In case (4b), (5.5) holds for

$$\begin{aligned} ar_1r_3 + de^{2i\psi}r_2r_4 &= 0 \\ ar_2r_4 + de^{-2i\psi}r_1r_3 &= 0, \end{aligned} \tag{5.12}$$

which for  $Q \neq 0$  implies  $a^2 = d^2$ . Generically, there are no solutions in case (4b) with  $Q \neq 0$ . In case (4c), (5.5) holds for

$$ar_1r_3 + de^{2i\psi}r_2^2 = 0 \tag{5.13a}$$

$$\nu + ar_2^2 + bN_2 + cN_1 + de^{-2i\psi}r_1r_3 = 0. \tag{5.13b}$$

Solving (5.13a) for  $r_1r_3e^{-2i\psi}$  and solving (4c) in (5.8) for  $\nu$ , we can write (5.13b) as

$$(c - b)N_1 = \left( c - b + \frac{d^2 - a^2}{2a} \right) N_2. \tag{5.14}$$

Here we replaced  $r_2^2$  by  $N_2/2$ . The real and imaginary parts of (5.14) give two equations for the ratio  $N_1/N_2$  so that generically there are no new solutions in case (4c).

In summary we have found that generically there are either five or six nontrivial periodic solution branches in the case of the cubic truncation of (4.2). The existence of the sixth solution depends on the values of the cubic coefficients in (4.2). The five solutions that always bifurcate from the trivial solution at  $\lambda = 0$  are those guaranteed by the equivariant Hopf theorem as discussed in section 3. The corresponding horizontal planforms are depicted in Figure 4. The sixth solution which may exist is SCR. This solution is a superposition of standing rolls aligned with the  $\hat{x}_1$  and  $\hat{x}_2$  directions. The relative phases and amplitudes of these standing rolls depend on the cubic coefficients through equations (5.10). We note that the standing wave solutions, which satisfy  $r_1 = r_3$ ,  $r_2 = r_4$  (*i.e.*, SR, SS, AR, SCR), were found by Swift [11] who studied the Hopf bifurcation with  $D_4$  symmetry. The TR and SR solutions were found in the study of Hopf bifurcations with  $O(2)$  symmetry [6]. The travelling wave solution TS, which exist because of the spatial translation symmetry  $T^2$ , is new.

## 6. Solution Stability.

In this section we determine the linear stability of the solutions in Table 3 with respect to perturbations periodic on the square lattice. Since the solutions are neutrally stable to perturbations that merely translate them along their group orbit, we will consider the stability of the group orbit of a solution to small perturbations transverse to the group orbit. If such perturbations decay exponentially, then the solutions are linearly orbitally stable.

Let  $\mathbf{D}\phi|_z$  denote the Jacobian matrix associated with (5.2) evaluated on a solution branch  $(z(\lambda), \tau(\lambda))$  of  $\phi = \mathbf{0}$ . A solution is linearly orbitally stable if all eigenvalues of

$\mathbf{D}\phi|_{\mathbf{z}}$ , not forced to be zero by symmetry, have negative real parts. The corresponding periodic solution of (4.2) then has Floquet multipliers lying inside the unit circle (except for those forced to be one by symmetry) [6]. If the solution  $\mathbf{z}$  has symmetry  $\Sigma_{\mathbf{z}}$  then the group orbit has the dimension of  $(\Gamma_s \times S^1)/\Sigma_{\mathbf{z}}$  and the number of zero eigenvalues of  $\mathbf{D}\phi|_{\mathbf{z}}$  is

$$n_{\mathbf{z}} = 3 - \dim(\Sigma_{\mathbf{z}}). \quad (6.1)$$

Thus if the symmetry group  $\Sigma_{\mathbf{z}}$  is discrete, there are three zero eigenvalues: two associated with the  $T^2$  translation symmetries and one associated with the  $S^1$  phase shift symmetry. This is the case for the standing wave solutions SS, AR and SCR. In contrast  $\mathbf{D}\phi$  has two zero eigenvalues when evaluated on the TS and SR solution branches and one zero eigenvalue on the TR branch.

The form of the Jacobian matrix is restricted for solutions with nontrivial symmetry. In particular,  $\mathbf{D}\phi|_{\mathbf{z}}$  commutes with all elements of the isotropy subgroup  $\Sigma_{\mathbf{z}}$ , *i.e.*,

$$\sigma \mathbf{D}\phi|_{\mathbf{z}} = \mathbf{D}\phi|_{\mathbf{z}} \sigma \quad \text{for all } \sigma \in \Sigma_{\mathbf{z}}. \quad (6.2)$$

This restriction greatly simplifies the calculation of  $\mathbf{D}\phi|_{\mathbf{z}}$  and its eigenvalues, especially in the case where  $\mathbf{z}$  lies in a two-dimensional fixed point subspace. For these five solutions (TR, TS, SR, SS, AR) it is possible to put the Jacobian matrix into block diagonal form with each block consisting of a  $2 \times 2$  real matrix. This is most readily done by decomposing  $\mathbb{C}^4$  into subspaces, each of which is invariant under a different representation of the isotropy subgroup  $\Sigma_{\mathbf{z}}$ . Specifically, we form the isotypic decomposition [14]

$$\mathbb{C}^4 = W_0 \oplus W_1 \oplus \cdots \oplus W_k, \quad (6.3)$$

where each isotypic component  $W_i$  may be further decomposed into subspaces  $U_i^j$ , each of which transforms according to the  $i^{\text{th}}$  irreducible representation of  $\Sigma_{\mathbf{z}}$ :

$$W_i = U_i^1 \oplus U_i^2 \oplus \cdots \oplus U_i^{n_i}. \quad (6.4)$$

Thus  $W_i$  contains all irreducible subspaces of  $\mathbb{C}^4$  which are  $\Sigma_{\mathbf{z}}$ -isomorphic, meaning that there exists an isomorphism  $A : U_i^j \rightarrow U_i^l$  such that

$$A(\sigma u) = \sigma(Au) \quad \text{for all } u \in U_i^j \text{ and } \sigma \in \Sigma_{\mathbf{z}}. \quad (6.5)$$

The isotypic components in (6.3) are invariant under the mapping  $\mathbf{D}\phi|_{\mathbf{z}} : \mathbb{C}^4 \rightarrow \mathbb{C}^4$  [14], *i.e.*,

$$\mathbf{D}\phi|_{\mathbf{z}}(W_j) \subset W_j, \quad j = 1, \dots, k. \quad (6.6)$$

Hence the Jacobian matrix is block diagonal in these coordinates. Table 4 gives the isotypic decomposition of  $\mathbb{C}^4$  for each of the solutions in Table 3, together with the appropriate

representation of  $\Sigma_{\mathbf{z}}$  that acts on each isotypic component. Note that for each solution  $W_0 = \text{Fix}(\Sigma_{\mathbf{z}})$  so that  $W_0$  transforms according to the identity representation of  $\Sigma_{\mathbf{z}}$ . In addition the orbit of  $\mathbf{z}$  under the action of  $S^1$  (i.e.,  $\mathbf{z} \rightarrow e^{i\varphi}\mathbf{z}$ ) is in  $W_0$  so that there is a zero eigenvalue in  $\mathbf{D}\phi|_{W_0}$  for each solution  $\mathbf{z}$ . Here  $\mathbf{D}\phi|_{W_0}$  is  $\mathbf{D}\phi|_{\mathbf{z}}$  restricted to the  $W_0$  subspace. Table 5 gives the stabilities of the solutions to perturbations in each  $W_j$  for each solution with  $W_0 \approx \mathbb{C}$ .

Orbit Representative	Isotypic Components of $\mathbb{C}^4$	Representation of Isotropy Subgroup
Travelling Rolls (TR) $\mathbf{z} = (v_1, 0, 0, 0)$ $v_1 \neq 0$	$W_0^{TR} = \{(z_0, 0, 0, 0)\}$ $W_1^{TR} = \{(0, 0, z_1, 0)\}$ $W_2^{TR} = \{(0, z_1, 0, z_2)\}$	I $S^1$ $O(2) \times S^1$
Travelling Squares (TS) $\mathbf{z} = (v_1, v_1, 0, 0)$ $v_1 \neq 0$	$W_0^{TS} = \{(z_0, z_0, 0, 0)\}$ $W_1^{TS} = \{(z_1, -z_1, 0, 0)\}$ $W_2^{TS} = \{(0, 0, z_1, z_2)\}$	I $Z_2 = \langle \pi \rangle$ $Z_2 \times S^1 : Z_2 = \langle \kappa_0 \rangle$
Standing Rolls (SR) $\mathbf{z} = (v_1, 0, v_1, 0)$ $v_1 \neq 0$	$W_0^{SR} = \{(z_0, 0, z_0, 0)\}$ $W_1^{SR} = \{(z_1, 0, -z_1, 0)\}$ $W_2^{SR} = \{(0, z_1, 0, z_2)\}$	I $Z_2 = \langle \pi \rangle$ $O(2)$
Standing Squares (SS) $\mathbf{z} = (v_1, v_1, v_1, v_1)$ $v_1 \neq 0$	$W_0^{SS} = \{(z_0, z_0, z_0, z_0)\}$ $W_1^{SS} = \{(z_1, -z_1, z_1, -z_1)\}$ $W_2^{SS} = \{(z_1, z_2, -z_1, -z_2)\}$	I $Z_2 = \langle \pi \rangle$ $D_4$
Alternating Rolls (AR) $\mathbf{z} = (v_1, iv_1, v_1, iv_1)$ $v_1 \neq 0$	$W_0^{AR} = \{(z_0, iz_0, z_0, iz_0)\}$ $W_1^{AR} = \{(z_1, -iz_1, z_1, -iz_1)\}$ $W_2^{AR} = \{(z_1, -iz_2, -z_1, iz_2)\}$	I $Z_2 = \langle \pi \rangle$ $D_4$
Standing Cross-Rolls (SCR) $\mathbf{z} = (v_1, v_2, v_1, v_2)$ $v_1 \neq 0, v_2 \neq 0,  v_1  \neq  v_2 $	$W_0^{SCR} = \{(z_1, z_2, z_1, z_2)\}$ $W_1^{SCR} = \{(z_1, z_2, -z_1, -z_2)\}$	I $Z_2^2 = \langle \kappa_1, \kappa_2 \rangle$

Table 4: Isotypic decomposition of  $\mathbb{C}^4$  for each isotropy subgroup associated with a solution in Table 3. Here,  $\mathbf{z} = (v_1, v_2, w_1, w_2)$  and  $S^1 : \mathbf{z} \rightarrow e^{i\varphi}\mathbf{z}$ ,  $\pi : \mathbf{z} \rightarrow -\mathbf{z}$ ,  $O(2) = \langle \kappa_0, \theta \rangle$ , and  $D_4 = \langle \kappa_0, \kappa_1 \rangle$ , where  $\theta : (z_1, z_2) \rightarrow (e^{i\theta}z_1, e^{-i\theta}z_2)$ ,  $\kappa_0 : (z_1, z_2) \rightarrow (z_2, z_1)$ ,  $\kappa_1 : (z_1, z_2) \rightarrow (z_1, -z_2)$ , and  $\kappa_2 : (z_1, z_2) \rightarrow (-z_1, z_2)$ .



For those solutions with a two-dimensional fixed point subspace  $W_0$ , the nonzero eigenvalue of  $\mathbf{D}\phi|W_0$  is given by its trace. The eigenvector associated with this eigenvalue is in the plane of the limit cycle and transverse to it. The stability of the limit cycle to a perturbation in  $W_0$ , in this “radial direction”, depends on the direction of bifurcation of the solution branch. For the third order truncation (5.3), the radial eigenvalue is  $-2\lambda$ , just as in the Hopf bifurcation without symmetry. This is readily checked by computing  $\text{Tr}(\mathbf{D}\phi|W_0)$ . Consider, for example, travelling rolls for which  $(v_1, v_2, w_1, w_2) = (z_0, 0, 0, 0) \in W_0^{TR}$ . Treating  $z_0, \bar{z}_0$  as independent perturbations in  $W_0$  we obtain

$$(\mathbf{D}\phi|W_0^{TR}) \begin{pmatrix} z_0 \\ \bar{z}_0 \end{pmatrix} = \begin{pmatrix} \frac{\partial\phi_1}{\partial v_1} & \frac{\partial\phi_1}{\partial \bar{v}_1} \\ \frac{\partial\bar{\phi}_1}{\partial v_1} & \frac{\partial\bar{\phi}_1}{\partial \bar{v}_1} \end{pmatrix}_{TR} \begin{pmatrix} z_0 \\ \bar{z}_0 \end{pmatrix}. \quad (6.7)$$

Then

$$\text{Tr}(\mathbf{D}\phi|W_0^{TR}) = 2 \text{Re} \left( \frac{\partial\phi_1}{\partial v_1} \right)_{TR}, \quad (6.8)$$

where the TR subscript indicates that the Jacobian matrix is evaluated on the TR solution branch. We evaluate (6.8) using the third order truncation  $\Phi$  given by (5.3):

$$2 \text{Re} \left( \frac{\partial\Phi_1}{\partial v_1} \right)_{TR} = 2b_r |v_1|^2 = -2\lambda, \quad (6.9)$$

where the  $r$  subscript on  $b$  specifies the real part of  $b$  (similarly,  $b_i \equiv \text{Im}(b)$ ). Similar calculations for the solutions TS, SR, SS and AR give the same result. Specifically,  $\text{Tr}(\mathbf{D}\Phi|W_0) = -2\lambda$ . The remaining calculations performed in compiling Table 5 follow below. Throughout, we give first a general expression for the eigenvalues associated with a solution in terms of partial derivatives of the general equivariant vector field (5.2). These expressions are then evaluated using the third order truncation (5.3).

Let  $z_1, \bar{z}_1$  be independent perturbations in  $W_1 \approx \mathbb{C}$ , then  $\mathbf{D}\phi|W_1$  has the form

$$(\mathbf{D}\phi|W_1) \begin{pmatrix} z_1 \\ \bar{z}_1 \end{pmatrix} = \begin{pmatrix} \alpha & \beta \\ \bar{\beta} & \bar{\alpha} \end{pmatrix} \begin{pmatrix} z_1 \\ \bar{z}_1 \end{pmatrix}, \quad \alpha, \beta \in \mathbb{C}. \quad (6.10)$$

The eigenvalues of (6.10) are determined by computing the trace and determinant, where

$$\text{Tr}(\mathbf{D}\phi|W_1) = 2 \text{Re}(\alpha), \quad \det(\mathbf{D}\phi|W_1) = |\alpha|^2 - |\beta|^2. \quad (6.11)$$

It follows from (6.2) that the matrix in (6.10) commutes with the action of  $\Sigma_z$  on  $(z_1, \bar{z}_1)$ ,  $z_1, \bar{z}_1 \in W_1$ . For example, the action of  $\Sigma_{TR}$  on  $W_1^{TR}$  is that of  $S^1 = [(-\varphi, 0), \varphi]$  on  $(v_1, v_2, w_1, w_2) = (0, 0, z_1, 0)$ , namely

$$S^1 : W_1^{TR} \rightarrow W_1^{TR} : \begin{pmatrix} z_1 \\ \bar{z}_1 \end{pmatrix} \mapsto \begin{pmatrix} e^{2i\varphi} & 0 \\ 0 & e^{-2i\varphi} \end{pmatrix} \begin{pmatrix} z_1 \\ \bar{z}_1 \end{pmatrix}. \quad (6.12)$$

The condition (6.2) is

$$\begin{pmatrix} e^{2i\varphi} & 0 \\ 0 & e^{-2i\varphi} \end{pmatrix} \begin{pmatrix} \alpha & \beta \\ \bar{\beta} & \bar{\alpha} \end{pmatrix} = \begin{pmatrix} \alpha & \beta \\ \bar{\beta} & \bar{\alpha} \end{pmatrix} \begin{pmatrix} e^{2i\varphi} & 0 \\ 0 & e^{-2i\varphi} \end{pmatrix}, \quad (6.13)$$

for all  $\varphi \in S^1$ . It follows that  $\beta = 0$ . The real parts of the eigenvalues  $\alpha, \bar{\alpha}$  are

$$\text{Re}(\alpha)_{TR} = \text{Re}\left(\frac{\partial\phi_3}{\partial w_1}\right)_{TR} = a_r |v_1|^2. \quad (6.14)$$

For TS, SR, SS and AR, the action of the appropriate isotropy subgroup on each  $W_1$  is generated by a reflection  $(z_1, \bar{z}_1) \rightarrow -(z_1, \bar{z}_1)$ , where  $z_1$  is given in each case by

$$\begin{aligned} (\text{TS}) : (v_1, v_2, w_1, w_2) &= (z_1, -z_1, 0, 0) \in W_1^{TS} \\ (\text{SR}) : (v_1, v_2, w_1, w_2) &= (z_1, 0, -z_1, 0) \in W_1^{SR} \\ (\text{SS}) : (v_1, v_2, w_1, w_2) &= (z_1, -z_1, z_1, -z_1) \in W_1^{SS} \\ (\text{AR}) : (v_1, v_2, w_1, w_2) &= (z_1, -iz_1, z_1, -iz_1) \in W_1^{AR}. \end{aligned} \quad (6.15)$$

This representation commutes trivially with  $\mathbf{D}\phi|_{W_1}$  since it is a scalar multiple of the identity. The stability of each solution to perturbations in  $W_1$  is thus determined by calculating both the trace and determinant of  $\mathbf{D}\phi|_{W_1}$ . However, for both the TS and SR solutions, there is a null eigenvector in the corresponding  $W_1$  subspace. For travelling squares, the null eigenvector is in the direction of the group orbit of the solution  $(v_1, v_1, 0, 0)$  under the action of  $(\theta_1, -\theta_1) \in T^2$ . It is given by

$$\left. \frac{d}{d\theta_1} \right|_{\theta_1=0} \begin{pmatrix} e^{i\theta_1} v_1 \\ e^{-i\theta_1} v_1 \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} iv_1 \\ -iv_1 \\ 0 \\ 0 \end{pmatrix} \in W_1^{TS}, \quad (6.16)$$

and corresponds to a perturbation that translates the travelling squares pattern along the diagonal ( $x_1 = -x_2$ ). For standing rolls, the null eigenvector is associated with a translation of the pattern in the  $\hat{x}_1$ -direction. Hence it is determined from the group orbit of the solution  $(v_1, 0, v_1, 0)$  under  $(\theta_1, 0) \in T^2$  by

$$\left. \frac{d}{d\theta_1} \right|_{\theta_1=0} \begin{pmatrix} e^{i\theta_1} v_1 \\ 0 \\ e^{-i\theta_1} v_1 \\ 0 \end{pmatrix} = \begin{pmatrix} iv_1 \\ 0 \\ -iv_1 \\ 0 \end{pmatrix} \in W_1^{SR}. \quad (6.17)$$

The nonzero eigenvalues of  $\mathbf{D}\phi|_{W_1^{TS}}$  and  $\mathbf{D}\phi|_{W_1^{SR}}$  are then

$$\text{Tr}(\mathbf{D}\phi|_{W_1^{TS}}) = 2 \text{Re}(\alpha)_{TS} = 2\text{Re}\left(\frac{\partial\phi_1}{\partial v_1} - \frac{\partial\phi_1}{\partial v_2}\right)_{TS} = 2(b_r - c_r)|v_1|^2 \quad (6.18a)$$

$$\text{Tr}(\mathbf{D}\phi|_{W_1^{SR}}) = 2 \text{Re}(\alpha)_{SR} = 2\text{Re}\left(\frac{\partial\phi_1}{\partial v_1} - \frac{\partial\phi_1}{\partial w_1}\right)_{SR} = -2a_r |v_1|^2. \quad (6.18b)$$

The stability of standing squares to perturbations in  $W_1^{SS}$  depends on both  $\alpha$  and  $\beta$  in (6.10), where

$$\alpha = \left( \frac{\partial \phi_1}{\partial v_1} - \frac{\partial \phi_1}{\partial v_2} + \frac{\partial \phi_1}{\partial w_1} - \frac{\partial \phi_1}{\partial w_2} \right)_{SS} = (f - 3d)|v_1|^2 \quad (6.19a)$$

$$\beta = \left( \frac{\partial \phi_1}{\partial \bar{v}_1} - \frac{\partial \phi_1}{\partial \bar{v}_2} + \frac{\partial \phi_1}{\partial \bar{w}_1} - \frac{\partial \phi_1}{\partial \bar{w}_2} \right)_{SS} = (f + d)v_1^2 \quad (6.19b)$$

(recall  $f \equiv a + 2b - 2c$ ). Combining (6.11) and (6.19) gives

$$\text{Tr}(\mathbf{D}\phi|W_1^{SS}) = 2(f_r - 3d_r)|v_1|^2 \quad (6.20a)$$

$$\det(\mathbf{D}\phi|W_1^{SS}) = 8(|d|^2 - \text{Re}(\bar{d}f))|v_1|^4. \quad (6.20b)$$

The eigenvalues of  $\mathbf{D}\phi|_{AR}$  may be inferred from those of  $\mathbf{D}\phi|_{SS}$ . This is because of a parameter symmetry [11] in (5.3). Specifically, (5.3) is equivariant under the transformation

$$(v_1, v_2, w_1, w_2; d) \rightarrow (v_1, iv_2, w_1, iw_2; -d). \quad (6.21)$$

Thus the AR solution is obtained from the SS solution by letting  $d \rightarrow -d$  in (5.3), *i.e.*,

$$(v_1, v_1, v_1, v_1; d) \rightarrow (v_1, iv_1, v_1, iv_1; -d). \quad (6.22)$$

Combining (6.20) and (6.22) gives

$$\text{Tr}(\mathbf{D}\phi|W_1^{AR}) = 2(f_r + 3d_r)|v_1|^2 \quad (6.23a)$$

$$\det(\mathbf{D}\phi|W_1^{AR}) = 8(|d|^2 + \text{Re}(\bar{d}f))|v_1|^4. \quad (6.23b)$$

It remains to determine the stability of the solutions relative to perturbations in  $W_2 \approx \mathbb{C}^2$ . Since  $z_1, \bar{z}_1, z_2, \bar{z}_2$  are independent perturbations in  $W_2$ ,  $\mathbf{D}\phi|W_2$  has the form

$$(\mathbf{D}\phi|W_2) \begin{pmatrix} z_1 \\ \bar{z}_1 \\ z_2 \\ \bar{z}_2 \end{pmatrix} = \begin{pmatrix} A_1 & A_2 \\ A_3 & A_4 \end{pmatrix} \begin{pmatrix} z_1 \\ \bar{z}_1 \\ z_2 \\ \bar{z}_2 \end{pmatrix}, \quad (6.24a)$$

$$A_j = \begin{pmatrix} \alpha_j & \beta_j \\ \bar{\beta}_j & \bar{\alpha}_j \end{pmatrix}, \quad \alpha_j, \beta_j \in \mathbb{C}, \quad j = 1, 2, 3, 4. \quad (6.24b)$$

Let  $(z_1, z_2) \in \mathbb{C}^2$  be defined for each solution in Table 5 by

$$(TR), (SR) : (v_1, v_2, w_1, w_2) = (0, z_1, 0, z_2) \quad (6.25a)$$

$$(TS) : (v_1, v_2, w_1, w_2) = (0, 0, z_1, z_2) \quad (6.25b)$$

$$(SS), (AR) : (v_1, v_2, w_1, w_2) = (z_1, z_2, -z_1, -z_2). \quad (6.25c)$$

The action of  $\sigma_v \in O(2) \subset \Sigma_{TR}, \Sigma_{SR}$  on  $(z_1, z_2) \in W_2^{TR}, W_2^{SR}$  is the same as the action of  $(r_{\pi/2}\sigma_v) \in \Sigma_{TS}, \Sigma_{SS}$  on  $(z_1, z_2) \in W_2^{TS}, W_2^{SS}$ . Specifically,  $\sigma_v$  and  $(r_{\pi/2}\sigma_v)$  act on the appropriate  $W_2$  subspaces by reflection  $\kappa_0 : (z_1, z_2) \rightarrow (z_2, z_1)$ . It follows that  $\mathbf{D}\phi|_{W_2}$  commutes with

$$\begin{pmatrix} 0 & \mathbf{I} \\ \mathbf{I} & 0 \end{pmatrix}, \quad (6.26)$$

where  $\mathbf{I}$  and  $0$  are the  $2 \times 2$  identity and zero matrices, respectively. Hence, (6.24a) has the form

$$\mathbf{D}\phi|_{W_2} = \begin{pmatrix} A_1 & A_2 \\ A_2 & A_1 \end{pmatrix}. \quad (6.27)$$

Note that this matrix is similar to

$$\begin{pmatrix} A_1 + A_2 & 0 \\ 0 & A_1 - A_2 \end{pmatrix} = S(\mathbf{D}\phi|_{W_2})S^{-1}, \quad (6.28a)$$

where

$$S = S^{-1} = \frac{1}{\sqrt{2}} \begin{pmatrix} \mathbf{I} & \mathbf{I} \\ \mathbf{I} & -\mathbf{I} \end{pmatrix}. \quad (6.28b)$$

In addition to the reflection symmetry,  $O(2) \subset \Sigma_{TR}, \Sigma_{SR}$  contains a translational symmetry  $\theta : (z_1, z_2) \rightarrow (e^{i\theta}z_1, e^{-i\theta}z_2)$ . Thus  $\mathbf{D}\phi|_{W_2^{TR}}$  and  $\mathbf{D}\phi|_{W_2^{SR}}$  commute with

$$\begin{pmatrix} R_\theta & 0 \\ 0 & R_{-\theta} \end{pmatrix}, \quad R_\theta = \begin{pmatrix} e^{i\theta} & 0 \\ 0 & e^{-i\theta} \end{pmatrix}. \quad (6.29)$$

Therefore  $A_1, A_2$  in (6.27) satisfy

$$A_1 R_\theta = R_\theta A_1, \quad A_2 R_\theta = R_{-\theta} A_2, \quad (6.30)$$

and hence have the form

$$A_1 = \begin{pmatrix} \alpha & 0 \\ 0 & \bar{\alpha} \end{pmatrix}, \quad A_2 = \begin{pmatrix} 0 & \beta \\ \bar{\beta} & 0 \end{pmatrix}. \quad (6.31)$$

The eigenvalues of  $\mathbf{D}\phi|_{W_2^{SR}}$  are those of

$$(A_1 \pm A_2)_{SR} = \begin{pmatrix} \alpha & \pm\beta \\ \pm\bar{\beta} & \bar{\alpha} \end{pmatrix}_{SR}. \quad (6.32)$$

The stability of SR to perturbations in  $W_2^{SR}$  is therefore determined by

$$\text{Tr}(A_1 \pm A_2)_{SR} = 2 \text{Re}(\alpha)_{SR}, \quad \det(A_1 \pm A_2)_{SR} = (|\alpha|^2 - |\beta|^2)_{SR}, \quad (6.33)$$

where

$$(\alpha)_{SR} = \left( \frac{\partial \phi_2}{\partial v_2} \right)_{SR} = -f|v_1|^2, \quad (\beta)_{SR} = \left( \frac{\partial \phi_2}{\partial \bar{w}_2} \right)_{SR} = dv_1^2. \quad (6.34)$$

The eigenvalues of  $\mathbf{D}\phi|W_2^{SR}$  are degenerate with

$$\text{Tr}(A_1 \pm A_2)_{SR} = -2f_r|v_1|^2 \quad (6.35a)$$

$$\det(A_1 \pm A_2)_{SR} = (|f|^2 - |d|^2)|v_1|^4. \quad (6.35b)$$

The  $W_2^{TR}$  subspace has one additional symmetry; it is invariant under the  $S^1$  action  $[(-\varphi, 0), \varphi]$  that phase shifts  $(z_1, z_2)$ . Thus  $\mathbf{D}\phi|W_2^{TR}$  commutes with

$$\begin{pmatrix} R_\varphi & 0 \\ 0 & R_\varphi \end{pmatrix}. \quad (6.36)$$

where  $R_\varphi$  is defined as in (6.29). In this case conditions (6.30) are supplemented by

$$A_2 R_\varphi = R_\varphi A_2. \quad (6.37)$$

Then  $\beta = 0$  in (6.31) and  $\mathbf{D}\phi|W_2^{TR}$  is diagonal with repeated eigenvalues  $\alpha, \bar{\alpha}$ , where

$$(\alpha)_{TR} = \left( \frac{\partial \phi_2}{\partial v_2} \right)_{TR} = (c - b)|v_1|^2. \quad (6.38)$$

The  $W_2^{TS}$  subspace is invariant under the action of (6.36), but not under the action of (6.29). It follows that  $A_1, A_2$  in (6.27) have the form

$$A_j^{TS} = \begin{pmatrix} \alpha_j & 0 \\ 0 & \bar{\alpha}_j \end{pmatrix}_{TS}, \quad j = 1, 2. \quad (6.39)$$

Thus the eigenvalues are those of

$$(A_1 \pm A_2)_{TS} = \begin{pmatrix} \alpha_1 \pm \alpha_2 & 0 \\ 0 & \bar{\alpha}_1 \pm \bar{\alpha}_2 \end{pmatrix}_{TS}, \quad (6.40)$$

where

$$(\alpha_1)_{TS} = \left( \frac{\partial \phi_3}{\partial w_1} \right)_{TS} = a|v_1|^2, \quad (\alpha_2)_{TS} = \left( \frac{\partial \phi_3}{\partial w_2} \right)_{TS} = d|v_1|^2. \quad (6.41)$$

Hence the real parts of the eigenvalues of  $\mathbf{D}\phi|W_2^{TS}$  are

$$\text{Re}(\alpha_1 \pm \alpha_2)_{TS} = (a_r \pm d_r)|v_1|^2. \quad (6.42)$$

Finally,  $\mathbf{D}\phi|W_2^{SS}$  commutes with a reflection  $(z_1, z_2) \rightarrow (z_1, -z_2)$  so that  $A_2 = 0$  in (6.27) and the eigenvalues of  $\mathbf{D}\phi|W_2^{SS}$  have multiplicity two. One degenerate pair of eigenvalues is zero since the null eigenvectors associated with the group orbit of the SS solution under the  $T^2$  action reside in  $W_2^{SS}$ . These eigenvectors are

$$\left. \frac{d}{d\theta_1} \right|_{\theta_1=0} \begin{pmatrix} e^{i\theta_1} v_1 \\ v_1 \\ e^{-i\theta_1} v_1 \\ v_1 \end{pmatrix} = \begin{pmatrix} i v_1 \\ 0 \\ -i v_1 \\ 0 \end{pmatrix}, \quad \left. \frac{d}{d\theta_2} \right|_{\theta_2=0} \begin{pmatrix} v_1 \\ e^{i\theta_2} v_1 \\ v_1 \\ e^{-i\theta_2} v_1 \end{pmatrix} = \begin{pmatrix} 0 \\ i v_1 \\ 0 \\ -i v_1 \end{pmatrix}. \quad (6.43)$$

Solution	Null Eigenvectors	Eigenvalues	Stability
Travelling Rolls (TR) $\mathbf{z} = (v_1, 0, 0, 0)$	$(iv_1, 0, 0, 0) \in W_0^{TR}$	$W_0^{TR} : \{2 \operatorname{Re}(\frac{\partial \phi_1}{\partial v_1}), 0\}$ $W_1^{TR} : \{\mu_1, \bar{\mu}_1\}; \operatorname{Re}(\mu_1) = \operatorname{Re}(\frac{\partial \phi_2}{\partial w_1})$ $W_2^{TR} : \{\mu_2, \mu_2, \bar{\mu}_2, \bar{\mu}_2\}; \operatorname{Re}(\mu_2) = \operatorname{Re}(\frac{\partial \phi_2}{\partial v_2})$	$\operatorname{sgn}(b_r)$ $\operatorname{sgn}(a_r)$ $\operatorname{sgn}(a_r - f_r)$
Travelling Squares (TS) $\mathbf{z} = (v_1, v_1, 0, 0)$	$(iv_1, iv_1, 0, 0) \in W_0^{TS}$ $(iv_1, -iv_1, 0, 0) \in W_1^{TS}$	$W_0^{TS} : \{2 \operatorname{Re}(\frac{\partial \phi_1}{\partial v_1} + \frac{\partial \phi_1}{\partial v_2}), 0\}$ $W_1^{TS} : \{2 \operatorname{Re}(\frac{\partial \phi_1}{\partial v_1} - \frac{\partial \phi_1}{\partial v_2}), 0\}$ $W_2^{TS} : \{\mu_1, \mu_2, \bar{\mu}_1, \bar{\mu}_2\}; \operatorname{Re}(\mu_1) = \operatorname{Re}(\frac{\partial \phi_1}{\partial w_1} + \frac{\partial \phi_2}{\partial w_2}),$ $\operatorname{Re}(\mu_2) = \operatorname{Re}(\frac{\partial \phi_2}{\partial w_1} - \frac{\partial \phi_2}{\partial w_2})$	$\operatorname{sgn}(a_r + 4b_r - f_r)$ $\operatorname{sgn}(f_r - a_r)$ $\operatorname{sgn}(q_r + d_r)$ $\operatorname{sgn}(a_r - d_r)$
Standing Rolls (SR) $\mathbf{z} = (v_1, 0, v_1, 0)$	$(iv_1, 0, iv_1, 0) \in W_0^{SR}$ $(iv_1, 0, -iv_1, 0) \in W_1^{SR}$	$W_0^{SR} : \{2 \operatorname{Re}(\frac{\partial \phi_1}{\partial v_1} + \frac{\partial \phi_1}{\partial w_1}), 0\}$ $W_1^{SR} : \{2 \operatorname{Re}(\frac{\partial \phi_1}{\partial v_1} - \frac{\partial \phi_1}{\partial w_1}), 0\}$ $W_2^{SR} : \{\mu_1, \mu_1, \mu_2, \mu_2\}; \mu_1 + \mu_2 = 2 \operatorname{Re}(\frac{\partial \phi_2}{\partial v_2}),$ $\mu_1 \mu_2 =  \frac{\partial \phi_2}{\partial v_2} ^2 -  \frac{\partial \phi_2}{\partial w_2} ^2$	$\operatorname{sgn}(a_r + 2b_r)$ $\operatorname{sgn}(-a_r)$ $\operatorname{sgn}(-f_r)$ $\operatorname{sgn}( d ^2 -  f ^2)$

(Table 5 continued on the next page)

Solution	Null Eigenvectors	Eigenvalues	Stability
Standing Squares (SS) $\mathbf{z} = (v_1, v_1, v_1, v_1)$	$(iv_1, iv_1, iv_1, iv_1) \in W_0^{SS}$	$W_2^{SS} : \{2 \operatorname{Re}(\frac{\partial \phi_1}{\partial v_1} + \frac{\partial \phi_1}{\partial v_2} + \frac{\partial \phi_1}{\partial w_1} + \frac{\partial \phi_1}{\partial w_2}), 0\}$ $W_1^{SS} : \{\mu_1, \mu_2\}; \mu_1 + \mu_2 = 2 \operatorname{Re}(\alpha),$ $\mu_1 \mu_2 =  \alpha ^2 -  \beta ^2;$ $\alpha = (\frac{\partial \phi_1}{\partial v_1} - \frac{\partial \phi_1}{\partial v_2} + \frac{\partial \phi_1}{\partial w_1} + \frac{\partial \phi_1}{\partial w_2} - \frac{\partial \phi_1}{\partial w_2})$ $\beta = (\frac{\partial \phi_1}{\partial v_1} - \frac{\partial \phi_1}{\partial v_2} + \frac{\partial \phi_1}{\partial w_1} + \frac{\partial \phi_1}{\partial w_2} - \frac{\partial \phi_1}{\partial w_2})$ $W_2^{SS} : \{\mu, \mu, 0, 0\}; \mu = 2 \operatorname{Re}(\frac{\partial \phi_1}{\partial v_1} - \frac{\partial \phi_1}{\partial w_1})$	$\operatorname{sgn}(2a_r + 4b_r + d_r - f_r)$ $\operatorname{sgn}(f_r - 3d_r)$ $\operatorname{sgn}(\operatorname{Re}(\bar{d}f) -  d ^2)$ $\operatorname{sgn}(-d_r - a_r)$
Alternating Rolls (AR) $\mathbf{z} = (v_1, iv_1, v_1, iv_1)$	$(iv_1, -v_1, iv_1, -v_1) \in W_0^{AR}$	$W_0^{AR} : \{2 \operatorname{Re}(\frac{\partial \phi_1}{\partial v_1} + i \frac{\partial \phi_1}{\partial v_2} + \frac{\partial \phi_1}{\partial w_1} + \frac{\partial \phi_1}{\partial w_2} + i \frac{\partial \phi_1}{\partial w_2}), 0\}$ $W_1^{AR} : \{\mu_1, \mu_2\}; \mu_1 + \mu_2 = 2 \operatorname{Re}(\alpha),$ $\mu_1 \mu_2 =  \alpha ^2 -  \beta ^2;$ $\alpha = (\frac{\partial \phi_1}{\partial v_1} - i \frac{\partial \phi_1}{\partial v_2} + \frac{\partial \phi_1}{\partial w_1} + \frac{\partial \phi_1}{\partial w_2} - i \frac{\partial \phi_1}{\partial w_2})$ $\beta = (\frac{\partial \phi_1}{\partial v_1} + i \frac{\partial \phi_1}{\partial v_2} + \frac{\partial \phi_1}{\partial w_1} + \frac{\partial \phi_1}{\partial w_2} + i \frac{\partial \phi_1}{\partial w_2})$ $W_2^{AR} : \{\mu, \mu, 0, 0\}; \mu = 2 \operatorname{Re}(\frac{\partial \phi_1}{\partial v_1} - \frac{\partial \phi_1}{\partial w_1})$	$\operatorname{sgn}(2a_r + 4b_r + d_r - f_r)$ $\operatorname{sgn}(f_r + 3d_r)$ $\operatorname{sgn}(-\operatorname{Re}(\bar{d}f) -  d ^2)$ $\operatorname{sgn}(d_r - a_r)$

Table 5: Eigenvalues of the Jacobian matrix  $D\phi$  evaluated on the solution branches guaranteed by the equivariant Hopf theorem. A solution to the third order truncation (5.3) is stable if the signed quantities in the fourth column are all negative. The SCR solution, not listed above, is unstable. (Recall  $f \equiv a + 2b - 2c$ .)

Hence the stability of the SS solution to perturbations in  $W_2^{SS}$  is determined by  $\text{Tr}(\mathbf{D}\phi|W_2^{SS}) = 4 \text{Re}(\alpha)_{SS}$ , where

$$(\alpha)_{SS} = \left( \frac{\partial\phi_1}{\partial v_1} - \frac{\partial\phi_1}{\partial w_1} \right)_{SS} = -(d+a)|v_1|^2. \quad (6.44)$$

It follows from the parameter symmetry (6.22) that the stability of the AR solution to perturbations in  $W_2^{AR}$  depends on

$$\text{Tr}(\mathbf{D}\phi|W_2^{AR}) = 4(d_r - a_r)|v_1|^2. \quad (6.45)$$

So far we have considered the stability of those solutions guaranteed by the equivariant Hopf theorem, *i.e.*, those with  $W_0 \approx \mathbb{C}$ . However, for coefficient values in (5.3) satisfying

$$|f|^2 > |d|^2 > |\text{Re}(\bar{d}f)|, \quad (6.46)$$

there is an additional solution (SCR) for which  $W_0^{SCR} \approx \mathbb{C}^2$ . The SCR solution was determined to be unstable in the  $W_0^{SCR}$  subspace by Swift [11]. His stability argument hinges on the observation that the SCR existence conditions (6.46) are the same as the conditions for (6.20b), (6.23b) and (6.35b) to be positive. Specifically, consider the bifurcations that occur as  $d$  and  $f$  are varied through a point in the  $(d, f)$ -parameter space satisfying one of the following degeneracy conditions:

$$\begin{aligned} |d|^2 &= \text{Re}(\bar{d}f) \\ |d|^2 &= -\text{Re}(\bar{d}f) \\ |d|^2 &= |f|^2. \end{aligned} \quad (6.47)$$

These are surfaces in the  $(d, f)$ -parameter space where pitchfork bifurcations occur along the SS, AR and SR solution branches, respectively. Two SCR solutions (both on the same group orbit) are created at the pitchfork bifurcations. Swift invokes the exchange of stability at pitchfork bifurcations to infer the instability of the SCR solution branch. We now calculate the eigenvalues of  $\mathbf{D}\Phi|_{SCR}$  directly for the cubic truncation (5.3).

The isotropy subgroup  $\Sigma_{SCR}$  acts trivially on the  $W_0^{SCR}$  subspace and hence does not restrict the form of  $\mathbf{D}\phi|W_0^{SCR}$ . However, two of the four eigenvalues are known since they are associated with eigenvectors in the phase and radial directions of the limit cycle. Thus the eigenvalues of  $\mathbf{D}\Phi|W_0^{SCR}$  are  $0, -2\lambda, \mu_1, \mu_2$  and the characteristic polynomial associated with  $\mathbf{D}\Phi|W_0^{SCR}$  may be formally factored

$$\mu^4 + a_1\mu^3 + a_2\mu^2 + a_3\mu = \mu(\mu + 2\lambda)(\mu - \mu_1)(\mu - \mu_2) = 0. \quad (6.48)$$

The sum and product of  $\mu_1$  and  $\mu_2$  may then be expressed in terms of  $\lambda, a_1, a_2$  as follows

$$\mu_1 + \mu_2 = 2\lambda - a_1, \quad (6.49a)$$

$$\mu_1\mu_2 = 2\lambda(2\lambda - a_1) + a_2. \quad (6.49b)$$



These quantities were computed using the computer algebra program, Macsyma:

$$\mu_1 + \mu_2 = - \left[ \frac{3d_i(|f|^2 - |d|^2)}{\text{Im}(\bar{d}f)} + 2f_r \right] (|v_1|^2 + |v_2|^2) \quad (6.50a)$$

$$\mu_1 \mu_2 = \frac{8(|d|^2 - \text{Re}(\bar{d}f))(|d|^2 + \text{Re}(\bar{d}f))(|f|^2 - |d|^2)|v_1|^4}{|d|^2(|f|^2 - |d|^2) - 4(\text{Im}(\bar{d}f))^2|v_1|^2/(|v_1|^2 + |v_2|^2)}. \quad (6.50b)$$

It follows from the inequalities (6.46) that the numerator of (6.50b) is positive. The denominator is negative, as demonstrated by the following chain of inequalities

$$\begin{aligned} \text{denominator} &< |d|^2(|f|^2 - |d|^2) - 2(\text{Im}(\bar{d}f))^2 \\ &= -[|d|^2(|f|^2 + |d|^2) - 2(\text{Re}(\bar{d}f))^2] \\ &< -2[|d|^4 - (\text{Re}(\bar{d}f))^2] < 0, \end{aligned} \quad (6.51)$$

where, without loss of generality, we have assumed  $(|v_1|^2 + |v_2|^2) < 2|v_1|^2$  in (6.50b). This explicit calculation confirms Swift's result that SCR are unstable.

We now calculate the remaining eigenvalues of  $\mathbf{D}\Phi|_{SCR}$ . Let  $(z_1, z_2) \in W_1^{SCR}$  be defined by  $(v_1, v_2, w_1, w_2) = (z_1, z_2, -z_1, -z_2)$ ; then  $\Sigma_{SCR}$  acts on  $(z_1, z_2)$  by the reflections  $\kappa_1, \kappa_2$ , where  $\kappa_1 : (z_1, z_2) \rightarrow (z_1, -z_2)$  and  $\kappa_2 : (z_1, z_2) \rightarrow (-z_1, z_2)$ . The matrix  $\mathbf{D}\phi|_{W_1^{SCR}}$ , of the form (6.24), commutes with matrices

$$\begin{pmatrix} \epsilon \mathbf{I} & 0 \\ 0 & \delta \mathbf{I} \end{pmatrix}, \quad (6.52)$$

where  $\epsilon, \delta$  take on values of  $\pm 1$ . It follows that

$$\mathbf{D}\phi|_{W_1^{SCR}} = \begin{pmatrix} A_1 & 0 \\ 0 & A_4 \end{pmatrix}, \quad A_j = \begin{pmatrix} \alpha_j & \beta_j \\ \bar{\beta}_j & \bar{\alpha}_j \end{pmatrix}, \quad j = 1, 4. \quad (6.53)$$

As in the case of  $\mathbf{D}\phi|_{W_2^{SS}}$ , there are two zero eigenvalues in  $\mathbf{D}\phi|_{W_1^{SCR}}$  associated with translations of the pattern under  $T^2$ . The null eigenvectors, determined by (6.43), are  $z_1 = (iv_1, 0)$ ,  $z_2 = (0, iv_2)$ . Hence  $A_1$  and  $A_4$  each contain a zero eigenvalue. The stability of SCR to perturbations in  $W_1^{SCR}$  is determined by

$$\text{Tr}(A_j)_{SCR} = 2\text{Re}(\alpha_j)_{SCR}, \quad j = 1, 4, \quad (6.54)$$

where

$$(\alpha_1)_{SCR} = \left( \frac{\partial \phi_1}{\partial v_1} - \frac{\partial \phi_1}{\partial w_1} \right)_{SCR} = \nu + bN_1 + cN_2 \quad (6.55a)$$

$$(\alpha_4)_{SCR} = \left( \frac{\partial \phi_2}{\partial v_2} - \frac{\partial \phi_2}{\partial w_2} \right)_{SCR} = \nu + bN_2 + cN_1. \quad (6.55b)$$

Substituting for  $N_1, N_2$  on the SCR solution branch we obtain the following two nonzero eigenvalues of  $\mathbf{D}\Phi|W_1^{SCR}$ :

$$-\left[ f_r + a_r + \frac{d_i(|f|^2 - |d|^2) \pm (f_r - a_r)|d|^2 \sin(2\psi)}{\text{Im}(\bar{d}f)} \right] (|v_1|^2 + |v_2|^2), \quad (6.56)$$

where  $\sin(2\psi)$  is given in Table 3.

## 7. Bifurcation Diagrams.

The previous section determined that the solution stabilities depend on the following six nonlinear coefficients

$$a_r, b_r, d_r, f_r, d_i, f_i, \quad (7.1)$$

where  $f$  has been chosen in favor of  $c = \frac{1}{2}(a + 2b - f)$ . In this section we neglect the unstable SCR solution in our classification of the bifurcation problems; the stability results for the remaining five solutions, summarized in Table 5, suggest the following nondegeneracy conditions on (5.3):

$$b_r \neq 0; \quad a_r \neq 0, f_r, \pm d_r, f_r - 4b_r, -2b_r, -2b_r + \frac{1}{2}(f_r \pm d_r); \quad (7.2a)$$

$$|f|^2 \neq |d|^2 \neq \pm \text{Re}(\bar{d}f); \quad (7.2b)$$

$$f_r \neq 0, \pm 3d_r. \quad (7.2c)$$

The condition  $f_r = 0$  defines a degeneracy in (5.3) only if  $|f|^2 > |d|^2$ . In this case the eigenvalues  $\mu_1, \mu_2$  of  $\mathbf{D}\phi|W_2^{SR}$  are purely imaginary with  $\mu_1 = -\mu_2$ . Similarly, the nondegeneracy conditions  $f_r \neq 3d_r, -3d_r$  are only necessary if  $|d|^2 > \text{Re}(\bar{d}f), -\text{Re}(\bar{d}f)$ , respectively.

The inequalities (7.2) divide the six-dimensional parameter space (7.1) into regions characterized by distinct bifurcation diagrams. The dependence of the bifurcation diagrams on  $f_i, d_i$  is summarized in Figure 5 where  $d$  is plotted in the complex  $f$  plane. For the remainder of this section we restrict our analysis to the case  $d_r > 0$ . The parameter symmetry (6.21) allows us to infer the results for  $d_r < 0$  from those obtained for positive  $d_r$ . It remains to determine the bifurcation diagrams in the various regions of the  $(a_r, b_r, f_r)$ -parameter space defined by the inequalities (7.2a). First, however, we review the bifurcation results for the four-dimensional invariant subspaces (*i.e.*, entries VII-XI in Table 1).

As noted in section 3, equation (5.3) restricted to subspace VII is  $D_4 \times S^1$ -equivariant; it has the form

$$\begin{aligned} 0 &= \nu v_1 + (a + 2b - f)(|v_1|^2 + |v_2|^2)v_1 + f|v_1|^2 v_1 + dv_2^2 \bar{v}_1 \\ 0 &= \nu v_2 + (a + 2b - f)(|v_1|^2 + |v_2|^2)v_2 + f|v_2|^2 v_2 + dv_1^2 \bar{v}_2. \end{aligned} \quad (7.3)$$




Figure 5: The complex coefficient  $d$  is plotted in the complex  $f$  plane. The nondegeneracy conditions (7.2b) divide this plane into five regions, each labeled by its distinguishing feature.

The nontrivial solutions of (7.3) are SR, SS, AR and SCR. The bifurcation diagrams for the SR, SS and AR solutions are given in Figure 6 for the various regions in the  $(a_r + 2b_r, f_r)$ -plane. Here, stability is determined relative to perturbations in subspace VII only. The results for  $d_r < 0$  are obtained by interchanging the SS and AR labels on the bifurcation diagrams in Figure 6. Note that if any two solution branches are subcritical, then all three are unstable. If, on the other hand, all three branches are supercritical then it is possible for two of the three solutions to be stable.

In each of the subspaces VIII-XI, the restricted form of (5.3) on  $\mathbb{C}^2$  is  $O(2) \times S^1$ -equivariant [6]:

$$\begin{aligned} 0 &= \nu z_1 + \alpha |z_2|^2 z_1 + \beta (|z_1|^2 + |z_2|^2) z_1 \\ 0 &= \nu z_2 + \alpha |z_1|^2 z_2 + \beta (|z_1|^2 + |z_2|^2) z_2. \end{aligned} \tag{7.4}$$




Figure 6: Bifurcation diagrams associated with (7.3). The solid lines in the  $(a_r + 2b_r, f_r)$ -plane are  $(a_r + 2b_r) = 0, (f_r \pm |d_r|)/2$ , the dashed lines are  $f_r = 0, \pm 3|d_r|$ . (See the text for a discussion of when the dashed lines represent degeneracies.) The bifurcation diagrams give the direction of bifurcation and stability of the solution branches for each region of the  $(a_r + 2b_r, f_r)$ -plane. Dashed bifurcation curves indicate unstable solutions. An asterisk on a solution indicates that it is possibly stable (stability depends on where  $f$  lies in Figure 5). The bifurcation diagrams are plotted for  $d_r > 0$ .

Subspace	$\alpha$	$\beta$	S	R
VIII. $(z_1, z_2) = (v_1, v_2)$ $w_1 = w_2 = 0$	$\frac{1}{2}(a - f)$	$b$	TS	TR
IX. $(z_1, z_2) = (v_1, w_1)$ $v_2 = w_2 = 0$	$a$	$b$	SR	TR
X. $(z_1, z_2) = (v_1, w_1)$ $v_1 = v_2, w_1 = w_2$	$a + d$	$\frac{1}{2}(a + 4b - f)$	SS	TS
XI. $(z_1, z_2) = (v_1, w_1)$ $v_1 = -iv_2, w_1 = -iw_2$	$a - d$	$\frac{1}{2}(a + 4b - f)$	AR	TS*

Table 6: Correspondence between the coefficients and solution branches of (7.4) and those of (5.3) restricted to the fixed point subspaces VIII-XI. Here, TS\* is a solution on the group orbit of TS.




Figure 7: Bifurcation diagrams in  $(\alpha_r, \beta_r)$ -plane.

There are two nontrivial solution types in this subspace:



$$\begin{aligned} \text{(S)} : |z_1| = |z_2| \neq 0 \\ \text{(R)} : |z_1| \neq 0, z_2 = 0. \end{aligned} \tag{7.5}$$

The stabilities of these solutions depend on  $\alpha_r, \beta_r$ . In Table 6 the correspondence between  $(\alpha, \beta)$  and the cubic coefficients of (5.3) is given for each of the subspaces VIII–XI, together with the solutions corresponding to S and R in (7.5). In Figure 7 the bifurcation diagrams are given in the  $(\alpha_r, \beta_r)$ -plane. Note that both solutions are unstable if either is subcritical and that one and only one is stable if both branches bifurcate supercritically. Hence, for TS to be stable, we need the TS, TR, SS and AR branches all to bifurcate supercritically. In addition, if the TS branch is stable, then the TR, SS, and AR solutions are unstable. By similar reasoning, the TR branch is unstable if TR, SR or TS is subcritical. Note that the travelling wave solutions TS and TR cannot both be stable.



Finally, we summarize the bifurcation results in the  $(a_r, b_r, f_r)$ -coefficient space for all five nontrivial, maximally symmetric solutions (TR, TS, SR, SS, AR). All possible bifurcation diagrams are obtained by considering six slices through the  $(a_r, b_r, f_r)$ -space at fixed  $f_r$  satisfying the inequalities:

$$\begin{aligned} \text{(I)} \quad & f_r < -3d_r < 0 \\ \text{(II)} \quad & -3d_r < f_r < -d_r < 0 \\ \text{(III)} \quad & -d_r < f_r < 0 \\ \text{(IV)} \quad & 0 < f_r < d_r \\ \text{(V)} \quad & 0 < d_r < f_r < 3d_r \\ \text{(VI)} \quad & 0 < 3d_r < f_r. \end{aligned} \tag{7.6}$$

The  $(a_r, b_r)$ -planes are divided into 33 regions each by the nine degeneracy lines defined by (7.2a) (see Figure 8). The bifurcation diagrams appropriate to the various regions are given in Figure 9. Note that it is possible to have two stable solutions bifurcate from the trivial solution when all five branches are supercritical (regions 5, 20 and 27 of Figure 9). It is also possible for all five branches to bifurcate supercritically with none being stable (regions 6 and 35 of Figure 9). The dynamics associated with these regions of the coefficient space has not been fully determined. However, in the next section we indicate why we expect the behavior to be complex and explore the possibility of a primary bifurcation to a structurally stable heteroclinic cycle connecting three of the periodic solutions found in section 5. The structural stability of the cycle is a consequence of symmetry (see, for example, Guckenheimer and Holmes [16]).



(Figure 8 continued on the next page)



(Figure 8 continued on the next page)







Figure 8: The  $(a_r, b_r)$ -plane partitioned by the nondegeneracy conditions (7.2a) for  $d_r$  positive, and  $f_r$  satisfying the inequalities (7.6) as labeled. The labeled lines are (A)  $a_r = -2b_r + (f_r - d_r)/2$ , (B)  $a_r = -2b_r + (f_r + d_r)/2$ , (C)  $a_r = -2b_r$ , and (D)  $a_r = -4b_r + f_r$ . The bifurcation diagrams are given in Figure 9. The darkened regions and those regions labeled with a superscript  $a$  or  $b$  are discussed in section 8.




Figure 9: Bifurcation diagrams indicating direction of bifurcation and stability for the numbered regions in Figure 8. Solid lines correspond to stable or possibly stable solutions; as in Figure 6, an asterisk means that the solution stability depends on where  $f$  lies in Figure 5.

## 8. Heteroclinic Cycles.

In this section we investigate some of the complicated dynamics expected in certain regions of the coefficient space associated with the cubic truncation of (4.2). We begin by indicating why we expect a nontrivial attractor to be associated with regions of the coefficient space where all branches are supercritical and unstable (regions 6 and 35 in Figure 9). In this analysis we also assume that one of the following inequalities holds:

$$a_r < 0 \quad \text{or} \quad a_r + 2b_r - f_r < 0. \quad (8.1)$$

The second of these conditions is satisfied for regions labeled 35 in Figure 8 and for those labeled 6 in cross-section (III) of Figure 8 (*i.e.*, when  $f_r > -|d_r|$ ). The inequalities (8.1) are not satisfied in the blackened portions of the regions labeled 6 in Figure 8 (I), (II). Note that for SS (SR) to be unstable in region 6 (35), we are necessarily in a part of the coefficient space where the SCR solution does not exist (*cf.* Figure 5).

Let the radius of a hypersphere in  $\mathbf{R}^8$  be denoted by  $\mathcal{R} \geq 0$ , where

$$\mathcal{R}^2 \equiv |v_1|^2 + |v_2|^2 + |w_1|^2 + |w_2|^2. \quad (8.2)$$

We wish to show that  $\dot{\mathcal{R}} < 0$  for sufficiently large  $\mathcal{R}$  in the regions specified above, in which case trajectories starting near the origin cannot escape to infinity. We call the resulting attractor nontrivial since it is neither the fixed point at the origin nor one of the periodic orbits created at the Hopf bifurcation, these being unstable in the regions of interest. From the cubic truncation of (4.2), it follows that

$$\dot{\mathcal{R}} = \lambda + \frac{1}{\mathcal{R}} [2a_r(|v_1|^2|w_1|^2 + |v_2|^2|w_2|^2) + b_r\mathcal{R}^4 + (a_r - f_r)N_1N_2 + 4d_r \operatorname{Re}(Q)] \quad (8.3a)$$

$$\leq \lambda + \frac{1}{\mathcal{R}} [2a_r(|v_1|^2|w_1|^2 + |v_2|^2|w_2|^2) + b_r\mathcal{R}^4 + (a_r - f_r)N_1N_2 + 4|d_rQ|], \quad (8.3b)$$

for  $\mathcal{R} \neq 0$ . We introduce the following nonlinear coordinates:

$$\mathcal{Q} \equiv |Q| \geq 0, \quad (8.4a)$$

$$\mathcal{T} \equiv (|v_1||v_2| - |w_1||w_2|)^2 + (|v_1||w_2| - |w_1||v_2|)^2 \geq 0, \quad (8.4b)$$

$$\mathcal{U} \equiv (|v_1||w_1| - |v_2||w_2|)^2 \geq 0, \quad (8.4c)$$

$$\mathcal{V} \equiv [(|v_1| - |w_1|)^2 - (|v_2| - |w_2|)^2] [(|v_1| + |w_1|)^2 - (|v_2| + |w_2|)^2]. \quad (8.4d)$$

These quantities are of the same order (in the amplitudes  $|v_j|$ ,  $|w_j|$ ) as  $\mathcal{R}^4$ . Note that  $\mathcal{T}$  vanishes on all solution branches except for TS,  $\mathcal{U}$  vanishes on all but SR,  $\mathcal{V}$  vanishes on all

but TR, and  $\mathcal{Q}$  vanishes on all but SS and AR. The inequality (8.3b), expressed in terms of the coordinates (8.4), is

$$\dot{\mathcal{R}} \leq \lambda + \frac{1}{\mathcal{R}} [b_r \mathcal{V} + (a_r + 4b_r - f_r) \mathcal{T} + 2(a_r + 2b_r) \mathcal{U} + 4(2a_r + 4b_r - f_r + |d_r|) \mathcal{Q}] \quad (8.5a)$$

$$= \lambda + \frac{1}{\mathcal{R}} [b_r (\mathcal{V} + 4\mathcal{U}) + (a_r + 4b_r - f_r) \mathcal{T} + 2a_r \mathcal{U} + 4(2a_r + 4b_r - f_r + |d_r|) \mathcal{Q}] \quad (8.5b)$$

$$= \lambda + \frac{1}{\mathcal{R}} [b_r (\mathcal{V} + 2\mathcal{T}) + (a_r + 2b_r - f_r) \mathcal{T} + 2(a_r + 2b_r) \mathcal{U} + 4(2a_r + 4b_r - f_r + |d_r|) \mathcal{Q}]. \quad (8.5c)$$

A consequence of our choice of coordinates (8.4) is that the coefficients of the nonconstant terms on the right-hand-side of (8.5a) determine the direction of branching of the solutions (*e.g.*, TS is supercritical when the coefficient of  $\mathcal{T}$  is negative). For the case of interest here, all branches are supercritical and hence the coefficients are all negative:

$$b_r < 0, \quad a_r + 4b_r - f_r < 0, \quad a_r + 2b_r < 0, \quad 2a_r + 4b_r - f_r + |d_r| < 0. \quad (8.6)$$

That  $\dot{\mathcal{R}}$  is negative for sufficiently large  $\mathcal{R}$ , and for coefficient values satisfying (8.1) and (8.6), follows from the observation that  $\mathcal{V} + 4\mathcal{U}$  in (8.5b) and  $\mathcal{V} + 2\mathcal{T}$  in (8.5c) are nonnegative:

$$\mathcal{V} + 4\mathcal{U} = (|v_1|^2 + |w_1|^2 - |v_2|^2 - |w_2|^2)^2 \geq 0 \quad (8.7a)$$

$$\mathcal{V} + 2\mathcal{T} = (|v_1|^2 - |w_1|^2)^2 + (|v_2|^2 - |w_2|^2)^2 \geq 0. \quad (8.7b)$$

Hence each of the nonconstant terms on the right-hand-side of (8.5b) is negative if the first of the inequalities (8.1) is satisfied. (Similarly, each is negative in (8.5c) if the second inequality holds.) Since these terms dominate at large radius,  $\mathcal{R}$  cannot grow without bound. We now investigate the possibility of a heteroclinic cycle being associated with certain of these regions of the coefficient space.

Given the isotropy subgroup lattice in Figure 3, it is straightforward to enumerate all possible heteroclinic cycles. In particular, it is clear that the cycles must contain the TS solution; these possibilities are depicted in Figure 10. Note that in specifying a heteroclinic cycle we identify solutions with conjugate isotropy subgroups, *i.e.*, solutions on the same group orbit. The observation that the vector field (4.2) restricted to any of the fixed point subspaces VIII–XI is  $O(2) \times S^1$ -equivariant may be used to determine the conditions for existence of an orbit connecting the periodic solutions residing in these subspaces. In particular, one can analyze the two-dimensional real vector field (4.12) to determine that generically there is a saddle–sink connection whenever the branches bifurcate supercritically (see, for example, Melbourne *et al.* [17]). The problem of proving that there exists a heteroclinic cycle is thereby reduced to that of showing there is an orbit connecting the appropriate solutions in the SCR subspace. In order to investigate this possibility, we recall certain results of




Figure 10: Possible heteroclinic cycles. Three additional possibilities are obtained by interchanging AR and SS. These may be treated using the parameter symmetry.

Swift on the  $D_4 \times S^1$ -equivariant Hopf bifurcation problem [11], which applies in the SCR subspace. In particular, Swift was able to project the dynamics associated with the third order truncation on to the surface of a sphere and in this way to obtain a two-dimensional flow. Specifically, the following coordinate transformation of  $(v_1, v_2) \in \mathbb{C}^2$  was introduced:

$$\begin{aligned} v_1 &= \sqrt{r} \cos(\theta/2) e^{i(\phi+\psi)/2} \\ v_2 &= \sqrt{r} \sin(\theta/2) e^{i(-\phi+\psi)/2}. \end{aligned} \tag{8.8}$$

When this coordinate transformation is applied to the cubic truncation of (4.2), restricted to the SCR subspace, we obtain

$$\begin{aligned}\dot{r} &= r \{ 2\lambda + r [ 2(a_r + 2b_r - f_r) + f_r(1 + \cos^2 \theta) + d_r \sin^2 \theta \cos 2\phi ] \} \\ \dot{\theta} &= r \sin \theta [ \cos \theta (-f_r + d_r \cos 2\phi) - d_i \sin 2\phi ] \\ \dot{\phi} &= r [ \cos \theta (f_i - d_i \cos 2\phi) - d_r \sin 2\phi ].\end{aligned}\tag{8.9}$$

As noted by Swift, the radial dependence in the  $(\dot{\theta}, \dot{\phi})$  equation may be removed by rescaling time  $t \rightarrow t/r$ . In this case, the  $(\theta', \phi')$  equation decouples from the  $r'$  equation (where the prime indicates differentiation with respect to the scaled time). The flow may then be studied on  $S^2$ . The fixed points of the “associated spherical system” may be identified with the SR, SS, and AR solutions (see Figure 11a).

Before investigating the possibility of heteroclinic cycle I in Figure 10 we rule out possibilities II and III. This is accomplished by noting that there does not exist a connecting orbit between the solutions in the SCR subspace whenever the other connections are present. For example, it follows from Table 6 and Figure 7 that there are saddle–sink connections  $\text{SR} \rightarrow \text{TR}$ ,  $\text{TR} \rightarrow \text{TS}$ , and  $\text{TS} \rightarrow \text{SS}$  in subspaces IX, VIII, and X, respectively, if and only if

$$\begin{aligned}a_r < 0, \quad b_r < 0, \quad a_r - f_r > 0, \quad a_r + 4b_r - f_r < 0, \\ a_r + d_r > 0, \quad a_r + 2b_r + \frac{1}{2}(d_r - f_r) < 0.\end{aligned}\tag{8.10a}$$

These inequalities confine us to regions of the cubic coefficient space where bifurcation diagram 6 of Figure 9 applies (*cf.* Figure 8). In order for there to be a connecting orbit  $\text{SS} \rightarrow \text{SR}$  in the fixed point subspace SCR, the SS solution must be unstable, which is the case only if

$$\text{Re}(f\bar{d}) > |d|^2.\tag{8.10b}$$

However, in this parameter regime SR is unstable, in the SCR subspace, in directions transverse to the “radial direction”. Specifically, in the associated spherical system, the SS solution is a saddle and the SR solution is a source. Thus there cannot be a connecting orbit from the SS solution to the SR solution; heteroclinic cycle II does not appear (generically) in a primary bifurcation. A similar argument rules out the possibility of a primary bifurcation to heteroclinic cycle III.

In order for heteroclinic cycle I to exist, the SS solution must have an unstable direction in the SCR subspace and the AR solution must have at least one stable eigenvalue (in addition to the radial eigenvalue). The possibility of a heteroclinic cycle with a source  $\rightarrow$  sink connection in the associated spherical system may be excluded since it is not consistent with  $\text{AR} \rightarrow \text{TS} \rightarrow \text{SS}$  connections. We will also not pursue the possibility of heteroclinic cycle I having a source  $\rightarrow$  saddle connection in the associated spherical system since we do not expect






Figure 11: (a) Coordinate system for associated spherical system, shown in perspective, and in the  $(\phi, \theta)$ -plane. The SR, SS, and AR solutions appear as pairs of fixed-points (one on the group orbit of the other). (b) Phase portrait showing a saddle-sink connection in the associated spherical system (reproduced from [11]). (c) Phase portrait showing limit cycle in associated spherical system. This corresponds to a quasiperiodic solution in the SCR subspace (also from [11]).

such a cycle to be asymptotically stable. The final possibility of a saddle→sink connecting orbit in the associated spherical system exists for coefficient values in cross-section (I) of Figure 8. The condition that there be connections  $AR \rightarrow TS \rightarrow SS$  confines our analysis further

to those regions of  $(I)$  labeled 6. (Note that for SS to be a saddle inequality (8.10b) must hold.) In the associated spherical system the SR solution is then identified with a source at the north pole of the sphere (the source at the south pole is on its group orbit), the SS (AR) solution is a saddle (sink) located on the equator of the sphere at  $\phi = 0$  ( $\phi = \pi/2$ ). In Figure 11b, we reproduce a phase portrait from Swift's paper in which there appears to be a connecting orbit from the SS solution to the AR solution. This connection is certainly expected for  $d_r$  and  $d_i$  sufficiently small. This is seen by noting that at the degenerate point  $d = 0$  there is an invariant circle of equilibria on the equator of the spherical system. Specifically, at  $d = 0$  the associated spherical system is

$$\begin{aligned}\theta' &= -f_r \sin \theta \cos \theta \\ \phi' &= f_i \cos \theta.\end{aligned}\tag{8.11}$$

When  $d$  is perturbed from zero we expect the invariant set to persist. In this case it consists of four equilibria (two each of the SS and AR solution types) and the heteroclinic orbits linking them.

So far we have indicated just one possibility for the dynamics associated with regions of the coefficient space where all branches are supercritical and unstable. This applied to the regions labeled 6 in cross-section  $(I)$  of Figure 8. We close this section by suggesting possibilities for the dynamics associated with the other regions of the coefficient space where all branches are supercritical and unstable. The simplest scenario applies if the SCR subspace is attracting in which case the results of Swift on the  $D_4$ -equivariant Hopf bifurcation apply. In particular, Swift showed that there are open regions of the coefficient space where there exists a quasiperiodic solution [11]. Specifically, the Poincaré–Bendixson theorem was used to show that there exists a pair of limit cycles in the associated spherical system (one on the group orbit of the other). He further conjectured that there is at most one such pair of limit cycles. With  $d_r > 0$ , we expect the limit cycles to exist and be stable in the SCR subspace in the following regions of the coefficient space:

$$(II) : \quad \operatorname{Re}(|d|^2 - f\bar{d}) < 0, \quad (\text{SS, AR, SR}) = (\text{Sa, QP, So}) \tag{8.12a}$$

$$(III) : \quad \operatorname{Re}(|d|^2 - f\bar{d}) < 0, \quad (\text{SS, AR, SR}) = (\text{Sa, So, QP}) \tag{8.12b}$$

$$(VI) : \quad |f|^2 - |d|^2 < 0, \quad (\text{SS, AR, SR}) = (\text{QP, So, Sa}). \tag{8.12c}$$

(The parameter symmetry may be used to infer the results for  $d_r < 0$ .) Here the roman numerals refer to the cross-section of Figure 8 and the notation  $(\text{SS, AR, SR}) = (\text{Sa, QP, So})$  specifies that in the associated spherical system the SS solution is a saddle, the SR solution is a source, and that the quasiperiodic solution appears as a limit cycle surrounding the source AR (*cf.* Figure 11c). We now focus on the regions of cross-sections  $(II)$ ,  $(III)$  and  $(VI)$  where all branches are supercritical and unstable (regions 6 and 35); a superscript



$a$  indicates that the SCR subspace is attracting on neighborhoods of each of the AR, SS and SR solutions (*i.e.*, all eigenvalues associated with eigendirections transverse to the SCR subspace and its group orbit have negative real part). In these cases it is plausible that the SCR *subspace* is attracting and that the quasiperiodic solution is asymptotically stable. In the regions labeled with a superscript  $b$  the SCR subspace is not attracting and the dynamics associated with these regions may be quite complicated. In particular, it may be possible to get heteroclinic cycles involving not only the periodic solutions created at the Hopf bifurcation, but also the quasiperiodic solution in the SCR subspace. For example, with  $a_r < 0$  in region  $6^b$  of cross-section (III) there are heteroclinic connections  $SR \rightarrow TR \rightarrow TS \rightarrow SS$ , but as already noted there is no connection  $SS \rightarrow SR$ . However, in this region of coefficient space, in the associated spherical system the SS solution is a saddle and the SR solution is surrounded by a (presumably isolated) stable limit cycle. Hence one might expect a heteroclinic cycle involving the quasiperiodic solution rather than the SR solution (*i.e.*,  $QP \rightarrow TR \rightarrow TS \rightarrow SS \rightarrow QP$ ).

## 9. Conclusion.

In this paper we have employed the techniques of equivariant bifurcation theory to analyze the generic Hopf bifurcation problem with  $D_4 \times_s T^2$  symmetry. The problem is motivated by the Hopf bifurcation of a trivial (homogeneous) equilibrium in a continuous translation-invariant system. In the presence of periodic boundary conditions the problem has the spatial symmetry  $G \times_s T^2$ , where  $G$  is the symmetry of the unit cell of the resulting lattice and the two-torus  $T^2$  describes translation in two independent directions. We have focused on the case  $G = D_4$ , *i.e.*, on patterns periodic on a square lattice. For this problem the center manifold is eight-dimensional, and the appropriate action of  $D_4 \times_s T^2$  is described by (2.4). The resulting  $D_4 \times_s T^2$ -equivariant amplitude equations were assumed to be in Poincaré-Birkhoff normal form and hence to commute with an additional  $S^1$  phase shift symmetry. All isotropy subgroups were determined, thereby classifying the possible solutions by their symmetries. The  $S^1$  symmetry of the normal form introduced the possibility of patterns characterized by spatio-temporal symmetries, as well as by purely spatial symmetries. The equivariant Hopf theorem [6] was then invoked to determine five nontrivial periodic solutions that generically bifurcate from the trivial solution at the Hopf bifurcation. These solutions, depicted in Figure 4, correspond to two travelling wave patterns and three standing wave patterns.

The general form of a smooth  $D_4 \times_s T^2 \times S^1$ -equivariant vector field on  $C^4$  was derived in section 4. All small amplitude periodic solutions of the third order truncation of this vector field were determined on a neighborhood of the Hopf bifurcation. In addition to the five solutions guaranteed by the equivariant Hopf theorem, a sixth solution was found in an open region of the coefficient space. This solution resides in a four-dimensional invariant

subspace and has submaximal symmetry. It is unstable to perturbations in the fixed point subspace of its isotropy subgroup.

In section 6 the linear orbital stabilities of the maximally symmetric solutions were determined with respect to perturbations periodic on the square lattice. Full advantage was taken of the symmetries of the solutions in calculating the eigenvalues of the associated  $8 \times 8$  Jacobian matrices. This was done explicitly by forming the isotypic decomposition of  $\mathbb{C}^4$  for each isotropy subgroup. General expressions were given for the eigenvalues of the Jacobian matrix evaluated on the solution branches with maximal symmetry. In the generic bifurcation problem, solution stabilities depend on six of the eight (real) cubic coefficients. All possible bifurcation diagrams were determined within this coefficient space. This paper thus provides an exhausting classification of the generic possibilities for Hopf bifurcation on a square lattice. In section 8 attention was drawn to the interesting dynamics expected to take place when all the primary branches bifurcate supercritically but none are stable. In particular, we presented evidence for the existence of a primary bifurcation to a structurally stable heteroclinic cycle in a particular region of the coefficient space. We also indicated in which regions of the coefficient space there exists a quasiperiodic solution. The existence of heteroclinic cycles involving quasiperiodic solutions is currently under investigation. We hope in the future to report on these more complicated cycles, as well as on the asymptotic stability of the heteroclinic cycles and quasiperiodic solutions.

The theory described in the present paper can be applied to any large aspect ratio system undergoing a Hopf bifurcation from a trivial solution. As noted in section 3, aspects of the theory also apply to small aspect ratio systems defined on square domains when Neumann boundary conditions are used. Convection in a binary fluid mixture provides us with a typical example where our bifurcation analysis applies. If the separation ratio characterizing such a mixture is sufficiently negative the trivial state, corresponding to pure conduction, loses stability at a Hopf bifurcation as the Rayleigh number is raised. With periodic boundary conditions in one direction it is well known that travelling rolls (TR) are generally selected over standing rolls (SR). These conclusions are based on explicit computation of the coefficients  $a_r$  and  $b_r$  in (5.3) with both idealized [18] and experimentally realistic [19] boundary conditions, and accords with both numerical and experimental observations. The remaining four coefficients necessary to make predictions for the selection of two-dimensional patterns have not, however, been computed. The present paper should serve as a guide to efficiently performing such calculations. In an interesting paper Pismen [20] observed that if in addition to no-mass-flux boundary conditions at the top and bottom nearly thermally insulating boundary conditions are employed the wavelength of the initial instability will be long. In this case a simple equation for the two-dimensional planform of the pattern can be derived. This equation takes the form (5.3) and the pattern selection problem can be solved. In one dimension Pismen shows that TR are selected; in two dimensions the solution AR is stable.

These results are a special case of our more general analysis. Other applications along these lines can be easily imagined. In particular we mention here the application of the analysis of the Hopf bifurcation with  $D_6 \times_s T^2$  symmetry [12] to the convective instability of two superposed fluids heated from below [21].

**Acknowledgements.** We have benefitted from discussions with P. Chossat, M. Golubitsky, I. Melbourne, J.W. Swift and D.H. Wagner. This work was supported in part by NSF/DARPA under grant DMS-8814702.

## REFERENCES

- [1] M. Golubitsky, J.W. Swift and E. Knobloch, *Symmetries and pattern selection in Rayleigh-Bénard convection*, Physica **10D** (1984), pp. 249–276.
- [2] M. Silber and E. Knobloch, *Pattern selection in ferrofluids*, Physica **30D** (1988), pp. 83–98.
- [3] E. Knobloch, *Pattern selection in binary-fluid convection at positive separation ratios*, Phys. Rev. A **40** (1989), pp. 1549–1559.
- [4] J.W. Swift, *Bifurcation and symmetry in convection*, Ph.D. Thesis, University of California, Berkeley (1984).
- [5] L.M. Pismen, *Nonlinear dynamics of dissipative wave patterns near a bifurcation point*, Dynamics and Stability of Systems **1** (1986), pp. 97–113.
- [6] M. Golubitsky and I. Stewart, *Hopf bifurcation in the presence of symmetry*, Arch. Ration. Mech. Anal. **87** (1985), pp. 107–165.
- [7] E. Knobloch, *On the degenerate Hopf bifurcation with  $O(2)$  symmetry*, Contemp. Math. vol. **56**, ed. by M. Golubitsky and J. Guckenheimer (1986), pp. 193–201.
- [8] M. Golubitsky and M. Roberts, *A classification of degenerate Hopf bifurcations with  $O(2)$  symmetry*, J. Diff. Eq. **69** (1987), pp. 216–264.
- [9] J.D. Crawford and E. Knobloch, *Classification and unfolding of degenerate Hopf bifurcations with  $O(2)$  symmetry: no distinguished parameter*, Physica **31D** (1988), pp. 1–48.
- [10] M. Golubitsky and I. Stewart, *Hopf bifurcation with dihedral group symmetry: coupled nonlinear oscillators*, Contemp. Math. vol. **56**, ed. by M. Golubitsky and J. Guckenheimer (1986), pp. 131–173.
- [11] J.W. Swift, *Hopf bifurcation with the symmetry of the square*, Nonlinearity **1** (1988), pp. 333–377.
- [12] M. Roberts, J.W. Swift and D.H. Wagner, *The Hopf bifurcation on a hexagonal lattice*, Contemp. Math. vol. **56**, ed. by M. Golubitsky and J. Guckenheimer (1986), pp. 283–318.
- [13] J.E. Marsden and M. McCracken, *The Hopf bifurcation and its applications*, Springer Series in Appl. Math. Sci. vol **19** (1976).
- [14] M. Golubitsky, I. Stewart and D.G. Schaeffer, *Singularities and groups in bifurcation theory, volume II*, Springer Series in Appl. Math. Sci. vol. **69** (1988).
- [15] J.D. Crawford, M. Golubitsky, M.G.M. Gomes, E. Knobloch and I.N. Stewart, *Boundary conditions as symmetry constraints*, Proc. Warwick Symp. on Singularity Theory and its Applications (1990).
- [16] J. Guckenheimer and P. Holmes, *Structurally stable heteroclinic cycles*, Math. Proc. Camb. Phil. Soc. **103** (1988), pp. 189–192.
- [17] I. Melbourne, P. Chossat and M. Golubitsky, *Heteroclinic cycles involving periodic solutions in mode interactions with  $O(2)$  symmetry*, Proc. Roy. Soc. Edinburgh **113A** (1989), pp. 315–345.
- [18] E. Knobloch, *Oscillatory convection in binary mixtures*, Phys. Rev. A **34** (1986), pp. 1538–1549.

- [19] W. Schöpf and W. Zimmermann, *Multicritical behaviour in binary fluid convection*, Europhys. Lett. **8** (1989), pp. 41–46.
- [20] L.M. Pismen, *Selection of long-scale oscillatory convective patterns*, Phys. Rev. A **38** (1988), pp. 2564–2572.
- [21] M. Renardy and Y. Renardy, *Bifurcating solutions at the onset of convection in the Bénard problem for two fluids*, Physica **32D** (1988), pp. 227–252.

#	Author/s	Title
595	Mark J. Friedman and Eusebius J. Doedel,	Numerical computation and continuation of invariant manifolds connecting fixed points
596	Scott J. Spector,	Linear Deformations as Global Minimizers in Nonlinear Elasticity
597	Denis Serre,	Richness and the classification of quasilinear hyperbolic systems
598	L. Preziosi and F. Rosso,	On the stability of the shearing flow between pipes
599	Avner Friedman and Wenxiong Liu,	A system of partial differential equations arising in electrophotography
600	Jonathan Bell, Avner Friedman, and Andrew A. Lacey,	On solutions to a quasilinear diffusion problem from the study of soft tissue
601	David G. Schaeffer and Michael Shearer,	Loss of hyperbolicity in yield vertex plasticity models under nonproportional loading
602	Herbert C. Kranzer and Barbara Lee Keyfitz,	A strictly hyperbolic system of conservation laws admitting singular shocks
603	S. Laederich and M. Levi,	Qualitative dynamics of planar chains
604	Milan Miklavčič,	A sharp condition for existence of an inertial manifold
605	Charles Collins, David Kinderlehrer, and Mitchell Luskin,	Numerical approximation of the solution of a variational problem with a double well potential
606	Todd Arbogast,	Two-phase incompressible flow in a porous medium with various nonhomogeneous boundary conditions
607	Peter Poláčik,	Complicated dynamics in scalar semilinear parabolic equations in higher space dimension
608	Bei Hu,	Diffusion of penetrant in a polymer: a free boundary problem
609	Mohamed Sami ElBialy,	On the smoothness of the linearization of vector fields near resonant hyperbolic rest points
610	Max Jodeit, Jr. and Peter J. Olver,	On the equation $\text{grad } f = M \text{ grad } g$
611	Shui-Nee Chow, Kening Lu, and Yun-Qiu Shen,	Normal form and linearization for quasiperiodic systems
612	Prabir Daripa,	Theory of one dimensional adaptive grid generation
613	Michael C. Mackey and John G. Milton,	Feedback, delays and the origin of blood cell dynamics
614	D.G. Aronson and S. Kamin,	Disappearance of phase in the Stefan problem: one space dimension
615	Martin Krupa,	Bifurcations of relative equilibria
616	D.D. Joseph, P. Singh, and K. Chen,	Couette flows, rollers, emulsions, tall Taylor cells, phase separation and inversion, and a chaotic bubble in Taylor-Couette flow of two immiscible liquids
617	Artemio González-López, Niky Kamran, and Peter J. Olver,	Lie algebras of differential operators in two complex variables
618	L.E. Fraenkel,	On a linear, partly hyperbolic model of viscoelastic flow past a plate
619	Stephen Schecter and Michael Shearer,	Undercompressive shocks for nonstrictly hyperbolic conservation laws
620	Xinfu Chen,	Axially symmetric jets of compressible fluid
621	J. David Logan,	Wave propagation in a qualitative model of combustion under equilibrium conditions
622	M.L. Zeeman,	Hopf bifurcations in competitive three-dimensional Lotka-Volterra Systems
623	Allan P. Fordy,	Isospectral flows: their Hamiltonian structures, Miura maps and master symmetries
624	Daniel D. Joseph, John Nelson, Michael Renardy, and Yuriko Renardy,	Two-Dimensional cusped interfaces
625	Avner Friedman and Bei Hu,	A free boundary problem arising in electrophotography
626	Hamid Bellout, Avner Friedman and Victor Isakov,	Stability for an inverse problem in potential theory
627	Barbara Lee Keyfitz,	Shocks near the sonic line: A comparison between steady and unsteady models for change of type
628	Barbara Lee Keyfitz and Gerald G. Warnecke,	The existence of viscous profiles and admissibility for transonic shocks
629	P. Szmolyan,	Transversal heteroclinic and homoclinic orbits in singular perturbation problems
630	Philip Boyland,	Rotation sets and monotone periodic orbits for annulus homeomorphisms
631	Kenneth R. Meyer,	Apollonius coordinates, the N-body problem and continuation of periodic solutions
632	Chjan C. Lim,	On the Poincaré-Whitney circuit space and other properties of an

- incidence matrix for binary trees
- 633 **K.L. Cooke and I. Györi**, Numerical approximation of the solutions of delay differential equations on an infinite interval using piecewise constant arguments
- 634 **Stanley Minkowitz and Matthew Witten**, Periodicity in cell proliferation using an asynchronous cell population
- 635 **M. Chipot and G. Dal Maso**, Relaxed shape optimization: The case of nonnegative data for the Dirichlet problem
- 636 **Jeffery M. Franke and Harlan W. Stech**, Extensions of an algorithm for the analysis of nongeneric Hopf bifurcations, with applications to delay-difference equations
- 637 **Xinfu Chen**, Generation and propagation of the interface for reaction–diffusion equations
- 638 **Philip Korman**, Dynamics of the Lotka–Volterra systems with diffusion
- 639 **Harlan W. Stech**, Generic Hopf bifurcation in a class of integro-differential equations
- 640 **Stephane Laederich**, Periodic solutions of non linear differential difference equations
- 641 **Peter J. Olver**, Canonical Forms and Integrability of BiHamiltonian Systems
- 642 **S.A. van Gils, M.P. Krupa and W.F. Langford**, Hopf bifurcation with nonsemisimple 1:1 Resonance
- 643 **R.D. James and D. Kinderlehrer**, Frustration in ferromagnetic materials
- 644 **Carlos Rocha**, Properties of the attractor of a scalar parabolic P.D.E.
- 645 **Debra Lewis**, Lagrangian block diagonalization
- 646 **Richard C. Churchill and David L. Rod**, On the determination of Ziglin monodromy groups
- 647 **Xinfu Chen and Avner Friedman**, A nonlocal diffusion equation arising in terminally attached polymer chains
- 648 **Peter Gritzmann and Victor Klee**, Inner and outer  $j$ - Radii of convex bodies in finite-dimensional normed spaces
- 649 **P. Szmolyan**, Analysis of a singularly perturbed traveling wave problem
- 650 **Stanley Reiter and Carl P. Simon**, Decentralized dynamic processes for finding equilibrium
- 651 **Fernando Reitich**, Singular solutions of a transmission problem in plane linear elasticity for wedge-shaped regions
- 652 **Russell A. Johnson**, Cantor spectrum for the quasi-periodic Schrödinger equation
- 653 **Wenxiong Liu**, Singular solutions for a convection diffusion equation with absorption
- 654 **Deborah Brandon and William J. Hrusa**, Global existence of smooth shearing motions of a nonlinear viscoelastic fluid
- 655 **James F. Reineck**, The connection matrix in Morse–Smale flows II
- 656 **Claude Baesens, John Guckenheimer, Seunghwan Kim and Robert Mackay**, Simple resonance regions of torus diffeomorphisms
- 657 **Willard Miller, Jr.**, Lecture notes in radar/sonar: Topics in Harmonic analysis with applications to radar and sonar
- 658 **Calvin H. Wilcox**, Lecture notes in radar/sonar: Sonar and Radar Echo Structure
- 659 **Richard E. Blahut**, Lecture notes in radar/sonar: Theory of remote surveillance algorithms
- 660 **D.V. Anosov**, Hilbert’s 21st problem (according to Bolibruch)
- 661 **Stephane Laederich**, Ray–Singer torsion for complex manifolds and the adiabatic limit
- 662 **Geneviève Raugel and George R. Sell**, Navier-Stokes equations in thin 3d domains: Global regularity of solutions I
- 663 **Emanuel Parzen**, Time series, statistics, and information
- 664 **Andrew Majda and Kevin Lamb**, Simplified equations for low Mach number combustion with strong heat release
- 665 **Ju. S. Il’yashenko**, Global analysis of the phase portrait for the Kuramoto–Sivashinsky equation
- 666 **James F. Reineck**, Continuation to gradient flows
- 667 **Mohamed Sami Elbialy**, Simultaneous binary collisions in the collinear  $N$ -body problem
- 668 **John A. Jacquez and Carl P. Simon**, Aids: The epidemiological significance of two different mean rates of partner-change
- 669 **Carl P. Simon and John A. Jacquez**, Reproduction numbers and the stability of equilibria of SI models for heterogeneous populations
- 670 **Matthew Stafford**, Markov partitions for expanding maps of the circle
- 671 **Ciprian Foias and Edriss S. Titi**, Determining nodes, finite difference schemes and inertial manifolds
- 672 **M.W. Smiley**, Global attractors and approximate inertial manifolds for abstract dissipative equations
- 673 **M.W. Smiley**, On the existence of smooth breathers for nonlinear wave equations
- 674 **Hitay Özbay and Janos Turi**, Robust stabilization of systems governed by singular integro-differential equations
- 675 **Mary Silber and Edgar Knobloch**, Hopf bifurcation on a square lattice
- 676 **Christophe Golé**, Ghost circles for twist maps
- 677 **Christophe Golé**, Ghost tori for monotone maps
- 678 **Christophe Golé**, Monotone maps of  $T^n \times R^n$  and their periodic orbits