

ORIGINAL ARTICLE

Horizontal transfer of the nitrogen fixation gene cluster in the cyanobacterium *Microcoleus chthonoplastes*

Henk Bolhuis, Ina Severin, Veronique Confurius-Guns, Ute I A Wollenzien and Lucas J Stal
Department of Marine Microbiology, Netherlands Institute of Ecology (NIOO-KNAW), Centre for Estuarine and Marine Ecology, Yerseke, The Netherlands

The filamentous, non-heterocystous cyanobacterium *Microcoleus chthonoplastes* is a cosmopolitan organism, known to build microbial mats in a variety of different environments. Although most of these cyanobacterial mats are known for their capacity to fix dinitrogen, *M. chthonoplastes* has not been assigned as a diazotrophic organism. None of the strains that were correctly identified as *M. chthonoplastes* has been shown to fix dinitrogen and it has repeatedly been reported that these organisms lacked the cyanobacterial *nifH*, the structural gene for dinitrogenase reductase. In this study, we show that a complete *nif*-gene cluster is present in the genome of *M. chthonoplastes* PCC 7420 and that the three structural nitrogenase genes, *nifHDK*, are present in a collection of axenic strains of *M. chthonoplastes* from distant locations. Phylogenetic analysis of *nifHDK* revealed that they cluster with the *Deltaproteobacteria* and that they are closely related to *Desulfovibrio*. The *nif* operon is flanked by typical cyanobacterial genes, suggesting that it is an integral part of the *M. chthonoplastes* genome. In this study, we provide evidence that the *nif* operon of *M. chthonoplastes* is acquired through horizontal gene transfer. Moreover, the presence of the same *nif*-cluster in *M. chthonoplastes* isolates derived from various sites around the world suggests that this horizontal gene transfer event must have occurred early in the evolution of *M. chthonoplastes*. We have been unable to express nitrogenase in cultures of *M. chthonoplastes*, but we show that these genes were expressed under natural conditions in the field.

The ISME Journal (2010) 4, 121–130; doi:10.1038/ismej.2009.99; published online 10 September 2009

Subject Category: microbial ecology and functional diversity of natural habitats

Keywords: *Cyanobacteria*; *Microcoleus*; nitrogen fixation; horizontal gene transfer; microbial mat

Introduction

Many cyanobacterial mats are formed by the cosmopolitan filamentous non-heterocystous cyanobacterium *Microcoleus chthonoplastes* (Stal, 2000). This species is easily recognizable by its occurrence in bundles of trichomes, enveloped by a common polysaccharide sheath, although this property may be lost in culture, which has sometimes led to the misidentification of cultured isolates of morphological similar opportunists (Garcia-Pichel *et al.*, 1996; Siegesmund *et al.*, 2008).

The fixation of atmospheric dinitrogen is a property of all cyanobacterial mats that have been investigated for it (Severin and Stal, 2009). The first isolate of an aerobic dinitrogen-fixing filamentous non-heterocystous cyanobacterium was originally

assigned to *M. chthonoplastes* (Pearson *et al.*, 1979; Malin and Pearson, 1988), but was later re-assigned to *Symploca* sp. (Janson *et al.*, 1998). Similarly, *M. chthonoplastes* ‘strain 11’, isolated from a diazotrophic microbial mat from the Wadden Sea (Southern North Sea), was reported to be capable of anaerobic dinitrogen fixation (Stal and Krumbein, 1985), but was also re-assigned in this case to the genus *Geitlerinema* (Garcia-Pichel *et al.*, 1996; Siegesmund *et al.*, 2008), a genus to which also the famous anoxygenic phototrophic Solar Lake strain *Oscillatoria limnetica* belongs that can fix dinitrogen anaerobically. Other reports of dinitrogen-fixing *M. chthonoplastes* and a *Microcoleus* sp. were by Dubinin *et al.* (1992) and Sroga (1997), respectively, but it is unclear whether their assignments were correct.

Pure culture studies with different true *M. chthonoplastes* strains did not reveal nitrogenase activity or cyanobacterial *nifH* gene expression. Rippka *et al.* (1979) were unable to detect nitrogenase activity in the type strain of *M. chthonoplastes* PCC7420, not even under strictly anaerobic conditions. Villbrandt and Stal (unpublished results) confirmed this result and also showed that none of

Correspondence: Henk Bolhuis, Department of Marine Microbiology, Netherlands Institute of Ecology (NIOO-KNAW), Centre for Estuarine and Marine Ecology, Korrिंगaweg 7, 4401 NT Yerseke, The Netherlands.

E-mail: H.bolhuis@nioo.knaw.nl

Received 16 June 2009; revised 23 July 2009; accepted 28 July 2009; published online 10 September 2009

the 'true' *M. chthonoplastes* of the collection of Garcia-Pichel *et al.* (1996) possessed nitrogenase activity, even when assayed under anaerobic conditions (according to the method described in Rippka and Waterbury (1977)). Moreover, Steppe *et al.* (1996) were unable to detect cyanobacterial *nifH*, the gene coding for dinitrogenase reductase, in four strains of *M. chthonoplastes*, including the type strain PCC7420. Instead, they amplified a non-cyanobacterial *nifH* from these cultures, which they attributed to contaminants. The amplified *nifH* genes were closely related to *Gammaproteobacteria*, *Clostridium pasteurianum* and *Desulfovibrio gigas*, sequences that were also found in microbial mats dominated by *M. chthonoplastes*. Studies in *M. chthonoplastes*-dominated microbial mats revealed similar results, amplifying *Gamma- or Delta-proteobacterial nifH* sequences rather than cyanobacterial *nifH* genes (Zehr *et al.*, 1995; Olson *et al.*, 1999; Omoregie *et al.*, 2004). In contrast, mats containing another filamentous non-heterocystous cyanobacterium, *Lyngbya* sp., did reveal cyanobacterial *nifH* sequences, showing that it was not impossible to retrieve cyanobacterial *nifH* sequences from environmental samples. These results were generally interpreted as that *M. chthonoplastes* is not capable of fixing dinitrogen because it lacked the required nitrogen fixation genes. Moreover, finding bacterial *nifH* genes in *M. chthonoplastes*-dominated mats rather than cyanobacterial genes was dedicated to the presence of diazotrophic heterotrophic bacteria living in a consortium with *M. chthonoplastes*. It was also concluded that *Cyanobacteria* might be less important for dinitrogen fixation in these mats.

In another marine *M. chthonoplastes* mat it was shown, by combining metabolic inhibitor studies with the analyses of mRNA by reverse transcriptase PCR (RT-PCR), that sulfate-reducing bacteria were potentially important dinitrogen fixers and that most of the *nifH* genes expressed clustered with sulfate-reducing bacteria of the *Deltaproteobacteria* and other anaerobic bacteria (Steppe and Paerl, 2002). It has been suggested that these dinitrogen-fixing heterotrophic bacteria occur as epiphytes on the mucilaginous sheaths of *M. chthonoplastes* with which they presumably interact, and this interaction has been conceptualized as consortial dinitrogen fixation (Steppe *et al.*, 1996; Paerl *et al.*, 2000).

In this study, we report the existence of a complete nitrogenase operon in the genome of *M. chthonoplastes* PCC7420 that has been sequenced through the Moore Foundation Marine Microbiology project (<http://www.moore.org/microgenome/>).

Materials and methods

Strains and growth conditions

The strains used in this study were obtained from the Culture Collection Yerseke (CCY) and are listed

in Table 1. All strains were grown in artificial seawater medium ASN3 (Rippka *et al.*, 1979). The dinitrogen-fixing *Symploca* strains were grown in medium lacking combined nitrogen (NO_3^-) (ASN3 $^\circ$). *Microcoleus* strains were grown at 23 °C and a photon flux density of 70 $\mu\text{mol m}^{-2} \text{s}^{-1}$ in an orbital shaker at 120 rpm and a light/dark cycle of 14 h light and 10 h dark. *Symploca* strains were grown in an orbital shaker at 120 rpm and 18 °C, with a photon flux density of 20 $\mu\text{mol m}^{-2} \text{s}^{-1}$ and a light/dark cycle of 16 h light and 8 h dark. Nitrogenase activity in the other strains was induced following the method of Rippka and Waterbury (1977). Briefly, a growing culture was transferred to medium devoid of combined nitrogen (ASN3 $^\circ$) and incubation was continued until the first sign of bleaching was observed. Bleaching is caused by the degradation of the phycobiliproteins and indicates the onset of nitrogen starvation. Nitrogen-starved cultures were subsequently concentrated by centrifugation and re-suspended in 10 ml of ASN3 $^\circ$ medium to which 10 $^{-5}$ M DCMU (3-(3,4-dichlorophenyl)-1,1-dimethyl urea) was added to prevent oxygen evolution by the inhibition of photosystem II. The suspension was flushed with dinitrogen to achieve anaerobic conditions. The 30-ml flask was sealed and 15% acetylene gas was added with a syringe as over-pressure and the suspensions were incubated in front of 2 × 18 Watt daylight fluorescent tubes with a photon flux density of 50–60 $\mu\text{mol m}^{-2} \text{s}^{-1}$. Ethylene was assayed gas chromatographically (Stal, 1988) at regular intervals, starting 3 h after establishing anaerobic conditions. After 24 h, the incubation was terminated and the cells were mixed immediately with RNAlater (Ambion Inc., Austin, TX, USA)

Table 1 List of species used in this study

Species	CCY strain ID	Place of isolation
<i>Symploca atlantica</i> PCC 8002	CCY9617	UK
<i>Symploca</i> sp.	CCY0030	Rottneest Island, Australia
<i>Geitlerinema</i> sp.	CCY9619	Mellum, Germany
<i>Microcoleus chthonoplastes</i> PCC7420 ^a	CCY9604	Woods Hole, USA
<i>Microcoleus</i> sp. ^a	CCY0002	Schiermonnikoog, The Netherlands
<i>Microcoleus</i> sp. ^a	CCY9602	Chile
<i>Microcoleus</i> sp. ^a	CCY9603	Solar Lake, Egypt
<i>Microcoleus</i> sp. ^a	CCY9605	Mellum, Germany
<i>Microcoleus</i> sp. ^a	CCY9606	St Peter-Ording, Germany
<i>Microcoleus</i> sp. ^a	CCY9707	North Carolina, USA
<i>Microcoleus</i> sp. ^a	CCY9608	Wismar, Germany
<i>Geitlerinema</i> sp.	CCY9412	Schiermonnikoog, The Netherlands
<i>Geitlerinema</i> sp.	CCY0102	Krim, Russia
<i>Microcoleus</i> sp. ^a	CCY0602	Negev Desert, Israel

^aStrains proposed to be renamed to *Coleofasculus* according to Siegesmund *et al.* (2008)

following the manufacturer's instructions. These samples were subsequently used for extraction of RNA and RT-PCR of *nifHDK*. Natural samples were taken in the summer of 2007 from a *M. chthonoplastes* containing microbial mat located at the North Sea beach of Schiermonnikoog, The Netherlands (Severin and Stal, 2008). These samples were flash frozen in liquid nitrogen and kept at -80°C until use.

Isolation of nucleic acids

DNA was isolated from cultures and natural mixed mat samples using the UltraClean Soil DNA Isolation Kit (MoBio Laboratories Inc., Carlsbad, CA, USA) according to the manufacturer's instructions for maximal yields. RNA was extracted from cultures and natural mixed mat samples using the RNeasy Mini Kit (Qiagen Inc., Valencia, CA, USA) following the manufacturer's instructions. RNA samples were treated with DNase I (Invitrogen Corp., Carlsbad, CA, USA) to remove residual DNA, and cDNA was generated using Super-Script II reverse transcriptase (Invitrogen Corp.) following the manufacturer's instructions and using random hexamer primers.

PCR amplification, fragment cloning and DNA sequencing.

Oligonucleotide primers designed and used in this study are listed in Table 2. A total of 25 μl of PCR mixture consisted of 200 μM of dNTPs (Roche Applied Science, Indianapolis, IN, USA), 200 nM of primers, 1 \times HotStarPCR buffer (Qiagen Inc.), 10% v/v of dimethyl sulfoxide (Sigma-Aldrich, Munich, Germany), 0.2 mg ml $^{-1}$ of bovine serum albumin (Fermentas, Hanover, MD, USA) and 0.6 U of HotStarTaq DNA polymerase (Qiagen Inc.). The reactions were run on a thermal cycler (Thermal Cycler 2720, Applied Biosystem, Foster City, CA, USA). For the MC-*nifHDK* primers the following program was used: 15 min at 94 $^{\circ}\text{C}$; 35 cycles of 30 s at 94 $^{\circ}\text{C}$, 30 s at 48 $^{\circ}\text{C}$, 60 s at 72 $^{\circ}\text{C}$; and a final extension step for 7 min at 72 $^{\circ}\text{C}$. For the amplification of the 16S rRNA gene, primers B27F and U1492R (Table 2) were used with the following program: 15 min at 94 $^{\circ}\text{C}$; 35 cycles of 30 s at 94 $^{\circ}\text{C}$,

30 s at 55 $^{\circ}\text{C}$, 110 s at 72 $^{\circ}\text{C}$ and a final extension step for 7 min at 72 $^{\circ}\text{C}$. PCR products were separated by electrophoresis on a 1% w/v agarose (Sigma-Aldrich) gel and stained with SYBR Gold (Invitrogen Corp.). Amplicon size was estimated by comparison with a MassRuler DNA Ladder (Fermentas). Amplicons were cloned using the TOPO-TA cloning Kit (Invitrogen Corp.) following the manufacturer's instructions. White colonies were selected and suspended in 10 μl of sterile MilliQ water, boiled for 10 min and used as template in a PCR with the vector primers T7 and T3 to amplify the inserted gene fragments. The PCR mixture (25 μl) consisted of 200 μM of dNTPs (Roche), 200 nM of primers, standard Taq buffer (1 \times , New England BioLabs Inc., Ipswich, MA, USA) and 0.6 U of Taq DNA polymerase (New England Biolabs Inc.). The PCR reaction was run at 95 $^{\circ}\text{C}$ for 2 min; 35 cycles 30 s at 94 $^{\circ}\text{C}$, 30 s at 55 $^{\circ}\text{C}$, 60 s at 72 $^{\circ}\text{C}$ and finally 10 min at 72 $^{\circ}\text{C}$. Amplicons containing insert DNA of the appropriate size were purified using Sephadex G-50 Superfine (Sigma-Aldrich) and DNA concentrations were determined spectrophotometrically. Amplicons were sequenced using the BigDye Terminator chemistry (Big Dye Terminator v3.1 Cycle Sequencing Kit, Applied Biosystem) according to the manufacturer's instructions. The sequence products were analyzed with a 3130 Genetic Analyzer (Applied Biosystem).

Gene expression analysis by qRT-PCR

On the basis of the *nifH* sequence information of *M. chthonoplastes* CCY0002, a reverse primer (MC-*nifH*-QRTr; Table 2) was designed manually that could be combined with the MC-*nifH*-f forward primer in a qRT-PCR (quantitative RT-PCR) reaction. The specificity of this primer pair was checked *in silico* as well as by PCR amplification, cloning and sequencing 48 amplification products from environmental samples. All sequenced amplicons were 100% identical to part of the *nifH* gene of *M. chthonoplastes* CCY0002, confirming the specificity of the chosen primer set for qRT-PCR analysis. Quantitative RT-PCR was run on a Corbett Rotor-Gene 6000 (Corbett Life Science, Sydney, NSW,

Table 2 List of oligonucleotides used in this study

Name	Sequence 5'-3'	Gene	Position within gene
MC- <i>nifH</i> -f	TTTACGGTAAAGGTGGAATCG	<i>nifH</i>	17–37
MC- <i>nifH</i> -r	TGCAAATCAAACCGCTAAACG	<i>nifH</i>	550–529
MC- <i>nifD</i> -f	CAAGCCGTTTCAGGAAGGCTA	<i>nifD</i>	424–443
MC- <i>nifD</i> -r	GAGCGGTGACACATGACCAA	<i>nifD</i>	873–854
MC- <i>nifK</i> -f	GTCAGCCTATTGGTGAATG	<i>nifK</i>	68–87
MC- <i>nifK</i> -r	CAGAACACGAGACGGAACCT	<i>nifK</i>	811–792
MC- <i>nifH</i> -QRTr	ATCTAACACACTCTTTTGGTGC	<i>nifH</i>	153–174
B27F	AGAGTTTGATCMTGGCTCAG	Bact—16S	8–27
U1492R	GTTTACCTTGTTACGACTT	Univ—16S	1510–1492

MC oligonucleotides are identical to their matching sequences in the targeted genes of *Microcoleus chthonoplastes* PCC7420.

Australia). *M. chthonoplastes* CCY0002 was isolated from the microbial mats of Schiermonnikoog from which the environmental DNA was obtained. Environmental DNA as well as reverse-transcribed cDNA reverse transcribed from environmental RNA was used as a template. Cycling conditions were the following: 15 min at 95 °C followed by 40 cycles of 15 s at 95 °C, 20 s at 55 °C and 20 s at 72 °C. The reaction was directly followed by a melting curve analysis (72–95 °C). Environmental samples were run in triplicate and each run included two independent dilution series of linearized plasmid containing the target *nifH* insert as standard curves, as well as non-template and non-target controls. The 15 µl qPCR reactions consisted of 7.5 µl of a commercially available reaction mixture (Absolute QPCR SYBR Mix, Thermo Fisher Scientific, Waltham, MA, USA) containing all the components necessary to perform quantitative PCR, with the exception of template and primers, 333 nM of the forward and reverse primer, 5.5 µl of MilliQ water and 1 µl of template. The standard curves obtained from the linearized *nifH* containing plasmids allowed us to determine the number of gene copies and the number of transcripts per ng of extracted nucleic acid, and expression levels were calculated as number of transcripts per number of *nifH* copies. The number of *nifH* copies in the standard curve ranged from 2.2×10^4 to 2.2×10^7 per reaction. Comparison of the *ct* values of the transcripts and *nifH* gene copies with that of the standard curve by least square linear regression analyses allowed us to quantify the original amount of target RNA or DNA molecules in the sample. Efficiencies for all reactions were $85 \pm 2\%$. *NifH* expression levels were compared with a 24 h cycle of nitrogenase activity.

Nitrogenase activity

Nitrogenase activity was measured in pure cultures and microbial mat samples using the acetylene reduction assay and normalized per milligram of chlorophyll-a as previously described (Severin and Stal, 2008). Light response curves of nitrogenase activity and daily dinitrogen fixation were calculated from ethylene production rates in the microbial mat sample as previously described (Severin and Stal, 2008).

DNA sequence and phylogenetic analysis

The Molecular Evolutionary Genetics Analysis (MEGA 4.0) software package (Tamura *et al.*, 2007) was used to analyze and correct ABI trace files and to conduct phylogenetic analysis. DNA and protein sequences were aligned using the ClustalW module of MEGA 4.0 using the default settings and phylogenetic trees were constructed by applying the neighbor-joining method with a 1000 replicates bootstrap analysis. DNA sequences described in this study were deposited in GenBank under accession

numbers GQ397255–GQ397274 for the *nifH*, *nifD* and *nifK* gene sequences and GQ402014–GQ402026 for the 16S rRNA gene sequences.

Results

Genetic analysis

In June 2007, the partial sequenced and automatically annotated genome of *M. chthonoplastes* PCC 7420 became publicly available at <https://moore.jcvi.org/research/>. On analysis of contig #11036-59003591 we discovered a complete *nif*-gene cluster. The *nif*-gene cluster (Figure 1a) contains the genes encoding the structural proteins (*nifHDK*) and genes for biosynthesis of the iron–molybdenum cofactor (FeMo-co; *nifB'SU* and *nifENB*). In addition, the *nif*-cluster contains two genes proposed to be involved in nitrogen regulation, *P-II* and *P-II'*, (Martin and Reinhold-Hurek, 2002) and a gene encoding a ferredoxin protein (*fd*) mediating electron transfer (Rubio and Ludden, 2008). Two copies of *nifB* were found, of which the first seems to be a partial and possibly non-functional gene (*nifB'*). Further downstream of the *nif*-cluster, genes were found that encode binding- and transport proteins involved in the uptake of the molybdate cofactor (*mopI*, *modDABC*) (Figure 1b).




Figure 1 Comparative analysis of *nif*-cluster organization in different organisms. (a) Cyanobacterial genes are presented as white arrows and *Deltaproteobacterial/Chlorobia* type *nif*-genes are presented as black arrows. AV, *A. variabilis* ATCC29413; AV', *A. variabilis* ATCC29413 (anaerobic *nif*-cluster including insertion element plus integrase); CY, *Cyanothece* PCC7426; CT, *Chlorobium tepidum*; SF, *S. fumaroxidans* MPOB; DV, *D. vulgaris* DP4; MC, *M. chthonoplastes* PCC 7420. (b) G + C content of part of contig #1103659003591 of *M. chthonoplastes* PCC7420; revealing the depressions in G + C content (vertical arrows) flanking the predicted horizontally acquired *nif*-gene cluster (black arrows).

Phylogenetic analysis of the *nif*-genes revealed that only the initial three genes (*nifB'US*) are related to cyanobacterial homologs (Figure 2). Phylogenetic analysis of *nifHDK*, *P-II*, *P-II'*, *fd* and *nifENB* (hereafter called the *nifHDKENB* cluster), revealed clustering with *nif*-genes from the family of *Chlorobiaceae* and the *Deltaproteobacteria* families of *Syntrophobacteraceae* and especially with the *Desulfovibrionaceae* (Figure 2). The *M. chthonoplastes nifH* gene contains a six nucleotide deletion resulting in the absence of two amino acids after position 64 in the amino acid sequence, a feature not shared by other cyanobacterial *nifH* genes but common in deltaproteobacterial *nifH*. Despite clustering in the *Deltaproteobacteria* group, *nifH* has highest identity to Ava_4046, one of the five *nifH* genes of *Anabaena variabilis* ATCC 29413. This *nifH* is the only *nif*-gene in *A. variabilis* that clusters with the *Deltaproteobacteria/Chlorobia* group. The other four *nifH* genes of *A. variabilis* cluster with cyanobacterial homologs. To exclude the possibility that this non-typical cyanobacterial gene clustering is caused by a contamination in the original DNA

sample used for the genome sequencing, we studied genes directly flanking the *nif*-cluster and found that they are typical cyanobacterial. Moreover, we PCR-amplified DNA fragments of the *nifH*, *nifD* and *nifK* genes from *M. chthonoplastes* PCC 7420, that is kept axenic in our CCY culture collection, and from eight other axenic strains of *M. chthonoplastes* originating from different regions around the world (Table 1), using specific primers designed on the basis of the sequences of the *M. chthonoplastes* genome. The resulting DNA fragments were of the expected length and their sequence revealed high similarity to the genome sequences of *M. chthonoplastes* PCC7420. Neighbor-joining trees based on the nucleotide sequences of the three *nif*-genes from the CCY strains showed good congruency with their respective 16S rRNA genes (Figure 3). At the amino acid sequence level, the identity was 98%, 91% and 91% for NifH, NifD and NifK, respectively. Moreover, most mutations consist of neutral substitutions not causing an amino acid change. Using these *M. chthonoplastes*-specific *nifHDK* primers, we failed to obtain products from *Symploca* or *Geitlerinema*




Figure 2 Phylogenetic analysis of NifU (left) and NifH (right) of *M. chthonoplastes* PCC7420, revealing their cyanobacterial and non-cyanobacterial descent respectively. The neighbor-joining tree was constructed using bootstrap analysis with 1000 iterations (bootstrap values are shown at nodes). Shading indicates the phylogenetic Classes wherein NifH and NifU of *M. chthonoplastes* (bold and underlined) clusters.




Figure 3 Phylogenetic analysis of 16S, *nifH*, *nifD* and *nifK* of CCY strains reveals congruency in tree topology. Neighbor-joining trees were generated using maximum likelihood algorithm with 1000 bootstrap iterations (bootstrap values are shown at nodes). *nifH* copyDNA (cDNA) obtained by reverse transcriptase PCR (RT-PCR) from a natural microbial mat is included in the *nifH* tree. CCY, Culture Collection Yerseke.

strains. As we can now exclude contamination of the genome sequence, horizontal gene transfer of the *nifHDKENB* cluster from a *Deltaproteobacteria/Chlorobia* donor is the most likely explanation of this observation.

Evidence for horizontal gene transfer

Analysis of average G+C content revealed two G+C poor stretches (29.5% and 34.3% G+C) exactly flanking the predicted horizontally acquired *nif*-cluster (Figure 1b). The average G+C content of the *nifHDKENB* cluster is in the same range (47.1% G+C) as that of the whole contig (45.1% G+C). Codon usage frequencies for each gene in the contig was calculated and compared by cluster analysis using a correlation matrix in the statistical software package Past (Hammer *et al.*, 2001). The resulting dendrogram (Figure 4) shows that based on their codon usage, the *nifHDKENB* genes form a distinct cluster, suggesting that their codon usage is more similar to each other than to that of the other genes in the same contig. Putative promoter regions are not well resolved for *M. chthonoplastes* and no obvious deviations were found from the promoter regions of the typical cyanobacterial genes in the contig (data not shown).

Expression of nitrogenase

A total of eight strains of *M. chthonoplastes* (Table 1) were cultivated and subsequently starved for




Figure 4 Cluster analysis dendrogram of codon usage frequencies of the genes within contig #1103659003591 of *M. chthonoplastes* PCC7420. Genes form clusters when codon usage frequencies are similar. The genes of the horizontally transferred *nif*-gene cluster are in bold. The other open reading frames in the contig are arbitrarily numbered.

nitrogen. In so-called anaerobically dinitrogen-fixing *Cyanobacteria*, nitrogenase is normally induced when nitrogen-deprived cultures are incubated under anaerobic conditions while oxygenic photosynthesis is blocked through inhibition of PS-II by DCMU (Lavergne, 1982). However, none of the eight *M. chthonoplastes* strains showed any nitrogenase activity after 24 h of incubation. In addition, gene expression studied by RT-PCR analysis revealed no synthesis of *nifH*, *nifK* or *nifD* specific mRNA in any of the strains.

In contrast to the pure cultures, gene expression was observed in environmental samples of the *M. chthonoplastes*-dominated microbial mat from which strain CCY0002 was originally isolated. In total, 12 samples were taken from a 24 h sampling campaign from which RNA was extracted. RT-PCR using the *M. chthonoplastes*-specific *nifH* primers revealed positive bands for each sample. Bands were of the expected size and sequence analysis on two clones revealed 100% identity with the *nifH* gene from strain CCY0002 (Figure 3, cDNA 26-5-19 and cDNA 26-5-4). Quantitative RT-PCR revealed that the number of *nifH* transcripts per copy varied between the time points (Figure 5; vertical bars). In general, higher ratios were observed at daytime and ranged between virtually zero to 0.06 *nifH* transcripts per copy. The pattern was in agreement with the daily variations of nitrogenase activity. Peaks in nitrogenase activity were observed throughout the day but were higher during the day than at night reaching 0.23 and 0.13 $\mu\text{mol C}_2\text{H}_4$ per mg chlorophyll per hour, respectively.

Discussion

Analysis of *nifH* sequences is widely used as measure for the potential of a system to fix dinitrogen and for the diversity of diazotrophic




Figure 5 Natural 24 h cycle of nitrogenase activity (filled circles) and *nifH* gene expression (columns) measured in microbial mat samples. The standard deviation of three independent experiments is indicated by error bars. The horizontal bar at the top of the graph depicts the natural light (white) and dark (black) period of the 24 h cycle.

organisms in that ecosystem. The assumption is that the phylogeny of a gene can predict the phylogeny of its host. This would be true if all genes are only transferred to its successor by linear descent. However, there are many cases known in which DNA is acquired from other species through horizontal gene transfer and subsequently becomes fixed in the genome of a host organism through adaptive evolution. The acquisition of foreign DNA may occur through specific DNA uptake systems, the active uptake of free extracellular DNA, through conjugation with other microorganisms followed by active transfer of DNA or after transduction through phages (Koonin *et al.*, 2001). Horizontal gene transfer of DNA or RNA may occur between unrelated organisms and has been shown in several members of every domain of life.

For the study of the diversity of *nifH*, often the degenerated primers are used that were developed by Zehr and McReynolds (1989). These primers were initially designed to target *nifH* genes from marine dinitrogen-fixing organisms, particularly *Cyanobacteria*, but in several studies using these degenerated primers many non-cyanobacterial *nifH* genes are targeted. Using these primers, *M. chthonoplastes*-dominated microbial mats seem to be low in cyanobacterial specific *nifH* genes. As described in the introduction, this observation, in combination with the failure to detect nitrogenase activity in pure cultures of *M. chthonoplastes*, led to the general assumption that this organism lacked the genes for dinitrogen fixation and, hence, lacked the ability to fix dinitrogen (Zehr *et al.*, 1995; Steppe and Paerl, 2002; de Wit *et al.*, 2005).

Our results, however, show that all strains of *M. chthonoplastes* tested by us contained *nifH*, *nifD* and *nifK*, and the genome sequence of *M. chthonoplastes* PCC7420 shows that a complete *nif*-gene cluster is present (Rubio and Ludden, 2008). As *M. chthonoplastes* forms large bundles of filaments enclosed in a thick extracellular polysaccharide sheath, it was difficult to grow this organism axenically. Other prokaryotes have been found tightly associated with or even within the polysaccharide sheath, including members of the family of *Desulfovibrionaceae* (D'Amelio *et al.*, 1987). A putative contamination of the DNA used for genome sequencing can therefore not be excluded. However, as the same sequence was found in eight different strains of *M. chthonoplastes* isolated from various regions around the world and were brought into pure culture by different laboratories, the chance of catching the same contamination in all strains is highly unlikely. Therefore, the most likely explanation is that *M. chthonoplastes* acquired the *nif*-gene cluster through horizontal gene transfer. This is substantiated by the following data. Phylogenetic analysis shows that the genes of the *nifHDKENB* cluster group with *Deltaproteobacteria* rather than with *Cyanobacteria* (Figure 2). The *nifHDKENB* cluster is flanked by two steep depressions in

average G + C content, which is often found to be involved in DNA integration (Kleckner, 1990). Possibly, here an A + T rich sequence functioned as site of integration for the horizontally acquired DNA fragment. In fact, given the presence of the typical cyanobacterial *nifB'SU* genes directly upstream of the acquired *nifHDKENB* genes, it is likely that a putative original *nif*-gene cluster may have been replaced by the new *nif*-cluster. Such a replacement could be mediated by insertion sequences. However, except for the A + T rich sequence, no remnants of an IS (insertion sequence) element transfer were detected. The fact that the insertion itself does not deviate in G + C content suggests that either the donor organism had a similar G + C content or that the G + C content of DNA insert adapted to that of the host organism to ensure smooth replication and transcription. In addition, the codon usage can be a marker of putative horizontal gene transfer events. Overall the average codon usage is species or even strain specific, and is regarded as an adaptation to its pool of tRNA molecules (Ikemura, 1985). The preference for certain codons over others encoding the same amino acid in a protein depends on the relative expression profiles of the different tRNAs giving rise to optimal protein synthesis machinery. In the case of the *Microcoleus nifHDKENB* cluster, the codon usage of the genes in the suspected DNA insert differs from that of the flanking genes in the contig (Figure 4), which is in support of acquisition of these genes through horizontal gene transfer.

The filamentous cyanobacterium *A. variabilis* contains five NifH homologs, one of which (Ava_4046) is at the amino acid level 72% identical to NifH from *M. chthonoplastes* and also clusters in the *Deltaproteobacteria/Chlorobia* group. However, this particular *nifH* is not part of a *nif*-gene cluster in *A. variabilis*. This organism possesses three *nif*-clusters, one of which is interrupted by an insertion sequence including an integrase (Brusca *et al.*, 1989). Under anaerobic conditions, this insertion sequence is excised, leaving an intact *nif* operon that functions in vegetative cells (Thiel *et al.*, 1995). The third *nif*-cluster in *A. variabilis* encodes an alternative, vanadium-dependent nitrogenase that functions under aerobic conditions in the heterocyst when molybdate is unavailable. Otherwise, the regular molybdate nitrogenase is expressed in heterocysts. Possibly, *nifH* (Ava_4046) of *A. variabilis* is also obtained through horizontal gene transfer but its function in this organism is not known. We did not encounter any other putative *nif*-genes in the genome of *M. chthonoplastes*.

One of the reasons why *M. chthonoplastes* thus far has been tagged as a non-diazotroph is that it does not grow without combined nitrogen and it lacks nitrogenase activity in pure isolates in the lab, even under fully anaerobic conditions. In this assay, *Cyanobacteria* that possess the genetic capacity of synthesizing nitrogenase but are unable to provide anaerobic conditions will express nitrogenase

(Rippka and Waterbury, 1977). Although a highly identical nitrogenase cluster was found in all *M. chthonoplastes* strains in our culture collection, none showed nitrogenase activity or expressed any of the three structural *nif*-genes. In contrast, we showed the expression of *nifH*, *nifD* and *nifK* in natural samples taken from a *M. chthonoplastes*-dominated marine microbial mat using primers that were specifically designed for the *M. chthonoplastes* strain that was isolated from that mat. It is possible that after maintaining the cultures of *M. chthonoplastes* for many years in the lab, growing on combined nitrogen, it might have lost the capacity of expressing nitrogenase. However, this is not the case with many other species of *Cyanobacteria* that express nitrogenase only under anaerobic conditions. Another possibility is that *M. chthonoplastes* needs different conditions to express nitrogenase. For instance, the inhibitor DCMU may have other negative side effects in this organism that prevents it from expressing nitrogenase. Sulfide or far-red light are other means for inhibiting photosystem II while allowing photosystem I activity (Jørgensen *et al.*, 1986). It is also possible that regulation of nitrogenase activity in *M. chthonoplastes* is completely different from that in other *Cyanobacteria* due to the presence of the PII genes rather than *nifX* and *nifW* typically found in *Cyanobacteria*. Alternatively, nitrogenase expression might even depend on external signals from closely associated sulfate-reducing bacteria to be activated.

Quantitative RT-PCR revealed that *M. chthonoplastes*-specific *nifH* expression in the microbial mat was highest during day and declined during night. Nitrogenase activity, as measured by the acetylene reduction test, reveals a similar pattern suggesting that the expressed genes indeed result in the translation into active nitrogenase proteins. Many non-heterocystous diazotrophic *Cyanobacteria* separate dinitrogen fixation temporally from oxygenic photosynthesis and confine the former to the night. This was not obvious in this *M. chthonoplastes* mat, which showed activity both during the day and night. Possibly, dinitrogen fixation occurs only at greater depths in the mat in which the sulfide concentrations are higher and only far-red light penetrates (Jørgensen *et al.*, 1987).

We have shown here that *M. chthonoplastes* contains a functional *nif*-gene cluster, which is not typical cyanobacterial and which is most likely acquired through horizontal gene transfer from a member of the *Deltaproteobacteria*. Therefore, several previous studies may have misinterpreted the observed microbial diversity, and underestimated cyanobacterial dinitrogen fixation and overestimated the role of other bacteria in dinitrogen fixation in cyanobacterial mats (Wawer *et al.*, 1997).

Members of the family of *Desulfovibrionaceae*, such as *D. vulgaris* or *D. gigas*, are the most likely candidate donor organisms for the *nif*-genes found in *M. chthonoplastes*. First, the gene and protein

sequences of the *M. chthonoplastes nif*-cluster showed highest identity with that of the *Desulfovibrionaceae* and second, these sulfate-reducing bacteria are frequently found in *M. chthonoplastes*-dominated microbial mats (Zehr *et al.*, 1995; Sigalevich *et al.*, 2000). Moreover, members of the *Desulfovibrionaceae* are known to be capable of transducing DNA fragments through phages (Rapp and Wall, 1987). Finding the highly conserved *nif*-genes amongst *M. chthonoplastes* strains isolated from various locations around the world suggests that these genes were obtained early in the evolution of this species. 16S rRNA analysis of the strains used in this study revealed a clade consisting of species related to *Geitlerinema* sp., a clade with species related to *Symploca* sp. and one clade with species closely related to the *M. chthonoplastes* type strain (Figure 3). Recently, Siegesmund *et al.* (2008) studied 16S rRNA genes and ribosomal ITS (internal transcribed spacer) sequences of a large number of *Cyanobacteria* originally assigned to *Microcoleus* and related genera. They found that *Microcoleus* fell into two clades, one with taxa belonging to the *Oscillatoriaceae* and the other to the *Phormidiaceae*. As the terrestrial type strain *M. vaginatus* for this genus belongs to the *Oscillatoriaceae*, a new genus *Coleofasciculus* was proposed for the strains belonging to the *Phormidiaceae* (Siegesmund *et al.*, 2008). According to the proposed new nomenclature, the strains used in this study that are closely related to *M. chthonoplastes* belong to this new genus of *Coleofasciculus* (see Table 1).

Congruency between the 16S rRNA gene tree with the *nifH*, *nifD* and *nifK* trees (Figure 3) of the CCY strains are in agreement with an early horizontal acquisition of the *nif*-gene cluster followed by strictly vertical inheritance, and it is therefore tempting to speculate that this horizontal gene transfer event may have formed the basis for the speciation of *M. chthonoplastes*.

Acknowledgements

We thank Wolfgang Hess and Jan Mitschke of the University of Freiburg, Germany for their help in promoter search. Publication 4589 Netherlands Institute of Ecology (NIOO-KNAW).

References

- Brusca JS, Hale MA, Carrasco CD, Golden JW. (1989). Excision of an 11-kilobase-pair DNA element from within the *nifD* gene in *Anabaena variabilis* heterocysts. *J Bacteriol* **171**: 4138–4145.
- D'Amelio ED, Cohen Y, Des Marais DJ. (1987). Association of a new type of gliding, filamentous, purple phototrophic bacterium inside bundles of *Microcoleus chthonoplastes* in hypersaline cyanobacterial mats. *Arch Microbiol* **147**: 213–220.

- de Wit R, I. Falcón L, Charpy-Roubaud C. (2005). Heterotrophic dinitrogen fixation (acetylene reduction) in phosphate-fertilised *Microcoleus chthonoplastes* microbial mat from the hypersaline inland lake 'la Salada de Chiprana' (NE Spain). *Hydrobiologia* **534**: 245–253.
- Dubinin AV, Gerasimenko LM, Zavarzin GA. (1992). Nitrogen fixation by cyanobacterium *Microcoleus chthonoplastes* from hypersaline lagoons of lake Sivash. *Microbiology* **61**: 593–597.
- Garcia-Pichel F, Prufert-Bebout L, Muyzer G. (1996). Phenotypic and phylogenetic analyses show *Microcoleus chthonoplastes* to be a cosmopolitan cyanobacterium. *Appl Environ Microbiol* **62**: 3284–3291.
- Hammer O, Harper DAT, Ryan PD.. (2001). Paleontological Statistics Software Package for Education and Data Analysis. *Palaeontologia Electronica* **4**: 9.
- Ikemura T. (1985). Codon usage and tRNA content in unicellular and multicellular organisms. *Mol Biol Evol* **2**: 13–34.
- Janson S, Matveyev A, Bergman B. (1998). The presence and expression of *hetR* in the non-heterocystous cyanobacterium *Symploca* PCC 8002. *FEMS Microbiol Lett* **168**: 173–179.
- Jørgensen BB, Cohen Y, Des Marais DJ. (1987). Photosynthetic action spectra and adaptation to spectral light distribution in a benthic cyanobacterial mat. *Appl Environ Microbiol* **53**: 879–886.
- Jørgensen BB, Cohen Y, Revsbech NP. (1986). Transition from anoxygenic to oxygenic photosynthesis in a *Microcoleus chthonoplastes* cyanobacterial Mat. *Appl Environ Microbiol* **51**: 408–417.
- Kleckner N. (1990). Regulation of transposition in bacteria. *Annu Rev Cell Biol* **6**: 297–327.
- Koonin EV, Makarova KS, Aravind L. (2001). Horizontal gene transfer in prokaryotes: quantification and classification. *Annu Rev Microbiol* **55**: 709–742.
- Lavergne J. (1982). Mode of action of 3-(3,4-dichlorophenyl)-1,1-dimethylurea. Evidence that the inhibitor competes with plastoquinone for binding to a common site on the acceptor side of photosystem II. *Biochimica et Biophysica Acta* **682**: 345–353.
- Malin G, Pearson HW. (1988). Aerobic nitrogen fixation in aggregate-forming cultures of the nonheterocystous cyanobacterium *Microcoleus chthonoplastes*. *J Gen Microbiol* **134**: 1755–1763.
- Martin DE, Reinhold-Hurek B. (2002). Distinct roles of PII-like signal transmitter proteins and *amtB* in regulation of *nif* gene expression, nitrogenase activity, and posttranslational modification of NifH in *Azoarcus* sp. Strain BH72. *J Bacteriol* **184**: 2251–2259.
- Olson JB, Litaker RW, Paerl HW. (1999). Ubiquity of heterotrophic diazotrophs in marine microbial mats. *Aquat Microb Ecol* **19**: 29–36.
- Omeregíe EO, Crumbliss LL, Bebout BM, Zehr JP. (2004). Comparison of diazotroph community structure in *Lyngbya* sp. and *Microcoleus chthonoplastes* dominated microbial mats from Guerrero Negro, Baja, Mexico. *FEMS Microbiol Ecol* **47**: 305–319.
- Paerl HW, Pinckney JL, Steppe TF. (2000). cyanobacterial-mat consortia: examining the functional unit of microbial survival and growth in extreme environments. *Environ Microbiol* **2**: 11–26.
- Pearson HW, Howsley R, Kjeldsen CK, Walsby AE. (1979). Aerobic nitrogenase activity associated with a non-heterocystous filamentous cyanobacterium. *FEMS Microbiol Lett* **5**: 163–167.
- Rapp BJ, Wall JD. (1987). Genetic transfer in *Desulfovibrio desulfuricans*. *Proc Natl Acad Sci USA* **84**: 9128–9130.
- Rippka R, Deruelles J, Waterbury JB, Herdman M, Stanier RY. (1979). Generic assignments, strain histories and properties of pure cultures of *Cyanobacteria*. *J Gen Microbiol* **111**: 1–61.
- Rippka R, Waterbury JB. (1977). The synthesis of nitrogenase by non-heterocystous *Cyanobacteria*. *FEMS Microbiol Lett* **2**: 83–86.
- Rubio LM, Ludden PW. (2008). Biosynthesis of the iron-molybdenum cofactor of nitrogenase. *Annu Rev Microbiol* **62**: 93–111.
- Severin I, Stal LJ. (2008). Light dependency of nitrogen fixation in a coastal cyanobacterial mat. *ISME J* **2**: 1077–1088.
- Severin I, Stal LJ. (2009). Diazotrophic microbial mats. In: Seckbach J (ed) *Microbial Mats*. Springer, (in press).
- Siegesmund MA, Johansen JR, Karsten U, Friedl T. (2008). *Coleofasciculus* Gen. Nov (*Cyanobacteria*): morphological and molecular criteria for revision of the Genus *Microcoleus* Gomont. *J Phycol* **44**: 1572–1585.
- Sigalevich P, Meshorer E, Helman Y, Cohen Y. (2000). Transition from anaerobic to aerobic growth conditions for the sulfate-reducing bacterium *Desulfovibrio oxyclinae* results in flocculation. *Appl Environ Microbiol* **66**: 5005–5012.
- Sroga GE. (1997). Regulation of nitrogen fixation by different nitrogen sources in the filamentous non-heterocystous cyanobacterium *Microcoleus* sp. *FEMS Microbiol Lett* **153**: 11–15.
- Stal LJ. (1988). Nitrogen fixation in cyanobacterial mats. In: Packer L, Glazer AN (eds) *Methods in Enzymology, Cyanobacteria*, vol. 167. Academic Press: San Diego, pp 474–484.
- Stal LJ. (2000). Cyanobacterial mats and stromatolites. In: Whitton BA, Potts M (eds). *The Ecology of Cyanobacteria*. Kluwer: Dordrecht, pp 61–120.
- Stal LJ, Krumbein WE. (1985). Isolation and characterization of *Cyanobacteria* from a marine microbial mat. *Botanic Marina* **28**: 351–365.
- Steppe TF, Olson JB, Paerl HW, Litaker RW, Belnap J. (1996). Consortial N₂ fixation: a strategy for meeting nitrogen requirements of marine and terrestrial cyanobacterial mats. *FEMS Microbiol Ecol* **21**: 149–156.
- Steppe TF, Paerl HW. (2002). Potential N₂ fixation by sulfate-reducing bacteria in a marine intertidal microbial mat. *Aquat Microb Ecol* **28**: 1–12.
- Tamura K, Dudley J, Nei M, Kumar S. (2007). MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) software version 4.0. *Mol Biol Evol* **24**: 1596–1599.
- Thiel T, Lyons EM, Erker JC, Ernst A. (1995). A second nitrogenase in vegetative cells of a heterocyst-forming cyanobacterium. *Proc Natl Acad Sci USA* **92**: 9358–9362.
- Wawer C, Jetten M, Muyzer G. (1997). Genetic diversity and expression of the [NiFe] hydrogenase large-subunit gene of *Desulfovibrio* spp. in environmental samples. *Appl Environ Microbiol* **63**: 4360–4369.
- Zehr JP, McReynolds LA. (1989). Use of degenerate oligonucleotides for amplification of the *nifH* gene from the marine cyanobacterium *Trichodesmium thiebautii*. *Appl Environ Microbiol* **55**: 2522–2526.
- Zehr J, Mellon M, Braun S, Litaker W, Steppe T, Paerl H. (1995). Diversity of heterotrophic nitrogen fixation genes in a marine cyanobacterial mat. *Appl Environ Microbiol* **61**: 2527–2532.