



UNIVERSITY OF LEEDS

This is a repository copy of *How Should Beta-Diversity Inform Biodiversity Conservation?*.

White Rose Research Online URL for this paper:

<http://eprints.whiterose.ac.uk/103035/>

Version: Accepted Version

Article:

Socolar, JB, Gilroy, JJ, Kunin, WE et al. (1 more author) (2016) How Should Beta-Diversity Inform Biodiversity Conservation? *Trends in Ecology and Evolution*, 31 (1). pp. 67-80.
ISSN 0169-5347

<https://doi.org/10.1016/j.tree.2015.11.005>

© 2016, Elsevier. This is an author produced version of a paper published in *Trends in Ecology and Evolution*. Uploaded in accordance with the publisher's self-archiving policy.

Reuse

Unless indicated otherwise, fulltext items are protected by copyright with all rights reserved. The copyright exception in section 29 of the Copyright, Designs and Patents Act 1988 allows the making of a single copy solely for the purpose of non-commercial research or private study within the limits of fair dealing. The publisher or other rights-holder may allow further reproduction and re-use of this version - refer to the White Rose Research Online record for this item. Where records identify the publisher as the copyright holder, users can verify any specific terms of use on the publisher's website.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

How should beta-diversity inform biodiversity conservation?

5 **Jacob B. Socolar¹, James J. Gilroy², William E. Kunin³, and David P. Edwards⁴**

¹Department of Ecology and Evolutionary Biology, Princeton University, USA

10 ²School of Environmental Sciences, University of East Anglia, UK

³School of Biology, University of Leeds, UK

⁴Department of Animal and Plant Sciences, University of Sheffield, UK

15 Corresponding authors: Socolar, J.B. (jacob.socolar@gmail.com) and Edwards, D.P. (david.edwards@sheffield.ac.uk)

20 **Abstract**

To design robust protected area networks, accurately measure species losses, or understand the processes that maintain species diversity, conservation science must consider the organization of biodiversity in space. Central is beta-diversity—the component of regional diversity that accumulates from compositional differences between local species assemblages. We review how beta-diversity is impacted by human activities, including farming, selective-logging, urbanisation, species invasions, overhunting, and climate change. Beta-diversity increases, decreases or remains unchanged by these impacts, depending on the balance of processes that cause species composition to become more different (biotic heterogenization) or more similar (biotic homogenization) between sites. While maintaining high beta-diversity is not always a desirable conservation outcome, understanding beta-diversity is essential for protecting regional diversity and can directly assist conservation planning.

Key words: biodiversity conservation; biotic homogenization; alpha-diversity; beta-diversity; gamma-diversity; diversity partitioning; pairwise dissimilarities; species-area relationships; spatial scaling

Conservation targets at multiple spatial scales

As we enter the Anthropocene, humankind is reorganizing the biosphere [1]. Processes ranging from overhunting of large-bodied vertebrates [2] and moving alien species across biogeographic barriers [3] to wholesale clearing of natural habitats for agriculture [4] continue to erode biodiversity. Society values biodiversity at multiple spatial scales, with concerns ranging from local provision of ecosystem services [5] to global preservation of the intrinsic and instrumental value of species [6]. For example, the United Nations Convention on Biological Diversity seeks to stem biodiversity loss at regional, national, and global levels [7].

To assess how best to conserve biodiversity across spatial scales, we need to understand the relationship between locally-collected monitoring data and regional diversity dynamics, and how the mechanisms that maintain diversity vary from local to regional spatial scales [8-11]. Conservationists thus face a key question: how do changes measured locally scale up to regional impacts [12]?

The solution lies in our understanding of “beta-diversity”—the component of regional biodiversity (“gamma-diversity”; see Glossary) that accumulates due to inter-site differences between local species assemblages (“alpha-diversity”; Glossary) [13]. Measured appropriately, changes in beta-diversity provide the scaling factor that allows us to predict changes in gamma-diversity from measured changes in alpha-diversity. Furthermore, new techniques can shed light on the mechanisms responsible for beta-diversity maintenance from alpha-scale survey data, thereby revealing not only how biodiversity is organized in space [14-16], but also the mechanistic underpinnings of these patterns [9,17-19]. Such techniques provide conservation-relevant insights about the maintenance of diversity over large areas.

65 Conservation scientists can incorporate beta-diversity into management
decision-making in a variety of ways. Studies of beta-diversity can quantify
biodiversity loss [20] and inform the placement of protected areas [21,22], the
management of biological invasions [23], and the design of wildlife-friendly
landscapes [24-27]. However, the existence of many different metrics for beta-
diversity, each suited to subtly different questions, has been a source of confusion
and debate in the ecological literature (Box 1) [13,28-30].

70 If beta-diversity is to reliably aid conservationists, it is crucial that we define
and measure it appropriately. We must carefully match appropriate metrics to
specific problems, and judiciously choose spatial scales to measure local and
regional patterns. At its best, beta-diversity is a clarifying concept that unifies
conservation science across spatial scales. But at its worst, it can be a mathematical
abstraction that is easy to misapply and misinterpret. In this review, we highlight
75 recent advances and potential pitfalls in the application of beta-diversity to
conservation science.

Using beta-diversity for conservation: metrics, opportunities, and pitfalls

Diversity loss and spatial scaling

80 Conservation planning requires detailed biodiversity data to inform actions
ranging from land purchases and management [31-33] to agricultural policy and
international carbon payments [34-36]. However, our understanding of the
magnitude of biodiversity loss (or gain) depends on the scale at which we measure
it [37]. For example, local-scale patterns often suggest that diversity is maintained
85 [38,39], whereas global-scale patterns indicate that we are in the midst of an
unprecedented extinction event [40,41]. Scale-dependent differences in diversity
maintenance can even be apparent even within single landscapes or study sites
[20,23]. Therefore, we must understand biodiversity loss at spatial scales relevant to
conservation goals.

90 Carefully interpreted metrics of beta-diversity can help to meet this challenge
by revealing the scaling relationship between alpha- and gamma-diversity.
Diversity partitioning beta-diversity metrics (Box 1) directly provide the scaling
factors that relate alpha- and gamma-diversity, but their calculation requires prior
knowledge of gamma-diversity. Pairwise dissimilarity measures (Box 1) are more
95 widely applicable, but they convey unreliable information about these scaling
relationships [42]. Null model approaches, which are widely used to remove the
neutral component of beta-diversity, tend obscure the scaling relationship between
local and regional diversity [43](Box 2). However, new techniques using the
distance-decay of pairwise similarity provide a potentially powerful tool for
100 extrapolating small studies to larger landscapes [44](Box 3).

Even when the scaling factor is measured appropriately, the conservation
significance of a change in beta-diversity is not straightforward. Maximizing beta-
diversity is not necessarily desirable for gamma-diversity conservation, because
damaging anthropogenic impacts can cause the similarity of local communities to
105 increase, decrease, or remain unchanged, depending on the relative balance of
homogenization and heterogenization processes at the site level (Figure 1). During
the initial stages of anthropogenic impacts, localized species losses and invader

establishment might cause beta-diversity to increase (Figure 1b). Even when beta-diversity decreases, compensatory changes in alpha-diversity can buffer gamma-diversity against declines in beta-diversity (Figure 1a; [45]). For example, increases in the site-occupancy of rare species cause beta-diversity to decline, but bode well for gamma-diversity conservation. Furthermore, different beta-diversity metrics can change in opposite directions in response to a single disturbance event [13]. Therefore, any measured changes in beta-diversity must be interpreted with care.

115

Identifying mechanisms of diversity maintenance

Null model controls and pairwise beta-diversity metrics (Box 1, Box 2) provide mechanistic insight about the maintenance of gamma-diversity, allowing the development of landscape-specific conservation actions (e.g. designing protected area networks). Null model controls can help us partition beta diversity between the component expected by chance (neutral sampling effects) and that driven by environmental or dispersal filters [9,43]. Such distinctions may help guide management, but even 'neutral' beta diversity may be of conservation interest (Box 2). Pairwise dissimilarities can identify key spatial or environmental gradients where turnover occurs [8,18,46], and analysis of pairwise dissimilarities before and after disturbance could pinpoint the environmental gradients along which beta-diversity has been lost, thereby directing the preservation or restoration of key features. By calculating multiple metrics, studies can arrive at a more general understanding of how beta-diversity responds to disturbance (Box 4). A growing body of literature has begun the task of applying these various metrics to assess human impacts on beta-diversity, but it remains critical to recognize what each metric emphasises.

120

125

130

How humans have impacted beta-diversity

When human impacts are patchy in space, beta-diversity is likely to increase at the landscape scale [47,48]. However, human activities often generate completely novel landscapes, with unpredictable changes to alpha-, beta-, and gamma-diversity. Here, we review the beta-diversity impacts of five globally ubiquitous conservation issues. Our primary goal is to extract rules of thumb for interpreting alpha-scale studies of human disturbance. However, patterns of beta-diversity differ not only between disturbance types, but also among taxa and geographic locations. In some cases, patterns of beta-diversity are well established, with clear mechanistic underpinnings, but in many cases patterns are seemingly idiosyncratic or contradictory, without obvious mechanistic explanations. Where rules are not apparent, we advise caution in interpreting the implications of alpha-scale studies for landscape- or regional-scale biodiversity. These areas are frontiers for additional research.

135

140

145

Farming, tree plantations, and selective logging

Intensive pasture and mechanized agriculture usually erode beta-diversity compared to natural habitats and wildlife-friendly agriculture, as the spatially uniform conditions within intensive farmland are tolerable to only a small subset of abundant native species [24,25,49]. For example, Japanese larch plantations have

150

155 homogeneous leaf-litter compared to natural forests, and thus lower beta-diversity
among litter-dwelling invertebrates [50]. In European farms, pesticide-use
decreases multiplicative beta-diversity for many invertebrate taxonomic groups
[51] (but see [25]). Reductions in beta-diversity can persist following cessation of
160 agriculture due to species filtering based on dispersal ability. Understory plant
communities regenerating on abandoned agricultural land tend to have reduced
beta-diversity because they are dominated by dispersal-adapted species compared
to ancient controls [26].

When high-intensity land use reduces the total abundance of the many
species across the assemblage, beta-diversity can increase as species become rarer
(Fig. 1), even though species-environment relationships tend to weaken (Box 4)[20].
165 This increase is identified by null model controls as being driven by neutral
sampling effects of rarity (Fig. 1b). Conversely, when land use intensification leads
to high abundances and local richness of a particular group (e.g. hoverflies in Europe
[44], bees and wasps in Ecuador [52]), beta-diversity is likely to decline.

170 Although there are some clear generalities, many changes appear
idiosyncratic. For example, forest clearance homogenizes soil bacteria and
decreases bacterial beta-diversity in the Brazilian Amazon, but has the opposite
effect in Borneo [53,54]. In Europe, plants and spiders show opposite trends
(increasing and decreasing beta-diversity, respectively) across conventional and
organic wheat fields in Europe [25], while in Borneo, bacteria and birds show
175 different trends following logging (increases and minimal change, respectively)
[33,54]. In Egypt, gardening increases plant heterogeneity compared to natural
habitats by introducing novel crop species, but pollinator heterogeneity remains
unchanged [55]. These findings reflect marked variability in the scaling and
processes of diversity maintenance between regions and taxonomic groups. Further
180 research is needed to better understand how and why these differences arise.

Urbanisation

Urbanization consistently reduces between-city beta-diversity, compounding
severe declines in alpha-diversity [56,57]. Among birds, urbanization decreases the
185 distance decay of compositional similarity between cities [58,59]. In insects,
urbanization reduces beta-diversity because heterogeneous assemblages of
specialists disappear from cities, while consistent suites of tolerant species persist.
For example, in Switzerland diverse assemblages of birch-specialist true bugs and
leafhoppers show high turnover in rural areas, but are absent from cities [60]. In
190 southern California, urbanization increases soil moisture, which permits the
establishment of an invasive ant that homogenizes native ant communities by
excluding all but a few species [61].

Native plant assemblages tend to become more homogeneous with
urbanization [62], and cities often support numerous invasive species, which tend to
195 have lower turnover than natives [63]. Recently established exotics, however, can
show higher beta-diversity than natives [62,64], suggesting a short-term
heterogenizing process prior to the more widespread establishment of invasives
(Figure 1).

200 Despite ubiquitous declines in beta-diversity between cities, there is a more
complicated pattern within cities. Distance decay in Australian birds is high among
both the most urbanized and least urbanized neighbourhoods, but is lower at
intermediate levels of urbanization [59]. Similarly within Berlin, 100 m² tree plots in
the most and least urbanized areas are more dissimilar than intermediately
205 urban species-area relationships are as steep as rural relationships [58].

Biological invasions

Biological invasions affect diversity in two ways: by adding non-native
species and by excluding natives. Both processes can initially heterogenize
210 communities as the invader spreads, but can ultimately result in biotic
homogenization once the invader is ubiquitous (additive homogenization due to the
invader's ubiquity, and subtractive homogenization if natives are extirpated; figure
1)[63,64,66]. Conservationists are mainly concerned with minimizing the impacts of
invasives on native species, rather than maximizing the total diversity of invaded
215 communities (i.e. including both invasives and natives), so the subtractive processes
are of primary interest. We consider two important cases where the impacts of
invasive species can depend on spatial scale: competition effects, exemplified by
plant invasions; and predator-prey effects, exemplified by predator introductions on
oceanic islands.

220 Although the presence of exotic plants can increase alpha-diversity [67],
plant invasions often decrease the diversity of *natives*, at least over small spatial
extents [3]. However, few plant extinctions are attributable to competition from
invaders [68], and recent work suggests that they do not generally drive declines in
gamma-diversity, even of natives [23,69]. Extensive plant surveys from Great
225 Britain suggest that invasives are not causing island-wide extinctions [69]. Targeted
surveys of heavily invaded sites from the United States (Florida, Missouri, and
Hawaii) reveal that invasive plants sharply reduce diversity at very fine spatial
scales (1m² quadrats), but not at moderate scales (500m² plots; figure 2) [23]. Thus,
where invasive plants reduce native diversity at local scales, beta-diversity tends to
230 increase and gamma-diversity is maintained (but see [70]).

Unlike introductions of invasive plants, the introduction of rats, cats and
other predators on oceanic islands has decimated both alpha- and gamma-diversity
of island species [71]. The catastrophic loss of avian diversity in the Pacific, where as
many as 2000 species have disappeared since the arrival of man [72], precipitated a
235 huge decline in beta-diversity, because island-adapted endemic species were
disproportionately likely to go extinct, whereas insular populations of widespread
species typically retained better defences to invasive predators [71]. Among
nonpasserine birds, wide-ranging species were 24 times more likely to persist on
islands than single-island endemics, causing the subtractive homogenization of
240 island communities across large scales [73].

Hunting and fishing

Scant information is available about the impact of hunting and fishing (exploitation)
on beta-diversity. Exploitation often targets large-bodied species, and these tend to

245 range widely. Thus, exploitation may magnify local differences in species
assemblages and increase beta-diversity. For instance, fishing in the northwest
Atlantic targets large-bodied species and increases the decay of community
similarity with distance, an undesirable process of subtractive heterogenization [74].
250 Exploitation also has indirect effects on beta-diversity. For example, bottom-
trawling homogenizes benthic communities by destroying microhabitats on the sea
floor [75]. In tropical forests, hunting removes seed dispersers, causing
hyperdiverse tree assemblages to give way to depauperate communities of species
capable of recruiting near conspecific adults [76]. This subset of trees is likely to be
replicated across space, leading to declines in beta- and gamma-diversity [11].

255

Climate change

Rapid climate change is already causing dramatic shifts in eco-climatic conditions,
threatening species diversity globally [77]. Although community turnover is often
organized along climatic gradients [78], few studies have asked how climate change
260 affects beta-diversity. Available evidence suggests that recent climate change has
increased beta-diversity in some systems by decreasing local richness, without
driving regional extinctions. For example, in California's Sierra Nevada, climate
change has yet to cause regional extinctions of birds or mammals, but has caused
ubiquitous declines in local richness, increasing the neutral component of beta-
265 diversity [79,80]. The long-term effects of climate change on turnover remain to be
seen. Species differ widely in their capacity to track shifting climate envelopes,
depending in part on their dispersal capacity [81], suggesting that future
communities may be dominated by more dispersive taxa. Alongside predicted
climate-driven declines in specialists [77], this could result in increasing
270 homogenization of communities with ongoing climate change, in both natural and
anthropogenic landscapes.

Applications to conservation management

By revealing the spatial scaling of diversity loss and the mechanistic underpinnings
275 of diversity maintenance, beta-diversity has much to offer conservation science.
Here, we discuss the application of beta-diversity to specific longstanding problems
in conservation management.

Protected area selection

280 Given that conservation is underfunded, protected area selection is an
exercise of optimisation [82]. Landscape patterns of beta-diversity have obvious
implications for the SLOSS debate (is it better to create a Single Large Or Several
Small reserves?)[83]. In landscapes with high species turnover along spatial or
environmental gradients, protected area networks must successfully capture this
285 variation or risk losing species [84]. Thus, turnover might favour multiple spatially
disjunct reserves over single large parks [22,85,86]. On the other hand, high neutral
beta diversity or nestedness (richness differences along spatial gradients) imply
that conservation could better focus on diverse sites at the expense of species-poor
areas.

290 When null models suggest that turnover results primarily from neutral
processes in a well-mixed community, conservation should maximize the total area
protected, with less emphasis on protected areas' geographic locations. For example,
only large protected area networks will encompass a full complement of tropical
forest trees, even in well-connected tracts of environmentally homogeneous forest
295 [11].

Corridors and dispersal facilitation

Habitat fragmentation can increase beta diversity via dispersal limitation and
neutral processes [87]. Such patterns may indicate a need to increase between-
300 patch connectivity via corridor creation [88]. Successful corridors might decrease
beta-diversity, at least in the short term, by allowing better mixing between patches.
However, corridor creation is likely to benefit all forms of biodiversity in the long
term, by reducing the likelihood that local and regional extinction debts are realized
[88].

305 Corridors could also facilitate species range-shifts in response to climate
change [81]. These shifts tend to follow poleward and upward temperature shifts in
terrestrial biomes, and downward shifts in aquatic biomes, although all species
within a community will not necessarily shift in the same direction [89,90]. Species
differ markedly in rates of climate-driven movement, reflecting variation in
310 dispersal capacity and phenotypic plasticity [81]. Creating habitat corridors along
climate gradients is likely to aid the movement of poor dispersers, reducing the
potential homogenizing effect of climate change. In the short term, such corridors
may still decrease beta-diversity by facilitating mixing between currently isolated
communities, but the long-term effect is likely to be positive due to avoided
315 extinctions. Because species respond to multiple climatic factors including
temperature and precipitation, identifying the environmental determinants of
species' range limits can help us optimize the location and orientation of such
corridors [18,91].

Land-sharing versus land-sparing agriculture

Agriculture is a major driver of the global extinction crisis [4]. Strategies for
minimizing biodiversity impacts range from implementing low-yielding wildlife-
friendly practices over large areas (land-sharing) to promoting intensification whilst
saving natural habitat for conservation (land-sparing). Quantitative studies of
325 biodiversity loss can reveal the relative merits of the two approaches [34]. However,
most studies have focused on much smaller spatial extents than the regional biota
that they seek to conserve. Encouragingly, the few studies that have looked at beta-
diversity in a land-sparing versus land-sharing context have largely found that
alpha-scale conclusions are unchanged [27,92]. Classically, land-sharing and land-
330 sparing are compared on the basis of population changes across species, rather than
species richness [34]. Therefore, conclusions depend more heavily on whether
compositional change is subtractive or additive than on whether it is homogenizing
or heterogenizing (Figure 1). The loss of specialist species in low-intensity
agriculture is likely to simultaneously favour land-sparing and to drive subtractive

335 homogenization [20,35].

Beta-diversity carries two additional implications for the land-sharing versus land-sparing debate. First, land-sharing is inherently farm-based, whereas intensification on one farm can theoretically spare land at disparate locations. Therefore, the homogenizing effects of farm intensification under land-sparing could be counterbalanced if spared areas are located in an area where beta-diversity is higher or more prone to homogenization. Second, uniformity of agricultural practices and policies over large areas is inherently likely to reduce beta-diversity at coarse scales. This might even be true for agri-environmental management, if the same management practices are applied across large areas, and those practices favour a particular suite of species. A heterogeneous landscape that includes agri-environmental management as well as natural habitat and high-intensity agriculture (land-sparing) might better maintain alpha-, beta- and gamma-diversity at coarse scales, suggesting a need for land-use policy to incorporate diverse options and incentives.

350

Concluding remarks and future directions

Much of what we know in conservation science comes from studies conducted in small areas. Detailed gamma-diversity data will likely never exist for most of the globe, especially at regional scales and in understudied remote tropical regions. Therefore, we need to better understand processes and patterns in the maintenance and loss of beta-diversity in order effectively conserve gamma-diversity [12]. We need to examine how beta-diversity responds during land-use change to understand when and how alpha-scale research can be scaled up to gamma-scale problems and how management interventions can improve gamma-diversity conservation. We cannot afford to abandon the insights gained from alpha-scale studies, but we must update these insights using knowledge about our impacts on species turnover. Recent basic and applied advances in the study of beta-diversity are a welcome beginning [9,20,26].

Current research priorities are myriad (Box 5). Expanding on existing analyses [15,16], we need to document baseline patterns of beta-diversity at increasingly fine resolution. We need a broader and deeper understanding of the effects of land-use and climate change on beta-diversity, sufficient to develop robust methods of extrapolation for interpreting alpha-scale studies. Doing so will require that numerous studies of human biodiversity impacts report consistent metrics for beta-diversity (e.g. 'true' beta-diversity). At the same time, we must identify imperilled natural processes that have historically maintained beta-diversity, and target management to ensure their ongoing operation.

Whereas alpha- and gamma-diversity insights have long underpinned both local-scale conservation actions and regional to global scale policy initiatives [7], the adoption of beta-diversity research into conservation remains a critical frontier. Conservation scientists need to better engage with policy-makers and practitioners to communicate what losses or gains of beta-diversity mean for the global biodiversity extinction crisis, and in turn, how those results should inform decision making. Contemporary threats and solutions in conservation occur at all spatial

380 scales. Beta-diversity provides the link that integrates conservation insights across them all.

Glossary

385 **Alpha-diversity** The species diversity of a relatively small area, frequently expressed as species richness or other low-order Hill number [29]. For the purposes of this review, alpha-diversity corresponds to the scales at which biodiversity studies commonly make measurements (i.e. diversity of a quadrat, plot, or study site).

390 **Beta-diversity** The component of gamma-diversity (see below) that accumulates as a result of differences between sites. Includes heterogeneity resulting from including stochastic variation within a single habitat, differences between habitats along environmental gradients, and changes in species composition between biogeographic provinces [13]. Unlike alpha- and gamma-diversity, beta-diversity does not correspond to the total diversity of some region of space. See box 1.

395 **Gamma-diversity** The species diversity of a relatively large area, expressed in the same units as alpha-diversity (see above). For the purposes of this review, gamma-diversity corresponds to the regional-to-global scales over which society wants to maintain biodiversity (i.e. diversity of a landscape, ecoregion, nation, or planet).

400 **Nestedness** The component of beta-diversity that reflects differences in alpha-diversity between sites when species assemblages at different sites are nested subsets of one another [93]. Contrasts with *turnover* (see below).

Neutral sampling The random assortment of species into samples, either due to sampling errors (e.g. random failure to detect species in a sample) or due to random community assembly in nature.

405 **Similarity (also Compositional similarity)** A metric of how similar the species assemblages of two (or more) sites are. Numerous similarity metrics each define a different formulation of compositional similarity (see box 1).

410 **Turnover** The component of beta-diversity that reflects the replacement of species at some sites by different species at other sites [93]. Contrasts with *nestedness* (see above).

BOX 1: Measuring beta-diversity for conservation science

415 The literature contains numerous beta-diversity metrics [13,29]. All relate to
compositional heterogeneity, but they have subtly distinct biological meanings and
conservation significance.

Diversity partitioning

True beta-diversity (the ratio gamma-diversity divided by mean alpha-diversity)
partitions gamma-diversity into multiplicative alpha and beta components [29,94]
420 (Figure I). Likewise, *additive beta-diversity* partitions gamma into additive
components [21,95]. These exact partitions are ideal for revealing the spatial scaling
of diversity loss, but they can only be used if gamma diversity is known.

Pairwise dissimilarities


Various indices quantify compositional dissimilarity between pairs of sites [28].
425 *Pairwise beta-diversity* is the average dissimilarity across all such pairs within a
region [13] (Figure I). In addition to yielding summary metrics for beta-diversity,
the pairwise dissimilarities are useful for identifying environmental or geographic
features that structure beta-diversity, since the magnitude of the compositional
dissimilarity between two sites should correlate with between-site differences in
430 these features [13].

The choice of dissimilarity metric has been widely discussed [28], but rarely in
a conservation context. While abundance-based measures [30] depend less on
sample size than presence-absence measures, they achieve this precisely by giving
less weight to rare species. Thus, presence-absence measures might be most
435 appropriate for biodiversity conservation geared towards rare species (despite their
sample size dependence), while abundance-based measures might be preferred in
analyses of ecosystem service provision, which is dominated by common species.
 β_{sim} [28] is a widely recommended presence-absence measure that is nearly as
insensitive to sample size as the best abundance-based measures, and measures
440 only turnover (not nestedness; see below). For these reasons, it is especially apt for
identifying spatial and environmental gradients where rare-species turnover occurs,
when sampling is sparse or uneven.

Turnover and nestedness

Whether measuring beta-diversity using diversity-partitioning or using
445 pairwise dissimilarities, ecologists often distinguish between two patterns of beta-
diversity termed *turnover* and *nestedness* [93]. Turnover occurs when species
present at one site are absent at another site, but are replaced by other species
absent from the first. Nestedness occurs when species present at one site are absent
at another, but are not replaced by additional species. Turnover across natural sites
450 implies that conservation must target multiple sites, while nestedness suggests that
conservation might target the richest sites.

Box 1, Fig I



455

Figure I. When local sites harbour different species, beta-diversity can be calculated either by comparing the average alpha-diversity to the total gamma-diversity (diversity partitioning), or by assessing pairwise dissimilarities between local sites.

460

BOX 2: Null models for beta-diversity

465 Even in homogeneous communities, sampling effects can lead to heterogeneity among sites [9]. Rare species will typically be absent from some local samples, and samples from a species-rich assemblage will contain different subsets of the assemblage, even if the community is well mixed. Whenever gamma-diversity exceeds alpha-diversity, these neutral sampling effects ensure that the average pairwise dissimilarity between sites is nonzero, even if species sort randomly into sites.

470 Null models are available to distinguish neutral sampling effects from beta-diversity that exists due to ecological mechanisms such as dispersal limitation or environmental filtering [9,99,100]. Null models randomly reshuffle species identities among the local biodiversity samples, generating a null expectation for beta-diversity under a random assembly process that controls for alpha- and
475 gamma-diversity. These models are important when using beta-diversity to infer mechanisms of community assembly, because they calculate the amount of beta-diversity attributable purely to neutral assembly processes [43]. Frequently, this null expectation is subtracted from the observed beta-diversity metrics as a correction to remove the neutral sampling effects.

480 While null models are highly useful for distinguishing mechanisms of community assembly, they are not helpful for understanding the scaling relationship between the local samples and gamma-diversity. To do so, it is crucial to include scaling that results from neutral assembly processes. Some incidence-based null models fully control for the difference between alpha- and gamma-diversity [99],
485 and therefore cannot yield meaningful estimates of the scaling relationship between the two.

Abundance-based null models also obscure the scaling relationship. To illustrate, consider a forest with 20 ubiquitous common species and 20 rare species. A typical tree plot contains 19-20 of the common species, but only 1-2 rare species.
490 Plots differ due to sampling effects driven by the rare species. If every rare species goes extinct, plot-scale diversity changes by only 5-10%, but gamma-diversity is halved. The null model correctly concludes that only the null component of beta-diversity has changed—yet this component matters crucially to the scaling relationship.

495

Box 3: Upscaling biodiversity


Regional-scale conservation decisions often rely on estimates of gamma-diversity that are extrapolated from a sparse set of local biodiversity samples. Prominent techniques for estimating gamma-diversity assume that local samples are
500 independently drawn from a single regional community [96], but within-region dispersal limitation or environmental filtering violate this assumption. Therefore, measurements of beta-diversity are important for understanding the spatial scaling of species richness.

Traditional pairwise methods do not automatically yield estimates of gamma-diversity or the spatial scaling between alpha and gamma, because they fail to
505 account for patterns of co-occurrence among more than two sites [42]. Recently Hui and McGeoch [97] proposed a new method, generalising beta-diversity to examine overlap in trios, quartets, and larger collections of samples. “Zeta-diversity” (ζ_i) describes the species shared by any collection of i samples, such that ζ_1 is the
510 average richness of a single sample, ζ_2 is the average number of species shared by pairs of samples, ζ_3 is the average shared by trios, and so on. Zeta inevitably declines with i , but the functional form of this decline may vary (most frequently either as power-law or exponential). This approach effectively constructs a
515 “collector’s curve” of increasing sample number, and allows it to be partitioned into the contribution of successively higher levels of overlap. However, unless one has an exhaustive set of samples covering the whole region of interest, one can only assess gamma-diversity by extrapolation.

A different approach is to take advantage of the rich spatial information contained in pairwise dissimilarities to directly estimate gamma-diversity (and
520 indeed the full species-area relationship). This can be done for specific idealised models of community structure [98], but until recently no general formulation was available. Azaele et al. [44] use a general pair correlation function (PCF) to empirically fit the spatial turnover of species as a function of distance (Figure I). The
525 technique has so far been tested in a limited number of systems, but represents an important general approach for multi-scale biodiversity monitoring. Initial tests on well-studied forest plot data suggest this offers a useful new approach to link the beta-diversity of local samples to the gamma-diversity of the region from which they are drawn. The approach can be adapted to incorporate virtually any species-abundance distribution or PCF, or to incorporate environmental as well as spatial
530 distances. Such extensions should provide a new and powerful toolbox for investigating beta-diversity and spatial scaling.

Box 3, Fig. I

535



540

545

550

555

Figure I. (A) Power-law zeta-diversity (ζ_i) curves for trees sampled by 20 x 20 m quadrats on Barro Colorado Island, Panama (+), and birds sampled in quarter-degree cells in southern Africa (X) [97]. Zeta-diversity describes higher order overlaps between sites, but has not been used as a tool for inferring gamma-diversity. **(B,C)** The similarity of hoverfly assemblages in England decays more rapidly with distance on organic farms **(B)** than on conventional farms **(C)**, and this is reflected in the respective pair correlation functions (PCF). Dots represent empirical PCF ($\pm 1SD$) and curves are best-fitted curves. **(D)** As a consequence, the upscaled species-area relationships inferred by Azaele et al. [44] cross; thus while organic farms have fewer species in a local sample, they are predicted to have more species at landscape-scales larger than ~ 4 hectares. **(E)** Conventional farms are dominated by highly mobile species with larvae that feed on cereal aphids, while organic farms exhibit a large fraction species belonging to other feeding guilds that show higher turnover, such as this *Chrysotoxum sp.* Data figures from [97] (A) and [44] (B-D) and photograph reprinted with permission from William Kunin.

Box 4: Multiple beta-diversity metrics in Costa Rican agriculture

560 Low-intensity agriculture in Costa Rica supports local bird communities that are nearly as diverse as forest, but intensification erodes this diversity [35]. To better balance farming and nature in tropical countryside, it is important to understand how agriculture impacts beta-diversity. Karp et al [20] used multiple beta-diversity metrics to understand how beta-diversity changes across three land-use categories: forest, low-intensity agriculture (LIA), and high-intensity agriculture (HIA; Figure I).

565 Countervailing the decline in alpha-diversity, both pairwise beta-diversity and true beta-diversity are slightly higher in LIA than forest, and substantially higher in HIA. Additive beta-diversity (from data in [35]) reveals a similar picture, except that forest and LIA are indistinguishable, with HIA much higher. Thus, disturbance impacts gamma-diversity much less than alpha-diversity.


570 However, pairwise dissimilarities contain additional information. When Karp et al examined pairs of sites from environmentally disparate areas, HIA had significantly lower dissimilarities (i.e. higher beta-diversity) than either forest or LIA. Regressing pairwise dissimilarities on geographic distance between sites revealed that the similarity decreased with distance more rapidly in forest and LIA than HIA. Thus, HIA did not appear to maintain the compositional difference between disparate locations as well as other land uses.

580 Seeking to explain HIA's very high beta-diversity, Karp et al. noticed that total bird abundances were very low in these habitats. Thus, they implemented a null model control to calculate the expected beta diversity if birds sorted randomly into local samples. They found that this neutral sorting accounted for a large portion of HIA beta-diversity, a sizeable portion of LIA beta-diversity, and a much lower portion of forest beta-diversity. "Null" beta-diversity was therefore maintaining gamma-diversity in agricultural habitats, even as bird communities were driven to low total abundance.

585 What are we to conclude from these disparate patterns? First, because the scaling factor from alpha to gamma-diversity is by far highest in HIA, gamma-diversity consequences of intensification could be less dire than alpha-scale measurements might suggest (although gamma consequences may be time-lagged due to extinction debt). Second, intensification homogenizes bird diversity at large spatial scales, which bodes ill for the prospects for tropical diversity in massive swaths of high-intensity agriculture. Third, HIA (and to a lesser extent LIA) maintain high beta-diversity largely through neutral sampling effects and not through species-environment relationships, which could indicate trouble for habitat specialists. Finally, multiple measures of beta-diversity paint a fuller picture of change than any single metric.

595

Box 4, Fig I



600

Figure I. Impacts of farming in Costa Rica on beta-diversity. **(A)** Natural forest, **(B)** low-intensity agriculture, and **(C)** high-intensity agriculture studied by Karp et al. **(D)** Point estimates of true beta-diversity show very high beta-diversity in high-intensity agriculture. **(E)** Null model controls reveal that much of the beta-diversity in high-intensity agriculture is due to neutral sampling effects. **(F)** The distance decay of similarity is lowest in high-intensity agriculture, so distant agricultural sites are more similar to each other than are distant forest sites from each other. Photos courtesy of D. Karp and J. Zook; data figures from [20], reprinted with permission from John Wiley and Sons.

610

Box 5: Outstanding questions

Research questions

615 *Upscaling from samples to gamma-diversity:* The estimation of gamma-diversity
from a set of samples is a complicated problem, and even more so when spatial
structure means that samples are not independent. A novel approach offers
substantial promise [44] (Box 3), but has yet to be repeatedly tested and improved.
Modifying the framework to include environmental distances between sites and
geographic barriers is a frontier, and validating the framework's predictions will be
620 a key challenge.

Developing rules of thumb: As yet, we cannot say with confidence how beta-diversity
is likely to respond to most human activities (urbanization is a notable exception),
although experience to date suggests that the effects may depend critically on the
625 extent (in space and time) of the modification. Rules of thumb based on replicated
empirical studies reporting standard beta-diversity metrics would allow
conservation to qualitatively extrapolate alpha-scale data to larger regions, even
when quantitative upscaling is not possible. This could have far-reaching
consequences for contemporary questions such as the land-sharing versus land-
630 sparing debate.

Data gaps

Taxa: The available data on how beta-diversity responds to human impacts is biased
towards plants and birds. Since other organisms have dramatically different modes
635 and patterns of dispersal, we must better understand how their beta-diversity is
changing in the Anthropocene.

Systems: Studies of beta-diversity are biased towards terrestrial systems in
temperate and tropical latitudes. Very little information is available regarding beta-
640 diversity change in aquatic systems, and essentially no reports have assessed the
effect of anthropogenic change on beta-diversity at high latitudes, which generally
show less natural beta-diversity (i.e., in the absence of human impacts) than at
lower latitudes [15].

645 *Disturbances:* While the beta-diversity consequences of farming and invasions are
comparatively well studied, the literature contains very little on the consequences of
climate change and hunting, and even less on myriad other disturbances ranging
from surface pollution to alternative energy development. Even for a specific mode
of disturbance, the spatial pattern and extent of disturbance might further influence
650 beta-diversity.

Replication: Even in relatively data-rich taxa, systems, and disturbances, we so far
lack the replication of results necessary to separate signal from noise and build rules
of thumb. We suggest that almost any biodiversity study that incorporates land-use
655 could beneficially calculate and report beta-diversity metrics, thus contributing to
an emerging understanding of biodiversity loss across spatial scales.

Acknowledgments

We thank Paul Elsen, Xingli Giam, Bethanne Bruninga-Socolar, David Wilcove, and three anonymous reviewers for helpful discussions and comments. JBS was supported by an NSF GRFP; JJG by a Norwegian Research Council fellowship awarded to DPE; WEK was supported by the EU FP7 EU-BON project.

References

- 1 Corlett, R.T. (2015) The Anthropocene concept in ecology and conservation. *Trends in Ecology & Evolution* 30, 36–41
- 665 2 Dirzo, R. *et al.* (2014) Defaunation in the anthropocene. *Science* 345, 20150552
- 3 Vila, M. *et al.* (2011) Ecological impacts of invasive alien plants: a meta-analysis of their effects on species, communities and ecosystems. *Ecol Letters* 14, 702–708
- 670 4 Laurance, W.F. *et al.* (2014) Agricultural expansion and its impacts on tropical nature. *Trends in Ecology & Evolution* 29, 107–116
- 5 Gamfeldt, L. *et al.* (2013) Higher levels of multiple ecosystem services are found in forests with more tree species. *Nat Comms* 4, 1340
- 6 Cafaro, P. and Primack, R. (2014) Biological Conservation. *Biological Conservation* 170, 1–2
- 675 7 Secretariat of the Convention on Biological Diversity. (2014) *Global biodiversity outlook 4*
- 8 McClain, C.R. *et al.* (2012) Dispersal, environmental niches and oceanic-scale turnover in deep-sea bivalves. *Proceedings of the Royal Society B: Biological Sciences* 279, 1993–2002
- 680 9 Kraft, N.J.B. *et al.* (2011) Disentangling the drivers of β diversity along latitudinal and elevational gradients. *Science* 333, 1755–1758
- 10 Chesson, P. and Kuang, J.J. (2008) The interaction between predation and competition. *Nature* 456, 235–238
- 685 11 Terborgh, J. (2012) Enemies Maintain Hyperdiverse Tropical Forests. *Am. Nat.* 179, 303–314
- 12 Gardner, T.A. *et al.* (2013) A social and ecological assessment of tropical land uses at multiple scales: the Sustainable Amazon Network. *Philosophical Transactions of the Royal Society B: Biological Sciences* 368, 20120166–20120166
- 690 13 Anderson, M.J. *et al.* (2011) Navigating the multiple meanings of β diversity: a roadmap for the practicing ecologist. *Ecol Letters* 14, 19–28
- 14 Koleff, P. *et al.* (2003) Are there latitudinal gradients in species turnover? *Global Ecology and Biogeography* 12, 483–498
- 695 15 McKnight, M.W. *et al.* (2007) Putting Beta-Diversity on the Map: Broad-Scale Congruence and Coincidence in the Extremes. *PLoS Biology* 5, e272
- 16 Buckley, L.B. and Jetz, W. (2008) Linking global turnover of species and environments. *Proceedings of the National Academy of Sciences of the United States of America* 105, 17836–17841

- 700 17 Veech, J.A. and Crist, T.O. (2007) Habitat and climate heterogeneity maintain
beta-diversity of birds among landscapes within ecoregions. *Global Ecology
and Biogeography* 16, 650–656
- 18 Jankowski, J.E. *et al.* (2009) Beta diversity along environmental gradients:
implications of habitat specialization in tropical montane landscapes. *J Anim
Ecol* 78, 315–327
- 705 19 Stegen, J.C. *et al.* (2012) Stochastic and deterministic drivers of spatial and
temporal turnover in breeding bird communities. *Global Ecology and
Biogeography* 22, 202–212
- 20 Karp, D.S. *et al.* (2012) Intensive agriculture erodes β -diversity at large
710 scales. *Ecol Letters* 15, 963–970
- 21 Gering, J.C. *et al.* (2003) Additive partitioning of species diversity across
multiple spatial scales: implications for regional conservation of biodiversity.
Conservation Biology 17, 488–499
- 22 Wiersma, Y.F. and Urban, D.L. (2005) Beta diversity and nature reserve
715 system design in the Yukon, Canada. *Conservation Biology* 19, 1262–1272
- 23 Powell, K.I. *et al.* (2013) Invasive plants have scale-dependent effects on
diversity by altering species-area relationships. *Science* 339, 316–318
- 24 Gabriel, D. *et al.* (2006) Beta diversity at different spatial scales: plant
communities in organic and conventional agriculture. *Ecol Appl* 16, 2011–
720 2021
- 25 Clough, Y. *et al.* (2007) Alpha and beta diversity of arthropods and plants in
organically and conventionally managed wheat fields. *Journal of Applied
Ecology* 44, 804–812
- 26 Vellend, M. *et al.* (2007) Homogenization of forest plant communities and
725 weakening of species-environment relationships via agricultural land use.
Journal of Ecology 95, 565–573
- 27 Edwards, D.P. *et al.* (2014) Land-sharing versus land-sparing logging:
reconciling timber extraction with biodiversity conservation. *Global Change
Biology* 20, 183–191
- 730 28 Koleff, P. *et al.* (2003) Measuring beta diversity for presence–absence data. *J
Anim Ecol* 72, 367–382
- 29 Tuomisto, H. (2010) A diversity of beta diversities: straightening up a
concept gone awry. Part 1. Defining beta diversity as a function of alpha and
gamma diversity. *Ecography* 33, 2–22
- 735 30 Barwell, L.J. *et al.* (2015) Measuring β -diversity with species abundance data.
J Anim Ecol 84, 1112–1122
- 31 Giam, X. *et al.* (2015) Mitigating the impact of oil-palm monoculture on
freshwater fishes in Southeast Asia. *Conservation Biology* 29, 1357–1367
- 32 Clough, Y. *et al.* (2011) Combining high biodiversity with high yields in
740 tropical agroforests. *Proceedings of the National Academy of Sciences* 108,
8311–8316
- 33 Edwards, D. *et al.* (2010) Degraded lands worth protecting: the biological
importance of Southeast Asia's repeatedly logged forests. *Proceedings of the
Royal Society B: Biological Sciences* 278, 82–90
- 745 34 Phalan, B. *et al.* (2011) Reconciling food production and biodiversity

- Conservation: land sharing and land sparing compared. *Science* 333, 1289–1291
- 35 Frishkoff, L.O. *et al.* (2014) Loss of avian phylogenetic diversity in
neotropical agricultural systems. *Science* 345, 1343–1346
- 750 36 Gilroy, J.J. *et al.* (2014) Cheap carbon and biodiversity co-benefits from
forest regeneration in a hotspot of endemism. *Nature Climate Change* 4,
503–507
- 37 McGill, B.J. *et al.* (2015) Fifteen forms of biodiversity trend in the
Anthropocene. *Trends in Ecology & Evolution* 30, 104–113
- 755 38 Vellend, M. *et al.* (2013) Global meta-analysis reveals no net change in local-
scale plant biodiversity over time. *Proceedings of the National Academy of
Sciences* 110, 19456–19459
- 39 Dornelas, M. *et al.* (2014) Assemblage time series reveal biodiversity change
but not systematic loss. *Science* 344, 296–299
- 760 40 Barnosky, A.D. *et al.* (2011) Has the Earth's sixth mass extinction already
arrived? *Nature* 470, 51–57
- 41 Pimm, S.L. *et al.* (2014) The biodiversity of species and their rates of
extinction, distribution, and protection. *Science* 344, 1246752
- 42 Baselga, A. (2013) Multiple site dissimilarity quantifies compositional
765 heterogeneity among several sites, while average pairwise dissimilarity may
be misleading. *Ecography* 36, 124–128
- 43 Tucker, C.M. *et al.* (2015) Differentiating between niche and neutral
assembly in metacommunities using null models of β -diversity. *Oikos* DOI:
10.1111/oik.02803
- 770 44 Azaele, S. *et al.* (2015) Towards a unified descriptive theory for spatial
ecology: predicting biodiversity patterns across spatial scales. *Methods in
Ecology and Evolution* 6, 324–332
- 45 McCune, J.L. and Vellend, M. (2013) Gains in native species promote biotic
homogenization over four decades in a human-dominated landscape. *Journal
775 of Ecology* 101, 1542–1551
- 46 Pomara, L.Y. *et al.* (2014) Avian species composition across the Amazon
River: the roles of dispersal limitation and environmental heterogeneity.
Journal of Biogeography 41, 784–796
- 47 Gutiérrez-Cánovas, C. *et al.* (2013) Contrasting effects of natural and
780 anthropogenic stressors on beta diversity in river organisms. *Global Ecology
and Biogeography* 22, 796–805
- 48 Kessler, M. *et al.* (2009) Alpha and beta diversity of plants and animals along
a tropical land-use gradient. *Ecol Appl* 19, 2142–2156
- 49 Solar, R.R. de C. *et al.* (2015) How pervasive is biotic homogenization in
785 human-modified tropical forest landscapes? *Ecol Letters* 18, 1108–1118
- 50 Mori, A.S. *et al.* (2014) Biotic homogenization and differentiation of soil
faunal communities in the production forest landscape: taxonomic and
functional perspectives. *Oecologia* 177, 533–544
- 51 Dormann, C.F. *et al.* (2007) Effects of landscape structure and land-use
790 intensity on similarity of plant and animal communities. *Global Ecology and
Biogeography* 16, 774–787

- 52 Tylianakis, J.M. *et al.* (2006) Spatial scale of observation affects alpha, beta
and gamma diversity of cavity-nesting bees and wasps across a tropical
land-use gradient. *Journal of Biogeography* 33, 1295–1304
- 795 53 Rodrigues, J.L.M. *et al.* (2013) Conversion of the Amazon rainforest to
agriculture results in biotic homogenization of soil bacterial communities.
Proceedings of the National Academy of Sciences 110, 988–993
- 54 Lee-Cruz, L. *et al.* (2013) Impact of logging and forest conversion to oil palm
plantations on soil bacterial communities in Borneo. *Applied and*
800 *Environmental Microbiology* 79, 7290–7297
- 55 Norfolk, O. *et al.* (2015) Contrasting patterns of turnover between plants,
pollinators and their interactions. *Diversity and Distributions* 21, 405–415
- 56 Aronson, M.F.J. *et al.* (2014) A global analysis of the impacts of urbanization
on bird and plant diversity reveals key anthropogenic drivers. *Proceedings of*
805 *the Royal Society B: Biological Sciences* 281, 20133330–20133330
- 57 Mckinney, M.L. (2006) Urbanization as a major cause of biotic
homogenization. *Biological Conservation* 127, 247–260
- 58 Ferenc, M. *et al.* (2013) Are cities different? Patterns of species richness and
beta diversity of urban bird communities and regional species assemblages
810 in Europe. *Global Ecology and Biogeography* 23, 479–489
- 59 Luck, G.W. and Smallbone, L.T. (2010) The impact of urbanization on
taxonomic and functional similarity among bird communities. *Journal of*
Biogeography 38, 894–906
- 60 Knop, E. (2015) Biotic homogenization of three insect groups due to
815 urbanization. *Global Change Biology* DOI: 10.1111/gcb.13091
- 61 Holway, D.A. and Suarez, A.V. (2006) Homogenization of ant communities in
mediterranean California: The effects of urbanization and invasion.
Biological Conservation 127, 319–326
- 62 Kühn, I. and Klotz, S. (2006) Urbanization and homogenization – Comparing
820 the floras of urban and rural areas in Germany. *Biological Conservation* 127,
292–300
- 63 La Sorte, F.A. *et al.* (2014) Beta diversity of urban floras among European
and non-European cities. *Global Ecology and Biogeography* 23, 769–779
- 64 Ricotta, C. *et al.* (2011) Phylogenetic beta diversity of native and alien
825 species in European urban floras. *Global Ecology and Biogeography* 21, 751–
759
- 65 Trentanovi, G. *et al.* (2013) Biotic homogenization at the community scale:
disentangling the roles of urbanization and plant invasion. *Diversity and*
Distributions 19, 738–748
- 830 66 McKinney, M. and Lockwood, J. (1999) Biotic homogenization: a few winners
replacing many losers in the next mass extinction. *Trends in Ecology &*
Evolution 14, 450–453
- 67 Rosenzweig, M.L. (2001) The four questions: what does the introduction of
exotic species do to diversity? *Evol Ecol Res* 3, 361–367
- 835 68 Sax, D.F. *et al.* (2002) Species invasions exceed extinctions on islands
worldwide: a comparative study of plants and birds. *Am. Nat.* 160, 766–783
- 69 Thomas, C.D. and Palmer, G. (2015) Non-native plants add to the British

flora without negative consequences for native diversity. *Proceedings of the National Academy of Sciences* 112, 4387–4392

840 70 Stohlgren, T.J. and Rejmanek, M. (2014) No universal scale-dependent impacts of invasive species on native plant species richness. *Biology Letters* 10, 20130939–20130939

71 Blackburn, T.M. *et al.* (2004) Avian extinction and mammalian introductions on oceanic islands. *Science* 305, 1955–1958

845 72 Steadman, D.W. (1995) Prehistoric extinctions of Pacific island birds: biodiversity meets zooarchaeology. *Science* 267, 1123–1131

73 Duncan, R.P. *et al.* (2013) Magnitude and variation of prehistoric bird extinctions in the Pacific. *Proceedings of the National Academy of Sciences* 110, 6436–6441

850 74 Shackell, N.L. *et al.* (2012) Spatial scale of similarity as an indicator of metacommunity stability in exploited marine systems. *Ecol Appl* 22, 336–348

75 Hewitt, J.E. *et al.* (2005) The importance of small-scale habitat structure for maintaining beta diversity. *Ecology* 86, 1619–1626

855 76 Terborgh, J. *et al.* (2008) Tree recruitment in an empty forest. *Ecology* 89, 1757–1768

77 Urban, M.C. (2015) Accelerating extinction risk from climate change. *Science* 348, 571–573

78 Guerin, G.R. *et al.* (2013) Spatial modelling of species turnover identifies climate ecotones, climate change tipping points and vulnerable taxonomic groups. *Ecography* 36, 1086–1096

860 79 Moritz, C. *et al.* (2008) Impact of a century of climate change on small-mammal communities in Yosemite National Park, USA. *Science* 322, 261–264

80 Tingley, M.W. and Beissinger, S.R. (2013) Cryptic loss of montane avian richness and high community turnover over 100 years. *Ecology* 94, 598–609

865 81 Pearson, R.G. (2006) Climate change and the migration capacity of species. *Trends in Ecology & Evolution* 21, 111–113

82 Venter, O. *et al.* (2014) Targeting global protected area expansion for imperiled biodiversity. *PLoS Biology* 12, e1001891

870 83 Simberloff, D.S. and Abele, L.G. (1976) Island biogeography theory and conservation practice. *Science* 191, 285–286

84 Tuomisto, H. *et al.* (2003) Dispersal, environment, and floristic variation of western Amazonian forests. *Science* 299, 241–244

85 Tschardt, T. *et al.* (2002) Contribution of small habitat fragments to conservation of insect communities of grassland-cropland landscapes. *Ecol Appl* 12, 354–363

875 86 Quinn, J.F. and Harrison, S.P. (1988) Effects of habitat fragmentation and isolation on species richness: evidence from biogeographic patterns. *Oecologia* 75, 132–140

880 87 Püttker, T. *et al.* (2014) Ecological filtering or random extinction? Beta-diversity patterns and the importance of niche-based and neutral processes following habitat loss. *Oikos* 124, 206–215

88 Fagan, W.F. (2002) Connectivity, fragmentation, and extinction risk in

dendritic metapopulations. *Ecology* 83, 3243-3249

885 89 VanDerWal, J. (2012) Focus on poleward shifts in species' distribution underestimates the fingerprint of climate change. *Nature Climate change* 3, 239-243

90 Gillings, S. *et al.* (2015) Directionality of recent bird distribution shifts and climate change in Great Britain. *Global Change Biology* 21, 2155-2168

890 91 Tingley, M.W. *et al.* (2009) Birds track their Grinnellian niche through a century of climate change. *Proceedings of the National Academy of Sciences* 106, 19637-19643

92 Egan, J.F. and Mortensen, D.A. (2012) A comparison of land-sharing and land-sparing strategies for plant richness conservation in agricultural

895 landscapes. *Ecol Appl* 22, 459-471

93 Baselga, A. (2010) Partitioning the turnover and nestedness components of beta diversity. *Global Ecology and Biogeography* 19, 134-143

94 Jost, L. *et al.* (2009) Partitioning diversity for conservation analyses. *Diversity and Distributions* 16, 65-76

900 95 Crist, T.O. and Veech, J.A. (2006) Additive partitioning of rarefaction curves and species-area relationships: unifying α -, β - and γ -diversity with sample size and habitat area. *Ecol Letters* 9, 923-932

96 Chao, A. *et al.* (2014) Rarefaction and extrapolation with Hill numbers: a framework for sampling and estimation in species diversity studies. *Ecol Monogr* 84, 45-67

905 97 Hui, C. and McGeoch, M.A. (2014) Zeta diversity as a concept and metric that unifies incidence-based biodiversity patterns. *Am. Nat.* 184, 684-694

98 Rosindell, J. and Cornell, S.J. (2007) Species-area relationships from a spatially explicit neutral model in an infinite landscape. *Ecol Letters* 10, 586-595

910 99 Chase, J.M. *et al.* (2011) Using null models to disentangle variation in community dissimilarity from variation in α -diversity. *Ecosphere* 2, art24


100 Mori, A.S. *et al.* (2014) Null model approaches to evaluating the relative role of different assembly processes in shaping ecological communities.

915 *Oecologia* 178, 261-273

920

Figures

Figure 1



925 **Figure 1:** Beta-diversity can change in a variety of ways following human impacts.
930 **(A)** Beta-diversity can change as a result of local species gains (additive processes) and local species losses (subtractive processes). Beta-diversity increases when ubiquitous species disappear from some or all sites (1) or when new species arrive at some sites, but do not become ubiquitous (2). Beta-diversity decreases when rare, non-ubiquitous species go extinct (3) or when formerly rare or absent species become widespread (4). **(B)** A conceptual trajectory for typical patterns of beta-diversity change as human disturbance persists and intensifies. (i) Many native species become rarer, but few go extinct (subtractive heterogenization). Invasive species establish, but few become ubiquitous (additive heterogenization). (ii) Rare species disappear entirely (subtractive homogenization); generalists and invaders begin to dominate (additive homogenization). (iii) Homogeneous environments with little between-site variation. (iv) If assemblages are driven to very low abundance, the neutral component of beta-diversity may again increase.

940 **Figure 2**




Figure 2: (A) An example species-area relationship from hardwood hammocks in Florida, USA. Uninvaded sites (B) have shallower slopes than sites invaded by *Dianella ensifolia* (C). At larger sample areas, the number of species between uninvaded and invaded sites converges. Photos courtesy of K. Powell. Panel (A) from [23]. Reprinted with permission from AAAS.

950