

—Review—

How to Improve the Success Rate of Mouse Cloning Technology

Nguyen Van THUAN¹⁾, Satoshi KISHIGAMI²⁾ and Teruhiko WAKAYAMA³⁾

¹⁾Department of Animal Bioscience and Biotechnology, Konkuk University, Seoul 143-701, South Korea, ²⁾Department of Genetic Engineering, KINKI University, Wakayama 649-6493 and ³⁾Center for Developmental Biology, RIKEN Kobe, Kobe 650-0047, Japan

Abstract. It has now been 13 years since the first cloned mammal Dolly the sheep was generated from somatic cells using nuclear transfer (SCNT). Since then, this technique has been considered an important tool not only for animal reproduction but also for regenerative medicine. However, the success rate is still very low and the mechanisms involved in genomic reprogramming are not yet clear. Moreover, the NT technique requires donated fresh oocyte, which raises ethical problems for production of human cloned embryo. For this reason, the use of induced pluripotent stem cells for genomic reprogramming and for regenerative medicine is currently a hot topic in this field. However, we believe that the NT approach remains the only valid way for the study of reproduction and basic biology. For example, only the NT approach can reveal dynamic and global modifications in the epigenome without using genetic modification, and it can generate offspring from a single cell or even a frozen dead body. Thanks to much hard work by many groups, cloning success rates are increasing slightly year by year, and NT cloning is now becoming a more applicable method. This review describes how to improve the efficiency of cloning, the establishment of clone-derived embryonic stem cells and further applications.

Key words: Clone, ntES cell, Nuclear Transfer, Reprogramming

(J. Reprod. Dev. 56: 20–30, 2010)

Since it was first reported in 1997 that a sheep had been successfully cloned [1], the mouse [2], bovine [3], goat [4], pig [5, 6], gaur [7], mouflon [8], domestic cat [9], rabbit [10], horse [11], mule [12], rat [13], African wildcat [14], dog [15], ferret [16], wolf [17] red deer [18], sand cat [19], ibex [20] and camel [21] are among the mammalian species that have been generated successfully as cloned animals using somatic cell nuclear transfer (SCNT). While cloning efficiencies can range from 0 to 20%, efficiency rates of just 1–2% are typical in mice. Moreover, many abnormalities in mice cloned from somatic cells have been reported, including abnormal gene expression in embryos [22–24] abnormal placentas [25, 26], obesity [27, 28], and early death [29]. Such abnormalities notwithstanding, success in generating cloned offspring has still opened new avenues of investigation.

If cloning success rates depend on the donor cell type, we should be able to determine what is most important for reprogramming. For this reason, many different donor cell types have been examined in efforts to understand genomic reprogramming and how to improve the success rate of cloning by SCNT. The most popular cell types as nuclear donors are cumulus cells [2], tail tip cells (probably fibroblasts) [25, 30], Sertoli cells [31] and embryonic stem (ES) cells [32]. As completely differentiated cells, natural killer T cells [33] and granulocyte [34] have been used. As partially undifferentiated cells, fetal neuronal cells [35], hematopoietic stem cells [36], neuronal stem cells [37, 38] and keratinocyte stem cells [39] have also been used. Immature primordial germ cells were also examined [40]. Unfortunately, all cell types show very

low success rates for full-term cloning.

If we could produce higher success rates using the same donor cell type, this would suggest that genomic reprogramming could be enhanced artificially. Therefore, a variety of different methods have been trialed, such as oocyte activation [41], cell fusion vs. nuclear injection [30], enucleation timing [42], and inhibition of cytokinesis [43] (for review see [44]). However, again, most of these trials have failed to improve the success rate of cloning mice. Recently, we found that the efficiency of mouse cloning could be enhanced by up to five-fold through the addition of the histone deacetylation inhibitor trichostatin A (TSA) into the oocyte activation medium [45], even though high concentrations of TSA are toxic to embryonic development [46]. This result suggests that nuclear reprogramming might be enhanced by chemical treatment, and this method will provide a new approach for practical improvements in mouse cloning techniques and new insights into the genomic reprogramming of somatic cell nuclei.

On the other hand, we have shown that nuclear transfer-derived ES cell lines possess the same potential as do ES cells produced from fertilized blastocysts [47]. To distinguish nuclear transfer-derived ES cell lines from those lines derived from fertilized embryos, the former are referred to as ntES cell lines [48]. Although ntES cells have been shown to be equivalent to ES cells, there are ethical objections concerning human cells because fresh oocytes must be donated by healthy women and the resulting cloned embryos would be deprived of their potential to develop into a complete human being. To avoid these ethical objections, several approaches have been attempted and some problems have been solved. For example, aged oocytes from failed *in vitro* fertilization (IVF) attempts were used instead of freshly donated oocytes [49].

Received: December 15, 2009

Accepted: December 20, 2009

©2010 by the Society for Reproduction and Development

Correspondence: T Wakayama (e-mail: teru@cdb.riken.jp)

The use of induced pluripotent stem (iPS) cells for cloning has no ethical problems and is a relatively easy technique, whereas ntES cells pose serious ethical problems and cloning from them is extremely difficult. Therefore, although there are several reports proposing the use of ntES cells as a model of regenerative medicine [50, 51], the use of these cells in preliminary medical research is waning. However, for basic biology, the ntES cell techniques can be applied to characterize very rare and specialized cell types in the body, such as olfactory neurons. In theory, 5 to 10 donor cells can establish one ntES cell line and, once established, these cells will propagate indefinitely. Another example is to generate cloned animal from ntES cell lines using a second round of NT [48]. Even in infertile and ‘unclonable’ strains of mice, we can generate offspring from somatic cells by combining cloning with ntES technology [52–54]. Moreover, cloned offspring can be generated potentially even from the nuclei of dead bodies or cells [55, 56], such as from an extinct frozen animal. Currently, only the ntES technology is available for this purpose, because all other techniques, including iPS cell derivation, require significant numbers of living donor cells. Previously we have reviewed and described details of the features of cloning and ntES cell technology for both basic and applied investigations up to 2007 [44]. Here, we will describe recent updates.

Abnormalities Found in Cloned Mice

The current SCNT method causes numerous abnormalities in cloned mice that are probably related to the efficiency of successful cloning. We review these clone-specific abnormalities briefly below before considering how we can improve the efficiency of cloning.

Abnormalities in embryonic and extraembryonic lineages

Most cloned embryos die during the pre- and peri-implantation periods of gestation [57] and almost all clones showed abnormal placentas when full-term offspring were examined (Fig. 1A). When imprinted gene expression was examined in full-term placentas, some of the genes showed aberrant expression [58]. Therefore, it was thought that the abnormal establishment of extraembryonic lineages within the developing embryo caused the inefficiency of cloning. However, loss of and reduced pluripotency in the inner cell mass (ICM) lineage of cloned blastocysts has also been reported [22]. We have previously examined the trophoblast (TE) and ICM lineages using Cdx2 and Oct4, key molecules for the specification of TE and ICM fate, respectively [24]. The frequency of Oct4 expressing blastocysts was as low as 50%, as previously reported [22]; however, regardless of Oct4 expression, the majority of the cloned blastocysts (>90%) expressed Cdx2 normally. This result was supported by a recent report in which trophoblast stem (ntTS) cells from cloned mouse blastocysts were established with an efficiency as high as that from fertilized blastocysts [59]. A comprehensive analysis of the transcriptional and epigenetic traits demonstrated that these ntTS cells were indistinguishable from control TS cells. These results suggest that even though SCNT cloned embryos have a reduced potential to produce the ICM lineage, the TE lineage can be established and maintained relatively




Fig. 1. Abnormalities in cloned mice. (A) All cloned mice were born with an extremely large placenta and this was the heaviest in our records. (B) A few cloned mice showed several other abnormalities. This pup was born alive but showed intestinal herniation and lacked a skull.

normal.

Ogura’s group also obtained similar results using a different approach [60]. Embryos aggregated with diploid (2n) cloned and tetraploid (4n) fertilized embryos developed hyperplastic placentas. By contrast, placentas of the reciprocal combination—2n fertilized and 4n cloned embryos—were less hyperplastic. This result suggests that embryonic rather than extraembryonic tissues had more impact on the onset of placental hyperplasia, and that the abnormal placentation in clones occurs in a non cell-autonomous manner.

Epigenetic abnormalities in cloned offspring and their placentas

Recent molecular analyses of cloned embryos have revealed abnormal epigenetic modifications such as DNA methylation and histone modifications [61–63]. Moreover, several abnormalities were found not only in early embryos, but also in full-term offspring (Fig. 1B). Therefore, many cloned animals died soon after birth from respiratory failure or from other abnormalities [32, 64] and many cloned mice die prematurely because of liver necrosis, tumors, and pneumonia [29]. These abnormalities presumably reflect a dysregulation of gene function in cells, which may be

accounted for by aberrant genomic reprogramming—the failure to completely reset the differentiated state of the host cell [58, 65–67]. These findings indicate that a cloned mouse is not a perfect copy of the original mouse in terms of placental development, body weight, or the methylation status of its genomic DNA [63].

However, even though most cloned mice are abnormal and show placentomegaly or skewed X-chromosome inactivation in different tissues or individuals [68], some of them are able to develop into adulthood, which indicates that perfect reprogramming or maintenance of the methylation state is not essential for the overall success of mammalian cloning. Interestingly, those epigenetic abnormalities were corrected with increased age and became similar to those seen in normal mice [69]. Hence, some clones survive as long as non-clones. For example, the first cloned mouse, Cumulina, died at the age of 2 years and 7 months, which is slightly longer than the lifespan of an average mouse (News in Brief; Nature 2000, 405:268). Importantly, those cloned mice-specific abnormalities are not heritable and are absent from the progeny of both clones mated with clones and clones mated with wild type mice [27, 70]. This implies that epigenetic abnormalities, such as imprinting and/or reprogramming phenomena caused by the SCNT procedure, can be corrected during gametogenesis. Therefore, even if cloned animals from endangered species die from their abnormalities, we still have the possibility to rescue them by collecting the gonads because the gametes are probably epigenetically normal [71].

Genetic abnormalities in cloned mice and their offspring

As mentioned above, it was thought that most abnormalities in cloned animals arise epigenetically. However, Kawasumi *et al.* [72] reported that nuclear fragmentation was observed in 30% of 2-cell cloned mouse embryos, which suggests that genetic abnormalities might also cause cloning inefficiency. Recently two important papers were published. Balbach *et al.* found a short tailed mouse when they did cumulus cell NT cloning [73]. Although they failed to identify the genetic abnormality, this phenotype was inherited to the naturally mated offspring. Inoue *et al.* found more clear evidence of genetic abnormality caused by NT [74]. They reported the accidental birth of a female mouse following SCNT using a male donor. This ‘male-derived female’ clone grew into a normal adult and produced offspring by natural mating with a littermate. Chromosomal analysis revealed that the female clone had a 39, X karyotype, indicating that the Y chromosome had been deleted in the donor cell or at some early step during the SCNT procedure. For these reasons, we began to examine the genetic constitution of cloned embryos using a newly developed live cell imaging system [75], and found that more than half of the cloned embryos showed aberrant chromosome segregation during preimplantation development (unpublished). If this result is common to all cloned embryos, it may suggest that the inefficient cloning success rate is not only caused by epigenetically incomplete reprogramming but also arises from genetic damage caused by the SCNT procedure or occurring in the somatic cell donor.

Where are the Genomic Reprogramming Factors?

Zygotes are inappropriate recipients

Historically, the first report of cloned mammals was with mice, which were generated by the coordinated microinjection of ICM nuclei into zygotes and the immediate removal of the preexisting pronuclei [76]. However, this replacement procedure has been controversial as all subsequent attempts failed to reproduce the result [77]. Therefore, enucleated metaphase II oocytes have been the preferred recipients in almost all cloning procedures performed during the past 20 years. We have demonstrated that the use of zygotes causes serious damage to the transferred donor nuclei, and we have concluded that there is no firm reason for accepting the claim of mouse cloning success using zygotes [78].

Enucleation removes factors beneficial for developmental potential from the zygote

Modlinski *et al.* [79] demonstrated clearly that critical genomic reprogramming factors are present inside the zygotic pronuclei rather than in the cytoplasm and that these reprogramming factors are removed upon enucleation. The authors developed an alternative method for enucleation that allows for the removal of the pronuclear envelope with the chromatin attached and leaves the reprogramming factors in the zygotic cytoplasm. Egli *et al.* [80] and Piliszek *et al.* [81] confirmed this finding with different methods, but thus far no clones have been born from adult somatic cells using this method, suggesting that reprogramming factors are present in zygotes but that their efficiency is lower than the factors in oocytes.

Ogushi *et al.* [82] reported that there is a vital factor inside the zygote nuclei that is essential for any embryo development, regardless of the integrity of the reprogramming. This suggests that if the nuclei are removed from a zygote, the essential factors are also removed and therefore the subsequent cloned embryo fails to develop even if complete reprogramming occurs. Modlinski’s group confirmed this using selective enucleation and concluded that nuclear material is essential for the cleavage division [83]. Moreover, we found that the cytoplasmic extract from enucleated oocytes promoted somatic cell reprogramming, and the success rate of producing cloned offspring increased [84]. This suggests that at least some genomic reprogramming factors are located in the cytoplasm and that a vital factor essential for full-term development is present in the nucleus.

Can we enhance the oocyte’s reprogramming capacity?

One possible explanation of the low success rate of cloning is that the reprogramming factor level of each oocyte is insufficient or not properly adapted for the receipt of a somatic cell nucleus, because it is naturally prepared only for the receipt of a haploid male gamete. If this is simply caused by an insufficiency of the reprogramming factor(s), cloning success rates might be improved by adding supplementary oocyte cytoplasm before the nuclear transfer procedure. For this reason, we constructed oocytes with volumes two to nine times greater than normal by the electrofusion or mechanical fusion of enucleated oocytes, and injected somatic cell nuclei (Fig. 2). However, none of these giant oocytes could




Fig. 2. Large oocytes and embryos generated by electrofusion. (A) One to five times normally sized oocyte. (B) Fertilized giant embryos at four days of culture *in vitro*.

support the development of mice cloned by SCNT. These results suggest that oocytes with extra cytoplasm do not have enhanced reprogramming potential [85].

ntES Cells: intrinsic problems and solution

The first success in generating ES-like cell lines by SCNT was initially performed in the bovine [86] and then subsequently in the mouse [87, 88]. We have previously shown that these ES-like cell lines are capable of differentiating into all three germ layers *in vitro* or into spermatozoa and oocytes in chimeric mice [48]. This was the first demonstration that ntES-like cells have the same potential as ES cells from fertilized blastocysts. Such ntES cell lines can be established with success rates 10 times higher than from 'conventional' reproductive cloning (Fig. 3, Somatic clone *vs.* ntES establish) [47, 48, 54, 89]. Interestingly, although almost all the cloned mice showed some abnormalities and ntES cells were established using same nuclear transfer procedure, ntES cells are transcriptionally and functionally indistinguishable from ES cells derived from fertilized embryos [47, 90]. Recently, Chang *et al.* reported that abnormal methylation profiles of certain imprinted genes could be observed in both ntES and ES cell lines after long-term culture *in vitro* [91]. We also discovered that the methylation




Fig. 3. Success rate of cloning mice and rate of establishing nuclear transfer embryonic stem (ntES) cell lines in different individuals. In this experiment, seven mice were used as donors and the cloned mice were obtained from four of them (No. 1 to 4). The ntES cell lines were established from all donor mice at the same time with a very high success rate. Using a second round of nuclear transfer with ntES cells as the donors, cloned mice were obtained from two out of the remaining three donor mice (No. 5 and 6). Only one donor mouse (No. 7) failed to produce a cloned mouse by either somatic cell nuclear transfer (SCNT) or ntES cell cloning.

status at the upstream differentially methylated region of U2af1-rs1 was changed significantly in ntES cells, but not in ES cells [92]. These results suggest that NT affects the epigenetic status of a few gene regions in common and that a change in the methylation status of U2af1-rs1 allows us to distinguish ntES cells from ES cells.

Solution of the ethical problem for human ntES cell derivation

The use of human ntES cells has raised ethical objections because fresh oocytes must be donated by healthy women and the cloned embryos would be deprived of their potential to develop into a complete human being. Several different approaches have been proposed to overcome these problems. For example, human IVF is now routinely performed in infertility clinics with high successful fertilization rates (60–70% in humans, greater than 70% in mice) [93–95], but some oocytes still fail to fertilize for unknown reasons. However, if such aged fertilization failure (AFF) oocytes could be used as recipient oocytes to generate ntES cell lines, then this would reduce or even eliminate any ethical concerns over oocyte donation and embryo destruction. In mouse experiments, the efficiency of producing cloned mice decreased with increased oocyte age [96] and the fetuses never developed to full term when AFF oocytes were used [49]. However, surprisingly, ntES cell lines could be established from AFF oocytes with the same efficiency as fresh oocytes even when stored for 24 h after IVF (Table 1), and those ntES cell lines showed the differentiation potential to be normal, the same as ES cells [49]. Therefore, the use of AFF oocytes for NT can reduce or eliminate ethical concerns regarding oocyte donation and embryo destruction.

Egli *et al.* also developed a similar method but they used enucleated polyspermic mouse zygotes, which also cannot develop

Table 1. Success rate of cloned mice and ntES cell lines from IVF failed (AFF) oocytes

	Preserved period after IVF (Temperature)	Percent of survived oocyte after NT	Percent of blastocyst	Percent of established ntES cell lines	Percent of cloned offspring
Fresh control	—	81	56	16	2
AFF oocyte	6 h (37 C)	61	46	20	0
	24 h (25 C)	27	15	29	0

because they have a triploid phenotype [80]. Alternatively, ES cell lines have been established nondestructively by removing single blastomeres or polar bodies from fertilized embryos [97, 98]. However, this can only be applied to subsequent generations, not to direct therapy for the donor.

New Attempts to Improve the Efficiency of Mouse Cloning

We previously reviewed many different approaches to improve the efficiency of mouse cloning [44] but most of those methods had no effect on the overall success rate. However, nearly ten years ago, we accidentally discovered that 1% DMSO could significantly improve the frequency of development to the blastocyst stage *in vitro* [43]. Although it is not clear why DMSO enhanced the developmental potential of cloned embryos, it was the first demonstration that nuclear reprogramming can be enhanced artificially using chemical treatment.

Epigenetic alterations by chemical treatment

As mentioned above, a recent molecular analysis of cloned embryos revealed abnormal epigenetic modifications such as aberrant DNA methylation and histone modification [61–63, 99]. Therefore, the prevention of epigenetic errors is expected to improve the success rate of animal cloning. Recently, Iwatani *et al.* discovered that DMSO treatment affects DNA methylation status at multiple loci [100]. Probably, DMSO enhances the epigenetic reprogramming of cloned embryo during *in vitro* culture. Enright *et al.* have tried to alter the epigenetic status of donor nuclei used for cow cloning by using two chemicals: 5-azacytidine, an inhibitor of DNA methylation, and trichostatin A (TSA), a histone deacetylase inhibitor (HDACi) [101]. Although the *in vitro* developmental potential with TSA was increased significantly, the cells did not demonstrate full-term development. Because those epigenetically affecting drugs are very toxic [102, 103], each drug must be tested pharmacologically for its appropriate exposure, timing, concentration, and duration.

Discovery of optimal treatment of HDACi for mouse cloning

In this situation, by trial and error, one of us (SK) discovered the optimum concentration, timing and period of TSA treatment for cloned mouse embryos. Eventually this method led to a greater than five-fold increase in the success rate of mouse cloning (Fig. 4) and a doubling in the rate of establishing ntES cell lines [45]. These results were independently and concurrently obtained by Tsunoda's laboratory [104], showing that the effects of TSA are reproducible. On the other hand, previously, most cloned mice



Fig. 4. Production of cloned mice using the histone deacetylase inhibitor (HDACi) trichostatin A (TSA). This picture shows 13 cloned mice generated in just one experiment from a single donor mouse.

have only been generated from hybrid strains and have never been cloned from outbred or inbred strains [105, 106]. Although we found that TSA could be used to produce cloned mice even from an outbred supposedly 'unclonable' strain [107], most of the important mouse strains have never been cloned successfully.

Although TSA application resulted in great improvements in SCNT cloning in mice, the effects of TSA treatment on cloning efficiency are controversial in the bovine [108, 109], pig [110, 111], rabbit [112, 113] and rat [114]. Moreover, some groups have reported that TSA treatment had detrimental effects on the *in vitro* and *in vivo* development of the SCNT embryos [109, 112]. To our knowledge, the effects of TSA treatment on full-term development have not been determined in any species other than the mouse. In experiments on rabbits, all cloned offspring treated with TSA died within 19 days after birth, whereas the untreated control clones grew to adulthood [112]. On the other hand, it is known that the drug scriptaid acts as an HDACi but is less toxic than TSA [115]. Using this drug, Zhao *et al.* could improve the success rate of pig cloning to full term [116]. At the same time, one of us (NVT) found that scriptaid treatment could increase cloned embryo development not only in hybrid but also in inbred strains, and this allowed us to generate full-term offspring from several inbred mouse strains, such as C57BL/6 and C3H/He (Fig. 5) [117]. These results suggest that although the use of HDACi drugs can enhance reprogramming in cloned embryos, because of their toxicity, the effects depend on the sensitivity of the donor strain or species.




Fig. 5. Effects of TSA and scriptaid treatment on inbred mouse cloning. Without HDACi treatment, cloned mice could be obtained from hybrid and 129/Sv strains, but with a low success rate. Only one cloned mouse was obtained from the DBA/2 strain, but this animal never reproduced. When TSA was used, the success rates for hybrid and outbred strains were increased but we have never succeeded in producing full-term cloned mice from inbred strains. However, when scriptaid was used, the overall success rate was increased even from inbred strains.

How does HDACi treatment enhance reprogramming?

Although the underlying mechanism of how HDACi treatment improves cloning efficiency remains unknown, it is thought that it can induce hyperacetylation of the core histones, resulting in structural changes in chromatin that permit transcription and enhanced DNA demethylation of the somatic cell-derived genome after SCNT [45], which is a necessary part of genetic reprogramming [118]. In fact, several reports clearly showed that HDACi treatment improved histone acetylation [111, 119], nascent mRNA production [117] and gene expression [120] in a manner similar to that in normally fertilized embryos. However, more study is required to understand the mechanisms. On the other hand, another HDACi, APHA (3-(1-methyl-4-phenylacetyl-1H-2-pyrrolyl)-N-hydroxy-2-propanamide) had no effect [117] and sirtinol had very little effect [121] for mouse cloning. In general, HDACs are divided into five categories: class I (HDAC 1, 3 and 8), class IIa (HDAC 4, 5, 7 and 9), class IIb (HDAC 6 and 10), class III (SIRT1 to 7) and class IV (HDAC 11). It is known that TSA and scriptaid can inhibit the class I and IIa/b HDAC forms [122], whereas APHA can inhibit only class I [123] and sirtinol can inhibit only class III [124]. This suggests that inhibition of the class II HDAC is very important for the success of mouse cloning.

Effects of serial cloning

We and others have demonstrated that re-cloned animals can be generated from the somatic cells of cloned animals [125–129]. However, in most reports, the overall trend of the cloning efficiency is downward: in the mouse, six successive generations is the upper limit, whereas in bovines, the second generation is the upper limit. It is unclear why subsequent cloning is not possible. At the cellular level, there was no evidence of telomere shortening in the peripheral blood lymphocytes of cloned mice [126]. One possibil-

ity is the extremely low success rate of cloning animals. If the success rate is only few percent, then cloning is very unlikely to succeed in the production of the next generation, purely by chance. However, now we have raised the average production rate of cloned animal using HDACi. Therefore, we tried serial cloning experiments again using TSA treatment, and we have obtained 15 successive generations of cloned mice without decreasing the success rate (ongoing and unpublished results). We now believe that animal cloning can be repeated indefinitely.

Combination of ntES Cell Approaches for Mouse Cloning

We proposed previously that ntES techniques could be applied to the biological sciences as a novel investigative tool. Potential investigations could be made using only this technique and not iPS technology. For example, Li *et al.* and Eggan *et al.* generated ntES cells from the nucleus of a single olfactory sensory neuron and then demonstrated that the odorant receptor gene choice is reset by SCNT and is not accompanied by genomic alterations [130, 131]. Similarly, monoclonal mice have been generated from ntES cells derived from lymphocyte nuclei indirectly using tetraploid complementation [132]. Thus, ntES cell techniques can be applied to the characterization of very rare cells in the body. Once ntES cells from these rare cell nuclei are established, the cells can proliferate indefinitely. However, here we propose another application of ntES cells, in which they can be used for reproduction from infertile or dead animals.

Producing offspring from individual mice

The genetically modified mouse is a powerful tool for research in the fields of medicine and biology but they are often infertile after genetic modification. Overcoming this infertile phenotype is a challenge worth undertaking, as the ability to maintain such types of mutant mice as genetic resources would afford numerous advantages crucial to research in human infertility and to the biology of reproduction. Unfortunately, the success rate of somatic cell cloning is very low. Even in cases in which the cloning of a sterile mouse is successful, because of their infertile phenotype it will still be necessary to clone all subsequent generations. On the other hand, the rate of establishing ntES cells is nearly 10 times higher than the success rate of producing cloned mice (Fig. 2) [48, 89]. Converting from somatic cells to ntES cells does not improve the overall success rate of cloning by a second round of nuclear transfer [48, 53, 89]. However, we recommend the establishment of ntES cell lines at the same time to preserve the donor genome, because these lines can then be used as an unlimited source of donor nuclei for NT [54].

For example, because of the low success rate in producing cloned mice, only a few individual donor mice generated clones. However, we were ultimately able to obtain cloned mice from six out of seven individual donors by using either direct somatic cells or serial nuclear transfer of ntES cells (Figs. 2, 6) [54]. On the other hand, senescent mice are often infertile, and the cloning success rate decreases with age, making it almost impossible to produce cloned progeny directly from such mice. However, we




Fig. 6. Three different approaches to generate offspring from infertile, 'unclonable' or dead frozen mice.

succeeded in establishing ntES cell lines from aged mice (nearly 3 years old) regardless of sex or strain. The cloned mice were obtained from these ntES cells by a second round of NT (Fig. 7) [53]. We also achieved success using a mutant, hermaphroditic, sterile mouse found in our ICR mouse breeding colony [52]. Although the mutant mouse died accidentally soon after tail-tip biopsy and we failed to produce cloned mice from those cells, we could establish several ntES cell lines. Using those ntES cell lines, we generated chimeric mice by injecting them into normal embryos, and finally one diploid chimeric male transmitted most of its mutant mouse genes to the next generation via the ntES cells (Fig. 6). Thus, this technique is potentially applicable for the propagation of a variety of animals or important mutant genes, regardless of age or fertile potential.

The possibility of resurrecting an extinct animal

Cloning animals by SCNT provides an opportunity to preserve endangered mammalian species. When live cells can be collected from frozen bodies, it is possible to generate cloned animals [133]. However, the 'resurrection' of extinct species from permafrost (such as the woolly mammoth) is thought to be impractical, because no live cells could be available. On the other hand, it is known that 'dead' sperm from freeze-drying treatments [134] or from a whole frozen body [135] still possess the complete haploid genome and when such sperm are injected into oocytes, the resulting embryos can develop to full-term healthy offspring. Recently, successful SCNT using freeze-dried cells was reported. Loi *et al.*




Fig. 7. Production of offspring from aged infertile mice. The outcomes for two aged mice (both 2 years and 9 months) were compared between direct SCNT cloning, the ntES cell establishment rate from those cloned blastocysts and ntES cell cloning. In both animals, direct cloning failed, probably because of the difficulty in genomic reprogramming of aged somatic cells. However, several ntES cell lines were established from both animals, and many healthy cloned mice were obtained by successive rounds of NT using the ntES cell lines.

generated cloned bovine blastocysts from freeze-dried somatic cells preserved for three years at room temperature [136]. This report was the first to demonstrate that even freeze-dried somatic




Fig. 8. Cloned mice produced from frozen dead bodies. (A) As a model of extinct animals frozen in permafrost, we used frozen dead mouse bodies as nuclear donors. These had been kept at -20°C for 16 years without any cryoprotectant. Although we could not produce cloned mice directly from this material by SCNT, ntES cell lines could be established (B). (C) Healthy cloned mice were obtained after a second round of NT using these ntES cell lines. (D) Although the rate of establishment of ntES cell lines decreased with increased storage period, these tissues frozen for 16 years retained the genetic potential to establish ntES cell lines.

Table 2. Difference between ntES cell and iPS cells

	Difficulty	Genetic modification	Ethics problem	Minimum no. donor cells	Donor cell condition
ntES cell	Difficult	No	Yes	<10 cell	Any cell, even dead
iPS cell	Easy	Yes?	No	A lot	Live cell only

cells have developmental potential after NT, but the authors could not determine whether the blastocysts were normal. On the other hand, we established ntES cell lines from freeze-dried mouse cells [55]. Importantly, after chimera construction experiments, these ntES cell lines could contribute into all organs including germ cells, which suggests that the genomic integrity of somatic cells can be maintained after freeze-drying, and that it is possible to produce offspring from such cells using SCNT.

In addition, we have attempted to produce cloned mice from bodies kept frozen at -20°C for up to 16 years without any cryoprotection (Fig. 8). These conditions are similar to those of a frozen body recovered from permafrost. Although we could not produce cloned offspring from the somatic cell directly, several ntES cell lines were established from the cell nuclei of most organ cells, irrespective of the duration of preservation. Surprisingly, frozen brain was the best tissue for generating cloned embryos, even though fresh brain tissue has proven to be one of the most difficult sources

of nuclei for the successful production of cloned mice [35, 38]. Finally, healthy cloned mice were produced from these ntES cells by a second round of NT (Fig. 8) [56]. Thus, these techniques could be used to resurrect animals or to maintain genome stocks from tissues that have been frozen for prolonged periods or even when no live cells are available.

Perspectives

The mouse is a most popular experimental animal and the advent of mouse cloning from adult-derived cells in 1998 marked a new departure in the study of key biological problems in cloning biology. Unfortunately, the success rate of SCNT cloning is still very low and the mechanisms involved in reprogramming the epigenome are not yet clear. Therefore, the SCNT method has been thought of as a 'black box approach' and inadequate to determine the detail of how genomic reprogramming occurs. However, only

the NT approach can reveal dynamic and global modifications in the epigenome without using genetic modification and can give important hints to the reprogramming mechanism. Therefore, the use of iPS cells for genomic reprogramming and for regenerative medicine is currently a 'hot topic' in this field, but we still believe that the NT approach, far from being outmoded, remains the only valid way to study biology, especially in generating offspring from a single cell or even dead cell nuclei (Table 2). Using this system, we have demonstrated that nuclear DNA is much more stable than we thought, and it looks as though we might be closer to realizing a big dream: the resurrection of extinct species by cloning. On the other hand, the abnormalities seen in cloned animals pose a major obstacle for application but this may be resolved when the mechanisms of genomic reprogramming are better understood. We believe that the mechanisms of reprogramming will be clarified when cloning efficiency is improved through technical advances.

References

- Wilmot I, Schnieke AE, McWhir J, Kind AJ, Campbell KH. Viable offspring derived from fetal and adult mammalian cells. *Nature* 1997; 385: 810–813.
- Wakayama T, Perry AC, Zuccotti M, Johnson KR, Yanagimachi R. Full-term development of mice from enucleated oocytes injected with cumulus cell nuclei. *Nature* 1998; 394: 369–374.
- Kato Y, Tani T, Sotomaru Y, Kurokawa K, Kato J, Doguchi H, Yasue H, Tsunoda Y. Eight calves cloned from somatic cells of a single adult. *Science* 1998; 282: 2095–2098.
- Baguisi A, Behboodi E, Melican DT, Pollock JS, Destrempes MM, Cammuso C, Williams JL, Nims SD, Porter CA, Midura P, Palacios MJ, Ayres SL, Denniston RS, Hayes ML, Ziomek CA, Meade HM, Godke RA, Gavin WG, Overstrom EW, Echelard Y. Production of goats by somatic cell nuclear transfer. *Nat Biotechnol* 1999; 17: 456–461.
- Onishi A, Iwamoto M, Akita T, Mikawa S, Takeda K, Awata T, Hanada H, Perry AC. Pig cloning by microinjection of fetal fibroblast nuclei. *Science* 2000; 289: 1188–1190.
- Polejaeva IA, Chen SH, Vaught TD, Page RL, Mullins J, Ball S, Dai Y, Boone J, Walker S, Ayares DL, Colman A, Campbell KH. Cloned pigs produced by nuclear transfer from adult somatic cells. *Nature* 2000; 407: 86–90.
- Lanza RP, Cibelli JB, Diaz F, Moraes CT, Farin PW, Farin CE, Hammer CJ, West MD, Damiani P. Cloning of an endangered species (*Bos gaurus*) using interspecies nuclear transfer. *Cloning* 2000; 2: 79–90.
- Loi P, Ptak G, Barboni B, Fulka J, Jr., Cappai P, Clinton M. Genetic rescue of an endangered mammal by cross-species nuclear transfer using post-mortem somatic cells. *Nat Biotechnol* 2001; 19: 962–964.
- Shin T, Kraemer D, Pryor J, Liu L, Rugila J, Howe L, Buck S, Murphy K, Lyons L, Westhusin M. A cat cloned by nuclear transplantation. *Nature* 2002; 415: 859.
- Chesne P, Adenot PG, Viglietta C, Baratte M, Boulanger L, Renard JP. Cloned rabbits produced by nuclear transfer from adult somatic cells. *Nat Biotechnol* 2002; 20: 366–369.
- Galli C, Lagutina I, Crotti G, Colleoni S, Turini P, Ponderato N, Duchi R, Lazzari G. Pregnancy: a cloned horse born to its dam twin. *Nature* 2003; 424: 635.
- Woods GL, White KL, Vanderwall DK, Li GP, Aston KI, Bunch TD, Meerdo LN, Pate BJ. A mule cloned from fetal cells by nuclear transfer. *Science* 2003; 301: 1063.
- Zhou Q, Renard JP, Le Fric G, Brochard V, Beaujean N, Cherifi Y, Fraichard A, Cozzi J. Generation of fertile cloned rats by regulating oocyte activation. *Science* 2003; 302: 1179.
- Gomez MC, Pope CE, Giraldo A, Lyons LA, Harris RF, King AL, Cole A, Godke RA, Dresser BL. Birth of African Wildcat cloned kittens born from domestic cats. *Cloning Stem Cells* 2004; 6: 247–258.
- Lee BC, Kim MK, Jang G, Oh HJ, Yuda F, Kim HJ, Hossein MS, Kim JJ, Kang SK, Schatten G, Hwang WS. Dogs cloned from adult somatic cells. *Nature* 2005; 436: 641.
- Li Z, Sun X, Chen J, Liu X, Wisely SM, Zhou Q, Renard JP, Leno GH, Engelhardt JF. Cloned ferrets produced by somatic cell nuclear transfer. *Dev Biol* 2006; 293: 439–448.
- Kim MK, Jang G, Oh HJ, Yuda F, Kim HJ, Hwang WS, Hossein MS, Kim JJ, Shin NS, Kang SK, Lee BC. Endangered wolves cloned from adult somatic cells. *Cloning Stem Cells* 2007; 9: 130–137.
- Berg DK, Li C, Asher G, Wells DN, Oback B. Red deer cloned from antler stem cells and their differentiated progeny. *Biol Reprod* 2007; 77: 384–394.
- Gomez MC, Pope CE, Kutner RH, Ricks DM, Lyons LA, Ruhe M, Dumas C, Lyons J, Lopez M, Dresser BL, Reiser J. Nuclear transfer of sand cat cells into enucleated domestic cat oocytes is affected by cryopreservation of donor cells. *Cloning Stem Cells* 2008; 10: 469–483.
- Folch J, Cocero MJ, Chesne P, Alabart JL, Dominguez V, Cogne Y, Roche A, Fernandez-Arias A, Marti JJ, Sanchez P, Echegoyen E, Beckers JF, Bonastre AS, Vignon X. First birth of an animal from an extinct subspecies (*Capra pyrenaica pyrenaica*) by cloning. *Theriogenology* 2009; 71: 1026–1034.
- Wani NA, Wernery U, Hassan FA, Wernery R, Skidmore JA. Production of the first cloned camel by somatic cell nuclear transfer. *Biol Reprod* 2009.
- Boiani M, Eckardt S, Scholer HR, McLaughlin KJ. Oct4 distribution and level in mouse clones: consequences for pluripotency. *Genes Dev* 2002; 16: 1209–1219.
- Bortvin A, Eggan K, Skaletsky H, Akutsu H, Berry DL, Yanagimachi R, Page DC, Jaenisch R. Incomplete reactivation of Oct4-related genes in mouse embryos cloned from somatic nuclei. *Development* 2003; 130: 1673–1680.
- Kishigami S, Hikichi T, Van Thuan N, Ohta H, Wakayama S, Bui HT, Mizutani E, Wakayama T. Normal specification of the extraembryonic lineage after somatic nuclear transfer. *FEBS Lett* 2006; 580: 1801–1806.
- Wakayama T, Yanagimachi R. Cloning of male mice from adult tail-tip cells. *Nat Genet* 1999; 22: 127–128.
- Tanaka S, Oda M, Toyoshima Y, Wakayama T, Tanaka M, Yoshida N, Hattori N, Ohgane J, Yanagimachi R, Shiota K. Placentomegaly in cloned mouse concepti caused by expansion of the spongiotrophoblast layer. *Biol Reprod* 2001; 65: 1813–1821.
- Tamashiro KL, Wakayama T, Akutsu H, Yamazaki Y, Lachey JL, Wortman MD, Seeley RJ, D'Alessio DA, Woods SC, Yanagimachi R, Sakai RR. Cloned mice have an obese phenotype not transmitted to their offspring. *Nat Med* 2002; 8: 262–267.
- Tamashiro KL, Wakayama T, Blanchard RJ, Blanchard DC, Yanagimachi R. Postnatal growth and behavioral development of mice cloned from adult cumulus cells. *Biol Reprod* 2000; 63: 328–334.
- Ogonuki N, Inoue K, Yamamoto Y, Noguchi Y, Tanemura K, Suzuki O, Nakayama H, Doi K, Ohtomo Y, Satoh M, Nishida A, Ogura A. Early death of mice cloned from somatic cells. *Nat Genet* 2002; 30: 253–254.
- Ogura A, Inoue K, Takano K, Wakayama T, Yanagimachi R. Birth of mice after nuclear transfer by electrofusion using tail tip cells. *Mol Reprod Dev* 2000; 57: 55–59.
- Ogura A, Inoue K, Ogonuki N, Noguchi A, Takano K, Nagano R, Suzuki O, Lee J, Ishino F, Matsuda J. Production of male cloned mice from fresh, cultured, and cryopreserved immature Sertoli cells. *Biol Reprod* 2000; 62: 1579–1584.
- Wakayama T, Rodriguez I, Perry AC, Yanagimachi R, Mombaerts P. Mice cloned from embryonic stem cells. *Proc Natl Acad Sci USA* 1999; 96: 14984–14989.
- Inoue K, Wakao H, Ogonuki N, Miki H, Seino K, Nambu-Wakao R, Noda S, Miyoshi H, Koseki H, Taniguchi M, Ogura A. Generation of cloned mice by direct nuclear transfer from natural killer T cells. *Curr Biol* 2005; 15: 1114–1118.
- Sung LY, Gao S, Shen H, Yu H, Song Y, Smith SL, Chang CC, Inoue K, Kuo L, Lian J, Li A, Tian XC, Tuck DP, Weissman SM, Yang X, Cheng T. Differentiated cells are more efficient than adult stem cells for cloning by somatic cell nuclear transfer. *Nat Genet* 2006.
- Yamazaki Y, Makino H, Hamaguchi-Hamada K, Hamada S, Sugino H, Kawase E, Miyata T, Ogawa M, Yanagimachi R, Yagi T. Assessment of the developmental totipotency of neural cells in the cerebral cortex of mouse embryo by nuclear transfer. *Proc Natl Acad Sci USA* 2001; 98: 14022–14026.
- Inoue K, Ogonuki N, Miki H, Hirose M, Noda S, Kim JM, Aoki F, Miyoshi H, Ogura A. Inefficient reprogramming of the hematopoietic stem cell genome following nuclear transfer. *J Cell Sci* 2006; 119: 1985–1991.
- Inoue K, Noda S, Ogonuki N, Miki H, Inoue S, Katayama K, Mekada K, Miyoshi H, Ogura A. Differential developmental ability of embryos cloned from tissue-specific stem cells. *Stem Cells* 2007; 25: 1279–1285.
- Mizutani E, Ohta H, Kishigami S, Van Thuan N, Hikichi T, Wakayama S, Kosaka M, Sato E, Wakayama T. Developmental ability of cloned embryos from neural stem cells. *Reproduction* 2006; 132: 849–857.
- Li J, Greco V, Guasch G, Fuchs E, Mombaerts P. Mice cloned from skin cells. *Proc Natl Acad Sci USA* 2007; 104: 2738–2743.
- Miki H, Inoue K, Kohda T, Honda A, Ogonuki N, Yuzuriha M, Mise N, Matsui Y, Baba T, Abe K, Ishino F, Ogura A. Birth of mice produced by germ cell nuclear transfer. *Genesis* 2005; 41: 81–86.
- Kishikawa H, Wakayama T, Yanagimachi R. Comparison of oocyte-activating agents for mouse cloning. *Cloning* 1999; 1: 153–159.
- Wakayama S, Cibelli JB, Wakayama T. Effect of timing of the removal of oocyte chromosomes before or after injection of somatic nucleus on development of NT embryos. *Cloning Stem Cells* 2003; 5: 181–189.
- Wakayama T, Yanagimachi R. Effect of cytokinesis inhibitors, DMSO and the timing of oocyte activation on mouse cloning using cumulus cell nuclei. *Reproduction* 2001; 122: 49–60.

44. Wakayama T. Production of cloned mice and ES cells from adult somatic cells by nuclear transfer: how to improve cloning efficiency? *J Reprod Dev* 2007; 53: 13–26.
45. Kishigami S, Mizutani E, Ohta H, Hikichi T, Thuan NV, Wakayama S, Bui HT, Wakayama T. Significant improvement of mouse cloning technique by treatment with trichostatin A after somatic nuclear transfer. *Biochem Biophys Res Commun* 2006; 340: 183–189.
46. Svensson K, Mattsson R, James TC, Wentzel P, Pilartz M, MacLaughlin J, Miller SJ, Olsson T, Eriksson UJ, Ohlsson R. The paternal allele of the H19 gene is progressively silenced during early mouse development: the acetylation status of histones may be involved in the generation of variegated expression patterns. *Development* 1998; 125: 61–69.
47. Wakayama S, Jakt ML, Suzuki M, Araki R, Hikichi T, Kishigami S, Ohta H, Van Thuan N, Mizutani E, Sakaide Y, Senda S, Tanaka S, Okada M, Miyake M, Abe M, Nishikawa S, Shiota K, Wakayama T. Equivalency of nuclear transfer-derived embryonic stem cells to those derived from fertilized mouse blastocysts. *Stem Cells* 2006; 24: 2023–2033.
48. Wakayama T, Tabar V, Rodriguez I, Perry AC, Studer L, Mombaerts P. Differentiation of embryonic stem cell lines generated from adult somatic cells by nuclear transfer. *Science* 2001; 292: 740–743.
49. Wakayama S, Suetsugu R, Thuan NV, Ohta H, Kishigami S, Wakayama T. Establishment of mouse embryonic stem cell lines from somatic cell nuclei by nuclear transfer into aged, fertilization-failure mouse oocytes. *Curr Biol* 2007; 17: R120–121.
50. Rideout WM, 3rd, Hochedlinger K, Kyba M, Daley GQ, Jaenisch R. Correction of a genetic defect by nuclear transplantation and combined cell and gene therapy. *Cell* 2002; 109: 17–27.
51. Tabar V, Tomishima M, Panagiotakos G, Wakayama S, Menon J, Chan B, Mizutani E, Al-Shamy G, Ohta H, Wakayama T, Studer L. Therapeutic cloning in individual parkinsonian mice. *Nat Med* 2008; 14: 379–381.
52. Wakayama S, Kishigami S, Van Thuan N, Ohta H, Hikichi T, Mizutani E, Yanagimachi R, Wakayama T. Propagation of an infertile hermaphrodite mouse lacking germ cells by using nuclear transfer and embryonic stem cell technology. *Proc Natl Acad Sci USA* 2005; 102: 29–33.
53. Mizutani E, Ono T, Li C, Maki-Suetsugu R, Wakayama T. Propagation of senescent mice using nuclear transfer embryonic stem cell lines. *Genesis* 2008; 46: 478–483.
54. Wakayama S, Mizutani E, Kishigami S, Thuan NV, Ohta H, Hikichi T, Bui HT, Miyake M, Wakayama T. Mice cloned by nuclear transfer from somatic and nES cells derived from the same individuals. *J Reprod Dev* 2005; 51: 765–772.
55. Ono T, Mizutani E, Li C, Wakayama T. Nuclear transfer preserves the nuclear genome of freeze-dried mouse cells. *J Reprod Dev* 2008; 54: 486–491.
56. Wakayama S, Ohta H, Hikichi T, Mizutani E, Iwaki T, Kanagawa O, Wakayama T. Production of healthy cloned mice from bodies frozen at –20 degrees C for 16 years. *Proc Natl Acad Sci USA* 2008; 105: 17318–17322.
57. Wakayama T, Yanagimachi R. Cloning the laboratory mouse. *Semin Cell Dev Biol* 1999; 10: 253–258.
58. Inoue K, Kohda T, Lee J, Ogonuki N, Mochida K, Noguchi Y, Tanemura K, Kaneko-Ishino T, Ishino F, Ogura A. Faithful expression of imprinted genes in cloned mice. *Science* 2002; 295: 297.
59. Oda M, Tanaka S, Yamazaki Y, Ohta H, Iwatani M, Suzuki M, Ohgane J, Hattori N, Yanagimachi R, Wakayama T, Shiota K. Establishment of trophoblast stem cell lines from somatic cell nuclear-transferred embryos. *Proc Natl Acad Sci USA* 2009; 106: 16293–16297.
60. Miki H, Wakisaka N, Inoue K, Ogonuki N, Mori M, Kim JM, Ohta A, Ogura A. Embryonic rather than extraembryonic tissues have more impact on the development of placental hyperplasia in cloned mice. *Placenta* 2009; 30: 543–546.
61. Dean W, Santos F, Stojkovic M, Zakhartchenko V, Walter J, Wolf E, Reik W. Conservation of methylation reprogramming in mammalian development: aberrant reprogramming in cloned embryos. *Proc Natl Acad Sci USA* 2001; 98: 13734–13738.
62. Kang YK, Koo DB, Park JS, Choi YH, Chung AS, Lee KK, Han YM. Aberrant methylation of donor genome in cloned bovine embryos. *Nat Genet* 2001; 28: 173–177.
63. Ohgane J, Wakayama T, Kogo Y, Senda S, Hattori N, Tanaka S, Yanagimachi R, Shiota K. DNA methylation variation in cloned mice. *Genesis* 2001; 30: 45–50.
64. Renard JP, Chastant S, Chesne P, Richard C, Marchal J, Cordonnier N, Chavatte P, Vignon X. Lymphoid hypoplasia and somatic cloning. *Lancet* 1999; 353: 1489–1491.
65. Humpherys D, Eggan K, Akutsu H, Friedman A, Hochedlinger K, Yanagimachi R, Lander ES, Golub TR, Jaenisch R. Abnormal gene expression in cloned mice derived from embryonic stem cell and cumulus cell nuclei. *Proc Natl Acad Sci USA* 2002; 99: 12889–12894.
66. Kohda T, Inoue K, Ogonuki N, Miki H, Naruse M, Kaneko-Ishino T, Ogura A, Ishino F. Variation in gene expression and aberrantly regulated chromosome regions in cloned mice. *Biol Reprod* 2005; 73: 1302–1311.
67. Humpherys D, Eggan K, Akutsu H, Hochedlinger K, Rideout WM, 3rd, Biniszkiwicz D, Yanagimachi R, Jaenisch R. Epigenetic instability in ES cells and cloned mice. *Science* 2001; 293: 95–97.
68. Senda S, Wakayama T, Yamazaki Y, Ohgane J, Hattori N, Tanaka S, Yanagimachi R, Shiota K. Skewed X-inactivation in cloned mice. *Biochem Biophys Res Commun* 2004; 321: 38–44.
69. Senda S, Wakayama T, Arai Y, Yamazaki Y, Ohgane J, Tanaka S, Hattori N, Yanagimachi R, Shiota K. DNA methylation errors in cloned mice disappear with advancement of aging. *Cloning Stem Cells* 2007; 9: 293–302.
70. Shimozawa N, Ono Y, Kimoto S, Hioki K, Araki Y, Shinkai Y, Kono T, Ito M. Abnormalities in cloned mice are not transmitted to the progeny. *Genesis* 2002; 34: 203–207.
71. Ohta H, Wakayama T. Generation of normal progeny by intracytoplasmic sperm injection following grafting of testicular tissue from cloned mice that died postnatally. *Biol Reprod* 2005; 73: 390–395.
72. Kawasumi M, Anzai M, Takehara T, Mitani T, Kato H, Saeki K, Iritani A, Matsu-moto K, Hosoi Y. Abnormal distribution of chromosomes in the first division of nuclear transferred mouse embryos. *J Reprod Dev* 2007; 53: 615–622.
73. Balbach ST, Jauch A, Bohm-Steuer B, Cavaleri FM, Han YM, Boiani M. Chromosome stability differs in cloned mouse embryos and derivative ES cells. *Dev Biol* 2007; 308: 309–321.
74. Inoue K, Ogonuki N, Mekada K, Yoshiki A, Sado T, Ogura A. Sex-reversed somatic cell cloning in the mouse. *J Reprod Dev* 2009; 55: 566–569.
75. Yamagata K, Suetsugu R, Wakayama T. Long-term, six-dimensional live-cell imaging for the mouse preimplantation embryo that does not affect full-term development. *J Reprod Dev* 2009; 55: 343–350.
76. Illmensee K, Hoppe PC. Nuclear transplantation in *Mus musculus*: developmental potential of nuclei from preimplantation embryos. *Cell* 1981; 23: 9–18.
77. McGrath J, Solter D. Nuclear transplantation in the mouse embryo by microsurgery and cell fusion. *Science* 1983; 220: 1300–1302.
78. Wakayama T, Tateno H, Mombaerts P, Yanagimachi R. Nuclear transfer into mouse zygotes. *Nat Genet* 2000; 24: 108–109.
79. Greda P, Karasiewicz J, Modlinski JA. Mouse zygotes as recipients in embryo cloning. *Reproduction* 2006; 132: 741–748.
80. Egli D, Rosains J, Birkhoff G, Eggan K. Developmental reprogramming after chromosome transfer into mitotic mouse zygotes. *Nature* 2007; 447: 679–685.
81. Piliszek A, Modlinski JA, Pysniak K, Karasiewicz J. Foetal fibroblasts introduced to cleaving mouse embryos contribute to full-term development. *Reproduction* 2007; 133: 207–218.
82. Ogushi S, Palmieri C, Fulka H, Saitou M, Miyano T, Fulka J, Jr. The maternal nucleolus is essential for early embryonic development in mammals. *Science* 2008; 319: 613–616.
83. Mohammed AA, Karasiewicz J, Modlinski JA. Developmental potential of selectively enucleated immature mouse oocytes upon nuclear transfer. *Mol Reprod Dev* 2008; 75: 1269–1280.
84. Bui HT, Wakayama S, Kishigami S, Kim JH, Van Thuan N, Wakayama T. The cytoplasm of mouse germinal vesicle stage oocytes can enhance somatic cell nuclear reprogramming. *Development* 2008; 135: 3935–3945.
85. Wakayama S, Kishigami S, Van Thuan N, Ohta H, Hikichi T, Mizutani E, Thuy BH, Miyake M, Wakayama T. Effect of volume of oocyte cytoplasm on embryo development after parthenogenetic activation, intracytoplasmic sperm injection, or somatic cell nuclear transfer. *Zygote* 2008; 16: 211–222.
86. Cibelli JB, Stice SL, Golueke PJ, Kane JJ, Jerry J, Blackwell C, Ponce de Leon FA, Robl JM. Transgenic bovine chimeric offspring produced from somatic cell-derived stem-like cells. *Nat Biotechnol* 1998; 16: 642–646.
87. Kawase E, Yamazaki Y, Yagi T, Yanagimachi R, Pedersen RA. Mouse embryonic stem (ES) cell lines established from neuronal cell-derived cloned blastocysts. *Genesis* 2000; 28: 156–163.
88. Munsie MJ, Michalska AE, O'Brien CM, Trounson AO, Pera MF, Mountford PS. Isolation of pluripotent embryonic stem cells from reprogrammed adult mouse somatic cell nuclei. *Curr Biol* 2000; 10: 989–992.
89. Wakayama S, Ohta H, Kishigami S, Thuan NV, Hikichi T, Mizutani E, Miyake M, Wakayama T. Establishment of male and female nuclear transfer embryonic stem cell lines from different mouse strains and tissues. *Biol Reprod* 2005; 72: 932–936.
90. Brambrink T, Hochedlinger K, Bell G, Jaenisch R. ES cells derived from cloned and fertilized blastocysts are transcriptionally and functionally indistinguishable. *Proc Natl Acad Sci USA* 2006; 103: 933–938.
91. Chang G, Liu S, Wang F, Zhang Y, Kou Z, Chen D, Gao S. Differential methylation status of imprinted genes in nuclear transfer derived ES (NT-ES) cells. *Genomics* 2009; 93: 112–119.
92. Hikichi T, Kohda T, Wakayama S, Ishino F, Wakayama T. Nuclear transfer alters the DNA methylation status of specific genes in fertilized and parthenogenetically activated mouse embryonic stem cells. *Stem Cells* 2008; 26: 783–788.
93. Chen HL, Copperman AB, Grunfeld L, Sandler B, Bustillo M, Gordon JW. Failed

- fertilization *in vitro*: second day micromanipulation of oocytes versus reinsemination. *Fertil Steril* 1995; 63: 1337–1340.
94. Nagy ZP, Joris H, Liu J, Staessen C, Devroey P, Van Steirteghem AC. Intracytoplasmic single sperm injection of 1-day-old unfertilized human oocytes. *Hum Reprod* 1993; 8: 2180–2184.
 95. Toyoda Y, Yokoyama M, Hoshi T. Studies on the fertilization of mouse eggs *in vitro*. *Jpn J Anim Reprod* 1971; 16: 152–157.
 96. Liu G, Kato Y, Tsunoda Y. Aging of recipient oocytes reduces the development of cloned embryos receiving cumulus cells. *J Reprod Dev* 2007; 53: 785–790.
 97. Chung Y, Klimanskaya I, Becker S, Marh J, Lu SJ, Johnson J, Meisner L, Lanza R. Embryonic and extraembryonic stem cell lines derived from single mouse blastomeres. *Nature* 2006; 439: 216–219.
 98. Wakayama S, Hikichi T, Suetsugu R, Sakaide Y, Bui HT, Mizutani E, Wakayama T. Efficient establishment of mouse embryonic stem cell lines from single blastomeres and polar bodies. *Stem Cells* 2007; 25: 986–993.
 99. Santos F, Zakhartchenko V, Stojkovic M, Peters A, Jenuwein T, Wolf E, Reik W, Dean W. Epigenetic marking correlates with developmental potential in cloned bovine preimplantation embryos. *Curr Biol* 2003; 13: 1116–1121.
 100. Iwatani M, Ikegami K, Kremenska Y, Hattori N, Tanaka S, Yagi S, Shiota K. Dimethyl Sulfoxide (DMSO) increases expression of Dnmt3as and affects genome-wide DNA methylation profiles in mouse embryoid body. *Stem Cells* 2006.
 101. Enright BP, Kubota C, Yang X, Tian XC. Epigenetic characteristics and development of embryos cloned from donor cells treated by trichostatin A or 5-aza-2'-deoxycytidine. *Biol Reprod* 2003; 69: 896–901.
 102. Kishigami S, Ohta H, Mizutani E, Wakayama S, Bui HT, Thuan NV, Hikichi T, Suetsugu R, Wakayama T. Harmful or not: Trichostatin A treatment of embryos generated by ICSI or ROSI. *Central European Journal of Biology* 2006; 1: 376–385.
 103. Gurvich N, Berman MG, Wittner BS, Gentleman RC, Klein PS, Green JB. Association of valproate-induced teratogenesis with histone deacetylase inhibition *in vivo*. *Faseb J* 2005; 19: 1166–1168.
 104. Rybouchkin A, Kato Y, Tsunoda Y. Role of histone acetylation in reprogramming of somatic nuclei following nuclear transfer. *Biol Reprod* 2006; 74: 1083–1089.
 105. Inoue K, Ogonuki N, Mochida K, Yamamoto Y, Takano K, Kohda T, Ishino F, Ogura A. Effects of donor cell type and genotype on the efficiency of mouse somatic cell cloning. *Biol Reprod* 2003; 69: 1394–1400.
 106. Wakayama T, Yanagimachi R. Mouse cloning with nucleus donor cells of different age and type. *Mol Reprod Dev* 2001; 58: 376–383.
 107. Kishigami S, Bui HT, Wakayama S, Tokunaga K, Van Thuan N, Hikichi T, Mizutani E, Ohta H, Suetsugu R, Sata T, Wakayama T. Successful mouse cloning of an outbred strain by trichostatin A treatment after somatic nuclear transfer. *J Reprod Dev* 2007; 53: 165–170.
 108. Iager AE, Ragina NP, Ross PJ, Beyhan Z, Cunniff K, Rodriguez RM, Cibelli JB. Trichostatin A improves histone acetylation in bovine somatic cell nuclear transfer early embryos. *Cloning Stem Cells* 2008; 10: 371–379.
 109. Wu X, Li Y, Li GP, Yang D, Yue Y, Wang L, Li K, Xin P, Bou S, Yu H. Trichostatin A improved epigenetic modifications of transfected cells but did not improve subsequent cloned embryo development. *Anim Biotechnol* 2008; 19: 211–224.
 110. Li J, Svarcova O, Villemoes K, Kragh PM, Schmidt M, Bogh IB, Zhang Y, Du Y, Lin L, Purup S, Xue Q, Bolund L, Yang H, Maddox-Hyttel P, Vajta G. High *in vitro* development after somatic cell nuclear transfer and trichostatin A treatment of reconstructed porcine embryos. *Theriogenology* 2008; 70: 800–808.
 111. Yamanaka KI, Sugimura S, Wakai T, Kawahara M, Sato E. Acetylation level of histone H3 in early embryonic stages affects subsequent development of miniature pig somatic cell nuclear transfer embryos. *J Reprod Dev* 2009.
 112. Meng Q, Polgar Z, Liu J, Dinnyes A. Live birth of somatic cell-cloned rabbits following trichostatin A treatment and cotransfer of parthenogenetic embryos. *Cloning Stem Cells* 2009; 11: 203–208.
 113. Shi LH, Miao YL, Ouyang YC, Huang JC, Lei ZL, Yang JW, Han ZM, Song XF, Sun QY, Chen DY. Trichostatin A (TSA) improves the development of rabbit-rabbit intraspecies cloned embryos, but not rabbit-human interspecies cloned embryos. *Dev Dyn* 2008; 237: 640–648.
 114. Sterthaus O, Skoczylas E, De Geyter C, Burki K, Ledermann B. Evaluation of *in vitro* cultured rat oocytes, from different strains, by spindle morphology and maturation-promoting-factor activity combined with nuclear-transfer experiments. *Cloning Stem Cells* 2009; 11: 463–472.
 115. Su GH, Sohn TA, Ryu B, Kern SE. A novel histone deacetylase inhibitor identified by high-throughput transcriptional screening of a compound library. *Cancer Res* 2000; 60: 3137–3142.
 116. Zhao J, Ross JW, Hao Y, Spate LD, Walters EM, Samuel MS, Rieke A, Murphy CN, Prather RS. Significant improvement in cloning efficiency of an inbred miniature pig by histone deacetylase inhibitor treatment after somatic cell nuclear transfer. *Biol Reprod* 2009; 81: 525–530.
 117. Van Thuan N, Bui HT, Kim JH, Hikichi T, Wakayama S, Kishigami S, Mizutani E, Wakayama T. The histone deacetylase inhibitor scriptaid enhances nascent mRNA production and rescues full-term development in cloned inbred mice. *Reproduction* 2009; 138: 309–317.
 118. Simonsson S, Gurdon J. DNA demethylation is necessary for the epigenetic reprogramming of somatic cell nuclei. *Nat Cell Biol* 2004; 6: 984–990.
 119. Wang F, Kou Z, Zhang Y, Gao S. Dynamic reprogramming of histone acetylation and methylation in the first cell cycle of cloned mouse embryos. *Biol Reprod* 2007; 77: 1007–1016.
 120. Tsuji Y, Kato Y, Tsunoda Y. The developmental potential of mouse somatic cell nuclear-transferred oocytes treated with trichostatin A and 5-aza-2'-deoxycytidine. *Zygote* 2009; 17: 109–115.
 121. Hirata S, Fukasawa H, Tagaya H, Shoda T, Wakayama T, Hoshi K. Effects of Sirtinol on early development of the cloned murine embryos. *Yamanashi Med J* 2008; 23: 97–107.
 122. Chuang DM, Leng Y, Marinova Z, Kim HJ, Chiu CT. Multiple roles of HDAC inhibition in neurodegenerative conditions. *Trends Neurosci* 2009; 32: 591–601.
 123. Mai A, Massa S, Ragno R, Cerbara I, Jesacher F, Loidl P, Brosch G. 3-(4-Aroyl-1-methyl-1H-2-pyrrolyl)-N-hydroxy-2-alkylamides as a new class of synthetic histone deacetylase inhibitors. 1. Design, synthesis, biological evaluation, and binding mode studies performed through three different docking procedures. *J Med Chem* 2003; 46: 512–524.
 124. Denu JM. The Sir 2 family of protein deacetylases. *Curr Opin Chem Biol* 2005; 9: 431–440.
 125. Kubota C, Tian XC, Yang X. Serial bull cloning by somatic cell nuclear transfer. *Nat Biotechnol* 2004; 22: 693–694.
 126. Wakayama T, Shinkai Y, Tamashiro KL, Niida H, Blanchard DC, Blanchard RJ, Ogura A, Tanemura K, Tachibana M, Perry AC, Colgan DF, Mombaerts P, Yanagimachi R. Cloning of mice to six generations. *Nature* 2000; 407: 318–319.
 127. Cho SK, Kim JH, Park JY, Choi YJ, Bang JI, Hwang KC, Cho EJ, Sohn SH, Uhm SJ, Koo DB, Lee KK, Kim T, Kim JH. Serial cloning of pigs by somatic cell nuclear transfer: restoration of phenotypic normality during serial cloning. *Dev Dyn* 2007; 236: 3369–3382.
 128. Kurome M, Hisatomi H, Matsumoto S, Tomii R, Ueno S, Hiruma K, Saito H, Nakamura K, Okumura K, Matsumoto M, Kaji Y, Endo F, Nagashima H. Production efficiency and telomere length of the cloned pigs following serial somatic cell nuclear transfer. *J Reprod Dev* 2008; 54: 254–258.
 129. Yin XJ, Lee HS, Yu XF, Kim LH, Shin HD, Cho SJ, Choi EG, Kong IK. Production of second-generation cloned cats by somatic cell nuclear transfer. *Theriogenology* 2008; 69: 1001–1006.
 130. Eggan K, Baldwin K, Tackett M, Osborne J, Gogos J, Chess A, Axel R, Jaenisch R. Mice cloned from olfactory sensory neurons. *Nature* 2004; 428: 44–49.
 131. Li J, Ishii T, Feinstein P, Mombaerts P. Odorant receptor gene choice is reset by nuclear transfer from mouse olfactory sensory neurons. *Nature* 2004; 428: 393–399.
 132. Hochedlinger K, Jaenisch R. Monoclonal mice generated by nuclear transfer from mature B and T donor cells. *Nature* 2002; 415: 1035–1038.
 133. Hoshino Y, Hayashi N, Taniguchi S, Kobayashi N, Sakai K, Otani T, Iritani A, Saeki K. Resurrection of a bull by cloning from organs frozen without cryoprotectant in a –80 degrees c freezer for a decade. *PLoS One* 2009; 4: e4142.
 134. Wakayama T, Yanagimachi R. Development of normal mice from oocytes injected with freeze-dried spermatozoa. *Nat Biotechnol* 1998; 16: 639–641.
 135. Ogonuki N, Mochida K, Miki H, Inoue K, Fray M, Iwaki T, Moriwaki K, Obata Y, Morozumi K, Yanagimachi R, Ogura A. Spermatozoa and spermatids retrieved from frozen reproductive organs or frozen whole bodies of male mice can produce normal offspring. *Proc Natl Acad Sci USA* 2006; 103: 13098–13103.
 136. Loi P, Matsukawa K, Ptak G, Clinton M, Fulka J, Jr., Nathan Y, Arav A. Freeze-dried somatic cells direct embryonic development after nuclear transfer. *PLoS One* 2008; 3: e2978.