

**How well do we need to measure the Higgs boson mass and self-coupling?**Rick S. Gupta,<sup>1</sup> Heidi Rzehak,<sup>2,\*</sup> and James D. Wells<sup>2,3</sup><sup>1</sup>*IFAE, Universidad Autònoma de Barcelona, 08193 Bellaterra, Spain*<sup>2</sup>*CERN Theoretical Physics, CH-1211 Geneva 23, Switzerland*<sup>3</sup>*Physics Department, University of Michigan, Ann Arbor, Michigan 48109, USA*

(Received 4 July 2013; published 27 September 2013)

Much of the discussion regarding future measurements of the Higgs boson mass and self-coupling is focused on how well various collider options can do. In this article we ask a physics-based question of how well do we need colliders to measure these quantities to have an impact on discovery of new physics or an impact in how we understand the role of the Higgs boson in nature. We address the question within the framework of the Standard Model and various beyond the Standard Model scenarios, including supersymmetry and theories of composite Higgs bosons. We conclude that the LHC's stated ability to measure the Higgs boson to better than 150 MeV will be as good as we will ever need to know the Higgs boson mass in the foreseeable future. On the other hand, we estimate that the self-coupling will likely need to be measured to better than 20% to see a deviation from the Standard Model expectation. This is a challenging target for future collider and upgrade scenarios.

DOI: [10.1103/PhysRevD.88.055024](https://doi.org/10.1103/PhysRevD.88.055024)

PACS numbers: 12.60.Jv, 12.60.-i, 12.60.Cn, 14.65.Ha

**I. INTRODUCTION**

The ATLAS and CMS experiments have recently announced the discovery of a boson consistent with the Higgs boson having a mass around 126 GeV [1]. The defining property of the Higgs field is that it acquires a vacuum expectation value and thus leads to electroweak symmetry breaking (EWSB). In the Standard Model (SM), for instance, the Higgs potential

$$V = m_{\Phi_{\text{SM}}}^2 |\Phi_{\text{SM}}|^2 + \lambda_{\text{SM}} |\Phi_{\text{SM}}|^4 \quad (1)$$

has a minimum at  $v \neq 0$ . Here, and in the rest of this paper,  $\Phi_{\text{SM}}$  is the SM Higgs doublet. One way to directly check that the discovered boson is indeed a component of the Higgs field responsible for electroweak symmetry is to check the relationship

$$m_h^2 = 2\lambda_{\text{SM}} v^2, \quad (2)$$

which is obtained by minimizing the above potential and finding the physical Higgs boson ( $h$ ) mass at the minimum. Here  $v \simeq 246$  GeV. The above relationship can be experimentally verified by independently measuring the left- and right-hand sides. Whereas the Higgs mass on the left-hand side can be measured by resonant Higgs boson production, the self-coupling on the right-hand side can be measured by its contribution to the double Higgs production process [2]:  $gg \rightarrow h^* \rightarrow hh$ .

The relationship in Eq. (2) is of course true only in the SM. In theories beyond the SM (BSM) there can be deviations of the self-coupling from its SM value  $m_h^2/(2v^2)$ . This can happen, for instance, because of the existence of more than one scalar (as in the case of theories with mixed-in

singlets and supersymmetric models) or because of the presence of effective higher-dimension operators in the Higgs potential (as in composite Higgs models and theories with first-order electroweak phase transition). How accurately do we need to measure the left- and right-hand sides of Eq. (2)? We use the same criterion that was used in Ref. [3] to answer this question. The self-coupling needs to be measured at least to an accuracy equal to the maximum allowed deviation in its value [relative to the SM expectation  $\sim m_h^2/(2v^2)$ ] in different BSM theories, assuming no other EWSB states are accessible at the LHC. This is because if other EWSB states are observed we will already know that the Higgs sector is exotic,<sup>1</sup> whereas if no such states are seen the measurement of the Higgs coupling deviations would be the primary evidence that the Higgs sector is nonstandard. We will call this maximum allowed deviation the physics target for the Higgs self-coupling measurement precision. In computing this maximum possible deviation in a given BSM theory, we will also ensure that all existing direct and indirect constraints are obeyed.

Regarding the Higgs boson mass, there is a separate reason why we might want to accurately measure it. In some theories the precise value of the Higgs mass tells us something about UV physics. For instance, it is well known that with a Higgs mass less than  $\sim 125$  GeV the SM cannot be correct up to arbitrarily high scales because the self-coupling becomes negative below the Planck scale due to renormalization-group running. This would mean that our Universe is in a metastable vacuum. Thus the accuracy of determination of the scale where the self-coupling

<sup>1</sup>In the case of the minimal supersymmetric Standard Model, we also discuss the deviations depending on the mass of the relevant superpartners to take into account a possible discovery of them.

\*On leave from Albert-Ludwigs-Universität Freiburg, Physikalisches Institut, Freiburg, Germany.

becomes negative, the so-called triviality scale, would depend in part on the Higgs boson mass measurement precision. Another such example is supersymmetry where the Higgs boson mass is related to the scale of superpartners. In the first section we will discuss how accurately the Higgs mass needs to be measured to determine the relevant UV physics in these scenarios. In the subsequent sections we compute the Higgs self-coupling measurement precision needed according to the criterion explained in the previous paragraph in theories with mixed-in singlets, composite Higgs models, theories with first-order electro-weak phase transition, the minimal supersymmetric Standard Model (MSSM) and finally the next to minimal supersymmetric Standard Model (NMSSM). Our conclusion summarizes the results.

## II. HOW WELL DO WE NEED TO MEASURE THE HIGGS BOSON MASS?

Measuring the mass of any newly discovered elementary particle is an important scientific endeavor. The ability to test the SM and to test ideas of physics beyond the SM requires us to know them. The ATLAS and CMS detectors are equipped well to find the precise value of the Higgs boson mass. For a Higgs boson that has a mass within a few GeV of 126 GeV, one expects through the  $H \rightarrow \gamma\gamma$  channel to determine its mass to within 0.1%,<sup>2</sup> implying  $\Delta m_h < 150$  MeV.

The conjecture we make here is that there is no conceivable need to measure the Higgs boson mass better than what the LHC can already do. A proof of that statement is not possible, but we will do the next best thing and show circumstances where it is important to know the Higgs boson mass, and ask if in these circumstances knowing the Higgs boson mass better than 150 MeV would be helpful. In each case the answer is no, but let us follow the details.

In the first illustration, we consider the question of the fate of the Universe [6]. At leading order the Higgs boson self-coupling is related to its mass by  $m_h^2 = 2\lambda_{\text{SM}}v^2$ . The effective potential that leads to spontaneous symmetry breaking, and therefore our current vacuum state with vacuum expectation value  $v = 246$  GeV, depends on this coupling as

$$V(h) = \frac{\lambda_{\text{SM}}(Q)}{4} h^4 + \dots, \quad (3)$$

where the additional terms are small for large field values of  $h$ . In a renormalization-group-improved potential the coupling  $\lambda_{\text{SM}}$  depends on the scale  $Q$ . If  $\lambda_{\text{SM}}(Q) < 0$ , then the Higgs vacuum can be destabilized by the “turning over” of the potential, and the Universe can tunnel to a new catastrophic ground state sometime in the future. The renormalization-group equation for  $\lambda_{\text{SM}}$  at leading order is [7]

$$\frac{d\lambda_{\text{SM}}}{d \ln Q} = \frac{1}{16\pi^2} \left[ \frac{9}{8} \left( \frac{3}{25} g_1^4 + \frac{2}{5} g_1^2 g_2^2 + g_2^4 \right) - \left( \frac{9}{5} g_1^2 + 9g_2^2 + 12y_t^2 \right) \lambda_{\text{SM}} - 6y_t^4 + 24\lambda_{\text{SM}}^2 \right],$$

where  $g'$  and  $g$  are the gauge couplings of  $U(1)_Y$  and  $SU(2)_L$ , respectively, and  $y_t$  is the top quark Yukawa coupling. The QCD gauge coupling,  $g_s$  with  $\alpha_s = g_s^2/(4\pi)$ , comes into the renormalization-group flow at two loops and also from the one-loop feedback in the top Yukawa coupling flow.

When the Higgs mass is small ( $\lambda_{\text{SM}} \ll 1$ ) the  $-6y_t^4$  term dominates and tends to push  $\lambda_{\text{SM}}$  to negative values at higher  $Q$ , whereas when the Higgs mass is large ( $\lambda_{\text{SM}} \gtrsim 1$ ) the  $24\lambda_{\text{SM}}^2$  term dominates and tends to increase the value of  $\lambda_{\text{SM}}$  with scale, keeping the vacuum stable. There is an interface between  $\lambda_{\text{SM}}(Q)$  going negative versus going positive before  $Q = M_{\text{Pl}}$ , and that value determines the critical Higgs mass on this interface. The equation for the critical Higgs mass needed for stability is [8]

$$m_h > 129.4 \text{ GeV} + 1.4 \text{ GeV} \left( \frac{m_t - 173.2 \text{ GeV}}{0.7 \text{ GeV}} \right) - 0.5 \text{ GeV} \left( \frac{\alpha_s(M_Z) - 0.1184}{0.0007} \right) \pm 1 \text{ GeV}, \quad (4)$$

where 1 GeV is the theoretical uncertainty, which arises mainly due to the uncertainty in the low-energy matching scale for the Higgs quartic,  $\lambda_{\text{SM}}$ . Given that the Higgs mass is probably between 125 and 126 GeV, and given the errors expressed in the above equation, it is not totally clear if we are in an unstable or stable vacuum. What is clear is that measuring the Higgs boson mass with accuracy better than the 150 MeV target that LHC can achieve would be of no help. From direct top quark uncertainties alone, we would have to know the top quark mass  $m_t$  to much better than 100 MeV, which will not happen given the inherent QCD uncertainties in top quark measurements. At the LHC the best one can hope for is  $\delta m_t \sim 1$  GeV [9], and even in the cleaner  $e^+e^-$  linear collider environment the best one can hope for is  $\delta m_t \sim 100$  MeV [10]—a measurement that is a generation away and still not good enough for these purposes.<sup>3</sup> Thus, a better measurement of the Higgs mass is of no value for addressing the vacuum stability of the Universe question.

The Higgs boson mass computation in supersymmetry is another example where a precise determination of the Higgs boson mass can be helpful to test and constrain a theory. One can define the Higgs boson mass calculation in the MSSM exactly [12] using the definition

<sup>2</sup>See, for example, Fig. 10.37 of CMS technical design report (TDR) [4] for  $30 \text{ fb}^{-1}$  of integrated luminosity at  $\sqrt{s} = 14$  TeV. See also Figs. 19–45 of the ATLAS TDR [5].

<sup>3</sup>One could also hope for direct measurement of  $y_t$  in  $t\bar{t}h$  production at LHC and an  $e^+e^-$  linear collider, but this determination accuracy would be well below (see [11]) the extractions directly from the top mass measurement assuming the SM.

$$m_h^2 = m_Z^2 \cos^2 2\beta + \frac{3G_F m_t^4}{\sqrt{2}\pi^2} \ln \frac{\Delta_S^2}{m_t^2}, \quad (5)$$

where  $\tan \beta$  is the ratio of vacuum expectation values of the two Higgs doublets of supersymmetry,  $G_F$  is the Fermi constant with  $v = (2^{1/4} \sqrt{G_F})^{-1} \approx 246$  GeV, and  $\Delta_S^2 = m_{\tilde{t}_1} m_{\tilde{t}_2}$  is the leading-order radiative correction with  $m_{\tilde{t}_1}$  ( $m_{\tilde{t}_2}$ ) being the lighter (heavier) top squark mass. The above expression is subject to threshold corrections, the most important one of these being the correction due to top squark mixing (see, e.g., [13]):

$$\delta m_h^2 = \frac{3G_F m_t^4}{2\sqrt{2}\pi^2} \frac{X_t^2}{M_S^2} \left(1 - \frac{X_t^2}{12M_S^2}\right), \quad (6)$$

where  $X_t \equiv A_t - \mu/\tan \beta$  is the top squark mixing parameter and  $M_S^2 = \tilde{m}_{\tilde{t}_1} \tilde{m}_{\tilde{t}_2}$  is the geometric mean of the top squark masses. In the language of Eq. (5) this is equivalent to

$$\Delta_S^2 = m_{\tilde{t}_1} m_{\tilde{t}_2} + m_t^2 \exp \left[ \frac{X_t^2}{2M_S^2} \left(1 - \frac{X_t^2}{12M_S^2}\right) \right]. \quad (7)$$

There are many unknowns in this equation at the moment, but if supersymmetry is discovered and the particle content is measured precisely then a test can be made of the consistency of the prediction for the lightest Higgs boson mass and the measured value.

To demonstrate how unimportant a more precise measurement of the Higgs mass is to test supersymmetry, let us suppose that  $\tan \beta$  is large and known exactly, such that  $\cos 2\beta = 1$  is a good estimate. Let us then suppose that  $m_t$  can be measured to within 100 MeV, which as we have described above will not be possible in the current generation but might be possible in  $t\bar{t}$  threshold scans at a future linear collider. Let us further suppose that  $\Delta_S$  can be measured to within 0.5%. Even at  $1\sigma$  uncertainty this is an extraordinarily optimistic assumption, since  $\Delta_S$ , taking higher-order corrections into account, is a complicated combination of many different superpartner mass and supersymmetry mixing angles. We use 0.5% because that could be the  $1\sigma$  statistical (only) error of right-handed squark mass measurements in  $e^+e^- \rightarrow \tilde{q}_R \tilde{\bar{q}}_R \rightarrow q\bar{q}\chi\chi$  [14]. The difference between the Higgs mass value when both  $m_t$  and  $\Delta_S$  are at their central values to both being  $1\sigma$  lower than their central values, under these optimistic assumptions, is  $\delta m_h = 149$  MeV, and for  $2\sigma$  shifts in  $m_t$  and  $\Delta_S$  the value is  $\delta m_h = 297$  MeV. This is right near and above the uncertainty in the Higgs mass measurement already anticipated at the LHC, and thus we can conclude that there is no need to measure the Higgs mass to better accuracy for this test within supersymmetry.

Finally, measurement of the Higgs boson self-coupling is a physics measurement goal in next generation experiments. One reason to perform this measurement is to determine if this coupling that governs double Higgs production matches the prediction that one can make for it in

the SM which depends directly on the mass of the Higgs boson. Within the SM, the relative uncertainty on the prediction of  $\lambda_{\text{SM}}$  compared to the uncertainty in the Higgs mass is  $\delta\lambda_{\text{SM}}/\lambda_{\text{SM}} \approx 1.6\%$  ( $\delta m_h/1$  GeV), with the coefficient subject to small higher-order corrections. There is no known measurement of the relative Higgs boson self-coupling at the LHC, its upgrades or any  $e^+e^-$  collider discussed that could even come close to approaching the subpercent value needed to test self-consistency with the Higgs boson mass measurement of 150 MeV anticipated at the LHC. The best estimates for the LHC upgrades are that  $\delta\lambda_{\text{SM}}/\lambda_{\text{SM}}$  might be measured to within +30% and -20%  $1\sigma$  accuracy with over 3000  $\text{fb}^{-1}$  of integrated luminosity at 14 TeV center of mass energy [15].

In a subsequent section we will ask a logically separate and more general question of how well do we need to measure  $\lambda_{\text{SM}}$  for it to be interesting and valuable. The narrow conclusion just above is only that an anticipated measurement of  $\lambda_{\text{SM}}$  in principle can be compared to the Higgs boson mass measurement as one test of the SM, but given the poor quality of determination of the self-coupling further improvements of the mass measurement would not be helpful.

The three examples provided here illustrate our general conjecture that the inherent uncertainties in experimental measurements of observables and higher-order computations of theory imply that there is no obvious physics justification for pursuing a Higgs boson mass measurement better than what the LHC can provide.

### III. HOW WELL DO WE NEED TO MEASURE THE HIGGS BOSON SELF-COUPLING?

In this section we will find the maximal self-coupling deviation from its SM value in different BSM theories. Before delving into the BSM scenarios let us first gain our bearings by expanding out the SM Higgs potential to show the relationships between the Higgs boson mass, vacuum expectation value, and self-coupling. The tree-level BSM scenarios will decouple to the SM results when the BSM decoupling parameters, like supersymmetry (SUSY) breaking mass or compositeness scale, go to infinity.

The SM Higgs potential is

$$V = m_{\Phi_{\text{SM}}}^2 |\Phi_{\text{SM}}|^2 + \lambda_{\text{SM}} |\Phi_{\text{SM}}|^4, \quad (8)$$

which has a minimum at nonzero value of the field when  $m_{\Phi_{\text{SM}}}^2 < 0$ . In this case one finds

$$\langle \Phi_{\text{SM}}^\dagger \Phi_{\text{SM}} \rangle = \frac{v^2}{2}, \quad \text{where } v^2 = \frac{-m_{\Phi_{\text{SM}}}^2}{\lambda_{\text{SM}}}. \quad (9)$$

At the minimum the Higgs doublet can be expanded about its vacuum expectation value:

$$\Phi_{\text{SM}} = \left( \begin{array}{c} G^\pm \\ \frac{h+v}{\sqrt{2}} + iG^0 \end{array} \right), \quad (10)$$

where  $G^{\pm,0}$  are the Goldstone bosons eaten by the  $W^{\pm}, Z^0$  gauge bosons and  $h$  is the physical propagating Higgs boson.

The interaction Lagrangian of the Higgs boson, when expanded about the minimum, is

$$\Delta \mathcal{L} = -\frac{m_h^2}{2} h^2 - \frac{g_{hhh}}{3!} h^3 - \frac{g_{hhhh}}{4!} h^4, \quad (11)$$

where

$$m_h^2 \equiv 2\lambda_{\text{SM}} v^2, \quad (12)$$

$$g_{hhh} \equiv 6\lambda_{\text{SM}} v = \frac{3m_h^2}{v}, \quad (13)$$

$$g_{hhhh} \equiv 6\lambda_{\text{SM}} = \frac{3m_h^2}{v^2}. \quad (14)$$

In the subsequent sections we will use the normalization convention of Eq. (11) to define what is meant by the three-point ( $g_{hhh}$ ) Higgs self-interaction coupling. The Feynman rules for the three-point Higgs interaction is

$$hhh: -ig_{hhh} = -3i\frac{m_h^2}{v}. \quad (15)$$

Now we discuss the deviations of the  $hhh$  couplings from the above value in three different BSM models. It should be noted that the  $hhh$  coupling in the Standard Model,  $g_{hhh}^{\text{SM}}$ , as well as in the BSM models,  $g_{hhh}^{\text{BSM}}$ , is affected by higher-order corrections with a  $m_t^4$  contribution being the leading one-loop term [see Eq. (A9)]. Although this contribution leads to a sizable correction of the triple Higgs coupling, it affects both the numerator and denominator of  $g_{hhh}^{\text{BSM}}/g_{hhh}^{\text{SM}}$  leading to an overall correction to the ratio that is small. Thus we have neglected this effect for computation of the relative triple Higgs coupling deviation:  $g_{hhh}^{\text{BSM}}/g_{hhh}^{\text{SM}} - 1$ . For the deviations in the MSSM, we have explicitly checked that the  $m_t^4$  contribution does not change our results significantly (see the Appendix).

We should keep in mind that the processes by which the  $hhh$  coupling is measured [16] at  $pp$  and  $e^+e^-$  colliders,  $gg \rightarrow hh$  and  $VV \rightarrow hh$ , can have additional contributions beyond the deviations in the  $hhh$  vertex. For example, a heavier Higgs mixing with the lighter Higgs boson can generate an  $Hhh$  coupling which can contribute to  $gg \rightarrow H^* \rightarrow hh$ . Or new physics can contribute to the effective gluon-gluon- $h$ - $h$  four-point interaction [17]. Furthermore, as with the  $Hhh$  example, additional dynamics in electro-weak symmetry breaking can produce non-SM local vertices in the effective theory  $t\bar{t}hh$ ,  $W^+W^-hh$ , and  $ZZhh$  that contribute to the  $hh$  production cross section. All of these potential extra contributions are not considered, or are taken to be subdominant, in the discussion below. We are interested only in what deviations different theories put on  $hhh$  and leave the details of how one measures it, and

how it may need to be separated from other dynamics, to another study.

### A. Mixed-in singlets

Let us consider a theory with an extra singlet where the singlet mixes with the SM Higgs through a renormalizable operator like  $|\Phi_{\text{SM}}|^2|\Phi_H|^2$  [18,19]. The Lagrangian for our model is

$$\begin{aligned} \mathcal{L}_{\text{Higgs}} = & |\mathcal{D}_\mu \Phi_{\text{SM}}|^2 + |\mathcal{D}_\mu \Phi_H|^2 - m_{\Phi_{\text{SM}}}^2 |\Phi_{\text{SM}}|^2 \\ & - m_{\Phi_H}^2 |\Phi_H|^2 - \lambda |\Phi_{\text{SM}}|^4 - \rho |\Phi_H|^4 \\ & - \eta |\Phi_{\text{SM}}|^2 |\Phi_H|^2. \end{aligned}$$

The physical component fields are written as

$$\Phi_{\text{SM}} = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ \phi_{\text{SM}} + v \end{pmatrix}, \quad \Phi_H = \frac{1}{\sqrt{2}} (\phi_H + v'), \quad (16)$$

where  $v$  ( $\approx 246$  GeV) and  $v'$  are vacuum expectation values around which the  $\Phi_{\text{SM}}$  and  $\Phi_H$  are expanded. After diagonalizing the mass matrix, we rotate from the gauge eigenstates  $\phi_{\text{SM}}, \phi_H$  to mass eigenstates  $h, H$ :

$$\phi_{\text{SM}} = \cos \theta_h h + \sin \theta_h H, \quad (17)$$

$$\phi_H = -\sin \theta_h h + \cos \theta_h H. \quad (18)$$

The mixing angle  $\theta_h$  and the mass eigenvalues are given by

$$\begin{aligned} \tan \theta_h = & \frac{\eta v v'}{(-\lambda v^2 + \rho v'^2) + \sqrt{(\lambda v^2 - \rho v'^2)^2 + \eta^2 v^2 v'^2}}, \\ m_{h,H}^2 = & (\lambda v^2 + \rho v'^2) \pm \sqrt{(\lambda v^2 - \rho v'^2)^2 + \eta^2 v^2 v'^2}. \end{aligned}$$

Using the above equations we can express  $\lambda, \rho$  and  $\eta$  in terms of  $m_h^2, m_H^2, \theta_h, v$  and  $v'$ :

$$\begin{aligned} \lambda = & \frac{c_h^2 m_h^2 + s_h^2 m_H^2}{2v^2}, \quad \eta = c_h s_h \frac{m_H^2 - m_h^2}{v v'}, \\ \rho = & \frac{c_h^2 m_H^2 + s_h^2 m_h^2}{2v'^2}, \end{aligned} \quad (19)$$

where  $c_h \equiv \cos \theta_h$  and  $s_h \equiv \sin \theta_h$ . The above Lagrangian gives the  $h^3$  interaction

$$-\left( \lambda c_h^3 v - \rho s_h^3 v' - \frac{\eta}{2} c_h^2 s_h v' + \frac{\eta}{2} c_h s_h^2 v \right) h^3. \quad (20)$$

Using Eq. (19) we can now rewrite the Lagrangian terms for the cubic Higgs interaction as

$$g_{hhh} = \frac{3m_h^2}{v} \left( c_h^3 - s_h^3 \frac{v}{v'} \right). \quad (21)$$

Recall that the SM value for the Higgs cubic coupling is  $3m_h^2/v$ .

The maximum deviation of the tri-Higgs boson coupling from its SM value occurs for the maximum allowed  $s_h^2$  value. This can be found from Fig. 1. The region below the






FIG. 1 (color online). The area below the dashed line in the  $s_h^2 - m_H$  plane is allowed by electroweak precision tests (EWPT) at the 90% C.L. in the presence of a mixed-in singlet. The area above, and to the left, of the solid line is the area where the heavy mixed-in singlet Higgs boson is detectable with  $100 \text{ fb}^{-1}$  data at the 14 TeV LHC. The maximum allowed  $s_h^2$  value that can evade detection is thus given by the intersection of the two lines and is  $s_h^2 = 0.12$ .

dashed curve is the region allowed by electroweak precision tests and the region above the solid curve is the region where  $H$  is detectable with  $100 \text{ fb}^{-1}$  data. For details about how these curves were made see Ref. [3]. Clearly the maximum allowed value of  $s_h^2$  given that  $H$  is not detectable is

$$(s_h^2)_{\text{max}} = 12\%. \quad (22)$$

This tells us that the maximum fractional deviation of the coupling with respect to the SM value Eq. (21) is

$$(c_h^3 - 1) - s_h^3 \frac{v}{v'} \lesssim -18\% - 4\% \frac{v}{v'}. \quad (23)$$

Although  $v'$  is not known, the second term above is small compared to the first one ( $\sim s_h^3$ ) and can be safely ignored. This gives

$$\frac{\Delta g_{hhh}^{\text{target}}}{g_{hhh}^{\text{SM}}} = -18\%. \quad (24)$$

This sets the target for the Higgs boson self-coupling measurement in the context of the mixed-in singlet scenario.

## B. Higher-dimensional operators and composite models

We want to consider the effect of higher-dimensional operators on Higgs physics. Such theories are low-energy effective theories of a more fundamental theory. The quintessential example of this is the prospect of a composite Higgs boson. That will be our primary focus in this section, with some reference later to another theory of dimension-six operators that gives rise to a possible first-order phase transition for electroweak symmetry breaking.

Let us consider composite Higgs models where the Higgs is a pseudo-Nambu-Goldstone boson and thus its

mass is much lighter than the strong scale. Explicit models realizing this are little Higgs models [20] and holographic composite Higgs models [21]. An effective field theory for such a strongly interacting light Higgs (SILH) has been developed in Ref. [22]. The SILH Lagrangian contains higher-dimensional operators involving SM fields that supplement the SM Lagrangian. It is characterized by two independent parameters: the mass of the new resonances  $m_\rho$  and their coupling  $g_\rho$ . The ‘‘pion’’ decay constant  $f$  is given by

$$m_\rho = g_\rho f, \quad (25)$$

where  $g_\rho \leq 4\pi$ .

Here we do not list all the operators in the SILH Lagrangian but only those relevant to us, i.e. those that affect the triple Higgs coupling:

$$\begin{aligned} \mathcal{L}_{\text{SILH}} = & \frac{c_H}{2f^2} \partial^\mu (\Phi_{\text{SM}}^\dagger \Phi_{\text{SM}}) \partial_\mu (\Phi_{\text{SM}}^\dagger \Phi_{\text{SM}}) \\ & + \frac{c_6 \lambda}{f^2} (\Phi_{\text{SM}}^\dagger \Phi_{\text{SM}})^3. \end{aligned}$$

The coefficients of the above operators have been estimated using naive dimensional analysis [22,23] such that the couplings  $c_H$  and  $c_6$  are expected to be  $\mathcal{O}(1)$  numbers. Electroweak precision constraint from the measurement of the  $S$  parameter<sup>4</sup> requires at 90% C.L. (see [22,24])

$$m_\rho \gtrsim 2.6 \text{ TeV}. \quad (26)$$

Precision constraints also require at 90% C.L. [3]

$$c_H \xi \lesssim 0.15, \quad (27)$$

where  $\xi = v^2/f^2$ .

Let us now review how to obtain the triple Higgs coupling expression in the presence of these additional operators. The Higgs potential is modified by the operator with coupling  $c_6$  as follows:

$$\begin{aligned} V(H) = & \mu^2 (\Phi_{\text{SM}}^\dagger \Phi_{\text{SM}}) + \lambda (\Phi_{\text{SM}}^\dagger \Phi_{\text{SM}})^2 \\ & + \frac{c_6 \lambda}{f^2} (\Phi_{\text{SM}}^\dagger \Phi_{\text{SM}})^3. \end{aligned}$$

We can eliminate  $\mu^2$  using the minimization condition:

$$\mu^2 = -\lambda v^2 - \frac{3}{4} c_6 \lambda \xi v^2 \quad (28)$$

and then find the mass of the Higgs boson at the minima:

$$m_h^2 = 2\lambda v^2 + 3c_6 \lambda \xi v^2, \quad (29)$$

where we have used Eq. (28).

<sup>4</sup>Note that the constraint from the  $T$  parameter is more severe but this is avoided by imposing custodial symmetry in specific composite Higgs models.

Now let us write down the cubic Higgs terms from the SM and the operators above:

$$\left(-\lambda - \frac{5}{2}c_6\lambda\xi\right)h^3v. \quad (30)$$

Substituting  $\lambda$  from Eq. (29) results in

$$-\frac{m_h^2}{2v^2}(1 + c_6\xi)v h^3. \quad (31)$$

Finally we must consider the effect of the operator proportional to  $c_H$  which leads to the following derivative couplings:

$$\frac{c_H\xi}{2}\left(1 + \frac{h}{v}\right)^2\partial_\mu h\partial^\mu h. \quad (32)$$

These terms can be eliminated by the following redefinition of the Higgs field [22]:

$$h \rightarrow h - \frac{c_H\xi}{2}h\left(1 + \frac{h}{v} + \frac{h^2}{3v^2}\right). \quad (33)$$

Note that the above redefinition shifts  $m_h$  but not the Higgs vacuum expectation value. Thus we recover the expression for the triple Higgs coupling,  $g_{hhh}/3!$ , correct up to order  $\xi$ , already presented in Ref. [22]:

$$-\frac{m_h^2}{2v}\left(1 + c_6\xi - \frac{3}{2}c_H\xi\right)h^3. \quad (34)$$

In the above equation  $\xi \rightarrow 0$  is the SM limit. Using the constraint in Eq. (27) we can thus derive the self-coupling target

$$\frac{\Delta g_{hhh}}{g_{hhh}} = \frac{c_6}{c_H}(c_H\xi)_{\max} - \frac{3}{2}(c_H\xi)_{\max} \approx 0.15\frac{c_6}{c_H} - 0.23. \quad (35)$$

In the past in a different context with a different theoretical motivation, the operator  $(\Phi_{\text{SM}}^\dagger\Phi_{\text{SM}})^3$  has been studied. With such an operator one can achieve a first-order electroweak phase transition [25] that may be relevant for the problem of baryogenesis. It was shown that one needs

$$\frac{f}{\sqrt{c_6}} > 550 \text{ GeV} \quad (36)$$

to ensure that the expansion of the Universe does not disallow percolation of bubbles of metastable vacuum. This corresponds to  $c_6\xi < 0.2$ . This would mean that the maximum coupling deviation allowed in such a scenario would be

$$\frac{\Delta g_{hhh}}{g_{hhh}} = c_6\xi = 0.2. \quad (37)$$

Equations (35) and (37) set a target of  $\sim 20\%$  needed for measuring the Higgs boson self-coupling interaction in these higher-dimensional operator theories, including

composite Higgs models and models with Higgs potentials that allow first-order electroweak phase transitions.

### C. Minimal supersymmetric Standard Model

The MSSM exhibits an extended Higgs sector with two Higgs boson doublets,  $H_d$  and  $H_u$ , which couple to down- and up-type quarks, respectively. The two neutral,  $CP$ -even components of the Higgs boson doublets  $H_d^0$ ,  $H_u^0$  mix and form the mass eigenstates  $H$  and  $h$ . For our discussion, we will assume that the lighter  $CP$ -even Higgs boson  $h$  is the SM-like one. In addition to the Higgs sector extension with respect to the SM, the MSSM is comprised of superpartners to the particles of the two Higgs doublet model which might be discovered at the LHC depending on the scenario realized in nature. For the investigation of the Higgs sector, the top squarks, superpartners of the top quarks, are most relevant.

In the following, we will discuss the maximal possible deviation of the MSSM triple Higgs coupling with respect to the SM one if no other Higgs boson will be discovered at the LHC. We model the LHC discovery potential according to Fig. 1.21 of [14] inspired from Chap. 19 in [5]. To take into account the possible discovery of superpartners at the LHC we will discuss the maximal deviations, noting its dependence of the mass of the most relevant superpartners, the top squarks.

At tree level the tri-Higgs boson coupling is

$$g_{hhh}^{\text{tree}} = \frac{3m_Z^2}{v}\cos(2\alpha)\sin(\alpha + \beta) \quad (38)$$

with  $v = (2^{1/4}\sqrt{G_F})^{-1} \approx 246 \text{ GeV}$  and  $G_F$  being the Fermi constant. The angle  $\alpha$  describes the transformation from the Higgs boson interaction eigenstates to the mass eigenstates. The  $Z$  boson mass is denoted by  $m_Z$  and the angle  $\beta$  is defined as the ratio of the Higgs vacuum expectation values,  $\tan\beta = v_u/v_d$ . In the decoupling limit of  $m_A \rightarrow \infty$  one finds that  $\alpha \rightarrow \beta - \pi/2$  and  $m_h^2 = m_Z^2\cos^2 2\beta$ . Substituting this into Eq. (38) gives  $g_{hhh}^{\text{tree}} \rightarrow 3m_h^2/v$ , matching the SM result as expected.

At tree level, the mass of the lighter  $CP$ -even Higgs boson has an upper theoretical bound at  $m_Z$ ; however, large radiative corrections shift the mass to higher values. In order to compare the triple Higgs coupling in the MSSM and the SM, it is important to apply the same approximations in both cases—which includes calculating the Higgs mass and the triple Higgs coupling to the same order. It is clear that we have to take into account higher-order effects. However, we cannot make use of all the higher-order corrections known for the computation of the mass of the lightest MSSM Higgs boson when the analogous corrections for the triple Higgs coupling are not known.

In Ref. [26] renormalization-group-improved corrections to the effective potential are presented, taking into account dominant one- and two-loop contributions. For our investigation, using this effective potential and deriving the

Higgs mass and the triple coupling from there has the advantage of automatically applying the same approximation for the Higgs mass and the triple Higgs coupling. Also, the two-loop contributions have a sizable effect on the coupling deviations. See also the discussion in the Appendix with results of various other more limiting approximations.

The effective potential, used in Ref. [26], is defined according to the conventions of Ref. [27] as

$$V^{\text{eff}} = \frac{\lambda_1}{2}|H_d|^4 + \frac{\lambda_2}{2}|H_u|^4 + \lambda_3|H_d|^2|H_u|^2 + \lambda_4|H_d^\dagger H_u|^2 + \left[ \frac{\lambda_5}{2}|H_d^\dagger H_u|^2 + (\lambda_6|H_d|^2 + \lambda_7|H_u|^2)H_d^\dagger H_u + \text{c.c.} \right],$$

where the coefficients  $\lambda_i$ ,  $i = 1, \dots, 7$  contain also the loop corrections. Explicit expressions for the coefficients  $\lambda_i$  can be found in the Appendix. With  $H_d$  and  $H_u$  developing a vacuum expectation value, the effective potential can be expanded about these vacuum expectation values, and the triple Higgs vertices in terms of interaction eigenstates can be derived:

$$g_{H_d^0 H_d^0 H_d^0} = 3v[\lambda_1 \cos \beta + \lambda_6 \sin \beta], \quad (39)$$

$$g_{H_d^0 H_d^0 H_u^0} = v[3\lambda_6 \cos \beta + (\lambda_3 + \lambda_4 + \lambda_5) \sin \beta], \quad (40)$$

$$g_{H_u^0 H_u^0 H_u^0} = v[3\lambda_7 \sin \beta + (\lambda_3 + \lambda_4 + \lambda_5) \cos \beta], \quad (41)$$

$$g_{H_u^0 H_u^0 H_d^0} = 3v[\lambda_2 \sin \beta + \lambda_7 \cos \beta]. \quad (42)$$

The transformation into mass eigenstates can be done with the help of the effective mixing angle  $\alpha$ , obtained by diagonalizing the Higgs mass matrix while including the corresponding higher-order corrections. This leads to

$$g_{hhh} = -\sin^3 \alpha g_{H_d^0 H_d^0 H_d^0} + 3\sin^2 \alpha \cos \alpha g_{H_d^0 H_d^0 H_u^0} - 3\cos^2 \alpha \sin \alpha g_{H_d^0 H_u^0 H_u^0} + \cos^3 \alpha g_{H_u^0 H_u^0 H_u^0}. \quad (43)$$

In the decoupling limit, the above potential gives a Higgs mass

$$m_h^2 = v^2[\lambda_1 \cos^4 \beta + \lambda_2 \sin^4 \beta + 2(\lambda_3 + \lambda_4 + \lambda_5) \times \sin^2 \beta \cos^2 \beta + 4\sin \beta \cos \beta (\lambda_6 \cos^2 \beta + \lambda_7 \sin^2 \beta)], \quad (44)$$

and the triple Higgs coupling reduces to the correct SM value.

The coefficients  $\lambda_i$  given in Ref. [26] comprise the dominant one-loop  $\mathcal{O}(y_t^4)$  contributions, stemming from the large top Yukawa coupling  $y_t$ , as well as one-loop corrections of order  $\mathcal{O}(m_Z^2/v^2 y_t^2)$ ,  $\mathcal{O}(y_t^2 y_b^2)$ ,  $\mathcal{O}(y_b^4)$  and  $\mathcal{O}(m_Z^2/v^2 y_b^2)$ , where the latter ones are proportional to the bottom Yukawa coupling  $y_b$ . The dominant two-loop contributions are of the order  $\mathcal{O}(y_t^4 \alpha_s)$  and  $\mathcal{O}(y_t^6)$ . They are

included into the  $\lambda_i$  as well as two-loop terms proportional to  $y_b$  of order  $\mathcal{O}(y_b^4 \alpha_s)$ ,  $\mathcal{O}(y_b^6)$ ,  $\mathcal{O}(y_t^4 y_b^2)$  and  $\mathcal{O}(y_b^4 y_t^2)$ .

For the investigation of the deviation of the triple Higgs coupling from the SM one we performed a scan over several parameters:  $\tan \beta$  from 2 to 45 and the  $CP$ -odd Higgs boson mass  $m_A$  from 200 to 800 GeV, which are the free parameters in the Higgs sector at tree level, the diagonal soft breaking parameters of the top squark mass matrices  $M_{L_{\tilde{Q}_3}} = M_{R_t}$  from 100 GeV to 3 TeV, the Higgs superfield mixing parameter  $\mu$  between  $\pm 1$  TeV and the top squark mixing parameter  $X_t = A_t - \mu/\tan \beta$  between  $\pm 150$  GeV  $\cdot n_{\text{max}}$ , where  $n_{\text{max}}$  is the nearest bigger integer to  $\sqrt{6}M_{L_{\tilde{Q}_3}}/(150 \text{ GeV})$  ( $A_t$  being the top soft breaking trilinear coupling).

The top Yukawa coupling is determined as  $y_t = \sqrt{2}\bar{m}_t/(v \sin \beta)$  with the running top quark mass  $\bar{m}_t = m_t/[1 + 4\alpha_s/(3\pi)]$ ,  $m_t = 173.2$  GeV. The bottom Yukawa coupling is

$$y_b = \frac{\sqrt{2}}{v \cos \beta} \frac{\bar{m}_b}{1 + \Delta_b}, \quad (45)$$

where [28]

$$\Delta_b = \frac{2\alpha_s}{3\pi} m_{\tilde{g}} \mu \tan \beta I(m_{\tilde{b}_1}, m_{\tilde{b}_2}, m_{\tilde{g}}) + \frac{y_t^2}{16\pi^2} A_t \mu \tan \beta I(m_{\tilde{t}_1}, m_{\tilde{t}_2}, \mu).$$

The function  $I$  is defined as

$$I(m_1, m_2, m_3) = -\frac{m_1^2 m_2^2 \log \frac{m_1^2}{m_2^2} + m_2^2 m_3^2 \log \frac{m_2^2}{m_3^2} + m_3^2 m_1^2 \log \frac{m_3^2}{m_1^2}}{(m_1^2 - m_2^2)(m_2^2 - m_3^2)(m_3^2 - m_1^2)}.$$

The running bottom quark mass is chosen as  $\bar{m}_b = 2.9$  GeV, the gluino mass as  $m_{\tilde{g}} = 1$  TeV and the diagonal right-handed soft-SUSY-breaking parameter in the sbottom-mass matrix as  $M_{\tilde{b}} = 1.2$  TeV. The resulting sbottom masses are denoted as  $m_{\tilde{b}_1}$  and  $m_{\tilde{b}_2}$ .

The deviations of the triple Higgs coupling,  $\Delta g_{hhh}/g_{hhh}^{\text{SM}}$  with  $\Delta g_{hhh} = g_{hhh} - g_{hhh}^{\text{SM}}$ , are shown in Fig. 2 versus  $m_A$  and  $\tan \beta$  in the left and the right plot, respectively. The deviations are calculated taking into account all the corrections given in Ref. [26]. The parameter points either fulfill the mass constraint  $122 \text{ GeV} \leq m_h \leq 129 \text{ GeV}$  or a relaxed mass constraint. In the case of the relaxed mass constraint the Higgs boson mass is calculated neglecting the terms  $\mathcal{O}(m_Z^2/v^2 y_t^2)$ . If the resulting Higgs mass fulfills the constraint  $122 \text{ GeV} \leq m_h \leq 129 \text{ GeV}$ , the parameter point is kept (the deviations however are calculated with the full expressions). The terms of  $\mathcal{O}(m_Z^2/v^2 y_t^2)$  lead to a Higgs boson mass reduction. This means that parameter points for which only the relaxed mass constraint is fulfilled actually correspond to a Higgs boson mass below 122 GeV within the used approximation. Points with a




FIG. 2 (color online). The triple Higgs coupling deviations in the MSSM as a function of  $m_A$  in the renormalization-group-improved leading-log approximation, with  $y_t$  defined via the running top quark mass. The red (plus) region corresponds to points in the  $\tan\beta - m_A$  plane lying in the several Higgs boson discovery region, while all the other points lie in the single Higgs boson discovery region. The light blue (square), yellow (circle) and green (cross) points correspond to mass values of the lighter top squark of  $m_{\tilde{t}_1} < 1.0$  TeV, the  $1.0 \text{ TeV} \leq m_{\tilde{t}_1} < 2.5$  TeV and  $m_{\tilde{t}_1} \geq 2.5$  TeV, respectively. The plot on the right is the same but versus  $\tan\beta$ .

relaxed mass constraint are especially those with small  $\tan\beta$  and in particular all points with  $\tan\beta \leq 5$ . We know that other approximations including two-loop order terms lead to Higgs boson mass values within the interval  $122 \text{ GeV} \leq m_h \leq 129 \text{ GeV}$  for  $\tan\beta = 5$  and we observed that the deviations are of similar size independent on whether the  $\mathcal{O}(m_{\tilde{Z}}^2/v^2 y_t^2)$  terms are taken into account or not [compare with Fig. 4 (right) in the Appendix]. Therefore we include also the points with a relaxed mass constraint.

In Fig. 2 the red points (plus) indicate that for this parameter point several Higgs bosons are expected to be discovered at the LHC while all the other points correspond to the single Higgs boson discovery region. The colors light blue (square), yellow (circle) and green (cross) indicate the mass of the lighter top squark as below 1 TeV, between 1 TeV and 2.5 TeV and above 2.5 TeV, respectively.

The largest deviations are found for  $m_A = 200$  GeV and  $\tan\beta = 5$  and amount to roughly  $-15\%$ , with slightly larger deviations for the light top squark mass in the interval  $1.0 \text{ TeV} \leq m_{\tilde{t}_1} < 2.5$  TeV than for the light top squark heavier than 2.5 TeV. Taking only points which fulfill the Higgs mass constraint  $122 \text{ GeV} \leq m_h \leq 129 \text{ GeV}$  within the applied approximation leads to a maximal deviation of  $-3\%$  to  $-4\%$  for  $m_A \approx 240$  GeV,  $\tan\beta \approx 7$  and the lighter top squark mass in the interval  $1.0 \text{ TeV} \leq m_{\tilde{t}_1} < 2.5$  TeV; for the light top squark being heavier than 2.5 TeV the maximal deviations in this case are about  $-2\%$  for  $m_A \approx 280$  GeV and  $\tan\beta \approx 8$ .

Including the  $y_b$  terms does not have a substantial impact on the allowed deviations. In Fig. 2 only points with a  $\Delta\rho$  contribution from top squarks and bottoms (see [29]) smaller than 0.001 are included [30]—however this constraint does not change the result qualitatively given that direct constraints are generally more constraining for superpartners than precision electroweak constraints.

In summary, we have pointed out that the answer to the question of how large deviations of the triple Higgs coupling can be in the MSSM is quite sensitive to the applied approximation (see also the Appendix). For large  $m_A$ , the decoupling effect leads to small deviations, below  $\sim -2\%$ , while for  $m_A \approx 200$  GeV and small  $\tan\beta$  the deviations can be as high as order  $\mathcal{O}(-15\%)$ . This sets the target for the self-coupling measurement in the MSSM context.

#### D. Next to minimal supersymmetric Standard Model

We now consider triple Higgs coupling deviations in the NMSSM [31]. In this section we will assume that the  $CP$ -even singlet in the NMSSM is very heavy and thus undetectable at the LHC. If the singlet soft mass  $m_S$  is made very large (while other soft parameters are not changed), the NMSSM Higgs sector reduces to a MSSM-like two Higgs doublet model and a decoupled singlet.<sup>5</sup> If  $m_A$  and  $\tan\beta$  are taken as input parameters, effectively this limit amounts to taking an additional term in the scalar potential:

$$\Delta V = \lambda_S^2 (H_d^0)^2 (H_u^0)^2, \quad (46)$$

where  $H_u^0$  and  $H_d^0$  are the neutral components of the up- and down-type Higgs doublets, respectively. This leads to a contribution to the triple Higgs vertex<sup>6</sup>

<sup>5</sup>This can be checked for instance by looking at the scalar and pseudoscalar mass matrices given in Ref. [31], where the most general superpotential has been considered, and taking the limit  $m_S \rightarrow \infty$ .

<sup>6</sup>One can derive triple Higgs vertex including singlet-doublet mixing effects, in a similar way, by using the effective potential given in Ref. [32], which includes corrections due to singlet-doublet mixing. For our investigation, this would require a more detailed study of the discovery potential of the singletlike Higgs boson at the LHC in order to find the target in this model. This, however, goes beyond the scope of this paper.






FIG. 3 (color online). The triple Higgs coupling target in the NMSSM with  $M_S = 500$  GeV as a function of  $m_A$  for various values of  $\tan \beta$  where the singlet coupling remains perturbative below 10 TeV.

$$\Delta g_{hhh} = -\frac{3v}{2} \lambda_S^2 \sin 2\alpha \cos(\alpha + \beta), \quad (47)$$

in addition to the MSSM tree-level contribution. Once again in the decoupling limit we have  $m_A \rightarrow \infty$  and  $\alpha \rightarrow \beta - \frac{\pi}{2}$  so that  $g_{hhh}$  again reduces to its SM value  $3m_h^2/v$  where, for the NMSSM in the decoupling limit, we have

$$m_h^2 = m_Z^2 \cos^2 2\beta + \frac{1}{2} \lambda_S^2 v^2 \sin^2 2\beta + \Delta_t, \quad (48)$$

where  $\Delta_t$  is the loop contribution due to the top squarks discussed in detail in the last subsection.

In Fig. 3 we show the Higgs self-coupling deviation as a function of  $m_A$  for  $\tan \beta = 2, 5$  and  $7.5$ . We have assumed  $M_S = 500$  GeV and  $X_t = 0$  for the top squark loop contribution.<sup>7</sup> The self-coupling deviation increases with  $\tan \beta$  because the value of  $\lambda_S$  required to raise the Higgs mass to 126 GeV increases [this can be understood from the  $\sin^2 2\beta$  factor in Eq. (48)]. For  $\tan \beta = 2$  we find  $\lambda_S \leq 0.7$ , which satisfies the condition for perturbativity up to the grand unification scale [33] ( $M_{\text{GUT}} \sim 2 \times 10^{16}$  GeV), whereas for  $\tan \beta = 7.5$  we find  $\lambda_S \leq 2$ , the upper value ( $\lambda_S = 2$ ) leading to a divergence in  $\lambda_S$  at  $\sim 10$  TeV [34]. For  $\tan \beta > 7.5$  we find that the condition for perturbativity up to 10 TeV,  $\lambda_S < 2$ , is not satisfied. Thus the maximum possible deviation, if we require perturbativity up to 10 TeV, is about  $-25\%$  for  $\tan \beta = 7.5$ ,  $m_A = 500$  GeV.

Now we come to the question, would the heavier Higgs remain undetected by the LHC for this point  $\tan \beta = 7.5$ ,  $m_A = 500$  GeV? In the case of the MSSM this point lies outside the LHC reach of heavy supersymmetric Higgs searches (see Fig. 1.21 of Ref. [14]). In the NMSSM the coupling of the heavier Higgs bosons to down-type quarks and vector bosons is the same up to the percent level while the coupling to up-type quarks is reduced with respect to

<sup>7</sup>For higher top squark masses, lower values of  $\lambda_S$  would have to be chosen for the same Higgs boson mass and hence the model would be more MSSM-like.

TABLE I. Summary of the physics-based targets for the triple Higgs boson coupling. The target is based on scenarios where no other exotic electroweak symmetry breaking state (e.g., new Higgs bosons or “ $\rho$  particle”) is found at the LHC except one: the  $\sim 126$  GeV SM-like Higgs boson. Percentages quoted are approximate maximal deviations for each model based on the discussion in the text. For the  $\Delta g_{hhh}/g_{hhh}^{\text{SM}}$  values of supersymmetry, superscript *a* refers to the case of high  $\tan \beta > 10$  and no superpartners are found at the LHC, and superscript *b* refers to all other cases, with the maximum value of  $-15\%$  reached for the special case of  $\tan \beta \approx 5$ . In the last row, the best estimates for the  $1\sigma$  accuracy of the measurement of the triple Higgs coupling at the LHC with  $3 \text{ ab}^{-1}$  integrated luminosity is given. It is assumed here that no additional dynamics or operators contribute to non-SM shifts in  $pp \rightarrow hh$  except the self-coupling.

Model	$\Delta g_{hhh}/g_{hhh}^{\text{SM}}$
Mixed-in singlet	$-18\%$
Composite Higgs	Tens of %
Minimal supersymmetry	$-2\%^a - 15\%^b$
NMSSM	$-25\%$
LHC $3 \text{ ab}^{-1}$ [15]	$[-20\%, +30\%]$

the MSSM. This means that we expect similar (in processes controlled by heavy Higgs boson couplings to down-type fermions like  $bb \rightarrow H \rightarrow \tau\tau$ ) or smaller cross sections (if the process involves, for instance, gluon fusion where coupling to the top would be suppressed relative to the MSSM). Thus we would expect that if a point like  $\tan \beta = 7.5$ ,  $m_A = 500$  GeV is beyond LHC reach for the MSSM the same would hold for the NMSSM too, given our construction. Thus  $\tan \beta = 7.5$ ,  $m_A = 500$  GeV indeed represents a point where the self-coupling deviation from SM is maximal, and the heavy Higgs bosons are beyond the LHC reach. The self-coupling deviation for this point,  $-25\%$ , is thus the target in the case of the NMSSM.

#### IV. CONCLUSIONS

To summarize, we have found that the 150 MeV uncertainty on the Higgs boson mass that ATLAS and CMS are scheduled to achieve is likely to be better than we will ever need to know it in the foreseeable future. Better determinations yield no obvious advantage in testing any proposed question about nature that we can formulate today.

On the other hand, we have shown that in beyond the SM scenarios there can be significant gains in our understanding of nature—including discovering and discerning new physics—if the self-coupling measurement can be performed much more accurately than LHC projections afford. Generally speaking, measurements need to determine the Higgs self-coupling to better than 20% in order to have the chance to see effects of new physics in this important channel, if no other dynamics associated with electroweak symmetry breaking are seen. Our results for different BSM theories are summarized in Table I. This is a challenging

target for future LHC upgrades and proposed  $e^+e^-$  linear collider options of the future.

## APPENDIX

The result for the tri-Higgs coupling presented in Sec. III C turns out to be sensitive to the choice of higher-order terms included in the computation. It is conceivable that adding more terms self-consistently to the tri-Higgs coupling and mass, and recomputing would also yield a change of similar size. For this reason we have included this Appendix to show the changes in the size of the tri-Higgs coupling for various approximations that exist in the literature.

We start our discussion taking into account radiative corrections to the Higgs triple coupling of the order of  $m_t^4$ . In the literature, two approximations can be found: one including all  $m_t^4$  contributions [35–37] and the other one based on a renormalization-group-improved leading-log approximation [26]. In the latter approach also two-loop leading-log terms are provided which are taken into account in Sec. III C.

In the leading-log approach we evaluate the coefficients  $\lambda_i$  with  $i = 1, \dots, 7$  of the effective potential Eq. (38), given in Ref. [26] as (showing here explicitly only the tree-level part and the  $m_t^4$  one-loop contributions)

$$\lambda_1 = \frac{m_Z^2}{v^2} - \frac{y_t^4}{32\pi^2} \frac{\mu^4}{M_{\text{SUSY}}^4}, \quad (\text{A1})$$

$$\lambda_2 = \frac{m_Z^2}{v^2} + \frac{3y_t^4}{8\pi^2} \left[ \log \frac{m_t^2}{M_{\text{SUSY}}^2} - \frac{A_t^2}{M_{\text{SUSY}}^2} \left( 1 - \frac{A_t^2}{12M_{\text{SUSY}}^2} \right) \right], \quad (\text{A2})$$

$$\lambda_3 = -\frac{m_Z^2}{v^2} (1 - 2c_W^2) + \frac{y_t^4}{32\pi^2} \frac{\mu^2}{M_{\text{SUSY}}^2} \left( 3 - \frac{A_t^2}{M_{\text{SUSY}}^2} \right), \quad (\text{A3})$$

$$\lambda_4 = -2\frac{m_Z^2}{v^2} c_W^2 + \frac{y_t^4}{32\pi^2} \frac{\mu^2}{M_{\text{SUSY}}^2} \left( 3 - \frac{A_t^2}{M_{\text{SUSY}}^2} \right), \quad (\text{A4})$$

$$\lambda_5 = -\frac{y_t^4}{32\pi^2} \frac{\mu^2 A_t^2}{M_{\text{SUSY}}^4}, \quad (\text{A5})$$

$$\lambda_6 = -\frac{y_t^4}{32\pi^2} \frac{\mu^3 A_t}{M_{\text{SUSY}}^4}, \quad (\text{A6})$$

$$\lambda_7 = \frac{y_t^4}{32\pi^2} \frac{\mu}{M_{\text{SUSY}}} \left( \frac{A_t^3}{M_{\text{SUSY}}^3} - 6 \frac{A_t}{M_{\text{SUSY}}} \right) \quad (\text{A7})$$

with  $c_W = m_W/m_Z$  being the cosine of the weak mixing angle and  $m_W$  the  $W$  mass. The parameter  $M_{\text{SUSY}}^2$  is defined as the average of the top squark masses squared:  $M_{\text{SUSY}}^2 = (m_{\tilde{t}_1}^2 + m_{\tilde{t}_2}^2)/2$ . The top soft breaking trilinear coupling is denoted as  $A_t$  and the top Yukawa coupling as  $y_t$ .

In Fig. 4 the relative deviations of the triple Higgs coupling,

$$\Delta g_{hhh}/g_{hhh}^{\text{SM}}, \quad \Delta g_{hhh} = g_{hhh} - g_{hhh}^{\text{SM}}, \quad (\text{A8})$$

are shown versus  $m_A$  with the MSSM coupling,  $g_{hhh}$ , as given in Eq. (43). The parameter scan is performed as described in Sec. III C. Only points with a Higgs boson mass of  $122 \text{ GeV} \leq m_h \leq 129 \text{ GeV}$ , evaluated in the particular approximations, are taken into account.




FIG. 4 (color online). The triple Higgs coupling deviations in the MSSM as a function of  $m_A$  at  $\mathcal{O}(y_t^4)$  in the renormalization-group-improved leading-log approximation, with  $y_t$  defined via the on-shell top quark mass,  $y_t = \sqrt{2}m_t/(v \sin \beta)$  with  $m_t = 173.2 \text{ GeV}$  (left plot) and including additionally  $\mathcal{O}(y_t^4 \alpha_s)$  and  $\mathcal{O}(y_t^6)$  terms, with  $y_t$  defined via the running mass  $y_t = \sqrt{2}\tilde{m}_t/(v \sin \beta)$  (right plot). The red points (plus) correspond to parameter points for which several Higgs bosons can be discovered at the LHC. All other points belong to the single Higgs boson discovery region and are colored according to the mass value of the lighter top squark  $m_{\tilde{t}_1}$ : light blue (square) for  $m_{\tilde{t}_1} < 1.0 \text{ TeV}$ , yellow (circle) for  $1.0 \leq m_{\tilde{t}_1} < 2.5 \text{ TeV}$ , and green ( $\times$ ) for  $m_{\tilde{t}_1} \geq 2.5 \text{ TeV}$ . In contrast to Fig. 2 in both plots, the contributions of the order of  $\mathcal{O}(m_Z^2/v^2 y_t^2)$ ,  $\mathcal{O}(y_t^2 y_b^2)$ ,  $\mathcal{O}(y_b^4)$ ,  $\mathcal{O}(m_Z^2/v^2 y_b^2)$ ,  $\mathcal{O}(y_b^4 \alpha_s)$ ,  $\mathcal{O}(y_b^6)$ ,  $\mathcal{O}(y_t^4 y_b^2)$  and  $\mathcal{O}(y_b^4 y_t^2)$  are neglected and the mass constraint of  $122 \text{ GeV} \leq m_h \leq 129 \text{ GeV}$  has been strictly applied.

In the left plot of Fig. 4 the deviations are calculated taking only into account the one-loop  $m_t^4$  terms; see Eqs. (A1)–(A7). The top Yukawa coupling is chosen as  $y_t = \sqrt{2}m_t/(v \sin \beta)$  with the top quark mass  $m_t = 173.2$  GeV. The red region (+) corresponds to parameter points in the  $m_A - \tan \beta$  plane with a several Higgs boson discovery potential at the LHC according to Fig. 1.21 of Ref. [14]. All other points belong to the region where only a single Higgs boson can be discovered. The largest deviations can be found for low  $m_A$  and can be larger than 30% in this approximation. The color coding of the single Higgs boson discovery region is the following: light blue (■) for the mass of the lighter top squark  $m_{\tilde{t}_1}$  below 1 TeV, yellow (•) for  $m_{\tilde{t}_1}$  between 1.0 and 2.5 TeV, and green (×) for  $m_{\tilde{t}_1}$  larger than 2.5 TeV. Within this leading-log approximation the largest deviations occur only for rather light top squark masses, and only for large  $m_A$  do we find points corresponding to top squark masses larger than 2.5 TeV. This is mainly due to the fact that the one-loop corrections to the Higgs mass are very large and for large top squark masses the light Higgs boson becomes too heavy with mass values above 129 GeV. The two-loop corrections to the Higgs boson mass decrease the mass so that it is to be expected that at two-loop order for a wider range of deviations large top squark masses can be found.

Applying  $y_t = \sqrt{2}\tilde{m}_t/(v \sin \beta)$  with the running top quark mass  $\tilde{m}_t = m_t/[1 + 4\alpha_s/(3\pi)]$ , as used in Ref. [26], instead of the top Yukawa coupling expressed by the top quark pole mass, leads to a rescaling of the one-loop contributions by  $\tilde{m}_t^4/m_t^4$ . As  $\tilde{m}_t < m_t$  this decreases the one-loop contributions to the triple Higgs coupling as well as to the Higgs mass. As a result the size of the coupling deviations is reduced, for  $m_A = 200$  GeV from roughly 0.35 to 0.25, while the regions where large top squark masses can be found are enlarged.

Using the exact  $m_t^4$  approximation for  $g_{hhh}$  as given in Refs. [35–37] gives a similar picture to the one shown in the left plot of Fig. 4 if the same approximation is used in the SM. The complete  $m_t^4$  contributions include a constant term which is also present in the SM:

$$g_{hhh}^{\text{SM}} = \frac{3}{v} \left( m_h^2 - \frac{\sqrt{2}G_F}{\pi^2} m_t^4 \right) \quad (\text{A9})$$

and which has to be taken into account to ensure a proper decoupling [36,38] ( $m_h$  is also understood to be calculated in the complete  $m_t^4$  approximation, of course). We find slight positive deviations, below 1% (using the top quark pole mass), in the complete  $m_t^4$  approximation which are not present in the renormalization-group-improved result, which is a rather small effect. However, going to larger  $X_t$  values, larger than the ones chosen in the scan, we find rather big deviations and the approximations seem to not work properly anymore.

In Fig. 4 (right), two-loop contributions to the effective potential of the order  $\mathcal{O}(y_t^4 \alpha_s)$  and  $\mathcal{O}(y_t^6)$  have been taken into account [the contributions of the order of  $\mathcal{O}(m_Z^2/v^2 y_t^2)$ ,  $\mathcal{O}(y_t^2 y_b^2)$ ,  $\mathcal{O}(y_b^4)$ ,  $\mathcal{O}(m_Z^2/v^2 y_b^2)$ ,  $\mathcal{O}(y_b^4 \alpha_s)$ ,  $\mathcal{O}(y_b^6)$ ,  $\mathcal{O}(y_t^4 y_b^2)$  and  $\mathcal{O}(y_b^4 y_t^2)$  are neglected in contrast to the results presented in Fig. 2]. The mass constraint of  $122 \text{ GeV} \leq m_h \leq 129 \text{ GeV}$  has been strictly applied (in contrast to Fig. 2). The top Yukawa coupling again is evaluated with the running top quark mass. The color coding is the same as before. In Fig. 4 the deviations of the couplings are shown in dependence on  $m_A$ . A further decrease of the deviations and an increase of the region with very large top squark masses can be observed. Top squark mass values larger than 2.5 TeV cover almost the complete single Higgs boson discovery region.

These results can be compared with the results shown in Sec. III C and Fig. 2. Taking into account the complete set of corrections given in Ref. [26] particularly reduces the Higgs mass and, therefore, keeping strictly the mass constraint of  $122 \text{ GeV} \leq m_h \leq 129 \text{ GeV}$  leads to a sizable reduction of the possible deviations as no parameter points are found for  $\tan \beta = 5$  and hence no points for  $m_A = 200$  GeV in the single Higgs boson discovery region, for which before we found the largest deviations. Relaxing however the mass constraint leads to a similar size of the deviations (see Sec. III C and Fig. 2).

- 
- [1] ATLAS Collaboration, *Phys. Lett. B* **716**, 1 (2012); CMS Collaboration, *Phys. Lett. B* **716**, 30 (2012).  
 [2] See, e.g., J. Baglio, A. Djouadi, R. Gröber, M. M. Mühlleitner, J. Quevillon, and M. Spira, *J. High Energy Phys.* **04** (2013) 151; M. J. Dolan, C. Englert, and M. Spannowsky, *J. High Energy Phys.* **10** (2012) 112; *Phys. Rev. D* **87**, 055002 (2013); M. Gillioz, R. Gröber, C. Grojean, M. Mühlleitner, and E. Salvioni, *J. High Energy Phys.* **10** (2012) 004; M. Gouzevitch, A. Oliveira, J. Rojo, R. Rosenfeld, G. Salam, and V. Sanz, *J. High Energy Phys.* **07** (2013) 148; J. Cao, Z. Heng, L.

- Shang, P. Wan, and J. M. Yang, *J. High Energy Phys.* **04** (2013) 134.  
 [3] R. S. Gupta, H. Rzehak, and J. D. Wells, *Phys. Rev. D* **86**, 095001 (2012).  
 [4] CMS Collaboration, *J. Phys. G* **34**, 995 (2007).  
 [5] ATLAS Collaboration, Report No. CERN-LHCC-99-15, 1999.  
 [6] For other similar considerations, see, for example, M. Holthausen, K. S. Lim, and M. Lindner, *J. High Energy Phys.* **02** (2012) 037.  
 [7] J. D. Wells, [arXiv:0909.4541](https://arxiv.org/abs/0909.4541).

- [8] G. Degrandi, S. Di Vita, J. Elias-Miro, J. R. Espinosa, G. F. Giudice, G. Isidori, and A. Strumia, *J. High Energy Phys.* **08** (2012) 098.
- [9] M. Beneke *et al.*, [arXiv:hep-ph/0003033](https://arxiv.org/abs/hep-ph/0003033).
- [10] J. A. Aguilar-Saavedra *et al.* (ECFA/DESY LC Physics Working Group Collaboration), [arXiv:hep-ph/0106315](https://arxiv.org/abs/hep-ph/0106315).
- [11] M. E. Peskin, [arXiv:1207.2516](https://arxiv.org/abs/1207.2516).
- [12] K. Tobe and J. D. Wells, *Phys. Rev. D* **66**, 013010 (2002).
- [13] G. F. Giudice and A. Strumia, *Nucl. Phys.* **B858**, 63 (2012).
- [14] L. Linssen, A. Miyamoto, M. Stanitzki, and H. Weerts, [arXiv:1202.5940](https://arxiv.org/abs/1202.5940).
- [15] F. Goertz, A. Papaefstathiou, L. L. Yang, and J. Zurita, *J. High Energy Phys.* **06** (2013) 016.
- [16] U. Baur, T. Plehn, and D. L. Rainwater, *Phys. Rev. D* **67**, 033003 (2003); **68**, 033001 (2003).
- [17] L. Wang, W. Wang, J. M. Yang, and H. Zhang, *Phys. Rev. D* **76**, 017702 (2007).
- [18] R. Schabinger and J. D. Wells, *Phys. Rev. D* **72**, 093007 (2005).
- [19] M. Bowen, Y. Cui, and J. D. Wells, *J. High Energy Phys.* **03** (2007) 036.
- [20] N. Arkani-Hamed, A. G. Cohen, E. Katz, and A. E. Nelson, *J. High Energy Phys.* **07** (2002) 034.
- [21] K. Agashe, R. Contino, and A. Pomarol, *Nucl. Phys.* **B719**, 165 (2005).
- [22] G. F. Giudice, C. Grojean, A. Pomarol, and R. Rattazzi, *J. High Energy Phys.* **06** (2007) 045.
- [23] A. Manohar and H. Georgi, *Nucl. Phys.* **B234**, 189 (1984).
- [24] D. Pappadopulo, A. Thamm, and R. Torre, *J. High Energy Phys.* **07** (2013) 058.
- [25] C. Delaunay, C. Grojean, and J. D. Wells, *J. High Energy Phys.* **04** (2008) 029.
- [26] M. S. Carena, J. R. Espinosa, M. Quiros, and C. E. M. Wagner, *Phys. Lett. B* **355**, 209 (1995).
- [27] H. E. Haber and R. Hempfling, *Phys. Rev. D* **48**, 4280 (1993).
- [28] M. S. Carena, D. Garcia, U. Nierste, and C. E. M. Wagner, *Nucl. Phys.* **B577**, 88 (2000).
- [29] M. Drees and K. Hagiwara, *Phys. Rev. D* **42**, 1709 (1990).
- [30] J. Beringer *et al.*, *Phys. Rev. D* **86**, 010001 (2012); J. Erler and P. Langacker, <http://pdg.lbl.gov/2013/reviews/rpp2012-rev-standard-model.pdf>.
- [31] U. Ellwanger, C. Hugonie, and A. M. Teixeira, *Phys. Rep.* **496**, 1 (2010).
- [32] R. S. Gupta, M. Montull, and F. Riva, *J. High Energy Phys.* **04** (2013) 132.
- [33] For a description of how larger  $\lambda_5$  can be compatible with unification, see E. Hardy, J. March-Russell, and J. Unwin, *J. High Energy Phys.* **10** (2012) 072.
- [34] L. J. Hall, D. Pinner, and J. T. Ruderman, *J. High Energy Phys.* **04** (2012) 131.
- [35] V. D. Barger, M. S. Berger, A. L. Stange, and R. J. N. Phillips, *Phys. Rev. D* **45**, 4128 (1992).
- [36] W. Hollik and S. Penaranda, *Eur. Phys. J. C* **23**, 163 (2002).
- [37] K. E. Williams, H. Rzehak, and G. Weiglein, *Eur. Phys. J. C* **71**, 1669 (2011).
- [38] A. Dobado, M. J. Herrero, W. Hollik, and S. Penaranda, *Phys. Rev. D* **66**, 095016 (2002).