

HPLC-DAD-ESI-MS/MS Characterization of Bioactive Secondary Metabolites from *Strelitzia nicolai* Leaf Extracts and Their Antioxidant and Anticancer Activities *In vitro*

Mosad Ghareeb¹, Amal Saad¹, Wafaa Ahmed^{1,2}, Laila Refahy¹, Sami Nasr³

Departments of ¹Medicinal Chemistry and ³Biochemistry and Molecular Biology, Theodor Bilharz Research Institute, Giza, Egypt, ²Department of Chemistry, College of Science and Arts, Sajir, Shaqra University, Shaqra, Kingdom of Saudi Arabia


ABSTRACT

Background: *Strelitzia nicolai* Regel and Körn (Strelitziaceae) is native to Southern Africa whose phytochemistry and pharmacology were slightly investigated. **Materials and Methods:** In the current work, different solvent extracts of *S. nicolai* were screened for their chemical profiles through high-performance liquid chromatography coupled with diode array detection and electrospray ionization mass spectrometry (HPLC-DAD-ESI-MS/MS) analyses. Furthermore, their *in vitro* antioxidant, cytotoxic, and anticancer activities were evaluated using 2,2'-diphenyl-1-picrylhydrazyl radical (DPPH), 2,2'-azino-bis (3-ethylbenzothiazoline-6-sulphonic acid) (ABTS) & ferric reducing antioxidant power (FRAP) and crystal violet staining (CVS) colorimetric assays, respectively. **Results:** HPLC-DAD-ESI-MS/MS analyses led to the identification of nineteen and eleven phenolic compounds from the ethyl acetate and *n*-butanol extracts, respectively including flavonoids (e.g., quercetin 3-(2-G-rhamnosylrutinoside, quercetin, quercetin-3-O-glucoside, kaempferol-3,7-O-dirhamnoside, isorhamnetin-3-O-rutinoside and kaempferol-3-O-glucoside), phenolic acids derivatives (e.g., chlorogenic acid glycoside, protocatechuic acid-O-glucoside and caftaric acid), chalcones (e.g., xanthoangelol), and phenylethanoids (e.g., ligstroside glucoside). Moreover, in the DPPH assay the IC₅₀ value of the most active ethyl acetate extract was 20.49 µg/mL, relative to 2.92 µg/mL of ascorbic acid. ABTS and FRAP results reinforced the results of DPPH assay. According to the National Cancer Institute criteria, the tested extracts showed weak to moderate cytotoxic activities with IC₅₀ values ranged from 65.23 to 451.29 µg/mL. Furthermore, the EtOAc and *n*-BuOH extracts showed a noticeable anticancer activity with CVS spectroscopic readings for liver hepatocellular carcinoma growth 0.806 and 0.684 at a concentration (125 µg/mL), as well as 0.730 and 0.618 at concentration (500 µg/mL), respectively against control at 1.022. **Conclusion:** The obtained results reveal the high efficacy of the phenolic-rich extracts from *S. nicolai* as naturally occurring antioxidant and anti-tumor agents.

Key words: Anticancer activity, antioxidant activity, cytotoxicity, HPLC-DAD-ESI-MS/MS, polyphenolics, *Strelitzia nicolai*

SUMMARY

- The current research work evaluated the biological activities of different solvent extracts of *Strelitzia nicolai* including antioxidant, anticancer, and cytotoxic activities
- Among the tested extracts, the ethyl acetate and *n*-butanol extracts are the most promising extracts
- High-performance liquid chromatography coupled with diode array detection and electrospray ionization mass spectrometry analyses of the most active extracts led to the characterization of certain polyphenolic compounds, the majority are flavonoids and phenolic acids.



Abbreviations Used: HPLC-DAD-ESI-MS/MS: High-performance liquid chromatography-diode array detection-electrospray ionization-mass/mass; DPPH: 2,2'-Diphenyl-1-picrylhydrazyl radical; ABTS: 2,2'-Azino-bis (3-ethylbenzothiazoline-6-sulphonic acid); FRAP: Ferric reducing antioxidant power; TPTZ: Tripyridyl-s-triazine; FE: Ferrous equivalents; DMEM: Dulbecco's modified eagle's medium; DMSO: Dimethyl sulfoxide; EDTA: Ethylenediaminetetraacetic acid; PBS: Phosphate buffered saline; HepG-2: Liver hepatocellular carcinoma; CVS: Crystal violet stain; NCI: National cancer institute; Glu: Glucose; Rha: Rhamnose.

Correspondence:

Dr. Mosad Ghareeb,
Department of Medicinal Chemistry, Theodor
Bilharz Research Institute, Kornish El-Nile Street,
Warrak El-Hader, Imbaba, P. O. 12615, Giza, Egypt.
E-mail: m.ghareeb@tbri.gov.eg
DOI: 10.4103/pr.pr_89_18

Access this article online

Website: www.phcogres.com

Quick Response Code:



INTRODUCTION

Strelitziaceae is a tropical monocotyledonous ornamental family famous by its bioactive compounds namely phenalenones.^[1] *Strelitzia nicolai* Regel and Körn are usually known as the white bird of paradise tree.^[2] This plant is native to Southern Africa and broadly growing in many tropical regions around the world.^[3] To the best of our knowledge, there is limited information available in the literature about the phytochemical and biological investigations were reported on the plant. Bilirubin-IX

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

Cite this article as: Ghareeb M, Saad A, Ahmed W, Refahy L, Nasr S. HPLC-DAD-ESI-MS/MS characterization of bioactive secondary metabolites from *Strelitzia nicolai* leaf extracts and their antioxidant and anticancer activities *In vitro*. *Phcog Res* 2018;10:368-78.

is orange pigment with cyclic tetrapyrrole nucleus was isolated from the arils of *S. nicolai*.^[4-7] Moreover, it was reported that this pigment showed potent antioxidant and anticancer activities.^[7] In addition, the main chemical ingredients of the essential oil isolated from the seed arils of the plant were categorized as follows: amine, ethers, ketones, hydrocarbons, aromatic compounds, alcohols, amides, and esters.^[3] Moreover, accumulation of the reactive species in our bodies led to oxidative stress, which is associated with several disorders like cancer and cardiovascular diseases.^[8] Cancer is a well-known global problem which represents the second cause of death and accounts for “7–8 million deaths” worldwide.^[9] The treatment of cancer is based on the use of synthesized chemotherapeutic drugs, using this drugs as chemopreventive agents accompanied by a series of health problems due to their side effects and low safety, this concept encourage scientists to discover the role of medicinal plants in cancer therapeutic as natural sources of naturally occurring anticancer agents.^[9,10] Recently, there is increasing interest in the chemical investigation and characterization of the different class of secondary metabolites especially polyphenolic compounds to establish their structure-activity relationship.^[11] Since there no adequate information has been documented on the chemical and biological profile of the plant, therefore, the aim of the current study is to characterize the polyphenolic compounds of the different solvent extracts from *S. nicolai* leaves via high-performance liquid chromatography coupled with diode array detection and electrospray ionization mass spectrometry (HPLC-DAD-ESI-MS/MS) analysis as well as their antioxidant and anticancer activities.

MATERIALS AND METHODS

Plant material

The fresh leaves *S. nicolai* were collected from Zoo Garden, Giza, Egypt in June 2014. The identity of the plant was established by Dr. Tearse Labib, Botany Specialist, Department of Flora and Taxonomy, El-Orman Botanical Garden, Giza, Egypt. A voucher specimen (No. S25/5/6) was kept at the herbarium of the garden.

Extraction and fractionation

Dry powdered leaves of *S. nicolai* (1.5 kg), were extracted with MeOH in room temperature with shaking day by day followed by filtration and again extraction for 4 times. The extract was filtered using Whatman filter paper No. 1 and concentrated using Rotatory evaporator (Buchi, Switzerland) at (40°C ± 2°C). The crude extract was collected and stored at room temperature in the dark for the further process. The methanolic crude extract (230 g) was defatted by washing several times with petroleum ether (60–80°C), then undergoing fractionation process using organic solvents such as CH₂Cl₂; EtOAc; and *n*-BuOH (5 mL × 150 mL solvent).

High-performance liquid chromatography coupled with diode array detection and electrospray ionization mass spectrometry conditions

The phytochemical analysis of polyphenolic compounds was done using HPLC-photodiode array (PDA)-MS/MS. The liquid chromatography system was Thermofingan (Thermo Electron Corporation, USA) coupled with an LCQ Duo ion trap MS with an ESI source (ThermoQuest). The separation was achieved by using a C18 reversed-phase column (Zorbax Eclipse XDB-C18, Rapid resolution, 4.6 mm × 150 mm, 3.5 μm, Agilent, USA). A gradient of water and acetonitrile (with 1% formic acid each in the positive mode) was applied from 5% to 50% ACN in 60 min with flow rate 1 mL/min throughout the whole run. The samples were injected automatically using autosampler surveyor ThermoQuest. The instrument

was controlled by Xcalibur software (Thermo Fisher Scientific Inc., USA) to collect the ultraviolet (UV) chromatogram using PDA mode. The MS operated in the negative mode with a capillary voltage of – 10 V, a source temperature of 200°C, and high-purity nitrogen as a sheath and auxiliary gas at a flow rate of 80 and 40 (arbitrary units), respectively. The ions were detected in a full scan mode and mass range of 50–2000 *m/z*.

Antioxidant assays

2,2'-Diphenyl-1-picrylhydrazyl radical free radical-scavenging assay

2,2'-Diphenyl-1-picrylhydrazyl radical (DPPH) assay was performed according to the method described by Ghareeb *et al.* (2018). Briefly, 200 μl of plant extract, diluted appropriately in methanol in a concentration range from 0.24 to 500 μg/mL, was mixed with 100 μl of 0.2 mM DPPH in methanol in wells of 96-well plates. The plates were kept in the dark for 15 min; thereafter, the absorbance of the solution was measured at 515 nm in a Biochrom Asys UVM 340 Microplate Reader. Appropriate blanks, methanol, and standards (ascorbic acid solutions in methanol) were analyzed simultaneously. The scavenging activity (in %) was calculated using the following equation:

$$\text{DPPH scavenging (\%)} = 100 \times [(\text{Abs sample} + \text{DPPH}) - (\text{Abs sample blank})] / [(\text{Abs DPPH}) - (\text{Abs methanol})]$$

The IC₅₀ value is defined as the amount of extract needed to scavenge 50% of DPPH radicals. All analyses were performed in triplicate.^[12]

2,2'-Azino-bis (3-ethylbenzothiazoline-6-sulphonic acid) assay

The samples were dissolved in water to prepare the stock solutions (1 mg/mL) from which a radical-scavenging activity was determined by the 2,2'-azino-bis (3-ethylbenzothiazoline-6-sulphonic acid) (ABTS)⁺ radical cation decolorization assay,^[13] over a concentration range of 0.24–500 μg/mL. The ABTS cation radical was prepared by reacting 7 mM aqueous solution of ABTS (15 mL) with 140 mM potassium persulfate (264 μl). The mixture was allowed to stand in dark at room temperature for 16 h before use. Before assay, the ABTS working reagent was diluted with methanol to give an absorbance of 0.70 ± 0.02 at 734 nm and was equilibrated at room temperature. The reaction mixtures in the 96-well plates consisted of sample (50 μl) and the ABTS methanol working solution (100 μl). The mixture was stirred and left to stand for 10 min in dark, and then the absorbance was determined at 734 nm against a blank. All determinations were performed in triplicate. The scavenging activity (in %) was calculated as follows:

$$\% \text{ scavenging rate} = [1 - (A_1 - A_2) / A_0] \times 100.$$

Where A₀ is the absorbance of the control (without sample), and A₁ is the absorbance in the presence of the sample, A₂ is the absorbance of the sample without ABTS working solution. The scavenging activity of the samples was expressed as IC₅₀ value, which is the effective concentration, at which 50% of ABTS radicals were scavenged. Trolox was used as a standard.

Ferric reducing antioxidant power assay

The ferric reducing antioxidant power (FRAP) assay was carried out according to the previously reported procedure,^[14] with minor modifications. Each sample was dissolved in methanol to prepare the stock solution (1 mg/mL). Briefly, the working FRAP reagent was prepared freshly by mixing 300 mM acetate buffer (pH 3.6), a solution of 10 mM 2,4,6-tripyridyl-s-triazine (TPTZ) in 40 mM hydrochloric acid and 20 mM ferric chloride at 10:1:1 (v/v/v). 20 μl of each extract was mixed with 180 μl FRAP reagent in wells of 96-well plates. The mixture was then incubated for 6 min at 37°C, and the absorbance was measured at 595 nm in a microplate reader (Biochrom Asys UVM 340).

Appropriate blanks of plant extract and FRAP reagent lacking TPTZ (to correct the colors of the extracts) were run, together with quercetin (in methanol), and ferrous sulfate heptahydrate ($\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$) was used as a standard. FRAP activity was calculated as ferrous equivalents, the concentration of extract/quercetin which produced an absorbance value equal to that of 1 mM FeSO_4 .

Evaluation of cytotoxic activities

Splenocytes were isolated from normal albino mouse according to Goodman *et al.*^[15] beginning with washing thoroughly with 70% alcohol whole body and cervical dislocation done after anesthesia, and then the abdominal cavity was incised, and the spleen was transferred to sterile petri dish and slice into small pieces.^[16] Fragments were placed onto a strainer attached to a 50-mL conical tube. Excised spleen pieces were pressed through a strainer using a plunger end of a syringe and cells have been washed with phosphate buffered saline (PBS) (Dulbecco's PBS, pH = 7.4). Cell suspension centrifuged at 1600 rpm for 5 min. Cells pellets were resuspended in 2 mL lysing buffer (0.15 M NH_4Cl , 1 mM KHCO_3 , and 0.1 mM ethylenediaminetetraacetic acid). The cells have been incubated in a 37°C water bath for 2 min, and 30 mL of PBS were added and cells centrifuged at 1600 rpm for 5 min. Pellet cells were resuspended in 3 mL growth Dulbecco's Modified Eagle's Medium, Lonza, Belgium), 30% FBS (Hyclone, USA), 1% Penicillin/Streptomycin (Lonza, Belgium), and 1% L-glutamine (Lonza, Belgium)^[17,18] at final concentration of 2000×10000 cells per mL, the cells were cultured in 96 well plates. Cell count was performed and viability checked using trypan blue and a hemacytometer. Cells were incubated with serial dilutions of eleven fractions dissolved in dimethyl sulfoxide (DMSO) as follow (1000 $\mu\text{g}/\text{mL}$, 500 $\mu\text{g}/\text{mL}$, 250 $\mu\text{g}/\text{mL}$, 125 $\mu\text{g}/\text{mL}$, 62.5 $\mu\text{g}/\text{mL}$, and 31.2 $\mu\text{g}/\text{mL}$) at 37°C in 5% CO_2 and 90% humidified atmosphere for 36 h.

Evaluation of anticancer activities

Liver hepatocellular carcinoma (HepG-2) cell line obtained from the holding company for biological products and vaccines, Egypt (VACSERA) passage number 80 \approx 85 has been cultured in a T25 flask with Roswell Park Memorial Institute medium contains 10% fetal bovine serum, 100 U/mL penicillin and 100 $\mu\text{g}/\text{mL}$ streptomycin and incubated in a humidified 5% CO_2 incubator at 37°C.^[19] Culture medium was removed after reaching 90% confluency and 0.25% trypsin (Gibco/Invitrogen) was added and flask observed under an inverted microscope until all cells were detached. Cells were cultured in 96-well plates at a density of 10000 cells per well. Incubation has been done for 48 h until sheet was observed. Serial dilutions of fractions in DMSO were added to cultured cells at concentrations of 1000 $\mu\text{g}/\text{mL}$, 500 $\mu\text{g}/\text{mL}$, 250 $\mu\text{g}/\text{mL}$, 125 $\mu\text{g}/\text{mL}$, 62.5 $\mu\text{g}/\text{mL}$, and 31.2 $\mu\text{g}/\text{mL}$ and incubated for 24 h. Media were decanted and plates were washed carefully with PBS (Dulbecco's PBS, pH = 7.4) and cells were stained with 20 μl of 0.5% crystal violet (Sigma-Aldrich Corp., St. Louis, MO, USA) in 30% ethanol for 10 min at room temperature. Plate wells were washed 3 times using distilled water^[20] and the absorbance was measured at optical density = 490 nm.

Statistical analysis and dose-response curve

A dose-response curve has been traced to find the equation and estimate IC_{50} using unpaired Student's *t*-test excel, referring to different concentrations, more than three to draw the curve which presented as mean standard deviation (SD) taking in consideration $P < 0.05$.

RESULTS AND DISCUSSION

Antioxidant activity

Different solvent extracts of *S. nicolai* were evaluated for their antioxidant activities using three techniques including; DPPH, ABTS, and FRAP. In DPPH assay, the IC_{50} values for the tested extracts ranged from 20.49 to 118.17 $\mu\text{g}/\text{mL}$ compared to ascorbic acid as standard with IC_{50} equal to 2.92 $\mu\text{g}/\text{mL}$. The results are in the order: EtOAc > n-BuOH > MeOH > H_2O > CH_2Cl_2 > Pet. ether. In addition, all tested extracts showed similar activity using ABTS assay with IC_{50} values arranged in the following order: EtOAc (9.18) > n-BuOH (12.43) > MeOH (15.73) > H_2O (23.18) > CH_2Cl_2 (29.89) > Pet. ether extracts (52.25) ($\mu\text{g}/\text{mL}$) compared to Trolox (IC_{50} = 1.63 $\mu\text{g}/\text{mL}$). In FRAP assay, the EtOAc fraction showed high reducing power activity with 22.19 mM FeSO_4 equivalent/mg extract, followed by n-BuOH (18.34), MeOH (15.46), H_2O (11.39), CH_2Cl_2 (7.92), and Pet. ether extracts (1.58), respectively, compared to quercetin (21.45). Results are documented in Table 1. In conclusion, the results of the three assays agree with each other and the high antioxidant activity of the EtOAc and n-BuOH extracts may be due to the presence of the tentatively identified polyphenolic compounds which are broadly reported in the literature by their antioxidant potential for instance; flavonoids (e.g. quercetin 3-(2-G-rhamnosylrutinoside), rutin, quercetin-3-O-glucoside, kaempferol-3,7-O-dirhamnoside, isorhamnetin-3-O-rutinoside, kaempferol-3-O-rutinoside, and kaempferol-3-O-glucoside)^[21,22] and phenolic acids (e.g. chlorogenic acid glycoside, sinapaldehyde, and caftaric acid).^[23,24]

In vitro cytotoxic and anticancer activities of different solvent extracts

Different solvent extracts of *S. nicolai* have been evaluated for their *in vitro* cytotoxic potential to murine spleen cells by visual counting for the vital cells after growing under the effect of serial dilutions of these extracts. The results revealed that methanol (1) and *n*-butanol (5) extracts showed very rare cytotoxic activities with IC_{50} values of 451.29 and 382.38 $\mu\text{g}/\text{mL}$, respectively. While pet. ether (2), dichloromethane (3), and ethyl acetate (4) extracts showed moderate cytotoxic effects with IC_{50} values of 68.03, 90.29, and 65.23 $\mu\text{g}/\text{mL}$, respectively [Table 2 and Figure 1], indicating that flagging cytotoxic effects against murine spleen cell have been observed. Moreover, the National Cancer Institute instructions demonstrate the criteria and the conditions of cytotoxic activity for a chemical complexes as the concentration of an inhibitor that is required for 50% inhibition of cells growth.^[25] IC_{50} where the values ≤ 20 $\mu\text{g}/\text{mL}$, is considered to be potentially cytotoxic, while IC_{50} values 21-100 $\mu\text{g}/\text{mL}$ = moderately cytotoxic, IC_{50} 101-200 $\mu\text{g}/\text{mL}$ = weakly cytotoxic and $\text{IC}_{50} > 501$ $\mu\text{g}/\text{mL}$ = no cytotoxic effect.^[26,27] On the other hand, significant anticancer effects against

Table 1: Antioxidant activities of different extracts of *S. nicolai*

Sample	DPPH (IC_{50} $\mu\text{g}/\text{mL}$)	ABTS (IC_{50} $\mu\text{g}/\text{mL}$)	FRAP (mM FeSO_4 equivalent/mg extract)
MeOH	24.89 \pm 2.73	15.73 \pm 1.22	15.46 \pm 0.81
Petroleum ether	118.17 \pm 3.39	52.25 \pm 1.35	1.58 \pm 1.47
CH_2Cl_2	65.28 \pm 5.14	29.89 \pm 0.56	7.92 \pm 0.49
EtOAc	20.49 \pm 1.59	9.18 \pm 0.73	22.19 \pm 1.56
<i>n</i> -BuOH	22.09 \pm 1.88	12.43 \pm 0.78	18.34 \pm 1.37
H_2O	56.70 \pm 5.06	23.18 \pm 1.18	11.39 \pm 0.67
Ascorbic acid	2.92 \pm 0.29	-	-
Quercetin	-	-	21.45 \pm 2.55
Trolox	-	1.63 \pm 0.46	-

Results are average of triplicate measurements ($n=3$) and expressed as mean \pm SD; IC_{50} values were calculated using a four-parameter logistic curve (SigmaPlot® 11.0). SD: Standard deviation; ABTS: Radical-scavenging activity; DPPH: Free radical-scavenging activity; FRAP: Ferric reducing antioxidant power

Table 2: Different solvent extracts of *S. nicolai* showing the variation in cytotoxic effect (IC₅₀) on murine spleen cells by visual counting under the inverted microscope

Concentration µg/ml	Extracts	1	2	3	4	5
1000	Concentration 1	390				210
500	Concentration 2	420				300
250	Concentration 3	510				450
125	Concentration 4	450	210	240	150	330
62.5	Concentration 5	450	300	330	300	450
31.2	Concentration 6		360	420	390	510
Control		600	600	600	600	600
-	Equation	$y = -0.0832x + 476.25$	$y = -1.5765x + 404.92$	$y = -1.8507x + 464.92$	$y = -2.536x + 464.88$	$y = -0.2627x + 461.19$
Results	Cytotoxicity IC ₅₀ µg/ml	451.29	68.03	90.29	65.23	382.38
-		Rare	Moderate	Moderate	Moderate	Rare

^ay: Is the concentration equivalent to IC₅₀ per µg and (x) is the control IC₅₀ (300); 1: MeOH; 2: Petroleum ether; 3: CH₂Cl₂; 4: EtOAc and 5: *n*-BuOH

Table 3: Crystal violet staining and colorimetric assay at optical density=490 nm for human hepatoma cell line under the effect of *S. nicolai* extracts showing cytotoxicity profile

Extract	CVS spectroscopic reading for HepG2 growth	
	Complex concentration (125 µg/mL)	Complex concentration (500 µg/mL)
1	0.883	0.684
2	0.931	0.611
3	0.753	0.702
4	0.806	0.730
5	0.684	0.618
Control	1.022	

Control: No added fraction. CVS: Crystal violet stain; HepG-2: Human hepatoma

HepG-2 cell line after 24 h exposure were obvious using colorimetric assay. The results revealed that the crystal violet stain spectroscopic readings for HepG-2 growth at complex concentration (125 µg/mL) were 0.883, 0.931, 0.753, 0.806, and 0.684, while at complex concentration (500 µg/mL) were 0.684, 0.611, 0.702, 0.730, and 0.618, respectively for MeOH, pet. ether, dichloromethane, ethyl acetate and *n*-butanol extracts against control at 1.022 [Table 3]. Results showed an intimate relationship between anticancer activity and concentration. In general, the anticancer activity of the polyphenolic-rich extracts may be due to the presence of such phenolic compounds.^[28-32] Moreover, these polyphenolic compounds are capable of inhibiting cancer cells through several modes of actions.^[26]

Characterization of the phenolic compounds in *Strelitzia nicolai*

The ethyl acetate and *n*-butanol extracts of *S. nicolai* were investigated for their polyphenolic constituents using HPLC-DAD-ESI-MS/MS technique. Nineteen compounds were detected and tentatively identified in the ethyl acetate extract were categorized as phenolic acids, flavonoids (glycosides and aglycones), chalcones, and other nucleus. On the other hand, eleven compounds were detected and tentatively identified in the *n*-butanol extract were categorized as phenolic acids and flavonoids (glycosides and aglycones). The identification of the phenolic compounds was based on comparing their fragmentation pattern using negative ion ionization mode with the available data in the literature. The chemical structures of some selected compounds and their full MS/MSⁿ pattern will be mentioned below Tables 4, 5 and Figures 2-7.

Phenolic compounds were detected in the EtOAc extract

Phenolic acids and their derivatives

Compound 1 was detected at *Rt* = 1.43 min, it showed a deprotonated ion [M-H]⁻ at *m/z* 515, it also generated a fragment ion as a base peak




Figure 1: Cytotoxic effect of different solvent extracts of *S. nicolai* on murine splenocytes after 36 h incubation

at *m/z* 341 corresponding to the loss of quinic acid moiety ($-m/z$ 174 u) [M-H-quinic acid moiety]⁻, further loss of glucosyl moiety ($-m/z$ 162 u) was confirmed by the appearance of a fragment ion at *m/z* 179 [M-H-quinic acid-glucose moiety]⁻. Therefore, compound 1 could be identified as chlorogenic acid glucoside as previously described.^[33] Compound 16 was detected at *Rt* = 28.92 min, it showed a deprotonated ion as a base peak [M-H]⁻ at *m/z* 177, it also generated a fragment ion at *m/z* 162 corresponding to the loss of methyl moiety ($-m/z$ 15) [M-H-CH₃]⁻, further loss of hydroxyl moiety was confirmed by the appearance of a fragment ion at *m/z* 145 [M-H-CH₃-OH]⁻, further neutral loss of CO moiety of the aldehyde group was confirmed by the appearance of a fragment ion at *m/z* 117 ($-m/z$ 28 u) [M-H-CH₃-OH-CO]⁻. Therefore, compound 16 could be identified as 3-(4-hydroxy-3-methoxyphenyl) prop-2-enal (coniferyl aldehyde).^[34]

Compound 17 was detected at *Rt* = 30.57 min, it showed a deprotonated ion [M-H]⁻ at *m/z* 207, it also generated a fragment ion as a base peak at *m/z* 192 corresponding to the presence of sinapoyl moiety and due to the loss of methyl moiety ($-m/z$ 15 u) [M-H-CH₃]⁻, further loss of methyl moiety ($-m/z$ 15 u) was confirmed through the fragment ion at *m/z* 177 [M-H-2CH₃]⁻. Thus, compound 17 could be identified as 3,5-dimethoxy-4-hydroxycinnamaldehyde (sinapaldehyde).^[35] Compound 19 was detected at *Rt* = 40.70 min,

it showed a deprotonated molecular ion $[M-H]^-$ at m/z 311, it also generated a fragment ion at m/z 179 corresponding to the loss of tartaric moiety ($-m/z$ 132 u) $[M-H-tartaric]^-$, in addition to a fragment ion was appeared at m/z 149 corresponding to the loss of caffeoyl moiety ($-m/z$ 162 u) $[M-H-caffeoyl]^-$. This fragmentation pattern was typically assigned to caftaric acid (*cis*-caffeoyl tartaric acid).^[35,36]




Figure 2: Negative HPLC-DAD-ESI-MS/MS profiles of phenolic compounds were detected in the EtOAc extract of *S. nicolai* leaves

Flavonoids

Compound 3 was detected at $R_t = 14.66$ min, it showed a deprotonated molecule $[M-H]^-$ at m/z 755, a MS/MS fragment was observed at m/z 609 was assigned to rutin molecule after the neutral loss of one rhamnosyl moiety ($-m/z$ 146 u) $[M-H-rhamnose\ moiety]^-$, a fragment ion was detected at m/z 591 due to further loss of water molecule ($-m/z$ 18 u) $[M-H-rhamnose\ moiety-H_2O]^-$, another fragment ion was also appeared at m/z 445 due to the loss of the second rhamnosyl moiety ($-m/z$ 146 u) $[M-H-2rhamnose\ moiety-H_2O]^-$, a diagnostic fragment ion was observed at m/z 301 was assigned to quercetin a glycone and can be explained by the release of the glucose moiety ($-m/z$ 162 u) $[M-H-2rhamnose\ moieties-glucose\ moiety]^-$, another key fragment ions of quercetin a glycone were observed at m/z 271 and 255. Therefore, compound 3 was identified as quercetin 3-(2 G-rhamnosylrutinoside) in comparison with the previously published data.^[37] Compound 5 was detected at $R_t = 16.18$ min, it displayed a deprotonated molecule $[M-H]^-$ at m/z 609, the neutral loss of rhamnosyl moiety ($-m/z$ 146 u) afford a fragment ion at m/z 463 $[M-H-rhamnose\ moiety]^-$, further loss of glucosyl moiety ($-m/z$ 162 u) was confirmed through a diagnostic fragment ion at m/z 301 $[M-H-rhamnosyl-glucosyl]^-$, which was assigned to quercetin a glycone and other key fragment ions of quercetin a glycone were detected at m/z 271, 255 and 179. Therefore, compound 5 was identified as quercetin-3-O- α -L-rhamnopyranosyl-(1 \rightarrow 6)- β -D-glucopyranose (rutin).^[36] Compound 6 was detected at $R_t = 17.18$ min; it displayed a deprotonated molecule as a base peak at m/z 301, which was assigned to quercetin a

Table 4: Phenolic compounds tentatively identified in the EtOAc extract of *S. nicolai* leaves by HPLC-DAD-ESI-MS/MS in negative ion mode

Peak number	R_t	MW	$[M-H]^-$ (m/z)	MS/MS fragments*	Proposed compounds	References
1	1.43	516	515	341, 179, 173	Chlorogenic acid glycoside	Abu-Reidah <i>et al.</i> (2013a)
2	8.57	454	453	407	Unknown	-
3	14.66	756	755	737, 609, 591, 489, 445, 301, 300, 271, 255	Quercetin 3-(2 G-rhamnosylrutinoside)	Abu-Reidah <i>et al.</i> (2013b)
4	15.78	612	611	491, 447, 303, 302, 300, 273, 255	Unknown	-
5	16.18	610	609	463, 300, 301, 271, 255, 179	Rutin	Abu-Reidah <i>et al.</i> (2015)
6	17.18	301	300	301, 300, 271, 269, 255, 229, 179, 169, 151	Quercetin	Abu-Reidah <i>et al.</i> (2014)
7	18.12	464	463	301, 271, 255, 179, 151	Quercetin-3-O-glucoside	Bravo <i>et al.</i> (2007); Abu-Reidah <i>et al.</i> (2012)
8	19.29	578	577	431, 285	Kaempferol-3,7-O-dirhamnoside (Kaempferitrin)	Abu-Reidah <i>et al.</i> (2012)
9	19.66	624	623	459, 315, 300, 271, 255, 179	Isorhamnetin-3-O-rutinoside	Abu-Reidah <i>et al.</i> (2012)
10	20.18	594	593	473, 447, 429, 285, 255, 227, 151		Al-Rawahi <i>et al.</i> (2014)
11	20.75	448	447	285, 284, 255, 227, 151	Kaempferol-3-O-glucoside	Bravo <i>et al.</i> (2007); Chen <i>et al.</i> (2014)
12	21.19	564	563	473, 443, 413, 383, 353, 285, 284	kaempferol-3-O- β -xylopyranosyl-(1 \rightarrow 3)- α -L-rhamnopyranosyl	NA
13	21.98	392	391	376, 359, 239, 161, 151, 135	Xanthoangelol	Kim <i>et al.</i> (2014)
14	24.35	462	461	446, 341, 299	Isoorientin-3'-O-methyl ether (Isoscoparin)	Abu-Reidah <i>et al.</i> (2012)
15	28.85	446	445	269, 193, 176, 175, 160	Unknown	-
16	28.92	178	177	162, 145, 117	3-(4-hydroxy-3-methoxyphenyl) prop-2-enal (coniferyl aldehyde)	Sanz <i>et al.</i> (2012)
17	30.57	208	207	192	3,5-dimethoxy-4 hydroxycinnamaldehyde (Sinapaldehyde)	Sanz <i>et al.</i> (2012)
18	32.25	328	327	309, 291, 239, 229, 221, 211, 171	1,7-bis-(3,4-dihydroxyphenyl)-4-hepten-3-one (hirsutenone)	Riethmller <i>et al.</i> (2013)
19	38.57	312	311	179, 149	caftaric acid	Chen <i>et al.</i> (2012); Abu-Reidah <i>et al.</i> (2015)
20	41.29	330	329	314, 300, 299, 285, 243	Quercetin-dimethyl-ether	Pellati <i>et al.</i> (2011)
21	42.05	315	314	300, 299, 285, 271	Unknown	-
22	44.84	314	313	298, 283	4',7'-dimethoxy luteolin (4',5-dihydroxy-3',7-dimethoxyflavone)	Simirgiotis <i>et al.</i> (2015)
23	49.38	344	343	328, 313, 300, 283, 181	Dihydroxy trimethoxy flavonol isomer	Abdel-Hameed <i>et al.</i> (2014)
24	51.05	522	521	505, 493, 453, 433, 398, 372, 303	Unknown	-

*Bold items referred to the main aglycones fragments. NA: Not available; MS: Mass spectrometer

Table 5: Phenolic compounds tentatively identified in the *n*-butanol extract of *S. nicolai* leaves by HPLC-DAD-ESI-MS/MS in negative ion mode

Peak number	R_t	MW	[M-H] ⁻ (m/z)	MS/MS fragments*	Proposed compounds	References
1	1.47	316	315	225, 163, 153 , 152, 109, 108	Protocatechuic acid- <i>O</i> -glucoside	Abu-Reidah <i>et al.</i> (2015)
2	12.58	576	577	503, 474, 473 , 383, 353	Unknown	-
3	14.70	624	623	608, 477, 315 , 300, 271, 255, 279	Isorhamnetin 3- <i>O</i> -rutinoside	Abu-Reidah <i>et al.</i> (2012)
4	15.04	756	755	737, 609, 591, 489, 445, 301, 300 , 271, 255	Quercetin 3-(2- <i>G</i> -rhamnosylrutinoside)	Abu-Reidah <i>et al.</i> (2013a)
5	15.80	740	739	593, 575 , 473, 429, 393, 285, 284, 257, 255, 227	Kaempferol 3- <i>O</i> -rutinoside-7- <i>O</i> -rhamnoside	Ágnes, 2013
6	16.61	770	769	605	Unknown	-
7	17.2	612	611	593, 374, 344, 301 , 271, 255, 179	Unknown	-
8	18.18	464	463	301 , 271, 255, 179, 151	Quercetin-3- <i>O</i> -glucoside	Bravo <i>et al.</i> (2007); Abu-Reidah <i>et al.</i> (2012); Al-Rawahi <i>et al.</i> (2014)
9	19.95	594	593	447, 429, 285 , 255, 227, 179, 169	Kaempferol 3- <i>O</i> -rutinoside	Al-Rawahi <i>et al.</i> (2014)
10	20.83	564	563	473 , 443, 413, 383, 353, 285, 284	Unknown	-
11	21.80	448	447	285, 284 , 255, 227, 151	Kaempferol-3- <i>O</i> -glucoside	Ágnes, 2013; Bravo <i>et al.</i> (2007); Chen <i>et al.</i> (2014)
12	24.84	610	609	463, 445, 300 , 301, 271, 255, 179	Quercetin-3- <i>O</i> -rutinoside (Rutin)	Abu-Reidah <i>et al.</i> (2015)
13	32.20	686	685	523 , 505, 477, 343	Ligstroside glucoside	Sanz <i>et al.</i> , 2012
14	42.70	312	311	179, 149	Monocaffeoyltartaric acid	Chen <i>et al.</i> (2012); Abu-Reidah <i>et al.</i> (2015)
15	42.90	330	329	314 , 285	Quercetin-dimethyl-ether	Pellati <i>et al.</i> (2011)

*Bold items referred to the main aglycones fragments. NA: Not available; MS: Mass spectrometer

glycone and other characteristic key fragment ions were appeared at m/z 271, 269, 255, 229, 179, and 151. Therefore, compound 6 was identified as 5,7,3',4'-flavon-3-ol (quercetin).^[38] Compound 7 was detected at $R_t = 18.12$ min, it displayed a deprotonated molecule [M-H]⁻ at m/z 463, the neutral loss of glucosyl moiety ($-m/z$ 162 u) was confirmed through a fragment ion as a base peak at m/z 301 [M-H-glucose]⁻ was assigned to quercetin a glycone. Other key fragment ions were also observed at m/z 271, 255, 179, and 151. This fragmentation pattern was typically assigned to quercetin-3-*O*- β -D-glucoside (isoquercetin).^[39-41] Compound 8 was detected at $R_t = 19.29$ min, it displayed a deprotonated molecule [M-H]⁻ at m/z 577, the neutral loss of rhamnosyl moiety ($-m/z$ 146 u) give a fragment ion as a base peak at m/z 431 [M-H-rhamnosyl moiety]⁻ was assigned to kaempferol-*O*-rhamnoside, further neutral loss of the second rhamnosyl moiety ($-m/z$ 146 u) was confirmed through the appearance of a fragment ion at m/z 285 which was attributed to kaempferol a glycone. Accordingly, compound 8 was identified as kaempferol-3,7-*O*-dirhamnoside (kaempferitrin).^[40] Compound 9 was detected at $R_t = 19.66$ min, it displayed a deprotonated molecule [M-H]⁻ at m/z 623, the neutral loss of methyl moiety ($-m/z$ 15 u) give a fragment ion at m/z 608 [M-H-CH₃]⁻, while the neutral loss of rhamnosyl moiety ($-m/z$ 146 u) give a fragment ion at m/z 477 [M-H-Rha]⁻, the loss of glucose moiety ($-m/z$ 162 u) was confirmed through a diagnostic fragment ion as a base peak at m/z 315 [M-H-rutinoside]⁻, which was assigned to isorhamnetin a glycone. Therefore, compound 9 was identified as isorhamnetin-3-*O*-rutinoside.^[40]

Compound 10 was detected at $R_t = 20.18$ min, it displayed a deprotonated molecule [M-H]⁻ at m/z 593, the neutral loss of rhamnosyl moiety ($-m/z$ 146 u) give a fragment ion [M-H-Rha]⁻ at m/z 447 was assigned to kaempferol glucoside, further loss of the glucosyl moiety ($-m/z$ 162 u) was confirmed through the appearance of a fragment ion as a base peak at m/z 285 [M-H-rhamnosyl-glucosyl]⁻ was assigned to kaempferol a glycone and due to total loss of rutinoside moiety. Therefore, compound 10 was identified as kaempferol-3-*O*-rutinoside.^[41] Compound 11 was detected at $R_t = 20.75$ min, it displayed a deprotonated molecule [M-H]⁻ at m/z 447 was assigned to kaempferol glucoside, the neutral loss of glucosyl moiety ($-m/z$ 162 u) give a fragment ion at m/z 285 [M-H-glucosyl]⁻ was assigned to kaempferol a glycone, and other key fragment ions were detected at m/z 255, 227, 179 and 151. Compound 11 could be identified as kaempferol-3-*O*-glucoside.^[32,39]

Compound 12 was detected at $R_t = 21.19$ min, it displayed a deprotonated molecule [M-H]⁻ at m/z 563, the neutral loss of rhamnosyl moiety ($-m/z$ 146 u) give a fragment ion at m/z 417 [M-H-rhamnosyl]⁻, further loss of xylosyl moiety ($-m/z$ 132) [M-H-rhamnosyl-xylosyl]⁻ was confirmed through the appearance of a diagnostic fragment ion at m/z 285 was assigned to kaempferol a glycone. Thus, compound 12 was tentatively identified as kaempferol-3-*O*- β -xylopyranosyl-(1 \rightarrow 3)- α -L-rhamnopyranosyl. Compound 14 was detected at $R_t = 22.67$ min, it displayed a deprotonated molecule [M-H]⁻ at m/z 461, the loss of methyl moiety ($-m/z$ 15 u) give a fragment ion at m/z 446 [M-H-CH₃]⁻, while neutral loss of glucosyl moiety ($-m/z$ 162 u) give a fragment ion as a base peak at m/z 299 [M-H-glu]⁻ was assigned to tri-hydroxy flavone nucleus. This fragmentation pattern was typically assigned to isoorientin-3'-*O*-methyl ether (isoscoparin).^[42] Compound 20 was detected at $R_t = 42.05$ min, it displayed a deprotonated molecule [M-H]⁻ at m/z 329, the loss of methyl moiety ($-m/z$ 15 u) give a fragment ion as a base peak at m/z 314 [M-H-CH₃]⁻, further neutral loss of methyl moiety ($-m/z$ 15 u) give a fragment ion at m/z 299 [M-H-2CH₃]⁻, while the neutral loss of CO₂ moiety ($-m/z$ 44 u) give a fragment ion at m/z 285 [M-H-CO₂]⁻, and another ion ($-m/z$ 42 u) at m/z 243 [M-H-C₂H₂O-CO₂]⁻, thus compound 20 could be identified as quercetin-dimethyl-ether.^[43]

Compound 22 was detected at $R_t = 49.38$ min, the MS/Ms spectrum displayed a deprotonated molecular ion [M-H]⁻ at m/z 313, while the loss of methyl moiety ($-m/z$ 15 u) give a fragment ion as a base peak at m/z 298 [M-H-CH₃]⁻, further neutral loss of methyl moiety ($-m/z$ 15 u) give a fragment ion at m/z 283 [M-H-2CH₃]⁻, therefore, compound 22 was identified as 3',7-dimethoxyluteolin (4',5-dihydroxy-3',7-dimethoxyflavone).^[44] Compound 23 was detected at $R_t = 51.05$ min, the MS/Ms spectrum displayed a deprotonated molecular ion [M-H]⁻ at m/z 343, the loss of methyl moiety ($-m/z$ 15 u) give a fragment ion as a base peak at m/z 328 [M-H-CH₃]⁻, further neutral loss of methyl moiety ($-m/z$ 15 u) give a fragment ion at m/z 313 [M-H-2CH₃]⁻, other fragment ions were observed at m/z 297 and 285. Therefore, compound 23 was identified as dihydroxy-trimethoxy flavanol.^[45]

Chalcones

Compound 13 was detected at $R_t = 21.98$ min, it displayed a deprotonated molecule [M-H]⁻ at m/z 391. The neutral loss of methyl moiety ($-m/z$ 15 u) give a fragment ion as a base peak at m/z 376 [M-H-CH₃]⁻ was assigned




Figure 3: Mass spectrometry spectra and postulated fragmentation pattern of some selected phenolic compounds detected in the EtOAc extract of *S. nicolai* leaves using HPLC-DAD-ESI-MS/MS in negative ionization mode; (a) quercetin [M-H]⁻ m/z = 301; (b) quercetin-3-O-glucoside [M-H]⁻ m/z = 463; (c) rutin [M-H]⁻ m/z = 609; (d) isohammetin-3-O-rutinoside [M-H]⁻ m/z = 623; (e) kaempferol-3-O-glucoside [M-H]⁻ m/z = 447; and (f) kaempferol-3-O-rutinoside [M-H]⁻ m/z = 593

to methylated chalcone, the appearance of a fragment ion at m/z 135 was assigned to *p*-hydroxycinnamic acid moiety and due to the loss of 2,6-octa-dienyll-4-hydroxyphenyl moiety. Therefore, compound 13 was tentatively identified as 3'-(3,7-dimethyl-2,6-Octadienyl)-2'-4,4'-trihydroxy chalcone.^[46]

Other detected compounds

Compound 18 was detected at R_t = 34.23 min, it displayed a deprotonated molecular ion [M-H]⁻ at m/z 327, the neutral loss of water moiety ($-m/z$ 18 u) give a fragment ion at m/z 309 [M-H-H₂O]⁻, the appearance of another fragment ion at m/z 291 due to further loss of another water moiety [M-H-2H₂O]⁻, other fragment ions were observed at m/z 239, 229, 221, 211 and 171. Therefore, compound 18 was tentatively identified

as 1,7-bis-(3,4-dihydroxyphenyl)-4-hepten-3-one (hirsutenone) in comparison with the previously published data.^[47]

Phenolic compounds were detected in the n-BuOH extract

Phenolic acids and their derivatives

Compound 1 was detected at R_t = 2.76 min, it showed a deprotonated molecular ion [M-H]⁻ at m/z 315, a diagnostic fragment ion as a base peak was detected at m/z 153 due to the loss of glucosyl moiety ($-m/z$ 162 u) and corresponding to protocatechuic acid [M-H-glucosyl moiety]⁻, further loss of CO₂ moiety ($-m/z$ 44 u) was confirmed through the appearance of fragment ion at m/z 109 [M-H-glucosyl moiety-CO₂]⁻.




Figure 4: A proposed fragmentation pattern of compound 5 (rutin) and compound 10 (Kaempferol-3-O-rutinoside)

Therefore, compound 1 could be identified as protocatechuic acid glucoside.^[36] Compound 14 was detected at $R_t = 42.08$ min, it showed a deprotonated molecular ion $[M-H]^-$ at m/z 311, it also generated a fragment ion at m/z 179 corresponding to the loss of tartaric moiety ($-m/z$ 132) $[M-H\text{-tartaric moiety}]^-$, another fragment ion was appeared at m/z 149 due to the loss of caffeoyl moiety ($-m/z$ 162) $[M-H\text{-caffeoyl moiety}]^-$, this fragmentation profile was assigned to moncaffeoyltartaric acid (caftaric).^[35,36]

Flavonoids

Compound 3 was detected at $R_t = 14.70$ min, it showed a deprotonated molecular ion $[M-H]^-$ at m/z 623, it also generated a fragment ion at m/z 608 corresponding to the loss of methyl moiety ($-m/z$ 15 u) $[M-H\text{-CH}_3]^-$, another fragment ion was observed at m/z 477 refer to the loss of rhamnosyl moiety ($-m/z$ 146 u) $[M-H\text{-rhamnosyl moiety}]^-$, it also generated a diagnostic fragment ion as a base peak at m/z 315 refer to further loss

of glucosyl moiety ($-m/z$ 162 u), which may represent isorhamnetin a glycone [M-H-rhamnosyl moiety-glucosyl moiety] $^-$, another fragment ion was detected at m/z 300 due to the loss of methyl moiety from the a glycone nucleus [M-H-rhamnosyl moiety-glucosyl moiety-CH₃] $^-$, this fragmentation was typically assigned to isorhamnetin 3-O-rutinoside.^[40] Compound 4 was detected at $R_t = 15.04$ min, it showed a deprotonated molecule [M-H] $^-$ at m/z 755, a MS/MS fragment was appeared at m/z 609 corresponding to rutin molecule after the neutral loss of one rhamnosyl

moiety ($-m/z$ 146 u) [M-H-rhamnose moiety] $^-$, a fragmentation was observed at m/z 445 due to the loss of another rhamnosyl and water moieties ($-m/z$ 146-18 u) [M-H-2 rhamnose moiety-H₂O] $^-$, a diagnostic fragment ion was detected at m/z 301 was assigned to quercetin a glycone and can be explained by the release of the glucose moiety ($-m/z$ 162 u) [M-H-2 rhamnose moieties-glucose moiety] $^-$, in addition to the appearance of key fragment ions of quercetin a glycone at m/z 271, and 255. Therefore, compound 4 was identified as quercetin 3-(2 G-rhamnosylrutinoside).^[37]

Compound 5 was detected at $R_t = 15.80$ min, it showed a deprotonated molecule [M-H] $^-$ at m/z 739, a MS/MS fragment ion was observed at m/z 593 due to the loss of rhamnosyl moiety ($-m/z$ 146 u) [M-H-rhamnose moiety] $^-$, further neutral loss of water molecule ($-m/z$ 18 u) led to generation of a fragment ion at m/z 575 [M-H-rhamnose moiety-H₂O] $^-$, further neutral loss of another rhamnosyl moiety ($-m/z$ 146 u) led to generation of a fragment ion at m/z 429 [M-H-2rhamnose moiety-H₂O] $^-$, the appearance of fragment ion at m/z 285 was accounted for neutral loss of glucosyl moiety ($-m/z$ 162 u) [M-H-2 rhamnose moiety-glucosyl moiety] $^-$ it was assigned to kaempferol a glycone with key fragment ions at m/z 257, 255, and 227. Therefore, compound 5 was identified as kaempferol 3-O-rutinoside-7-O-rhamnoside.^[41] Compound 8 was detected at $R_t = 18.18$ min, it showed a deprotonated molecule [M-H] $^-$ at m/z 463, a diagnostic MS/MS fragment ion as a base peak was detected at m/z 301 due to the loss of glucosyl moiety ($-m/z$ 162 u) [M-H-glucosyl moiety] $^-$ it was assigned to quercetin a glycone, another key a glycone fragments were appeared at m/z 271, 255, 179 and 151. Thus compound




Figure 5: Negative HPLC-DAD-ESI-MS/MS profiles of phenolic compounds were detected in the EtOAc extract of *S. nicolai* leaves





Figure 6: Mass spectrometry spectra and postulated fragmentation pattern of some selected phenolic compounds detected in the *n*-BuOH extract of *S. nicolai* leaves using HPLC-DAD-ESI-MS/MS in negative ionization mode; (a) protocatechuic acid-O-glucoside [M-H] $^-$ $m/z = 315$; (b) quercetin 3-(2 G-rhamnosylrutinoside) [M-H] $^-$ $m/z = 755$; (c) kaempferol 3-O-rutinoside-7-O-rhamnoside [M-H] $^-$ $m/z = 739$, and (d) ligstroside glucoside [M-H] $^-$ $m/z = 685$



8 could be characterized as quercetin-3-O- β -D-glucoside.^[39-41]

Compound 9 was detected at $R_t = 19.95$ min, it showed a deprotonated molecule $[M-H]^-$ at m/z 593, a fragment ion was observed at m/z 447 refer to the release of rhamnosyl moiety ($-m/z$ 146 u) $[M-H\text{-rhamnosyl moiety}]^-$, further elimination of water molecule ($-m/z$ 18 u) led to generation of fragment ion at m/z 429 $[M-H\text{-rhamnosyl moiety-H}_2\text{O}]^-$, while the release of glucosyl moiety ($-m/z$ 162 u) led to generation of a diagnostic MS/MS fragment ion as a base peak at m/z 285 $[M-H\text{-rhamnosyl moiety-glucosyl moiety}]^-$ it was assigned to kaempferol a glycone, another key a glycone fragments were detected at m/z 255, 227, 179, and 169. Therefore, compound 9 was identified as kaempferol 3-O-rutinoside.^[41]

Compound 11 was detected at $R_t = 21.80$ min, it showed a deprotonated molecule $[M-H]^-$ at m/z 447, while the release of glucosyl moiety ($-m/z$ 162 u) led to generation of a diagnostic MS/MS fragment ion as a base peak at m/z 285 $[M-H\text{-glucosyl moiety}]^-$ it was assigned to kaempferol a glycone, another key a glycone fragments were detected at m/z 255, 227, 179, 169 and 151. Therefore, compound 11 was identified as kaempferol-3-O- β -D-glucoside.^[39,42,48] Compound 12 was detected at R_t 27.82 min, it showed a deprotonated molecule $[M-H]^-$ at m/z 609, while the release of rhamnosyl moiety ($-m/z$ 146 u) led to generation of MS/MS fragment ion at m/z 463 $[M-H\text{-rhamnosyl moiety}]^-$ it was assigned to quercetin-O-glucoside, the removal of water molecule ($-m/z$ 18 u) produce a fragment ion at m/z 445, while the elimination of the

glucosyl moiety ($-m/z$ 162 u) was confirmed through a diagnostic fragment at m/z 301 was assigned to quercetin a glycone, another key a glycone fragments were observed at m/z 271, 255, and 179. Therefore, compound 12 was identified as quercetin-3-O-rutinoside (rutin).^[36] Compound 15 was detected at $R_t = 42.90$ min, it showed a deprotonated molecule $[M-H]^-$ at m/z 329, while the release of methyl moiety ($-m/z$ 15 u) led to generation of MS/MS fragment ion at m/z 314 $[M-H\text{-CH}_3]^-$ it was assigned to quercetin methyl ether, the removal of another methyl moiety ($-m/z$ 15 u) produce a fragment ion at m/z 299 $[M-H\text{-2CH}_3]^-$. Therefore, compound 15 was identified as quercetin-dimethyl-ether as previously reported.^[43]

Phenylethanoid derivatives

Compound 13 was detected at $R_t = 32.20$ min, it showed a deprotonated molecule $[M-H]^-$ at m/z 685, while the release of glucosyl moiety ($-m/z$ 162 u) led to generation of a diagnostic MS/MS fragment ion as a base peak at m/z 523 $[M-H\text{-glucosyl moiety}]^-$ it was assigned to ligstroside molecule, further removal of water molecule ($-m/z$ 18 u) produce a fragment ion at m/z 505 $[M-H\text{-glucosyl moiety-H}_2\text{O}]^-$ and the elimination of another glucosyl moiety ($-m/z$ 162 u) was confirmed through the appearance of a diagnostic fragment at m/z 343 $[M-H\text{-H}_2\text{O-2glucosyl moiety}]^-$. Therefore, compound 13 was identified as ligstroside glucoside as previously described.^[34]

CONCLUSION

In the current study, the HPLC-DAD-ESI-MS/MS analysis led to the tentative identification of a total 30 phenolic compounds in the ethyl acetate and *n*-butanol extracts of *S. nicolai* leaves based on the determination of the precise mass of the deprotonated ions $[M-H]^-$, which was obtained from the MS data and MSⁿ fragmentation pattern. Among the tentatively identified compounds, flavonoids were the major constituents. To the best of our knowledge, this research work is the first comprehensive study on the polyphenolic composition of the Egyptian *S. nicolai* species. Moreover, *S. nicolai* extracts demonstrated considerable antioxidant and anticancer activities. *S. nicolai* could be further studied to isolate its biologically-active constituents and to study their modes of actions.

Acknowledgment

The authors wish to express their gratitude to Dr. Tearse Labib, Botany Specialist, Department of Flora and Taxonomy, El-Orman Botanical Garden, Giza, Egypt, for identification and authentication of the plant. Also, we would like to thank Dr. Mansour Sobeh & Dr. Tamer Mohamed; Department of Pharmaceutical Biology, Institute of Pharmacy and Molecular Biotechnology, Heidelberg University, Heidelberg, Germany, for their kind cooperation to perform LC/MS analysis.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

- Hölscher D, Schneider B. Phenalenones from *Strelitzia reginae*. *J Nat Prod* 2000;63:1027-8.
- Dwarka D, Thaver V, Naidu M, Baijnath H. New insights into the presence of bilirubin in a plant species *Strelitzia nicolai* (strelitziaceae). *Afr J Tradit Complement Altern Med* 2017;14:253-62.
- Chalannavar RK, Venugopala KN, Baijnath H, Odhav B. Chemical composition of essential oil from the seed arils of *Strelitzia nicolai* regel and Koern from South Africa. *J Essent Oil Bearing Plants* 2014;17:1373-7.

4. Pirone C, Quirke JM, Priestap HA, Lee DW. Animal pigment bilirubin discovered in plants. *J Am Chem Soc* 2009;131:2830.
5. Pirone C, Johnson JV, Quirke JM, Priestap HA, Lee D. The animal pigment bilirubin identified in *Strelitzia reginae*, the bird of paradise flower. *Hort Sci* 2010;45:1411-5.
6. Pirone C. Bilirubin: An Animal Pigment in the *Zingiberales* and Diverse Angiosperm Orders. Dissertation, Florida International University, UAS; 2010.
7. Dwarka D, Thaver V, Naidu M, Koorbanally NA, Baijnath AH. *In vitro* chemo-preventative activity of *Strelitzia nicolai* aril extract containing bilirubin. *Afr J Tradit Complement Altern Med* 2017;14:147-56.
8. Ghareeb MA, Saad AM, Abdou AM, Refahy LA, Ahmed WS. A new kaempferol glycoside with antioxidant activity from *Chenopodium ambrosioides* growing in Egypt. *Orient J Chem* 2016;32:3053-61.
9. Mangal M, Sagar P, Singh H, Raghava GP, Agarwal SM. NPACT: Naturally occurring plant-based anti-cancer compound-activity-target database. *Nucleic Acids Res* 2013;41:D1124-9.
10. Cheung-Ong K, Giaefer G, Nislow C. DNA-damaging agents in cancer chemotherapy: Serendipity and chemical biology. *Chem Biol* 2013;20:648-59.
11. Ibrahim RM, El-Halawany AM, Saleh DO, El Naggar EB, El-Shabrawy AO, El-Hawary SS. HPLC-DAD-MS/MS profiling of phenolics from *Securigera securidaca* flowers and its anti-hyperglycemic and anti-hyperlipidemic activities. *Rev Bras Farmacogn* 2015;25:134-41.
12. Ghareeb MA, Mohamed T, Saad AM, Refahy LA, Sobeh M, Wink M. HPLC-DAD-ESI-MS/MS analysis of fruits from *Firmiana simplex* (L.) and evaluation of their antioxidant and antigenotoxic properties. *J Pharm Pharm* 2018;70:133-142.
13. Yang H, Dong Y, Du H, Shi H, Peng Y, Li X, *et al.* Antioxidant compounds from propolis collected in Anhui, China. *Molecules* 2011;16:3444-55.
14. Qader SW, Abdulla MA, Chua LS, Najim N, Zain MM, Hamdan S, *et al.* Antioxidant, total phenolic content and cytotoxicity evaluation of selected Malaysian plants. *Molecules* 2011;16:3433-43.
15. Goodman J, Chandna A, Roe K. Trends in animal use at US research facilities. *J Med Ethics* 2015;41:567-9.
16. Soliman RH, Ismail OA, Badr MS, Nasr SM. Resveratrol ameliorates oxidative stress and organ dysfunction in *Schistosoma mansoni* infected mice. *Exp Parasitol* 2017;174:52-8.
17. Dunham JH, Guthmiller P. Doing good science: Authenticating cell line identity. *Cell Notes* 2008;22:15-7.
18. Li L, Sharma N, Chippada U, Jiang X, Schloss R, Yarmush ML, *et al.* Functional modulation of ES-derived hepatocyte lineage cells via substrate compliance alteration. *Ann Biomed Eng* 2008;36:865-76.
19. Nasr SM, Ghareeb MA, Mohamed MA, Elwan NM, Abdel-Aziz AA, Abdel-Aziz MS. HPLC-Fingerprint analyses, *in vitro* cytotoxicity, antimicrobial and antioxidant activities of the extracts of two *Cestrum* species growing in Egypt. *Pharmacog Res* 2018;10:173-180.
20. Sliwka L, Wiktorska K, Suchocki P, Milczarek M, Mielczarek S, Lubelska K, *et al.* The comparison of MTT and CVS assays for the assessment of anticancer agent interactions. *PLoS One* 2016;11:e0155772.
21. Pietta PG. Flavonoids as antioxidants. *J Nat Prod* 2000;63:1035-42.
22. Arora A, Nair MG, Strasburg GM. Structure-activity relationships for antioxidant activities of a series of flavonoids in a liposomal system. *Free Radic Biol Med* 1998;24:1355-63.
23. Rice-Evans CA, Miller NJ, Paganga G. Structure-antioxidant activity relationships of flavonoids and phenolic acids. *Free Radic Biol Med* 1996;20:933-56.
24. Rice-Evans CA, Miller N, Paganga G. Antioxidant properties of phenolic compounds. *Trends Plant Sci* 1997;2:152-9.
25. Discher DE, Janmey P, Wang YL. Tissue cells feel and respond to the stiffness of their substrate. *Science* 2005;310:1139-43.
26. Sobeh M, Mahmoud MF, Hasan RA, Abdelfattah MAO, Sabry OM, Ghareeb MA, *et al.* Tannin-rich extracts from *Lannea stuhlmannii* and *Lannea humilis* (Anacardiaceae) exhibit hepatoprotective activities *in vivo* via enhancement of the anti-apoptotic protein Bcl-2. *Sci Rep* 2018;8:9343.
27. Rashdan HR, Nasr SM, El-Refai HA, Abdel-Aziz MS. A novel approach of potent antioxidant and antimicrobial agents containing coumarin moiety accompanied with cytotoxicity studies on the newly synthesized derivatives. *J Appl Pharm Sci* 2017;7:186-96.
28. Tanaka T, Kojima T, Kawamori T, Wang A, Suzui M, Okamoto K, *et al.* Inhibition of 4-nitroquinoline-1-oxide-induced rat tongue carcinogenesis by the naturally occurring plant phenolics caffeic, ellagic, chlorogenic and ferulic acids. *Carcinogenesis* 1993;14:1321-5.
29. Lee IR, Yang MY. Phenolic compounds from *Duchesnea chrysantha* and their cytotoxic activities in human cancer cell. *Arch Pharm Res* 1994;17:476-9.
30. Hertog MG, Kromhout D, Aravanis C, Blackburn H, Buzina R, Fidanza F, *et al.* Flavonoid intake and long-term risk of coronary heart disease and cancer in the seven countries study. *Arch Intern Med* 1995;155:381-6.
31. Yáñez J, Vicente V, Alcaraz M, Castillo J, Benavente-García O, Canteras M, *et al.* Cytotoxicity and antiproliferative activities of several phenolic compounds against three melanocytes cell lines: Relationship between structure and activity. *Nutr Cancer* 2004;49:191-9.
32. Abliz G, Mijit F, Hua L, Abdixkur G, Ablimit T, Amat N, *et al.* Anti-carcinogenic effects of the phenolic-rich extract from abnormal Savda Munziq in association with its cytotoxicity, apoptosis-inducing properties and telomerase activity in human cervical cancer cells (SiHa). *BMC Complement Altern Med* 2015;15:23.
33. Abu-Reidah IM, Arráez-Román D, Segura-Carretero A, Fernández-Gutiérrez A. Extensive characterisation of bioactive phenolic constituents from globe artichoke (*Cynara scolymus* L.) by HPLC-DAD-ESI-QTOF-MS. *Food Chem* 2013;141:2269-77.
34. Sanz M, de Simón BF, Cadahía E, Esteruelas E, Muñoz AM, Hernández T, *et al.* LC-DAD/ESI-MS/MS study of phenolic compounds in Ash (*Fraxinus excelsior* L. and *F. americana* L.) heartwood. Effect of toasting intensity at cooperage. *J Mass Spectrom* 2012;47:905-18.
35. Chen HJ, Inbaraj BS, Chen BH. Determination of phenolic acids and flavonoids in *Taraxacum fimosanum* kitam by liquid chromatography-tandem mass spectrometry coupled with a post-column derivatization technique. *Int J Mol Sci* 2012;13:260-85.
36. Abu-Reidah IM, Ali-Shtayah MS, Jamous RM, Arráez-Román D, Segura-Carretero A. HPLC-DAD-ESI-MS/MS screening of bioactive components from *Rhus coriaria* L. (Sumac) fruits. *Food Chem* 2015;166:179-91.
37. Abu-Reidah IM, Arráez-Román D, Lozano-Sánchez J, Segura-Carretero A, Fernández-Gutiérrez A. Phytochemical characterisation of green beans (*Phaseolus vulgaris* L.) by using high-performance liquid chromatography coupled with time-of-flight mass spectrometry. *Phytochem Anal* 2013;24:105-16.
38. Abu-Reidah IM, del Mar Contreras M, Arráez-Román D, Fernández-Gutiérrez A, Segura-Carretero A. UHPLC-ESI-QTOF-MS-based metabolic profiling of *Vicia faba* L. (Fabaceae) seeds as a key strategy for characterization in foodomics. *Electrophoresis* 2014;35:1571-81.
39. Bravo L, Goya L, Lecumberri E. LC/MS characterization of phenolic constituents of mate (*Ilex paraguariensis*, St. Hil.) and its antioxidant activity compared to commonly consumed beverages. *Food Res Int* 2007;40:393-405.
40. Abu-Reidah IM, Arráez-Román D, Quirantes-Piné R, Fernández-Arroyo S, Segura-Carretero A, Fernández-Gutiérrez A. HPLC-ESI-Q-TOF-MS for a comprehensive characterization of bioactive phenolic compounds in cucumber whole fruit extract. *Food Res Int* 2012;46:108-17.
41. Al-Rawahi AS, Edwards G, Al-Sibani M, Al-Thani G, Al-Harrasi AS, Rahman MS. Phenolic constituents of pomegranate peels (*Punica granatum* L.) cultivated in Oman. *Eur J Med Plants* 2014;4:315-31.
42. Chen F, Long X, Liu Z, Shao H, Liu L. Analysis of phenolic acids of Jerusalem artichoke (*Helianthus tuberosus* L.) responding to salt-stress by liquid chromatography/tandem mass spectrometry. *ScientificWorldJournal* 2014;2014:568043.
43. Pellati F, Orlandini G, Pinetti D, Benvenuti S. HPLC-DAD and HPLC-ESI-MS/MS methods for metabolite profiling of Propolis extracts. *J Pharm Biomed Anal* 2011;55:934-48.
44. Simirgiotis MJ, Benites J, Areche C, Sepúlveda B. Antioxidant capacities and analysis of phenolic compounds in three endemic Nolana species by HPLC-PDA-ESI-MS. *Molecules* 2015;20:11490-507.
45. Abdel-Hameed ES, Bazaid SA, Shohayeb MM. RP-HPLC-UV-ESI-MS phytochemical analysis of fruits of *Conocarpus erectus* L. *Chem Pap* 2014;68:1358-67.
46. Kim DW, Curtis-Long MJ, Yuk HJ, Wang Y, Song YH, Jeong SH, *et al.* Quantitative analysis of phenolic metabolites from different parts of *Angelica keiskei* by HPLC-ESI MS/MS and their xanthine oxidase inhibition. *Food Chem* 2014;153:20-7.
47. Riethmüller E, Alberti A, Tóth G, Béni S, Ortolano F, Kéry A, *et al.* Characterisation of diarylheptanoid- and flavonoid-type phenolics in *Corylus avellana* L. Leaves and bark by HPLC/DAD-ESI/MS. *Phytochem Anal* 2013;24:493-503.
48. Ágnes A. LC-ESI-MS/MS methods in profiling of flavonoid glycosides and phenolic acids in traditional medicinal plants: *Sempervivum tectorum* L. and *Corylus avellana* L. Dissertation, Semmelweis University, Hungary; 2013.