

1-16-2017

Hubble Parameter Measurement Constraints on the Redshift of the Deceleration-Acceleration Transition, Dynamical Dark Energy, and Space Curvature

Muhammad O. Farooq

Embry-Riddle Aeronautical University, muhammad.farooq@erau.edu

Foram R. Madiyar

Kansas State University and Embry Riddle Aeronautical University, madiyarf@erau.edu

Sara Crandall

Kansas State University

Bharat Ratra

Kansas State University,

Follow this and additional works at: <https://commons.erau.edu/publication>



Part of the [Cosmology, Relativity, and Gravity Commons](#)

Scholarly Commons Citation

Farooq, M. O., Madiyar, F. R., Crandall, S., & Ratra, B. (2017). Hubble Parameter Measurement Constraints on the Redshift of the Deceleration-Acceleration Transition, Dynamical Dark Energy, and Space Curvature. *The Astrophysical Journal*, 835(1). <https://doi.org/10.3847/1538-4357/835/1/26>

This Article is brought to you for free and open access by Scholarly Commons. It has been accepted for inclusion in Publications by an authorized administrator of Scholarly Commons. For more information, please contact commons@erau.edu.

HUBBLE PARAMETER MEASUREMENT CONSTRAINTS ON THE REDSHIFT OF THE DECELERATION-ACCELERATION TRANSITION, DYNAMICAL DARK ENERGY, AND SPACE CURVATURE

OMER FAROOQ¹, FORAM RANJEET MADIYAR¹, SARA CRANDALL², AND BHARAT RATRA²

¹Department of Physical Sciences, Embry-Riddle Aeronautical University, 600 Clyde Morris Boulevard, Daytona Beach, FL 32114, USA; farooqm@erau.edu

²Department of Physics, Kansas State University, 116 Cardwell Hall, Manhattan, KS 66506, USA

Draft version November 16, 2016

ABSTRACT

We compile an updated list of 38 measurements of the Hubble parameter $H(z)$ between redshifts $0.07 \leq z \leq 2.36$ and use them to place constraints on model parameters of constant and time-varying dark energy cosmological models, both spatially flat and curved. We use five models to measure the redshift of the cosmological deceleration-acceleration transition, z_{da} , from these $H(z)$ data. Within the error bars, the measured z_{da} are insensitive to the model used, depending only on the value assumed for the Hubble constant H_0 . The weighted mean of our measurements is $z_{\text{da}} = 0.72 \pm 0.05$ (0.84 ± 0.03) for $H_0 = 68 \pm 2.8$ (73.24 ± 1.74) $\text{km s}^{-1} \text{Mpc}^{-1}$ and should provide a reasonably model-independent estimate of this cosmological parameter. The $H(z)$ data are consistent with the standard spatially-flat Λ CDM cosmological model but do not rule out non-flat models or dynamical dark energy models.

Subject headings: cosmological parameters — cosmology: observations — dark energy

1. INTRODUCTION

In the standard scenario the currently accelerating cosmological expansion is a consequence of dark energy dominating the current cosmological energy budget; at earlier times non-relativistic (cold dark and baryonic) matter dominated the energy budget and powered the decelerating cosmological expansion.¹ Initial quantitative observational support for this picture came from “lower” redshift Type Ia supernova (SNIa) apparent magnitude observations and “higher” redshift cosmic microwave background (CMB) anisotropy measurements.

More recently, cosmic chronometric and baryon acoustic oscillation (BAO) techniques (see, e.g., [Simon et al. 2005](#); [Moresco et al. 2012](#); [Busca et al. 2013](#)) have resulted in the measurement of the cosmological expansion rate or Hubble parameter, $H(z)$, from the present epoch back to a redshift z exceeding 2, higher than currently probed by SNIa observations. This has resulted in the first mapping out of the cosmological deceleration-acceleration transition, the epoch when dark energy took over from non-relativistic matter, and the first measurement of the redshift of this transition (see, e.g., [Farooq & Ratra 2013b](#); [Farooq et al. 2013b](#); [Moresco et al. 2016](#)).²

$H(z)$ measurements have also been used to constrain some more conventional cosmological parameters, such as the density of dark energy and the density of non-relativistic matter (see, e.g., [Samushia & Ratra 2006](#); [Chen & Ratra 2011b](#); [Farooq & Ratra 2013a](#); [Akarsu et al. 2014](#); [Chimento & Richarte 2013](#); [Gruber & Luongo 2014](#); [Bamba et al. 2014](#); [Ferreira et al. 2013](#); [Forte 2014](#); [Chen et al. 2015](#); [Dankiewicz et al. 2014](#); [Capozziello et al. 2014](#); [Meng et al. 2015](#); [Guo & Zhang 2016](#); [Mukherjee & Banerjee 2016](#); [Alam et al. 2016](#)), typically providing constraints comparable to or better than those provided by SNIa data, but not as good as those from BAO or CMB anisotropy measurements. More recently, $H(z)$ data has been used to measure the Hubble constant H_0 ([Verde et al. 2014](#); [Chen et al. 2016a](#)), with the resulting H_0 value being more consistent with recent lower values determined from a median statistics analysis of Huchra’s H_0 compilation ([Chen & Ratra 2011a](#)), from CMB anisotropy data ([Hinshaw et al. 2013](#); [Sievers et al. 2013](#); [Ade et al. 2015](#)), from BAO measurements ([Aubourg et al. 2015](#); [Ross et al. 2015](#); [L’Huillier & Shafieloo 2016](#)), and from current cosmological data and the standard model of particle physics with only three light neutrino species (see, e.g., [Calabrese et al. 2012](#)).

In this paper, we put together an updated list of $H(z)$ measurements, compared to that of [Farooq & Ratra \(2013b\)](#), and use this compilation to constrain the redshift of the cosmological deceleration-acceleration transition, z_{da} , as well as other cosmological parameters. In the z_{da} analysis here we study more models than used by [Farooq & Ratra \(2013b\)](#) and [Farooq et al. \(2013b\)](#), now also allowing for non-zero spatial curvature in the Λ CDM parametrization of dynamical dark energy case and in the dynamical dark energy ϕ CDM model ([Pavlov et al. 2013](#)). The cosmological parameter constraints derived here are based on more, as well as more recent, $H(z)$ data than were used by [Farooq et al. \(2015\)](#) and we also explore a much larger range of parameter space in the non-flat ϕ CDM model than they did.

We find, from the likelihood analyses, that the z_{da} values measured from the $H(z)$ data agree within the error bars in all five models. They, however, depend more sensitively on the value of H_0 assumed in the analysis. These results are consistent with those found in [Farooq & Ratra \(2013b\)](#) and [Farooq et al. \(2013b\)](#). In addition, the binned $H(z)$

¹ For reviews of this picture, as well as of the alternate modified gravity scenario, see [Ratra & Vogeley \(2008\)](#), [Weinberg et al. \(2013\)](#), [Martin \(2012\)](#), [Joyce et al. \(2016\)](#), and references therein.

² See [Sutherland & Rothnie \(2015\)](#) and [Muthukrishna & Parkinson \(2016\)](#) for lower limits on this redshift derived using SNIa and other data. For upper limits on the transition redshift see [Rani et al. \(2015\)](#).

TABLE 1
HUBBLE PARAMETER VERSUS REDSHIFT DATA

z	$H(z)$ (km s ⁻¹ Mpc ⁻¹)	σ_H (km s ⁻¹ Mpc ⁻¹)	Reference ^a
0.070	69	19.6	5
0.090	69	12	1
0.120	68.6	26.2	5
0.170	83	8	1
0.179	75	4	3
0.199	75	5	3
0.200	72.9	29.6	5
0.270	77	14	1
0.280	88.8	36.6	5
0.352	83	14	3
0.380	81.5	1.9	10
0.3802	83	13.5	9
0.400	95	17	1
0.4004	77	10.2	9
0.4247	87.1	11.2	9
0.440	82.6	7.8	4
0.4497	92.8	12.9	9
0.4783	80.9	9	9
0.480	97	62	2
0.510	90.4	1.9	10
0.593	104	13	3
0.600	87.9	6.1	4
0.610	97.3	2.1	10
0.680	92	8	3
0.730	97.3	7	4
0.781	105	12	3
0.875	125	17	3
0.880	90	40	2
0.900	117	23	1
1.037	154	20	3
1.300	168	17	1
1.363	160	33.6	8
1.430	177	18	1
1.530	140	14	1
1.750	202	40	1
1.965	186.5	50.4	8
2.340	222	7	7
2.360	226	8	6

^a Reference numbers: 1. [Simon et al. \(2005\)](#), 2. [Stern et al. \(2010\)](#), 3. [Moresco et al. \(2012\)](#), 4. [Blake et al. \(2012\)](#), 5. [Zhang et al. \(2012\)](#), 6. [Font-Ribera et al. \(2014\)](#), 7. [Delubac et al. \(2015\)](#), 8. [Moresco \(2015\)](#), 9. [Moresco et al. \(2016\)](#), 10. [Alam et al. \(2016\)](#).

data in redshift space show qualitative visual evidence for the deceleration-acceleration transition, independent of how they are binned provided the bins are narrow enough, in agreement with that originally found by [Farooq et al. \(2013b\)](#). Given that the measured z_{da} are relatively model independent, it is not unreasonable to average the measured values to determine a reasonable summary estimate. We find, for a weighted mean estimate, $z_{\text{da}} = 0.72 \pm 0.05$ (0.84 ± 0.03) if we assume $H_0 = 68 \pm 2.8$ (73.24 ± 1.74) km s⁻¹ Mpc⁻¹.

The constraints on the more conventional cosmological parameters, such as the density of dark energy, derived from the likelihood analysis of the $H(z)$ data here, indicate that these data are quite consistent with the spatially-flat Λ CDM model, the standard model of cosmology where the cosmological constant Λ is the dark energy. These $H(z)$ data, however, do not rule out the possibility of dynamical dark energy or space curvature, especially when included simultaneously, in agreement with the conclusions of [Farooq et al. \(2015\)](#). Currently available SNIa, BAO, growth factor, CMB anisotropy, and other data can tighten the constraints on these parameters, and it will be interesting to study these data sets in conjunction with the $H(z)$ data we have compiled here, but this is beyond the scope of our paper. Near-future data will also result in interesting limits (see, e.g., [Podariu et al. 2001a](#); [Pavlov et al. 2012](#); [Basse et al. 2014](#); [Santos et al. 2013](#)).

The outline of our paper is as follows. In the next section, we discuss and tabulate our new $H(z)$ data compilation. In Sec. 3 we summarize how we bin the $H(z)$ data in redshift space and list binned $H(z)$ data. Section 4 summarizes the cosmological models we consider. In Sec. 5 we discuss how we compute and measure the deceleration-acceleration transition redshift and tabulate numerical values of z_{da} determined from the $H(z)$ measurements. Section 6 presents the constraints on cosmological parameters, and we conclude in the last section.

In Table 1 we collect 38 Hubble parameter $H(z)$ measurements from Simon et al. (2005), Stern et al. (2010), Moresco et al. (2012), Blake et al. (2012), Zhang et al. (2012), Font-Ribera et al. (2014), Delubac et al. (2015), Moresco (2015), Moresco et al. (2016), and Alam et al. (2016). These data are plotted in the top panel of Fig. 1.

These 38 $H(z)$ measurements are not completely independent. The three measurements taken from Blake et al. (2012) are correlated with each other and the three measurements of Alam et al. (2016) also are correlated. **Also, in these and other cases, when BAO observations are used to measure $H(z)$, one has to apply a prior on the radius of the sound horizon, $r_d = \int_{z_d}^{\infty} c_s(z)dz/H(z)$, evaluated at the drag epoch z_d , shortly after recombination, when photons and baryons decouple. This prior value of r_d is generally derived from CMB observations.**




FIG. 1.— The top panel shows the 38 $H(z)$ measurements of Table 1. All error bars are 1σ . The left (right) panel in the second row shows the binned $H(z)$ data with 3 or 4 (4 or 5) measurements per bin, combined using weighted mean statistics, listed in Table 2. In the last row, the left (right) panel shows binned $H(z)$ data with 5 or 7 (4, 5, or 6) measurements per bin, combined using weighted mean statistics, listed in Table 2. In all panels, there are five different colored solid (dot-dashed) best-fit model prediction lines for the two H_0 priors used in our analyses (see main text for details; NF stands for non-flat).

Table 1 here is based on Table 1 of [Farooq & Ratra \(2013b\)](#) with the following modifications. We drop older SDSS galaxy clustering $H(z)$ determinations from [Chuang & Wang \(2013\)](#) in favor of the more recent measurements from [Alam et al. \(2016\)](#). We have added the new [Moresco et al. \(2016\)](#) measurements. We have dropped the older [Busca et al. \(2013\)](#) Ly- α forest measurement in favor of the newer [Font-Ribera et al. \(2014\)](#) and [Delubac et al. \(2015\)](#) ones. We have also added two new measurements from [Moresco \(2015\)](#).

There are many other compilations of $H(z)$ data available in the literature (see, e.g., [Meng et al. 2015](#); [Cai et al. 2014](#); [Solà et al. 2016](#); [Yu & Wang 2016](#); [Duan et al. 2016](#); [Qi et al. 2016](#); [Nunes et al. 2016](#); [Zhang & Xia 2016](#)). We emphasize that our compilation here does not include older, less reliable, data, a few with a lot of weight because of anomalously small error bars.

3. BINNING OF HUBBLE PARAMETER DATA

There are two reasons to compute “average” $H(z)$ values for bins in redshift space. First, the weighted mean technique of binning data can indicate if the original unbinned data have error bars inconsistent with Gaussianity, an important consistency check. Second, data binned in redshift space can more clearly visually illustrate trends as a function of redshift, with the additional advantage of not having to assume a particular cosmological model.

The 38 Hubble parameter measurements in Table 1 are binned to ensure as many measurements as possible per bin, while also retaining as many (narrow) redshift bins as

possible. The ideal case is $\sqrt{38}$ measurements in each of $\sqrt{38}$ bins. Here we consider about 3-4, 4-5, 4-5-6, and 5-7 measurements per bin. The last four measurements are binned by twos in all but the 4-5-6 measurement per bin case. In all cases, data points in a given bin are not correlated with each other.

After binning the data, we use weighted mean statistics³ to find a representative central estimate for each bin. Following [Podariu et al. \(2001b\)](#) the weighted mean is given by,

$$\bar{H}(z) = \frac{\sum_{i=1}^N H(z_i)/\sigma_i^2}{\sum_{i=1}^N 1/\sigma_i^2}, \quad (1)$$

where $H(z_i)$ and σ_i are the Hubble parameter and one standard deviation of $i = 1, 2, 3 \dots N$ measurements in the bin. We also compute the weighted bin redshift using

$$\bar{z} = \frac{\sum_{i=1}^N z_i/\sigma_i^2}{\sum_{i=1}^N 1/\sigma_i^2}. \quad (2)$$

The associated weighted error is given by

$$\bar{\sigma} = \left(\sum_{i=1}^N 1/\sigma_i^2 \right)^{-1/2}. \quad (3)$$

A goodness-of-fit, χ^2 , can be found for each bin where the reduced χ^2 is

$$\chi_\nu^2 = \frac{1}{N-1} \sum_{i=1}^N \frac{[H(z_i) - \bar{H}(z)]^2}{\sigma_i^2}. \quad (4)$$

The number of standard deviations that χ_ν deviates from unity (the expected value) is given by

$$N_\sigma = |\chi_\nu - 1| \sqrt{2(N-1)}. \quad (5)$$

A large N_σ can be the result of non-Gaussian measurements, the presence of un-accounted for systematic errors, or correlations between measurements. Table 2 lists the weighted mean results for the binned $H(z)$ measurements.

The last column of Table 2 shows reasonably small N_σ for all binnings, and so suggests that the error bars of the $H(z)$ data of Table 1 are not inconsistent with Gaussianity. As in [Farooq et al. \(2013b\)](#), we find that the cosmological constraints that follow from the weighted mean binned data are almost identical to those derived using the unbinned data, while the median statistics binned data typically result in somewhat weaker constraints. A possible reason for this could be that some of the unbinned $H(z)$ data error bars might be a bit larger than they really should be. This would be consistent with the low reduced χ^2 shown in the last line of Table 1 in [Chen et al. \(2016a\)](#).

The binned data are plotted in the four lower panels of Fig. 1. It is reassuring that, independent of the binning used, all the binned data sets show clear visual qualitative evidence for the cosmological deceleration-acceleration transition, as in [Farooq et al. \(2013b\)](#). This is model-independent qualitative evidence for the existence of the cosmological deceleration-acceleration transition. We shall see, in Sec. 5, that all cosmological models we use in the analysis of

³ We also used median statistics to find central estimates, where the median is the value for which there is a 50% chance of finding a measurement above and below it. Since median statistics does not make use of individual measurement errors, the resultant central estimate error is larger than that for weighted mean statistics. For discussions and applications of median statistics, see [Gott et al. \(2001\)](#), [Chen & Ratra \(2003\)](#), [Hodge et al. \(2009\)](#), [Crandall & Ratra \(2014\)](#), [Crandall et al. \(2015\)](#), [Ding et al. \(2015\)](#), [Crandall & Ratra \(2015\)](#), and [Zheng et al. \(2016\)](#). As in [Farooq et al. \(2013b\)](#) for the earlier $H(z)$ data tabulated in [Farooq & Ratra \(2013b\)](#), all median statistics analyses results look reasonable and, since the weighted mean results are also all reasonable and more constraining, going forward we use only weighted mean results.

the $H(z)$ data to measure z_{da} result in z_{da} values that overlap within the error bars (for a given H_0 prior). This is additional model-independent evidence for the presence of the deceleration-acceleration transition.

TABLE 2
WEIGHTED MEAN RESULTS FOR 38 REDSHIFT MEASUREMENTS

Bin	N	z^a	$H(z)$ (km s ⁻¹ Mpc ⁻¹)	$H(z)$ (1 σ range) (km s ⁻¹ Mpc ⁻¹)	$H(z)$ (2 σ range) (km s ⁻¹ Mpc ⁻¹)	N_σ
3 or 4 measurements per bin						
1	3	0.0892	69.0	59.4-78.5	49.9-88.0	2.0
2	4	0.185	76.0	73.1-78.9	70.2-81.8	1.1
3	3	0.309	80.6	71.0-90.2	61.5-99.7	1.5
4	4	0.381	81.5	79.7-83.4	77.9-85.2	1.2
5	3	0.438	85.8	80.1-91.5	74.3-97.3	1.0
6	3	0.509	90.0	88.1-91.9	86.3-93.7	0.53
7	3	0.609	96.5	94.5-98.4	92.6-100	0.22
8	3	0.720	96.6	91.8-101	87.0-106	0.71
9	4	0.929	129	118-140	108-151	0.066
10	4	1.43	158	149-167	140-176	0.047
11	2	1.83	196	165-227	133-259	1.1
12	2	2.35	224	219-229	213-234	0.88
4 or 5 measurements per bin						
1	2	0.0846	69.0	58.8-79.2	48.5-89.5	1.4
2	5	0.184	75.9	73.1-78.8	70.2-81.7	1.4
3	5	0.377	81.5	79.7-83.3	77.8-85.2	2.3
4	5	0.427	84.6	79.8-89.4	75.0-94.2	1.1
5	5	0.518	90.1	88.3-91.8	86.6-93.6	0.66
6	4	0.628	97.2	95.3-99.1	93.3-101	1.2
7	4	0.929	129	118-140	108-151	0.066
8	4	1.43	158	149-167	140-176	0.047
9	2	1.83	196	165-227	133-259	1.1
10	2	2.35	224	219-229	213-234	0.88
4, 5, or 6 measurements per bin						
1	4	0.137	77.2	71.1-83.3	64.9-89.5	0.85
2	5	0.192	75.2	72.1-78.2	69.1-81.2	2.3
3	5	0.380	81.6	79.7-83.4	77.9-85.2	1.5
4	6	0.502	89.6	87.8-91.4	86.1-93.1	1.1
5	4	0.613	96.2	94.3-98.1	92.4-100	0.10
6	6	0.787	106	101-112	95.8-117	1.1
7	6	1.46	161	153-170	144-178	0.16
8	2	2.35	224	219-229	213-234	0.88
5 or 7 measurements per bin						
1	5	0.166	75.7	72.3-79.0	69.0-82.4	1.2
2	7	0.355	80.7	79.0-82.4	77.2-84.2	1.6
3	5	0.427	84.6	79.8-89.4	75.0-94.2	1.1
4	5	0.518	90.1	88.3-91.8	86.6-93.6	0.66
5	7	0.633	97.7	95.8-99.6	93.9-102	0.55
6	5	1.37	158	149-166	141-174	0.32
7	2	1.83	196	165-227	133-259	1.1
8	2	2.35	224	219-229	213-234	0.88

^a Weighted mean of z values of measurements in the bin.

4. COSMOLOGICAL MODELS

In this section we briefly describe the five models we use to analyze the $H(z)$ data. These are the Λ CDM model that allows for spatial curvature and where dark energy is the cosmological constant Λ (Peebles 1984), as well as the ϕ CDM model in which dynamical dark energy is represented by a slowly evolving scalar field ϕ (Peebles & Ratra 1988; Ratra & Peebles 1988). We also consider an incomplete, but popular, parameterization of dynamical dark energy, XCDM, where dynamical dark energy is represented by an X -fluid. In the ϕ CDM and XCDM cases, we consider both

spatially-flat and non-flat models (Pavlov et al. 2013).

In the Λ CDM model with spatial curvature the Hubble parameter is

$$H(z; H_0, \mathbf{p}) = H_0 [\Omega_{m0}(1+z)^3 + \Omega_\Lambda + (1 - \Omega_{m0} - \Omega_\Lambda)(1+z)^2]^{1/2}, \quad (6)$$

where we have made use of $\Omega_{K0} = 1 - \Omega_{m0} - \Omega_\Lambda$ to eliminate the current value of the space curvature energy density parameter in favor of the current value of the non-relativistic matter energy density parameter, Ω_{m0} , and the cosmological constant energy density parameter, Ω_Λ . Here $\mathbf{p} = (\Omega_{m0}, \Omega_\Lambda)$ are the two cosmological parameters that conventionally characterize Λ CDM and H_0 is the value of Hubble parameter at the present time and is called the Hubble constant.

It has become fashionable to parameterize dynamical dark energy as a spatially homogeneous X -fluid, with a constant equation of state parameter, $\omega_X = p_X/\rho_X < -1/3$ (here p_X and ρ_X are the pressure and energy density of the X -fluid respectively). For the spatially-flat XCDM parameterization, using $\Omega_{X0} = 1 - \Omega_{m0}$ (where Ω_{X0} is the current value of the X -fluid energy density parameter), we have

$$H(z; H_0, \mathbf{p}) = H_0 [\Omega_{m0}(1+z)^3 + (1 - \Omega_{m0})(1+z)^{3(1+\omega_X)}]^{1/2}. \quad (7)$$

In this spatially-flat case the two cosmological parameters are $\mathbf{p} = (\Omega_{m0}, \omega_X)$. The XCDM parameterization is incomplete as it cannot describe the evolution of energy density inhomogeneities. In the non-flat XCDM parametrization case, Ω_{K0} is the third free parameter and

$$H(z; H_0, \mathbf{p}) = H_0 [\Omega_{m0}(1+z)^3 + (1 - \Omega_{m0} - \Omega_{K0})(1+z)^{3(1+\omega_X)} + \Omega_{K0}(1+z)^2]^{1/2}, \quad (8)$$

where the three cosmological parameters are $\mathbf{p} = (\Omega_{m0}, \omega_X, \Omega_{K0})$. ϕ CDM is the simplest, complete and consistent dynamical dark energy model. Here dark energy is modeled as a slowly-rolling scalar field ϕ with an, e.g., inverse-power-law potential energy density $V(\phi) = \kappa m_p^2 \phi^{-\alpha}/2$, where m_p is the Planck mass and α is a non-negative parameter that determines the coefficient $\kappa(m_p, \alpha)$ (Peebles & Ratra 1988). The equation of motion of the scalar field is

$$\ddot{\phi} + 3\frac{\dot{a}}{a}\dot{\phi} - \frac{\kappa}{2}\alpha m_p^2 \phi^{-(\alpha+1)} = 0, \quad (9)$$

where an overdot represents a time derivative and a is the scale factor. For the spatially-flat ϕ CDM model

$$H(z; H_0, \mathbf{p}) = H_0 [\Omega_{m0}(1+z)^3 + \Omega_\phi(z, \alpha)]^{1/2}, \quad (10)$$

where the time-dependent scalar field energy density parameter is

$$\Omega_\phi(z, \alpha) = \frac{1}{12H_0^2} (\dot{\phi}^2 + \kappa m_p^2 \phi^{-\alpha}). \quad (11)$$

In this case the two cosmological parameters are $\mathbf{p} = (\Omega_{m0}, \alpha)$. In the non-flat ϕ CDM model

$$H(z; H_0, \mathbf{p}) = H_0 [\Omega_{m0}(1+z)^3 + \Omega_\phi(z, \alpha) + \Omega_{K0}(1+z)^2]^{1/2}, \quad (12)$$

and the three cosmological parameters are $\mathbf{p} = (\Omega_{m0}, \alpha, \Omega_{K0})$.

Solving the coupled differential equations of motion allows for a numerical computation of the Hubble parameter $H(z; H_0, \mathbf{p})$ (Peebles & Ratra 1988; Samushia 2009; Farooq 2013; Pavlov et al. 2013).⁴

In Sec. 6 we use these expressions for the Hubble parameter in conjunction with the $H(z)$ measurements in Table 1 to constrain the cosmological parameters of these models. In our analyses here we study the following parameter ranges: $0 \leq \Omega_{m0} \leq 1$, $0 \leq \Omega_\Lambda \leq 1.4$, $-2 \leq \omega_X \leq 0$, $0 \leq \alpha \leq 5$, and $-0.7 \leq \Omega_{K0} \leq 0.7$ for non-flat XCDM and $-0.4 \leq \Omega_{K0} \leq 0.4$ for non-flat ϕ CDM (which is double the Ω_{K0} range used in Farooq et al. 2015).

5. COSMOLOGICAL DECELERATION-ACCELERATION TRANSITION REDSHIFT

At the current epoch, dark energy dominates the cosmological energy budget and accelerates the cosmological expansion. At earlier times non-relativistic (baryonic and cold dark) matter dominated the energy budget and the cosmological expansion decelerated. The cosmological deceleration-acceleration transition redshift, z_{da} , is defined as the redshift at which $\ddot{a} = 0$, in the cosmological model under consideration. \ddot{a} is proportional to the active gravitational mass density, the sum of the energy densities and three times the pressure of the constituents.

For Λ CDM, setting $\ddot{a} = 0$ we find

$$z_{\text{da}} = \left(\frac{2\Omega_\Lambda}{\Omega_{m0}} \right)^{1/3} - 1. \quad (13)$$

For the case of the spatially-flat XCDM parameterization

$$z_{\text{da}} = \left(\frac{\Omega_{m0}}{(\Omega_{m0} - 1)(1 + 3\omega_X)} \right)^{1/3\omega_X} - 1, \quad (14)$$

⁴ For discussions of observational constraints on the ϕ CDM model see, e.g. Podariu & Ratra (2000), Chen & Ratra (2004), Samushia & Ratra (2010), Samushia et al. (2010), Campanelli et al. (2012), Pavlov et al. (2014), Avsajanishvili et al. (2014), Avsajanishvili et al. (2015), Lima et al. (2016), Gosenca & Coles (2015), and Chen et al. (2016b).

while for non-flat XCDM

$$z_{\text{da}} = \left(\frac{\Omega_{m0}}{(\Omega_{m0} + \Omega_{K0} - 1)(1 + 3\omega_X)} \right)^{1/3\omega_X} - 1. \quad (15)$$

For the spatially-flat ϕ CDM model, defining the time-dependent equation-of-state-parameter for the scalar field

$$\omega_\phi(z) = \frac{\frac{1}{2}\dot{\phi}^2 - V(\phi)}{\frac{1}{2}\dot{\phi}^2 + V(\phi)}, \quad (16)$$

the redshift $z_{\text{da}}(\Omega_{m0}, \alpha)$ is determined by numerically solving

$$\Omega_{m0}(1 + z_{\text{da}})^3 + \Omega_\phi(z_{\text{da}}, \alpha) [1 + 3\omega_\phi(z_{\text{da}})] = 0 \quad (17)$$

where $\Omega_{\phi0} = 1 - \Omega_{m0}$. In the non-flat ϕ CDM model $z_{\text{da}}(\Omega_{m0}, \alpha, \Omega_{K0})$ is determined by numerically solving the same equation, but now setting $\Omega_{\phi0} = 1 - \Omega_{m0} - \Omega_{K0}$.

TABLE 3
DECELERATION-ACCELERATION TRANSITION REDSHIFTS^a

Model	h Prior ^b	BF ^c	χ^2_{min}	$z_{\text{da}} \pm \sigma_{z_{\text{da}}}$ ^d	$z_{\text{da}} \pm \sigma_{z_{\text{da}}}$ ^e
Λ CDM	0.68 ± 0.028	$\Omega_{m0} = 0.23$ $\Omega_\Lambda = 0.60$	22.4	0.723 ± 0.089	0.690 ± 0.096
	0.7324 ± 0.0174	$\Omega_{m0} = 0.25$ $\Omega_\Lambda = 0.78$	24.2	0.832 ± 0.055	0.781 ± 0.067
Flat XCDM	0.68 ± 0.028	$\Omega_{m0} = 0.26$ $\omega_X = -0.86$	22.5	0.753 ± 0.091	0.677 ± 0.097
	0.7324 ± 0.0174	$\Omega_{m0} = 0.24$ $\omega_X = -1.06$	23.9	0.813 ± 0.062	0.696 ± 0.082
Flat ϕ CDM	0.68 ± 0.028	$\Omega_{m0} = 0.27$ $\alpha = 0.50$	22.9	0.703 ± 0.104	0.724 ± 0.148
	0.7324 ± 0.0174	$\Omega_{m0} = 0.25$ $\alpha = 0$	25.2	0.885 ± 0.056	0.850 ± 0.116
Non-flat XCDM	0.68 ± 0.028	$\Omega_{m0} = 0.15$ $\omega_X = -1.68$ $\Omega_{K0} = 0.45$	21.9	0.684 ± 0.117	...
	0.7324 ± 0.0174	$\Omega_{m0} = 0.13$ $\omega_X = -2$ $\Omega_{K0} = 0.41$	20.3	0.709 ± 0.090	...
Non-flat ϕ CDM	0.68 ± 0.028	$\Omega_{m0} = 0.23$ $\alpha = 0$ $\Omega_{K0} = 0.18$	22.6	0.690 ± 0.118	...
	0.7324 ± 0.0174	$\Omega_{m0} = 0.25$ $\alpha = 0$ $\Omega_{K0} = -0.03$	25.0	0.853 ± 0.053	...

^a Estimated using the unbinned data of Table 1.

^b Hubble constant in units of $100 \text{ km s}^{-1} \text{ Mpc}^{-1}$.

^c Best-fit parameter values.

^d Computed using Eqs. 13–18 of this work.

^e The deceleration-acceleration transition redshift in the model, as computed in Farooq et al. (2013b) Table 1. Note that the best-fit cosmological parameter values found in Farooq et al. (2013b) differ from those found here and listed in this Table.

To compute the expected values $\langle z_{\text{da}} \rangle$ and $\langle z_{\text{da}}^2 \rangle$ for the two-parameter models we use

$$\langle z_{\text{da}} \rangle = \frac{\int \int z_{\text{da}}(\mathbf{p}) \mathcal{L}(\mathbf{p}) d\mathbf{p}}{\int \int \mathcal{L}(\mathbf{p}) d\mathbf{p}}, \quad \langle z_{\text{da}}^2 \rangle = \frac{\int \int z_{\text{da}}^2(\mathbf{p}) \mathcal{L}(\mathbf{p}) d\mathbf{p}}{\int \int \mathcal{L}(\mathbf{p}) d\mathbf{p}}. \quad (18)$$

Here $\mathcal{L}(\mathbf{p})$ is the $H(z)$ data likelihood function after marginalization over the Gaussian H_0 prior in the two-parameter model under consideration, as explained in [Farooq et al. \(2013a\)](#) and [Farooq et al. \(2015\)](#) but this time accounting for the non-diagonal correlation matrices of the [Blake et al. \(2012\)](#) and the [Alam et al. \(2016\)](#) measurements, which have a small effect. $\mathcal{L}(\mathbf{p})$ depends only on the model parameters $(\Omega_{m0}, \Omega_\Lambda)$ for Λ CDM, (Ω_{m0}, ω_X) for flat XCDM, and (Ω_{m0}, α) for flat ϕ CDM. The generalization for the three-parameter models is straightforward. The standard deviation in z_{da} is computed from the standard formula $\sigma_{z_{\text{da}}} = \sqrt{\langle z_{\text{da}}^2 \rangle - \langle z_{\text{da}} \rangle^2}$. The results of this computation are summarized in [Table 3](#).

[Table 3](#) shows best-fit cosmological parameter values and the corresponding minimum χ^2 for the five different cosmological models and for the two Gaussian H_0 priors. The second last column in [Table 3](#) shows the average deceleration-acceleration transition redshift with corresponding standard deviation for each model. It is very reassuring that the z_{da} values we measure in the five different models (for a given H_0 prior) overlap reasonably well. (The main effect on the measured z_{da} value is the assumed H_0 prior value.) Given that the measured z_{da} are almost independent of the other model parameters, within the errors, we may conclude that to leading order we have measured a model-independent z_{da} value. However, it is useful to have a single summary value for this cosmological parameter.

By taking the simple average of the penultimate column z_{da} values and computing the population standard deviation for the five values in this column, we find $z_{\text{da}} = 0.71 \pm 0.03$ (0.82 ± 0.06) for $H_0 \pm \sigma_{H_0} = 68 \pm 2.8$ (73.24 ± 1.74) $\text{km s}^{-1} \text{Mpc}^{-1}$. Using all ten z_{da} values in the penultimate column of [Table 3](#) we find $z_{\text{da}} = 0.76 \pm 0.07$.

A more reliable summary value of the deceleration-acceleration transition redshift is determined from a weighted mean analysis. Using [Eqs. 1–3](#), we find $z_{\text{da}} = 0.72 \pm 0.05$ (0.84 ± 0.03) for $H_0 \pm \sigma_{H_0} = 68 \pm 2.8$ (73.24 ± 1.74) $\text{km s}^{-1} \text{Mpc}^{-1}$, and using all ten values in the penultimate column of [Table 3](#) we get $z_{\text{da}} = 0.80 \pm 0.02$. **By looking at the fourth and the fifth columns of [Table 3](#) it appears that all the five models discussed here fit better with the lower value of H_0 while the uncertainty in z_{da} is more sensitive to σ_{H_0} .**

These results are listed in [Table 4](#) and compared with the previously computed summary values of [Farooq et al. \(2013b\)](#). Note that only three models (Λ CDM, flat XCDM, and flat ϕ CDM) were considered in [Farooq et al. \(2013b\)](#). Here we also consider non-flat XCDM and non-flat ϕ CDM. We see that there is good agreement between the old and new weighted mean z_{da} for $h = 0.68$, less so for $h = 0.7324$. From [Table 4](#) we see that for a given H_0 the weighted average values of z_{da} for all five models and for the two sets of (non-nested) triplets of models agree to within the error bars.

TABLE 4
 z_{da} SUMMARY

	$h \pm \sigma_h = 0.68 \pm 0.028^{\text{a}}$		$h \pm \sigma_h = 0.7324 \pm 0.0174^{\text{a}}$		Total ^b	
	Here ^c	Previous ^d	Here ^c	Previous ^d	Here ^c	Previous ^d
Simple Averages	0.71 ± 0.03	0.70 ± 0.02	0.82 ± 0.06	0.78 ± 0.06	0.76 ± 0.07	0.74 ± 0.06
Weighted Averages	0.72 ± 0.05	0.69 ± 0.06	0.84 ± 0.03	0.76 ± 0.05	0.80 ± 0.02	0.74 ± 0.04
Simple Averages from Λ CDM and Flat Models	0.73 ± 0.02	...	0.84 ± 0.03	...	0.78 ± 0.06	...
Weighted Averages from Λ CDM and Flat Models	0.73 ± 0.05	...	0.85 ± 0.03	...	0.81 ± 0.03	...
Simple Averages from Non-Flat Models	0.70 ± 0.02	...	0.80 ± 0.06	...	0.75 ± 0.07	...
Weighted Averages from Non-Flat Models	0.70 ± 0.06	...	0.82 ± 0.04	...	0.79 ± 0.03	...

^a Hubble constant in units of $100 \text{ km s}^{-1} \text{Mpc}^{-1}$.

^b Combination of results from both H_0 priors.

^c Estimated using the unbinned data of 38 $H(z)$ measurements from [Table 1](#).

^d Results from [Farooq et al. \(2013b\)](#). We have corrected typos in that paper here.

In this section, we use the 38 Hubble parameter measurements (over $0.07 \leq z \leq 2.36$) listed in Table 1 to determine constraints on the parameters of the five different cosmological models. We use the technique of Farooq et al. (2015) to find constraints on $(\Omega_{m0}, \Omega_{\Lambda})$ in the Λ CDM model, (Ω_{m0}, ω_X) for the spatially-flat XCDM parameterization, (Ω_{m0}, α) in the spatially-flat ϕ CDM model, $(\Omega_{m0}, \omega_X, \Omega_{K0})$ for the XCDM parameterization with space curvature, and $(\Omega_{m0}, \alpha, \Omega_{K0})$ in the ϕ CDM model with space curvature. For the $H(z)$ cosmological test, cosmological parameter constraints depend on the value of the Hubble constant (see, e.g., Samushia et al. 2007). We use two different Gaussian priors for the Hubble constant; the lower value is $68 \pm 2.8 \text{ km s}^{-1} \text{ Mpc}^{-1}$ and the higher is $73.24 \pm 1.74 \text{ km s}^{-1} \text{ Mpc}^{-1}$. The lower value is from a median statistics analysis (Gott et al. 2001) of 553 measurements of H_0 tabulated by Huchra (Chen & Ratra 2011a). It agrees with earlier median statistics estimates of H_0 from smaller compilations (Gott et al. 2001; Chen et al. 2003) and is consistent with a number of other recent determinations of H_0 from Wilkinson Microwave Anisotropy Probe, Atacama Cosmology Telescope, and Planck CMB anisotropy data (Hinshaw et al. 2013; Sievers et al. 2013; Ade et al. 2015; Addison et al. 2016), from BAO measurements (Aubourg et al. 2015; Ross et al. 2015; L’Huillier & Shafieloo 2016), from Hubble parameter data (Chen et al. 2016a), and with what is expected in the standard model of particle physics with only three light neutrino species given current cosmological data (see, e.g. Calabrese et al. 2012). The higher value is a relatively local measurement, based on Hubble Space Telescope data (Riess et al. 2016). It is consistent with other recent local measurements of H_0 (Riess et al. 2011; Freedman et al. 2012; Efstathiou 2014).

We compute the likelihood function $\mathcal{L}(\mathbf{p})$ for the models under discussion using Eq. (18) of Farooq et al. (2013a) for the ranges of the cosmological parameters listed at the end of Sec. 4. We need these likelihood functions for the z_{da} computation of the previous section, which is the main result of the paper. In this section, we use these likelihood functions to constrain cosmological parameters such as the dark energy density.

For the two-parameter models, maximizing the likelihood function $\mathcal{L}(\mathbf{p})$ is performed by minimizing the corresponding $\chi^2(\mathbf{p}) \equiv -2\ln[\mathcal{L}(\mathbf{p})]$ following the procedure of Farooq et al. (2015). The corresponding minimum values of χ^2 and best-fit parameter values for the two-parameter models are summarized in Table 3. 1σ , 2σ , and 3σ confidence contours are computed following the procedure of Farooq et al. (2015) and results are shown in Fig. 2. The generalization of this procedure for the three-parameter models is straightforward and best-fit three-dimensional parameter values and minimum χ^2 are also summarized in Table 3.

For the three-parameter models we next compute three two-dimensional likelihood functions by marginalizing the three-dimensional likelihood function over each of the three parameters (assuming flat priors) in turn. These three two-dimensional likelihood functions are maximized as above and the corresponding best-fit parameter values and minimum χ^2 are listed in Table 5. The confidence contours for these two-dimensional likelihood functions are shown in Fig. 3 for the non-flat XCDM parametrization and in Fig. 4 for the non-flat ϕ CDM model.

To get two one-dimensional likelihood functions from each of the two-dimensional likelihood functions, we marginalize (with a flat prior) over each parameter in turn. We then determine the best-fit parameter values by maximizing each one-dimensional likelihood function and compute 1σ , and 2σ intervals for each parameter in each model and for both H_0 priors. The best-fit parameter values and 1σ and 2σ intervals for the two-parameter models are given in Table 6 and for the three-parameter models, these are given in Table 7.




FIG. 2.— The three panels (from left to right) show 1σ , 2σ , and 3σ red solid (blue dot-dashed) constraint contours for the lower (higher) H_0 prior, for Λ CDM, flat XCDM, and flat ϕ CDM respectively. Red solid (blue empty) circles are the best-fit points for the lower (higher) H_0 prior. The straight dashed lines in the first and second panels correspond to spatially-flat Λ CDM models, the dotted lines demarcate zero-acceleration models, and the shaded area in the upper left-hand corner of the left panel is the region for which there is no big bang. For quantitative parameter best-fit values and ranges see Tables 3 and 6.

The best-fit (two- and three-dimensional) model predictions are shown in Fig. 1, for the five different cosmological models, Λ CDM in red, flat XCDM in blue, flat ϕ CDM in green, non-flat XCDM in orange, and non-flat ϕ CDM in brown, for the two H_0 priors, with $H_0 \pm \sigma_{H_0} = 68 \pm 2.8 \text{ km s}^{-1} \text{ Mpc}^{-1}$ in solid lines and $H_0 \pm \sigma_{H_0} = 73.24 \pm 1.74 \text{ km s}^{-1} \text{ Mpc}^{-1}$

$\text{s}^{-1} \text{Mpc}^{-1}$ in dot-dashed lines.

TABLE 5
TWO-DIMENSIONAL BEST-FIT PARAMETERS FOR THREE-PARAMETER, NON-FLAT MODELS

Model	h Prior ^a	Marginalized Parameter	BF ^b	χ^2_{min}
Non-flat XCDM	0.68 ± 0.028	Ω_{K0}	$\Omega_{m0} = 0.38$ $\omega_X = -0.64$	25.3
		ω_X	$\Omega_{m0} = 0.16$ $\Omega_{K0} = 0.43$	22.5
		Ω_{m0}	$\omega_X = -1.80$ $\Omega_{K0} = 0.47$	27.3
	0.7324 ± 0.0174	Ω_{K0}	$\Omega_{m0} = 0.13$ $\omega_X = -2$	25.0
		ω_X	$\Omega_{m0} = 0.15$ $\Omega_{K0} = 0.39$	22.1
		Ω_{m0}	$\omega_X = -2$ $\Omega_{K0} = 0.41$	26.8
Non-flat ϕ CDM	0.68 ± 0.028	Ω_{K0}	$\Omega_{m0} = 0.28$ $\alpha = 1.33$	25.7
		α	$\Omega_{m0} = 0.26$ $\Omega_{K0} = -0.02$	22.4
		Ω_{m0}	$\alpha = 0.01$ $\Omega_{K0} = 0.19$	28.7
	0.7324 ± 0.0174	Ω_{K0}	$\Omega_{m0} = 0.25$ $\alpha = 0.01$	29.0
		α	$\Omega_{m0} = 0.28$ $\Omega_{K0} = -0.19$	26.6
		Ω_{m0}	$\alpha = 0.01$ $\Omega_{K0} = -0.04$	31.6

^a Hubble constant in units of $100 \text{ km s}^{-1} \text{ Mpc}^{-1}$.

^b Best-fit parameter values.




FIG. 3.— The three panels (from left to right) show 1σ , 2σ , and 3σ two-dimensional constraint contours for the three-parameter, non-flat XCDM parameterization, computed after marginalizing over each of the three parameters in turn. Red (blue) solid lines are for the lower (higher) H_0 prior. Left, center, and right panels correspond to marginalizing over Ω_{K0} , ω_X , and Ω_{m0} respectively. Red (blue) solid circles are the best-fit points for the lower (higher) H_0 prior. Red (blue) dot-dashed lines in the left panel are 1σ , 2σ , and 3σ constraint contours for the lower (higher) H_0 prior for spatially-flat XCDM (see the central panel of Fig. 2). For quantitative parameter best-fit values and ranges see Tables 3, 5, and 7.

While the main purpose of our paper was to improve on the characterization of the deceleration-acceleration transition studied in Farooq & Ratra (2013b) and Farooq et al. (2013b), we see from Fig. 2 and the left panels of Figs. 3 and 4 that the $H(z)$ data by themselves indicate that the cosmological expansion is currently accelerating.

From these figures, it is clear that the $H(z)$ data of Table 1 are very consistent with the standard spatially-flat Λ CDM cosmological model, although even for the two-parameter model constraint contours shown in Fig. 2 there is a large range of dynamical dark energy models as well as spatially-curved models that are consistent with the data. In Figs. 3 and 4 for the non-flat dynamical dark energy models, it is clear that allowing for non-zero space curvature considerably broadens the dynamical dark energy options and vice versa. It is interesting to note that in the non-flat ϕ CDM model Chen et al. (2016b) find that the cosmological data bound on the sum of neutrino masses is considerably weaker than if the model were spatially flat.

While the error bars are large, it is curious that Table 7 entries show that the non-flat XCDM parametrization mildly favors open spatial hypersurfaces while the non-flat ϕ CDM model mildly prefers closed ones.

TABLE 6
ONE-DIMENSIONAL BEST-FIT PARAMETERS AND INTERVALS FOR TWO-PARAMETER MODELS

Model	h Prior ^a	Marginalization Range	BF ^b	1σ intervals	2σ intervals
ACDM	0.68 ± 0.028	$0 \leq \Omega_\Lambda \leq 1.4$	0.23	$0.19 \leq \Omega_{m0} \leq 0.27$	$0.15 \leq \Omega_{m0} \leq 0.30$
		$0 \leq \Omega_{m0} \leq 1$	0.58	$0.46 \leq \Omega_\Lambda \leq 0.69$	$0.32 \leq \Omega_\Lambda \leq 0.80$
	0.7324 ± 0.0174	$0 \leq \Omega_\Lambda \leq 1.4$	0.26	$0.22 \leq \Omega_{m0} \leq 0.29$	$0.19 \leq \Omega_{m0} \leq 0.32$
		$0 \leq \Omega_{m0} \leq 1$	0.79	$0.71 \leq \Omega_\Lambda \leq 0.86$	$0.63 \leq \Omega_\Lambda \leq 0.93$
Flat XCDM	0.68 ± 0.028	$-2 \leq \omega_X \leq 0$	0.27	$0.25 \leq \Omega_{m0} \leq 0.29$	$0.22 \leq \Omega_{m0} \leq 0.31$
		$0 \leq \Omega_{m0} \leq 1$	-0.85	$-0.98 \leq \omega_X \leq -0.73$	$-1.11 \leq \omega_X \leq -0.59$
	0.7324 ± 0.0174	$-2 \leq \omega_X \leq 0$	0.25	$0.23 \leq \Omega_{m0} \leq 0.26$	$0.22 \leq \Omega_{m0} \leq 0.28$
		$0 \leq \Omega_{m0} \leq 1$	-1.07	$-1.17 \leq \omega_X \leq -0.98$	$-1.27 \leq \omega_X \leq -0.89$
Flat ϕ CDM	0.68 ± 0.028	$0 \leq \alpha \leq 5$	0.26	$0.23 \leq \Omega_{m0} \leq 0.28$	$0.20 \leq \Omega_{m0} \leq 0.30$
		$0 \leq \Omega_{m0} \leq 1$	0.53	$0.09 \leq \alpha \leq 1.29$	$0 \leq \alpha \leq 2.9$
	0.7324 ± 0.0174	$0 \leq \alpha \leq 5$	0.24	$0.23 \leq \Omega_{m0} \leq 0.26$	$0.21 \leq \Omega_{m0} \leq 0.28$
		$0 \leq \Omega_{m0} \leq 1$	0	$0 \leq \alpha \leq 0.15$	$0 \leq \alpha \leq 0.46$

^a Hubble constant in units of $100 \text{ km s}^{-1} \text{ Mpc}^{-1}$.

^b Best-fit parameter values.

TABLE 7
ONE-DIMENSIONAL BEST-FIT PARAMETERS AND INTERVALS FOR THREE-PARAMETER, NON-FLAT MODELS

Model	h Prior ^a	Marginalization Range ^b	BF	1σ intervals	2σ intervals
Non-flat XCDM	0.68 ± 0.028	$0 \leq \Omega_{m0} \leq 1$ $-2 \leq \omega_X \leq 0$	0.45	$0.32 \leq \Omega_{K0} \leq 0.55$	$-0.06 \leq \Omega_{K0} \leq 0.66$
		$0 \leq \Omega_{m0} \leq 1$ $-0.7 \leq \Omega_{K0} \leq 0.7$	-0.71	$-1.28 \leq \omega_X \leq -0.59$	$-2 \leq \omega_X \leq -0.49$
		$-2 \leq \omega_X \leq 0$ $-0.7 \leq \Omega_{K0} \leq 0.7$	0.26	$0.21 \leq \Omega_{m0} \leq 0.33$	$0.16 \leq \Omega_{m0} \leq 0.39$
	0.7324 ± 0.0174	$0 \leq \Omega_{m0} \leq 1$ $-2 \leq \omega_X \leq 0$	0.36	$0.28 \leq \Omega_{K0} \leq 0.43$	$0.12 \leq \Omega_{K0} \leq 0.50$
		$0 \leq \Omega_{m0} \leq 1$ $-0.7 \leq \Omega_{K0} \leq 0.7$	-0.98	$-1.54 \leq \omega_X \leq -0.91$	$-2 \leq \omega_X \leq -0.70$
		$-2 \leq \omega_X \leq 0$ $-0.7 \leq \Omega_{K0} \leq 0.7$	0.22	$0.18 \leq \Omega_{m0} \leq 0.26$	$0.10 \leq \Omega_{m0} \leq 0.35$
Non-flat ϕ CDM	0.68 ± 0.028	$0 \leq \Omega_{m0} \leq 1$ $0 \leq \alpha \leq 5$	-0.28	$-0.4 \leq \Omega_{K0} \leq 0.10$	$-0.4 \leq \Omega_{K0} \leq 0.38$
		$0 \leq \Omega_{m0} \leq 1$ $-0.4 \leq \Omega_{K0} \leq 0.4$	0.087	$0 \leq \alpha \leq 2.03$	$0 \leq \alpha \leq 4.07$
		$0 \leq \alpha \leq 5$ $-0.4 \leq \Omega_{K0} \leq 0.4$	0.26	$0.21 \leq \Omega_{m0} \leq 0.30$	$0.17 \leq \Omega_{m0} \leq 0.33$
	0.7324 ± 0.0174	$0 \leq \Omega_{m0} \leq 1$ $0 \leq \alpha \leq 5$	-0.35	$-0.4 \leq \Omega_{K0} \leq -0.08$	$-0.4 \leq \Omega_{K0} \leq 0.08$
		$0 \leq \Omega_{m0} \leq 1$ $-0.4 \leq \Omega_{K0} \leq 0.4$	0	$0 \leq \alpha \leq 0.71$	$0 \leq \alpha \leq 1.25$
		$0 \leq \alpha \leq 5$ $-0.4 \leq \Omega_{K0} \leq 0.4$	0.28	$0.25 \leq \Omega_{m0} \leq 0.31$	$0.22 \leq \Omega_{m0} \leq 0.34$

^a Hubble constant in units of $100 \text{ km s}^{-1} \text{ Mpc}^{-1}$.

^b The three-parameter-dependent likelihood function is integrated over the two parameters in the ranges given in this column and the corresponding best-fit and 1σ and 2σ intervals of the third parameter is computed and listed in the fourth, fifth, and sixth columns respectively of the same row.




FIG. 4.— The three panels (from left to right) show 1σ , 2σ , and 3σ two-dimensional constraint contours for the three-parameter, non-flat ϕ CDM model, computed after marginalizing over each of the three parameters in turn. Red (blue) solid lines are for the lower (higher) H_0 prior. Left, center, and right panels correspond to marginalizing over Ω_{K0} , α , and Ω_{m0} respectively. Red (blue) solid circles are the best-fit points for the lower (higher) H_0 prior. Red (blue) dot-dashed lines in the left panel are 1σ , 2σ , and 3σ constraint contours for the lower (higher) H_0 prior for the spatially-flat ϕ CDM model (see the right panel of Fig. 2). For quantitative parameter best-fit values and ranges see Tables 3, 5, and 7.

7. CONCLUSION

From the new list of $H(z)$ data we have compiled, we find evidence for the cosmological deceleration-acceleration transition to have taken place at a redshift $z_{\text{da}} = 0.72 \pm 0.05$ (0.84 ± 0.03), depending on the value of $H_0 = 68 \pm 2.8$ (73.24 ± 1.74) $\text{km s}^{-1} \text{Mpc}^{-1}$, but otherwise only mildly dependent on other cosmological parameters. In addition, the binned $H(z)$ data in redshift space show qualitative visual evidence for the deceleration-acceleration transition, independent of how they are binned provided the bins are narrow enough, in agreement with that originally found by Farooq et al. (2013b). These $H(z)$ data are consistent with the standard spatially-flat Λ CDM cosmological model but do not rule out non-zero space curvature or dynamical dark energy, especially in models that allow for both. Other data, such as currently available SNIa, BAO, growth factor, or CMB anisotropy data can tighten the constraints on these parameters (see, e.g. Farooq et al. 2015), and it is of interest to study how the other data constrains parameters when used in conjunction with the $H(z)$ data we have compiled here.

ACKNOWLEDGEMENTS

O.F. and M.F. gratefully acknowledge the partial funding from the Department of Physical Sciences, Embry-Riddle Aeronautical University. S.C. and B.R. were supported in part by DOE grant DE-SC0011840.

REFERENCES

- Addison, G. E., et al. 2016, *ApJ*, 818, 132 [arXiv:1511.00055] 6
- Ade, P. A. R. et al. 2015, arXiv:1502.01589 1, 6
- Akarsu, O., Dereli, T., Kumar, S., & Xu, L. 2014, *Eur. Phys. J. Plus*, 129, 22 [arXiv:1305.5190] 1
- Alam, U., Bag, S., & Sahni, V. 2016, arXiv:1605.04707 1
- Alam, S., et al. 2016, arXiv:1607.03155 1, 2, 2, 5
- Aubourg, E. et al. 2015, *Phys. Rev. D*, 92, 123516 [arXiv:1411.1074] 1, 6
- Avsajanishvili, L., Arkhipova, N. A., Samushia, L., & Kahniashvili, T. 2014, *Eur. Phys. J. C*, 79, 3127 [arXiv:1406.0407] 4
- Avsajanishvili, L., Samushia, L., Arkhipova, N. A., & Kahniashvili, T. 2015, arXiv:1511.09317 4
- Bamba, K. et al. 2014, *Phys. Rev. D*, 89, 083518 [arXiv:1309.6413] 1
- Basse, T., et al. 2014, *JCAP*, 1405, 021 [arXiv:1304.2321] 1
- Blake, C. et al. 2012, *MNRAS*, 425, 405 [arXiv:1204.3674] 1, 2, 5
- Busca, N. G. et al. 2013, *A & A*, 552, A96 [arXiv:1211.2616] 1, 2
- Cai, R.-G., Guo, Z.-K., & Yang, T. 2015, *Phys. Rev. D*, 93, 43517 [arXiv:1509.06283] 2
- Calabrese, E., Archidiacono, M., Melchiorri, A., & Ratra, B. 2012, *Phys. Rev. D*, 86, 043520 [arXiv:1205.6753] 1, 6
- Campanelli, L. et al. 2012, *Eur. Phys. J. C*, 72, 2218 [arXiv:1110.2310] 4
- Capozziello, S., Farooq, O., Luongo, O., & Ratra, B. 2014, *Phys. Rev. D*, 90, 044016 [arXiv:1403.1421] 1
- Chen, G., & Ratra, B. 2003, *PASP*, 115, 1143 [arXiv:astro-ph/0302002] 3
- Chen, G., Gott, J. R., & Ratra, B. 2003, *PASP*, 115, 1269 [arXiv:astro-ph/0308099] 6
- Chen, G., & Ratra, B. 2004, *ApJ*, 612, L1 [arXiv:astro-ph/0405636] 4
- Chen, G., & Ratra, B. 2011a, *PASP*, 123, 1127 [arXiv:1105.5206] 1, 6
- Chen, Y., Kumar, S., & Ratra, B. 2016a, arXiv:1606.07316 1, 3, 6
- Chen, Y., & Ratra, B. 2011b, *Phys. Lett. B*, 703, 406 [arXiv:1106.4294] 1
- Chen, Y., et al. 2015, *JCAP*, 1502, 010 [arXiv:1312.1443] 1
- Chen, Y., et al. 2016b, [arXiv:1603.07115] 4, 6
- Chimento, L. P., & Richarte, M. G. 2013, *Eur. Phys. J. C*, 73, 2497 [arXiv:1308.0860] 1
- Chuang, C.-H., & Wang, Y. 2013, *MNRAS*, 435, 255 [arXiv:1209.0210] 2
- Crandall, S., Houston, S., & Ratra, B. 2015, *Mod. Phys. Lett. A*, 30, 1550123 [arXiv:1409.7332] 3
- Crandall, S., & Ratra, B. 2014, *Phys. Lett. B*, 732, 330 [arXiv:1311.0840] 3
- Crandall, S., & Ratra, B. 2015, *ApJ*, 815, 87 [arXiv:1507.07940] 3
- Dankiewicz, T., Dabrowski, M. P., Martins, C. J. A. P., & Vielzeuf, P. E. 2014, *Phys. Rev. D*, 89, 083514 [arXiv:1402.0520] 1
- Delubac, T., et al. 2015, *A&A*, 574, A59 [arXiv:1404.1801] 1, 2, 2
- Ding, X., et al. 2015, *ApJ*, 803, L22 [arXiv:1503.04923] 3
- Duan, X.-W., Zhou, M., & Zhang, T.-J. 2016, arXiv:1605.03947 2
- Efstathiou, G. 2014, *MNRAS*, 440, 1138 [arXiv:1311.3461] 6
- Farooq, O. 2013, Kansas State University PhD Thesis [arXiv:1309.3710] 4
- Farooq, O., Crandall, S., & Ratra, B. 2013b, *Phys. Lett. B*, 726, 72 [arXiv:1305.1957] 1, 1, 3, 3, 3, 5, 4, 6, 7
- Farooq, O., Mania, D., & Ratra, B. 2013a, *ApJ*, 764, 138 [arXiv:1211.4253] 5, 6
- Farooq, O., Mania, D., & Ratra, B. 2015, *ApSS*, 357, 11 [arXiv:1308.0834] 1, 1, 4, 5, 6, 7
- Farooq, O., & Ratra, B. 2013a, *Phys. Lett. B*, 723, 1 [arXiv:1212.4264] 1
- Farooq, O., & Ratra, B. 2013b, *ApJ*, 766, L7 [arXiv:1301.5243] 1, 1, 2, 3, 6
- Ferreira, P. C., Pavón, D., & Carvalho, J. C. 2013, *Phys. Rev. D*, 88, 083503 [arXiv:1310.2160] 1
- Font-Ribera, A., et al. 2014, *JCAP*, 1405, 027 [arXiv:1311.1767] 1, 2, 2
- Forte, M. 2014, *Gen. Rel. Grav.*, 46, 1811 [arXiv:1311.3921] 1
- Freedman, W. L., et al. 2012, *ApJ*, 758, 24 [arXiv:1208.3281] 6
- Gosena, M., & Coles, P. 2015, arXiv:1502.04020 4
- Gott, J. R., Vogeley, M. S., Podariu, S., & Ratra, B. 2001, *ApJ*, 549, 1 [arXiv:astro-ph/0006103] 3, 6
- Gruber, C., & Luongo, O. 2014, *Phys. Rev. D*, 89, 103516 [arXiv:1309.3215] 1
- Guo, R.-Y., & Zhang, X. 2016, *Eur. Phys. J. C*, 76, 163 [arXiv:1512.07703] 1
- Hinshaw, G., et al. 2013, *ApJS*, 208, 19 [arXiv:1212.5226] 1, 6
- Hodge, J. A., Zeimann, G. R., Becker, R. H., & White, R. L. 2009, *AJ*, 138, 900 [arXiv:0907.1981] 3
- L'Huillier, B., & Shafieloo, A. 2016, [arXiv:1606.06832] 1, 6
- Joyce, A., Lombriser, L., & Schmidt, F. 2016, arXiv:1601.06133 1
- Lima, N. A., Liddle, A. R., Sahlén, M., & Parkinson, D. 2016, *Phys. Rev. D*, 93, 063506 [arXiv:1501.02678] 4
- Martin, J., 2012, *C. R. Physique*, 13, 566 [arXiv:1205.3365] 1
- Meng, X.-L., Wang, X., Li, S.-Y., & Zhang, T.-J. 2015, arXiv:1507.02517 1, 2
- Moresco, M., 2015, *MNRAS*, 450, L16 [arXiv:1503.01116] 1, 2, 2
- Moresco, M., et al. 2012, *JCAP*, 1208, 006 [arXiv:1201.3609] 1, 1, 2
- Moresco, M., et al. 2016, *JCAP*, 1605, 014 [arXiv:1601.01701] 1, 1, 2, 2
- Mukherjee, A. & Banerjee, N. 2016, *Phys. Rev. D*, 93, 043002 [arXiv:1601.05172] 1
- Muthukrishna, D., & Parkinson, D. 2016, arXiv:1607.01884 2
- Nunes, R. C., Pan, S., & Saridakis, E. N. 2016, arXiv:1606.04359 2
- Pavlov, A., Farooq, O., & Ratra, B. 2014, *Phys. Rev. D*, 90, 023006 [arXiv:1312.5285] 4
- Pavlov, A., Samushia, L., & Ratra, B. 2012, *ApJ*, 760, 19 [arXiv:1206.3123] 1
- Pavlov, A., Westmoreland, S., Saaidi, K., & Ratra, B. 2013, *Phys. Rev. D*, 88, 123513 [arXiv:1307.7399] 1, 4, 4
- Peebles, P. J. E. 1984, *ApJ*, 284, 439 4
- Peebles, P. J. E., & Ratra, B. 1988, *ApJ*, 325, L17 4, 4, 4
- Podariu, S., Nugent, P., & Ratra, B. 2001a, *ApJ*, 553, 39 [arXiv:astro-ph/0008281] 1
- Podariu, S., et al. 2001b, *ApJ*, 559, 9 [arXiv:astro-ph/9910527] 3
- Podariu, S., & Ratra, B. 2000, *ApJ*, 532, 109 [arXiv:astro-ph/0102264] 4

- Podariu, S., et al. 2001, ApJ, 559, 9 [arXiv:astro-ph/9910527]
- Qi, J.-Z., Zhang, M.-J., & Liu, W.-B. 2016, arXiv:1606.00168 [2](#)
- Rani, N., et al. 2015, JCAP, 1512, 045 [arXiv:1503.08543] [2](#)
- Ratra, B., & Peebles, P. J. E. 1988, Phys. Rev. D, 37, 3406 [4](#)
- Ratra, B., & Vogeley, M. 2008, PASP, 120, 235 [arXiv:0706.1565] [1](#)
- Riess, A. G., et al. 2011, ApJ, 730, 119 [arXiv:1103.2976] [6](#)
- Riess, A. G., et al. 2016, arXiv:1604.01424 [6](#)
- Ross, A. J. et al. 2015, MNRAS, 449, 835 [arXiv:1409.3242] [1](#), [6](#)
- Samushia, L. 2009, Kansas State University PhD Thesis [arXiv:0908.4597] [4](#)
- Samushia, L., Chen, G., & Ratra, B. 2007, arXiv:0706.1963 [6](#)
- Samushia, L., Dev, A., Jain, D., & Ratra, B. 2010, Phys. Lett. B, 693, 509 [arXiv:0906.2734] [4](#)
- Samushia, L., & Ratra, B. 2006, ApJ, 650, L5 [arXiv:astro-ph/0607301] [1](#)
- Samushia, L., & Ratra, B. 2010, ApJ, 714, 1347 [arXiv:0905.3836] [4](#)
- Santos, L., Cabella, P., Balbi, A., & Vittorio, N. 2013, Phys. Rev. D, 88, 043505 [arXiv:1307.8919] [1](#)
- Sievers, J. L., et al. 2013, JCAP, 1310, 060 [arXiv:1301.0824] [1](#), [6](#)
- Simon, J., Verde, L., & Jimenez, R. 2005, Phys. Rev. D, 71, 123001 [arXiv:astro-ph/0412269] [1](#), [1](#), [2](#)
- Solà, J., Gómez-Valent, A., & de Cruz Pérez, J. 2016, arXiv:1602.02103 [2](#)
- Stern, D., et al. 2010, JCAP 1002 (2010) 008 [arXiv:0907.3152] [1](#), [2](#)
- Sutherland, W., & Rothnie, P. 2015, MNRAS, 446, 3863 [arXiv:1503.01122] [2](#)
- Verde, L., Protopapas, P., & Jimenez, R. 2014, Phys.Dark Univ., 5-6, 307 [arXiv:1403.2181] [1](#)
- Weinberg, D. H., et al. 2014, Phys.Rept., 530, 87 [arXiv:1201.2434] [1](#)
- Yu, H. & Wang, F. Y. 2016, arXiv:1605.02483 [2](#)
- Zhang, C., et al. 2012, arXiv:1207.4541 [astro-ph.CO] [1](#), [2](#)
- Zhang, C., et al. 2014, Res. Astron. Astrophys., 14, 1221 [arXiv:1207.4541]
- Zhang, M.-J., & Xia, J.-Q. 2016, arXiv:1606.04398 [2](#)
- Zheng, X., et al. 2016, arXiv:1604.07910 [3](#)