



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



ELSEVIER

Available online at www.sciencedirect.com

SciVerse ScienceDirect

journal homepage: <http://www.e-biomedicine.com>

Review article

Human coronaviruses: Clinical features and phylogenetic analysis

Shih-Wen Li, Cheng-Wen Lin*

Department of Medical Laboratory Science and Biotechnology, China Medical University, Taichung, Taiwan

ARTICLE INFO

Article history:

Received 17 October 2012

Received in revised form

23 November 2012

Accepted 19 December 2012

Available online 1 February 2013

Keywords:

human betacoronavirus 2c EMC2012

human coronavirus

phylogenetic tree

severe acute respiratory syndrome

coronavirus (SARS-CoV)

ABSTRACT

Strains of human coronavirus (HCoV), namely HCoV-OC43, HCoV-229E, HCoV-NL63, and HCoV-HKU1, primarily infect the upper respiratory and gastrointestinal tracts and are the most common cause of non-rhinovirus-induced common cold in humans. Although the manifestations of coronavirus infection (i.e., rhinorrhea, sneezing, cough, nasal obstruction, and bronchitis) are generally self-limiting in healthy adults, certain strains such as HCoV-NL63 and HCoV-HKU1 can cause severe lower respiratory tract infection and febrile seizure, especially in infants, people of advanced age, and immunocompromised hosts. In 2003, a novel HCoV strain was identified as the causative agent of the severe acute respiratory syndrome (SARS) epidemic that began in Asia in 2002. The strain has hence been referred to as SARS-CoV. In addition, as recently as September 2012, another novel HCoV, human betacoronavirus 2c EMC2012, was identified as being the cause of fever, renal failure, pneumonia, and severe respiratory distress in two patients in the Middle East. Phylogenetic analysis has revealed highly conserved sequences of ORF1ab, spike, nucleocapsid, and envelope protein genes, but not membrane protein genes, between human betacoronavirus 2c EMC2012 and SARS-CoV. This review focuses on the differences in the genomes of certain HCoV strains, the pathogenesis of said strains, and recent developments in the establishment of therapeutic agents that might aid in the treatment of patients with such infections.

Copyright © 2013, China Medical University. Published by Elsevier Taiwan LLC. All rights reserved.

1. Coronaviruses

Coronaviruses, which are enveloped, single-stranded, positive-sense RNA viruses belonging to the subfamily Coronavirinae in the family Coronaviridae, primarily cause respiratory and enteric diseases in mammals and birds [1–9]. The genus is subdivided into three main groups (alpha, beta, and gamma) based on the genetic and serological properties of

each virus [10–13]. Most coronaviruses in the three groups infect animals or birds only. Of the viruses that infect human hosts, human coronavirus (HCoV)-229E and HCoV-OC43, both alphacoronaviruses, are the best characterized and are responsible for most non-rhinovirus-induced cases of the common cold [14]. Other respiratory pathogens in the coronavirus genera that have recently been characterized include the alpha virus HCoV-NL63 and the beta virus HCoV-HKU1.

* Corresponding author. Department of Medical Laboratory Science and Biotechnology, China Medical University, Number 91, Hsueh-Shih Road, Taichung 404, Taiwan.

E-mail address: cwlin@mail.cmu.edu.tw (C.-W. Lin).

2211-8020/\$ – see front matter Copyright © 2013, China Medical University. Published by Elsevier Taiwan LLC. All rights reserved.
<http://dx.doi.org/10.1016/j.biomed.2012.12.007>

Those two viruses tend to be isolated from young patients, patients of advanced age, and immunocompromised hosts [13,15,16]. Recently, however, more virulent strains have emerged, including severe acute respiratory syndrome (SARS)-associated coronavirus (SARS-CoV), a betacoronavirus that was responsible for the outbreak of SARS in Asia and other countries in 2002–2003, and a novel HCoV, 2c EMC2012, which was identified as the causative agent of rapidly progressive acute respiratory infection in two men from the Middle East in 2012 [17–19].

Coronavirus virions typically range in size from 80 nm to 100 nm in diameter, contain an RNA genome of approximately 26–32 kb in size, and are characterized microscopically by their crown- or halo-shaped appearance [20–26]. The genome of most coronaviruses encodes two replicases, expressed in the form of two polyproteins, open reading frame (ORF)1a, measuring approximately 450 kDa, and ORF1ab, measuring approximately 750 kDa. These polyproteins are processed into a number of non-structural (NS) proteins and four structural proteins, namely spike (S) protein, envelope (E) protein, membrane (M) protein, and nucleocapsid (N) protein (Fig. 1) [3,27]. ORF1a and ORF1ab are predecessors of NS proteins, being cleaved by papain-like protease (PLpro, nsp3) and 3C-like protease (3CLpro, nsp5) to generate the final NS proteins [28,29]. Some members of betacoronavirus subgroup A also code for a shorter spike-like protein called hemagglutinin esterase [30–32]. Studies have shown that, at least in SARS-CoV, a receptor-binding domain on the S protein mediates the attachment of said virus to its cellular receptor, angiotensin-converting enzyme 2 (ACE2) [33,34]. Lin et al found that HCoV-NL63 also makes use of the S protein-ACE2 mechanism for cell entry [35].

Very recently, van Boheemen et al. and Xingyi Ge et al. constructed a phylogenetic tree of coronaviruses using the polymerase gene nsp12 and found that coronaviruses could be grouped according to the presence of ORFs, namely Groups 1, 2a, 2b, 2c, 2d, and 3 [36,37]. According to their data, HCoV-229E, HCoV-NL63, porcine epidemic diarrhea virus, and transmissible gastroenteritis virus belong to Group 1; HCoV-OC43 and mouse hepatitis virus belong to Group 2a; SARS-CoV belongs to Group 2b; human betacoronavirus 2c EMC2012 belongs to Group 2c; and avian infectious bronchitis virus belongs to Group 3 [36,37]. Other phylogenetic trees that have been constructed in our laboratory have revealed that 53% of the ORF1ab sequences, 33% of the envelope protein

sequences, 45% of the nucleocapsid protein sequences, 34% of the spike protein sequences, and 41% of the membrane protein sequences in SARS-CoV are conserved in human betacoronavirus 2c EMC2012, indicating that SARS-CoV and betacoronavirus 2c EMC2012 are very similar (Fig. 2) [36–39].

2. Epidemiology and pathogenesis of SARS coronavirus

SARS-CoV is the causative agent of SARS and causes an atypical pneumonia that spread rapidly throughout parts of Asia, North America, and Europe during 2002–2003 [37,38]. Shortly after the first case of SARS was reported in the Chinese province of Guangdong in November 2002, the disease quickly spread worldwide, with cases having been reported in some 30 countries in 2003. By the time the outbreak had been contained, over 8000 cases had been confirmed and 916 people had died due to the disease [40–45]. According to the World Health Organization, the mortality rate associated with SARS was >10% [46,47]. People of advanced age were at the greatest risk of death, with a case-fatality rate approaching 50% in people >65 years who were infected by the virus [48].

The major route of transmission of SARS-CoV is close person-to-person contact, primarily via contact with aerosolized droplets or other bodily fluids. Many studies have shown that this virus can survive for more than 3 days outside the human body in sputum and feces, resulting in the potential for continuous infection during an epidemic. In addition to causing respiratory symptoms, chills, headache, muscle ache, and fever, infection with SARS-CoV has been shown to result in severe diffuse pneumonia, pulmonary fibrosis, and in the most severe cases, death. Pathological examinations of specimens taken from patients with SARS-related acute lung injury, acute respiratory distress syndrome, and pulmonary fibrosis show bronchial epithelial cell exfoliation, cilia loss, multinucleated syncytia cells, squamous epithelial tissue deformation, and increased levels of immune response cells in lung tissue [49,50]. In addition, clinical laboratory tests have shown that patients with SARS often present with thrombocytopenia and leukopenia [51,52] as well as overexpression of serum inflammatory cytokines, including interferon (IFN)- γ , interleukin (IL)-18, transforming




Fig. 1 – Coronavirus genome. ORF1a and 1b are located at the 5'-terminal 2/3 gene of the coronavirus and encode two polyproteins, namely pp1a (~450 kDa) and pp1ab (~750 kDa). The four structural proteins in coronavirus include the spike (S) protein, envelope (E) protein, membrane (M) protein, and nucleocapsid (N) protein.

growth factor (TGF)- β 1, tumor necrosis factor (TNF)- α , and IL-6 [50,53].

SARS-CoV infection can cause bronchial epithelial cell peeling, cilia damage, the formation of multinucleated giant cells, squamous cell aplasia, alveolar interstitial fiber cell hyperplasia, and fibrotic lung disease [54,55]. Although blood tests in some patients with SARS-CoV infection show a normal white blood cell count, most show evidence of leukopenia, lymphopenia, and thrombocytopenia. Other biochemical signs typically exhibited in blood samples from patients with SARS include abnormal levels of proteins indicative of abnormal liver function, increased C-reactive protein levels, increased levels of creatine phosphokinase and lactate dehydrogenase, overexpression of inflammatory cytokines IL-1, IL-6, IL-8, IL-12, IL-18, TGF- β 1, TNF- α , IFN- γ -inducible Protein 10 (IP-10), IFN- γ , monokine induced by IFN- γ (MIG) as well as overexpression of chemokines chemokine [C-C motif] ligand 2 (CCL2)/monocyte chemoattractant protein-1 (MCP-1), CCL3/MIP-1 α , and chemokine [C-X-C motif] (CXCL10)/IP-10 [50,53,56–62]. In the early stage of infection, TGF- β 1 has been shown to be markedly elevated in serum and in lung tissue [50,63,64]. In addition, patients with late-phase SARS-related acute respiratory distress syndrome and pulmonary fibrosis often present with activated T-helper type-1 cells [53,65], neutrophil aggregation, the accumulation of immune response cells, such as monocytes, macrophages, T cells, and B cells in the lungs [50], and rapid depletion of T cells [57–62].

The most effective treatment for patients with SARS-CoV infection is the combination of ribavirin and steroids; however, this treatment is associated with a number of side effects. Ribavirin can cause red blood cell rupture, anemia, and death, while steroid treatment can cause stomach ulcers, edema, cardiac stress, and osteoporosis [66]. Controlled trials of drugs for the treatment of patients with SARS are urgently needed.

The innate immune response of type I IFN has a vital role in anti-viral replication, but SARS-CoV was demonstrated to antagonize type I IFN in mouse models [67,68]. Mice lacking innate immunity suffer from signal transducer and activator of transcription-1 (STAT1) or myeloid primary response gene 88 (MyD88) deficiency and higher SARS-CoV infection rates [67,68]. In addition, it has been reported that SARS-CoV-infected macrophages release CXCL10/IFN- γ -inducible protein 10 and CCL2/monocyte chemoattractant protein 1, but do not induce the production of IFN- β [69].

Several SARS-CoV structural and NS proteins and genomes have been shown to participate in immune regulation and to induce cytokine expression. For example, the nucleocapsid of the SARS-CoV virion has been shown to induce the expression of plasminogen activator-1 via a Smad3-dependent induction of TGF- β 1 expression [70–72], where Smad3 is an intracellular protein that transduces extracellular signals from TGF- β ligands to the nucleus, where they activate downstream TGF- β gene transcription. Nucleocapsid protein was also identified as antagonizing IFN- β [73]. Furthermore, baculovirus-synthesized SARS-CoV spike protein has been shown to induce activator protein-1 (AP-1) and IL-8 upregulation by activating mitogen-activated protein kinases (MAPKs) in lung cells [74]. Moreover, it was found that spike protein induces innate immune response through the activation of the nuclear factor-kappaB (NF- κ B) pathway [75]. In addition, Law et al

found that SARS-CoV NSP1 induces CCL5, CXCL10, and CCL3 mRNA expression via the activation of NF- κ B [76]. The 3a DNA vaccine is reported to stimulate IFN- γ production mainly by stimulating the production of CD8⁺ T cells, and IL-2 mainly by stimulating the production of CD4⁺ T cells [77]. Overall, structural and NS proteins and also SARS genomes are demonstrated to interfere with cell immunity.

Unlike other coronaviruses, the SARS-CoV genome only codes for one papain-like protease (PLpro). PLpro recognizes the amino acid fragments of LNGG and cleaves polyprotein replicase 1a at NS1/2, NS2/3, and NS3/4 boundaries, resulting in a mature viral protein [29,78]. In a cell model of SARS-CoV infection, Speigal et al showed that PLpro de-ubiquitinating/de-ISGylating activity was associated with the inhibition of type I IFN expression [79]. Other studies have demonstrated that PLpro inhibits polyI:C-induced activation of IFN- β , NF- κ B, and IFN-stimulated response element by blocking the export of phosphorylated regulatory factor 3 into the nucleus, thereby preventing the biosynthesis of type I IFN [80,81]. In addition, Li et al showed that elevated levels of PLpro result in an attenuation of type I IFN-induced activation of the IFN-stimulated response element and AP-1, and that PLpro downregulates ERK1 (MAPK3) by upregulating the ubiquitin–proteasome system and suppressing the interaction between ERK1 and STAT1, resulting in type I IFN antagonism of SARS-CoV PLpro [82]. Furthermore, PLpro has been shown to induce TGF- β 1, IL-1 α , and CCL5 expression as well as triggering TGF- β 1 production via ubiquitin proteasome-, p38 MAPK-, and ERK1/2-mediated signaling [46]. Therefore, it appears that PLpro plays an important role in the maturation process of SARS-CoV and in avoiding detection by the innate immune response system.

3. HCoV-NL63

HCoV-NL63, a Group 1 coronavirus, was first identified in a child with bronchiolitis in the Netherlands in 2004. Since then, cases of HCoV-NL63 have been documented worldwide. Infection is most prevalent during the winter months, tends to affect children under the age of 6 years, normally manifests as cough, fever, sore throat, rhinitis, expectoration, and upper and lower respiratory tract infection, such as bronchitis, bronchiolitis or pneumonia and has been shown to be associated with croup [83–85].

4. HCoV-HKU1

HCoV-HKU1, a Group 2 coronavirus that was first identified in an adult with chronic pulmonary disease in Hong Kong in 2005 [86,87], causes rhinorrhea, fever, coughing, and wheezing, and can result in bronchiolitis and pneumonia if left untreated [15,88].

5. Human betacoronavirus 2c EMC/2012

Human betacoronavirus 2c EMC2012 first appeared in the Middle East in April 2012 and was laboratory proven in nine




Fig. 2 – Phylogenetic trees. Phylogenetic trees were constructed with (A) ORF1ab, (B) nucleocapsid, (C) membrane, (D) spike, and (E) envelope protein. The phylogenetic trees were constructed using MEGA5 software [36–39].

patients by December 30, 2012 [17–19,89–92]. All of the laboratory-confirmed cases have been reported in Qatar (two cases), Saudi Arabia (five cases) and Jordan (two cases) [92]. All patients were severely ill, and five have died. The first confirmed case of betacoronavirus 2c EMC2012 infection occurred in Saudi Arabia. The patient presented with symptoms and signs of acute respiratory illness (high fever, cough, shortness of breath, and difficulty in breathing) and died in hospital. The second confirmed case was diagnosed in a man of Qatari origin who had recently travelled to Saudi Arabia. The patient presented with respiratory disease and renal failure. He was sent to London for treatment and recovered from the disease in hospital [89,90]. The recent three confirmed cases in Saudi Arabia are epidemiologically linked and occurred in one family living within the same household in Saudi Arabia, and two of these have died [92]. In these three cases, at least two family members with direct personal contact increases the suspicion that person-to-person transmission may have occurred [92]. Two confirmed cases in Jordan were discovered through the testing of stored samples from a cluster of pneumonia cases in healthcare workers that occurred in April 2012. Both of these patients died [92].

This novel coronavirus was identified by real-time polymerase chain reaction based on pan-coronavirus primers [93–95] and there is evidence that the virus might have originated from bats [94]. It is not clear whether the symptoms and signs in these patients are typical of infection with this virus and no specific treatment recommendations have been made thus far. Acute respiratory support is advised for hospitalized patients with severe symptoms.

Like other coronaviruses, 2c EMC2012 is a very fragile virus, with a survival time not exceeding 24 hours outside the body. The pathogenic function of human betacoronavirus 2c EMC2012 PLpro(s) is not clear. Preliminary studies suggest that maturation of viral replication involves the cleavage of polyprotein replicase 1a at NS1/2, NS2/3, and NS3/4 boundaries, a mechanism shared by other coronaviruses [29,78]. Moreover, phylogenetic tree analyses have shown that human betacoronavirus 2c EMC2012 and SARS-CoV have a similar ORF1ab subgenome (Fig. 2). Further studies are needed to differentiate between 2c EMC2012 and SARS-CoV.

6. Conclusion

HCoVs cause upper respiratory and gastrointestinal tract problems and the common cold in humans in worldwide, 2012 HCoVs cause upper respiratory and gastrointestinal tract problems and the common cold in humans in worldwide. The major method of transmission of HCoVs is from person-to-person and by droplet infection. Since the first two HCoVs, HCoV-229E and HCoV-OC43, were identified and studied extensively from the 1960s to 1980s, SARS-CoV has caused a mortality rate >10% globally in 2003 [14,45]. HCoV-NL63 and HCoV-HKU1 were identified in 2004 and 2005 [13,15,16]. Since then, a novel coronavirus, human betacoronavirus 2c EMC2012, appeared in Saudi Arabia in April 2012, that has so far infected nine people and resulted in five deaths [92]. In the phylogenetic trees constructed with ORF1ab, E, N, S, and M, human betacoronavirus 2c EMC2012, has shown high

similarities with SARS-CoV. All of these facts prove that novel coronaviruses have appeared in recent years and are becoming more severe, causing high mortality in humans.

REFERENCES

- [1] Lai MM. Coronavirus: organization, replication and expression of genome. *Ann Rev Microbiol* 1990;44:303–33.
- [2] Lai MM. Background paper. Transcription and replication of coronavirus RNA: a 1989 update. *Adv Exp Med Biol* 1990;276:327–33.
- [3] Gorbalenya AE, Koonin EV, Donchenko AP, Blinov VM. Coronavirus genome: prediction of putative functional domains in the non-structural polyprotein by comparative amino acid sequence analysis. *Nucleic Acids Res* 1989;17:4847–61.
- [4] Prentice E, Denison MR. The cell biology of coronavirus infection. *Adv Exp Med Biol* 2001;494:609–14.
- [5] Loa CC, Lin TL, Wu CC, Bryan TA, Thacker HL, Hooper T, et al. Purification of turkey coronavirus by Sephacryl size-exclusion chromatography. *J Virol Methods* 2002;104:187–94.
- [6] Wunschmann A, Frank R, Pomeroy K, Kapil S. Enteric coronavirus infection in a juvenile dromedary (*Camelus dromedarius*). *J Vet Diagn Invest* 2002;14:441–4.
- [7] Theamboonlers A, Samransamruajkit R, Thongme C, Amonsin A, Chongsrisawat V, Poovorawan Y. Human coronavirus infection among children with acute lower respiratory tract infection in Thailand. *Intervirology* 2007;50:71–7.
- [8] Pedersen NC, Allen CE, Lyons LA. Pathogenesis of feline enteric coronavirus infection. *J Feline Med Surg* 2008;10:529–41.
- [9] van der Hoek L, Ihorst G, Sure K, Vabret A, Dijkman R, de Vries M, et al. Burden of disease due to human coronavirus NL63 infections and periodicity of infection. *J Clin Virol* 2010;48:104–8.
- [10] Cavanagh D. Coronaviruses in poultry and other birds. *Avian Pathol* 2005;34:439–48.
- [11] Esper F, Weibel C, Ferguson D, Landry ML, Kahn JS. Coronavirus HKU1 infection in the United States. *Emerg Infect Dis* 2006;12:775–9.
- [12] Vijgen L, Keyaerts E, Zlateva K, Van Ranst M. Identification of six new polymorphisms in the human coronavirus 229E receptor gene (aminopeptidase N/CD13). *Int J Infect Dis* 2004;8:217–22.
- [13] van der Hoek L, Pyrc K, Jebbink MF, Vermeulen-Oost W, Berkhout RJ, Wolthers KC, et al. Identification of a new human coronavirus. *Nat Med* 2004;10:368–73.
- [14] Hamre D, Procknow JJ. A new virus isolated from the human respiratory tract. *Proc Soc Exp Biol Med* 1966;121:190–3.
- [15] Sloots TP, McErlean P, Speicher DJ, Arden KE, Nissen MD, Mackay IM. Evidence of human coronavirus HKU1 and human bocavirus in Australian children. *J Clin Virol* 2006;35:99–102.
- [16] Woo PC, Lau SK, Huang Y, Tsoi HW, Chan KH, Yuen KY. Phylogenetic and recombination analysis of coronavirus HKU1, a novel coronavirus from patients with pneumonia. *Arch Virol* 2005;150:2299–311.
- [17] Health Protection Agency, HPA Press release. Acute respiratory illness associated with a new virus identified in the UK. London: HPA. Available at: <http://www.hpa.org.uk/NewsCentre/NationalPressReleases/2012PressReleases/120923acuterespiratoryillnessidentified/>; 2012 [accessed 06.01.13].
- [18] European Centre for Disease Prevention and Control, Rapid Risk Assessment: Severe respiratory disease associated with a novel coronavirus, ECDPC, September 24, 2012. Available

- at: <http://ecdc.europa.eu/en/publications/Publications/RRA-Novel-coronavirus-final20120924.pdf> [accessed 06.01.13].
- [19] Centers for Disease Control and Prevention (CDC). Severe respiratory illness associated with a novel coronavirus - Saudi Arabia and Qatar, 2012. *MMWR Morb Mortal Wkly Rep* 2012;61:820.
- [20] Herold J, Raabe T, Siddell S. Molecular analysis of the human coronavirus (strain 229E) genome. *Arch Virol* 1993; 7:63–74.
- [21] Herold J, Raabe T, Siddell SG. Characterization of the human coronavirus 229E (HCV 229E) gene 1. *Adv Exp Med Biol* 1993; 342:75–9.
- [22] Lai MM, Liao CL, Lin YJ, Zhang X. Coronavirus: how a large RNA viral genome is replicated and transcribed. *Infect Agents Dis* 1994;3:98–105.
- [23] Haijema BJ, Volders H, Rottier PJ. Switching species tropism: an effective way to manipulate the feline coronavirus genome. *J Virol* 2003;77:4528–38.
- [24] Thiel V, Ivanov KA, Putics A, Hertzog T, Schelle B, Bayer S, et al. Mechanisms and enzymes involved in SARS coronavirus genome expression. *J Gen Virol* 2003;84:2305–15.
- [25] Havlickova M. [Newly identified representatives of the Coronaviridae family.]. *Klin Mikrobiol Infekc Lek* 2005;11:70.
- [26] Gonzalez JM, Gomez-Puertas P, Cavanagh D, Gorbalenya AE, Enjuanes L. A comparative sequence analysis to revise the current taxonomy of the family Coronaviridae. *Arch Virol* 2003;148:2207–35.
- [27] Woo PC, Lau SK, Chu CM, Chan KH, Tsoi HW, Huang Y, et al. Characterization and complete genome sequence of a novel coronavirus, coronavirus HKU1, from patients with pneumonia. *J Virol* 2005;79:884–95.
- [28] Rota PA, Oberste MS, Monroe SS, Nix WA, Campagnoli R, Icenogle JP, et al. Characterization of a novel coronavirus associated with severe acute respiratory syndrome. *Science* 2003;300:1394–9.
- [29] Ziebhuh J. Molecular biology of severe acute respiratory syndrome coronavirus. *Curr Opin Microbiol* 2004;7:412–9.
- [30] Zhang XM, Kousoulas KG, Storz J. The hemagglutinin/esterase gene of human coronavirus strain OC43: phylogenetic relationships to bovine and murine coronaviruses and influenza C virus. *Virology* 1992;186:318–23.
- [31] Klausegger A, Strobl B, Regl G, Kaser A, Luytjes W, Vlasak R. Identification of a coronavirus hemagglutinin-esterase with a substrate specificity different from those of influenza C virus and bovine coronavirus. *J Virol* 1999;73:3737–43.
- [32] Popova R, Zhang X. The spike but not the hemagglutinin/esterase protein of bovine coronavirus is necessary and sufficient for viral infection. *Virology* 2002;294:222–36.
- [33] Li W, Choe H, Farzan M. Insights from the association of SARS-CoV S-protein with its receptor, ACE2. *Adv Exp Med Biol* 2006;581:209–18.
- [34] Li W, Wong SK, Li F, Kuhn JH, Huang IC, Choe H, et al. Animal origins of the severe acute respiratory syndrome coronavirus: insight from ACE2-S-protein interactions. *J Virol* 2006;80:4211–9.
- [35] Lin HX, Feng Y, Tu X, Zhao X, Hsieh CH, Griffin L, et al. Characterization of the spike protein of human coronavirus NL63 in receptor binding and pseudotype virus entry. *Virus Res* 2011;160:283–93.
- [36] Ge X, Li Y, Yang X, Zhang H, Zhou P, Zhang Y, et al. Metagenomic analysis of viruses from bat fecal samples reveals many novel viruses in insectivorous bats in China. *J Virol* 2012;86:4620–30.
- [37] van Boheemen S, de Graaf M, Lauber C, Bestebroer TM, Raj VS, Zaki AM, Osterhaus ADME, et al. Genomic characterization of newly discovered coronavirus associated with acute respiratory distress syndrome in humans. *mBio* 2012;3.
- [38] Gholizadeh S, Djadid ND, Nouroozi B, Bekmohammadi M. Molecular phylogenetic analysis of *Anopheles* and *Cellia* subgenus *Anophelines* (Diptera: Culicidae) in temperate and tropical regions of Iran. *Acta Tropica* 2012 [epub ahead of print].
- [39] Guzmán C, Caballero L, Martín LM, Alvarez JB. Waxy genes from spelt wheat: new alleles for modern wheat breeding and new phylogenetic inferences about the origin of this species. *Ann Bot* 2012;110:1161–71.
- [40] Booth CM, Matukas LM, Tomlinson GA, Rachlis AR, Rose DB, Dwosh HA, et al. Clinical features and short-term outcomes of 144 patients with SARS in the greater Toronto area. *JAMA* 2003;289:2801–9.
- [41] Peiris JS. Severe acute respiratory syndrome (SARS). *J Clin Virol* 2003;28:245–7.
- [42] Peiris JS, Chu CM, Cheng VC, Chan KS, Hung IF, Poon LL, et al. Clinical progression and viral load in a community outbreak of coronavirus-associated SARS pneumonia: a prospective study. *Lancet* 2003;361:1767–72.
- [43] Peiris JS, Lai ST, Poon LL, Guan Y, Yam LY, Lim W, et al. Coronavirus as a possible cause of severe acute respiratory syndrome. *Lancet* 2003;361:1319–25.
- [44] Rickerts V, Wolf T, Rottmann C, Preiser W, Drosten C, Jakobi V, et al. [Clinical presentation and management of the severe acute respiratory syndrome (SARS)]. *Dtsch Med Wochenschr* 2003;128:1109–14.
- [45] World Health Organization. Summary table of SARS cases by country, 1 November 2002 - 7 August 2003. Geneva: WHO. Available at: http://www.who.int/csr/sars/country/country2003_08_15.pdf; 2003 [accessed 06.01.13].
- [46] Li SW, Yang TC, Wan L, Lin YJ, Tsai FJ, Lai CC, et al. Correlation between TGF-beta1 expression and proteomic profiling induced by severe acute respiratory syndrome coronavirus papain-like protease. *Proteomics* 2012;12: 3193–205.
- [47] Lai CC, Jou MJ, Huang SY, Li SW, Wan L, Tsai FJ, et al. Proteomic analysis of up-regulated proteins in human promonocyte cells expressing severe acute respiratory syndrome coronavirus 3C-like protease. *Proteomics* 2007;7: 1446–60.
- [48] Chan-Yeung M, Xu RH. SARS: epidemiology. *Respirology* 2003;8(Suppl):5.
- [49] Nicholls JM, Poon LL, Lee KC, Ng WF, Lai ST, Leung CY, et al. Lung pathology of fatal severe acute respiratory syndrome. *Lancet* 2003;361:1773–8.
- [50] He L, Ding Y, Zhang Q, Che X, He Y, Shen H, et al. Expression of elevated levels of pro-inflammatory cytokines in SARS-CoV-infected ACE2+ cells in SARS patients: relation to the acute lung injury and pathogenesis of SARS. *J Pathol* 2006; 210:288–97.
- [51] Yan H, Xiao G, Zhang J, Hu Y, Yuan F, Cole DK, et al. SARS coronavirus induces apoptosis in Vero E6 cells. *J Med Virol* 2004;73:323–31.
- [52] Wang WK, Chen SY, Liu JJ, Chen YC, Chen HL, Yang CF, et al. Detection of SARS-associated coronavirus in throat wash and saliva in early diagnosis. *Emerg Infect Dis* 2004;10:1213–9.
- [53] Huang KJ, Su IJ, Theron M, Wu YC, Lai SK, Liu CC, et al. An interferon-gamma-related cytokine storm in SARS patients. *J Med Virol* 2005;75:185–94.
- [54] Nicholls J, Dong XP, Jiang G, Peiris M. SARS: clinical virology and pathogenesis. *Respirology* 2003;8(Suppl):S6–8.
- [55] Lang Z, Zhang L, Zhang S, Meng X, Li J, Song C, et al. Pathological study on severe acute respiratory syndrome. *Chin Med J (Engl)* 2003;116:976–80.
- [56] Wang M, Xu HF, Mo ZY, Zheng BJ, Gu J, Qin PZ, et al. Anti-SARS virus antibody responses against human SARS-associated coronavirus and animal SARS-associated coronavirus-like virus. *Chin Med J (Engl)* 2004;117:1723–5.

- [57] Cameron MJ, Ran L, Xu L, Danesh A, Bermejo-Martin JF, Cameron CM, et al. Interferon-mediated immunopathological events are associated with atypical innate and adaptive immune responses in patients with severe acute respiratory syndrome. *J Virol* 2007;81:8692–706.
- [58] Ding Y, He L, Zhang Q, Huang Z, Che X, Hou J, et al. Organ distribution of severe acute respiratory syndrome (SARS) associated coronavirus (SARS-CoV) in SARS patients: implications for pathogenesis and virus transmission pathways. *J Pathol* 2004;203:622–30.
- [59] Drosten C, Günther S, Preiser W, van der Werf S, Brodt HR, Becker S, et al. Identification of a novel coronavirus in patients with severe acute respiratory syndrome. *N Engl J Med* 2003;348:1967–76.
- [60] Gu J, Gong E, Zhang B, Zheng J, Gao Z, Zhong Y, et al. Multiple organ infection and the pathogenesis of SARS. *J Exp Med* 2005;202:415–24.
- [61] Li T, Qiu Z, Zhang L, Han Y, He W, Liu Z, et al. Significant changes of peripheral T lymphocyte subsets in patients with severe acute respiratory syndrome. *J Infect Dis* 2004;189:648–51.
- [62] Perlman S, Dandekar AA. Immunopathogenesis of coronavirus infections: implications for SARS. *Nat Rev Immunol* 2005;5:917–27.
- [63] Pang C, Gu DL. [Some problems about detecting the suspected cases of SARS according to the local skin temperatures on face]. *Space Med Med Eng (Beijing)* 2003;16:231–4.
- [64] Baas T, Taubenberger JK, Chong PY, Chui P, Katze MG. SARS-CoV virus-host interactions and comparative etiologies of acute respiratory distress syndrome as determined by transcriptional and cytokine profiling of formalin-fixed paraffin-embedded tissues. *J Interferon Cytokine Res* 2006;26:309–17.
- [65] Wang WK, Chen SY, Liu JJ, Kao CL, Chen HL, Chiang BL, et al. Temporal relationship of viral load, ribavirin, interleukin (IL)-6, IL-8, and clinical progression in patients with severe acute respiratory syndrome. *Clin Infect Dis* 2004;39:1071–5.
- [66] Morgenstern B, Michaelis M, Baer PC, Doerr HW, Cinatl Jr J. Ribavirin and interferon-beta synergistically inhibit SARS-associated coronavirus replication in animal and human cell lines. *Biochem Biophys Res Commun* 2005;326:905–8.
- [67] Hogan RJ, Gao G, Rowe T, Bell P, Flieder D, Paragas J, et al. Resolution of primary severe acute respiratory syndrome-associated coronavirus infection requires Stat1. *J Virol* 2004;78:11416–21.
- [68] Sheahan T, Morrison TE, Funkhouser W, Uematsu S, Akira S, Baric RS, et al. MyD88 is required for protection from lethal infection with a mouse-adapted SARS-CoV. *PLoS Pathog* 2008;4:e1000240.
- [69] Cheung CY, Poon LL, Ng IH, Luk W, Sia SF, Wu MH, et al. Cytokine responses in severe acute respiratory syndrome coronavirus-infected macrophages in vitro: possible relevance to pathogenesis. *J Virol* 2005;79:7819–26.
- [70] Lin X, Chen Y, Meng A, Feng X. Termination of TGF-beta superfamily signaling through SMAD dephosphorylation—a functional genomic view. *J Genet Genomics* 2007;34:1–9.
- [71] Ray S, Broor SL, Vaishnav Y, Sarkar C, Girish R, Dar L, et al. Transforming growth factor beta in hepatitis C virus infection: in vivo and in vitro findings. *J Gastroenterol Hepatol* 2003;18:393–403.
- [72] Zhao X, Nicholls JM, Chen YG. Severe acute respiratory syndrome-associated coronavirus nucleocapsid protein interacts with Smad3 and modulates transforming growth factor-beta signaling. *J Biol Chem* 2008;283:3272–80.
- [73] Lu X, Pan J, Tao J, Guo D. SARS-CoV nucleocapsid protein antagonizes IFN-beta response by targeting initial step of IFN-beta induction pathway, and its C-terminal region is critical for the antagonism. *Virus Genes* 2011;42:37–45.
- [74] Chang YJ, Liu CY, Chiang BL, Chao YC, Chen CC. Induction of IL-8 release in lung cells via activator protein-1 by recombinant baculovirus displaying severe acute respiratory syndrome-coronavirus spike proteins: identification of two functional regions. *J Immunol* 2004;173:7602–14.
- [75] Dosch SF, Mahajan SD, Collins AR. SARS coronavirus spike protein-induced innate immune response occurs via activation of the NF-kappaB pathway in human monocyte macrophages in vitro. *Virus Res* 2009;142:19–27.
- [76] Law AH, Lee DC, Cheung BK, Yim HC, Lau AS. Role for nonstructural protein 1 of severe acute respiratory syndrome coronavirus in chemokine dysregulation. *J Virol* 2007;81:416–22.
- [77] Lu B, Tao L, Wang T, Zheng Z, Li B, Chen Z, et al. Humoral and cellular immune responses induced by 3a DNA vaccines against severe acute respiratory syndrome (SARS) or SARS-like coronavirus in mice. *Clin Vaccine Immunol* 2009;16:73–7.
- [78] Sulea T, Lindner HA, Purisima EO, Menard R. Deubiquitination, a new function of the severe acute respiratory syndrome coronavirus papain-like protease? *J Virol* 2005;79:4550–1.
- [79] Spiegel M, Pichlmair A, Martínez-Sobrido L, Cros J, García-Sastre A, Haller O, et al. Inhibition of Beta interferon induction by severe acute respiratory syndrome coronavirus suggests a two-step model for activation of interferon regulatory factor 3. *J Virol* 2005;79:2079–86.
- [80] Frieman M, Ratia K, Johnston RE, Mesecar AD, Baric RS. Severe acute respiratory syndrome coronavirus papain-like protease ubiquitin-like domain and catalytic domain regulate antagonism of IRF3 and NF-kappaB signaling. *J Virol* 2009;83:6689–705.
- [81] Devaraj SG, Wang N, Chen Z, Chen Z, Tseng M, Barretto N, et al. Regulation of IRF-3-dependent innate immunity by the papain-like protease domain of the severe acute respiratory syndrome coronavirus. *J Biol Chem* 2007;282:32208–21.
- [82] Li SW, Lai CC, Ping JF, Tsai FJ, Wan L, Lin YJ, et al. Severe acute respiratory syndrome coronavirus papain-like protease suppressed alpha interferon-induced responses through downregulation of extracellular signal-regulated kinase 1-mediated signalling pathways. *J Gen Virol* 2011;92:1127–40.
- [83] Hofmann H, Pyrc K, van der Hoek L, Geier M, Berkhout B, Pöhlmann S. Human coronavirus NL63 employs the severe acute respiratory syndrome coronavirus receptor for cellular entry. *Proc Natl Acad Sci U S A* 2005;102:7988–93.
- [84] Han TH, Chung JY, Kim SW, Hwang ES. Human Coronavirus-NL63 infections in Korean children, 2004–2006. *J Clin Virol* 2007;38:27–31.
- [85] Vabret A, Mourez T, Dina J, van der Hoek L, Gouarin S, Petitjean J, et al. Human coronavirus NL63, France. *Emerg Infect Dis* 2005;11:1225–9.
- [86] Woo PC, Huang Y, Lau SK, Tsoi HW, Yuen KY. In silico analysis of ORF1ab in coronavirus HKU1 genome reveals a unique putative cleavage site of coronavirus HKU1 3C-like protease. *Microbiol Immunol* 2005;49:899–908.
- [87] Pyrc K, Berkhout B, van der Hoek L. The novel human coronaviruses NL63 and HKU1. *J Virol* 2007;81:3051–7.
- [88] Lau SK, Woo PC, Yip CC, Tse H, Tsoi HW, Cheng VC, et al. Coronavirus HKU1 and other coronavirus infections in Hong Kong. *J Clin Microbiol* 2006;44:2063–71.
- [89] Health Protection Agency, Novel Coronavirus 2012, HPA, 2012. Available at: <http://www.hpa.org.uk/Topics/InfectiousDiseases/InfectionsAZ/NovelCoronavirus2012/> [accessed 06.01.13].
- [90] World Health Organization. World Health Organization: global alert and response (GAR) - Novel coronavirus infection 2012. WHO. Available at: http://www.who.int/csr/don/2012_09_25/en/index.html; 2012 [accessed 06.01.13].

- [91] Danielsson N, ECDC Internal Response Team Catchpole M. Novel coronavirus associated with severe respiratory disease: case definition and public health measures. *Euro Surveill* 2012;17(39).
- [92] World Health Organization. Background and summary of novel coronavirus infection – as of 21 December 2012. WHO. Available at: http://www.who.int/csr/disease/coronavirus_infections/update_20121221/en/index.html; 2012 [accessed 06.01.13].
- [93] Corman VM, Eckerle I, Bleicker T, Zaki A, Landt O, Eschbach-Bludau M, et al. Detection of a novel human coronavirus by real-time reverse-transcription polymerase chain reaction. *Euro Surveill* 2012;17(39).
- [94] Balboni A, Gallina L, Palladini A, Prosperi S, Battilani M. A real-time PCR assay for bat SARS-like coronavirus detection and its application to Italian greater horseshoe bat faecal sample surveys. *ScientificWorldJournal* 2012;2012:989514.
- [95] Moës E, Vijgen L, Keyaerts E, Zlateva K, Li S, Maes P, et al. A novel pancoronavirus RT-PCR assay: frequent detection of human coronavirus NL63 in children hospitalized with respiratory tract infections in Belgium. *BMC Infect Dis* 2005;5:6.