

REVIEW

Open Access

Human interphase chromosomes: a review of available molecular cytogenetic technologies

Svetlana G Vorsanova^{1,2}, Yuri B Yurov^{1,2}, Ivan Y Iourov^{1,2*}

Abstract

Human karyotype is usually studied by classical cytogenetic (banding) techniques. To perform it, one has to obtain metaphase chromosomes of mitotic cells. This leads to the impossibility of analyzing all the cell types, to moderate cell scoring, and to the extrapolation of cytogenetic data retrieved from a couple of tens of mitotic cells to the whole organism, suggesting that all the remaining cells possess these genomes. However, this is far from being the case inasmuch as chromosome abnormalities can occur in any cell along ontogeny. Since somatic cells of eukaryotes are more likely to be in interphase, the solution of the problem concerning studying postmitotic cells and larger cell populations is interphase cytogenetics, which has become more or less applicable for specific biomedical tasks due to achievements in molecular cytogenetics (i.e. developments of fluorescence *in situ* hybridization – FISH, and multicolor banding – MCB). Numerous interphase molecular cytogenetic approaches are restricted to studying specific genomic loci (regions) being, however, useful for identification of chromosome abnormalities (aneuploidy, polyploidy, deletions, inversions, duplications, translocations). Moreover, these techniques are the unique possibility to establish biological role and patterns of nuclear genome organization at suprachromosomal level in a given cell. Here, it is to note that this issue is incompletely worked out due to technical limitations. Nonetheless, a number of state-of-the-art molecular cytogenetic techniques (i.e. multicolor interphase FISH or interphase chromosome-specific MCB) allow visualization of interphase chromosomes in their integrity at molecular resolutions. Thus, regardless numerous difficulties encountered during studying human interphase chromosomes, molecular cytogenetics does provide for high-resolution single-cell analysis of genome organization, structure and behavior at all stages of cell cycle.

Introduction

Currently, it is estimated that no fewer than 1 million cytogenetic and molecular cytogenetic analyses are performed per year representing the standard of care in several fields of medicine and the routine clinical work-up for numerous patients suffering from congenital malformations, mental diseases, cancers, or reproductive problems [1]. Molecular cytogenetic techniques have been repeatedly proven effective in diagnostics and have been recognized as a valuable addition or even alternative to chromosomal banding [2-4]. Furthermore, contemporary basic biomedical research widely applies molecular cytogenetic technologies [5-7]. Browsing the most popular scientific resources would undoubtedly return several tens of thousands of articles, which mention at least one molecular cytogenetic technique (for

more details refer to [3] and web page about multicolor fluorescence *in situ* hybridization at <http://www.med.uni-jena.de/fish/mFISH/mFISHlit.htm> managed by Dr. Thomas Liehr, Jena, Germany). Thus, one can be certain that it is hard to overestimate the role of molecular cytogenetics in current biomedicine.

There are two main advantages that molecular cytogenetics possesses: (i) the ability to provide either an on-chip scan of the whole genome at extremely high resolution or visualization of single peculiar genomic loci [4,6,8]; (ii) the capability to analyze genome organization, structure and behavior in single cells at the DNA (RNA) sequence level [7,9,10]. Both are continuously used for biomedical research and molecular diagnosis of chromosome abnormalities in humans [2-13]. The first advantage is appreciable when analyzing mixed DNA isolated from large amount of cells. Therefore, it is unsurprising that such approaches are rarely used for single-cell analysis [10,14]. The second advantage of

* Correspondence: ivan_iourov@yahoo.com

¹Institute of Pediatrics and Children Surgery, Rosmedtechnologii, Moscow, 127412, Russia

molecular cytogenetic techniques is consistently emphasized [3,5-13], but is used more commonly for studying mitotic cells *via* analyzing metaphase chromosomes [3,7,10,12]. However, cells of eukaryotes are more likely to be in interphase. Therefore, during surveys of genome organization, structure and behavior, essential part of cellular life is usually fallen out of researchers' scope. As to molecular diagnosis of chromosome abnormalities, one can notice that interphase analysis is uncommonly applied, as well. The explanation of leaving interphase cytogenetics aside from diagnostics and research might be a suggested lack of reproducibility and low resolution. A brief look through studies of genome architecture in interphase nuclei [15-19] and somatic genomic variations [7,10,12,20-29] as well as developments in interphase cytogenetics [30-35] will reveal such assumptions unsupported and will show that laboratories elaborating such techniques are able to solve different practical and research tasks without major difficulties [3,7,12-35]. It seems thereby that preferences to use interphase molecular cytogenetic techniques suffer rather from "insufficient publicity" than from "technological underdevelopment".

Looking through the voluminous amount of reviews dedicated to molecular cytogenetics, we have found occasional descriptions of both technological and theoretical side of visualizing human chromosomes in interphase. Consequently, we were forced to conclude that undeservedly little attention is paid to interphase molecular cytogenetics in modern biomedical literature. Additionally, technical side of the application is even more rarely addressed. To fill this gap, we have attempted to give an overview of currently applied molecular cytogenetic techniques with a special emphasis on their technological abilities for studying human interphase chromosomes.

Molecular cytogenetic techniques, their resolution and potential for single-cell analysis of interphase chromosomes

The overwhelming majority of molecular cytogenetic techniques are based on hybridization. There are currently two essential platforms available for developments in molecular cytogenetics: fluorescence in situ hybridization (FISH) including comparative genomic hybridization (CGH) [3,36] and peptide nucleic acid (PNA) probing for analysis of chromosomal DNA [37,38]. Alternatively, another technique uses primed in situ labelling (PRINS) reaction [37,38]. The resolution and level of excellence of all these techniques are established against cytogenetic banding analysis, which remains the golden standard in this instance [36]. Single-cell molecular cytogenetic analysis can be performed either through analysis of metaphase plates or through analysis of

interphase nuclei. Studying metaphase plates has been long described to be successful using several detection technologies (i.e. spectral karyotyping – SKY or multicolor FISH – MFISH) and different DNA probe sets (chromosome-enumeration/centromeric, site-specific, whole-painting, microdissected) [2,3,5-7,9-13,30,36,39-46]. In general, if modified, almost all these techniques can be applied to interphase cells, but this "transfer of technology" requires significant efforts [2,3,7,10,12,13,30-33,35,47]. Generally, all molecular cytogenetic assays that provide for visualization of genomic loci in an interphase nucleus are termed interphase FISH or I-FISH [35]. Table 1 gives an overview of molecular cytogenetic techniques that are used for metaphase and interphase analysis with special attention to the resolution and to the modifications for studying single cells. The impossibility of listing all known molecular cytogenetic approaches seems to be apparent, but even a short description of such techniques (Table 1) shows molecular cytogenetics able to perform high-resolution analysis of chromosomal structure and behavior at all stages of cell cycle, being, nevertheless, more frequently used to detect metaphase chromosome imbalances and rearrangements or to operate with total DNA for probing in CGH analysis [2-7,10-14,19-54]. Further, we attempt to review each aforementioned approach in context of applications to single-cell chromosomal analysis.

FISH

FISH offers numerous possibilities to study either the whole genome or specific genomic loci (regions) [2-7,10-13,36,39-41]. The probes mainly determine the resolution of molecular cytogenetic techniques [3]. Regardless molecular peculiarities and pattern of sequence modifications (i.e. LNA (locked-nucleic acid) or PNA probes, for more details see [3,13,37,38]), probes for molecular cytogenetic assays can be classified according to the pattern of detected DNA sequences. Such classification includes repetitive-sequence DNA (centromeric and telomeric), site-specific, whole chromosome painting (wcp) probes [3,55].

FISH, which paints repetitive genomic sequences, can be performed with either centromeric (chromosome enumeration or chromosome-specific) or telomeric DNA probes. Analysis of telomeres is an important area of biomedical research [56]. Usually, DNA or PNA probes possessing TTAGGG repetitive sequence motifs are used [3,56]. These assays are needed to cover large area of cancer and aging research (telomere biology), but seem to be poorly applicable for diagnosis [3]. I-FISH analysis using telomeric probes was only described in few nuclear organization studies [57]. Contrariwise, applications of I-FISH with centromeric DNA probes are an integral part of diagnostics in medical genetics, oncology and reproductive medicine [1-3,5,7,10,12,13,20-30,35-38,41,42,44,46,55,

Table 1 Molecular cytogenetic techniques, their resolution and validity for single-cell analysis of interphase/metaphase chromosomes (for more details see text)

Approach	Resolution	MA*	IA**	SCA [^]	PVC ^{^^}	Refs
Cytogenetic banding analysis ("golden standard")	5-7 Mb	+	-	+	+	[1]
FISH/MFISH/SKY						
FISH/MFISH/SKY with centromeric probes	>0.3-1 Mb	+/-	+	+	+/-	[20-26,30,32,35,41,42,44,46]
FISH/MFISH/SKY with site-specific probes	~0.1-2 Mb	+/-	+/-	+/-	+/-	[45,47-50]
FISH/MFISH/SKY with whole-painting probes	>5-10 Mb	+	-	+	+	[2,3,5,6,13,36,39,40]
MCB						
Metaphase MCB	~2-5 Mb	+	-	+	+	[2,13,43,45]
ICS-MCB	~2-5 Mb	-	+	+	+	[23,24,26,28,29,31-35]
Fiber FISH	>2.3 (2-3) kb	na	na	+	+	[51,52]
Single-cell CGH						
Standard CGH	2-5 Mb	na	na	+	-	[53]
Array CGH	0.03-1 Mb	na	na	+	-	[14,54]

* - analysis of metaphase chromosomes (MA - metaphase analysis); ** - analysis of interphase chromosomes (IA - interphase analysis); [^] - possibility to perform single cell analysis (SCA); ^{^^} - possibility to visualize chromosomes or chromosomal loci (PVC - possibility to visualize chromosomal loci); na - not applicable;

58-61]. Moreover, application of these probes has been long demonstrated to be extremely valuable for research in fields of chromosome biology studying genome organization (chromosomal and nuclear), evolution, behavior and variation in health and disease [2,3,7,10,12,13,20-30,35,41,42,44,55,62-67]. The popularity of these DNA probes is usually attributed to near 100% hybridization efficiency because of painting highly repetitive DNAs as well as to chromosome-specificity of centromeric human DNAs allowing analysis of individual homologous chromosome pairs in interphase [7,10,30,35]. Moreover, due to the extreme interindividual variations of pericentromeric heterochromatic DNA, such assays allow application of quantitative FISH (QFISH) that can be useful for solving numerous problems encountered during metaphase and interphase analysis of chromosomes [32,35,59]. The potential of related assays is poorly determined by its genomic resolution (Table 1), inasmuch as applications of centromeric DNA probes suggest the analysis of phenomena encompassing significantly larger genomic loci as to visualized ones [3,7,10]. As to interphase cytogenetics, I-FISH with chromosome-enumeration probes makes possible to detect numerical chromosome imbalances (aneuploidy and polyploidy) in vast cell populations [7,10,12,20-30,35,41,42,60]. In a limited amount of cases, similar approaches are applicable for metaphase cytogenetic analysis of chromosome abnormalities [58,63,64,68]. Numbers of signals for these probes are supposed to be identical to numbers of homologous chromosomes per interphase nucleus [3,7,10,20-30,33,35,41,42,46]. However, this is not always the case [7,10,23-33,35]. This is the main disadvantage of I-FISH with centromeric DNA probes, which is, however, successfully solved by means of FISH with site-specific DNA probes (locus-specific BAC probes or BAC-probe contigs) [13].

FISH using site-specific DNA probes (YACs, BACs, PACs, cosmids) is usually used to map chromosomal regions, within which a breakpoint is located [3,5,13,61]. Additionally, these probes can be used for diagnosing known microdeletion and microduplication syndromes [1,3,13,27], aneuploidy and recurrent chromosome abnormalities during preimplantation genetic diagnosis [48-50], prenatal diagnosis [3,13,47], oncocyto-genetic analysis [1-3,5,13,36,27,50], and precision of copy number variations [8]. Being valuable approach for studying genomic loci smaller than 1 Mb, I-FISH with site-specific probes is frequently used for studying nuclear organization of genes and its impact on the transcriptional activity [16-18,69]. Nevertheless, relatively moderate hybridization efficiency (<70%) hinders the application of such approaches in numerous areas of biomedical research and diagnosis [7,10]. The latter does not concern a number of diagnostic FISH procedures applying these types of probes (for instance, in cases of routine oncohematological and tumor diagnostics). For diagnostic issues, such approaches has cut-offs between 92 and 98% [13].

FISH using wcp is a basis for MFISH (24-color FISH) and SKY [2,13,39,40]. These methods are valuable for cancer cytogenetics and, in some cases, for diagnosis of constitutional chromosome abnormalities [2,5,6,13,36]. For analysis of interphase chromosomes, MFISH/SKY is hardly applicable. Nevertheless, a study has visualized simultaneously all chromosomes in interphase nuclei of fibroblasts and prometaphase rosettes by 24-color MFISH [70]. Afterwards, such approaches have not been ever considered for related analysis. Two-to-five-color assays with wcp probes have been repeatedly used for molecular cytogenetic diagnosis of structural alterations to metaphase chromosomes [1-3,5-7,13,36,61] and

investigation of genome organization in interphase nuclei [15,57,66,70-72]. I-FISH with wcp probes is too problematic to be competitive with other techniques of interphase molecular cytogenetic diagnosis [7,10,33].

By microdissection of chromosomal loci for obtaining a set of probes that produce multicolor pseudo-G-banding, a high-resolution molecular cytogenetic technique for analysis of metaphase chromosomes termed MCB (multicolor banding) was proposed [43]. The latter has been consistently shown to be applicable for the identification of structural chromosome abnormalities and genome organization [2,13,36,43,45,61,73]. A modification of this technique, called recently interphase chromosome-specific MCB (ICS-MCB) that generates MCB of a homologous chromosome pair on single nuclei, has been demonstrated effective for studying human interphase chromosome organization and variations (somatic genomic variations and chromosome instability in health and disease) [3,7,10,13,19,23,24,26-29,31,33-35,74,75]. Apart from impossibility to analyze simultaneously several homologous chromosome pairs and relative complexity of the analysis, ICS-MCB does not possess major limitations. Moreover, this is the unique way to obtain a view on the entire interphase chromosome in its integrity [23,33,35].

The highest molecular cytogenetic resolution is achieved by fiber FISH (~2.3 kb) [52,76]. This approach was originally designed for mapping cloned DNA fragments at high resolution. The latter was found useful for investigation of genomic organization (on metaphase chromosomes), stalled transcription and genomic rearrangements (including large deletions within gene sequences) [51,52,76]. Although this technique is based on obtaining DNA fibers from interphase nuclei, it cannot be attributed to I-FISH. Single-cell molecular cytogenetic analysis by fiber FISH (especially, analysis of large cell populations) is highly complicated.

CGH

Since CGH compares quantitative differences between individual genomes, its applications are restricted to analysis of losses/gains of chromosomal (genomic) loci without direct visualization of chromosomes [4,77]. Array CGH can provide for a resolution up to nucleotide level, but still is poorly applicable for studying chromosomes of a cell. Nevertheless, several reports have demonstrated either standard CGH or array CGH on microdissected interphase nuclei to detect chromosome aberrations in single cells of preimplantation embryos [14,53,54]. Such approaches are applicable for unbalanced genomic rearrangements being useless for other areas of chromosome biology, which requires visualization of chromosomal DNA [10]. The potential of CGH-based single-cell analysis for molecular diagnosis and for

surveys of somatic genomic variations remains to be estimated.

PNA and PRINS

Both PNA and PRINS can be successfully applied for studying human chromosomes [3,7,10,13,37,38]. PNAs are suggested to have several advantages over conventional molecular cytogenetic DNA probes, which are the result of their smaller size [38]. Notwithstanding, poor availability does not allow researchers to evaluate *in situ* hybridization with PNA probes for either metaphase or interphase molecular cytogenetics. Moreover, these probes are usually restricted to studying centromeric and telomeric repetitive chromosomal DNA.

In contrast to FISH and CGH, PRINS is based on another biochemical process (polymerase reaction) [37]. This makes it useful for case-control studies of newly discovered phenomena to exclude hypothetical errors that might be produced by hybridization [24]. Usually, PRINS shows almost the same results as FISH. Therefore, there is no apparent interest to substitute FISH-based techniques by PRINS, especially taking into account its essential limitation: available probes are oligonucleotides for pericentromeric/heterochromatic and few euchromatic regions (poorly reproducible!) [3,7,24,37].

The key process of all the studies aimed to analyze interphase chromosomes is visualization. In other words, lacking of direct (microscopic) DNA visualization makes all such researches incomplete. This becomes even more evident for studying chromosome organization in single cells. As one can see, only FISH-based techniques offer possibilities to detect either whole chromosomes or specific genomic loci of extremely small size in single cells. Therefore, to perform a valid study of human interphase chromosomes, I-FISH protocols are to use. The next part of our review addresses areas of I-FISH applications as well as its advantages and limitations.

I-FISH: advantages and limitations

I-FISH as all other FISH-based methods roughly requires three steps to be performed: (i) obtaining cells suspensions or performing another preparations of biopsies for the analysis; (ii) denaturation/hybridization; (iii) microscopic visual/digital analysis of hybridization results [13,35,78]. The first stage is not associated with any limitation of I-FISH, because any cell type of a human organism can be processed for such analyses [7,35,78]. This is considered the essential advantage of interphase molecular cytogenetic techniques in contrast to classical cytogenetics (analysis of metaphase chromosomes) – the ability to analyze chromosomes in all the tissue (cell) types [3,7,13,20-36]. Classically, I-FISH was suggested to be limited to analyses of specific genomic

loci [2,3,7,13]. However, some modifications such as ICS-MCB allow to get a view of interphase chromosomes in their integrity [23,24,26-29,31,33-35,74]. As mentioned before, ICS-MCB still have a limitation that is referred to the possibility of studying only one homologous chromosome pair per analysis (metaphase chromosomal analysis allows to visualize all chromosomes of a cell), being, however, the unique way to visualize the whole banded chromosome in a nucleus [23,33]. Denaturation and hybridization steps of I-FISH are performed identically to metaphase FISH-based approaches [13,35]. Therefore, no additional drawbacks can be attributed to these procedures during interphase molecular cytogenetic studies. Scoring of I-FISH results is usually performed *via* conventional visual analysis [35,79]. However, there are numerous possibilities to apply digital analysis for studying interphase chromosomes. These include, but are not restricted to, QFISH, analysis of signal co-localization (oncocytogetic studies of gene fusions because of translocations in interphase nuclei), ICS-MCB (visualization of chromosomal structures), increasing of FISH result visibility, automatic signal detection [79]. Furthermore, digital analysis is a need for multicolor FISH-based assays (SKY, MFISH, multiprobe interphase FISH or mFISH), which are usually applied to increase the potential of FISH applications through simultaneous analysis of multiple targets [2,3,12,13,20-30,35,36,80]. Combining several aforementioned techniques mFISH with 2-5 probes (colors) per assay, QFISH and ICS-MCB has become a basis for an integrated approach proven to be of highest efficiency for molecular diagnosis and genome/chromosome researches at supramolecular level in interphase [7,10,12,13,20-30,32,33,35,41,60,78,81]. The type of FISH result evaluation (i.e. visual or digital) is determined by the type of assay or, more precisely, by features of DNA probes (amount of probes per reaction and DNA sequence affinity) and detection. Therefore, to get an overview it is to subdivide I-FISH techniques this way. Table 2 gives such overview.

I-FISH with centromeric probes

I-FISH with centromeric probes is highly applicable for different areas of biomedical research and diagnosis [7,10,13,20-22,30,35,41,58-60,81,82]. The most frequent application of the method is the identification of numerical chromosome abnormalities (aneuploidy and polyploidy) in interphase nuclei (Figure 1). The latter is required for pre-/postnatal diagnosis, cancer diagnosis/prognosis, somatic genomic variation surveys [7,10,20-22,30,35,82]. As one can see from table 2, near 100% hybridization efficiency of centromeric DNA probes [7,10,30,35] and chromosome-specific DNA sequences forming pericentromeric/heterochromatic

chromosomal regions (apart from shared alphoid DNA of chromosomes 5 and 19, 13 and 21, 14 and 22) [30,35,41,42,44,83] are the essential source of advantages that this technique possesses. Heteromorphisms of pericentromeric DNAs can produce the lack of a signal leading, thereby, to impossibility of the I-FISH assay application. Fortunately, such extreme heteromorphisms (centromeric DNA variations) are rare in the general population [32,35,59,84-86].

I-FISH with site-specific probes

Interphase molecular cytogenetic studies by I-FISH with site-specific probes are commonly applied in preimplantation, prenatal and postnatal diagnosis as well as in cancer cytogenetics (Figure 2) [2,3,13,36,47-50]. Although repeatedly noted to be of significant importance for detecting gene fusions resulting from interchromosomal translocations (cancer biomarkers) [49,87-89] and to be useful for preimplantation diagnosis [48-50], such I-FISH modifications has considerable disadvantages. Firstly, hybridization efficiency of site-specific probes is usually between 40 and 70% [7,10]. This has the potential to produce false-positive or false-negative data [7,28]. Additionally, it requires to use probes for "well-characterized" genomic DNA sequences (i.e. mapped oncogenes, genes/genomic loci within microdeletion or microduplication regions) [3]. Therefore, it is not surprising that there are only few approaches using these DNA probes that are performed to detect well-known chromosomal rearrangements in cancer cells [87-89] and, more rarely, deletions/duplications in clinical populations [1,3,8,50,90-92]. However, FISH using site-specific probes is almost the unique way to visualize DNA sequences smaller than 1 Mb in interphase nuclei. Simultaneous use of centromeric and site-specific probes in an mFISH assay (Figure 3) is sometimes useful for diagnostics and survey of intercellular (somatic) genomic variations [7,20,28,46,48].

I-FISH with wcp

It is generally recognized that FISH chromosomal painting using wcp is completely useless for identification of number and structure of interphase chromosomes (Figure 4) [3,7,10,13,33,35,80]. However, basic research of chromosome architecture in interphase is usually performed using I-FISH with wcp. These probes allows to visualize chromosome territories and their positioning relative to nuclear compartments (Figure 4B) [57,70-72,85,93]. For the last two decades, I-FISH-wcp approaches were almost the unique way to study genomic organization in interphase [72]. Some studies proposed to use the complete wcp set in an interphase MFISH reaction [70,93]. Nonetheless, these techniques are all limited in their abilities to paint chromosome territories (volumes) only (Table 2) [33].

Table 2 Overview of I-FISH techniques

Technique	Brief description	Advantages	Limitations	Refs
I-FISH with centromeric probes	I-FISH on interphase nuclei painting pericentromeric (heterochromatic) regions	High hybridization efficiency, chromosome specificity (apart few chromosomes)	Signal associations, impossible to analyze chromosomes 5,13, 14, 19, 21, 22; heteromorphisms	[7,10,20-22,30,35,41,58-60,81,82]
I-FISH with site-specific probes	I-FISH painting specific euchromatic genomic loci	Small specific genomic loci are visualized	Low hybridization efficiency, numerous artifacts	[8,13,28,42,47-50,69]
I-FISH with wcp	I-FISH painting chromosome territories	Identification of nuclear chromosome territories	Chromosome territories are ambiguous, no additional information	[57,70-72,93]
mFISH	Multicolor I-FISH with >2 probes labeled by different fluorochromes/ligands	Analysis of several targeted genomic loci	Difficulty to distinguish between artifacts and aneuploidy/ployploidy	[7,10,20-22,30,35]
mFISH/QFISH	mFISH + QFISH digitalization of FISH signals	Distinguishes between FISH artifacts and aneuploidy (ployploidy)	Same as mFISH	[7,10,24-29,32,35]
MFISH	Simultaneous visualization of the complete set of chromosomes in an interphase nucleus	All chromosome territories are simultaneously seen	Exceedingly sophisticated analysis; data poorly interpretable	[70,93]
ICS-MCB	Chromosome-specific MCB generated on interphase nuclei	Visualization of whole banded interphase chromosomes in their integrity	A pair of homologous chromosomes is studied per assay; relative complexity of the analysis	[7,10,13,19,23,24,26-29,31,33-35,74,75]

ICS-MCB

To visualize a homologous pair of interphase chromosomes in their integrity, one has to generate MCB. Interphase banded chromosomes appear as metaphase ones when ICS-MCB is applied. Therefore, this I-FISH approach solves the long-standing limitation of cytogenetics that refers to obtaining metaphase chromosomes [23,31-35]. Figure 5 gives an example of aneuploidy detection in an interphase nucleus isolated from the Alzheimer's disease brain [28]. ICS-MCB can be widely applied for basic research of somatic genomic variations, chromosome structural and functional organization in interphase, supramolecular disease mechanisms [3,7,10,12,13,19,23,24,26-29,31,33-36,73-75,79-81]. Apparently, the sole disadvantage of this technique is the impossibility to analyze more than one homologous chromosome pair at once [23,33].

There are several general problems that surround I-FISH application. As we have already mentioned, differences of hybridization efficiency complicate simultaneous applications of different probe sets [7]. For instance, signals of site-specific probes can be missed because of high brightness of wcp or centromeric probe signals. Here, the most apparent solution is ICS-MCB application [33,35]. However, some interphase protocols, mostly associated with molecular oncocyto-genetics, are proven to be valid for diagnostic purposes [1,13,36,87,88]. DNA replication during S phase of cell

cycle is another major problem of I-FISH applications [7,47]. Despite of recommendations concerning this type of I-FISH artifacts in the available literature, FISH analysis can be hindered by replicative signal appearance. This is mainly related to site-specific DNA probes, being, however, noticed during I-FISH with centromeric probes, as well [7,10,22,35,47] (Figure 6A-C). Additional source of numerous artifacts that can be considered as false-positive chromosome abnormalities in interphase is nuclear organization. In this context, the most problematic pattern of chromosome arrangement in the nucleus is related to chromosomal loci associations [94,95]. This significantly affects I-FISH results becoming even more important taking into account that numerous cell types are prone to exhibit intranuclear associations/pairing of genomic loci (Figure 6D) [20,32,35,95]. Regardless frequent occurrence of related difficulties, the problem is easily solved by QFISH (Figure 6E) [23,24,28,32,35,95].

Finishing the list of interphase FISH-based techniques, it is to mention Immuno-FISH. This method combines immunohistochemical detection of proteins and FISH for visualization of DNA (RNA) targets [96-98]. Immuno-FISH is found applicable in cancer research/diagnosis (simultaneous immunophenotyping and single-cell genetic analysis), studies of chromosome structure and organization, transplantation research, and identification of supramolecular disease mechanisms




Figure 1 Two- and three-color I-FISH with centromeric DNA probes. (A) normal diploid nucleus with two signals for chromosome 1 and chromosome 15; (B) monosomic nucleus with two signals for chromosome 1 and one signal for chromosome 15; (C) trisomic nucleus with two signals for chromosome 1 and three signals for chromosome 15; (D) normal diploid nucleus with two signals for chromosome 1, chromosome 9 and chromosome 16; (E) monosomic nucleus with two signals for chromosome 1 and chromosome 9 and one signal for chromosome 16; (F) trisomic nucleus with two signals for chromosome 1 and chromosome 16 and three signals for chromosome 9; (G) triploid nucleus with three signals for chromosome 16 and chromosome 18; (H) tetraploid nucleus with two signals for chromosome X and chromosome Y; (I) tetraploid nucleus with two signals for chromosome X and chromosome Y, and four signals for chromosome 1.

[28,29,96-100]. Figure 7 demonstrates Immuno-FISH used for studying interphase chromosomes in neuronal cells of the adult human brain [28,29].

After listing the most known methods of interphase molecular cytogenetics, it is to focus on their specific applications. Currently, there are three main biomedical areas requiring the use of I-FISH: analysis of intranuclear chromosome (genome) organization; identification of somatic (intercellular and intertissular) genomic

variations; molecular cytogenetic diagnosis. Below, a brief description of these applications is given.

Genome organization in interphase

Spatial chromosome organization in interphase has been repeatedly shown to be a driving force for numerous crucial intracellular processes. It is suggested that specific arrangement of interphase chromosomes is likely to associate with genome activity, normal/abnormal cell




Figure 2 I-FISH with site-specific DNA probes. (A) normal diploid nucleus with two signals for chromosome 21; (B) trisomic nucleus with three signals for chromosome 21; (C) interphase nucleus exhibiting co-localization of *ABL* and *BCR* genes probably due to t(9;22)/Philadelphia chromosome.

division, chromosome rearrangements occurring during meiosis and mitosis [7,15-17,69,19,70-72,75,93,100,101]. To get an integrated view of genome organization in interphase, numerous approaches should be applied. The leading role in these studies is played by I-FISH [7,80,72,93]. There could be several applications of I-FISH approaches for interphase chromosome analysis on this occasion: (i) identification of chromosome

positioning and its relation to other nuclear compartments (nucleolus, Cajal bodies, nuclear speckles etc.) – I-FISH with wcp, interphase MFISH or ICS-MCB [19,31,23,33,35,34,70-72,74,93]; (ii) studying correlation between positioning of specific genomic loci in relation to each other (i.e. association of whole chromosomes or their regions) and their behavior (transcriptional/replicative activity) for elucidating functional meaning of




Figure 3 Five-color I-FISH (mFISH) with DNA probes for chromosomes 18, X and Y (centromeric probes) as well as 13 and 21 (site-specific probes). a presumably normal (diploid) male nucleus isolated from the adult human brain.




Figure 4 I-FISH with two wcp for chromosomes 7 and 21. **(A)** ambiguous chromosome territories provide information neither about number of chromosomes nor about structure of chromosomes (chromosome 7 – green signal; chromosome 21 – red signals), whereas this individual presented with regular unbalanced t(7;21); more details are given in Vorsanova et al. 2008 [64]; **(B)** chromosome territories in an interphase nucleus of a cell isolated from the ataxia-telangiectasia brain (chromosome 7 – green signals; chromosome 14 – red signal); note the impossibility to identify number of chromosomes 14.

nuclear organization and its driving forces – I-FISH with centromeric, site-specific and wcp, mFISH/QIFSH or ICS-MCB [7,19,31,23,33,35,32,34,57,66,69,72,74,75,93-95,100]; (iii) analysis of chromosome behavior in relation to genome, epigenome and proteome changes for delineation of possible consequences of specific interphase chromosome architecture (i.e. occurring of somatic chromosomal mutations in cancers) – I-FISH with centromeric, site-specific and wcp, mFISH/QIFSH, ICS-MCB and Immuno-FISH [7,15-19,34,69,71,72,74,75,93-95,100,101]. Additional complication of I-FISH analysis of spatial chromosome organization is associated with structural preservation of nuclei. It is to note, that some researchers report about dependence of fixation type on I-FISH results [72,93], whereas others do not [71]. Regardless these debates, an alternative for I-FISH spatial genome analysis could be a suspension FISH (S-FISH) technique [102]. The advantage of this approach is related to possibility of studying three-dimensional (3D) preserved nuclei from any human tissue, whereas other 3D preservation techniques require specific conditions of cell cultivation. The latter makes I-FISH to lose its main advantage. Together, it is to conclude that comprehensive description of functional significance of nuclear organization requires application of almost all known interphase molecular cytogenetic techniques.

Somatic genomic variations

During the last half decade, genomic variations – a source of human healthy and pathological diversity – have become a major focus of current biomedical research. Being involved in evolutionary and disease pathways, variations of the human genome are considered the main target of researches aimed to uncover disease mechanisms and species origins [103]. Soon after description of high rate of interindividual genomic diversification, it has been hypothesized that related processes – somatic genomic variations – lie at the origin of intercellular genomic differences. Moreover, somatic variability of cellular genomes was proposed as a mechanism for complex human diseases [7,10,12]. The latter has been partially confirmed by high-resolution interphase molecular cytogenetic (molecular neurocytogenetic) studies of neurological and psychiatric diseases [7,20-29]. The growing evidence for contribution of somatic genomic variations to the key physiological processes has been used for further hypothesizing about the emerging role of cell-to-cell genome variability in normal/abnormal human intrauterine development (including exogenous effects), cancerization, tissue-specific pathology (i.e. targeted neurodegeneration), sex differences in complex diseases, responses to molecular therapy of debilitating neurological disorders [21,22,24,28,29,104-109]. Altogether, this forms a basis for




Figure 5 ICS-MCB with chromosome 21-specific probe. Monosomy (loss) of chromosome 21 in a nucleus isolated from the Alzheimer's disease brain.

forthcoming researches in the field of single-cell biology. All these achievements were the result of numerous developments in interphase molecular cytogenetics. To prove it, we would like to refer to determination of stochastic (sporadic or background) aneuploidy level in human tissues (Table 3) [20-24,59,28,29,35,81,109-112]. Looking through these data, it is hard to avoid the conclusion that aneuploidy rates become more reasonable if high-resolution I-FISH approaches are applied. Additionally, interindividual genomic variations can be detected in interphase by a parent-of-origin-determination FISH (pod-FISH) technique [113]. Together, I-FISH can be proposed as a required addition for studying genomic variations at microscopic and submicroscopic levels.

Molecular cytogenetic diagnosis

Molecular cytogenetic identification of chromosomal aberrations by I-FISH has been already mentioned in

this review. Here, we would like to make some additional comments related to more specific problems of medical cytogenetics and to show again that studying chromosomes in interphase nuclei has profound effects on molecular cancer and prenatal diagnosis [114,115]. It is obvious that it is almost impossible to refer all the studies that used I-FISH. Here, we have preferred to describe several difficulties encountered during I-FISH introduction and usage for diagnostic purposes. Newly introduced interphase techniques (i.e. ICS-MCB) were used for research purposes only and, therefore, have not been tested for diagnostic validity. Despite of limiting practical application of these I-FISH protocols, related drawbacks can be easily eliminated by forthcoming studies. Another problem comes from the diagnosis of chromosomal mosaicism. There do not exist commonly accepted guidelines or criteria for mosaicism definition [7,10,35]. Regardless some attempts (for details see [35]), there is still no consensus concerning this topic. The




Figure 6 Problems of I-FISH with centromeric/site-specific DNA probes. (A) and (B) replication of specific genomic loci (LSI21 probe) – some nuclei exhibit replicated signals, whereas in some nuclei it is not apparent; note the distance between signals can be more than a diameter of a signal; (C) asynchronous replication of a signal (DXZ1) in case of tetrasomy of chromosome X; note the difficulty to make a definitive conclusion about number of signals in the right nucleus; (D) Two-color FISH with centromeric/site-specific DNA probes for chromosome 1 shows chromosomal associations in a nucleus isolated from the adult human brain; note the impossibility to identify number of chromosomes; (E) QFISH demonstrating an association of centromeric regions of homologous chromosomes 9, but not a monosomy or chromosome loss (for more details see [32]).

solution would be a large-scale study aimed to uncover somatic genomic variations in unaffected human tissues. Hopefully, similar studies have been already launched [20-24,59,28,29,35,81]. Finally, there are still no data or recommendations concerning correlation between metaphase and interphase diagnostic analysis of the same individual. In other words, it is still poorly understood what data is more valid. The structural point of view insists that metaphase analysis of chromosomes is more

precise. From the other hand, mosaics require large cell populations to be analyzed. It becomes even more difficult to solve this problem when cases of complex, hidden (cryptic) or dynamic mosaicism are attempted to be described. Metaphase analysis in these case is indispensable for thorough definition of all cell lines, because simple I-FISH analyses are unable to precise a percentage of each cell line [116,117]. Moreover, some studies require additional data to obtain, i.e. parental origin of




Figure 7 Immuno-FISH. I-FISH using centromeric probe for chromosome Y (DYZ3) with immunostaining by NeuN (neuron-specific antibody) performed for the analysis of cells isolated from the human brain.

Table 3 Data on sporadic aneuploidy in different human tissues (presumably normal) depending on techniques used for the evaluation

Tissue	Technique	Aneuploidy rate	Refs
Ovarian tissues	I-FISH with site-specific probes	Statistically significant proportion of aneuploid cells (trisomy 21)	[109]
Chorionic villi	mFISH/QFISH ICS-MCB	~24% (~1% per chromosome)	[21,24,35]
Fetal human brain	mFISH/QFISH ICS-MCB	~30% (~1.5% per chromosome) ~35% + confined mosaicism	[22,24,35]
Blood	I-FISH with centromeric probes	Chromosome X: 1.5%-2.5% and 4.5%-5%*; Autosomes: 1.2% and 1%	[110]
	mFISH/QFISH	Chromosome X: 1.11%; Autosomes: 0,73%	[25]
Skin	mFISH	2,2% and 4,4%* (whole genome – over 50%)	[111]
Liver	mFISH	~3% (whole genome – over 50%)	[112]
Adult human brain	mFISH/QFISH ICS-MCB	~10% (~0.5% per chromosome)	[20,22,23,26,28,29]

* - in young and old individuals, respectively;

chromosomes or some epigenetic features for more thorough confirmational or exclusive diagnosis. To get this opportunity, it is to apply QFISH [32] or pod-FISH [112].

It is widely accepted that molecular cytogenetic diagnosis should be performed using a panel of techniques [1-10]. It could be either a combination of molecular cytogenetic techniques that use different platforms (i.e. FISH+CGH) or consecutive metaphase and interphase FISH analyses in cases of complex mosaics or balanced structural chromosome abnormalities. Thus, regardless significant developments in the field of molecular interphase cytogenetics, I-FISH techniques remain an addition to metaphase cytogenetics or whole genome screening approaches based on array CGH. The exception is few targeted assays for identification of known cancer-associated translocations in interphase and preimplantation genetic diagnosis. Consequently, I-FISH should be more thoroughly analyzed in terms of the diagnostic potential to take a well-deserved place among genetic testing procedures.

Conclusions and future directions

Structural and behavioral properties of human interphase chromosomes in different tissue/cell types in health and disease remain largely unknown. To date, only fragmentary data on distantly related areas of interphase chromosome biology are available without an integral view of chromosome behavior and arrangement along cell cycle. An overview of molecular cytogenetic techniques for visualizing chromosomes in interphase evidences that a strong technological basis does exist for high-resolution analyses of chromosomes of almost all human tissues. Three main directions of I-FISH application has been advanced by developments in interphase molecular cytogenetics which has provided for possibilities to define functional consequences of spatiotemporal

chromosome arrangement in the nuclei, to elucidate the role of such immense intercellular genomic diversity (somatic genomic variations), to propose new diagnostic solutions for medical genetics and oncology. I-FISH is the unique way to study variations and behavior of the genome in all the cell types of human organism, at all stages of cell cycle and at molecular and supramolecular resolutions. Thus, developments in interphase molecular cytogenetics open numerous prospects for genetics, cellular and molecular biology, genomic/molecular medicine. Taking into account data on technological aspects of studying human interphase chromosomes, we conclude that this biomedical direction has the potential to provide revolutionary solutions for basic and applied biomedical research in fields of human genetics and cell biology. This would be undoubtedly the result of combination of interphase molecular cytogenetic techniques (i.e. mFISH, QFISH, ICS-MCB, S-FISH, pod-FISH, Immuno-FISH etc), which has already given rise to several discoveries in current biomedicine.

Acknowledgements

This review is dedicated to Dr. Ilya V Soloviev. Dr. Thomas Liehr (Jena, Germany) is acknowledged for our fruitful collaboration that allowed us to present examples of ICS-MCB (Figure 5). Authors are supported by DLR/BMBF (RUS 09/006) and Philip Morris USA, Inc.

Author details

¹Institute of Pediatrics and Children Surgery, Rosmedtechnologii, Moscow, 127412, Russia. ²National Research Center of Mental Health, Russian Academy of Medical Sciences, Moscow 119152, Russia.

Authors' contributions

SGV, YBY and IYI wrote the manuscript. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 22 December 2009

Accepted: 11 January 2010 Published: 11 January 2010

References

- Gersen SL, Keagle MB: *The principles of clinical cytogenetics* Totowa, NJ: Humana Press, 2005.
- Liehr T, Claussen U: **Multicolor-FISH approaches for the characterization of human chromosomes in clinical genetics and tumor cytogenetics.** *Curr Genomics* 2002, **3**:231-235.
- Iourov IY, Vorsanova SG, Yurov YB: **Recent patents on molecular cytogenetics.** *Recent Pat DNA Gene Seq* 2008, **2**:6-15.
- Bejjani BA, Shaffer LG: **Clinical utility of contemporary molecular cytogenetics.** *Annu Rev Genomics Hum Genet* 2008, **9**:71-86.
- Liehr T, Starke H, Weise A, Lehrer H, Claussen U: **Multicolor FISH probe sets and their applications.** *Histol Histopathol* 2004, **19**:229-237.
- Speicher MR, Carter NP: **The new cytogenetics: blurring the boundaries with molecular biology.** *Nat Rev Genet* 2005, **6**:782-792.
- Iourov IY, Vorsanova SG, Yurov YB: **Chromosomal variations in mammalian neuronal cells: known facts and attractive hypotheses.** *Int Rev Cytol* 2006, **249**:143-191.
- Carter NP: **Methods and strategies for analyzing copy number variation using DNA microarrays.** *Nat Genet* 2007, **39**:S16-S21.
- Levsky JM, Singer RH: **Gene expression and the myth of the average cell.** *Trends Cell Biol* 2003, **13**:4-6.
- Iourov IY, Vorsanova SG, Yurov YB: **Intercellular genomic (chromosomal) variations resulting in somatic mosaicism: mechanisms and consequences.** *Curr Genomics* 2006, **7**:435-446.
- Emanuel BS, Saitta SC: **From microscopes to microarrays: dissecting recurrent chromosomal rearrangements.** *Nat Rev Genet* 2007, **8**:869-883.
- Iourov IY, Vorsanova SG, Yurov YB: **Chromosomal mosaicism goes global.** *Mol Cytogenet* 2008, **1**:26.
- Liehr T: *Fluorescence in situ hybridization (FISH) – Application guide* Berlin, Heidelberg: Springer 2009.
- Vanneste E, Voet T, Le Caignec C, Ampe M, Konigs P, Mellote C, Debrock S, Amyere M, Vikkula M, Schuit F, Fryns JP, Verbeke G, D'Hooghe T, Moreau Y, Vermeesch JR: **Chromosome instability is common in human cleavage-stage embryos.** *Nat Med* 2009, **15**:577-583.
- Cremer T, Cremer M, Dietzel S, Muller S, Solovei I, Fakan S: **Chromosome territories - a functional nuclear landscape.** *Curr Opin Cell Biol* 2006, **18**:307-316.
- Goetze S, Mateos-Langerak J, van Driel R: **Three-dimensional genome organization in interphase and its relation to genome function.** *Semin Cell Dev Biol* 2007, **18**:707-714.
- Schneider R, Grosschedl R: **Dynamics and interplay of nuclear architecture, genome organization, and gene expression.** *Genes Dev* 2007, **21**:3027-3043.
- Gondor A, Ohlsson R: **Chromosome crosstalk in three dimensions.** *Nature* 2009, **461**:212-217.
- Manvelyan M, Hunstig F, Bhatt S, Mrasek K, Pellestor F, Weise A, Simonian I, Aroutiounian R, Liehr T: **Chromosome distribution in human sperm - a 3D multicolor banding-study.** *Mol Cytogenet* 2008, **1**:25.
- Yurov YB, Vostrikov VM, Vorsanova SG, Monakhov VV, Iourov IY: **Multicolor fluorescent in situ hybridization on post mortem brain in schizophrenia as an approach for identification of low-level chromosomal aneuploidy in neuropsychiatric diseases.** *Brain Dev* 2001, **23**(Suppl 1):186-190.
- Vorsanova SG, Kolotii AD, Iourov IY, Monakhov VV, Kirillova EA, Soloviev IV, Yurov YB: **Evidence for high frequency of chromosomal mosaicism in spontaneous abortions revealed by interphase FISH analysis.** *J Histochem Cytochem* 2005, **53**:375-380.
- Yurov YB, Iourov IY, Monakhov VV, Soloviev IV, Vostrikov VM, Vorsanova SG: **The variation of aneuploidy frequency in the developing and adult human brain revealed by an interphase FISH study.** *J Histochem Cytochem* 2005, **53**:385-390.
- Iourov IY, Liehr T, Vorsanova SG, Kolotii AD, Yurov YB: **Visualization of interphase chromosomes in postmitotic cells of the human brain by multicolour banding (MCB).** *Chromosome Res* 2006, **14**:223-229.
- Yurov YB, Iourov IY, Vorsanova SG, Liehr T, Kolotii AD, Kutsev SI, Pellestor F, Beresheva AK, Demidova IA, Kravets VS, Monakhov VV, Soloviev IV: **Aneuploidy and confined chromosomal mosaicism in the developing human brain.** *PLoS ONE* 2007, **2**:e558.
- Yurov YB, Vorsanova SG, Iourov IY, Demidova IA, Beresheva AK, Kravetz VS, Monakhov VV, Kolotii AD, Voinova-Ulas VY, Gorbachevskaya NL: **Unexplained autism is frequently associated with low-level mosaic aneuploidy.** *J Med Genet* 2007, **44**:521-535.
- Yurov YB, Iourov IY, Vorsanova SG, Demidova IA, Kravetz VS, Beresheva AK, Kolotii AD, Monakhov VV, Uranova NA, Vostrikov VM, Soloviev IV, Liehr T: **The schizophrenia brain exhibits low-level aneuploidy involving chromosome 1.** *Schizophr Res* 2008, **98**:137-147.
- Iourov IY, Vorsanova SG, Yurov YB: **Molecular cytogenetics and cytogenomics of brain diseases.** *Curr Genomics* 2008, **9**:452-465.
- Iourov IY, Vorsanova SG, Liehr T, Yurov YB: **Aneuploidy in the normal, Alzheimer's disease and ataxia-telangiectasia brain: differential expression and pathological meaning.** *Neurobiol Dis* 2009, **34**:212-220.
- Iourov IY, Vorsanova SG, Liehr T, Kolotii AD, Yurov YB: **Increased chromosome instability dramatically disrupts neural genome integrity and mediates cerebellar degeneration in the ataxia-telangiectasia brain.** *Hum Mol Genet* 2009, **18**:2656-2669.
- Yurov YB, Soloviev IV, Vorsanova SG, Marçais B, Roizes G, Lewis R: **High resolution fluorescence in situ hybridization using cyanine and fluorescein dyes: ultra-rapid chromosome detection by directly fluorescently labeled alphoid DNA probes.** *Hum Genet* 1996, **97**:390-398.
- Lemke J, Claussen J, Michel S, Chudoba I, Muhlig P, Westermann P, Sperling K, Rubtsov N, Grummt UW, Ullmann P, Kromeyer-Hauschild K, Liehr T, Claussen U: **The DNA-based structure of human chromosome 5 in interphase.** *Am J Hum Genet* 2002, **71**:1051-1059.
- Iourov IY, Soloviev IV, Vorsanova SG, Monakhov VV, Yurov YB: **An approach for quantitative assessment of fluorescence in situ hybridization (FISH) signals for applied human molecular cytogenetics.** *J Histochem Cytochem* 2005, **53**:401-408.
- Iourov IY, Liehr T, Vorsanova SG, Yurov YB: **Interphase chromosome-specific multicolor banding (ICS-MCB): a new tool for analysis of interphase chromosomes in their integrity.** *Biomol Eng* 2007, **24**:415-417.
- Manvelyan M, Hunstig F, Mrasek K, Bhatt S, Pellestor F, Weise A, Liehr T: **Position of chromosomes 18, 19, 21 and 22 in 3D-preserved interphase nuclei of human and gorilla and white hand gibbon.** *Mol Cytogenet* 2008, **1**:9.
- Iourov IY, Vorsanova SG, Soloviev IV, Yurov YB: **Interphase FISH: detection of intercellular genomic variations and somatic chromosomal mosaicism.** *Fluorescence in situ hybridization (FISH) - Application guide* Berlin, Heidelberg: Springer Verlag/Liehr T 2009, 301-311.
- Liehr T, Claussen U: **Current developments in molecular cytogenetic techniques.** *Curr Mol Med* 2002, **2**:283-297.
- Pellestor F, Ed: **PRINS and In Situ PCR protocols.** *Methods Mol Biol*, 2006, **334**:1-243.
- Pellestor F, Paulasova P, Andreo B, Lefort F, Hamamah S: **Multicolor PRINS and multicolor PNA.** *Cytogenet Genome Res* 2006, **114**:263-269.
- Schrock E, du Manoir S, Veldman T, Schoell B, Weinberg J, Ferguson-Smith MA, Ning Y, Ledbetter DH, Bar-Am I, Soenksen D, Garini Y, Ried T: **Multicolor spectral karyotyping of human chromosomes.** *Science* 1996, **273**:494-497.
- Speicher MR, Ballard GS, Ward DC: **Karyotyping human chromosomes by combinatorial multi-fluor FISH.** *Nat Genet* 1996, **12**:368-375.
- Soloviev IV, Yurov YB, Vorsanova SG, Malet P, Zerova TE, Buzhievskaya TI: **Double color in situ hybridization of alpha-satellite chromosome 13, 21 specific cosmid clones for a rapid screening of their specificity.** *Tsitol Genet* 1998, **32**:60-64.
- Soloviev IV, Yurov YB, Vorsanova SG, Marçais B, Rogaev EI, Kapanadze BI, Brodiansky VM, Yankovsky NK, Roizes G: **Fluorescent in situ hybridization analysis of α -satellite DNA in cosmid libraries specific for human chromosomes 13, 21 and 22.** *Rus J Genet* 1998, **34**:1247-1255.
- Liehr T, Heller A, Starke H, Rubtsov N, Trifonov V, Mrasek K, Weise A, Kuechler A, Claussen U: **Microdissection based high resolution multicolor banding for all 24 human chromosomes.** *Int J Mol Med* 2002, **9**:335-339.
- Nietzel A, Rocchi M, Starke H, Heller A, Fiedler W, Wlodarska I, Loncarevic IF, Beensen V, Claussen U, Liehr T: **A new multicolor-FISH approach for the characterization of marker chromosomes: centromer-specific multicolor-FISH (cenM-FISH).** *Hum Genet* 2001, **108**:199-204.
- Liehr T, Weise A, Heller A, Starke H, Mrasek K, Kuechler A, Weier H-UG, Claussen U: **Multicolor chromosome banding (MCB) with YAC/BAC-based probes and region specific microdissection DNA libraries.** *Cytogenet Genome Res* 2002, **97**:43-50.
- Baumgartner A, Weier JF, Weier H-UG: **Chromosome-specific DNA repeat probes.** *J Histochem Cytochem* 2006, **54**:1363-1370.
- Soloviev IV, Yurov YB, Vorsanova SG, Fayet F, Roizes G, Malet P: **Prenatal diagnosis of trisomy 21 using interphase fluorescence in situ**

- hybridization of postreplicated cells with site-specific cosmid and cosmid contig probes. *Prenat Diagn* 1995, **15**:237-248.
48. Fung J, Weier H-UG, Pedersen RA: **Detection of structural and numerical chromosome abnormalities in interphase cells using spectral imaging.** *J Histochem Cytochem* 2001, **49**:797-798.
49. Stumm M, Wegner R-D, Bloechle M, Eckel H: **Interphase M-FISH applications using commercial probes in prenatal and PGD diagnostics.** *Cytogenet Genome Res* 2006, **114**:296-301.
50. Lu CM, Kwan J, Baumgartner A, Weier JF, Wang M, Escudero T, Munne S, Zitzelsberger HF, Weier H-UG: **DNA probe pooling for rapid delineation of chromosomal breakpoints.** *J Histochem Cytochem* 2009, **57**:587-597.
51. Weier H-UG, Wang M, Mullikin JC, Zhu Y, Cheng JF, Greulich KM, Bensimon A, Gray JW: **Quantitative DNA fiber mapping.** *Hum Mol Genet* 1995, **4**:1903-1910.
52. Weier H-UG: **DNA fiber mapping techniques for the assembly of high-resolution physical maps.** *J Histochem Cytochem* 2001, **49**:939-948.
53. Wells D, Delhanty JDA: **Comprehensive chromosomal analysis of human preimplantation embryos using whole genome amplification and single cell comparative genomic hybridization.** *Mol Hum Reprod* 2000, **6**:1055-1062.
54. Le Caignec C, Spits C, Sermon K, De Rycke M, Thienpont B, Debrock S, Staessen C, Moreau Y, Fryns JP, Van Steirteghem A, Liebaers I, Vermeesch JR: **Single-cell chromosomal imbalances detection by array CGH.** *Nucleic Acids Res* 2006, **34**:e68.
55. McNeil N, Ried T: **Novel molecular cytogenetic techniques for identifying complex chromosomal rearrangements: technology and applications in molecular medicine.** *Expert Rev Mol Med* 2000, **2**:1-14.
56. Aubert G, Lansdorp PM: **Telomeres and aging.** *Physiol Rev* 2008, **88**:557-579.
57. Weierich C, Brero A, Stein S, von Hase J, Cremer C, Cremer T, Solovoi I: **Three-dimensional arrangements of centromeres and telomeres in nuclei of human and murine lymphocytes.** *Chromosome Res* 2003, **11**:485-502.
58. Vorsanova SG, Yurov YB, Ulas VY, Demidova IA, Kolotii AD, Gorbachevskaia NL, Beresheva AK, Soloviev IV: **Cytogenetic and molecular-cytogenetic studies of Rett syndrome (RTT): a retrospective analysis of a Russian cohort of RTT patients (the investigation of 57 girls and three boys).** *Brain Dev* 2001, **23**:S196-S201.
59. Vorsanova SG, Iourov IY, Beresheva AK, Demidova IA, Monakhov VV, Kravets VS, Bartseva OB, Goyko EA, Soloviev IV, Yurov YB: **Non-disjunction of chromosome 21, alloid DNA variation, and sociogenetic features of Down syndrome.** *Tsitol Genet* 2005, **39**(6):30-36.
60. Vorsanova SG, Iourov IY, Demidova IA, Kirillova EA, Soloviev IV, Yurov YB: **Chimerism and multiple numerical chromosome imbalances in a spontaneously aborted fetus.** *Tsitol Genet* 2006, **40**(5):28-30.
61. Vorsanova SG, Iourov IY, Voinova-Ulas VY, Weise A, Monakhov VV, Kolotii AD, Soloviev IV, Novikov PV, Yurov YB, Liehr T: **Partial monosomy 7q34-qter and 21pter-q22.13 due to cryptic unbalanced translocation t(7;21) but not monosomy of the whole chromosome 21: a case report plus review of the literature.** *Mol Cytogenet* 2008, **1**:13.
62. Yurov YB: **In situ hybridization of cloned repeating DNA sequences and differential staining of human chromosomes.** *Bull Exp Biol Med* 1984, **97**:643-647.
63. Vorsanova SG, Yurov YB, Alexandrov IA, Demidova IA, Mitkevich SP, Tirskaia AF: **18p- syndrome: an unusual case and diagnosis by in situ hybridization with chromosome 18-specific alloid DNA sequence.** *Hum Genet* 1986, **72**:185-187.
64. Vorsanova SG, Yurov YB, Kurbatov MB, Kazantseva LZ: **Translocation t(1;17)(q12;q25) with a clinical picture like a proximal deletion of 1q: identification by in situ hybridization with chromosome 1-specific satellite DNA probe.** *Hum Genet* 1990, **86**:173-174.
65. She X, Horvath JE, Jiang Z, Liu G, Furey TS, Christ R, Graves T, Gulden CL, Alkan C, Bailey JA, Sahinalp C, Rocchi M, Haussler D, Wilson RK, Miller W, Schwartz S, Eichler EE: **The structure and evolution of centromeric transition regions within the human genome.** *Nature* 2004, **430**:857-864.
66. Solovoi I, Schermelleh L, Düring K, Engelhardt A, Stein S, Cremer C, Cremer T: **Differences in centromere positioning of cycling and postmitotic human cell types.** *Chromosoma* 2004, **112**:410-423.
67. Kaname T, McGuigan A, Georgiou A, Yurov Y, Osoegawa K, De Jong PJ, Ioannou P, Huxley C: **Alloid DNA from different chromosomes forms de novo minichromosomes with high efficiency.** *Chromosome* 2005, **13**:411-422.
68. Vorsanova SG, Iourov IY, Demidova IA, Kolotii AD, Soloviev IV, Yurov YB: **Pericentric inversion inv(7)(p11q21.1): report on two cases and genotype-phenotype correlations.** *Tsitol Genet* 2006, **40**(3):45-48.
69. Shopland LS, Lynch CR, Peterson KA, Thornton K, Keeper N, van Hase J, Stein S, Vincent S, Molloy KR, Kreth G, Cremer C, Bult CJ, O'Brien TP: **Folding and organization of a contiguous chromosome region according to the gene distribution pattern in primary genomic sequence.** *J Cell Biol* 2006, **174**:27-38.
70. Bolzer A, Kreth G, Solovoi I, Koehler D, Saracoglu K, Fauth C, Müller S, Ellis R, Cremer C, Speicher MR, Cremer T: **Three-dimensional maps of all chromosomes in human male fibroblast nuclei and prometaphase rosettes.** *PLoS Biol* 2005, **3**:e157.
71. Croft JA, Bridger JM, Boyle S, Perry P, Teague P, Bickmore WA: **Differences in the location and morphology of chromosomes in the human nucleus.** *J Cell Biol* 1999, **145**:1119-1131.
72. Cremer T, Cremer C: **Rise, fall and resurrection of chromosome territories: a historical perspective. Part II. Fall and resurrection of chromosome territories during the 1950s to 1980s. Part III. Chromosome territories and the functional nuclear architecture: experiments and models from the 1990s to the present.** *Eur J Histochem* 2006, **50**:223-272.
73. Kosyakova N, Weise A, Mrasek K, Claussen U, Liehr T, Nelle H: **The hierarchically organized splitting of chromosomal bands for all human chromosomes.** *Mol Cytogenet* 2009, **2**:4.
74. Weise A, Starke H, Heller A, Claussen U, Liehr T: **Evidence for interphase DNA decondensation transverse to the chromosome axis: a multicolour banding analysis.** *Int J Mol Med* 2002, **9**:359-361.
75. Manvelyan M, Kempf P, Weise A, Mrasek K, Heller A, Liehr A, Hoffken K, Fricke HJ, Sayer HG, Liehr T, Mkrтчan H: **Preferred co-localization of chromosomes 8 and 21 in myeloid bone marrow cells detected by three dimensional molecular cytogenetics.** *Int J Mol Med* 2009, **24**:335-341.
76. Lu CM, Wang M, Greulich-Bode K, Weier JF, Weier HUG: **Quantitative DNA fiber mapping.** *Fluorescence in situ hybridization (FISH) - Application guide* Berlin, Heidelberg: Springer Verlag Liehr T 2009, 269-291.
77. Pinkel D, Albertson DG: **Comparative genomic hybridization.** *Annu Rev Genomics Hum Genet* 2005, **6**:331-354.
78. Iourov IY, Vorsanova SG, Pellestor F, Yurov YB: **Brain tissue preparations for chromosomal PRINS labeling.** *Methods Mol Biol* 2006, **334**:123-132.
79. Iourov IY: **Microscopy and imaging systems.** *Fluorescence in situ hybridization (FISH) - Application guide* Berlin, Heidelberg: Springer Verlag Liehr T 2009, 75-84.
80. Liehr T, Ed: **Multicolor FISH in human cytogenetics.** *Cytogenet Genome Res* 2006, **114**:183-389.
81. Iourov IY, Vorsanova SG, Yurov YB: **Detection of aneuploidy in neural stem cells of the developing and adult human brain.** *Electron J Biol* 2008, **4**:36-42.
82. Vorsanova SG, Yurov YB, Deryagin GV, Soloviev IV, Bytenskaya GA: **Diagnosis of aneuploidy by in situ hybridization: analysis of interphase nuclei.** *Bull Exp Biol Med* 1991, **112**:413-415.
83. Lee C, Wevrick R, Fisher RB, Ferguson-Smith MA, Lin CC: **Human centromeric DNAs.** *Hum Genet* 1997, **100**:291-304.
84. Verma RS, Luke S: **Variation in alloid DNA sequences escape detection of aneuploidy in interphase FISH technique.** *Genomics* 1992, **14**:113-116.
85. Liehr T, Pfeiffer RA, Trautmann U, Gebhart E: **Centromeric alloid DNA heteromorphisms of chromosome 22 as revealed by FISH-technique.** *Clin Genet* 1998, **53**:231-232.
86. Vorsanova SG, Yurov YB, Brusquant D, Carles E, Roizes G: **Two new cases of the christchurch (Ch1) chromosome 21: evidence for clinical consequences of de novo deletion 21p-.** *Tsitol Genet* 2002, **36**(1):46-49.
87. Sinclair PB, Green AR, Grace C, Nacheva EP: **Fluorescence in situ hybridization system improved sensitivity of BCR-ABL detection: a triple-probe three-color.** *Blood* 1997, **90**:1395-1402.
88. Mitelman F, Johanson B, Martens F: **The impact of translocations and gene fusions on cancer causation.** *Nat Rev Cancer* 2007, **7**:233-245.
89. Virgili A, Brazza D, Reid AG, Howard-Reeves J, Valgañón M, Chanalaris A, De Melo VAS, Marin D, Apperley JF, Grace C, Nacheva EP: **FISH mapping of Philadelphia negative BCR/ABL1 positive CML.** *Mol Cytogenet* 2008, **1**:14.

90. Kucheria K, Jobanputra V, Talwar R, Ahmad ME, Dada R, Sivakumaran TA: **Human molecular cytogenetics: diagnosis, prognosis, and disease management.** *Teratogenesis Carcinog Mutagen Suppl* 1:225-233.
91. Halder A, Jain M, Kabra M, Gupta N: **Mosaic 22q11.2 microdeletion syndrome: diagnosis and clinical manifestations of two cases.** *Mol Cytogenet* 2008, **1**:18.
92. Portnoi MF: **Microduplication 22q11.2: a new chromosomal syndrome.** *Eur J Med Genet* 2009, **52**:88-93.
93. Walter J, Joffe B, Bolzer A, Albiez H, Benedetti PA, Muller S, Speicher MR, Cremer T, Cremer M, Solovei I: **Towards many colors in FISH on 3D-preserved interphase nuclei.** *Cytogenet Genome Res* 2006, **114**:367-378.
94. Krueger C, Osborne CS: **Raising the curtains on interchromosomal interactions.** *Trends Genet* 2006, **22**:637-639.
95. Iourov IY, Vorsanova SG, Yurov YB: **Fluorescence intensity profiles of in situ hybridization signals depict genome architecture within human interphase nuclei.** *Tsitol Genet* 2008, **42**(5):3-8.
96. Speel EJM, Herbergs J, Ramaekers FCS, Hopman AHN: **Combined immunocytochemistry and fluorescence in situ hybridization for simultaneous tricolor detection of cell cycle, genomic, and phenotypic parameters of tumor cells.** *J Histochem Cytochem* 1994, **42**:961-966.
97. Marzais B, Vorsanova SG, Roizes G, Yurov YB: **Analysis of alphoid DNA variation and kinetochore size in human chromosome 21: evidence against pathological significance of alphoid satellite DNA diminutions.** *Tsitol Genet* 1999, **33**(1):25-31.
98. Dundas SR, Boyle S, Bellamy CO, Hawkins W, Garden OJ, Ross JA, Bickmore W: **Dual Y-chromosome painting and immunofluorescence staining of archival human liver transplant biopsies.** *J Histochem Cytochem* 2001, **49**:1321-1322.
99. Yang F, Shao C, Vedanarayanan V, Ehrlich M: **Cytogenetic and immuno-FISH analysis of the 4q subtelomeric region, which is associated with facioscapulohumeral muscular dystrophy.** *Chromosoma* 2004, **112**:350-359.
100. Zinner R, Teller K, Versteeg R, Cremer T, Cremer M: **Biochemistry meets nuclear architecture: multicolor immuno-FISH for co-localization analysis of chromosome segments and differentially expressed gene loci with various histone methylations.** *Adv Enzyme Regul* 2007, **47**:223-241.
101. Meaburn KJ, Gudla PR, Khan S, Lockett SJ, Misteli T: **Disease-specific gene repositioning in breast cancer.** *J Cell Biol* 2009, **187**:801-812.
102. Steinhäuser U, Starke H, Nietzel A, Lindenau J, Ullmann P, Claussen U, Liehr T: **Suspension (S)-FISH, a new technique for interphase nuclei.** *J Histochem Cytochem* 2002, **50**:1697-1698.
103. Zhang F, Gu W, Hurles ME, Lupski JR: **Copy number variation in human health, disease, and evolution.** *Annu Rev Genomics Hum Genet* 2009, **10**:451-81.
104. Iourov IY, Yurov YB, Vorsanova SG: **Mosaic X chromosome aneuploidy can help to explain the male-to-female ratio in autism.** *Med Hypotheses* 2008, **70**:456.
105. Vorsanova SG, Yurov YB, Iourov IY: **Maternal smoking as a cause of mosaic aneuploidy in spontaneous abortions.** *Med Hypotheses* 2008, **71**:607.
106. Iourov IY, Vorsanova SG, Yurov YB: **Developmental chromosome instability as a possible cause of childhood brain cancers.** *Med Hypotheses* 2009, **72**:615-616.
107. Yurov YB, Iourov IY, Vorsanova SG: **Neurodegeneration mediated by chromosome instability suggests changes in strategy for therapy development in ataxia-telangiectasia.** *Med Hypotheses* 2009, **73**:1075-1076.
108. Yurov YB, Vorsanova SG, Iourov IY: **GIN'n'CIN hypothesis of brain aging: deciphering the role of somatic genetic instabilities and neural aneuploidy during ontogeny.** *Mol Cytogenet* 2009, **2**:23.
109. Hulten MA, Patel SD, Tankimanova M, Westgren M, Papadogiannakis N, Johnson AM, Iwarsson E: **On the origin of trisomy 21 Down syndrome.** *Mol Cytogenet* 2008, **1**:21.
110. Guttenbach M, Koschorz B, Bernthaler U, Grimm T, Schmid M: **Sex chromosomes loss and aging: in situ hybridization studies on human interphase nuclei.** *Am J Hum Genet* 1995, **57**:1143-1150.
111. Geigl JB, Langer S, Barwisch S, Pfliegerhaer K, Lederer G, Speicher MR: **Analysis of gene expression patterns and chromosomal changes associated with aging.** *Cancer Res* 2004, **64**:8850-8857.
112. Wilkens L, Flemming P, Gebel M, Bleck J, Terkamp C, Wingen L, Kreipe H, Schelgelberger B: **Induction of aneuploidy by increasing chromosomal instability during dedifferentiation of hepatocellular carcinoma.** *Proc Natl Acad Sci USA* 2004, **101**:1309-1314.
113. Weise A, Gross M, Mrasek K, Mkrtychyan H, Horsthemke B, Jonsrud C, Von Eggeling F, Hinreiner S, Witthuhn V, Claussen U, Liehr T: **Parental-origin-determination fluorescence in situ hybridization distinguishes homologous human chromosomes on a single-cell level.** *Int J Mol Med* 2008, **21**:189-200.
114. Liehr T, Ziegler M: **Rapid prenatal diagnostics in the interphase nucleus: procedure and cut-off rates.** *J Histochem Cytochem* 2005, **53**:289-291.
115. Nicholson JM, Duesberg P: **On the karyotypic origin and evolution of cancer cells.** *Cancer Genet Cytogenet* 2009, **194**:96-110.
116. Carreira IM, Mascarenhas A, Matoso E, Couceiro AB, Ramos L, Dufke A, Mazauric M, Stressig R, Kosyakova N, Melo JB, Liehr T: **Three unusual but cytogenetically similar cases with up to five different cell lines involving structural and numerical abnormalities of chromosome 18.** *J Histochem Cytochem* 2007, **55**:1123-1128.
117. Iourov IY, Vorsanova SG, Liehr T, Monakhov VV, Soloviev IV, Yurov YB: **Dynamic mosaicism manifesting as loss, gain and rearrangement of an isodicentric Y chromosome in a male child with growth retardation and abnormal external genitalia.** *Cytogenet Genome Res* 2008, **121**:302-306.

doi:10.1186/1755-8166-3-1

Cite this article as: Vorsanova et al.: Human interphase chromosomes: a review of available molecular cytogenetic technologies. *Molecular Cytogenetics* 2010 **3**:1.

Submit your next manuscript to BioMed Central
and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

