

Hydrogen Peroxide Metabolism in Soybean Embryonic Axes at the Onset of Germination¹

Received for publication January 15, 1987 and in revised form April 30, 1987

SUSANA PUNTARULO, RODOLFO A. SÁNCHEZ, AND ALBERTO BOVERIS*
*Instituto de Química y Fisicoquímica Biológicas, Facultad de Farmacia y Bioquímica (S.P., A.B.),
Departamento de Ecología, Facultad de Agronomía R.A.S., Universidad de Buenos Aires, Junín 956,
1113 Buenos Aires, Argentina*

ABSTRACT

Hydrogen peroxide steady state levels of 5 micromolar were determined in soybean (*Glycine max*) embryonic axes incubated for 2 hours and in axes pretreated with aminotriazole or cyanide, where these levels were 50 and 1 micromolar, respectively. The activities of catalase (105 picomoles H₂O₂ per minute per axis), peroxidase (10–44 picomoles H₂O₂ per minute per axis), glutathione peroxidase (3 picomoles H₂O₂ per minute per axis) and superoxide dismutase (3.5 units per axis), were also determined. Catalase seems to be the most important H₂O₂ consuming enzyme at the physiological concentration of H₂O₂. A short treatment with aminotriazole, while substantially increasing H₂O₂ level, did not affect the growth of the axes. The production of superoxide anion by the mitochondria isolated from soybean axes was measured from the superoxide dismutase-sensitive rate of adrenochrome formation in the presence of NADH or succinate as substrate and amounted to 1.3 and 0.8 nanomole O₂⁻ per minute per milligram protein, respectively. According to the stoichiometry of O₂⁻ and H₂O₂ dismutation reactions, it is apparent that about 0.9 to 1.5% of the total oxygen uptake proceeds through the formation of the free intermediates of the partial reduction of oxygen.

in the germination process; (b) that catalase utilizes an important fraction of the H₂O₂ produced; and (c) that both H₂O₂ production and H₂O₂ utilization to oxidize NADPH are cyanide insensitive.

In this paper we report a measurement of H₂O₂ production and of the activities of the enzymes involved in H₂O₂ metabolism at an early stage of soybean embryonic axis imbibition.

MATERIALS AND METHODS

Soybean seeds (*Glycine max*, cv Hood) recently harvested, were air-dried to 12% humidity at room temperature and stored at 20°C in an evacuated desiccator until used; germination capacity remained over 92% throughout.

Measurements of H₂O₂. Embryonic axes incubated for 2 h at 26°C over water-saturated filter paper were used for the determination of H₂O₂ steady state levels. The diffusates were prepared by soaking 80 axes in 3 ml of 40 mM potassium phosphate buffer (pH 7.0). Two aliquots of 0.3 ml were taken at the indicated times; one of them was supplemented with 10 μM catalase for 10 min, and H₂O₂ was assayed in both aliquots by measuring the peak values of chemiluminescence produced in the reaction of H₂O₂ with luminol (28, 33). The reaction was carried out in a scintillation vial containing 2.70 ml of 40 mM potassium phosphate buffer (pH 7.8), 20 μM FeSO₄, and 10 μM luminol (Sigma). Light emission was measured, for 30 s, immediately after adding the aliquots to the luminol solution and analyzed in a Packard Tri-Carb model 3320 scintillation counter in the out-of-coincidence mode. Reagents and vials were dark-adapted before use.

When the embryonic axes were previously incubated with aminotriazole, aliquots of 0.03 ml were taken, diluted to 0.3 ml and processed as described before. When the axes were incubated in 16 or 160 μM H₂O₂ solutions, samples of 0.1 ml or 0.01 ml were diluted to 0.3 ml and processed as described.

Alternatively, H₂O₂ concentration in the diffusates was measured by formation of the enzyme-substrate complex of horseradish peroxidase (5).

Isolation of Mitochondria. Mitochondria were isolated from (1000–2000) soybean embryonic axes that were incubated on filter paper covering wet cotton for 2 h at 26°C following the procedure described by Bonner (4) and modified according to Beconi *et al.* (1). The axes were homogenized in 150 ml of 0.3 M mannitol, 1 mM EDTA, 4 mM cysteine, and 5 mM Tris-HCl (pH 7.0), in a mortar during 4 min and passed twice through a Potter-Elvehjem homogenizer. The homogenate was filtered through three layers of cheesecloth and centrifuged at 18,000g for 10 min in a Sorvall RC-2 centrifuge to sediment mitochondria. The sediment was suspended in 20 ml of 0.45 M mannitol, 1 mM EDTA, and 5 mM Tris-HCl (pH 7.0), homogenized and centrifuged at 270g for 10 min to sediment nuclei, cell debris,

The physiological and biochemical events taking place during the initial stages of seed imbibition have been the subject of numerous studies (3, 16). A number of processes such as membrane reorganization (29) and metabolic reactivation (2, 20, 26) occurring at this stage can have profound influence on seed germination and on the future growth of the seedling (21, 34).

Hydroperoxide metabolism in plants in general and particularly during seed germination has received little attention, although information about hydroperoxide production and utilization could be most relevant to understand important processes such as lipid peroxidation and NADPH redox changes. Several authors have indicated that an increased hydroperoxide production could be linked to a shift to the oxidized state of the NADPH/NADP⁺ couple (7, 15, 25, 31). Taylorson and Hendricks (32), in order to explain the role of cyanide in breaking seed dormancy, proposed that the inhibition of catalase by cyanide could result in a higher steady state level of H₂O₂ which would lead, through an enhanced peroxidase activity, to a lower NADPH/NADP⁺ ratio and to a stimulation of the pentose phosphate pathway. Such an interpretation has three implicit assumptions: (a) that there is a significant H₂O₂ production early

¹ Supported by grants from Consejo Nacional de Investigaciones Científicas y Técnicas, Argentina (CONICET). R. A. S. and A. B. are Career Investigators from CONICET.

cellulose fibers, and so forth. The supernatant was centrifuged at 14,500g for 10 min. The mitochondrial pellet was washed twice and the final suspension (8–12 mg of protein/ml) was made with the same mannitol-EDTA-Tris solution. All operations were performed at 0–2°C.

Enzyme Assays. For the measurement of enzyme activities, dissected embryonic axes were incubated for 2 h on wet filter paper at 26°C and then homogenized in 50 mM potassium phosphate buffer (pH 7.0), followed by centrifugation at 700g for 10 min, at 0 to 2°C. The supernatant was used as the source of enzymes. All enzymes were assayed at 26°C.

Superoxide dismutase activity was assayed spectrophotometrically at 480 nm by the method of Misra and Fridovich (17), in a reaction medium containing glycine-NaOH buffer 50 mM (pH 9.0) and 1 mM epinephrine. The amount of homogenate inhibiting by 50% the oxidation of epinephrine to adrenochrome was taken as 1 unit (adrenochrome assay) of superoxide dismutase activity. Under the conditions of the assay, this amounts to 0.12 μg of superoxide dismutase/ml.

Ascorbate peroxidase and dehydroascorbate reductase activities were assayed spectrophotometrically (E_{265} ascorbate = 15 mm⁻¹·cm⁻¹). The reaction medium for ascorbate peroxidase consisted of 50 mM potassium phosphate buffer (pH 7.0), 0.05 mM ascorbate, 0.1 mM H₂O₂, and 0.1 mM EDTA. A correction was made for the nonenzymic rate of ascorbate oxidation by H₂O₂ (19, 23). The reaction medium for dehydroascorbate reductase consisted of 50 mM potassium phosphate buffer (pH 7.0), 1.5 mM GSH, 0.2 mM dehydroascorbic acid, and 0.1 mM EDTA. Reaction rates were corrected for nonenzymic reduction of dehydroascorbate by GSH (18, 19).

Peroxidase activity was measured spectrophotometrically at 430 nm ($E = 2.47$ mm⁻¹ cm⁻¹) in a reaction medium consisting of 50 mM potassium phosphate buffer (pH 7.0), 5 mM pyrogallol, and 1 mM H₂O₂ (27).

Glutathione peroxidase activity was measured spectrophotometrically at 340 nm. The assay mixture contained 0.5 mM *t*-butyl-hydroperoxide, 1 mM GSH, 0.1 unit/ml glutathione reductase, and 200 μM NADPH in 50 mM potassium phosphate buffer (pH 7.0) (12).

Catalase was assayed by back titration with 0.05 N KMnO₄ of the remaining H₂O₂ in a reaction medium consisting of 50 mM phosphate buffer (pH 7.0), 60 mM sodium perborate. Aliquots of the assay mixture were taken after 30, 60, and 90 s and the reaction was stopped by addition of 1 M H₂SO₄ (11).

Other Determinations. Determination of O₂ uptake of isolated soybean embryonic axes were made polarographically using a Clark oxygen electrode, in a reaction medium consisting of 50 mM potassium phosphate buffer (pH 7.0).

The effect of H₂O₂ and aminotriazole on the growth of the seedlings was performed by incubating the axes with H₂O₂ and aminotriazole for 15 min; afterwards the axes were removed and placed on wet paper and weighed after 48 h.

Production of superoxide radical was determined as the superoxide dismutase-sensitive rate of adrenochrome formation, measured at 485 to 575 nm ($E = 2.97$ mm⁻¹ cm⁻¹) in a Perkin-Elmer dual wavelength spectrophotometer. The reaction medium consisted of 0.23 M mannitol, 0.07 M sucrose, 1 mM EDTA, 30 mM Tris-HCl (pH 7.0), and 1 mM epinephrine (5).

RESULTS

The light emission of the reaction of H₂O₂ and luminol allowed us to measure the amount of H₂O₂ diffused out of soybean embryonic axes. The kinetics of the reaction showed maximal luminescence in the first 30 s after the reactants were mixed (Fig. 1); therefore all determinations were made at this initial time interval and considering emission peak values. Preincubation of both H₂O₂ standard solutions and aliquots of the diffusate from

the axes with catalase removed most of the signal showing that H₂O₂ was the main reactant for light emission and that other molecular species capable of reacting with luminol were quantitatively less important (Fig. 1).

The catalase-sensitive luminol emission of diffusate aliquots




FIG. 1. Luminol-dependent chemiluminescence in H₂O₂ solutions and diffusate from soybean embryonic axes: 0.5 μM H₂O₂ (●); 2 min-incubated diffusate (▲); 2 min-incubated diffusate treated with 10 μM catalase (○). Inset: Luminol-dependent and catalase-sensitive chemiluminescence of H₂O₂ solutions. The abscissa values indicate H₂O₂ concentrations in the reaction mixture in the scintillation vials.




FIG. 2. H₂O₂ concentrations in diffusates from soybean embryonic axes and external incubation medium. A, H₂O₂ in the diffusate (●); H₂O₂ in the diffusate in the presence of 0.1 mM cyanide (▲); and H₂O₂ in the external medium when the soybean embryonic axes were initially added with 16 μM H₂O₂ (○). H₂O₂ in the diffusate measured by formation of horseradish peroxidase-H₂O₂ compound (■). B, H₂O₂ in the diffusate (●) and external incubation medium (○) when the soybean embryonic axes were pretreated with aminotriazole. Inset: H₂O₂ equilibrium concentration in the external medium as function of aminotriazole concentration.

Table I. Effect of Aminotriazole and H₂O₂ in the Growth of Soybean Seedlings

Treatment	Concentration	Seedling Weight
	mM	mg
Control		35.3 ± 0.7
Aminotriazole	0.1	36.1 ± 0.8
	10	35.7 ± 0.7
	50	36.7 ± 0.6
	100	33.3 ± 0.9
H ₂ O ₂	0.8	32.5 ± 0.5
	4.1	32.2 ± 0.4
	8.3	34.5 ± 0.8
	8300	13.2 ± 0.8

increased with the time of incubation up to a maximum equivalent to $5 \mu\text{M}$ H_2O_2 in the diffusate; this situation was reached in about 8 to 10 min. In this condition the diffusate apparently reached an equilibrium with the intracellular steady state level of H_2O_2 (Fig. 2A). Similar results were obtained when the H_2O_2 in the diffusate was measured by formation of the enzyme-substrate complex of horseradish peroxidase; again the steady state concentration of H_2O_2 was about $5 \mu\text{M}$. When the axes were incubated with $16 \mu\text{M}$ H_2O_2 in the external medium, the H_2O_2 concentration measured by the luminol assay decreased with time to reach a concentration of about $5 \mu\text{M}$ with a t one-half of 4 min (Fig. 2A). These results indicate that in this case H_2O_2 in the external medium reaches also a diffusion equilibrium with the cell internal medium and that the steady state concentration of H_2O_2 in the soybean axes is about $5 \mu\text{M}$.

Cyanide, which inhibits catalase activity as well as the mitochondrial production of H_2O_2 (see below), severely depressed the H_2O_2 steady state concentration (Fig. 2A).

The pretreatment of the axes during 15 min, with aminotriazole (an irreversible inhibitor of catalase [22]) before assaying for H_2O_2 , significantly increased the H_2O_2 in the diffusate (Fig. 2B). The H_2O_2 in the diffusate of aminotriazole-treated axes reached a level of about $50 \mu\text{M}$ which seems to be the steady state concentration of H_2O_2 in the axes when this inhibitor suppressed catalase activity. This effect of aminotriazole (Fig. 2B, inset) shows the importance of catalase in controlling the steady state level H_2O_2 . Aminotriazole-treated axes incubated in an external medium containing 0.16 mM H_2O_2 decreased this level to $50 \mu\text{M}$ H_2O_2 with a t one-half of 21 min (Fig. 2B).

The treatment with aminotriazole did not affect the growth of the axes (Table I). Treatment with up to 100 mM H_2O_2 did not affect axis growth, which may be explained by the rapid exodiffusion of H_2O_2 and by H_2O_2 utilization.

Figure 3A shows O_2^- production by the isolated mitochondria in the presence of NADH which functions as substrate of the NADH dehydrogenase of the external side of the inner mitochondrial membrane (10). The rate of O_2^- production was calculated from the superoxide dismutase-sensitive rate of adrenochrome formation.

The traces of Figure 3A also show that cyanide inhibits O_2^- production by mitochondria in agreement with similar observations in animal and plant mitochondria (13, 24). Figure 3B shows the rate of O_2^- production in the presence of succinate and glutamate as substrates.

The activities of catalase and other H_2O_2 consuming enzymes were measured in homogenates of soybean embryonic axes (Table II). Figure 4 gives an integrated picture of the various metabolic pathways related to the production and utilization of H_2O_2




FIG. 3. Production of O_2^- by mitochondria isolated from soybean embryonic axes; $0.59 \text{ mg protein/ml}$. The numbers near the traces indicate the rate of O_2^- production in $\text{nmol/min} \cdot \text{mg protein}$.

Table II. Activities of Enzymes Involved in H_2O_2 Metabolism

Enzyme	Activity <i>pmol H₂O₂/min · axis</i>
Catalase ^a	105
Peroxidase (pyrogallol)	44
Peroxidase (ascorbate)	10
Glutathione peroxidase	3
Superoxide dismutase ^b	115
Dehydroascorbate reductase	12
Oxygen uptake (<i>pmol O₂/min · axis</i>)	4200

^a Assuming H_2O_2 in the axes equal to $5 \mu\text{M}$. ^b Assuming O_2^- in the axes equal to 0.1 nM . Corresponds to 3.5 units/axis (17).

according to the results presented in this paper. The rate of mitochondrial O_2^- production ($230 \text{ pmol/min} \cdot \text{axis}$) was calculated from the specific rate of O_2^- production in the presence of external NADH as substrate (Fig. 3A) and the amount of mitochondrial membranes in the axis ($0.19 \text{ mg protein/axis}$). This rate of O_2^- production will provide a rate of $115 \text{ pmol H}_2\text{O}_2/\text{min} \cdot \text{axis}$ according to the stoichiometry of the dismutation reaction. Peroxidase activity using either pyrogallol or ascorbate as hydrogen donor could utilize either 44 or $10 \text{ pmol H}_2\text{O}_2/\text{min} \cdot \text{axis}$, respectively. Glutathione peroxidase could be able to use $3 \text{ pmol H}_2\text{O}_2/\text{min} \cdot \text{axis}$. All these rates were calculated assuming maximal levels of hydrogen donor.

In the case that mitochondrial H_2O_2 was the only significant H_2O_2 source and that glutathione peroxidase and peroxidase (ascorbate as donor), acted at maximal H_2O_2 -utilizing rates, catalase would use the rest of H_2O_2 production, *i.e.* $102 \text{ pmol H}_2\text{O}_2/\text{min} \cdot \text{axis}$. If utilization of H_2O_2 by catalase occurs according to $d[\text{H}_2\text{O}_2]/dt = k[\text{H}_2\text{O}_2][\text{Cat}]$ (7) with $k = 4.6 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$ and the steady state value of $[\text{H}_2\text{O}_2]$ equal to $5 \mu\text{M}$, the rate of H_2O_2 utilization amounts to $105 \text{ pmol H}_2\text{O}_2/\text{min} \cdot \text{axis}$.

DISCUSSION

The results of this study indicate that H_2O_2 -producing and H_2O_2 -consuming processes are active early in the germination of soybean embryonic axes. Mitochondrial electron transfer appears to be the most important H_2O_2 source. Mitochondrial H_2O_2 is generated through the primary production of O_2^- and its subsequent dismutation (13). The rate of O_2^- production measured in mitochondria isolated from soybean axes agrees with values found in submitochondrial particles from potato tubers (6), mung bean hypocotyls (24), and *Arum maculatum* spadices (14). These values also agree with the rates of H_2O_2 production measured in mitochondria isolated from potato tubers and mung bean hypocotyls (13, 23).

From the ratio of O_2^- production and total O_2 uptake (Table II) it can be estimated that about 3.6 to 5.9% of the electron flow in mitochondria (extramitochondrial O_2 uptake is not considered) results in univalent reduction of oxygen, depending on whether succinate-glutamate or external NADH are considered the physiological mitochondrial substrate. Due to the stoichiometry of O_2^- and H_2O_2 dismutations (6) it follows that about 0.9 to 1.5% of the total oxygen uptake proceeds through the formation of the free intermediates of the partial reduction of oxygen.

The concept of the mitochondrial membranes as an important source of intracellular H_2O_2 is supported by the cyanide sensitivity of H_2O_2 steady state levels, a property of the mitochondrial generation of O_2^- and H_2O_2 (24).

The large effect of aminotriazole (Fig. 3) on the H_2O_2 steady state level supports the view of a predominant utilization of H_2O_2 by catalase. Other H_2O_2 consuming enzymes like peroxidase and glutathione peroxidase are also active in the axes early on the imbibition phase. However, they could utilize at most 35% of




FIG. 4. Scheme of H₂O₂ metabolism in soybean embryonic axes. The numbers indicate the flow of metabolites in pmol/min·axis. UQ, ubiquinone; SOD, superoxide dismutase.

total H₂O₂, assuming saturation with their hydrogen donors (aromatic hydrogen donors, ascorbate, reduced glutathione, etc.). This result agrees with the effect of aminotriazole on the H₂O₂ levels of homogenates from germinating castor beans as reported by Warm and Laties (33).

The estimation of the rate of H₂O₂ utilization by catalase is hindered by insufficient information about the occurrence of extent of the phenomenon of latency (9) and the activity of extramitochondrial sources of H₂O₂ in the soybean axes. The rate of H₂O₂ utilization by catalase appears to be between 102 to 105 pmol/min·axis. The first value corresponds to a situation in which mitochondria are the only source of H₂O₂ and utilization of H₂O₂ by using aromatic hydrogen donors is negligible. The second value is calculated assuming that intracellular H₂O₂ is equal to 5 μM. At any rate, catalase will utilize between 91 and 97% of the intracellular H₂O₂.

The concentration of H₂O₂ measured in the diffusate (about 5 μM) by following both (a) the decrease of an added concentration of H₂O₂ and (b) the increase up to about the same value starting with no H₂O₂ in the medium, seems a reasonable indication of the intracellular H₂O₂ steady state level. The rapid establishment of a steady state situation similar to the intracellular level is in agreement with the large permeability of plant membranes to H₂O₂ (30) and with the rapid effects of aminotriazole and cyanide. The measured H₂O₂ levels are lower than those reported by Warm and Laties (33) in potato tubers, tomatoes, and germinating castor beans (224, 92, and 138 μmol/kg, fresh weight, respectively).

The steady state concentration of H₂O₂ is thought to have important physiological implications. A higher H₂O₂ level might affect the axis growth potential by increasing lipid peroxidation and other damaging reactions related to oxyradicals (15). However, the short treatment of the axes with aminotriazole, although substantially increasing the H₂O₂ level did not affect their growth (Table I).

The proposal of Taylorson and Hendricks (32) that the inhibition of catalase by cyanide would lead to an enhanced rate of NADPH oxidation and pentose phosphate activity was based on the implicit assumptions mentioned in the beginning of this paper. So far, no evidence was available to substantiate these assumptions (3). Our results show that H₂O₂ production is an early event and that catalase appears to be the predominant H₂O₂ consuming activity. Moreover, the measured glutathione peroxidase activity offers a cyanide insensitive pathway to oxidize

NADPH by H₂O₂. Therefore, an increase in the steady state concentration of H₂O₂ by a treatment with cyanide would only be possible if catalase were more sensitive to the inhibition than the H₂O₂ production system.

LITERATURE CITED

1. BECONI MT, RA SÁNCHEZ, A BOVERIS 1983 Alternative oxidase and cyanide- and antimycin-insensitive respiration in corn root mitochondria. *Plant Sci Lett* 32: 125-132
2. BERJACK P, TA VILLIERS 1972 Ageing in plant embryos II. Age-induced damage and its repair during early germination. *New Phytol* 71: 135-144
3. BEWLEY JD, M BLACK 1982 *In Physiology and Biochemistry of Seeds in Relation to Germination*, Vol 1. Springer-Verlag, Berlin
4. BONNER WD JR 1967 A general method for the preparation of plant mitochondria. *Methods Enzymol* 10: 126-133
5. BOVERIS A 1984 Oxygen radicals in biological systems. *Methods Enzymol* 105: 429-435
6. BOVERIS A, RA SÁNCHEZ, MT BECONI 1978 Antimycin- and cyanide-resistant respiration and superoxide anion production in fresh and aged potato tuber mitochondria. *FEBS Lett* 92: 333-338
7. CHANCE B, H SIES, A BOVERIS 1979 Hydroperoxide metabolism in mammalian organs. *Physiol Rev* 59: 527-605
8. DE CHATELET LR, G LONG, PS SHIRLEY, DA BASS, MJ THOMAS, FW HENDERSON, MS COHEN 1982 Mechanism of the luminol-dependent chemiluminescence of human neutrophils. *J Immunol* 129: 1589-1593
9. DE DUVE C, P BANDHUIN 1966 Peroxidase (Microbodies) and related particles. *Physiol Rev* 46, 323-355
10. DOUCE R, CA MANNELA, WD BONNER JR 1973 The external NADH dehydrogenase of intact plant mitochondria. *Biochim Biophys Acta* 292: 105-116
11. FEINSTEIN R 1949 Perborate as substrate in a new assay of catalase. *J Biol Chem* 180: 1197-1202
12. FLOHE L, WA GUNZLER 1984 Assays of glutathione peroxidase. *Methods Enzymol* 105: 114-121
13. FORMAN HJ, A BOVERIS 1982 Superoxide radical and hydrogen peroxide in mitochondria. *In* WA Pryor, ed, *Free Radicals in Biology*, Vol 5. Academic Press, New York, pp 65-87
14. HUO S, JM PALMER 1978 Superoxide and hydrogen peroxide production in cyanide-resistant *Arum maculatum* mitochondria. *Plant Sci Lett* 11: 351-358
15. LESHEM YY 1984 Interaction of cytokinins with lipid-associated oxy free radicals during senescence: a prospective mode of cytokinin action. *Can J Bot* 62: 2943-2949
16. MEYER AM 1977 Metabolic control of germination. *In* AA Khan, ed. *The Physiology and Biochemistry of Seed Development, Dormancy and Germination*. North Holland, Amsterdam, pp 357-384
17. MISRA H, I FRIDOVICH 1972 The role of superoxide anion in the autoxidation of epinephrine and a simile assay for superoxide dismutase. *J Biol Chem* 247: 3170-3175
18. NAKANO Y, K ASADA 1980 Spinach chloroplast scavenge hydrogen peroxide on illumination. *Plant Cell Physiol* 21: 1295-1307
19. NAKANO Y, K ASADA 1981 Hydrogen peroxide is scavenged by ascorbate specific peroxidase in spinach chloroplasts. *Plant Cell Physiol* 22: 867-880
20. OSBORNE D 1982 Deoxyribonucleic and integrity and repair in seed germina-

- tion. The importance in viability and survival. In AA Khan, ed, The physiology and Biochemistry of Seed Development Dormancy and Germination. Elsevier Biomedical Press, Amsterdam, pp 435-464
21. POLLOCK BM 1969 Imbibition temperature sensitivity of lima bean seeds controlled by initial moisture. *Plant Physiol* 44: 907-911
 22. PYFROM HT, D APPLEMAN, WG HEIN 1957 Catalase and chlorophyll depression by 3-amino-1,2,4-triazole. *Plant Physiol* 32: 674-676
 23. RICH P, A BOVERIS, WD BONNER JR, AL MOORE 1976 Hydrogen peroxide generation by the alternate oxidase of higher plants. *Biochem Biophys Res Commun* 71: 695-703
 24. RICH PR, WD BONNER JR 1978 The site of superoxide anion generation in higher plant mitochondria. *Arch Biochem Biophys* 188: 206-213
 25. ROBERTS EH 1973 Oxidative processes and the control of seed germination. In W Heydecker, ed, *Seed Ecology*, Pennsylvania State University Press, Pittsburgh, pp 189-216
 26. SÁNCHEZ RA, L DE MIGUEL 1983 Ageing of *Datura ferox* seeds embryos during dry storage and its reversal imbibition. *Z Pflanzenphysiol* 110: 319-329
 27. SAUNDERS BC, AC HOLMES-SIEDLE, BP STARK 1964 Peroxidase. Butterworths, Washington, DC, pp 160-161
 28. SEITZ WR 1978 Chemiluminescence detection of enzymically generated peroxide. *Methods Enzymol* 57: 445-462
 29. SIMON EW, RM RAJA HARUM 1978 Leakage during seed imbibition *J Exp Bot* 23: 1076-1085
 30. STADELMAN EJ 1969 Permeability of the plant cell. *Annu Rev Plant Physiol* 20: 585-606
 31. TAYLORSON RB 1984 Interaction of phytochrome and other factors in seed germination. In AA Khan, ed, *The Physiology and Biochemistry of Seed Development, Dormancy and Germination*. Elsevier Biomedical Press, Amsterdam, pp 324-346
 32. TAYLORSON RB, SB HENDRICKS 1977 Dormancy in seeds. *Annu Rev Plant Physiol* 28: 331-354
 33. WARM E, GG LATIES 1982 Quantification of hydrogen peroxide in plant extracts by the chemiluminescence reaction with luminol. *Phytochemistry* 21: 827-831
 34. WOODSTOCK LW, RB TAYLORSON 1981 Ethanol and acetaldehyde in imbibing soybean seeds in relation to deterioration. *Plant Physiol* 67: 424-428