



Hyers–Ulam stability of n th order linear differential equations

Tongxing Li^{a,b,*}, Akbar Zada^c, Shah Faisal^c

^a*School of Informatics, Linyi University, Linyi, Shandong 276005, P. R. China.*

^b*LinDa Institute of Shandong Provincial Key Laboratory of Network Based Intelligent Computing, Linyi University, Linyi, Shandong 276005, P. R. China.*

^c*Department of Mathematics, University of Peshawar, Peshawar 25000, Pakistan.*

Communicated by C. Park

Abstract

For n th order linear homogeneous and nonhomogeneous differential equations with nonconstant coefficients, we prove Hyers–Ulam stability by using open mapping theorem. The generalized Hyers–Ulam stability is also investigated. ©2016 All rights reserved.

Keywords: Hyers–Ulam stability, generalized Hyers–Ulam stability, n th order linear differential equation, open mapping theorem.

2010 MSC: 35B35.

1. Introduction

The theory of stability is an important branch of the qualitative theory of differential equations. In 1940, Ulam [24] raised a problem: “when can we assert that the solutions of an inequality are close to one of the exact solutions of the corresponding equation?” Since then, this question has attracted the attention of many researchers. We refer the reader to [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 25, 26, 27, 28] and the references cited therein. Note that the first solution to this question was given by Hyers [10] in 1941. Thereafter, Aoki [2], Bourgin [3], and Rassias [21] improved the result reported in [10].

*Corresponding author

Email addresses: litongx2007@163.com (Tongxing Li), zadababo@yahoo.com (Akbar Zada), shahfaisal8763@gmail.com (Shah Faisal)

To the best of our knowledge, papers by Obłozza [17, 18] were among the first contributions dealing with Hyers–Ulam stability of differential equations. Alsina and Ger [1] proved Hyers–Ulam stability of differential equation $y'(x) = y(x)$. Later, Takahasi et al. [23] extended results of [1] to the Banach space-valued differential equation $y'(x) = \lambda y(x)$ (Hyers–Ulam stability holds true for this equation). Using direct method, iteration method, fixed point method, and open mapping theorem, Huang and Li [9] investigated the Hyers–Ulam stability of some classes of functional differential equations. In the paper by Zada et al. [26], the concepts of Hyers–Ulam stability are generalized for a class of nonautonomous linear differential systems. Very recently, Choi and Jung [6] studied the generalized Hyers–Ulam stability of the linear differential equation

$$y''(x) + f(x)y'(x) + g(x)y(x) = r(x).$$

It should be noted that research in this paper was strongly motivated by the recent contributions of Choi and Jung [6] and Huang and Li [9]. Our principal goal is to prove Hyers–Ulam stability of

$$\sum_{i=0}^n \beta_{n-i}(x)y^{(n-i)}(x) = \alpha(x) \quad (1.1)$$

and the generalized Hyers–Ulam stability of

$$y^{(n)}(x) + \sum_{i=1}^n Q_{n-i}(x)y^{(n-i)}(x) = f(x) \quad (1.2)$$

by using different approaches.

2. Preliminaries

We present the following definitions to make this paper self-dependent.

Definition 2.1 ([12, 13]). Let \mathcal{W} be a normed space and I be an open interval. Then, equation (1.1) is said to have Hyers–Ulam stability on I if, for any function $g : I \rightarrow \mathcal{W}$ satisfying

$$\left\| \sum_{i=0}^n \beta_{n-i}(x)g^{(n-i)}(x) - \alpha(x) \right\| \leq \varepsilon$$

for all $x \in I$ and for some $\varepsilon \geq 0$, there exists a function $g_0 : I \rightarrow \mathcal{W}$ such that

$$\sum_{i=0}^n \beta_{n-i}(x)g_0^{(n-i)}(x) = \alpha(x)$$

and $\|g - g_0\| \leq K(\varepsilon)$ for all $x \in I$, where $K(\varepsilon)$ is an expression for ε only.

If the above statement is also true when we replace ε and $K(\varepsilon)$ with $\phi(x)$ and $\Phi(x)$, where $\phi, \Phi : I \rightarrow [0, \infty)$ are functions not depending on g and g_0 explicitly, then we say that the corresponding differential equation has the generalized Hyers–Ulam stability (or the Hyers–Ulam–Rassias stability).

Definition 2.2 ([9]). Let $\Upsilon : A \rightarrow B$ be an operator from a space A to another space B . We say that Υ has Hyers–Ulam stability if, for any $g \in \Upsilon(A)$, $\varepsilon \geq 0$, and $f \in A$ satisfying $\|\Upsilon f - g\| \leq \varepsilon$, there exist a constant K and an $f_0 \in A$ such that $\Upsilon f_0 = g$ and $\|f - f_0\| \leq K\varepsilon$, where K is called a Hyers–Ulam stability constant of operator Υ .

In order to establish our main results, we need the following auxiliary lemmas.

Lemma 2.3 ([9, Theorem 2.4]). *Let A and B be Banach spaces and $\Upsilon : A \rightarrow B$ be a bounded linear operator. Then the following statements are equivalent:*

- (a) Υ has Hyers–Ulam stability;
- (b) The range of Υ is closed.

Lemma 2.4 ([28]). Assume that c_1, c_2, \dots, c_n are arbitrary real-valued constants. If $y_c = \sum_{i=1}^n c_i y_i$ is the complementary solution of equation (1.2) on a common interval I , where $Q_j, f \in C(I, \mathbb{R}), j = 0, 1, \dots, n - 1$ and $f(x) \neq 0$ for all $x \in I$, then its general solution $y \in C^n(I, \mathbb{R})$ is given by

$$y(x) = y_c(x) + \sum_{i=1}^n y_i(x) \int_{x_i}^x \frac{P_i(t)}{W(t)} f(t) dt,$$

where $x, x_1, x_2, \dots, x_n \in I, W$ is the Wronskian of $y_1, y_2, \dots, y_n, P_i = W_i/f$, and $W_i, i = 1, 2, \dots, n$ are the Wronskian with i th column replaced with the vector $(0, 0, 0, \dots, f(x))$, i.e.,

$$W_i = \begin{vmatrix} y_1 & y_2 & y_3 & \dots & 0 & \dots & y_n \\ y_1' & y_2' & y_3' & \dots & 0 & \dots & y_n' \\ y_1'' & y_2'' & y_3'' & \dots & 0 & \dots & y_n'' \\ \cdot & \cdot & \cdot & \dots & 0 & \dots & \cdot \\ \cdot & \cdot & \cdot & \dots & 0 & \dots & \cdot \\ \cdot & \cdot & \cdot & \dots & 0 & \dots & \cdot \\ y_1^{(n-1)} & y_2^{(n-1)} & y_3^{(n-1)} & \dots & f & \dots & y_n^{(n-1)} \end{vmatrix}.$$

Remark 2.5. $P_i, i = 1, 2, \dots, n$ in Lemma 2.4 can be written in the form

$$P_i = \begin{vmatrix} y_1 & y_2 & y_3 & \dots & 0 & \dots & y_n \\ y_1' & y_2' & y_3' & \dots & 0 & \dots & y_n' \\ y_1'' & y_2'' & y_3'' & \dots & 0 & \dots & y_n'' \\ \cdot & \cdot & \cdot & \dots & 0 & \dots & \cdot \\ \cdot & \cdot & \cdot & \dots & 0 & \dots & \cdot \\ \cdot & \cdot & \cdot & \dots & 0 & \dots & \cdot \\ y_1^{(n-1)} & y_2^{(n-1)} & y_3^{(n-1)} & \dots & 1 & \dots & y_n^{(n-1)} \end{vmatrix}.$$

3. Hyers–Ulam stability of equation (1.1)

In what follows, $C[a, b]$ denotes the set of all continuous functions on $[a, b]$ and $C^n[a, b]$ stands for the set containing all n times continuously differentiable functions on $[a, b]$.

Theorem 3.1. If $\beta_i \in C[a, b]$ for $i = 0, 1, \dots, n$, then the homogeneous n th order linear differential equation

$$\sum_{i=0}^n \beta_{n-i}(x) y^{(n-i)}(x) = 0 \tag{3.1}$$

has Hyers–Ulam stability on $[a, b]$.

Proof. For every $y \in C^n[a, b]$, define a norm in $C^n[a, b]$ by

$$\|y\| := \max \left(\max_{1 \leq i \leq n} \sup_{x \in [a, b]} |y^{(i)}(x)|, \sup_{x \in [a, b]} |y(x)| \right).$$

It is not difficult to verify that $C[a, b]$ and $(C^n[a, b], \|\cdot\|)$ are Banach spaces (see Huang et al. [8] for details). For all $g \in C^n[a, b]$, define now an operator $\Upsilon : C^n[a, b] \rightarrow C[a, b]$ by

$$(\Upsilon g)(x) := \sum_{i=0}^n \beta_{n-i}(x) g^{(n-i)}(x).$$

Clearly, Υ is well-defined and is a linear operator. By virtue of

$$\begin{aligned} \|\Upsilon\| &= \sup_{\|g\|=1} \|\Upsilon g\| \\ &= \sup_{\|g\|=1} \sup_{x \in [a,b]} \left| \sum_{i=0}^n \beta_{n-i}(x) g^{(n-i)}(x) \right| \\ &\leq \sup_{\|g\|=1} \sup_{x \in [a,b]} \sum_{i=0}^n \left| \beta_{n-i}(x) g^{(n-i)}(x) \right| \\ &= \sup_{\|g\|=1} \sup_{x \in [a,b]} \sum_{i=0}^n |\beta_{n-i}(x)| \left| g^{(n-i)}(x) \right| \\ &\leq \sup_{x \in [a,b]} \sum_{i=0}^n |\beta_{n-i}(x)| \end{aligned}$$

and the facts that β_i are continuous on $[a, b]$ for $i = 0, 1, \dots, n$, we deduce that Υ is bounded. Next, we prove that the range of Υ is closed. Obviously, $\Upsilon g \in C[a, b]$ for every $g \in C^n[a, b]$. Conversely, using the basic theory of differential equations (see [27]), we conclude that, for every $y \in C[a, b]$, there are some $f \in C^n[a, b]$ such that $\Upsilon f = y$. Moreover, $C[a, b]$ is a Banach space, and so the range of Υ is closed. It follows now from Lemma 2.3 that Υ has Hyers–Ulam stability. Noticing that $0 \in \Upsilon(C^n[a, b])$ and using Definition 2.2, we obtain that, for any $\varepsilon \geq 0$ and $f \in C^n[a, b]$ satisfying

$$\left\| \sum_{i=0}^n \beta_{n-i}(x) f^{(n-i)}(x) \right\| \leq \varepsilon$$

for all $x \in [a, b]$, there exist a constant K and a solution $f_0 \in C^n[a, b]$ to (3.1) with the property

$$\|f - f_0\| \leq K\varepsilon.$$

By virtue of Definition 2.1, equation (3.1) has Hyers–Ulam stability on $[a, b]$. The proof is complete. \square

Similar as in the proof of Theorem 3.1, one can obtain the following result.

Corollary 3.2. *If $\beta_i, \alpha \in C[a, b]$ for $i = 0, 1, \dots, n$, then the n th order nonhomogeneous linear differential equation (1.1) has Hyers–Ulam stability on $[a, b]$.*

4. The generalized Hyers–Ulam stability of equation (1.2)

In this section, we prove the generalized Hyers–Ulam stability of an n th order nonhomogeneous linear differential equation (1.2) on a non-degenerate interval I .

Theorem 4.1. *Assume that $Q_j, f \in C(I, \mathbb{R})$, $j = 0, 1, \dots, n - 1$ and $f(x) \neq 0$ for all $x \in I$, and let $y_c = \sum_{i=1}^n c_i y_i$ be the complementary solution of equation (1.2) on I , where c_i are arbitrary real-valued constants. If a function $y \in C^n(I, \mathbb{R})$ satisfies the inequality*

$$\left| y^{(n)}(x) + \sum_{i=1}^n Q_{n-i}(x) y^{(n-i)}(x) - f(x) \right| \leq \phi(x) \tag{4.1}$$

for all $x \in I$, where $\phi : I \rightarrow [0, \infty)$ is given such that the following integrals exist, then there is a solution $y_0 \in C^n(I, \mathbb{R})$ of (1.2) satisfying

$$|y(x) - y_0(x)| \leq \sum_{i=1}^n |y_i(x)| \int_{x_i}^x \left| \frac{P_i(t)}{W(t)} \right| \phi(t) dt,$$

where $x, x_1, x_2, \dots, x_n \in I$, W, W_i , and P_i are defined as in Lemma 2.4.

Proof. Let $y \in C^n(I, \mathbb{R})$ be such that (4.1) holds. Define a function $r \in C(I, \mathbb{R})$ by

$$r(x) := y^{(n)}(x) + \sum_{i=1}^n Q_{n-i}(x)y^{(n-i)}(x). \quad (4.2)$$

It follows from (4.1) that, for all $x \in I$,

$$|r(x) - f(x)| \leq \phi(x). \quad (4.3)$$

By virtue of Lemma 2.4, the general solution of (4.2) is given by

$$y(x) = y_c(x) + \sum_{i=1}^n y_i(x) \int_{x_i}^x \frac{P_i(t)}{W(t)} r(t) dt,$$

where $W \neq 0$ for all $t \in I$ due to the fact that y_1, y_2, \dots, y_n are linearly independent on I . For every $x \in I$, define a new function $y_0 : I \rightarrow \mathbb{R}$ by

$$y_0(x) := y_c(x) + \sum_{i=1}^n y_i(x) \int_{x_i}^x \frac{P_i(t)}{W(t)} f(t) dt.$$

It follows now from Lemma 2.4 that y_0 is the solution of (1.2). An application of (4.3) implies that

$$\begin{aligned} |y(x) - y_0(x)| &= \left| \sum_{i=1}^n y_i(x) \int_{x_i}^x \frac{P_i(t)}{W(t)} r(t) dt - \sum_{i=1}^n y_i(x) \int_{x_i}^x \frac{P_i(t)}{W(t)} f(t) dt \right| \\ &\leq \sum_{i=1}^n |y_i(x)| \int_{x_i}^x \left| \frac{P_i(t)}{W(t)} \right| \phi(t) dt, \end{aligned}$$

which completes the proof. \square

Remark 4.2. Theorem 4.1 reduces to [6, Theorem 3.2] in the case where $n = 2$.

Acknowledgements

The authors express their sincere gratitude to the editors and anonymous referees for the careful reading of the original manuscript and useful comments that helped to improve the presentation of the results and accentuate important details. This research is supported by NNSF of P. R. China (Grant No. 61503171), CPSF (Grant No. 2015M582091), NSF of Shandong Province (Grant No. ZR2012FL06), DSRF of Linyi University (Grant No. LYDX2015BS001), and the AMEP of Linyi University, P. R. China.

References

- [1] C. Alsina, R. Ger, *On some inequalities and stability results related to the exponential function*, J. Inequal. Appl., **2** (1998), 373–380.1
- [2] T. Aoki, *On the stability of the linear transformation in Banach spaces*, J. Math. Soc. Japan, **2** (1950), 64–66.1
- [3] D. G. Bourgin, *Classes of transformations and bordering transformations*, Bull. Amer. Math. Soc., **57** (1951), 223–237.1
- [4] J. Brzdęk, D. Popa, B. Xu, *Remarks on stability of linear recurrence of higher order*, Appl. Math. Lett., **23** (2010), 1459–1463.1
- [5] M. Burger, N. Ozawa, A. Thom, *On Ulam stability*, Israel J. Math., **193** (2013), 109–129.1
- [6] G. Choi, S.-M. Jung, *Invariance of Hyers–Ulam stability of linear differential equations and its applications*, Adv. Difference Equ., **2015** (2015), 14 pages.1, 4.2
- [7] J. Chung, *Hyers–Ulam stability theorems for Pexider equations in the space of Schwartz distributions*, Arch. Math., **84** (2005), 527–537.1

- [8] J. Huang, S.-M. Jung, Y. Li, *On Hyers–Ulam stability of nonlinear differential equations*, Bull. Korean Math. Soc., **52** (2015), 685–697. 1, 3
- [9] J. Huang, Y. Li, *Hyers–Ulam stability of linear functional differential equations*, J. Math. Anal. Appl., **426** (2015), 1192–1200. 1, 2.2, 2.3
- [10] D. H. Hyers, *On the stability of the linear functional equation*, Proc. Nat. Acad. Sci. USA, **27** (1941), 222–224. 1
- [11] S.-M. Jung, *Hyers–Ulam stability of linear differential equations of first order*, Appl. Math. Lett., **17** (2004), 1135–1140. 1
- [12] S.-M. Jung, *Hyers–Ulam stability of linear differential equations of first order, III*, J. Math. Anal. Appl., **311** (2005), 139–146. 1, 2.1
- [13] S.-M. Jung, *Hyers–Ulam stability of linear differential equations of first order, II*, Appl. Math. Lett., **19** (2006), 854–858. 1, 2.1
- [14] Y. Li, Y. Shen, *Hyers–Ulam stability of nonhomogeneous linear differential equations of second order*, Int. J. Math. Math. Sci., **2009** (2009), 7 pages. 1
- [15] Y. Li, Y. Shen, *Hyers–Ulam stability of linear differential equations of second order*, Appl. Math. Lett., **23** (2010), 306–309. 1
- [16] T. Miura, S. Miyajima, S.-E. Takahasi, *A characterization of Hyers–Ulam stability of first order linear differential operators*, J. Math. Anal. Appl., **286** (2003), 136–146. 1
- [17] M. Obloza, *Hyers stability of the linear differential equation*, Rocznik Nauk.-Dydakt. Prace Mat., **13** (1993), 259–270. 1
- [18] M. Obloza, *Connections between Hyers and Lyapunov stability of the ordinary differential equations*, Rocznik Nauk.-Dydakt. Prace Mat., **14** (1997), 141–146. 1
- [19] D. Popa, *Hyers–Ulam–Rassias stability of a linear recurrence*, J. Math. Anal. Appl., **309** (2005), 591–597. 1
- [20] D. Popa, I. Raşa, *Hyers–Ulam stability of the linear differential operator with nonconstant coefficients*, Appl. Math. Comput., **219** (2012), 1562–1568. 1
- [21] Th. M. Rassias, *On the stability of the linear mapping in Banach spaces*, Proc. Amer. Math. Soc., **72** (1978), 297–300. 1
- [22] H. Rezaei, S.-M. Jung, Th. M. Rassias, *Laplace transform and Hyers–Ulam stability of linear differential equations*, J. Math. Anal. Appl., **403** (2013), 244–251. 1
- [23] S.-E. Takahasi, T. Miura, S. Miyajima, *On the Hyers–Ulam stability of the Banach space-valued differential equation $y' = \lambda y$* , Bull. Korean Math. Soc., **39** (2002), 309–315. 1
- [24] S. M. Ulam, *A Collection of Mathematical Problems*, Interscience Publishers, New York, (1960). 1
- [25] G. Wang, M. Zhou, L. Sun, *Hyers–Ulam stability of linear differential equations of first order*, Appl. Math. Lett., **21** (2008), 1024–1028. 1
- [26] A. Zada, O. Shah, R. Shah, *Hyers–Ulam stability of non-autonomous systems in terms of boundedness of Cauchy problems*, Appl. Math. Comput., **271** (2015), 512–518. 1
- [27] Z. Zheng, *Theory of Functional Differential Equations*, Anhui Education Press, (1994). (in Chinese) 1, 3
- [28] D. G. Zill, *A First Course in Differential Equations with Modeling Applications*, Brooks/Cole, USA, (2012). 1, 2.4