

# Hyper Relative Order $(p, q)$ of Entire Functions

Dibyendu Banerjee and Saikat Batabyal

**Abstract.** After the works of Lahiri and Banerjee [6] on the idea of relative order  $(p, q)$  of entire functions, we introduce in this paper hyper relative order  $(p, q)$  of entire functions where  $p, q$  are positive integers with  $p > q$  and prove sum theorem, product theorem and theorem on derivative.

**AMS Subject Classification (2010).** 30D20

**Keywords.** Entire functions, Hyper relative order  $(p, q)$ , Property(A), Composition, Order (Lower order)

## 1 Introduction and Definitions

Let  $f$  and  $g$  be non-constant entire functions and  $M_f(r) = \max\{|f(z)| : |z| = r\}$ ,  $M_g(r) = \max\{|g(z)| : |z| = r\}$ . Then  $M_f(r)$  is strictly increasing and continuous function of  $r$  and its inverse  $M_f^{-1} : (|f(0)|, \infty) \rightarrow (0, \infty)$  exists and  $\lim_{R \rightarrow \infty} M_f^{-1}(R) = \infty$ .

In 1988, Bernal [2] introduced the definition of relative order of  $f$  with respect to  $g$  as

$$\rho_g(f) = \inf\{\mu > 0 : M_f(r) < M_g(r^\mu)\}$$

for all  $r > r_0(\mu) > 0$ .

When  $g(z) = \exp(z)$ ,  $\rho_g(f)$  coincides with the classical definition of order ([15], p-248).

Following Sato [14], we write  $\log^{[0]} x = x$ ,  $\exp^{[0]} x = x$  and for positive integer  $m \geq 1$ ,  $\log^{[m]} x = \log(\log^{[m-1]} x)$ ,  $\exp^{[m]} x = \exp(\exp^{[m-1]} x)$ .

If  $p, q$  are positive integers,  $p \geq q$  then Juneja et.al., [7] defined  $(p, q)$ th order of  $f$  by

$$\rho^{(p,q)}(f) = \limsup_{r \rightarrow \infty} \frac{\log^{[p]} M_f(r)}{\log^{[q]} r}.$$

During the past decades, several authors (see for example [1], [7], [8], [9], [10], [11]) made close investigations on  $(p, q)$  order of entire functions. After this in 2005, Lahiri and Banerjee [6] introduced the concept of relative order  $(p, q)$  of entire functions as follows.

**Definition 1.1.** [6] *Let  $p$  and  $q$  be positive integers with  $p > q$ . The relative order  $(p, q)$  of  $f$  with respect to  $g$  is defined by*

$$\rho_g^{(p,q)}(f) = \inf\{\mu > 0 : M_f(r) < M_g(\exp^{[p-1]}(\mu \log^{[q]} r)) \text{ for all } r > r_0(\mu) > 0\}$$

$$= \limsup_{r \rightarrow \infty} \frac{\log^{[p-1]} M_g^{-1} M_f(r)}{\log^{[q]} r} = \limsup_{r \rightarrow \infty} \frac{\log^{[p-1]} M_g^{-1}(r)}{\log^{[q]} M_f^{-1}(r)}.$$

If  $g(z) = \exp(z)$  then  $\rho_g^{(p,q)}(f) = \rho^{(p,q)}(f)$ .

In the present paper we introduce the concept of hyper relative  $(p, q)$  order as follows.

**Definition 1.2.** *Let  $f$  and  $g$  be entire functions and  $p, q$  are positive integers with  $p > q$ . The hyper relative  $(p, q)$  order of  $f$  with respect to  $g$  is defined by*

$$\rho_g^{-(p,q)}(f) = \inf\{\mu > 0 : M_f(r) < M_g(\exp^{[p]}(\mu \log^{[q]} r)) \text{ for all } r > r_0(\mu) > 0\}.$$

$$\text{Clearly } \rho_g^{-(p,q)}(f) = \limsup_{r \rightarrow \infty} \frac{\log^{[p]} M_g^{-1} M_f(r)}{\log^{[q]} r}.$$

When  $p = 2, q = 1$  and  $g(z) = \exp(z)$ , then the definition coincides with the classical definition of hyper order of entire functions which have been investigated closely by several authors (see for example [4], [13] etc.).

The following definition of Bernal [2] will be needed.

**Definition 1.3.** [2] *A non-constant entire function  $g$  is said to have the property (A) if for any  $\sigma > 1$  and for all large  $r$ ,  $[M_g(r)]^2 \leq M_g(r^\sigma)$  holds.*

Examples of functions with or without the property (A) are available in [2]. Throughout the paper we shall assume  $f, g, h$  etc., are non-constant entire functions and  $M_f(r), M_g(r), M_h(r)$  etc., denote respectively their maximum modulus on  $|z| = r$ .

## 2 Lemmas

The following lemmas will be needed in the sequel.

**Lemma 2.1.** [2] *Let  $g$  be an entire function which satisfies the property (A), and let  $\sigma > 1$ . Then for any positive integer  $n$  and for all large  $r$ ,*

$$[M_g(r)]^n \leq M_g(r^\sigma)$$

*holds.*

**Lemma 2.2.** [2] *Suppose  $f$  is an entire function,  $\alpha > 1, 0 < \beta < \alpha, s > 1, 0 < \mu < \lambda$  and  $n$  is a positive integer. Then*

(a)  $M_f(\alpha r) > \beta M_f(r)$ .

(b) *There exists  $K = K(s, f) > 0$  such that  $[M_f(r)]^s \leq K M_f(r^s)$  for  $r > 0$ .*

(c)  $\lim_{r \rightarrow \infty} \frac{M_f(r^s)}{M_f(r)} = \infty = \lim_{r \rightarrow \infty} \frac{M_f(r^\lambda)}{M_f(r^\mu)}$ .

(d) *If  $f$  is transcendental, then*

$$\lim_{r \rightarrow \infty} \frac{M_f(r^s)}{r^n M_f(r)} = \infty = \lim_{r \rightarrow \infty} \frac{M_f(r^\lambda)}{r^n M_f(r^\mu)}$$

**Lemma 2.3.** ([12], p-21) *Let  $f(z)$  be holomorphic in the circle  $|z| = 2eR (R > 0)$  with  $f(0) = 1$  and  $\eta$  be an arbitrary positive number not exceeding  $\frac{3e}{2}$ . Then inside the circle  $|z| = R$ , but outside of a family of excluded circles the sum of whose radii is not greater than  $4\eta R$ , we have*

$$\log |f(z)| > -T(\eta) \log M_f(2eR)$$

for  $T(\eta) = 2 + \log \frac{3e}{2\eta}$ .

**Lemma 2.4.** [5] *Every entire function  $g$  satisfying the property (A) is transcendental.*

**Lemma 2.5.** [3] *Let  $f(z)$  and  $g(z)$  be entire functions with  $g(0) = 0$ . Let  $\alpha$  satisfy  $0 < \alpha < 1$  and let  $C(\alpha) = \frac{(1-\alpha)^2}{4\alpha}$ . Then for  $r > 0$*

$$M_{f \circ g}(r) \geq M_f(C(\alpha)M_g(\alpha r)).$$

*Further if  $g(z)$  is any entire function, then with  $\alpha = \frac{1}{2}$ , for sufficiently large values of  $r$*

$$M_{f \circ g}(r) \geq M_f\left(\frac{1}{8}M_g\left(\frac{r}{2}\right) - |g(0)|\right).$$

Clearly

$$M_{f \circ g}(r) \geq M_f\left(\frac{1}{16}M_g\left(\frac{r}{2}\right)\right). \quad (1)$$

On the other hand the opposite inequality

$$M_{f \circ g}(r) \leq M_f(M_g(r)) \quad (2)$$

is an immediate consequence of the definition.

**Lemma 2.6.** *If  $f$  is a polynomial of degree  $n$  and  $g$  is transcendental, then  $\rho_g^{-(p,q)}(f) = 0$ .*

*Proof.* For all large  $r$ ,  $M_f(r) \leq Nr^n$  where  $N (> 0)$  is a constant and  $M_g(r) > Kr^m$  where  $K (> 0)$  is a constant and  $m (> 0)$  is arbitrary.

Then

$$M_g^{-1}M_f(r) < \left(\frac{Nr^n}{K}\right)^{\frac{1}{m}} = \lambda r^{\frac{n}{m}},$$

where  $\lambda = \left(\frac{N}{K}\right)^{\frac{1}{m}}$ .

Now

$$\begin{aligned} \rho_g^{-(p,q)}(f) &= \limsup_{r \rightarrow \infty} \frac{\log^{[p]} M_g^{-1}M_f(r)}{\log^{[q]} r} \\ &\leq \limsup_{r \rightarrow \infty} \frac{\log^{[p]} (\lambda r^{\frac{n}{m}})}{\log^{[q]} r} = 0. \end{aligned}$$

□

### 3 Sum Theorem

**Theorem 3.1.** *If  $f_1, f_2, g$  and  $h$  are entire functions with  $0 < \lambda_h \leq \rho_h < \infty$ , then for  $p > 2$*

$$\rho_g^{-(p,q)}(f_1 \pm f_2) \leq \max\{\rho_{g \circ h}^{(p,q)}(f_1), \rho_{g \circ h}^{(p,q)}(f_2)\},$$

*the equality holding when  $\rho_{g \circ h}^{(p,q)}(f_1) \neq \rho_{g \circ h}^{(p,q)}(f_2)$ .*

*Proof.* We consider the theorem for  $f_1 + f_2$ . Let  $f = f_1 + f_2$  and suppose that

$$\rho_{goh}^{(p,q)}(f_1) \leq \rho_{goh}^{(p,q)}(f_2).$$

Let  $\varepsilon > 0$  be arbitrary. For all large  $r$ , we have

$$\begin{aligned} M_{f_1}(r) &< M_{goh} \left[ \exp^{[p-1]} \{ (\rho_{goh}^{(p,q)}(f_1) + \varepsilon) \log^{[q]} r \} \right] \\ &\leq M_{goh} \left[ \exp^{[p-1]} \{ (\rho_{goh}^{(p,q)}(f_2) + \varepsilon) \log^{[q]} r \} \right] \end{aligned}$$

$$\text{and } M_{f_2}(r) < M_{goh} \left[ \exp^{[p-1]} \{ (\rho_{goh}^{(p,q)}(f_2) + \varepsilon) \log^{[q]} r \} \right].$$

Now,

$$M_f(r) \leq M_{f_1}(r) + M_{f_2}(r)$$

$$< 2M_{goh} \left[ \exp^{[p-1]} \{ (\rho_{goh}^{(p,q)}(f_2) + \varepsilon) \log^{[q]} r \} \right]$$

$$< M_{goh} \left[ 3 \exp^{[p-1]} \{ (\rho_{goh}^{(p,q)}(f_2) + \varepsilon) \log^{[q]} r \} \right]$$

using Lemma 2.2(a) it follows by (2)

$$M_g \left[ M_h \{ 3 \exp^{[p-1]} \{ (\rho_{goh}^{(p,q)}(f_2) + \varepsilon) \log^{[q]} r \} \} \right].$$

So,

$$M_g^{-1} M_f(r) < M_h \left[ 3 \exp^{[p-1]} \{ (\rho_{goh}^{(p,q)}(f_2) + \varepsilon) \log^{[q]} r \} \right]$$

i.e.

$$\log M_g^{-1} M_f(r) < \log M_h \left[ 3 \exp^{[p-1]} \{ (\rho_{goh}^{(p,q)}(f_2) + \varepsilon) \log^{[q]} r \} \right]$$

$$< \left[ 3 \exp^{[p-1]} \{ (\rho_{goh}^{(p,q)}(f_2) + \varepsilon) \log^{[q]} r \} \right]^{(\rho_h + \varepsilon)}$$

i.e.

$$\log \log M_g^{-1} M_f(r) < (\rho_h + \varepsilon) \left[ \log 3 + \exp^{[p-2]} \{ (\rho_{goh}^{(p,q)}(f_2) + \varepsilon) \log^{[q]} r \} \right]$$

i.e.

$$\log^{[p]} M_g^{-1} M_f(r) < (\rho_{goh}^{(p,q)}(f_2) + \varepsilon) \log^{[q]} r + O(1)$$

i.e.

$$\rho_g^{-(p,q)}(f) \leq (\rho_{goh}^{(p,q)}(f_2) + \varepsilon).$$

Since  $\varepsilon > 0$  be arbitrary,

$$\bar{\rho}_g^{(p,q)}(f_1 + f_2) \leq \rho_{goh}^{(p,q)}(f_2) = \max\{\rho_{goh}^{(p,q)}(f_1), \rho_{goh}^{(p,q)}(f_2)\}. \quad (3)$$

Next let,  $\rho_{goh}^{(p,q)}(f_1) < \rho_{goh}^{(p,q)}(f_2)$ . Then for all large  $r$

$$M_{f_1}(r) < M_{goh} \left[ \exp^{[p-1]} \{(\rho_{goh}^{(p,q)}(f_1) + \varepsilon) \log^{[q]} r\} \right] \quad (4)$$

and there exists a sequence  $\{r_n\}$ ,  $r_n \rightarrow \infty$  such that

$$M_{f_2}(r_n) > M_{goh} \left[ \exp^{[p-1]} \{(\rho_{goh}^{(p,q)}(f_2) - \varepsilon) \log^{[q]} r_n\} \right] \text{ for } n = 1, 2, 3, \dots \quad (5)$$

From Lemma 2.2(c), we obtain

$$\lim_{r \rightarrow \infty} \frac{M_{goh} \left[ \exp^{[p-1]} \{(\rho_{goh}^{(p,q)}(f_2) - \varepsilon) \log^{[q]} r\} \right]}{M_{goh} \left[ \exp^{[p-1]} \{(\rho_{goh}^{(p,q)}(f_1) + \varepsilon) \log^{[q]} r\} \right]} = \infty.$$

Then for all large  $r$ ,

$$\begin{aligned} & M_{goh} \left[ \exp^{[p-1]} \{(\rho_{goh}^{(p,q)}(f_2) - \varepsilon) \log^{[q]} r\} \right] \\ & > 2M_{goh} \left[ \exp^{[p-1]} \{(\rho_{goh}^{(p,q)}(f_1) + \varepsilon) \log^{[q]} r\} \right]. \end{aligned}$$

So,

$$\begin{aligned} M_{f_2}(r_n) & > M_{goh} \left[ \exp^{[p-1]} \{(\rho_{goh}^{(p,q)}(f_2) - \varepsilon) \log^{[q]} r_n\} \right] \\ & > 2 M_{goh} \left[ \exp^{[p-1]} \{(\rho_{goh}^{(p,q)}(f_1) + \varepsilon) \log^{[q]} r_n\} \right]. \end{aligned}$$

Now for a sequence of values of  $\{r_n\}$ ,  $r_n \rightarrow \infty$  we get by using (4)

$$M_{f_2}(r_n) > 2M_{f_1}(r_n) \text{ for } n = 1, 2, 3, \dots$$

So,

$$\begin{aligned} M_f(r_n) & \geq M_{f_2}(r_n) - M_{f_1}(r_n) > M_{f_2}(r_n) - \frac{1}{2}M_{f_2}(r_n) = \frac{1}{2}M_{f_2}(r_n) \\ & > \frac{1}{2}M_{goh} \left[ \exp^{[p-1]} \{(\rho_{goh}^{(p,q)}(f_2) - \varepsilon) \log^{[q]} r_n\} \right], \end{aligned}$$

using (5)

$$> M_{goh} \left[ \frac{1}{3} \exp^{[p-1]} \{ (\rho_{goh}^{(p,q)}(f_2) - \varepsilon) \log^{[q]} r_n \} \right]$$

using Lemma 2.2(a)

$$\geq M_g \left[ \frac{1}{16} M_h \left( \frac{\exp^{[p-1]} \{ (\rho_{goh}^{(p,q)}(f_2) - \varepsilon) \log^{[q]} r_n \}}{6} \right) \right],$$

using (1)

i.e.,

$$M_g^{-1} M_f(r_n) > \frac{1}{16} M_h \left[ \frac{\exp^{[p-1]} \{ (\rho_{goh}^{(p,q)}(f_2) - \varepsilon) \log^{[q]} r_n \}}{6} \right]$$

i.e.,

$$\log M_g^{-1} M_f(r_n) \geq \log M_h \left[ \frac{\exp^{[p-1]} \{ (\rho_{goh}^{(p,q)}(f_2) - \varepsilon) \log^{[q]} r_n \}}{6} \right] + O(1)$$

$$> \left[ \frac{\exp^{[p-1]} \{ (\rho_{goh}^{(p,q)}(f_2) - \varepsilon) \log^{[q]} r_n \}}{6} \right]^{(\lambda_h - \varepsilon)} + O(1)$$

i.e.

$$\log \log M_g^{-1} M_f(r_n) > (\lambda_h - \varepsilon) \left[ \exp^{[p-2]} \{ (\rho_{goh}^{(p,q)}(f_2) - \varepsilon) \log^{[q]} r_n \} \right] + O(1).$$

So,

$$\log^{[p]} M_g^{-1} M_f(r_n) > (\rho_{goh}^{(p,q)}(f_2) - \varepsilon) \log^{[q]} r_n + O(1)$$

i.e.,

$$\bar{\rho}_g^{-(p,q)}(f) \geq (\rho_{goh}^{(p,q)}(f_2) - \varepsilon).$$

Since  $\varepsilon > 0$  is arbitrary,

$$\bar{\rho}_g^{-(p,q)}(f_1 + f_2) \geq \rho_{goh}^{(p,q)}(f_2) = \max\{\rho_{goh}^{(p,q)}(f_1), \rho_{goh}^{(p,q)}(f_2)\}. \tag{6}$$

From (3) and (6) we have

$$\bar{\rho}_g^{-(p,q)}(f_1 + f_2) = \max\{\rho_{goh}^{(p,q)}(f_1), \rho_{goh}^{(p,q)}(f_2)\}.$$

This proves the theorem. □

## 4 Product Theorems

**Theorem 4.1.** *Let  $P$  be a polynomial and  $f, g, h$  are entire functions with  $0 < \lambda_h \leq \rho_h < \infty$ , where  $f$  is transcendental. Then for  $p > 2$*

$$\bar{\rho}_g^{-(p,q)}(Pf) = \rho_{g \circ h}^{(p,q)}(f).$$

*Proof.* Let the degree of  $P(z)$  be  $m$ . Then there exists  $\alpha$ ,  $0 < \alpha < 1$  and a positive integer  $n$  ( $> m$ ) such that  $2\alpha < |P(z)| < r^n$  holds on  $|z| = r$ , for all large  $r$ . Now by Lemma 2.2(a)

$$M_f\left(\frac{1}{\alpha}\alpha r\right) > \frac{1}{2\alpha}M_f(\alpha r)$$

i.e.,  $M_f(\alpha r) < 2\alpha M_f(r)$ .

Let  $k(z) = P(z)f(z)$ . Then for all large  $r$  and  $s > 1$

$M_f(\alpha r) < 2\alpha M_f(r) \leq M_k(r) \leq r^n M_f(r) < M_f(r^s)$ , by Lemma 2.2(d).

Let  $\varepsilon > 0$  be arbitrary. Now for all large  $r$ ,

$$M_k(r) < M_f(r^s)$$

$$< M_{g \circ h} \left[ \exp^{[p-1]} \{ (\rho_{g \circ h}^{(p,q)}(f) + \varepsilon) \log^{[q]} r^s \} \right]$$

$$< M_g \left[ M_h \{ \exp^{[p-1]} \{ (\rho_{g \circ h}^{(p,q)}(f) + \varepsilon) \log^{[q]} r^s \} \} \right] \text{ by (2) .}$$

$$\text{So, } M_g^{-1}M_k(r) < M_h \left[ \exp^{[p-1]} \{ (\rho_{g \circ h}^{(p,q)}(f) + \varepsilon) \log^{[q]} r^s \} \right]$$

$$\text{i.e., } \log M_g^{-1}M_k(r) < \log M_h \left[ \exp^{[p-1]} \{ (\rho_{g \circ h}^{(p,q)}(f) + \varepsilon) \log^{[q]} r^s \} \right]$$

$$< \left[ \exp^{[p-1]} \{ (\rho_{g \circ h}^{(p,q)}(f) + \varepsilon) \log^{[q]} r^s \} \right]^{(\rho_h + \varepsilon)}$$

$$\text{i.e., } \log \log M_g^{-1}M_k(r) < (\rho_h + \varepsilon) \exp^{[p-2]} \{ (\rho_{g \circ h}^{(p,q)}(f) + \varepsilon) \log^{[q]} r^s \}$$

$$\text{i.e., } \log^{[p]} M_g^{-1}M_k(r) < (\rho_{g \circ h}^{(p,q)}(f) + \varepsilon) \log^{[q]} r^s + O(1).$$

$$\text{So, } \bar{\rho}_g^{-(p,q)}(k) \leq (\rho_{g \circ h}^{(p,q)}(f) + \varepsilon).$$

Since  $\varepsilon > 0$  be arbitrary,

$$\bar{\rho}_g^{-(p,q)}(k) \leq \rho_{g \circ h}^{(p,q)}(f). \quad (7)$$

On the other hand for a sequence of values of  $\{r_n\}$ ,  $r_n \rightarrow \infty$



$$M_k(r_n) > M_f(\alpha r_n) > M_{g \circ h} \left[ \exp^{[p-1]} \{ (\rho_{g \circ h}^{(p,q)}(f) - \varepsilon) \log^{[q]}(\alpha r_n) \} \right]$$

$$> M_g \left[ \frac{1}{16} M_h \left\{ \frac{\exp^{[p-1]} \{ (\rho_{g \circ h}^{(p,q)}(f) - \varepsilon) \log^{[q]}(\alpha r_n) \}}{2} \right\} \right] \text{ by (1).}$$

So,  $M_g^{-1} M_k(r_n) > \frac{1}{16} M_h \left[ \frac{\exp^{[p-1]} \{ (\rho_{g \circ h}^{(p,q)}(f) - \varepsilon) \log^{[q]}(\alpha r_n) \}}{2} \right]$

i.e,

$$\log M_g^{-1} M_k(r_n) > \log M_h \left[ \frac{\exp^{[p-1]} \{ (\rho_{g \circ h}^{(p,q)}(f) - \varepsilon) \log^{[q]}(\alpha r_n) \}}{2} \right] + O(1)$$

$$> \left[ \frac{\exp^{[p-1]} \{ (\rho_{g \circ h}^{(p,q)}(f) - \varepsilon) \log^{[q]}(\alpha r_n) \}}{2} \right]^{(\lambda_h - \varepsilon)} + O(1)$$

i.e,  $\log \log M_g^{-1} M_k(r_n) > (\lambda_h - \varepsilon) \left[ \exp^{[p-2]} \{ (\rho_{g \circ h}^{(p,q)}(f) - \varepsilon) \log^{[q]}(\alpha r_n) \} \right] + O(1)$

i.e,  $\log^{[p]} M_g^{-1} M_k(r_n) > (\rho_{g \circ h}^{(p,q)}(f) - \varepsilon) \log^{[q]}(\alpha r_n) + O(1)$

i.e.,  $\rho_g^{-(p,q)}(k) \geq (\rho_{g \circ h}^{(p,q)}(f) - \varepsilon)$ .

Since  $\varepsilon > 0$  is arbitrary,

$$\rho_g^{-(p,q)}(k) \geq \rho_{g \circ h}^{(p,q)}(f). \tag{8}$$

From (7) and (8) we have

$$\rho_g^{-(p,q)}(k) = \rho_{g \circ h}^{(p,q)}(f)$$

i.e.,  $\rho_g^{-(p,q)}(Pf) = \rho_{g \circ h}^{(p,q)}(f)$ . □

**Theorem 4.2.** *If  $n > 1$  be a positive integer and  $f, g$  and  $h$  are entire functions with  $0 < \lambda_h \leq \rho_h < \infty$ , then for  $p > 2$*

$$\rho_g^{-(p,q)}(f^n) = \rho_{g \circ h}^{(p,q)}(f).$$

*Proof.* From Lemmas 2.2(a) and 2.2(b), we obtain

$$\begin{aligned} & [M_f(r)]^n \leq KM_f(r^n) \\ & < M_f[(K+1)r^n] < M_{g \circ h} \left[ \exp^{[p-1]} \{ (\rho_{g \circ h}^{(p,q)}(f) + \varepsilon) \log^{[q]}((K+1)r^n) \} \right] \\ & < M_g \left[ M_h \{ \exp^{[p-1]} \{ (\rho_{g \circ h}^{(p,q)}(f) + \varepsilon) \log^{[q]}((K+1)r^n) \} \} \right], \text{ by(2)} \end{aligned}$$

where  $K = K(n, f) > 0, n > 1, r > 1$ .

So,

$$M_g^{-1} [M_f(r)]^n < M_h \left[ \exp^{[p-1]} \{ (\rho_{g \circ h}^{(p,q)}(f) + \varepsilon) \log^{[q]}((K+1)r^n) \} \right]$$

i.e.,

$$\log M_g^{-1} [M_f(r)]^n < \log M_h \left[ \exp^{[p-1]} \{ (\rho_{g \circ h}^{(p,q)}(f) + \varepsilon) \log^{[q]}((K+1)r^n) \} \right]$$

$$< \left[ \exp^{[p-1]} \{ (\rho_{g \circ h}^{(p,q)}(f) + \varepsilon) \log^{[q]}((K+1)r^n) \} \right]^{(\rho_h + \varepsilon)}$$

$$\text{i.e., } \log \log M_g^{-1} [M_f(r)]^n < (\rho_h + \varepsilon) \left[ \exp^{[p-2]} \{ (\rho_{g \circ h}^{(p,q)}(f) + \varepsilon) \log^{[q]}((K+1)r^n) \} \right]$$

$$\text{i.e., } \log^{[p]} M_g^{-1} [M_f(r)]^n < (\rho_{g \circ h}^{(p,q)}(f) + \varepsilon) \log^{[q]}((K+1)r^n) + O(1)$$

$$\text{i.e., } \limsup_{r \rightarrow \infty} \frac{\log^{[p]} M_g^{-1} [M_f(r)]^n}{\log^{[q]} r}$$

$$\leq \limsup_{r \rightarrow \infty} \frac{(\rho_{g \circ h}^{(p,q)}(f) + \varepsilon) \log^{[q]}((K+1)r^n)}{\log^{[q]}((K+1)r^n)} \limsup_{r \rightarrow \infty} \frac{\log^{[q]}((K+1)r^n)}{\log^{[q]} r^n}$$

$$\text{So, } \rho_g^{-(p,q)}(f^n) \leq (\rho_{g \circ h}^{(p,q)}(f) + \varepsilon).$$

Since  $\varepsilon > 0$  be arbitrary

$$\rho_g^{-(p,q)}(f^n) \leq \rho_{g \circ h}^{(p,q)}(f). \quad (9)$$

On the other hand for a sequence of values of  $r = r_n$

$$\begin{aligned} & [M_f(r_n)]^n > M_f(r_n) \\ & > M_{g \circ h} \left[ \exp^{[p-1]} \{ (\rho_{g \circ h}^{(p,q)}(f) - \varepsilon) \log^{[q]}(r_n) \} \right] \end{aligned}$$

$$> M_g \left[ \frac{1}{16} M_h \left\{ \frac{\exp^{[p-1]} \{ (\rho_{g \circ h}^{(p,q)}(f) - \varepsilon) \log^{[q]}(r_n) \}}{2} \right\} \right], \text{ by(1)}$$

$$\text{i.e., } M_g^{-1} [M_f(r_n)]^n > \frac{1}{16} M_h \left[ \frac{\exp^{[p-1]} \{ (\rho_{g \circ h}^{(p,q)}(f) - \varepsilon) \log^{[q]}(r_n) \}}{2} \right]$$

i.e.,

$$\log M_g^{-1} [M_f(r_n)]^n > \log M_h \left[ \frac{\exp^{[p-1]} \{ (\rho_{g \circ h}^{(p,q)}(f) - \varepsilon) \log^{[q]}(r_n) \}}{2} \right] + O(1)$$

$$> \left[ \frac{\exp^{[p-1]} \{ (\rho_{g \circ h}^{(p,q)}(f) - \varepsilon) \log^{[q]}(r_n) \}}{2} \right]^{(\lambda_h - \varepsilon)} + O(1)$$

$$\text{i.e., } \log \log M_g^{-1} [M_f(r_n)]^n > (\lambda_h - \varepsilon) \left[ \exp^{[p-2]} \{ (\rho_{g \circ h}^{(p,q)}(f) - \varepsilon) \log^{[q]}(r_n) \} \right] + O(1)$$

$$\text{i.e., } \log^{[p]} M_g^{-1} [M_f(r_n)]^n > (\rho_{g \circ h}^{(p,q)}(f) - \varepsilon) \log^{[q]}(r_n) + O(1).$$

$$\text{So, } \rho_g^{-(p,q)}(f^n) \geq (\rho_{g \circ h}^{(p,q)}(f) - \varepsilon).$$

Since  $\varepsilon > 0$  be arbitrary,

$$\rho_g^{-(p,q)}(f^n) \geq \rho_{g \circ h}^{(p,q)}(f). \tag{10}$$

From (9) and (10) we have,

$$\rho_g^{-(p,q)}(f^n) = \rho_{g \circ h}^{(p,q)}(f).$$

□

**Theorem 4.3.** *If  $f_1, f_2, g$  and  $h$  are entire functions with  $0 < \lambda_h \leq \rho_h < \infty$ , where  $g$  is transcendental and  $g \circ h$  has the property (A) then for  $p > 2$*

$$(i) \rho_g^{-(p,q)}(f_1 f_2) \leq \max\{\rho_{g \circ h}^{(p,q)}(f_1), \rho_{g \circ h}^{(p,q)}(f_2)\}.$$

$$(ii) \text{ Equality holds if } \rho_{g \circ h}^{(p,q)}(f_1) \neq \rho_{g \circ h}^{(p,q)}(f_2).$$

*Proof.* By Lemma 2.4,  $g \circ h$  is transcendental. We consider the following three cases.

Case(a).  $f_1$  and  $f_2$  both are polynomials. Then by Lemma 2.6

$$\rho_g^{-(p,q)}(f_1 f_2) \leq \max\{\rho_{g \circ h}^{(p,q)}(f_1), \rho_{g \circ h}^{(p,q)}(f_2)\}.$$

Case(b).  $f_1$  is polynomial and  $f_2$  is transcendental. Then by Theorem 4.1

$$\rho_g^{-(p,q)}(f_1 f_2) \leq \max\{\rho_{g \circ h}^{(p,q)}(f_1), \rho_{g \circ h}^{(p,q)}(f_2)\}.$$

Case(c).  $f_1$  and  $f_2$  both are transcendental. Let  $\rho_{g \circ h}^{(p,q)}(f_1) \leq \rho_{g \circ h}^{(p,q)}(f_2)$  and  $k = f_1 f_2$ . Let  $\varepsilon > 0$  be arbitrary. For all large  $r$ , we have

$$\begin{aligned} M_{f_1}(r) &< M_{g \circ h} \left[ \exp^{[p-1]} \left\{ (\rho_{g \circ h}^{(p,q)}(f_1) + \frac{\varepsilon}{2}) \log^{[q]} r \right\} \right] \\ &\leq M_{g \circ h} \left[ \exp^{[p-1]} \left\{ (\rho_{g \circ h}^{(p,q)}(f_2) + \frac{\varepsilon}{2}) \log^{[q]} r \right\} \right] \end{aligned}$$

$$\text{and } M_{f_2}(r) < M_{g \circ h} \left[ \exp^{[p-1]} \left\{ (\rho_{g \circ h}^{(p,q)}(f_2) + \frac{\varepsilon}{2}) \log^{[q]} r \right\} \right].$$

Now,

$$\frac{\exp^{[p-2]} \left\{ (\rho_{g \circ h}^{(p,q)}(f_2) + \varepsilon) \log^{[q]} r \right\}}{\exp^{[p-2]} \left\{ (\rho_{g \circ h}^{(p,q)}(f_2) + \frac{\varepsilon}{2}) \log^{[q]} r \right\}} \rightarrow \infty$$

as  $r \rightarrow \infty$ .

So for all large  $r$ , say  $r \geq r_1 > r_0$  the above expression is greater than

$$\frac{\exp^{[p-2]} \left\{ (\rho_{g \circ h}^{(p,q)}(f_2) + \varepsilon) \log^{[q]} r_0 \right\}}{\exp^{[p-2]} \left\{ (\rho_{g \circ h}^{(p,q)}(f_2) + \frac{\varepsilon}{2}) \log^{[q]} r_0 \right\}} = \sigma$$

(say).

Then  $\sigma > 1$ . For all large  $r$ , we have,

$$\begin{aligned} M_k(r) &\leq M_{f_1}(r) M_{f_2}(r) < \left( M_{g \circ h} \left[ \exp^{[p-1]} \left\{ (\rho_{g \circ h}^{(p,q)}(f_2) + \frac{\varepsilon}{2}) \log^{[q]} r \right\} \right] \right)^2 \\ &< \left( M_{g \circ h} \left[ \exp^{[p-1]} \left\{ (\rho_{g \circ h}^{(p,q)}(f_2) + \frac{\varepsilon}{2}) \log^{[q]} r \right\} \right] \right)^\sigma \end{aligned}$$

since  $g \circ h$  has the property (A) and  $\sigma > 1$

$$< M_{g \circ h} \left[ \exp^{[p-1]} \left\{ (\rho_{g \circ h}^{(p,q)}(f_2) + \varepsilon) \log^{[q]} r \right\} \right]$$

for  $r \geq r_1 > r_0$

$$\leq M_g \left[ M_h \left\{ \exp^{[p-1]} \left\{ (\rho_{g \circ h}^{(p,q)}(f_2) + \varepsilon) \log^{[q]} r \right\} \right\} \right] \text{ by (2).}$$

So,

$$M_g^{-1} M_k(r) \leq M_h \left[ \exp^{[p-1]} \left\{ (\rho_{g \circ h}^{(p,q)}(f_2) + \varepsilon) \log^{[q]} r \right\} \right]$$

i.e.,

$$\begin{aligned} \log M_g^{-1}M_k(r) &\leq \log M_h \left[ \exp^{[p-1]} \{ (\rho_{goh}^{(p,q)}(f_2) + \varepsilon) \log^{[q]} r \} \right] \\ &< \left[ \exp^{[p-1]} \{ (\rho_{goh}^{(p,q)}(f_2) + \varepsilon) \log^{[q]} r \} \right]^{(\rho_h + \varepsilon)} \end{aligned}$$

i.e.,

$$\log \log M_g^{-1}M_k(r) < (\rho_h + \varepsilon) \log \left[ \exp^{[p-1]} \{ (\rho_{goh}^{(p,q)}(f_2) + \varepsilon) \log^{[q]} r \} \right]$$

i.e.,

$$\log^{[p]} M_g^{-1}M_k(r) < (\rho_{goh}^{(p,q)}(f_2) + \varepsilon) \log^{[q]} r + O(1).$$

So,  $\limsup_{r \rightarrow \infty} \frac{\log^{[p]} M_g^{-1}M_k(r)}{\log^{[q]} r} \leq \rho_{goh}^{(p,q)}(f_2) + \varepsilon$ .

Since  $\varepsilon > 0$  be arbitrary,

$$\bar{\rho}_g^{(p,q)}(k) \leq \rho_{goh}^{(p,q)}(f_2) = \max\{\rho_{goh}^{(p,q)}(f_1), \rho_{goh}^{(p,q)}(f_2)\}. \tag{11}$$

(ii) Suppose  $\rho_{goh}^{(p,q)}(f_1) < \rho_{goh}^{(p,q)}(f_2)$ . Without loss of generality we may assume  $f_1(0) = 1$ .

We choose  $\mu, \lambda$  so that  $\rho_{goh}^{(p,q)}(f_1) < \mu < \lambda < \rho_{goh}^{(p,q)}(f_2)$ . There exists a sequence  $R_n \rightarrow \infty$  such that

$$M_{f_2}(R_n) > M_{goh}[\exp^{[p-1]}(\lambda \log^{[q]} R_n)], \text{ for } n = 1, 2, 3, \dots$$

Also for all large  $r$ ,

$$M_{f_1}(r) < M_{goh}[\exp^{[p-1]}(\mu \log^{[q]} r)].$$

In Lemma 2.3, we take  $f_1(z)$  for  $f(z), \eta = \frac{1}{16}, R = 2R_n$  and obtain

$$\log|f_1(z)| > -T(\eta) \log M_{f_1}(2e \cdot 2R_n),$$

where

$$T(\eta) = 2 + \log\left(\frac{3e}{2 \cdot \frac{1}{16}}\right) = 2 + \log(24e).$$

So,

$$\log|f_1(z)| > -(2 + \log(24e)) \log M_{f_1}(4eR_n)$$

holds within and on  $|z| = 2R_n$  but outside a family of excluded circles the sum of whose radii is not greater than  $4 \cdot \frac{1}{16} \cdot 2R_n$  i.e.,  $\frac{R_n}{2}$ .

If  $r_n \in (R_n, 2R_n)$  then on  $|z| = r_n, \log|f_1(z)| > -7 \log M_{f_1}(4eR_n)$ .

Now,

$$\begin{aligned} M_{f_2}(r_n) &> M_{f_2}(R_n) > M_{goh} \left[ \exp^{[p-1]}(\lambda \log^{[q]} R_n) \right] \\ &> M_{goh} \left[ \exp^{[p-1]}(\lambda \log^{[q]} \frac{r_n}{2}) \right]. \end{aligned}$$

Let  $z_r$  be a point on  $|z| = r_n$  such that  $M_{f_2}(r_n) = |f_2(z_r)|$ .  
Then

$$\begin{aligned} M_k(r_n) &= \max\{|k(z)| : |z| = r_n\} \\ &= \max\{|f_1(z)||f_2(z)| : |z| = r_n\} \\ &> [M_{f_1}(4eR_n)]^{-7} M_{goh}[\exp^{[p-1]}(\lambda \log^{[q]} \frac{r_n}{2})] \\ &> [M_{goh}\{\exp^{[p-1]}(\mu \log^{[q]}(4eR_n))\}]^{-7} M_{goh}[\exp^{[p-1]}(\lambda \log^{[q]} \frac{r_n}{2})] \\ &\geq [M_{goh}\{\exp^{[p-1]}(\mu \log^{[q]}(4er_n))\}]^{-7} M_{goh}[\exp^{[p-1]}(\lambda \log^{[q]} \frac{r_n}{2})], \end{aligned}$$

since  $r_n > R_n$ .

Now for all large  $n$ , we have  $\frac{\log^{[q]} 4er_n}{\log^{[q]} \frac{r_n}{2}} < \frac{\lambda}{\gamma}$  where  $\gamma \in (\mu, \lambda)$ . So,

$$M_k(r_n) > [M_{goh}\{\exp^{[p-1]}(\mu \log^{[q]}(4er_n))\}]^{-7} M_{goh}[\exp^{[p-1]}(\gamma \log^{[q]} 4er_n)]. \quad (12)$$

The expression

$$\frac{\exp^{[p-2]}(\gamma \log^{[q]}(4er_n))}{\exp^{[p-2]}(\mu \log^{[q]}(4er_n))}$$

tends to infinity as  $n \rightarrow \infty$ .

So for all large  $n$ ,  $r_n \geq r_1 > r_0$  we may write

$$\frac{\exp^{[p-2]}(\gamma \log^{[q]}(4er_n))}{\exp^{[p-2]}(\mu \log^{[q]}(4er_n))} > \frac{\exp^{[p-2]}(\gamma \log^{[q]}(4er_0))}{\exp^{[p-2]}(\mu \log^{[q]}(4er_0))} = \alpha$$

(say), then  $\alpha > 1$ .

From Lemma 2.1 and for all large  $r$ , we have

$$[M_{goh}\{\exp^{[p-1]}(\mu \log^{[q]}(4er_n))\}]^\alpha$$

$$> [M_{goh}\{exp^{[p-1]}(\mu log^{[q]}(4er_n))\}]^8. \tag{13}$$

Also for the above value of  $\alpha$ , one can easily verify that

$$\begin{aligned} & [M_{goh}\{exp^{[p-1]}(\gamma log^{[q]}(4er_n))\}] \\ & > [M_{goh}\{exp^{[p-1]}(\mu log^{[q]}(4er_n))\}^\alpha]. \end{aligned} \tag{14}$$

Therefore for all large  $n$ , we have from (12), (13) and (14)

$$\begin{aligned} M_k(r_n) & > M_{goh} [exp^{[p-1]}(\mu log^{[q]}(4er_n))] \\ & > M_{goh} [exp^{[p-1]}(\mu log^{[q]}r_n)] \\ & \geq M_g \left[ \frac{1}{16} M_h \left\{ \frac{exp^{[p-1]}(\mu log^{[q]}r_n)}{2} \right\} \right] \text{ by (1).} \end{aligned}$$

So,

$$M_g^{-1}M_k(r_n) > \frac{1}{16} M_h \left[ \frac{exp^{[p-1]}(\mu log^{[q]}r_n)}{2} \right]$$

i.e.,

$$\begin{aligned} \log M_g^{-1}M_k(r_n) & > \log M_h \left[ \frac{exp^{[p-1]}(\mu log^{[q]}r_n)}{2} \right] + O(1) \\ & > \left[ \frac{exp^{[p-1]}(\mu log^{[q]}r_n)}{2} \right]^{(\lambda_h - \epsilon)} + O(1) \end{aligned}$$

i.e.,

$$\log \log M_g^{-1}M_k(r_n) > (\lambda_h - \epsilon) [exp^{[p-2]}(\mu log^{[q]}r_n)] + O(1)$$

i.e.,  $\log^{[p]} M_g^{-1}M_k(r_n) > \mu \log^{[q]} r_n + O(1)$ .

So,  $\rho_g^{-(p,q)}(f_1 f_2) \geq \mu$ .

Now since  $\mu < \rho_{goh}^{(p,q)}(f_2)$  is arbitrary, we have

$$\rho_g^{-(p,q)}(f_1 f_2) \geq \rho_{goh}^{(p,q)}(f_2) = \max\{\rho_{goh}^{(p,q)}(f_1), \rho_{goh}^{(p,q)}(f_2)\}. \tag{15}$$

From (11) and (15), we have

$$\rho_g^{-(p,q)}(f_1 f_2) = \max\{\rho_{goh}^{(p,q)}(f_1), \rho_{goh}^{(p,q)}(f_2)\}.$$

This proves the theorem. □

## 5 Hyper Relative order $(p,q)$ of the derivative

**Theorem 5.1.** *Let  $f, g$  and  $h$  be entire transcendental with  $0 < \lambda_h \leq \rho_h < \infty$ . Then for  $p > 2$*

$$\rho_g^{-(p,q)}(f') = \rho_{goh}^{(p,q)}(f).$$

*Proof.* We write  $M_{f'}(r) = \max\{|f'(z)| : |z| = r\}$ ,  $M_g(r) = \max\{|g'(z)| : |z| = r\}$  and  $M_h(r) = \max\{|h'(z)| : |z| = r\}$ .

Without loss of generality we may assume that  $f(0) = 0$ . Otherwise we set  $f_1(z) = zf(z)$ . Then  $f_1(0) = 0$  and by Theorem 4.1

$$\rho_g^{-(p,q)}(f_1) = \rho_{goh}^{(p,q)}(f).$$

We may write  $f(z) = \int_0^z f'(t)dt$ , where the line of integration is the segment from  $z = 0$  to  $z = re^{i\theta_0}$ ,  $r > 0$ . Let  $z_1 = re^{i\theta_1}$  be such that  $|f(z_1)| = \max\{|f(z)| : |z| = r\}$ . Then

$$M_f(r) = |f(z_1)| = \left| \int_0^{z_1} f'(t)dt \right| \leq rM_{f'}(r). \quad (16)$$

Let  $C$  denote the circle  $|t - z_0| = r$ , where  $z_0, |z_0| = r$  is defined so that

$$|f'(z_0)| = \max\{|f'(z)| : |z| = r\}.$$

So,

$$M_{f'}(r) = \max\{|f'(z)| : |z| = r\} =$$

$$|f'(z_0)| = \left| \frac{1}{2\pi i} \oint_C \frac{f(t)}{(t - z_0)^2} dt \right| \leq \frac{1}{2\pi} \frac{M_f(2r)}{r^2} 2\pi r = \frac{M_f(2r)}{r}. \quad (17)$$

From (16) and (17) we obtain

$$\frac{M_f(r)}{r} \leq M_{f'}(r) \leq \frac{M_f(2r)}{r}, \text{ for } r > 0. \quad (18)$$

Let  $\sigma \in (0, 1)$ . From Lemma 2.2(d),  $\lim_{r \rightarrow \infty} \frac{M_f(r^s)}{r^n M_f(r)} = \infty$ , where  $s > 1$  and  $n$  is a positive integer and so  $M_f(r^s) > r^n M_f(r)$  for all large  $r$ . If we replace  $r$  by  $r^\sigma$  and  $s = \frac{1}{\sigma}$  then from above  $M_f(r^{s\sigma}) > r^{n\sigma} M_f(r^\sigma) \geq r M_f(r^\sigma)$ , where the positive integer  $n$  is such that  $n\sigma \geq 1$ .

So  $M_f(r) > r M_f(r^\sigma)$  for all large  $r$ .

From (18) and above, we have



$$M_f(r^\sigma) < \frac{M_f(r)}{r} \leq M_{f'}(r) \leq \frac{M_f(2r)}{r} < M_f(2r) \text{ for all large } r > 1. \quad (19)$$

So,  $M_{f'}(r) < M_f(2r)$

$$< M_{goh} \left[ \exp^{[p-1]} \{ (\rho_{goh}^{(p,q)}(f) + \varepsilon) \log^{[q]} 2r \} \right]$$

$$\leq M_g \left[ M_h \{ \exp^{[p-1]} \{ (\rho_{goh}^{(p,q)}(f) + \varepsilon) \log^{[q]} 2r \} \} \right], \text{ by (2)}$$

i.e.,

$$M_g^{-1} M_{f'}(r) < M_h \left[ \exp^{[p-1]} \{ (\rho_{goh}^{(p,q)}(f) + \varepsilon) \log^{[q]} 2r \} \right]$$

i.e.,

$$\log M_g^{-1} M_{f'}(r) < \log M_h \left[ \exp^{[p-1]} \{ (\rho_{goh}^{(p,q)}(f) + \varepsilon) \log^{[q]} 2r \} \right]$$

$$< \left[ \exp^{[p-1]} \{ (\rho_{goh}^{(p,q)}(f) + \varepsilon) \log^{[q]} 2r \} \right]^{(\rho_h + \varepsilon)}$$

i.e.,

$$\log \log M_g^{-1} M_{f'}(r) < (\rho_h + \varepsilon) \left[ \exp^{[p-2]} \{ (\rho_{goh}^{(p,q)}(f) + \varepsilon) \log^{[q]} 2r \} \right]$$

i.e.,

$$\log^{[p]} M_g^{-1} M_{f'}(r) < (\rho_{goh}^{(p,q)}(f) + \varepsilon) \log^{[q]} 2r + O(1).$$

So,

$$\limsup_{r \rightarrow \infty} \frac{\log^{[p]} M_g^{-1} M_{f'}(r)}{\log^{[q]} r} \leq \lim_{r \rightarrow \infty} \frac{(\rho_{goh}^{(p,q)}(f) + \varepsilon) \log^{[q]} 2r}{\log^{[q]} 2r} \cdot \lim_{r \rightarrow \infty} \frac{\log^{[q]} 2r}{\log^{[q]} r}$$

i.e.,

$$\rho_g^{-(p,q)}(f') \leq \rho_{goh}^{(p,q)}(f) + \varepsilon.$$

Since  $\varepsilon > 0$  be arbitrary,

$$\rho_g^{-(p,q)}(f') \leq \rho_{goh}^{(p,q)}(f). \quad (20)$$

Now for a sequence of values of  $r = r_n$ , we have from (19)

$$M_{f'}(r_n) > M_f(r_n^\sigma) > M_{goh} \left[ \exp^{[p-1]} \{ (\rho_{goh}^{(p,q)}(f) - \varepsilon) \log^{[q]} r_n^\sigma \} \right]$$

$$> M_g \left[ \frac{1}{16} M_h \left\{ \frac{\exp^{[p-1]} \{ (\rho_{goh}^{(p,q)}(f) - \varepsilon) \log^{[q]} r_n^\sigma \}}{2} \right\} \right], \text{ by (1)}$$

i.e.,

$$M_g^{-1} M_{f'}(r_n) > \frac{1}{16} M_h \left[ \frac{\exp^{[p-1]} \{ (\rho_{goh}^{(p,q)}(f) - \varepsilon) \log^{[q]} r_n^\sigma \}}{2} \right]$$

i.e.,

$$\log M_g^{-1} M_{f'}(r_n) > \log M_h \left[ \frac{\exp^{[p-1]} \{ (\rho_{goh}^{(p,q)}(f) - \varepsilon) \log^{[q]} r_n^\sigma \}}{2} \right] + O(1) >$$

$$\left[ \frac{\exp^{[p-1]} \{ (\rho_{goh}^{(p,q)}(f) - \varepsilon) \log^{[q]} r_n^\sigma \}}{2} \right]^{(\lambda_h - \varepsilon)} + O(1)$$

i.e.,

$$\log \log M_g^{-1} M_{f'}(r_n) > (\lambda_h - \varepsilon) \left[ \exp^{[p-2]} \{ (\rho_{goh}^{(p,q)}(f) - \varepsilon) \log^{[q]} r_n^\sigma \} \right] + O(1)$$

i.e.,

$$\log^{[p]} M_g^{-1} M_{f'}(r_n) > (\rho_{goh}^{(p,q)}(f) - \varepsilon) \log^{[q]} r_n^\sigma + O(1).$$

So,  $\rho_g^{-(p,q)}(f') \geq \rho_{goh}^{(p,q)}(f) - \varepsilon$ .

Since  $\varepsilon > 0$  be arbitrary,

$$\rho_g^{-(p,q)}(f') \geq \rho_{goh}^{(p,q)}(f). \quad (21)$$

From (20) and (21), we have

$$\rho_g^{-(p,q)}(f') = \rho_{goh}^{(p,q)}(f).$$

This proves the theorem.  $\square$

## 6 Conclusion

Our main goal through this paper is to enquire the basic relation between hyper relative (p,q) orders of entire function with respect to a single entire function and also of composition of entire functions which have not studied previously. But still there remains some problems to be investigated for future researchers in this field.

### Acknowledgement

The authors are thankful to referee for his/her valuable suggestions towards the improvement of the paper.

### References

- [1] **W. Bergweiler, G. Jank, and L. Volkmann**, Wachstumsverhalten Zusammengesetzter Funktionen, *Results in Mathematics*, **7**, (1984), 35–53.
- [2] **L. Bernal**, Orden relative de crecimiento de funciones enteras, *Collect. Math.*, **39**, (1988), 209-229.
- [3] **J. Clunie**, The composition of entire and meromorphic functions, *Mathematical Essays dedicated to Macintyre*, (1970), 75–92.
- [4] **S. K. Datta and R. Biswas**, A special type of differential polynomial and its comparative growth properties, *International Journal of Modern Engineering Research*, **3(5)**, (2013), 2606-2614.
- [5] **B. K. Lahiri and Dibyendu Banerjee**, Generalized relative order of entire functions, *Proc. Nat. Acad. Sci. India*, **72(A)** (IV), (2002), 351-371.
- [6] **B. K. Lahiri and Dibyendu Banerjee**, Entire functions of relative order  $(p, q)$ , *Soochow Journal of Mathematics*, **31** (4), (2005), 497–513.
- [7] **O. P. Juneja, G. P. Kapoor, and S. K. Bajpai**, On the  $(p, q)$ -order and lower  $(p, q)$ -order of an entire function, *J. Reine Angew. Math.*, **282**, (1976), 53-67.
- [8] **O. P. Juneja, G. P. Kapoor, and S. K. Bajpai**, On the  $(p, q)$ -type and lower  $(p, q)$ -type of an entire function, *J. Reine Angew. Math.*, **290**, (1977), 180-190.
- [9] **H. S. Kasana**, Existence theorem for proximate type of entire functions with index pair  $(p, q)$ , *Bull. Aus. Math. Soc.*, **35**, (1987), 35-42.
- [10] **H. S. Kasana**, On the coefficients of entire functions with index pair  $(p, q)$ , *Bull. Math.*, **32(80)** (3), (1988), 235-242.
- [11] **H. S. Kasana**, The generalized type of entire functions with index pair  $(p, q)$ , *Comment. Math.*, **2**, (1990), 215-222.
- [12] **B. J. Levin**, *Distribution of zeros of entire functions*, American Mathematical Society, publProvidence, 1980.
- [13] **F. Lu and J. Qi**, A note on the hyper order of entire functions, *Bull. Korean Math. Soc.*, **50** (4), (2013), 1209-1219.
- [14] **D. Sato**, On the rate of growth of entire functions of fast growth, *Bull. Amer. Math. Soc.*, **69**, (1963), 411-414.
- [15] **E. C. Titchmarsh**, *The theory of functions, 2nd ed.*, University Press, Oxford, 1968.

Dibyendu Banerjee  
Department of Mathematics  
Visva-Bharati University  
Santiniketan -731235  
India  
E-mail: [dibyendu192@rediffmail.com](mailto:dibyendu192@rediffmail.com)

Saikat Batabyal  
Annapurnapalli  
Kalikapur  
Bolpur -731204  
India  
E-mail: [santubatabyal@gmail.com](mailto:santubatabyal@gmail.com)

Received: 18.01.2017

Accepted: 2.07.2017