


RESEARCH

Open Access



Hyperglycemia enhances arsenic-induced platelet and megakaryocyte activation

Jonathan D. Newman^{1*} , Christina T. Echagarruga², Yoscar M. Ogando², Emilie Montenont², Yu Chen³, Edward A. Fisher^{1,2} and Jeffrey S. Berger^{1,2}

Abstract

Objective: Low to moderate inorganic arsenic (iAs) exposure is independently associated with cardiovascular disease (CVD), particularly for patients with diabetes mellitus (DM). The mechanism of increased CVD risk from iAs exposure in DM has not been adequately characterized. We evaluated whether increasing concentrations of glucose enhance the effects of iAs on platelet and megakaryocyte activity, key steps in atherothrombosis.

Methods: Healthy donor whole blood was prepared in a standard fashion and incubated with sodium arsenite in a range from 0 to 10 μM . iAs-induced platelet activation was assessed by platelet receptor CD62P (P-selectin) expression and monocyte-platelet and leukocyte-platelet aggregation (MPA and LPA, respectively) in the presence of increasing sodium arsenite and glucose concentrations. Megakaryocyte (Meg-01) cell adhesion and gene expression was assessed after incubation with or without iAs and increasing concentrations of D-glucose.

Results: Platelet activity markers increased significantly with 10 vs. 0 μM iAs ($P < 0.05$ for all) and with higher D-glucose concentrations. Platelet activity increased significantly following co incubation of 1 and 5 μM iAs concentrations with hyperglycemic D-glucose ($P < 0.01$ for both) but not after incubation with euglycemic D-glucose. Megakaryocyte adhesion was more pronounced after co incubation with iAs and hyperglycemic than euglycemic D-glucose, while gene expression increased significantly to iAs only after co incubation with hyperglycemic D-glucose.

Conclusion: We demonstrate that glucose concentrations common in DM potentiate the effect of inorganic arsenic exposure on markers of platelet and megakaryocyte activity. Our results support recent observational cohort data that DM enhances the vasculotoxic effects of arsenic exposure, and suggest that activation of the platelet-megakaryocyte hemostatic axis is a pathway through which inorganic arsenic confers atherothrombotic risk, particularly for patients with DM.

Keywords: Atherothrombosis, Platelet activity, Megakaryocyte adhesion, Inorganic arsenic, Diabetes mellitus, Environmental exposures

Background

The adverse cardiovascular and vasculotoxic effects of long-term exposure to high levels of inorganic arsenic in drinking water have been well characterized [1]. Recent studies have demonstrated an increased risk of

cardiovascular disease (CVD), ischemic heart disease (IHD) and mortality from low-moderate drinking water inorganic arsenic (iAs) exposure (10–20 $\mu\text{g/L}$) common in the United States (U.S.), particularly for patients with diabetes mellitus (DM) [2]. Recent prospective cohort study data indicates the vasculotoxicity and cardiovascular disease risk of environmental pollutants, including inorganic arsenic, may be greater for individuals with diabetes [2, 3]. However, the mechanism of this increased risk of environmental exposures for diabetic vasculopathy

*Correspondence: Jonathan.Newman@nyumc.org

¹ Division of Cardiology and the Center for the Prevention of Cardiovascular Disease, Department of Medicine, New York University School of Medicine, TRB rm. 853, New York, NY 10016, USA
Full list of author information is available at the end of the article

has not been studied. Pathological and clinical studies consistently demonstrate that platelets play a key role in atherothrombosis [4], and have shown the importance of the platelet-megakaryocyte hemostatic axis for vascular disease and CVD events [5–7]. Patients with DM exhibit increased platelet activity both in vitro and in vivo, and heightened platelet function may contribute to excess macrovascular risk in patients with DM [8]. A previous in vitro study of iAs and atherothrombosis used very high concentrations of sodium arsenite and did not examine the effects of hyperglycemia on thrombotic risk [9]. We examined whether glucose concentrations common in DM potentiate the effects of iAs on in vitro measures of platelet and megakaryocyte adhesion and activity.

Methods

Subjects

Whole blood was collected from healthy donors in the fasting state. Subjects were not on any antiplatelet therapy nor did they have any history of cardiovascular disease, metabolic syndrome or DM. All human experiments were performed in accordance with institutional and state guidelines. Phlebotomy was performed after 10 min of quiet rest. Blood was collected following a clean, problem-free venipuncture, using a 21-gauge needle after a 5 cc discard (a tourniquet was used to obtain access and was removed before blood collection). Blood was collected into vacutainer tubes containing 3.2% (0.105 mol/l) sodium citrate for platelet activity measurements. After collection, each tube was gently inverted 3 times and immediately transferred to the laboratory for processing.

Reagents

Sodium arsenite was dissolved in dH₂O for a stock concentration of 1000 μM then added to whole blood at a concentration of up to 10 μM for a total of 30 min, similar to prior studies [10, 11]. Similar procedures were performed to achieve concentrations of 0.1, 1, 5 μM sodium arsenite. D-glucose was dissolved in dH₂O for a stock concentration of 500 mM then added to whole blood and megakaryocytes at concentrations of 5, 15 or 25 mM to approximate euglycemia (5 mM D-glucose ≈ 90 mg/dl blood glucose) to a range of hyperglycemia common in DM (15 mM ≈ 270 mg/dl, 25 mM ≈ 450 mg/dl).

Flow cytometry

To examine the effect of iAs on platelet activity, we first measured platelet activation by assessing platelet P-selectin exposure and the presence of monocyte and lymphocyte platelet aggregates (MPA and LPA, respectively) in whole blood samples. We began with a 10 μM concentration of sodium arsenite used in prior in vitro models with

aortic endothelial [10, 11] and vascular smooth muscle cell cultures [12, 13], a concentration 50–75% less than that used in prior studies of arsenic and thrombosis [9]. P-selectin expression (CD62P) is a cell surface marker primarily expressed by activated platelets and involved in platelet adhesion. To identify platelet specific P-selectin, we performed flow cytometry on whole blood with CD42b and CD61 to constitutively expressed platelet glycoproteins 1b (GP1b) and IIIa (GPIIIa), respectively. Flow cytometric analysis was performed using the BD Accuri flow cytometer (C6 Flow Cytometer). Whole blood was incubated in the dark for 30 min at room temperature with APC-conjugated mouse antibody specific for CD42b (glycoprotein 1b) and FITC-conjugated mouse antibody specific for CD62P (P-selectin) (BD Biosciences) before the mean fluorescence intensity of P-selectin-bound antibody per 10,000 events was measured. P-selectin is a component of the alpha granule membrane of resting platelets that is only expressed on the platelet surface membrane after alpha granule secretion. In-vivo circulating degranulated platelets rapidly lose their surface P-selectin, but continue to circulate and function [14]. Monocyte and leukocyte platelet aggregates provide complementary information on in vivo platelet activation; are independently associated with cardiovascular disease events; [15, 16] and were assessed as events positive to markers CD14-APC and CD45-APC, respectively, in addition to platelet marker CD-61. MPAs were defined as events positive to both monocyte markers (CD14-APC [BD Biosciences]) and the platelet marker CD61-FITC (Dako). Monocytes were identified by their staining with CD14-APC and by their characteristic orthogonal light scatter. Monocytes with adherent platelets were identified by CD14-APC positivity. LPAs were defined as events positive to both leukocyte markers (CD45-APC [BD Biosciences]) and the same platelet marker CD61-FITC (Dako). The leukocytes with adherent platelets were identified by CD45-APC positivity. Appropriate color compensation was determined in singly labeled samples and matched nonspecific antibody controls (Mouse IgG1 FITC [BD Biosciences]). For the co-incubation experiments, whole blood was first incubated with 5 and 15 mM D-glucose for 30 min. We then used lower concentrations of sodium arsenite at 0, 0.1, 1 and 5 μM which were added to solution and incubated for an additional 30 min. P-selectin expression with unstimulated and stimulated with thrombin 0.025 IU/ml (Sigma) was then assessed.

Cell culture and megakaryocyte gene expression

Meg-01 cells were purchased from American Type Culture Collection (VA) and cultured in RPMI-1640 medium supplemented with 10% heat-inactivated fetal

bovine serum (FBS), 100 U/ml penicillin, and 100 µg/ml streptomycin (Invitrogen, CA, USA) at 37 °C in a 5% CO₂ humidified atmosphere, consistent with prior studies [17, 18]. For adhesion assays, 18 mm glass coverslips (Fisher Scientific) coated with collagen (Helena Laboratories, Beaumont, TX, USA) were blocked with 1% BSA in 12-well plates [17, 18]. Meg-01 cells were stained for 10 min with 1 µM DiOC6 (Fisher Scientific), washed and incubated at 2.5 × 10⁵ cells/ml for 3 h with and without addition of iAs (0, 1, 5 and 10 µM) in presence of 5 or 25 mM D-glucose. After the supernatant was aspirated, adherent cells were gently washed with FBS. For each well, five random fields were captured and area of coverage was quantified using Image J (National Institutes of Health, Bethesda, MD).

Nuclear transcription factor kappa B (NFκB) gene expression was measured because of its roles in inflammation, platelet activation, and arsenic vasculopathy [18–20]. Other genes measured include monocyte chemoattractant protein-1 (CCL2) and CD36 that have also been associated with platelet degranulation, diabetes and inflammation. To measure these genes, total RNA was isolated from Meg-01 cells using the Direct-zol RNA Miniprep kit (ZymoResearch, Irvine, CA, USA) and quantified using a Nanodrop ND-2000 spectrophotometer (Wilmington, DE, USA). RNA was converted to cDNA using the iScript cDNA synthesis kit (BioRad). Gene expression of GAPDH and NFκB1 using the Sso fast Evagreen Supermix (BioRad) was assessed with real-time PCR (iCycler Real-Time Detection System, Eppendorf). The sequences of the NFκB1, CCL2 and CD36 primers used for qRT-PCR were CAGATGG CCCATACCTTCAA and TTGCAGATTTTGACC TGAGGG, CCCAAAGAAGCTGTGATCTTCA and GCAGATTCCTGGGTTGTGGA, and CTATTGGGAA GGTCAGTCCGA and CAGGTCTCCCTTCTTTGC ATT, respectively.

Statistical analysis

All experimental values are represented as mean ± standard error of the mean (SEM). Differences in selected categorical variables between the respective comparison groups were analyzed with the χ^2 test of statistical significance. Unpaired two-tailed *t* tests and ANOVA were used to examine differences in continuous variables overall and at each time point under study in the different comparison groups. A value of *P* < 0.05 was considered statistically significant.

Results

We first examined the effect of a 10 µM iAs concentration used previously in endothelial and smooth muscle cell culture to assess the effects of inorganic arsenic exposure

[10–13]. There was a clear increase in platelets expressing P-selectin by flow cytometry following incubation with 10 µM iAs (Fig. 1a, b). Compared to 0 µM iAs, the mean fluorescence intensity of P-selectin expression increased significantly after incubation with 10 µM iAs for both unstimulated and thrombin-stimulated platelets (Fig. 1c, d). We subsequently examined the effect of iAs on monocyte and leukocyte platelet aggregation (MPA and LPA, respectively) a different measure of platelet activity predictive of CVD events [15]. Compared to 0 µM, incubation with 10 µM iAs significantly increased both MPA and LPA (Fig. 1e, f). These experiments demonstrate that sodium arsenite concentrations below those used in prior studies of platelet activation have significant effects on multiple measures of platelet activity [9, 21].

Consistent with prior data [8], platelet activity increased with increasing D-glucose concentrations (Fig. 2a, b). To investigate whether glucose and arsenic had a synergistic effect on platelet activation, we coincubated euglycemic (5 mM ≈ 90 mg/dl) and hyperglycemic (15 mM ≈ 270 mg/dl) concentrations of D-glucose with lower concentrations of sodium arsenite than used to demonstrate platelet activation without glucose coincubation. After incubation at hyperglycemic conditions, exposure to 0.1, 1 and 5 µM sodium arsenite led to marked increases in platelet activation. In contrast, these sodium arsenite concentrations did not potentiate platelet activation at euglycemic concentrations of D-glucose (Fig. 2a, b).

Hyperglycemia may induce prothrombotic changes in megakaryocyte function and platelet thrombogenesis [6]. To test whether glucose and iAs also had a synergistic effect on megakaryocyte adhesion, we coincubated megakaryocytes at euglycemic (5 mM) and hyperglycemic (25 mM) concentrations of D-glucose with 0, 1, 5 and 10 mM concentrations of sodium arsenite. Similar to the results observed for platelet activation, exposure to sub-threshold sodium arsenite concentrations below 10 µM induced significantly greater megakaryocyte adhesion after incubation with a hyperglycemic compared to a euglycemic concentration of D-glucose (Fig. 3a, b). Prior studies have demonstrated megakaryocyte nuclear transcription factor kappa B (NFκB) gene expression is an important regulator of inflammation and platelet activation [18, 19], and may also be an important transcriptional factor for the vascular effects of inorganic arsenic exposure [20]. To verify the prothrombotic effect of iAs, we measured the gene expression of NFκB1 in Meg-01 cells. monocyte chemoattractant protein-1 (CCL2) and CD36, genes involved in platelet activation and degranulation [22, 23], were also measured [22, 23]. Following coincubation of hyperglycemic D-glucose with 5 and 10 µM sodium arsenite, Meg-01 cells NFκB1 expression







Fig. 1 Unstimulated (a, c) and thrombin-stimulated (b, d) platelet activation by flow cytometry (a, b), mean fluorescence intensity (c, d) and platelet aggregation (e, f) to 0 and 10 μM sodium arsenite

increased significantly compared to coincubation with euglycemic D-glucose (Fig. 4). There were additional non-significant increases in MCP-1 (CCL2) and CD36 (data not shown). No deleterious effects on Meg-01 cell toxicity were observed within the range of concentrations of sodium arsenite used in this study (0–10 μM) up to concentrations 100-fold greater (Appendix, Fig. 5).

Discussion

There are four primary findings of this report. First, we show for the first time that a concentration of D-glucose common in DM potentiates sodium arsenite-induced platelet activation. Second, we demonstrate that hyperglycemia also potentiates the effects of sodium arsenite on megakaryocyte adhesion, a marker of atherothrombotic



risk [18]. Third, we demonstrate that lower concentrations of sodium arsenite than previously studied are associated with increased platelet activation and aggregation. Finally, we show that Meg01 NFκB transcription as a marker of megakaryocyte activation increases following exposure to hyperglycemia and sodium arsenite. These findings suggest that alterations in the platelet-megakaryocyte axis may be a pathway through which exposure to environmental toxicants such as iAs increase CVD risk, particularly for patients with DM.

Despite advances in effective medical therapy to reduce CVD events, nearly 70% of patients with DM will die of CVD [24]. The etiology of this excess CVD risk for DM patients remains unclear. The vasculotoxicity and

cardiovascular disease risk of environmental pollutants, including iAs, may be greater for individuals with diabetes [2, 3], and suggests that low-level environmental exposures may be a novel risk factor for CVD risk in DM. Environmental pollutants enhance inflammation and the generation of reactive oxygen species, steps also important in the pathogenesis of diabetic vasculopathy [25]. While prior studies have indicated that environmental exposures increase oxidative stress and platelet activation [26, 27], to our knowledge this is the first report to describe a potential link between diabetic hyperglycemia and enhanced atherothrombotic risk to iAs exposure.

There are a number of pathways of platelet activation shared between hyperglycemia and iAs exposure. Hyperglycemia and diabetes is associated with platelet hyperreactivity, and coupled with enhanced levels of thromboxane, may partially explain increases in cardiovascular disease morbidity and mortality seen among patients with DM [8]. High levels of drinking water inorganic arsenic (500 ppb) increase platelet thromboxane formation and adhesion protein expression [28]. Other synergistic pathways between hyperglycemia and iAs exposure include increases in aldose reductase activity and oxidative stress signaling. During hyperglycemia aldose reductase activity increases significantly, leading to abnormal activation of the polyol pathway and enhanced oxidative and osmotic stress [8]. In turn aldose reductase increases thromboxane formation and platelet activation [8]. Inorganic arsenic has also been shown to increase aldose reductase activity [29]. Taken together enhanced aldose reductase activity and thromboxane generation may represent a synergistic pathway of thrombotic risk for both hyperglycemia and inorganic arsenic exposure. Platelet and endothelial mitochondrial function may be another synergistic pathway of risk for iAs exposure in diabetes. Recent studies have indicated the importance of platelet mitochondrial function in cardiovascular disease [30], and have suggested that alterations in platelet mitochondrial function may increase the risk of diabetic atherothrombosis [31]. Inorganic arsenic has also been shown to alter endothelial cell mitochondrial function [13]. Future studies might consider the synergy of inorganic arsenic exposure and diabetes on mitochondrial function in platelets and vascular endothelium as novel pathways of cardiovascular disease risk.

Strengths of the current study include the use of multiple validated measures of the platelet-megakaryocyte axis associated with incident CVD; use of sodium arsenite concentrations below those used in previous models of iAs-induced atherothrombosis; and an investigation of the synergy between hyperglycemia and iAs exposure on atherothrombotic risk. Although we used a lower sodium arsenite concentration than previous

atherothrombosis studies [9, 21], we recognize the concentrations of sodium arsenite used may not correspond to current levels of iAs exposure in the U.S. Future studies should further investigate effects at very low concentrations corresponding to levels more prevalent in human populations. The discrepancy between exposure levels relevant to naturally contaminated drinking water and in vitro concentrations of sodium arsenite may reflect the lack of an accepted biomarker of internal iAs dose. Other limitations include the use of in vitro models and the inability to model in vivo differences in hyperglycemia and insulin resistance seen in type 1 and 2 diabetes. Further study is also needed to better estimate internal inorganic arsenic dose relevant for in vitro modeling; to examine the effect of environmental exposures on the platelet-megakaryocyte axis across the spectrum of diabetes control; and to study the effects of iAs and hyperglycemia on mitochondrial function in platelets and other relevant systems. Treatment studies could consider the use of aldose-reductase inhibitors to attenuate platelet activation and megakaryocyte adhesion [8].

Conclusion

Our findings suggest that increased platelet activation and megakaryocyte adhesion may be pathways through which hyperglycemia in DM can enhance the vasculotoxicity of inorganic arsenic exposure. While intensive glycemic control has failed to significantly reduce macrovascular risk in DM, exposure to environmental toxicants such as inorganic arsenic may represent a novel class of modifiable CVD risk factors, particularly for patients with diabetes. Future studies should investigate platelet activation in patients with and without diabetes, at varying levels of glycemic control, following exposure to environmentally relevant concentrations of inorganic arsenic and other environmental exposures.

Abbreviations

iAs: inorganic arsenic; CVD: cardiovascular disease; DM: diabetes mellitus; MPA: monocyte-platelet aggregation; LPA: leukocyte-platelet aggregation; Meg-01: megakaryocyte; NFκB: nuclear transcription factor kappa B; FBS: fetal bovine serum; RNA: ribonucleic acid; GAPDH: glyceraldehyde 3-phosphate dehydrogenase; PCR: polymerase chain reaction; SEM: standard error of the mean; ANOVA: analysis of variance; U.S.: United States.

Authors' contributions

JDN lead and corresponding author, had full access to the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis. He wrote the majority of the manuscript. CTE performed many of the experiments, made substantial critical revisions and aided with interpretation. YMO aided substantially with experimental conditions and data acquisition. EM aided substantially with experimental conditions and data acquisition. YC provided critical revisions to the manuscript and aided substantially in the preparation of the revised submission. EAF provided input into study design and made substantial critical revisions to the manuscript. JSB provided crucial laboratory support and input into experimental design and

analysis, and made substantial critical revisions to the manuscript. All authors read and approved the final manuscript.

Author details

¹ Division of Cardiology and the Center for the Prevention of Cardiovascular Disease, Department of Medicine, New York University School of Medicine, TRB rm. 853, New York, NY 10016, USA. ² Department of Medicine, Marc and Ruti Bell Program in Vascular Biology and Disease, New York University Medical Center, New York, NY, USA. ³ Departments of Medicine, Population Health and Environmental Medicine, New York University Medical Center, New York, NY, USA.

Acknowledgements

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Availability of data and materials


Data and materials will be held by the corresponding author, Dr. Jonathan Newman, and are available on request.

Funding

This study was funded by the National Institute of Diabetes and Digestive and Kidney Disease (NIDDK) of the National Institute of Health (NIH, U24DK076169-09, subcontract 25732-60). Dr. Newman was partially funded by the National Heart, Lung, and Blood Institute (NHLBI) of the NIH (K23HL125991) and the American Heart Association Mentored Clinical and Population Research Award (15MCRP24480132). Dr. Berger was partially funded by the NHLBI of the NIH (HL114978). Funders had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; and preparation, review, or approval of the article.

Appendix

See Fig. 5.



Received: 25 July 2016 Accepted: 17 February 2017

Published online: 06 March 2017

References

- Moon K, Guallar E, Navas-Acien A. Arsenic exposure and cardiovascular disease: an updated systematic review. *Curr Atheroscler Rep*. 2012;14:542–55.
- Moon KA, Guallar E, Umans JG. Association between exposure to low to moderate arsenic levels and incident cardiovascular disease: a prospective cohort study. *Ann Intern Med*. 2013;159:649–59.
- Tellez-Plaza M, Guallar E, Howard BV, Umans JG, Francesconi KA, Goessler W, et al. Cadmium exposure and incident cardiovascular disease. *Epidemiology*. 2013;24:421–9.
- Davi G, Patrono C. Platelet activation and atherothrombosis. *N Engl J Med*. 2007;357:2482–94.
- Pathansali R, Smith N, Bath P. Altered megakaryocyte–platelet haemostatic axis in hypercholesterolaemia. *Platelets*. 2009;12:292–7.
- Brown AS, Hong Y, de Belder A, Beacon H, Beeso J, Sherwood R, et al. Megakaryocyte ploidy and platelet changes in human diabetes and atherosclerosis. *Arterioscler Thromb Vasc Biol*. 1997;17:802–7.
- Rajagopalan S, McKay I, Ford I, Bachoo P, Greaves M, Brittenden J. Platelet activation increases with the severity of peripheral arterial disease: implications for clinical management. *J Vasc Surg*. 2007;46:485–90.
- Tang WH, Stitham J, Gleim S, Di Febbo C, Porreca E, Fava C, et al. Glucose and collagen regulate human platelet activity through aldose reductase induction of thromboxane. *J Clin Invest*. 2011;121:4462–76.
- Bae ON, Lim KM, Noh JY, Chung SM, Kim H, Lee CR, et al. Arsenite-enhanced procoagulant activity through phosphatidylserine exposure in platelets. *Chem Res Toxicol*. 2007;20:1760–8.
- Barchowsky A, Dudek EJ, Treadwell MD, Wetterhahn KE. Arsenic induces oxidant stress and NF-kappa B activation in cultured aortic endothelial cells. *Free Radic Biol Med*. 1996;21:783–90.
- Barchowsky A, Klei LR, Dudek EJ, Swartz HM, James PE. Stimulation of reactive oxygen, but not reactive nitrogen species, in vascular endothelial cells exposed to low levels of arsenite. *Free Radic Biol Med*. 1999;27:1405–12.
- Lynn S, Gurr JR, Lai HT, Jan KY. NADH oxidase activation is involved in arsenite-induced oxidative DNA damage in human vascular smooth muscle cells. *Circ Res*. 2000;86:514–9.
- Pace C, Banerjee TD, Welch B, Khalili R, Dagda RK, Angermann J. Monomethylarsonous acid, but not inorganic arsenic, is a mitochondria-specific toxicant in vascular smooth muscle cells. *Toxicol Vitro*. 2016;35:188–201.
- Michelson AD, Barnard MR, Krueger LA, Frelinger AL III, Furman MI. Evaluation of platelet function by flow cytometry. *Methods*. 2000;21:259–70.
- Furman MI, Barnard MR, Krueger LA, Fox ML, Shilale EA, Lessard DM, et al. Circulating monocyte-platelet aggregates are an early marker of acute myocardial infarction. *JACC*. 2001;38:1002–6.
- Michelson AD, Barnard MR, Krueger LA, Valeri CR, Furman MI. Circulating monocyte-platelet aggregates are a more sensitive marker of in vivo platelet activation than platelet surface P-selectin: studies in baboons, human coronary intervention, and human acute myocardial infarction. *Circulation*. 2001;104:1533–7.
- Beaulieu LM, Lin E, Morin KM, Tanriverdi K, Freedman JE. Regulatory effects of TLR2 on megakaryocytic cell function. *Blood*. 2011;117:5963–74.
- Beaulieu LM, Lin E, Mick E, Koupenova M, Weinberg EO, Kramer CD, et al. Interleukin 1 receptor 1 and interleukin 1β regulate megakaryocyte maturation, platelet activation, and transcript profile during inflammation in mice and humans. *Arterioscler Thromb Vasc Biol*. 2014;34:552–64.
- Spinelli SL, Casey AE, Pollock SJ, Gertz JM, McMillan DH, Narasipura SD, et al. Platelets and megakaryocytes contain functional nuclear factor-kappaB. *Arterioscler Thromb Vasc Biol*. 2010;30:591–8.
- Wei M, Liu J, Xu M, Rui D, Xu S, Feng G, et al. Divergent effects of arsenic on NF-κB signaling in different cells or tissues: a systematic review and meta-analysis. *Int J Environ Res Public Health*. 2016;13:163.
- Bae O-N, Lim KM, Noh JY, Chung SM, Kim SH, Chung JH. Trivalent methylated arsenical-induced phosphatidylserine exposure and apoptosis in platelets may lead to increased thrombus formation. *Toxicol Appl Pharmacol*. 2009;239:144–53.
- Gleissner CA, Hundelshausen P, Ley K. Platelet chemokines in vascular disease. *Arterioscler Thromb Vasc Biol*. 2008;28:1920–7.
- Englyst NA, Taube JM, Aitman TJ, Baglin TP, Byrne CD. A novel role for CD36 in VLDL-enhanced platelet activation. *Diabetes*. 2003;52:1248–55.
- Seshasai SRK, Kaptoge S, Thompson A, Di Angelantonio E, Gao P, Sarwar N, et al. Diabetes mellitus, fasting glucose, and risk of cause-specific death. *N Engl J Med*. 2011;364:829–41.
- Lamas GA, Navas-Acien A, Mark DB, Lee KL. Heavy metals, cardiovascular disease, and the unexpected benefits of chelation therapy. *J Am Coll Cardiol*. 2016;67:2411–8.

26. Bunderson M, Coffin JD, Beall HD. Arsenic induces peroxynitrite generation and cyclooxygenase-2 protein expression in aortic endothelial cells: possible role in atherosclerosis. *Toxicol Appl Pharmacol.* 2002;184:11–8.
27. Messner B, Knoflach M, Seubert A, Ritsch A, Pfaller K, Henderson B, et al. Cadmium is a novel and independent risk factor for early atherosclerosis mechanisms and in vivo relevance. *Arterioscler Thromb Vasc Biol.* 2009;29:1392–8.
28. Lee MY, Bae ON, Chung SM, Kang KT, Lee JY, Chung JH. Enhancement of platelet aggregation and thrombus formation by arsenic in drinking water: a contributing factor to cardiovascular disease. *Toxicol Appl Pharmacol.* 2002;179:83–8.
29. States JC, Srivastava S, Chen Y, Barchowsky A. Arsenic and cardiovascular disease. *Toxicol Sci.* 2009;107:312–23.
30. Baccarelli AA, Byun HM. Platelet mitochondrial DNA methylation: a potential new marker of cardiovascular disease. *Clin Epigenet.* 2015;7:e28.
31. Xin G, Wei Z, Ji C, Zheng H, Gu J, Ma L, et al. Metformin uniquely prevents thrombosis by inhibiting platelet activation and mtDNA release. *Scientific Rep.* 2016;6:36222.

Submit your next manuscript to BioMed Central
and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

