

Hypersurfaces with constant scalar or mean curvature in a unit sphere

Shichang Shu and Annie Yi Han

Abstract. Let M be an $n(n \geq 3)$ -dimensional complete connected hypersurface in a unit sphere $S^{n+1}(1)$. In this paper, we show that (1) if M has non-zero mean curvature and constant scalar curvature $n(n-1)r$ and two distinct principal curvatures, one of which is simple, then M is isometric to the Riemannian product $S^1(\sqrt{1-c^2}) \times S^{n-1}(c)$, $c^2 = \frac{n-2}{nr}$ if $r \geq \frac{n-2}{n-1}$ and $S \leq (n-1)\frac{n(r-1)+2}{n-2} + \frac{n-2}{n(r-1)+2}$. (2) if M has non-zero constant mean curvature and two distinct principal curvatures, one of which is simple, then M is isometric to the Riemannian product $S^1(\sqrt{1-c^2}) \times S^{n-1}(c)$, $c^2 = \frac{n-2}{nr}$ if one of the following conditions is satisfied: (i) $r \geq \frac{n-2}{n-1}$ and $S \leq (n-1)\frac{n(r-1)+2}{n-2} + \frac{n-2}{n(r-1)+2}$; or (ii) $r > 1 - \frac{2}{n}$, $r \neq \frac{n-2}{n-1}$ and $S \geq (n-1)\frac{n(r-1)+2}{n-2} + \frac{n-2}{n(r-1)+2}$, where S is the squared norm of the second fundamental form of M .

M.S.C. 2000: 53C42, 53A10.

Key words: complete hypersurface, scalar curvature, principal curvature.

1 Introduction

Let M be an n -dimensional hypersurface in a unit sphere $S^{n+1}(1)$ of dimension $n+1$. If scalar curvature $n(n-1)r$ of M is constant and $r \geq 1$. S. Y. Cheng and Yau [1] and Li [5] obtained some characterization theorems in terms of the sectional curvature or the squared norm of the second fundamental form of M respectively. We should notice that the condition $r \geq 1$ plays an essential role in their proofs of theorems. On the other hand, for any $0 < c < 1$, by considering the standard immersions $S^{n-1}(c) \subset R^n$, $S^1(\sqrt{1-c^2}) \subset R^2$ and taking the Riemannian product immersion $S^1(\sqrt{1-c^2}) \times S^{n-1}(c) \hookrightarrow R^2 \times R^n$, we obtain a hypersurface $S^1(\sqrt{1-c^2}) \times S^{n-1}(c)$ in $S^{n+1}(1)$ with constant scalar curvature $n(n-1)r$, where $r = \frac{n-2}{nc^2} > 1 - \frac{2}{n}$. Hence, not all Riemannian products $S^1(\sqrt{1-c^2}) \times S^{n-1}(c)$ appear in the results of [1] and [5]. Since the Riemannian product $S^1(\sqrt{1-c^2}) \times S^{n-1}(c)$ has only two distinct principal curvatures and its scalar curvature $n(n-1)r$ is constant and satisfies $r > 1 - \frac{2}{n}$,

Cheng[2] asked the following interesting problem:

Problem 1.1 ([2]). *Let M be an n -dimensional complete hypersurface with constant scalar curvature $n(n-1)r$ in $S^{n+1}(1)$. If $r > 1 - \frac{2}{n}$ and*

$$S \leq (n-1) \frac{n(r-1)+2}{n-2} + \frac{n-2}{n(r-1)+2},$$

then is M isometric to either a totally umbilical hypersurface or the Riemannian product $S^1(\sqrt{1-c^2}) \times S^{n-1}(c)$?

Cheng [2] said that when $r = \frac{n-2}{n-1}$, he answered the Problem 1.1 affirmatively. For the general case, Problem 1.1 is still open.

In this paper, we try to solve Problem 1.1 and give a partial affirmative answer. We obtain the following:

Theorem 1.1. *Let M be an $n(n \geq 3)$ -dimensional complete connected hypersurface in $S^{n+1}(1)$ with non-zero mean curvature and constant scalar curvature $n(n-1)r$ and with two distinct principal curvatures, one of which is simple. If $r \geq \frac{n-2}{n-1}$ and*

$$S \leq (n-1) \frac{n(r-1)+2}{n-2} + \frac{n-2}{n(r-1)+2},$$

then M is isometric to the Riemannian product $S^1(\sqrt{1-c^2}) \times S^{n-1}(c)$, where $c^2 = \frac{n-2}{nr}$.

If M has constant mean curvature, we can obtain the following:

Theorem 1.2. *Let M be an $n(n \geq 3)$ -dimensional complete connected hypersurface in $S^{n+1}(1)$ with non-zero constant mean curvature and with two distinct principal curvatures, one of which is simple. If $r \geq \frac{n-2}{n-1}$ and*

$$S \leq (n-1) \frac{n(r-1)+2}{n-2} + \frac{n-2}{n(r-1)+2},$$

then M is isometric to the Riemannian product $S^1(\sqrt{1-c^2}) \times S^{n-1}(c)$, where $c^2 = \frac{n-2}{nr}$.

Remark 1.1. We shall note that in [9], the author had given a topological answer to problem 1.1 when M is compact. In [3] and [4], Cheng and the author and Suh had given a partial affirmative answer to problem 1.1 when M is compact.

On the other hand, Cheng [2] also proved the following theorem:

Theorem 1.3 ([2]). *Let M be an n -dimensional complete hypersurface in $S^{n+1}(1)$ with constant scalar curvature $n(n-1)r$ and with two distinct principal curvatures, one of which is simple. Then $r > 1 - \frac{2}{n}$ and M is isometric to the Riemannian product $S^1(\sqrt{1-c^2}) \times S^{n-1}(c)$ if $r \neq \frac{n-2}{n-1}$ and*

$$S \geq (n-1) \frac{n(r-1)+2}{n-2} + \frac{n-2}{n(r-1)+2},$$

where $c^2 = \frac{n-2}{nr}$.

If M has constant mean curvature, we can obtain the following:

Theorem 1.4. *Let M be an $n(n \geq 3)$ -dimensional complete connected hypersurface in $S^{n+1}(1)$ with constant mean curvature and with two distinct principal curvatures, one of which is simple. If $r > 1 - \frac{2}{n}$, $r \neq \frac{n-2}{n-1}$ and*

$$S \geq (n-1) \frac{n(r-1)+2}{n-2} + \frac{n-2}{n(r-1)+2},$$

then M is isometric to the Riemannian product $S^1(\sqrt{1-c^2}) \times S^{n-1}(c)$, where $c^2 = \frac{n-2}{nr}$.

Remark 1.2. Recently, the authors also studied the complete hypersurfaces in a hyperbolic space with constant scalar curvature or with constant k -th mean curvature and with two distinct principal curvatures, one can see [7] and [8]. On the study of stable spacelike hypersurfaces with constant scalar curvature, one can see [6].

2 Preliminaries

Let M be an n -dimensional hypersurface in $S^{n+1}(1)$. We choose a local orthonormal frame e_1, \dots, e_{n+1} in $S^{n+1}(1)$ such that e_1, \dots, e_n are tangent to M . Let $\omega_1, \dots, \omega_{n+1}$ be the dual coframe. We use the following convention on the range of indices:

$$1 \leq A, B, C, \dots \leq n+1; \quad 1 \leq i, j, k, \dots \leq n.$$

The structure equations of $S^{n+1}(1)$ are given by

$$(2.1) \quad d\omega_A = \sum_B \omega_{AB} \wedge \omega_B, \quad \omega_{AB} + \omega_{BA} = 0,$$

$$(2.2) \quad d\omega_{AB} = \sum_C \omega_{AC} \wedge \omega_{CB} + \Omega_{AB},$$

where

$$(2.3) \quad \Omega_{AB} = -\frac{1}{2} \sum_{C,D} K_{ABCD} \omega_C \wedge \omega_D,$$

$$(2.4) \quad K_{ABCD} = \delta_{AC} \delta_{BD} - \delta_{AD} \delta_{BC}.$$

Restricting to M ,

$$(2.5) \quad \omega_{n+1} = 0,$$

$$(2.6) \quad \omega_{n+1i} = \sum_j h_{ij} \omega_j, \quad h_{ij} = h_{ji}.$$

The structure equations of M are

$$(2.7) \quad d\omega_i = \sum_j \omega_{ij} \wedge \omega_j, \quad \omega_{ij} + \omega_{ji} = 0,$$

$$(2.8) \quad d\omega_{ij} = \sum_k \omega_{ik} \wedge \omega_{kj} - \frac{1}{2} \sum_{k,l} R_{ijkl} \omega_k \wedge \omega_l,$$

$$(2.9) \quad R_{ijkl} = \delta_{ik}\delta_{jl} - \delta_{il}\delta_{jk} + h_{ik}h_{jl} - h_{il}h_{jk},$$

$$(2.10) \quad R_{ij} = (n-1)\delta_{ij} + nHh_{ij} - \sum_k h_{ik}h_{kj},$$

$$(2.11) \quad n(n-1)r = n(n-1) + n^2H^2 - S,$$

where $n(n-1)r$ is the scalar curvature, H is the mean curvature and S is the squared norm of the second fundamental form of M .

3 Proof of theorem

Let M be an $n(n \geq 3)$ -dimensional complete connected hypersurface in $S^{n+1}(1)$ with constant scalar curvature $n(n-1)r$ and with two distinct principal curvatures, one of which is simple. Without loss of generality, we may assume

$$(3.1) \quad \lambda_1 = \lambda_2 = \cdots = \lambda_{n-1} = \lambda, \quad \lambda_n = \mu,$$

where λ_i for $i = 1, 2, \dots, n$ are the principal curvatures of M . From (2.11) and (3.1), we have

$$(3.2) \quad n(n-1)(r-1) = (n-1)(n-2)\lambda^2 + 2(n-1)\lambda\mu.$$

If $\lambda = 0$ at a point of M , then from above equation, we obtain that $r = 1$ at this point. Since the scalar curvature $n(n-1)r$ is constant, we obtain $r \equiv 1$ on M . Since the principal curvatures λ and μ are continuous on M , by the same assertion we can deduce from (3.2) that $\lambda \equiv 0$ on M . By (2.9), we know that the sectional curvature of M is not less than 1. Therefore, we know that M is compact by use of Bonnet-Myers Theorem. According to Theorem 2 in Cheng and Yau [1], we have M^n is totally umbilical. This is impossible, therefore, we get $\lambda \neq 0$. From (3.2), we have

$$(3.3) \quad \mu = \frac{n(r-1)}{2\lambda} - \frac{n-2}{2}\lambda,$$

Since

$$\lambda - \mu = n \frac{\lambda^2 - (r-1)}{2\lambda} \neq 0,$$

we know that $\lambda^2 - (r-1) \neq 0$. If $\lambda^2 - (r-1) < 0$, we deduce that $r > 1$ and $\lambda^2 - \lambda\mu = \frac{n}{2}[\lambda^2 - (r-1)] < 0$. Therefore $\lambda\mu > \lambda^2$. From (2.9), we obtain the sectional curvature of M is not less than 1. Therefore, we know that M is compact by use of Bonnet-Myers Theorem. According to Theorem 2 in Cheng and Yau [1], we have M^n is totally umbilical. This is impossible, therefore, we get $\lambda^2 - (r-1) > 0$.

Let $\varpi = [\lambda^2 - (r-1)]^{-\frac{1}{n}}$. Cheng [2] proved the following:

Proposition 3.1 ([2]). *Let M be an $n(n \geq 3)$ -dimensional connected hypersurface with constant scalar curvature $n(n-1)r$ and with two distinct principal curvatures, and the space of principal vectors corresponding to one of them is of one*

dimension. Then M is a locus of moving $(n-1)$ -dimensional submanifold $M_1^{n-1}(s)$, along which the principal curvature λ of multiplicity $n-1$ is constant and which is locally isometric to an $(n-1)$ -dimensional sphere $S^{n-1}(c(s)) = E^n(s) \cap S^{n+1}(1)$ of constant curvature and $\varpi = [\lambda^2 - (r-1)]^{-\frac{1}{n}}$ satisfies the ordinary differential equation of order 2

$$(3.4) \quad \frac{d^2\varpi}{ds^2} - \varpi \left(\frac{n-2}{n} \varpi^{-n} - r \right) = 0,$$

where $E^n(s)$ is an n -dimensional linear subspace in the Euclidean space R^{n+2} which is parallel to a fixed $E^n(s_0)$.

The following Lemma 3.1 in Wei and Suh [10] is important to us.

Lemma 3.1 ([10]). *Equations (3.4) is equivalent to its first order integral*

$$(3.5) \quad \left(\frac{d\varpi}{ds} \right)^2 + r\varpi^2 + \frac{1}{\varpi^{n-2}} = C,$$

where C is a constant; for a constant solution equal to ϖ_0 , one has that $r > 0$ and $\varpi_0^n = \frac{n-2}{2r}$, so

$$(3.6) \quad C_0 = \frac{n}{2} \left(\frac{2r}{n-2} \right)^{(n-2)/n}.$$

Moreover, the constant solution of (3.4) corresponds to $S^1(\sqrt{1-c^2}) \times S^{n-1}(c)$, where $c^2 = \frac{n-2}{nr}$.

In [10], Wei and Suh proved the following:

Proposition 3.2 ([10]). *Let M be an $n(n \geq 3)$ -dimensional complete connected hypersurface in $S^{n+1}(1)$ with constant scalar curvature $n(n-1)r$ and with two distinct principal curvatures, one of which is simple. If $\lambda\mu + 1 \leq 0$, then M is isometric to $S^1(\sqrt{1-c^2}) \times S^{n-1}(c)$, where $c^2 = \frac{n-2}{nr}$.*

By the same method in [10], we can prove the following:

Proposition 3.3. *Let M be an $n(n \geq 3)$ -dimensional complete connected hypersurface in $S^{n+1}(1)$ with constant scalar curvature $n(n-1)r$ and with two distinct principal curvatures, one of which is simple. If $\lambda\mu + 1 \geq 0$, then M is isometric to the Riemannian product $S^1(\sqrt{1-c^2}) \times S^{n-1}(c)$, where $c^2 = \frac{n-2}{nr}$.*

Proof. From (3.3), we have

$$(3.7) \quad \lambda\mu + 1 = \frac{n(r-1)}{2} - \frac{n-2}{2}\lambda^2 + 1.$$

If $\lambda\mu + 1 \geq 0$, we have from (3.7), $\frac{n-2}{2}[\lambda^2 - (r-1)] \leq r$, it follows that

$$(3.8) \quad \frac{n-2}{2}\varpi^{-n} - r \leq 0.$$

From (3.4), we have $\frac{d^2\varpi}{ds^2} \leq 0$. Thus $\frac{d\varpi}{ds}$ is a monotonic function of $s \in (-\infty, +\infty)$. Therefore, $\varpi(s)$ must be monotonic when s tends to infinity. We see from (3.5) that the positive function of $\varpi(s)$ is bounded. Since $\varpi(s)$ is bounded and is monotonic when s tends infinity, we find that both $\lim_{s \rightarrow -\infty} \varpi(s)$ and $\lim_{s \rightarrow +\infty} \varpi(s)$ exist and then we have

$$(3.9) \quad \lim_{s \rightarrow -\infty} \frac{d\varpi(s)}{ds} = \lim_{s \rightarrow +\infty} \frac{d\varpi(s)}{ds} = 0.$$

By the monotonicity of $\frac{d\varpi}{ds}$, we see that $\frac{d\varpi}{ds} \equiv 0$ and $\varpi(s)$ is a constant. Then, by Lemma 3.1, it is easily see that M is isometric to the Riemannian product $S^1(\sqrt{1-c^2}) \times S^{n-1}(c)$, where $c^2 = \frac{n-2}{nr}$.

Since M has two distinct principal curvatures, we know that M has no umbilical points. From (3.1), we have

$$(3.10) \quad (n-1)\lambda + \mu = nH, \quad S = (n-1)\lambda^2 + \mu^2.$$

From (3.10) and (2.11), we have

$$(3.11) \quad \lambda\mu = (n-1)(r-1) - (n-2)H^2 + (n-2)H\sqrt{H^2 - (r-1)},$$

or

$$(3.12) \quad \lambda\mu = (n-1)(r-1) - (n-2)H^2 - (n-2)H\sqrt{H^2 - (r-1)}.$$

From (2.11), we obtain

$$(3.13) \quad \begin{aligned} \lambda\mu = & (r-1) - \frac{(n-2)}{n^2}[S - n(r-1)] \\ & + \frac{(n-2)}{n^2}\sqrt{[S + n(n-1)(r-1)][S - n(r-1)]}, \end{aligned}$$

or

$$(3.14) \quad \begin{aligned} \lambda\mu = & (r-1) - \frac{(n-2)}{n^2}[S - n(r-1)] \\ & - \frac{(n-2)}{n^2}\sqrt{[S + n(n-1)(r-1)][S - n(r-1)]}. \end{aligned}$$

Proof of theorem 1.1. If there exists a point x on M such that (3.13) and (3.14) hold at x , that is, we have $S = -n(n-1)(r-1)$ or $S = n(r-1)$ at x . If $S = -n(n-1)(r-1)$ at x , from (2.11), we have $H = 0$ at x , this is a contradiction to $H \neq 0$ on M . If $S = n(r-1)$ at x , from (2.11) we have $S = nH^2$ at x , that is, x is a umbilical point on M , this is a contradiction to M has no umbilical points. Therefore, we only consider two cases:

Case (1). If (3.13) holds on M , since $r \geq \frac{n-2}{n-1}$, then we get $r-1 \geq -\frac{1}{n-1}$ and $n(r-1) + 2 \geq \frac{n-2}{n-1}$. From

$$S \leq (n-1)\frac{n(r-1)+2}{n-2} + \frac{n-2}{n(r-1)+2},$$

we have

$$\begin{aligned}
(3.15) \quad & n+n(r-1) - \frac{n-2}{n}[S - n(r-1)] \\
& \geq n + 2(n-1)(r-1) - \frac{n-2}{n} \left[(n-1) \frac{n(r-1)+2}{n-2} + \frac{n-2}{n(r-1)+2} \right] \\
& = n + 2(n-1)(r-1) - \frac{n-1}{n}[n(r-1)+2] - \frac{(n-2)^2}{n} \frac{1}{n(r-1)+2} \\
& = \frac{n^2 - 2(n-1)}{n} + (n-1)(r-1) - \frac{(n-2)^2}{n} \frac{1}{n(r-1)+2} \\
& \geq \frac{n^2 - 2(n-1)}{n} - 1 - \frac{(n-2)^2}{n} \frac{n-1}{n-2} = 0,
\end{aligned}$$

From (3.13) and (3.15), obviously, we have $\lambda\mu + 1 \geq 0$. By Proposition 3.3, we obtain that M is isometric to the Riemannian product $S^1(\sqrt{1-c^2}) \times S^{n-1}(c)$, where $c^2 = \frac{n-2}{nr}$.

Case (2). If (3.14) holds on M , since $S \leq (n-1) \frac{n(r-1)+2}{n-2} + \frac{n-2}{n(r-1)+2}$, we know that this is equivalent to

$$\begin{aligned}
(3.16) \quad & \left\{ n + n(r-1) - \frac{n-2}{n}[S - n(r-1)] \right\}^2 \\
& \geq \frac{(n-2)^2}{n^2} \{n(n-1)(r-1) + S\} \{S - n(r-1)\}.
\end{aligned}$$

Let $f^2 = \sum_i (\lambda_i - H)^2 = S - nH^2$. Obviously, by (2.11), we have $n(n-1)(r-1) + S \geq 0$ and $f^2 = \frac{n-1}{n}[S - n(r-1)]$. Therefore, we know that $S - n(r-1) \geq 0$. From (3.15) and (3.16), we have

$$\begin{aligned}
(3.17) \quad & n + n(r-1) - \frac{n-2}{n}[S - n(r-1)] \\
& \geq \frac{n-2}{n} \sqrt{[n(n-1)(r-1) + S][S - n(r-1)]}.
\end{aligned}$$

Therefore, (3.14) and (3.17) imply that $\lambda\mu + 1 \geq 0$. By Proposition 3.3, we obtain that M is isometric to the Riemannian product $S^1(\sqrt{1-c^2}) \times S^{n-1}(c)$, where $c^2 = \frac{n-2}{nr}$. This completes the proof of Theorem 1.1.

In order to prove Theorem 1.2 and Theorem 1.4, we need the following Propositions due to [11].

Proposition 3.4 ([11]). *Let M be an n ($n \geq 3$)-dimensional connected hypersurface with constant mean curvature H and with two distinct principal curvatures λ and μ with multiplicities $(n-1)$ and 1 , respectively. Then M is a locus of moving $(n-1)$ -dimensional submanifold $M_1^{n-1}(s)$ along which the principal curvature λ of multiplicity $n-1$ is constant and which is locally isometric to an $(n-1)$ -dimensional sphere $S^{n-1}(c(s)) = E^n(s) \cap S^{n+1}(1)$ of constant curvature and $\varpi = |\lambda - H|^{-\frac{1}{n}}$ satisfies the ordinary differential equation of order 2*

$$(3.18) \quad \frac{d^2\varpi}{ds^2} + \varpi[1 + H^2 + (2-n)H\varpi^{-n} + (1-n)\varpi^{-2n}] = 0,$$

for $\lambda - H > 0$ or

$$(3.19) \quad \frac{d^2\varpi}{ds^2} + \varpi[1 + H^2 + (n-2)H\varpi^{-n} + (1-n)\varpi^{-2n}] = 0,$$

for $\lambda - H < 0$, where $E^n(s)$ is an n -dimensional linear subspace in the Euclidean space R^{n+2} which is parallel to a fixed $E^n(s_0)$.

Lemma 3.2 ([11]). Equation (3.18) or (3.19) is equivalent to its first order integral

$$(3.20) \quad \left(\frac{d\varpi}{ds}\right)^2 + (1 + H^2)\varpi^2 + 2H\varpi^{2-n} + \varpi^{2-2n} = C,$$

for $\lambda - H > 0$ or

$$(3.21) \quad \left(\frac{d\varpi}{ds}\right)^2 + (1 + H^2)\varpi^2 - 2H\varpi^{2-n} + \varpi^{2-2n} = C,$$

for $\lambda - H < 0$, where C is a constant. Moreover, the constant solution of (3.18) or (3.19) corresponds to the Riemannian product $S^1(\sqrt{1-c^2}) \times S^{n-1}(c)$.

We can prove the following:

Proposition 3.5. Let M be an $n(n \geq 3)$ -dimensional complete connected hypersurface in $S^{n+1}(1)$ with constant mean curvature H and with two distinct principal curvatures λ and μ with multiplicities $(n-1)$ and 1 , respectively. If $\lambda\mu + 1 \geq 0$, then M is isometric to the Riemannian product $S^1(\sqrt{1-c^2}) \times S^{n-1}(c)$.

Proof. Let λ and μ be the two distinct principal curvatures of M with multiplicities $(n-1)$ and 1 , respectively. Then, from $nH = (n-1)\lambda + \mu$, we have $\lambda\mu = nH\lambda - (n-1)\lambda^2$. Let $\varpi = |\lambda - H|^{-\frac{1}{n}}$. Then we have $\lambda = H + \varpi^{-n}$ for $\lambda - H > 0$ and $\lambda = H - \varpi^{-n}$ for $\lambda - H < 0$. If $\lambda - H > 0$, we have

$$\lambda\mu + 1 = 1 + H^2 + (2-n)H\varpi^{-n} + (1-n)\varpi^{-2n},$$

and if $\lambda - H < 0$, we have

$$\lambda\mu + 1 = 1 + H^2 + (n-2)H\varpi^{-n} + (1-n)\varpi^{-2n}.$$

Therefore, if $\lambda\mu + 1 \geq 0$, we obtain

$$1 + H^2 + (2-n)H\varpi^{-n} + (1-n)\varpi^{-2n} \geq 0,$$

for $\lambda - H > 0$ and

$$1 + H^2 + (n-2)H\varpi^{-n} + (1-n)\varpi^{-2n} \geq 0,$$

for $\lambda - H < 0$. From (3.18) and (3.19), we have $\frac{d^2\varpi}{ds^2} \leq 0$. Thus $\frac{d\varpi}{ds}$ is a monotonic function of $s \in (-\infty, +\infty)$. Therefore, $\varpi(s)$ must be monotonic when s tends to infinity. We see from (3.20) and (3.21) that the positive function of $\varpi(s)$ is bounded. Since $\varpi(s)$ is bounded and is monotonic when s tends infinity, we find that both $\lim_{s \rightarrow -\infty} \varpi(s)$ and $\lim_{s \rightarrow +\infty} \varpi(s)$ exist and then we have

$$(3.22) \quad \lim_{s \rightarrow -\infty} \frac{d\varpi(s)}{ds} = \lim_{s \rightarrow +\infty} \frac{d\varpi(s)}{ds} = 0.$$

By the monotonicity of $\frac{d\varpi}{ds}$, we see that $\frac{d\varpi}{ds} \equiv 0$ and $\varpi(s)$ is a constant. Then, by Lemma 3.2, it is easily see that M is isometric to the Riemannian product $S^1(\sqrt{1-c^2}) \times S^{n-1}(c)$. This completes the proof of Proposition 3.5.

On the other hand, if $\lambda\mu + 1 \leq 0$, from above, we can obtain $\frac{d^2\varpi}{ds^2} \geq 0$. We see from (3.20) and (3.21) that the positive function of $\varpi(s)$ is bounded. Combining $\frac{d^2\varpi}{ds^2} \geq 0$ with the boundedness of $\varpi(s)$, similar to the proof of Proposition 3.5, we know that $\varpi(s)$ is constant. Then, by Lemma 3.2, it is easily see that M is isometric to the Riemannian product $S^1(\sqrt{1-c^2}) \times S^{n-1}(c)$. Therefore, we have the following:

Proposition 3.6. *Let M be an $n(n \geq 3)$ -dimensional complete connected hypersurface in $S^{n+1}(1)$ with constant mean curvature H and with two distinct principal curvatures λ and μ with multiplicities $(n-1)$ and 1 , respectively. If $\lambda\mu + 1 \leq 0$, then M is isometric to the Riemannian product $S^1(\sqrt{1-c^2}) \times S^{n-1}(c)$.*

Proof of theorem 1.2. Since M has non-zero mean curvature, by the same assertion in the proof of Theorem 1.1, we only have two cases: (3.13) holds on M or (3.14) holds on M . If (3.13) holds on M , from the proof of Theorem 1.1, we have $\lambda\mu + 1 \geq 0$. By Proposition 3.5, we obtain that M is isometric to the Riemannian product $S^1(\sqrt{1-c^2}) \times S^{n-1}(c)$, where $c^2 = \frac{n-2}{nr}$. If (3.14) holds on M , from the proof of Theorem 1.1, we have $\lambda\mu + 1 \leq 0$. By Proposition 3.6, we obtain that M is isometric to the Riemannian product $S^1(\sqrt{1-c^2}) \times S^{n-1}(c)$, where $c^2 = \frac{n-2}{nr}$. This completes the proof of Theorem 1.2.

Proof of theorem 1.4. We firstly prove that $H \neq 0$. In fact, if $H = 0$, from (2.11) we have $S = -n(n-1)(r-1)$ on M . Since $S \geq (n-1)\frac{n(r-1)+2}{n-2} + \frac{n-2}{n(r-1)+2}$ is equivalent to

$$\begin{aligned} & \frac{(n-2)^2}{n^2} [S + n(n-1)(r-1)][S - n(r-1)] \\ & \geq \{n + n(r-1) - \frac{(n-2)}{n} [S - n(r-1)]\}^2, \end{aligned}$$

we have from $S = -n(n-1)(r-1)$ that

$$(3.23) \quad 0 \geq \{n + n(n-1)(r-1)\}^2.$$

From (2.23), we have $r = \frac{n-2}{n-1}$, this is a contradiction to the assumption that $r \neq \frac{n-2}{n-1}$.

If there exists a point x on M such that (3.13) and (3.14) hold at x , that is, we have $S = -n(n-1)(r-1)$ or $S = n(r-1)$ at x . If $S = -n(n-1)(r-1)$ at x , from (2.11), we have $H = 0$ at x , this is a contradiction to $H \neq 0$ on M . If $S = n(r-1)$ at x , from (2.11) we have $S = nH^2$ at x , that is, x is a umbilical point on M , this is a contradiction to M has no umbilical points. Therefore, we only consider two cases:

Case (1). If (3.13) holds on M , we can prove $\lambda\mu + 1 \geq 0$ on M . In fact, if $\lambda\mu + 1 < 0$ at a point of M , then at this point

$$\begin{aligned} & \frac{(n-2)}{n^2} \sqrt{[S + n(n-1)(r-1)][S - n(r-1)]} \\ & < -1 - (r-1) + \frac{(n-2)}{n^2} [S - n(r-1)]. \end{aligned}$$

Therefore, we have at this point

$$\begin{aligned} & \frac{(n-2)^2}{n^2} [S + n(n-1)(r-1)][S - n(r-1)] \\ & < \left\{ n + n(r-1) - \frac{(n-2)}{n} [S - n(r-1)] \right\}^2, \end{aligned}$$

this is equivalent to $S < (n-1) \frac{n(r-1)+2}{n-2} + \frac{n-2}{n(r-1)+2}$ at this point, we have a contradiction to $S \geq (n-1) \frac{n(r-1)+2}{n-2} + \frac{n-2}{n(r-1)+2}$ on M . Therefore, in case (1) we have $\lambda\mu + 1 \geq 0$. By Proposition 3.5, we obtain that M is isometric to the Riemannian product $S^1(\sqrt{1-c^2}) \times S^{n-1}(c)$, where $c^2 = \frac{n-2}{nr}$.

Case (2). If (3.14) holds on M , next we shall prove that $\lambda\mu + 1 \leq 0$ on M . We consider three subcases:

(i) If $1 + (r-1) - \frac{(n-2)}{n^2} [S - n(r-1)] \leq 0$ on M , then from (3.14), it is obvious that $\lambda\mu + 1 \leq 0$ on M .

(ii) If $1 + (r-1) - \frac{(n-2)}{n^2} [S - n(r-1)] > 0$ on M , from $S \geq (n-1) \frac{n(r-1)+2}{n-2} + \frac{n-2}{n(r-1)+2}$, we have

$$(n-2)[n(r-1) + 2]S \geq (n-1)n^2(r-1)^2 + 4n(n-1)(r-1) + n^2,$$

that is

$$\begin{aligned} & (n-2)\{4n(n-1)(r-1) + 2n^2 + (n-2)^2n(r-1)\}S \\ & \geq \{2n(n-1)(r-1) + n^2\}^2 + (n-2)^2n^2(n-1)(r-1)^2. \end{aligned}$$

Hence

$$\begin{aligned} (3.24) \quad & \left\{ n + n(r-1) - \frac{n-2}{n} [S - n(r-1)] \right\}^2 \\ & \leq \frac{(n-2)^2}{n^2} \{n(n-1)(r-1) + S\} \{S - n(r-1)\}. \end{aligned}$$

Since $1 + (r-1) - \frac{(n-2)}{n^2} [S - n(r-1)] > 0$ on M , from (3.24), we have

$$\begin{aligned} (3.25) \quad & n + n(r-1) - \frac{n-2}{n} [S - n(r-1)] \\ & \leq \frac{(n-2)}{n} \sqrt{[n(n-1)(r-1) + S][S - n(r-1)]}. \end{aligned}$$

From (3.14), we infer that $\lambda\mu + 1 \leq 0$ on M .

(iii) If $1 + (r-1) - \frac{(n-2)}{n^2} [S - n(r-1)] \leq 0$ at a point p of M and $1 + (r-1) - \frac{(n-2)}{n^2} [S - n(r-1)] > 0$ at other points of M , in this case, from (i) and (ii), we have at point p , $\lambda\mu + 1 \leq 0$ and at other points of M , also $\lambda\mu + 1 \leq 0$. Therefore, we obtain $\lambda\mu + 1 \leq 0$ on M .

Therefore, we know that if (3.14) holds on M , then $\lambda\mu + 1 \leq 0$ on M . By Proposition 3.6, we obtain that M is isometric to the Riemannian product $S^1(\sqrt{1-c^2}) \times S^{n-1}(c)$, where $c^2 = \frac{n-2}{nr}$. This completes the proof of Theorem 1.4.

Acknowledgements. Project supported by NSF of Shaanxi Province (SJ08A31) and NSF of Shaanxi Educational Committee (2008JK484). The authors would like to thank the referee for his many valuable suggestions that have really improved the paper.

References

- [1] S. Y. Cheng and S. T. Yau, *Hypersurfaces with constant scalar curvature*, Math. Ann. 225(1977), 195-204.
- [2] Q. M. Cheng, *Hypersurfaces in a unit sphere $S^{n+1}(1)$ with constant scalar curvature*, J. of London Math. Society, 64(2001), 755-768.
- [3] Q. M. Cheng, S. C. Shu and Y. J. Suh, *Compact hypersurfaces in a unit sphere*, Pro. of the Royal Soci. of Edinburgh, 135A(2005), 1129-1137.
- [4] Q. M. Cheng, *Compact hypersurfaces in a unit sphere with infinite fundamental group*, Pac. J. Math., 212(2003), 49-56.
- [5] H. Li, *Hypersurfaces with constant scalar curvature in space forms*, Math. Ann., 305(1996), 665-672.
- [6] X. Liu and B. Yang, *Stable space-like hypersurfaces with constant scalar curvature in generalized Roberston-Walker spacetimes*, Balkan J. of Geometry and Its Applications, 13(2008), 66-76.
- [7] S. C. Shu and Annie Y. Han, *Hypersurfaces in a hyperbolic space with constant k -th mean curvature*, Bull. Math. Soc. Sci. Math. Roumanie, 52(100)(2009), 65-78.
- [8] S. C. Shu, *Complete hypersurfaces with constant scalar curvature in a hyperbolic space*, Balkan J. of Geometry and Its Applications, 12(2007), 107-115.
- [9] S. C. Shu and S. Y. Liu, *The curvature and topological properties of hypersurfaces with constant scalar curvature*, Bull. Austral Math., 70(2004), 35-44.
- [10] G. Wei and Y. J. Suh, *Rigidity theorems for hypersurfaces with constant scalar curvature in unit sphere*, Glasgow Math. J., 49(2007), 235-241.
- [11] G. Wei, *Complete hypersurfaces with constant mean curvature in a unit sphere*, Monatsh. Math., 149(2006), 251-258.

Authors' addresses:

Shichang Shu
 Department of Mathematics, Xianyang Normal University,
 Xianyang 712000 Shaanxi, P.R.China.
 E-mail:shushichang@126.com

Annie Yi Han
 Department of Mathematics, Borough of Manhattan Community College
 CUNY 10007 N.Y. USA.
 E-mail:DrHan@nyc.rr.com