

Review

Hypertension in rheumatoid arthritis

V. F. Panoulas^{1,2}, G. S. Metsios¹, A. V. Pace¹, H. John¹, G. J. Treharne^{1,3}, M. J. Banks⁴ and G. D. Kitas^{1,5}

RA associates with an increased burden of cardiovascular disease, which is at least partially attributed to classical risk factors such as hypertension (HT) and dyslipidaemia. HT is highly prevalent, and seems to be under-diagnosed and under-treated among patients with RA. In this review, we discuss the mechanisms that may lead to increased blood pressure in such patients, paying particular attention to commonly used drugs for the treatment of RA. We also suggest screening strategies and management algorithms for HT, specific to the RA population, although it is clear that these need to be formally assessed in prospective randomized controlled trials designed specifically for the purpose, which, unfortunately, are currently lacking.

KEY WORDS: Systematic review, Rheumatoid arthritis, Cardiovascular, Hypertension, Inflammation, Physical inactivity, Medication, Non-steroidal anti-inflammatory drugs, Recommendations, Glucocorticosteroids.

Introduction and methods

RA associates with excessive morbidity and mortality from cardiovascular disease (CVD) [1], which may be due to multiple causes [2–6]. Several risk factors, such as hypertension (HT) [7–10], smoking [11–13], dyslipidaemia [14], as defined by National Cholesterol Education Program [15] and insulin resistance [16, 17] are thought to be more prevalent in RA and may be important contributors.

HT is one of the most important modifiable risk factors for the development of CVD in the general population [18]. It affects ~1 billion individuals worldwide [19], ~30% of the adult population in the United States [20, 21] and the United Kingdom [22]. HT may be a very important CVD risk factor in RA. Several studies among patients with RA have demonstrated that it associates with subclinical atherosclerosis [7, 23, 24] and is one of the most significant independent predictors of CVD, with relative risk ranking from 1.49 to 4.3 [1, 25, 26]. Using data from the Framingham Heart Study in the United States and the Third (US) National Health and Nutrition Examination Survey (NHANES III), Singh *et al.* [27] projected that a 20 mmHg increase in systolic blood pressure (SBP) in RA patients would associate with 1572 additional ischaemic heart disease events and 602 additional stroke events over 1 yr. The impact of HT on cardiovascular outcome is thought to be similar among patients with RA to those who do not have RA [28], but since CV mortality is higher in RA patients compared with matched, non-RA controls [29–31], the number of deaths attributed to HT may be higher amongst RA patients.

The reported prevalence of HT among patients with RA varies from 3.8% [32] to 73% [33]. In view of this very wide range, we conducted a systematic review of all studies reporting HT prevalence in patients with RA (Table 1). After taking into

consideration an evidence-based tool for literature searching specifically for RA [34], the databases Medline, Cochrane Library, Cumulative Index to Nursing and Allied Health research database (CINAHL) and Excerpta Medica database (EMBASE) were searched to identify publications from 1990 to 18 November 2007 in English regarding RA and HT. The Medical Subject Heading (MeSH) term 'rheumatoid arthritis' was employed in combination with 'hypertension' and 'blood pressure (BP)'. Initial searches identified 465 articles (i.e. 336 for 'rheumatoid arthritis' and 'hypertension' and 129 articles for 'rheumatoid arthritis' and 'BP'). Full articles were retrieved for assessment if the study fulfilled the following criteria: (i) studying HT prevalence in RA; (ii) studying risk factors for CVD, including HT; or (iii) studying BP changes in response to any intervention in RA, as part of the methodology. The final search revealed 48 articles of which 13 were excluded because of lack of reporting HT prevalence (as a categorical variable) [8, 35–46], 1 because of lack of HT definition [47] and 3 because of high selection bias due to very strict inclusion criteria {new onset coronary artery disease [48] and RA patients starting on DMARDs [49] or biologic agents [50]}. The final 31 studies included are listed in Table 1.

The very wide range of reported prevalence of HT in RA is explained by the different populations assessed, the varied sample sizes and significant differences in the definition of HT used. When using the current definition of HT [51] [i.e. SBP \geq 140 and/or diastolic BP (DBP) \geq 90 and/or the use of anti-hypertensive medication] prevalence of HT in RA in most large studies of unselected, community-based, RA populations lies between 52% and 73% [28, 33]; this appears to depend on the mean age, which ranges from 51 to 66 yrs. In the larger studies of RA samples from secondary care with a similar mean age (56–61.5 yrs), HT prevalence is slightly higher, from 62% to 70.5% [10, 52, 53]. In one of these studies [53], the reported prevalence of HT of 62% would have been higher if patients on lipid-lowering agents and therapy for diabetes had not been excluded. The above figures suggest that the prevalence of HT in the overall RA population is equal to, if not higher than the prevalence observed in the over 65 yrs population of England [22].

Direct evidence as to whether the prevalence of HT is greater among patients with RA than in the general population is contradictory [1, 9, 54, 55]; several studies support an increased prevalence amongst RA patients [7–9, 33, 36, 39, 43, 47], but in some cases this could be due to the higher age of the RA group compared with the controls used [33, 36, 43] or due to the bias arising from comparing RA patients recruited from secondary

¹Department of Rheumatology, Dudley Group of Hospitals NHS Trust, Russells Hall Hospital, Dudley, UK, ²Department of Internal Medicine, School of Medicine, University of Ioannina, Ioannina, Greece, ³Department of Psychology, University of Otago, Dunedin, New Zealand, ⁴Department of Cardiology, Dudley Group of Hospitals NHS Trust, Russells Hall Hospital, Dudley and ⁵ARC Epidemiology Unit, Manchester University, Manchester, UK.

Submitted 2 August 2007; revised version accepted 31 March 2008.

Correspondence to: G. D. Kitas, Department of Rheumatology, Dudley Group of Hospitals NHS Trust, Russells Hall Hospital, Pensnett Road, Dudley, West Midlands DY1 2HQ, UK. E-mail: gd.kitas@dgoh.nhs.uk; g.d.kitas@bham.ac.uk

TABLE 1. Detailed characteristics of the studies included in the systematic review

Reference	Group	HT definition	Group	No.	Age (yrs)	Female (%)	DD	GC (%)	HT (%)	Comments—exclusion criteria (or inclusion criteria where specified)
Dessein <i>et al.</i> [245, 246]	S	≥130/85 and/or on anti-HT	RA	74	57	86.5	11	15	49	Cross-sectional: lipid-lowering agents, anti-DM
Kravounaris <i>et al.</i> [247]	S	≥130/85 mmHg or self-reported anti-HT	RA	200	63	73.5	9.6	24	66 ^{††}	Matched case-control
Chung <i>et al.</i> [33]	C	≥130/85 and/or on anti-HT OR ≥140/90 and/or on anti-HT	E-RA	88	51 [†]	64	<2	53.2	56/39 [†]	Case-control: <18 yr
			L-RA	66	59 [†]	73	20	73/68 [†]		
Dessein <i>et al.</i> [53]	S	≥140/90 and/or on anti-HT	Ct	52	52 [†]	65			55/38 [†]	Cross-sectional: lipid-lowering agents, anti-DM
			RA	92	56	80.4	11	20	62	
Dessein <i>et al.</i> [16]	S	≥140/90 and/or on anti-HT	RA	79	52	83.5	8.5	13	50	Matched case-control: lipid-lowering agents, anti-DM
			OA	39	56				59	
Dessein and Jaffe [248]	S	≥140/90 and/or on anti-HT	RA	21	59	66	6	14	57.1	Cross-sectional: lipid-lowering agents, anti-DM and GC>3 months
Maradit-Kremers <i>et al.</i> [249–251]	C	≥140/90 and/or on anti-HT	RA	603	58	73.1	0	24.5	51.7	Matched case-control: RA incidence cohort
Gonzalez <i>et al.</i> [28]	S	≥140/90 and/or anti-HT	Ct	603	58	73.1			49	Cross-sectional: data source: Outcome of RA Longitudinal Evaluation (ORALE)
del Rincon <i>et al.</i> [52]			RA	631	58	72.3	14.1		62.8	
Roman <i>et al.</i> [23]	S	≥140/90 and/or anti-HT	RA	98	48	98	12	71*	18	Matched case-control: age<18, Creatinine >270 μmol/l, creatinine clearance ≤ 0.5 ml/s, current or recent (3 months) pregnancy
			Ct	98	47	98			22	
La Montagna <i>et al.</i> [252]	S	≥140/90 and/or on anti-HT	RA	45	54	93.3	12.4	51.1	22.2	Matched case-control
Pamuk <i>et al.</i> [253]	S	≥140/90 and/or on anti-HT	C	48	52	88.9			12.5	Matched case-control: atherosclerotic complications haemodialysis, malignancy, infections, DM
			RA	63	51	93.8	6	63.5	30.2	
Gerli <i>et al.</i> [7]	S	≥140/90 and/or on anti-HT	Ct	61	53	88.9			32.4	Case-control: CVD, HF, IHD, CVA, PVD
			RA	101	63	73	11	45.5	37 [†]	
Panoulas <i>et al.</i> [10, 254]	S	≥140/90 and/or on anti-HT	RA	400	61.5	73	10	31.3	70.5	Cross-sectional
Dessein <i>et al.</i> [255]	S	≥140/90 and/or on anti-HT	RA	94	55	66	6	14	51	Cross-sectional: impaired Glc metabolism prior RA onset, lipid- and glucose-lowering agents, hypothyroidism
Dessein <i>et al.</i> [256]	S	≥140/90 and/or on anti-HT	RA	77	54	83.1		14.2	44	Matched case-control: patients with raised ALT, AST or Hep B/Hep C, evidence of autoimmune hepatitis, lipid- or glucose-lowering agents or medication that can affect aminotransferase levels
Van Halm <i>et al.</i> [69]	S	On anti-HT	RA	613	62.6	70.3	9.6	30.3	22.5	Case-control
Singh <i>et al.</i> 2003 [54]	C	On anti-HT	RA	30 million					34	Cross-sectional
			OA						41	
Assous <i>et al.</i> [25]	S	On anti-HT (Q)	RA	239	56	82	11.6	88	34	Retrospective cohort
Wallberg-Jonsson <i>et al.</i> [26]	S	On anti-HT (at any point from baseline to death)	RA	211	52	59.7	<1	47**	32	Retrospective cohort: DD<1 yr at presentation included only
Wallberg-Jonsson <i>et al.</i> [257]	S	On anti-HT (at least for >1 yr)	RA	606	55	68	12.5	53**	29	Retrospective cohort: seropositive RA included only
Erb <i>et al.</i> [68]	S	On anti-HT	RA	150	60.8	65.3	11.2		28	Matched case-control
			Ct	100	61.7	61			24	
Roman <i>et al.</i> [258]	S	Physician's diagnosis	RA	80	45	99			3.8	Case-control
			Ct	105	47	95			4.8	
Wolfe <i>et al.</i> [1]	C	Physician's diagnosis (self-reported Q)	RA	9093	60 [†]	76.9			28.1 [†]	Cross-sectional
			OA	2479	66 [†]				37.4 [†]	
Solomon <i>et al.</i> [55]	C	Physician's diagnosis (self-reported Q)	RA	287	56 [†]	100			31.4	Case-control: data from: Nurses Health Study
			Ct	87019	58 [†]	100			29	
Wolfe and Michaud [259]	C	Physician's diagnosis (self-reported Q)	RA	13171	61	77	14.9	39	47	Cross-sectional
Han <i>et al.</i> [9]	C	ICD-9 401×	RA	28208	52	72.5			31 [†]	Matched case-control: <17, no interaction with health system
			Ct	11232	52	72.5			23.4 [†]	

S: secondary care; C: community; No.: number of participants; Ct: controls; F: female; DD: disease duration; E-RA: early RA; L-RA: long-standing RA; Q: questionnaire; DM: diabetes mellitus; HF: heart failure; IHD: ischaemic heart disease; CVA: cerebrovascular accident; PVD: peripheral vascular disease; ALT: alanine aminotransferase; AST: aspartate aminotransferase; [†]*P* < 0.05, ^{††}past/current use, ^{**}>1 yr use.

care (S) with community-derived controls (C) [8]. The most convincing evidence, comes from the large population study by Han *et al.* [9] on over 28 000 RA patients vs almost 113 000 age-matched controls in which prevalence of HT {defined as International Classification of Diseases 9th Revision, Clinical Modification (ICD-9) code CM 401× [56]} was significantly higher in RA (34% vs 23.4%). The lower overall rates in this, compared with other studies, are clearly due to the significantly different definitions used (physicians' diagnoses) [56] and the younger mean age of the participants.

Threshold BP levels for the diagnosis of HT are $\geq 140/90$ mmHg when using the auscultatory method in the clinic (with BP having been measured on at least 3–6 visits, spaced over a period of 3 months [57]); $\geq 125/80$ mmHg for ambulatory monitoring and $\geq 135/85$ mmHg for self-reported BP monitoring at home [58]. Algorithms for diagnosis and management are predominantly based on clinical measurements, with the higher value used for classification if systolic and diastolic values fall into different categories. According to British Hypertension Society (BHS) guidance, SBP/DBP thresholds in millimetres of mercury are as follows: optimal $<120/<80$, normal $<130/<85$, 'high normal' $130\text{--}139/85\text{--}89$, Grade 1 (mild HT) $140\text{--}159/90\text{--}99$, Grade 2 (moderate HT) $160\text{--}179/100\text{--}109$ and Grade 3 (severe HT) $>180/>110$. Isolated systolic HT is classified as Grade 1 ($140\text{--}159/<90$) or Grade 2 ($>160/<90$). The European Society of Hypertension/European Society of Cardiology (ESH/ESC) guidelines [58] and World Health Organization (WHO) guidelines [59] are virtually identical but the American Joint National Committee (JNC) [60] are slightly different by introducing the term 'pre-hypertension' to encompass normal and 'high normal' BP, and by merging Grades 2 and 3 HT in a single stage. This was based on evidence from the Framingham study [61, 62], which suggested an increased risk of future development of HT amongst individuals in the latter group(s). This classification was not adapted by the ESH/ESC [58] because it was felt that tagging individuals with a term of potentially ominous significance to the layman (i.e. 'pre-hypertension') may lead to an increased number of unnecessary visits and examinations [63].

Despite its high prevalence and the impact of its complications, control of HT is far from adequate both in the general population [60, 64–66] and in RA [10]. The poor control rates in the general population, where only a third of the people with HT have their BP under control [67], is attributed to poor access to health care and medications, and lack of adherence to long-term therapy for an usually asymptomatic condition. In a recent study [10], the rate of controlled HT in RA was significantly lower at 13.2%, than the 21–23% observed in the general population [65]. This is in line with several studies on RA patients, which report prevalence of treated HT between 22–34% [25, 26, 50, 54, 68, 69], a figure much lower than the aforementioned prevalence of HT in RA. Uncontrolled HT associates with premature CVD [20], increased risk for coronary heart disease (CHD) [70, 71], heart failure [23], cerebrovascular disease [71] and peripheral vascular disease (PVD). Left ventricular hypertrophy (LVH), a direct consequence of long-term increases in afterload due to HT, associates with increased incidence of heart failure, ventricular arrhythmias, fatal myocardial infarction (MI) or sudden cardiac death. In the general population, anti-hypertensive therapy has been associated with a 40% reduction in strokes, 20% in MIs and $>50\%$ in heart failure [72]: this emphasizes the importance of optimal BP control in any population, including RA. However, it must be emphasized that currently there are neither randomized controlled trials of the treatment of HT, nor any studies of the magnitude of benefit (if any) of HT control, specifically in patients with RA.

In the following sections, we discuss the multiple factors that may affect BP and its control in RA, particularly inflammation, physical inactivity and drugs. We then present the latest guidelines for the management of HT in the general population and suggest an adaptation for their use in patients with RA.

Inflammation

The first suspicion of an association between low-grade systemic inflammation and HT was raised in studies in the general population. Cross-sectional studies initially demonstrated higher levels of CRP measured by high-sensitivity (hs) assays amongst people with HT [73–78], while a prospective cohort study showed that increased hsCRP levels associated with future risk of developing HT [79]. A study in an asymptomatic sample of the general population [80] suggests that hsCRP associates inversely with large artery elasticity, thus increasing the systolic component of BP. Interestingly, vascular function is abnormal in RA compared with age- and sex-matched controls, with RA patients demonstrating reduced small- and large-artery elasticity and greater systemic vascular resistance [81]. Chronic inflammatory diseases, including RA, have been associated with increased arterial stiffness [32, 82, 83], which may subsequently lead to increased arterial BP [84] and partly explain the high prevalence of HT in RA.

There are several mechanisms by which systemic inflammation (reflected by high CRP levels) may promote the development of HT. High CRP can reduce nitric oxide production in endothelial cells (ECs), resulting in vasoconstriction, increased production of endothelin-1 [85, 86], leucocyte adherence, platelet activation, oxidation and thrombosis [87]. It can also lead to induction of plasminogen activator inhibitor-1 (PAI-1), which has been found raised in people with HT [88, 89], and may contribute to impaired fibrinolysis and atherothrombosis [86, 90]. CRP can also up-regulate the expression of angiotensin type-1 receptor, affect the renin-angiotensin system and further contribute to high BP [91].

However, the inverse has also been proposed. High BP, especially particular patterns of haemodynamic flow with low average but high oscillatory shear stress, are thought to cause greater expression of adhesion molecules and inflammatory genes by EC and initiate the inflammatory cascade in the arterial wall [92], including the production of pro-inflammatory cytokines; these may leak in the systemic circulation and may consequently induce an acute-phase response, including an elevation of CRP [93] (Fig. 1).

The relationship between systemic inflammation and BP has yet to be investigated in any detail in RA. In a recent cross-sectional study [10], there was no significant difference in RA activity and severity between hypertensive and normotensive RA patients. It is interesting to note that long-term (>6 months) oral daily prednisolone of ≥ 7.5 mg (i.e. medium dose according to recent nomenclature [94]), was significantly and independently associated with HT [95]. Given the cross-sectional design of the study, a causal relationship between long-term glucocorticosteroid (GC) and HT cannot be inferred nor can it be assumed that this association simply mirrors an association between severe cases of RA (which are more likely to be treated with steroids) and HT. It remains unknown whether good control of RA disease activity reduces the risk of developing HT in normotensive RA patients or contributes to improved BP control in hypertensives. Prospective longitudinal studies are required to answer these questions.

Physical inactivity

A prospective study of Harvard University male alumni started in the 1960s, demonstrated that the lack of moderately vigorous sports activities, being overweight and a parental history of HT independently increased the risk of developing HT, and this was a strong predictor of premature death [96]. The intensity of physical activity was more important than the quantity of energy output in deterring HT and preventing premature mortality, a finding that was confirmed for men but not for women in another cohort [97]. Several manifestations of RA, such as pain, stiffness and permanent joint damage [98], may compromise the physical activity and fitness levels of these patients compared with people




Fig. 1. Hypertension pathways in RA. Raised BP, particularly patterns of high oscillatory shear stress, initiate the inflammation cascade in the vascular wall which eventually leads to the production of acute-phase reactants from the liver, such as serum amyloid A and CRP. In turn, raised CRP levels lead to reduced NO/endothelin ratio and subsequent vasoconstriction, completing the vicious circle. Physical inactivity and obesity lead to increased propensity for the metabolic syndrome and subsequent BP elevations, whereas several anti-rheumatic medications (Fig. 2) have been associated with HT in RA. Even though the genetic element contribution to BP variation ranges from 30% to 50%, modern molecular approaches have not yet produced and substantiated any causative link between specific genes and HT neither in the general population, nor in RA. NO: nitric oxide; VCAM: vascular cell adhesion molecule; Th: T-helper lymphocytes.

of the same age and gender [99]. Together with fear for disease aggravation and the exercise restriction often recommended (unjustifiably) by rheumatology health professionals, this may account for the inactive, sedentary lifestyle of many RA patients [100]. Physical inactivity in turn may lead to obesity [101], which is highly prevalent [102] and associates independently with HT in RA [10]. Recent studies in the general population suggest that even mild physical exercise, such as walking to work, associates with a lower incidence of HT [103]. This could be recommended to patients with RA as part of the lifestyle changes required to reduce their CVD risk [104] while involvement in structured exercise programmes may provide even more pronounced benefits [99]. In the updated guidelines of the ACR [105] it is clearly stated that dynamic exercise is a safe and effective intervention for patients with RA of recent onset [106], or those with long-standing disease that is active [107] or inactive [108].

Medications

Polypharmacy is a characteristic of RA [109]: many of the drugs used routinely for the treatment of this disease may cause HT or interfere with its effective control. These include the non-selective NSAIDs, cyclo-oxygenase II inhibitors (coxibs), GCs and some

DMARDs. The use of these drugs in RA should always be considered in the context of co-morbid HT and its control.

Non-selective NSAIDs and coxibs

Since non-selective NSAIDs and coxibs are commonly used in RA, their effects on renal function and BP should be considered, especially for patients with pre-existing HT, renal impairment and in the elderly in whom side-effects are most pronounced [110]. In recent years, the absolute and relative contribution of non-selective NSAIDs/coxibs to adverse cardiovascular outcomes has been the subject of much controversy and multiple literature reviews [111–114]. In the present article, we concentrate only on published effects of non-selective NSAIDs/coxibs on BP and its control in people with RA.

A recent systematic review of randomized controlled trials (RCTs) [115] that has studied the effects on BP of non-selective NSAIDs used for at least 4 weeks, showed that there was a significant increase (from baseline to end of study) in mean BP values in ibuprofen (SBP/DBP: 3.54/1.16 mmHg) and indomethacin (SBP/DBP: 2.9/1.58 mmHg) users compared with placebo. Changes from baseline in SBP were positive but not statistically significant for naproxen, sulindac and nabumetone, while diclofenac users showed similar changes to placebo. Compared




Fig. 2. Mechanisms through which drugs used in RA affect BP. PG: prostaglandin.

with placebo, the risk ratio (95% CI) of HT was 2.85 (95% CI 1.44, 5.65; $P=0.003$) in two ibuprofen trials. In a large meta-analysis of 38 placebo-controlled and 12 head-to-head RCTs of non-selective NSAIDs, the latter, overall, have been shown to elevate the supine mean BP by 5 mmHg (95% CI 1.2, 8.7 mm Hg) [116]. This would be sufficient to cause a 15% increase in the risk for IHD and 67% for cerebrovascular accident [117]. In the latter meta-analysis, piroxicam, indomethacin and naproxen had the most marked effect, while aspirin (at anti-inflammatory doses), sulindac and flubiprofen caused the smallest BP elevation. The hypertensive effect of non-selective NSAIDs was more marked in people with HT taking anti-hypertensive therapy than in normotensive people [116].

Low (anti-platelet) doses of aspirin are not associated with BP elevation [118]. Two large prospective cohort studies [119, 120] of women in the 1976 Harvard Nurses' Health Study cohort [121] have shown that compared with non-users, regular non-selective NSAID and interestingly also acetaminophen users were twice as likely to be hypertensive. It was suggested that such analgesic use is responsible for the development of HT in 15% of middle-aged women [120]. However, in a large prospective cohort study of apparently healthy men in the 1982 Harvard Physicians' Health Study, analgesic use (non-selective NSAIDs, acetaminophen and aspirin) was not associated with subsequent HT [122]. As well as being gender-limited, these samples have narrow occupational group inclusion criteria that limit the generalizability, particularly given the effects of work status on subsequent CVD [123]. Even though several non-selective NSAIDs used in the above studies (such as indomethacin or piroxicam) are not any more frequently used in everyday clinical practice, overall data would suggest caution in the use of these agents, especially in the elderly, those who already have HT [124], patients with heart failure [125] or patients with renal impairment [126]. Patients belonging to these groups have increased activation of the renin-angiotensin and sympathetic nervous systems [127] and increased plasma

concentrations of prostaglandin E2 [128, 129], and are therefore more prone to BP increases when treated with non-selective NSAIDs. The concurrent use of non-selective NSAIDs with diuretics, angiotensin-converting enzyme inhibitors (ACE-I) and angiotensin II receptor blockers (ARBs) can be particularly nephrotoxic in the elderly [130]. Therefore, if the above combinations cannot be avoided, they should be accompanied by close clinical and biochemical monitoring. Sulindac was thought to be a better choice in patients at high risk, as in the past it had been shown to have little effect on BP, although this has recently been questioned [110].

Many studies have shown that non-selective NSAIDs attenuate the anti-hypertensive effects of diuretics [116, 131, 132], β -blockers [116, 132], ACE-I [133, 134], ARBs [133] and other vasodilators, such as prazosin [116, 135]. However, non-selective NSAIDs do not have an effect on the BP-decreasing effect of calcium channel blockers (CCBs) [136, 137], which may therefore be an appropriate choice of anti-hypertensive if a non-selective NSAID is necessitated and worsening HT is noted.

The picture is quite similar with regard to coxibs and HT. The UK National Institute of Health and Clinical Excellence (NICE) guidance recommends that these agents should not be used routinely in patients with RA or OA [138]. They should be used in preference to non-selective NSAIDs only in patients at 'high risk' of developing gastrointestinal adverse effects and should be avoided in patients with concomitant CVD [138]. A meta-analysis [139] of 19 RCTs compared the hypertensive effects of coxibs (celecoxib, rofecoxib and etoricoxib) with a non-selective NSAID (naproxen) and placebo. The relative risk for developing HT was higher for coxibs vs non-selective NSAID or placebo but not significantly so. However, with coxib use, there appears to be a disproportionate rise in SBP compared with DBP, which may increase CHD risk [140]. Rofecoxib caused more increase in BP compared with celecoxib or etoricoxib, a result seen in many studies [141–143]. In agreement with these findings are the results

of the most recent meta-analysis of 114 RCTs of coxib use by Zhang *et al.* [144]. Only rofecoxib was associated with HT (RR = 1.55; 95%CI 1.29, 1.85) whereas celecoxib was associated with a slightly lower risk of HT (RR = 0.83; 95%CI 0.71, 0.97), and the other agents (valdecoxib/parecoxib, etoricoxib and lumiracoxib) demonstrated no significant association [144]. A coxib class effect was not evident, as only rofecoxib, but not other coxibs were associated with increased risk of HT. However, in the recently published Multinational Etoricoxib and Diclofenac Arthritis Long-term (MEDAL) trial, participants receiving etoricoxib showed higher rates of discontinuation because of HT [145]. Taken together, this evidence would suggest an association of sulphone coxibs (rofecoxib, etoricoxib) with HT that could be explained by the pro-oxidant effects of these agents, which result to vasoconstriction and subsequent BP elevation [146].

Other studies have focused on non-selective NSAID and coxib effects on ambulatory BP and showed significant increases in SBP by rofecoxib but not celecoxib or naproxen [147]. The proportion of people with controlled HT at baseline who developed ambulatory HT by week 6 of the Sowers *et al.*'s [147] study was greater with rofecoxib (30%) than celecoxib (16%). In two other studies where ACE-I-treated people with HT were evaluated, the effects of celecoxib were comparable with placebo [148], whereas rofecoxib induced significant BP increases over placebo or indomethacin [149].

COX-2 is constitutively expressed in the kidney and expression is up-regulated during salt restriction indicating that COX-2 may be important in regulating renal function [150, 151], particularly in patients with low effective circulating volume, such as those with congestive heart failure, cirrhosis, renal insufficiency or users of diuretics [152]. In a comparative study of celecoxib *vs* naproxen, all participants experienced transient reduction in sodium and potassium excretion. Celecoxib at a dose of 400 mg BD lowered the glomerular filtration rate and renal plasma flow in contrast to naproxen, which had no such effect [153]. However, the Celecoxib Long-Term Arthritis Safety Study (CLASS) [154] in 7968 subjects, 2183 with RA and 5785 with OA, suggested an improved safety profile, regarding the renal effects of celecoxib compared with standard doses of ibuprofen or diclofenac. In cases with mild pre-renal azotemia, significantly fewer participants taking celecoxib exhibited clinically important reductions in renal function compared with diclofenac or ibuprofen. Celecoxib was associated with a similar incidence of HT or oedema to diclofenac, but significantly lower than ibuprofen [154]. However, despite the seemingly safe hypertensive profile of celecoxib, several studies have shown an association of the latter with increased risk of MI, consistent with a class effect for COX-2 specific inhibitors [155–157]. Therefore, use of celecoxib or any other coxib, should ideally be avoided in patients with RA, particularly when CVD factors or established CVD is present [158].

In the general population, concomitant use of low-dose aspirin is present in >20% of all people taking either non-selective NSAIDs or coxibs [159]. The interaction of non-selective NSAIDs/coxibs with low-dose aspirin may be of great importance: there is some evidence to suggest that ibuprofen (but not rofecoxib, diclofenac or acetaminophen) may antagonize the anti-platelet effect of aspirin and limit its cardioprotective effects [160]. There may also be an interaction between naproxen and aspirin, with naproxen inhibiting platelet COX-1 [161]. Pharmacodynamic data indicating an interaction between aspirin and NSAIDs have not translated to a consistent clinical effect in observational studies [162]. It remains unknown, currently, whether anti-platelet doses of aspirin decrease the cardiovascular risks afforded by chronic usage of coxibs or non-selective NSAIDs, although it may potentiate their gastrointestinal side-effects. [159]. For now, according to an FDA advisory [163] patients taking immediate release low-dose aspirin (not enteric coated) and ibuprofen 400 mg should take the latter at least 30 min after aspirin ingestion or at least 8 h before aspirin ingestion to avoid any potential

interaction. Expert recommendation is also that patients should take a non-enteric coated aspirin at least 15–30 min (if chewed) or 2 h (if swallowed) prior to taking an NSAID, and at least 6–8 h after the last dose of ibuprofen or 36–48 h after naproxen [164]. It would be reasonable to suggest that concomitant use of low-dose aspirin and NSAIDs should be avoided if at all possible, and the lowest NSAID dose should be used for the shortest period of time.

In summary, current evidence and common sense would indicate that non-selective NSAIDs and coxibs have largely similar effects on BP control and renal function, and their use in RA should be avoided. These agents should be particularly avoided in conditions with diminished intravascular volume or oedema, such as congestive heart failure, nephrotic syndrome or cirrhosis, and in patients with serum creatinine levels ≥ 2.5 mg/dl [126]. A recent statement from the American Heart Association (AHA) [165] recommends that for patients with known CVD or at risk for IHD and concomitant musculoskeletal symptoms, the least risky medication should be tried first (acetaminophen, acetylsalicylic acid, tramadol, short-term narcotic analgesics) with escalation only if the latter are ineffective (with order of preference: non-COX-2 selective NSAIDs being preferable to NSAIDs with some COX-2 activity, and these being preferable to COX-2 selective NSAIDs) and with realization that effective pain relief may come at the cost of a small but real increase in risk for cardiovascular or cerebrovascular complications. We suggest that if NSAID use is unavoidable, BP (and renal function) should be closely monitored and in cases of developing new or uncontrolled HT with a continuing requirement for non-selective NSAIDs/coxibs, anti-hypertensive treatment should be instituted, possibly with CCBs.

DMARDs

Some of the DMARDs can induce HT, so if RA is diagnosed in a person with HT (or indeed the converse, when a person with established RA develops HT), the choice of DMARD must be carefully considered. The concurrent use of NSAIDs/coxibs and/or steroids may further exacerbate these side-effects.

LEF

LEF-related HT is found in 2–4.7% of people who were prescribed LEF [166–168], in the absence of renal function abnormalities. A small longitudinal study of consecutive patients on stable doses of NSAIDs and corticosteroids [169] revealed significant increases in SBP and DBP occurring within the first 2–4 weeks of LEF therapy.

The mechanisms by which LEF induces HT include an increase in sympathetic drive and displacement of free fraction of concomitant diclofenac or ibuprofen from protein binding, increasing the NSAID's effect on the distribution of renal blood flow and retention of salt and water [170]. The latter is thought to be because the active malononitrilamide metabolite A771726 of LEF, which is formed in the intestinal mucosa and plasma, and is highly protein bound in plasma [171].

The British Society for Rheumatology (BSR) monitoring guidelines [172] recommend fortnightly BP measurements for the first 6 months of LEF therapy and every 2 months thereafter. Though LEF is not contraindicated in HT, other DMARD options should be considered first, while if HT occurs after commencing LEF, anti-hypertensives may be used, but a dose reduction or cessation of therapy may be required if BP control is not attained.

Cyclosporin

Cyclosporin has a number of side-effects including HT and nephrotoxicity [173–178] and is contraindicated in people with abnormal renal function or uncontrolled HT [172].

Several mechanisms have been suggested to explain cyclosporin-induced HT [179, 180], including: increased peripheral vascular resistance as a result of widespread endothelin-related vasoconstriction; impaired vasodilation due to a reduction in nitric oxide levels or suppression of vasodilatory prostaglandins such as prostacyclin; vasoconstriction in the renal circulation resulting in reduced glomerular filtration rate; and increased sodium retention.

After commencing cyclosporin, serum creatinine and BP should be measured fortnightly until the dose has been stable for 3 months, and continued at monthly intervals thereafter. A 30% rise in creatinine requires urgent adjustment or even cessation of therapy [181]. The new discovery of HT while on cyclosporin also requires action. CCBs, other than diltiazem, verapamil and nifedipine (that increase plasma cyclosporin levels), are the drugs of choice in this situation as they antagonize the vasoconstrictive effects and control effectively cyclosporin-induced HT [182]. A reduction in cyclosporin dose or even complete withdrawal may be necessary in cases of resistant HT.

Biologic anti-TNF- α therapies and other DMARDs

The ultimate effect of biologic anti-TNF therapies on cardiovascular risk is not yet known [183], although initial reports appear promising [184]. There are no reports of HT occurring in association with TNF- α blockers, even though in a minority of patients with RA they can cause or exacerbate heart failure [185]. Some DMARDs may have overall favourable effects on CVD risk factors and may therefore be particularly suitable for hypertensive RA patients. HCQ, for example, may be associated with a favourable lipid profile [186–188] and thus eliminate another potential risk factor for HT [189] among people with RA, but this remains to be proven.

GCs

In the general population, therapeutic use of supraphysiological doses of oral GCs (≥ 7.5 mg/day) may associate with increased rates of MI, stroke, heart failure and all-cause mortality [190, 191]. However, unmeasured confounders may be very important in all observational studies of this type, so such results should be seen with great caution.

It is widely held that GC therapy may raise BP in both normotensive and hypertensive people, but good quality evidence for this is limited and the possible mechanisms are not well understood [192, 193]. Endogenous plasma or urine cortisol levels are increased in hypertensive subjects compared with controls, implying potential hypertensive properties of this hormone when administered orally [194, 195] and there is a positive correlation between plasma cortisol and 'white-coat' HT [196]. Most of the reviews on GC side-effects [197–199] or on mechanisms of GC-induced HT [200] are citing very old references [201–203] when discussing the association of HT with exogenous steroid use. The regimes used and the design of these studies do not provide sufficient evidence to support such an association. This lack of evidence is also reported in the recent European League Against Rheumatism (EULAR) recommendations on the management of systemic GC therapy in rheumatic disease [204]; for Recommendation 5, which is referring to BP monitoring, authors state that 'There is no direct evidence from appropriately designed studies to support this proposition (category IV)'. A study from the early 1970s [205] reported that radiologically visible arteriosclerosis in the ankle joint region in RA patients on long-term GC was three times more frequent than in RA patients not receiving GC or in healthy controls. There was no information on GC dose and duration, and no adjustment for age, disease activity or severity. A prospective longitudinal study from the 1980s suggested that in people receiving 'low doses of GCs' (defined in that study as prednisone < 20 mg/day!), significant HT may be explained by age and initial BP rather than by the use of

GCs [206]. However, new thresholds were recently set to define 'low', 'medium' and 'high' dose prednisolone [94]. Participants in that study [206] could belong to the 'modern' low- or medium-dose category and this may have influenced the results. A recent small RCT [207] of GC use in RA, assessed the effects of low-dose prednisolone use (≤ 7.5 mg) on atherosclerosis, endothelial function and risk factors including HT. There was a trend for increased BP among participants who had been treated for at least 4 yrs with prednisolone 7.5 mg (157 ± 29 mmHg) compared with untreated participants (141 ± 28 mmHg, $P = 0.06$). A recent large cross-sectional study in RA [95] showed an increased prevalence of HT in patients on medium-dose (prednisolone ≥ 7.5 mg/day), long-term (> 6 months) GCs, with GCs appearing to be a risk factor for HT independent of other HT risk factors, RA activity or severity, but this finding needs to be confirmed in prospective longitudinal studies.

GC-induced HT is likely to be multifactorial [192, 193]. GCs inhibit extraneuronal uptake and catechol-*O*-methyltransferase (which breaks down noradrenaline) and thus raise noradrenaline levels in the synaptic cleft [208, 209]; they can also increase the number of α -1 adrenergic receptors [210] with the overall effect of increased peripheral vascular sensitivity to adrenergic agonists. GCs can also increase angiotensinogen production from adipose tissue [211] and inhibit prostaglandin production, thus leading to renal sodium retention and increase in blood volume [212].

So far, evidence suggests that GCs, when given in low doses (< 7.5 mg daily prednisolone) probably do not cause clinically significant BP increases [95, 199]; however, RA patients on higher doses of prednisolone should be regularly screened for HT and adequately treated, should the latter occur.

Guidelines for the management of HT

Risk assessment

People with high BP do not necessarily require treatment: in the general population, requirement for intervention is guided by risk stratification. There are several systems for this, including the ESH/ESC [58], which best lend themselves for adaptation in RA (Table 2). In the general population, the link between HT and inflammation has been established in prospective studies [79]. RA *per se*, a chronic disease of high inflammatory state, leads to increased arterial stiffness and carotid wall thickening compared with healthy controls [82, 213, 214] and could therefore be considered an independent risk factor for HT [84], although direct evidence for this is currently lacking. Furthermore, raised CRP levels (≥ 1 mg/dl), which are almost universally observed among RA patients, were considered an additional risk factor in the previous ESC/ESH guidelines [215]. We therefore recommend that, when stratifying the risk of a person with RA who is hypertensive (as per ESH/ESC), physicians should add '+1' in the total sum of risk factors and treat accordingly [58] (Table 2).

It is extremely important to remember that HT should not be addressed in isolation, but must be considered in the context of the overall cardiovascular risk of an individual. This risk can be calculated on the basis of risk factors (including HT), using risk calculators, such as the Joint British Societies calculator [216]. Table 3 outlines an overall therapeutic strategy for a patient, on the basis of their BP levels, 10-yr CVD risk and the presence of relevant comorbidity. RA patients who have 'compelling indications', including those with heart failure, post-MI, high CHD risk, diabetes, stroke or chronic kidney disease [19], require detailed consideration of their anti-hypertensive treatment and further management [140] by a cardiology specialist.

Next, we consider specific aspects to consider when managing HT in people with RA.

TABLE 2. Proposed risk stratification for RA patients according to BP levels and other CVD risk factors

Risk factors and disease history	Normal SBP 120–129 or DBP 80–84	High normal SBP 130–139 or DBP 85–89	Grade 1 SBP 140–159 or DBP 90–99	Grade 2 SBP 160–179 or DBP 100–109	Grade 3 SBP >180 or DBP >110
No other risk factor	Very low added risk	Very low added risk	Moderate added risk	Moderate added risk	Very high added risk
1 risk factor	Very low added risk	Very low added risk	Moderate added risk	Moderate added risk	Very high added risk
2 or more other risk factors	Moderate added risk	High added risk	High added risk	High added risk	High added risk
3 or more other risk factors or TOD or diabetes or MetS	Moderate added risk	High added risk	High added risk	High added risk	Very high added risk
Established CVD or renal disease	High added risk	Very high added risk	Very high added risk	Very high added risk	Very high added risk

RA patients in the shaded cells (bold line cut-off) may require prompt treatment. The dashed line represents cut-off levels for treatment initiation in the general population. Risk factors: age (M >55, F >65), smoking, dyslipidaemia (TC >5 mmol/l or LDL >3 mmol/l or HDL <1 mmol/l M or <1.2 mmol/l F or TG >1.7 mmol/l), fasting plasma glucose 5.6 < FPG < 7 mmol/l, abnormal glucose tolerance test, family history of premature CVD disease (<55 M, <65 F), abdominal obesity (>102 cm M, >88 cm F) Target organ damage (TOD): LVH, U/S evidence of arterial wall thickening (carotid IMT ≥ 0.9 mm or atherosclerotic plaque), carotid-femoral pulse wave velocity >12 m/s, ankle-brachial BP index <0.9, slight increase in serum creatinine (M: 115–133 μmol/l, F: 107–124 μmol/l), low estimated GFR (<60 ml/min/1.73 m²), microalbuminuria. CVD or renal disease: CVA, heart disease, renal disease (diabetic nephropathy, renal impairment, proteinuria), PVD, advanced retinopathy. M: Males; F: Females; TC: Total Cholesterol; TG: triglycerides; LDL: low density lipoprotein; HDL: high density lipoprotein; US: ultrasound; IMT: intima-media thickness; GFR: glomerular filtration rate; CVA: cerebrovascular accident.

TABLE 3. Risk factors and therapeutic targets for CVD prevention among people with RA

CVD risk <20% over the next 10 yrs		CVD risk >20% over the next 10 yrs		Atherosclerotic cardiovascular disease		Diabetes	
Normal BP (<140/90)	Raised BP (≥140/90)	Normal BP (<140/90)	Raised BP (≥140/90)	Normal BP (<130/80)	Raised BP (≥130/80)	Normal BP (<130/80)	Raised BP (≥130/80)
Lifestyle advice Monitoring at every clinical visit (at least 6 monthly) Anti-HT ^a	Lifestyle advice Anti-HT ^a	Lifestyle advice Aspirin	Lifestyle advice Aspirin ^b	Lifestyle advice Aspirin ^b	Lifestyle advice Aspirin ^b	Lifestyle advice Aspirin ^b	Lifestyle advice Aspirin ^b
		Statin ^c Anti-HT ^a	Statin ^c Anti-HT	Statin ^c ACE-I/ARBs ^{d,e} β-Blockers	Statin ^c ACE-I/ARBs β-Blockers Other anti-HT ^g	Statin ^c ACE-I/ARBs ^{e,f}	Statin ^c ACE-I/ARBs ^f Other anti-HT ^g

^aAnti-hypertensive therapy only if required from Table 2 risk stratification (see Table 4 for anti-hypertensive choice). ^bAvoid concomitant use of aspirin and NSAIDs. If no realistic alternative then use lowest dose possible and consider PPI use and *Helicobacter pylori* eradication. Anti-HT: anti-hypertensives; LDL: low density lipoprotein; PPI: protein pump inhibitor. ^cTargets are: total cholesterol <4 mmol/l or 25% reduction and LDL-C <2 mmol/l or 30% reduction. ^dIn people with heart failure or LV dysfunction and consider in people with coronary disease and normal LV function. ^eAccording to Table 2, these people fall into the high/moderate risk category. Therefore BP treatment should be initiated immediately. ACE-I/ARBs are a good initial choice because they also serve the compelling indication, diabetes and atherosclerosis. Consider CCB if the patient is on NSAIDs and unable to discontinue use. ^fIn people with renal dysfunction and microalbuminuria. ^gConsider if BP target not achieved with ACE-I, ARBs, β-blockers.

Lifestyle measures

Lifestyle modifications leading to improved BP control include maintenance of ideal weight (for RA patients to a BMI of <23 kg/m² for people with RA) [102], limiting daily sodium intake to <60 mmol/l (<3.8 g/day) [217] and alcohol consumption to 20–30 g ethanol per day for men and 10–20 g for women [218], smoking cessation [219], dietary patterns based on the Dietary Approaches to Stop HT (DASH) diet (rich in fruits and vegetables and low-fat dairy products, with a reduced content of dietary cholesterol as well as saturated and total fat) [220], high-dose omega-3-polyunsaturated fatty acid supplements [221], increased potassium, calcium and magnesium intake (even though more trials are needed to establish the benefits [222, 223] and last but not the least, regular aerobic exercise [99]. Relevant advice and leaflets can be found through the BHS [224] or British Heart Foundation [225] websites. Some of these lifestyle modifications, e.g. exercise [99] and smoking cessation [226] could also result in clear benefits in control of RA disease activity; there is no existing

evidence that any of the other lifestyle modifications would precipitate adverse outcomes of RA.

Anti-hypertensive drugs

The recent joint NICE and BHS 2006 guidelines for HT management provide an excellent framework for initial drug treatment and stepwise escalation depending on response [227], which could be applied to people with RA either in primary care or in the rheumatology clinic. There are no RCTs to guide HT management specifically among patients with RA. There are, however, several issues that should be taken into account when making treatment decisions for a person with RA who is hypertensive:

- requirement for anti-hypertensive therapy should always be considered in the context of the patient's anti-rheumatic therapy (Table 4);

TABLE 4. A practical approach to the prevention, diagnosis and management of hypertension in patients with RA

- (i) Diagnosis of hypertension
Establish diagnosis of hypertension as per new ESH/ESC guidelines.
- (ii) Identify the cause (essential or secondary)
Screen for current use of non-selective NSAIDs, coxibs, GCs, LEF, cyclosporin.
- (iii) Initial steps in hypertension management
- Remove cause if possible (e.g. NSAID) and re-assess (within 3 months).
 - Provide lifestyle advice and re-assess (within 3–6 months).
 - Decide on requirement for pharmacological therapy and target BP according to risk stratification (as per Table 2).
- (iv) Choice of anti-hypertensives
- If essential hypertension present, use ACE-I/ARBs as first choice anti-hypertensive treatment. Monitor renal function closely, particularly in the elderly or those on nephrotoxic medication (e.g. NSAIDs, cyclosporin).
 - If hypertension is due to non-selective NSAID/coxib, which cannot be withdrawn, use CCB as initial treatment option. If the person is taking GCs consider tapering dose.
 - If insulin resistance is present, avoid β -blockers/diuretics.
 - If RP is present avoid β -blockers and use CCB/ACE-I/ARBs as initial anti-hypertensive treatment.
- (v) Decide if other treatment is required in the context of calculated 10-yr CVD risk or presence of comorbidities (CVD or diabetes)
- Treat as per Table 3. If suspicion of compelling indication (heart failure, post-MI, stroke, etc.) refer to cardiologist.
 - If concomitant use of low-dose aspirin and NSAID necessary, consider using the lowest possible NSAID dose, PPI and *Helicobacter pylori* eradication.
- (vi) BP monitoring requirements in patients with RA
- Systematic BP monitoring every time a patient attends primary or secondary care, or at least every 6 months.
 - If patient started on non-selective NSAIDs/coxibs or GCs, monthly follow-up (either at the primary or secondary setting) for the first 6 months, as above (a) thereafter.
 - If patient initiated on LEF/cyclosporin therapy, fortnightly follow-up for the first 6 months, 2 monthly thereafter.
 - Review of continuing requirement for any anti-rheumatic therapy with BP-raising potential at every clinic visit, irrespective of whether patient is hypertensive or not.
 - If not required: stop
 - If alternatives available: change
 - If continuing requirement: monitor BP as above
 - In case of incident hypertension, monthly monitoring until BP target is reached; thereafter as above (a).

Coxib: cyclo-oxygenase; PPI: protein pump inhibitor.

- ACE-I may be particularly suited to patients with active RA, since they have increased sympathetic activity [228, 229], which may put them in a high renin state [230, 231]. Additionally, these agents may also have a beneficial effect on the course of RA disease *per se*, since they suppress mediators of inflammation, including reactive oxygen species and CRP and they increase the expression of inhibitory κ B (inhibitor of nuclear factor- κ B) [232];
- Increased systemic inflammation (due to over-production of TNF- α and IL-1) leads to down-regulation of various cardiovascular receptors including β -adrenergic [233] and L-type cardiac calcium channel [234] receptors. No such effect has been reported for the angiotensin II type 1 receptors. This may, in theory, influence the efficacy of β -blockers and some CCBs, but the anti-hypertensive effect of ARBs, such as valsartan [235] and losartan [236], in RA patients should remain unaltered. There are no clinical trials in RA to support or refute this possibility;
- Insulin resistance is a common feature in RA patients [237], especially among those on treatment with GCs [238]. In this setting, β -blockers and thiazide diuretics should be avoided, due to their dyslipidaemic and diabetogenic effects. ACE-I, ARBs or CCBs would be preferable for lowering BP [239, 240]; and
- If secondary RP is present then CCBs [241] or ACE-I/ARBs [242, 243] may be the preferred first-line anti-hypertensive treatment, while β -blockers should be avoided.

All of the above suggest that RA patients are more likely to benefit from ACE-I or ARBs, irrespective of their age, with CCBs being also useful in specific situations. An overall approach to the management of HT in patients with RA is presented in Table 4.

Summary

Awareness of the risk of CVD and aggressive screening and treatment strategies, including meticulous BP control [244], have greatly benefited patients with diabetes over the last decade. Over the same time period, awareness of the cardiovascular risk associated with RA has increased, but is yet to be translated in a systematic approach aiming to reduce comorbidity and mortality. Although direct evidence specifically in RA is

lacking, it seems reasonable to suggest that early detection and aggressive management of HT in patients with RA should form part of such a systematic approach. There is a need for specifically designed trials to identify the best approaches for HT management in patients with RA, and quantify the beneficial effect of HT control (if any) on the cardiovascular outcome of this group of people.

Rheumatology key messages

- HT is highly prevalent in RA.
- Use of anti-inflammatory analgesics and disease-modifying drugs with hypertensive potential, physical inactivity and yet to be determined inflammatory pathways, and genetic factors may synergistically lead to HT in patients with RA.
- Aggressive screening and low BP thresholds for anti-hypertensive treatment may be helpful, but these strategies need to be confirmed in prospective studies designed specifically for the purpose.

Acknowledgements

Funding: V.F.P. is supported by a PhD scholarship from Empirikon Institute, Athens, Greece. H.J. is in receipt of an Arthritis Research Campaign (ARC) Educational Research Fellowship (No. 17883). The Dudley Group of Hospitals NHS Trust Department of Rheumatology is in receipt of an infrastructure support grant from the Arthritis Research Campaign (No. 17682).

Disclosure statement: The authors have declared no conflicts of interest.

References

- Wolfe F, Freundlich B, Straus WL. Increase in cardiovascular and cerebrovascular disease prevalence in rheumatoid arthritis. *J Rheumatol* 2003;30:36–40.
- Dessein PH, Joffe BI. When is a patient with rheumatoid arthritis at risk for cardiovascular disease? *J Rheumatol* 2006;33:201–3.
- Kitas GD, Erb N. Tackling ischaemic heart disease in rheumatoid arthritis. *Rheumatology* 2003;42:607–13.
- Panoulas VF, Milionis HJ, Douglas KM *et al.* Association of serum uric acid with cardiovascular disease in rheumatoid arthritis. *Rheumatology* 2007;46:1466–70.
- Panoulas VF, Nikas SN, Smith JP *et al.* The Lymphotoxin 252A>G polymorphism is common and associates with myocardial infarction in patients with rheumatoid

- arthritis. *Ann Rheum Dis*. Advance Access published January 29, 2008, doi: 0:ard.2007.082594v1.
- 6 Mattey DL, Thomson W, Ollier WE *et al*. Association of DRB1 shared epitope genotypes with early mortality in rheumatoid arthritis: results of eighteen years of followup from the early rheumatoid arthritis study. *Arthritis Rheum* 2007;56:1408–16.
 - 7 Gerli R, Sherer Y, Vaudo G *et al*. Early atherosclerosis in rheumatoid arthritis: effects of smoking on thickness of the carotid artery intima media. *Ann NY Acad Sci* 2005;1051:281–90.
 - 8 McEntegart A, Capell HA, Creran D, Rumley A, Woodward M, Lowe GD. Cardiovascular risk factors, including thrombotic variables, in a population with rheumatoid arthritis. *Rheumatology* 2001;40:640–4.
 - 9 Han C, Robinson DW Jr, Hackett MV, Paramore LC, Fraeman KH, Bala MV. Cardiovascular disease and risk factors in patients with rheumatoid arthritis, psoriatic arthritis, and ankylosing spondylitis. *J Rheumatol* 2006;33:2167–72.
 - 10 Panoulas VF, Douglas KM, Milionis HJ *et al*. Prevalence and associations of hypertension and its control in patients with rheumatoid arthritis. *Rheumatology* 2007;46:1477–82.
 - 11 Symmons DP. Epidemiology of rheumatoid arthritis: determinants of onset, persistence and outcome. *Best Pract Res Clin Rheumatol* 2002;16:707–22.
 - 12 Harrison BJ. Influence of cigarette smoking on disease outcome in rheumatoid arthritis. *Curr Opin Rheumatol* 2002;14:93–7.
 - 13 Criswell LA, Merlino LA, Cerhan JR *et al*. Cigarette smoking and the risk of rheumatoid arthritis among postmenopausal women: results from the Iowa Women's Health Study. *Am J Med* 2002;112:465–71.
 - 14 Dessein PH, Joffe BI, Stanwix A, Botha AS, Moomal Z. The acute phase response does not fully predict the presence of insulin resistance and dyslipidemia in inflammatory arthritis. *J Rheumatol* 2002;29:462–6.
 - 15 Sempos CT, Cleeman JI, Carroll MD *et al*. Prevalence of high blood cholesterol among US adults. An update based on guidelines from the second report of the National Cholesterol Education Program Adult Treatment Panel. *J Am Med Assoc* 1993;269:3009–14.
 - 16 Dessein PH, Stanwix AE, Joffe BI. Cardiovascular risk in rheumatoid arthritis versus osteoarthritis: acute phase response related decreased insulin sensitivity and high-density lipoprotein cholesterol as well as clustering of metabolic syndrome features in rheumatoid arthritis. *Arthritis Res* 2002;4:R5.
 - 17 Despres JP, Lamarche B, Mauriege P *et al*. Hyperinsulinemia as an independent risk factor for ischemic heart disease. *N Engl J Med* 1996;334:952–7.
 - 18 Yusuf S, Hawken S, Ounpuu S *et al*. Effect of potentially modifiable risk factors associated with myocardial infarction in 52 countries (the INTERHEART study): case-control study. *Lancet* 2004;364:937–52.
 - 19 Kearney PM, Whelton M, Reynolds K, Muntner P, Whelton PK, He J. Global burden of hypertension: analysis of worldwide data. *Lancet* 2005;365:217–23.
 - 20 Hajjar J, Kotchen TA. Trends in prevalence, awareness, treatment, and control of hypertension in the United States, 1988–2000. *J Am Med Assoc* 2003;290:199–206.
 - 21 Fields LE, Burt VL, Cutler JA, Hughes J, Roccella EJ, Sorlie P. The burden of adult hypertension in the United States 1999 to 2000: a rising tide. *Hypertension* 2004;44:398–404.
 - 22 Health survey for England, 2003. http://www.dh.gov.uk/en/Publicationsandstatistics/Publications/PublicationsStatistics/DH_4098712 (18 April 2008, date last accessed).
 - 23 Roman MJ, Moeller E, Davis A *et al*. Preclinical carotid atherosclerosis in patients with rheumatoid arthritis. *Ann Intern Med* 2006;144:249–56.
 - 24 Dessein PH, Tobias M, Veller M. Metabolic syndrome and subclinical atherosclerosis in rheumatoid arthritis. *J Rheumatol* 2006;33:2425–32.
 - 25 Assou N, Touze E, Meune C, Kahan A, Allanore Y. Cardiovascular disease in rheumatoid arthritis: single-center hospital-based cohort study in France. *Joint Bone Spine* 2007;74:66–72.
 - 26 Wallberg-Jonsson S, Johansson H, Ohman ML, Rantapaa-Dahlqvist S. Extent of inflammation predicts cardiovascular disease and overall mortality in seropositive rheumatoid arthritis. A retrospective cohort study from disease onset. *J Rheumatol* 1999;26:2562–71.
 - 27 Singh G, Miller JD, Huse DM, Pettitt D, D'Agostino RB, Russell MW. Consequences of increased systolic blood pressure in patients with osteoarthritis and rheumatoid arthritis. *J Rheumatol* 2003;30:714–9.
 - 28 Gonzalez A, Maradit KH, Crowson CS *et al*. Do cardiovascular risk factors confer the same risk for cardiovascular outcomes in rheumatoid arthritis patients as in non-rheumatoid arthritis patients? *Ann Rheum Dis* 2008;67:64–9.
 - 29 Reilly PA, Cosh JA, Maddison PJ, Rasker JJ, Silman AJ. Mortality and survival in rheumatoid arthritis: a 25 year prospective study of 100 patients. *Ann Rheum Dis* 1990;49:363–9.
 - 30 Wolfe F, Mitchell DM, Sibley J, Fries JF. The mortality of rheumatoid arthritis. *Arthritis Rheum* 1994;37:481–94.
 - 31 Wallberg-Jonsson S, Ohman ML, Rantapaa-Dahlqvist S. Cardiovascular morbidity and mortality in patients with seropositive RA in Northern Sweden. *J Rheumatol* 1997;24:445–51.
 - 32 Roman MJ, Devereux RB, Schwartz JE *et al*. Arterial stiffness in chronic inflammatory diseases. *Hypertension* 2005;46:194–9.
 - 33 Chung CP, Oesser A, Solus JF *et al*. Prevalence of the metabolic syndrome is increased in rheumatoid arthritis and is associated with coronary atherosclerosis. *Atherosclerosis* 2008;196:756–63.
 - 34 McGowan J. Literature searching. In: Tugwell P, ed. Evidence-based rheumatology. London, UK: BMJ Publishing Group, 2005;3–18.
 - 35 Tanaka K, Inaba M, Goto H *et al*. Paraarticular trabecular bone loss at the ultradistal radius and increased arterial stiffening in postmenopausal patients with rheumatoid arthritis. *J Rheumatol* 2006;33:652–8.
 - 36 del Rincon ID, Williams K, Stern MP, Freeman GL, Escalante A. High incidence of cardiovascular events in a rheumatoid arthritis cohort not explained by traditional cardiac risk factors. *Arthritis Rheum* 2001;44:2737–45.
 - 37 Nagata-Sakurai M, Inaba M, Goto H *et al*. Inflammation and bone resorption as independent factors of accelerated arterial wall thickening in patients with rheumatoid arthritis. *Arthritis Rheum* 2003;48:3061–7.
 - 38 Dekkers JC, Geenen R, Evers AW, Kraaijaat FW, Bijlsma JW, Godaert GL. Biopsychosocial mediators and moderators of stress-health relationships in patients with recently diagnosed rheumatoid arthritis. *Arthritis Rheum* 2001;45:307–16.
 - 39 Alkaabi JK, Ho M, Levison R, Pullar T, Belch JJ. Rheumatoid arthritis and macrovascular disease. *Rheumatology* 2003;42:292–7.
 - 40 Del R, I, Williams K, Stern MP, Freeman GL, O'Leary DH, Escalante A. Association between carotid atherosclerosis and markers of inflammation in rheumatoid arthritis patients and healthy subjects. *Arthritis Rheum* 2003;48:1833–40.
 - 41 Del R, I, Haas RW, Pogossian S, Escalante A. Lower limb arterial incompressibility and obstruction in rheumatoid arthritis. *Ann Rheum Dis* 2005;64:425–32.
 - 42 Veldhuijzen van Zanten JJ, Ring C, Carroll D, Kitas GD. Increased C reactive protein in response to acute stress in patients with rheumatoid arthritis. *Ann Rheum Dis* 2005;64:1299–304.
 - 43 Dessein PH, Joffe BI, Singh S. Biomarkers of endothelial dysfunction, cardiovascular risk factors and atherosclerosis in rheumatoid arthritis. *Arthritis Res Ther* 2005;7:R634–43.
 - 44 Maki-Petaja KM, Hall FC, Booth AD *et al*. Rheumatoid arthritis is associated with increased aortic pulse-wave velocity, which is reduced by anti-tumor necrosis factor-alpha therapy. *Circulation* 2006;114:1185–92.
 - 45 Maki-Petaja KM, Booth AD, Hall FC *et al*. Ezetimibe and simvastatin reduce inflammation, disease activity, and aortic stiffness and improve endothelial function in rheumatoid arthritis. *J Am Coll Cardiol* 2007;50:852–8.
 - 46 Pedersen LM, Nordin H, Svensson B, Bliddal H. Microalbuminuria in patients with rheumatoid arthritis. *Ann Rheum Dis* 1995;54:189–92.
 - 47 Hakoda M, Oiwa H, Kasagi F *et al*. Mortality of rheumatoid arthritis in Japan: a longitudinal cohort study. *Ann Rheum Dis* 2005;64:1451–5.
 - 48 Warrington KJ, Kent PD, Frye RL *et al*. Rheumatoid arthritis is an independent risk factor for multi-vessel coronary artery disease: a case control study. *Arthritis Res Ther* 2005;7:R984–91.
 - 49 Gordon MM, Thomson EA, Madhok R, Capell HA. Can intervention modify adverse lifestyle variables in a rheumatoid population? Results of a pilot study. *Ann Rheum Dis* 2002;61:66–9.
 - 50 Hyrich K, Symmons D, Watson K, Silman A. Baseline comorbidity levels in biologic and standard DMARD treated patients with rheumatoid arthritis: results from a national patient register. *Ann Rheum Dis* 2006;65:895–8.
 - 51 Williams B, Poulter NR, Brown MJ *et al*. Guidelines for management of hypertension: report of the fourth working party of the British Hypertension Society, 2004-BHS IV. *J Hum Hypertens* 2004;18:139–85.
 - 52 Del Rincon I, Freeman GL, Haas RW, O'Leary DH, Escalante A. Relative contribution of cardiovascular risk factors and rheumatoid arthritis clinical manifestations to atherosclerosis. *Arthritis Rheum* 2005;52:3413–23.
 - 53 Dessein PH, Joffe BI, Stanwix AE, Christian BF, Veller M. Glucocorticoids and insulin sensitivity in rheumatoid arthritis. *J Rheumatol* 2004;31:867–74.
 - 54 Singh G, Miller JD, Huse DM, Pettitt D, D'Agostino RB, Russell MW. Consequences of increased systolic blood pressure in patients with osteoarthritis and rheumatoid arthritis. *J Rheumatol* 2003;30:714–9.
 - 55 Solomon DH, Curhan GC, Rimm EB, Cannuscio CC, Karlson EW. Cardiovascular risk factors in women with and without rheumatoid arthritis. *Arthritis Rheum* 2004;50:3444–9.
 - 56 International Classification of Diseases, Ninth Revision, Clinical Modification-9 (ICD-9-CM), hypertensive disease. <http://icd9cm.chrisendres.com/index.php?action=child&recordid=4066> (18 April 2008, date last accessed).
 - 57 Hartley RM, Velez R, Morris RW, D'Souza MF, Heller RF. Confirming the diagnosis of mild hypertension. *Br Med J (Clin Res Ed)* 1983;286:287–9.
 - 58 Mancia G, De Backer G, Dominiczak A *et al*. 2007 ESH-ESC Practice Guidelines for the Management of Arterial Hypertension: ESH-ESC Task Force on the Management of Arterial Hypertension. *J Hypertens* 2007;25:1751–62.
 - 59 Whitworth JA. 2003 World Health Organization (WHO)/International Society of Hypertension (ISH) statement on management of hypertension. *J Hypertens* 2003;21:1983–92.
 - 60 Chobanian AV, Bakris GL, Black HR *et al*. Seventh report of the joint national committee on prevention, detection, evaluation, and treatment of high blood pressure. *Hypertension* 2003;42:1206–52.
 - 61 Vasan RS, Beiser A, Seshadri S *et al*. Residual lifetime risk for developing hypertension in middle-aged women and men: The Framingham Heart Study. *J Am Med Assoc* 2002;287:1003–10.
 - 62 Vasan RS, Larson MG, Leip EP, Kannel WB, Levy D. Assessment of frequency of progression to hypertension in non-hypertensive participants in the Framingham Heart Study: a cohort study. *Lancet* 2001;358:1682–6.
 - 63 Mancia G, Grassi G. European, American and British Guidelines: similarities and differences. In: Black HR, Elliot WJ, eds. Hypertension. A companion to Braunwald's heart diseases. Amsterdam: Saunders-Elsevier, 2007;571–5.
 - 64 Oliveria SA, Lapuerta P, McCarthy BD, L'Italien GJ, Berlowitz DR, Asch SM. Physician-related barriers to the effective management of uncontrolled hypertension. *Arch Intern Med* 2002;162:413–20.
 - 65 Primates P, Poulter NR. Improvement in hypertension management in England: results from the health survey for England 2003. *J Hypertens* 2006;24:1187–92.
 - 66 Luepker RV, Arnett DK, Jacobs DR Jr *et al*. Trends in blood pressure, hypertension control, and stroke mortality: the Minnesota Heart Survey. *Am J Med* 2006;119:42–9.

- 67 Wang TJ, Vasan RS. Epidemiology of uncontrolled hypertension in the United States. *Circulation* 2005;112:1651–62.
- 68 Erb N, Pace AV, Douglas KM, Banks MJ, Kitaz GD. Risk assessment for coronary heart disease in rheumatoid arthritis and osteoarthritis. *Scand J Rheumatol* 2004;33:293–9.
- 69 van Halbeek H, Nurmohamed MT, Twisk JW, Dijkman BA, Voskuyl AE. Disease-modifying antirheumatic drugs are associated with a reduced risk for cardiovascular disease in patients with rheumatoid arthritis: a case control study. *Arthritis Res Ther* 2006;8:R151.
- 70 MacMahon S, Peto R, Cutler J *et al.* Blood pressure, stroke, and coronary heart disease. Part 1. Prolonged differences in blood pressure: prospective observational studies corrected for the regression dilution bias. *Lancet* 1990;335:765–74.
- 71 Prospective Studies Collaboration. Age specific relevance of usual blood pressure to vascular mortality: a meta-analysis of individual data for one million adults in 61 prospective studies. *Lancet* 2002;360:1903–13.
- 72 Neal B, MacMahon S, Chapman N. Effects of ACE inhibitors, calcium antagonists, and other blood-pressure-lowering drugs: results of prospectively designed overviews of randomised trials. *Blood Pressure Lowering Treatment Trialists' Collaboration*. *Lancet* 2000;356:1955–64.
- 73 Ford ES, Giles WH. Serum C-reactive protein and fibrinogen concentrations and self-reported angina pectoris and myocardial infarction: findings from national health and nutrition examination survey III. *J Clin Epidemiol* 2000;53:95–102.
- 74 Rohde LE, Hennekens CH, Ridker PM. Survey of C-reactive protein and cardiovascular risk factors in apparently healthy men. *Am J Cardiol* 1999;84:1018–22.
- 75 Bermudez EA, Rifai N, Buring J, Manson JE, Ridker PM. Interrelationships among circulating interleukin-6, C-reactive protein, and traditional cardiovascular risk factors in women. *Arterioscler Thromb Vasc Biol* 2002;22:1668–73.
- 76 Chae CU, Lee RT, Rifai N, Ridker PM. Blood pressure and inflammation in apparently healthy men. *Hypertension* 2001;38:399–403.
- 77 Bautista LE, Vera LM, Arenas IA, Gamarra G. Independent association between inflammatory markers (C-reactive protein, interleukin-6, and TNF-alpha) and essential hypertension. *J Hum Hypertens* 2005;19:149–54.
- 78 Bautista LE, Lopez-Jaramillo P, Vera LM, Casas JP, Otero AP, Guaracao AI. Is C-reactive protein an independent risk factor for essential hypertension? *J Hypertens* 2001;19:857–61.
- 79 Sesso HD, Buring JE, Rifai N, Blake GJ, Gaziano JM, Ridker PM. C-reactive protein and the risk of developing hypertension. *J Am Med Assoc* 2003;290:2945–51.
- 80 Duprez DA, Somasundaram PE, Sigurdsson G, Hoke L, Florea N, Cohn JN. Relationship between C-reactive protein and arterial stiffness in an asymptomatic population. *J Hum Hypertens* 2005;19:515–9.
- 81 Wong M, Toh L, Wilson A *et al.* Reduced arterial elasticity in rheumatoid arthritis and the relationship to vascular disease risk factors and inflammation. *Arthritis Rheum* 2003;48:81–9.
- 82 Klocke R, Cockcroft JR, Taylor GJ, Hall IR, Blake DR. Arterial stiffness and central blood pressure, as determined by pulse wave analysis, in rheumatoid arthritis. *Ann Rheum Dis* 2003;62:414–8.
- 83 Arosio E, De Marchi S, Rigoni A, Prior M, Delva P, Lechi A. Forearm haemodynamics, arterial stiffness and microcirculatory reactivity in rheumatoid arthritis. *J Hypertens* 2007;25:1273–8.
- 84 Franklin SS. Arterial stiffness and hypertension: a two-way street? *Hypertension* 2005;45:349–51.
- 85 Verma S, Li SH, Badiwala MV *et al.* Endothelin antagonism and interleukin-6 inhibition attenuate the proatherogenic effects of C-reactive protein. *Circulation* 2002;105:1890–6.
- 86 Devaraj S, Xu DY, Jialal I. C-reactive protein increases plasminogen activator inhibitor-1 expression and activity in human aortic endothelial cells: implications for the metabolic syndrome and atherothrombosis. *Circulation* 2003;107:398–404.
- 87 Verma S, Anderson TJ. The ten most commonly asked questions about endothelial function in cardiology. *Cardiol Rev* 2001;9:250–2.
- 88 Srikanth N, Brown NJ, Hopkins PN *et al.* PAI-1 in human hypertension: relation to hypertensive groups. *Am J Hypertens* 2002;15:683–90.
- 89 Poli KA, Tofler GH, Larson MG *et al.* Association of blood pressure with fibrinolytic potential in the Framingham offspring population. *Circulation* 2000;101:264–9.
- 90 Dawson S, Henney A. The status of PAI-1 as a risk factor for arterial and thrombotic disease: a review. *Atherosclerosis* 1992;95:105–17.
- 91 Wang CH, Li SH, Weisel RD *et al.* C-reactive protein upregulates angiotensin type 1 receptors in vascular smooth muscle. *Circulation* 2003;107:1783–90.
- 92 Dai G, Kaazempur-Mofrad MR, Natarajan S *et al.* Distinct endothelial phenotypes evoked by arterial waveforms derived from atherosclerosis-susceptible and -resistant regions of human vasculature. *Proc Natl Acad Sci USA* 2004;101:14871–6.
- 93 Hansson GK. Inflammation, atherosclerosis, and coronary artery disease. *N Engl J Med* 2005;352:1685–95.
- 94 Buttgerit F, da Silva JA, Boers M *et al.* Standardised nomenclature for glucocorticoid dosages and glucocorticoid treatment regimens: current questions and tentative answers in rheumatology. *Ann Rheum Dis* 2002;61:718–22.
- 95 Panoulas VF, Douglas KM, Stavropoulos-Kalinoglou A *et al.* Long-term exposure to medium-dose glucocorticoid therapy associates with hypertension in patients with rheumatoid arthritis. *Rheumatology* 2008;47:72–5.
- 96 Paffenbarger RS Jr, Lee IM. Intensity of physical activity related to incidence of hypertension and all-cause mortality: an epidemiological view. *Blood Press Monit* 1997;2:115–23.
- 97 Haapanen N, Miilunpalo S, Vuori I, Oja P, Pasanen M. Association of leisure time physical activity with the risk of coronary heart disease, hypertension and diabetes in middle-aged men and women. *Int J Epidemiol* 1997;26:739–47.
- 98 Lee DM, Weinblatt ME. Rheumatoid arthritis. *Lancet* 2001;358:903–11.
- 99 Metsios GS, Stavropoulos-Kalinoglou A, Veldhuijzen van Zanten JJ *et al.* Rheumatoid arthritis, cardiovascular disease and physical exercise: a systematic review. *Rheumatology* 2008;47:239–48.
- 100 Munneke M, de Jong Z, Zwinderman AH *et al.* High intensity exercise or conventional exercise for patients with rheumatoid arthritis? Outcome expectations of patients, rheumatologists, and physiotherapists. *Ann Rheum Dis* 2004;63:804–8.
- 101 Fox KR, Hillsdon M. Physical activity and obesity. *Obes Rev* 2007;8(Suppl 1):115–21.
- 102 Stavropoulos-Kalinoglou A, Metsios GS, Koutedakis Y *et al.* Redefining overweight and obesity in rheumatoid arthritis patients. *Ann Rheum Dis* 2007;66:1316–21.
- 103 Hayashi T, Tsumura K, Suematsu C, Okada K, Fujii S, Endo G. Walking to work and the risk for hypertension in men: the Osaka Health Survey. *Ann Intern Med* 1999;131:21–6.
- 104 John H, Hale ED, Treharne GJ, Kitaz GD. Patient education on cardiovascular aspects of rheumatoid disease: an unmet need. *Rheumatology* 2007;46:1513–6.
- 105 Guidelines for the management of rheumatoid arthritis: 2002 Update. *Arthritis Rheum* 2002;46:328–46.
- 106 Hakkinen A, Hakkinen K, Hannonen P. Effects of strength training on neuromuscular function and disease activity in patients with recent-onset inflammatory arthritis. *Scand J Rheumatol* 1994;23:237–42.
- 107 van den Ende CH, Breedveld FC, Le Cessie S, Dijkman BA, de Mug AW, Hazes JM. Effect of intensive exercise on patients with active rheumatoid arthritis: a randomised clinical trial. *Ann Rheum Dis* 2000;59:615–21.
- 108 van den Ende CH, Hazes JM, Le Cessie S *et al.* Comparison of high and low intensity training in well controlled rheumatoid arthritis. Results of a randomised clinical trial. *Ann Rheum Dis* 1996;55:798–805.
- 109 Treharne GJ, Douglas KM, Iwaszko J *et al.* Polypharmacy among people with rheumatoid arthritis: the role of age, disease duration and comorbidity. *Musculoskel Care* 2007;5:175–90.
- 110 Pope JE. Hypertension, NSAID, and lessons learned. *J Rheumatol* 2004;31:1035–7.
- 111 White WB. Cardiovascular risk, hypertension, and NSAIDs. *Curr Rheumatol Rep* 2007;9:36–43.
- 112 Bresalier RS, Sandler RS, Quan H *et al.* Cardiovascular events associated with rofecoxib in a colorectal adenoma chemoprevention trial. *N Engl J Med* 2005;352:1092–102.
- 113 Psaty BM, Weiss NS. NSAID trials and the choice of comparators—questions of public health importance. *N Engl J Med* 2007;356:328–30.
- 114 Fitzgerald GA. Coxibs and cardiovascular disease. *N Engl J Med* 2004;351:1709–11.
- 115 Morrison A, Ramey DR, van Adelsberg J, Watson DJ. Systematic review of trials of the effect of continued use of oral non-selective NSAIDs on blood pressure and hypertension. *Curr Med Res Opin* 2007;23:2395–404.
- 116 Johnson AG, Nguyen TV, Day RO. Do nonsteroidal anti-inflammatory drugs affect blood pressure? A meta-analysis. *Ann Intern Med* 1994;121:289–300.
- 117 Collins R, Peto R, MacMahon S *et al.* Blood pressure, stroke, and coronary heart disease. Part 2. Short-term reductions in blood pressure: overview of randomised drug trials in their epidemiological context. *Lancet* 1990;335:827–38.
- 118 Zanchetti A, Hansson L, Leonetti G *et al.* Low-dose aspirin does not interfere with the blood pressure-lowering effects of antihypertensive therapy. *J Hypertens* 2002;20:1015–22.
- 119 Curhan GC, Willett WC, Rosner B, Stampfer MJ. Frequency of analgesic use and risk of hypertension in younger women. *Arch Intern Med* 2002;162:2204–8.
- 120 Dedier J, Stampfer MJ, Hankinson SE, Willett WC, Speizer FE, Curhan GC. Nonnarcotic analgesic use and the risk of hypertension in US women. *Hypertension* 2002;40:604–8.
- 121 Nurses' Health Study cohort <http://www.channing.harvard.edu/nhs/history/index.shtml> (18 April 2008, date last accessed).
- 122 Kurth T, Hennekens CH, Sturmer T *et al.* Analgesic use and risk of subsequent hypertension in apparently healthy men. *Arch Intern Med* 2005;165:1903–9.
- 123 De Vogli R, Ferrie JE, Chandola T, Kivimaki M, Marmot MG. Unfairness and health: evidence from the Whitehall II Study. *J Epidemiol Community Health* 2007;61:513–8.
- 124 Houston MC. Nonsteroidal anti-inflammatory drugs and antihypertensives. *Am J Med* 1991;90:42S–7S.
- 125 Tobias D. Risk of admission for CHF is higher for rofecoxib and nonselective NSAID users. Commentary: Cyclo-oxygenase-2 inhibitors versus nonselective, nonsteroidal anti-inflammatory drugs and congestive heart failure outcomes in elderly patients: a population-based cohort study. *Consult Pharm* 2004;19:1038–42.
- 126 Whelton A. Renal and related cardiovascular effects of conventional and COX-2-specific NSAIDs and non-NSAID analgesics. *Am J Ther* 2000;7:63–74.
- 127 Fu Q, Zhang R, Witkowski S *et al.* Persistent sympathetic activation during chronic antihypertensive therapy: a potential mechanism for long term morbidity? *Hypertension* 2005;45:513–21.
- 128 London GM, Hornych A, Safar ME, Levenson JA, Simon AC. Plasma prostaglandins PGE2 and PGF2 alpha, total effective vascular compliance and renal plasma flow in essential hypertension. *Nephron* 1982;32:118–24.
- 129 Pope JE, Anderson JJ, Felson DT. A meta-analysis of the effects of nonsteroidal anti-inflammatory drugs on blood pressure. *Arch Intern Med* 1993;153:477–84.
- 130 Adhiaman V, Asghar M, Oke A, White AD, Shah IU. Nephrotoxicity in the elderly due to co-prescription of angiotensin converting enzyme inhibitors and nonsteroidal anti-inflammatory drugs. *J R Soc Med* 2001;94:512–4.
- 131 Klassen D, Goodfriend TL, Schuna AA, Young DY, Peterson CA. Assessment of blood pressure during treatment with naproxen or ibuprofen in hypertensive patients treated with hydrochlorothiazide. *J Clin Pharmacol* 1993;33:971–8.

- 132 Wong DG, Spence JD, Lamki L, Freeman D, McDonald JW. Effect of non-steroidal anti-inflammatory drugs on control of hypertension by beta-blockers and diuretics. *Lancet* 1986;1:997-1001.
- 133 Fogari R, Zoppi A, Carretta R, Veglio F, Salvetti A. Effect of indomethacin on the antihypertensive efficacy of valsartan and lisinopril: a multicentre study. *J Hypertens* 2002;20:1007-14.
- 134 Salvetti A, Pedrinelli R, Magagna A, Ugenti P. Differential effects of selective and nonselective prostaglandin synthesis inhibition on the pharmacological responses to captopril in patients with essential hypertension. *Clin Sci* 1982;63:261-3.
- 135 Rubin P, Jackson G, Blaschke T. Studies on the clinical pharmacology of prazosin. The influences of indomethacin and of propranolol on the action and disposition of prazosin. *Br J Clinical Pharmacol* 1980;10:33-9.
- 136 Klassen DK, Kane LH, Young DY, Peterson CA. Assessment of blood pressure during naproxen therapy in hypertensive patients treated with nifedipine. *Am J Hypertens* 1995;8:146-53.
- 137 Morgan T, Anderson A. Interaction of indomethacin with felodipine and enalapril. *J Hypertens Suppl* 1993;11:S338-9.
- 138 NICE guidelines regarding the use of COX-2 in RA. <http://www.nice.org.uk/guidance/index.jsp?action=byID&o=11687> (18 April 2008, date last accessed).
- 139 Aw TJ, Haas SJ, Liew D, Krum H. Meta-analysis of cyclooxygenase-2 inhibitors and their effects on blood pressure. *Arch Intern Med* 2005;165:490-6.
- 140 Franklin SS, Khan SA, Wong ND, Larson MG, Levy D. Is pulse pressure useful in predicting risk for coronary heart disease? The Framingham Heart Study. *Circulation* 1999;100:354-60.
- 141 Whelton A, White WB, Bello AE, Puma JA, Fort JG. Effects of celecoxib and rofecoxib on blood pressure and edema in patients > or =65 years of age with systemic hypertension and osteoarthritis. *Am J Cardiol* 2002;90:959-63.
- 142 Wolfe F, Zhao S, Pettitt D. Blood pressure destabilization and edema among 8538 users of celecoxib, rofecoxib, and nonselective nonsteroidal antiinflammatory drugs (NSAID) and nonusers of NSAID receiving ordinary clinical care. *J Rheumatol* 2004;31:1143-51.
- 143 Whelton A, Fort JG, Puma JA, Normandin D, Bello AE, Verburg KM. Cyclooxygenase-2-specific inhibitors and cardiorenal function: a randomized, controlled trial of celecoxib and rofecoxib in older hypertensive osteoarthritis patients. *Am J Ther* 2001;8:85-95.
- 144 Zhang J, Ding EL, Song Y. Adverse effects of cyclooxygenase 2 inhibitors on renal and arrhythmia events: meta-analysis of randomized trials. *J Am Med Assoc* 2006;296:1619-32.
- 145 Cannon CP, Curtis SP, FitzGerald GA *et al.* Cardiovascular outcomes with etoricoxib and diclofenac in patients with osteoarthritis and rheumatoid arthritis in the multinational etoricoxib and diclofenac arthritis long-term (MEDAL) programme: a randomised comparison. *Lancet* 2006;368:1771-81.
- 146 Ruschitzka F. Painful lessons: COX-2 inhibitors, NSAIDs, and hypertension. *Curr Hypertens Rep* 2007;9:41-4.
- 147 Sowers JR, White WB, Pitt B *et al.* The effects of cyclooxygenase-2 inhibitors and nonsteroidal anti-inflammatory therapy on 24-hour blood pressure in patients with hypertension, osteoarthritis, and type 2 diabetes mellitus. *Arch Intern Med* 2005;165:161-8.
- 148 White WB, Kent J, Taylor A, Verburg KM, Lefkowitz JB, Whelton A. Effects of celecoxib on ambulatory blood pressure in hypertensive patients on ACE inhibitors. *Hypertension* 2002;39:929-34.
- 149 Merck Research Laboratories. FDA Advisory Committee background information, cardiovascular-renal safety review. Rofecoxib NDA 21-042. 2001;57-9.
- 150 Harris RC, McKanna JA, Akai Y, Jacobson HR, Dubois RN, Breyer MD. Cyclooxygenase-2 is associated with the macula densa of rat kidney and increases with salt restriction. *J Clin Invest* 1994;94:2504-10.
- 151 Komhoff M, Grone HJ, Klein T, Seyberth HW, Nusing RM. Localization of cyclooxygenase-1 and -2 in adult and fetal human kidney: implication for renal function. *Am J Physiol* 1997;272:F460-8.
- 152 Sundry JS. COX-2 inhibitors in rheumatoid arthritis. *Curr Rheumatol Rep* 2001;3:86-91.
- 153 Rossat J, Maillard M, Nussberger J, Brunner HR, Burnier M. Renal effects of selective cyclooxygenase-2 inhibition in normotensive salt-depleted subjects. *Clin Pharmacol Ther* 1999;66:76-84.
- 154 Whelton A, Lefkowitz JL, West CR, Verburg KM. Cardiorenal effects of celecoxib as compared with the nonsteroidal anti-inflammatory drugs diclofenac and ibuprofen. *Kidney Int* 2006;70:1495-502.
- 155 Caldwell B, Aldington S, Weatherall M, Shirlcliffe P, Beasley R. Risk of cardiovascular events and celecoxib: a systematic review and meta-analysis. *J R Soc Med* 2006;99:132-40.
- 156 Mukherjee D, Nissen SE, Topol EJ. Risk of cardiovascular events associated with selective COX-2 inhibitors. *J Am Med Assoc* 2001;286:954-9.
- 157 Kearney PM, Baigent C, Godwin J, Halls H, Emberson JR, Patrono C. Do selective cyclo-oxygenase-2 inhibitors and traditional non-steroidal anti-inflammatory drugs increase the risk of atherothrombosis? Meta-analysis of randomised trials. *Br Med J* 2006;332:1302-8.
- 158 Scheiman JM, Fendrick AM. Practical approaches to minimizing gastrointestinal and cardiovascular safety concerns with COX-2 inhibitors and NSAIDs. *Arthritis Res Ther* 2005;7(Suppl 4):S23-9.
- 159 Sopena F, Lanas A. How to advise aspirin use in patients who need NSAIDs. *Curr Pharm Des* 2007;13:2248-60.
- 160 Catella-Lawson F, Reilly MP, Kapoor SC *et al.* Cyclooxygenase inhibitors and the antiplatelet effects of aspirin. *N Engl J Med* 2001;345:1809-17.
- 161 Capone ML, Sciulli MG, Tacconelli S *et al.* Pharmacodynamic interaction of naproxen with low-dose aspirin in healthy subjects. *J Am Coll Cardiol* 2005;45:1295-301.
- 162 Corman SL, Fedutes BA, Ansani NT. Impact of nonsteroidal antiinflammatory drugs on the cardioprotective effects of aspirin. *Ann Pharmacother* 2005;39:1073-9.
- 163 US Food and Drug Administration web site. Food and Drug Administration Science Paper. Concomitant use of ibuprofen and aspirin. http://www.fda.gov/cder/drug/infopage/ibuprofen/science_paper.htm (18 April 2008, date last accessed).
- 164 Steinhubl SR. The use of anti-inflammatory analgesics in the patient with cardiovascular disease: what a pain. *J Am Coll Cardiol* 2005;45:1302-3.
- 165 Antman EM, Bennett JS, Daugherty A, Furberg C, Roberts H, Taubert KA. Use of nonsteroidal antiinflammatory drugs: an update for clinicians: a scientific statement from the American Heart Association. *Circulation* 2007;115:1634-42.
- 166 Scott DL, Smolen JS, Kalden JR *et al.* Treatment of active rheumatoid arthritis with leflunomide: two year follow up of a double blind, placebo controlled trial versus sulfasalazine. *Ann Rheum Dis* 2001;60:913-23.
- 167 Strand V, Cohen S, Schiff M *et al.* Treatment of active rheumatoid arthritis with leflunomide compared with placebo and methotrexate. Leflunomide Rheumatoid Arthritis Investigators Group. *Arch Intern Med* 1999;159:2542-50.
- 168 Cohen S, Cannon GW, Schiff M *et al.* Two-year, blinded, randomized, controlled trial of treatment of active rheumatoid arthritis with leflunomide compared with methotrexate. Utilization of leflunomide in the treatment of rheumatoid arthritis trial investigator group. *Arthritis Rheum* 2001;44:1984-92.
- 169 Rozman B, Praprotnik S, Logar D *et al.* Leflunomide and hypertension. *Ann Rheum Dis* 2002;61:567-9.
- 170 Fox RI, Herrmann ML, Frangou CG *et al.* Mechanism of action for leflunomide in rheumatoid arthritis. *Clin Immunol* 1999;93:198-208.
- 171 Li EK, Tam LS, Tomlinson B. Leflunomide in the treatment of rheumatoid arthritis. *Clin Ther* 2004;26:447-59.
- 172 BSR Guidelines for Monitoring Second Line Drugs. http://www.rheumatology.org.uk/guidelines/guidelines_other/dmardmonitoring/view (18 April 2008, date last accessed).
- 173 Weinblatt ME, Coblyn JS, Fraser PA *et al.* Cyclosporin A treatment of refractory rheumatoid arthritis. *Arthritis Rheum* 1987;30:11-7.
- 174 Dougados M, Amor B. Cyclosporin A in rheumatoid arthritis: preliminary clinical results of an open trial. *Arthritis Rheum* 1987;30:83-7.
- 175 Kurki PT. Safety aspects of the long term cyclosporin A therapy. *Scand J Rheumatol Suppl* 1992;95:35-8.
- 176 Landewe RB, Goei The HS, van Rijthoven AW, Rietveld JR, Breedveld FC, Dijkmans BA. Cyclosporine in common clinical practice: an estimation of the benefit/risk ratio in patients with rheumatoid arthritis. *J Rheumatol* 1994;21:1631-6.
- 177 Marra CA, Esdaile JM, Guh D, Fisher JH, Chalmers A, Anis AH. The effectiveness and toxicity of cyclosporin A in rheumatoid arthritis: longitudinal analysis of a population-based registry. *Arthritis Rheum* 2001;45:240-5.
- 178 Kvien TK, Zeidler HK, Hannonen P *et al.* Long term efficacy and safety of cyclosporin versus parenteral gold in early rheumatoid arthritis: a three year study of radiographic progression, renal function, and arterial hypertension. *Ann Rheum Dis* 2002;61:511-6.
- 179 Textor SC, Canzanello VJ, Taler SJ *et al.* Cyclosporine-induced hypertension after transplantation. *Mayo Clin Proc* 1994;69:1182-93.
- 180 Taler SJ, Textor SC, Canzanello VJ, Schwartz L. Cyclosporin-induced hypertension: incidence, pathogenesis and management. *Drug Saf* 1999;20:437-49.
- 181 Dijkmans B, Gerards A. Cyclosporin in rheumatoid arthritis: monitoring for adverse effects and clinically significant drug interactions. *BioDrugs* 1998;10:437-45.
- 182 Schrama YC, Koornans HA. Interactions of cyclosporin A and amlodipine: blood cyclosporin A levels, hypertension and kidney function. *J Hypertens Suppl* 1998;16:S33-8.
- 183 Dixon WG, Symmons DP. What effects might anti-TNFalpha treatment be expected to have on cardiovascular morbidity and mortality in rheumatoid arthritis? A review of the role of TNFalpha in cardiovascular pathophysiology. *Ann Rheum Dis* 2007;66:1132-6.
- 184 Jacobsson LTH, Tureson C, Gulfe A *et al.* Treatment with tumor necrosis factor blockers is associated with a lower incidence of first cardiovascular events in patients with rheumatoid arthritis. *J Rheumatol* 2005;32:1213-8.
- 185 Ferraccioli GF, Gremese E. Autoantibodies and thrombophilia in RA: TNFalpha and TNFalpha blockers. *Ann Rheum Dis* 2004;63:613-5.
- 186 Munro R, Morrison E, McDonald AG, Hunter JA, Madhok R, Capell HA. Effect of disease modifying agents on the lipid profiles of patients with rheumatoid arthritis. *Ann Rheum Dis* 1997;56:374-7.
- 187 Mottonen TT, Hannonen PJ, Boers M. Combination DMARD therapy including corticosteroids in early rheumatoid arthritis. *Clin Exp Rheumatol* 1999;17:S59-65.
- 188 Petri M, Lakatta C, Magder L, Goldman D. Effect of prednisone and hydroxychloroquine on coronary artery disease risk factors in systemic lupus erythematosus: a longitudinal data analysis. *Am J Med* 1994;96:254-9.
- 189 Halperin RO, Sesso HD, Ma J, Buring JE, Stampfer MJ, Gaziano JM. Dyslipidemia and the risk of incident hypertension in men. *Hypertension* 2006;47:45-50.
- 190 Wei L, MacDonald TM, Walker BR. Taking glucocorticoids by prescription is associated with subsequent cardiovascular disease. *Ann Intern Med* 2004;141:764-70.
- 191 Souverein PC, Berard A, Van Staa TP *et al.* Use of oral glucocorticoids and risk of cardiovascular and cerebrovascular disease in a population based case-control study. *Heart* 2004;90:859-65.
- 192 Saruta T, Suzuki H, Handa M, Igarashi Y, Kondo K, Senba S. Multiple factors contribute to the pathogenesis of hypertension in Cushing's syndrome. *J Clin Endocrinol Metab* 1986;62:275-9.
- 193 Whitworth JA. Adrenocorticotrophin and steroid-induced hypertension in humans. *Kidney Int Suppl* 1992;37:S34-7.
- 194 Filipovsky J, Ducimetiere P, Eschwege E, Richard JL, Rosselin G, Claude JR. The relationship of blood pressure with glucose, insulin, heart rate, free fatty acids and

- plasma cortisol levels according to degree of obesity in middle-aged men. *J Hypertens* 1996;14:229–35.
- 195 Litchfield WR, Hunt SC, Jeunemaitre X *et al.* Increased urinary free cortisol: a potential intermediate phenotype of essential hypertension. *Hypertension* 1998;31:569–74.
- 196 Nystrom F, Aardal E, Ohman KP. A population-based study of the white-coat blood pressure effect: positive correlation with plasma cortisol. *Clin Exp Hypertens* 1998;20:95–104.
- 197 Townsend HB, Saag KG. Glucocorticoid use in rheumatoid arthritis: benefits, mechanisms, and risks. *Clin Exp Rheumatol* 2004;22:S77–82.
- 198 Maxwell SR, Moots RJ, Kendall MJ. Corticosteroids: do they damage the cardiovascular system? *Postgrad Med J* 1994;70:863–70.
- 199 Da Silva JA, Jacobs JW, Kirwan JR *et al.* Safety of low dose glucocorticoid treatment in rheumatoid arthritis: published evidence and prospective trial data. *Ann Rheum Dis* 2006;65:285–93.
- 200 Whitworth JA. Mechanisms of glucocorticoid-induced hypertension. *Kidney Int* 1987;31:1213–24.
- 201 Savage O, Copeman WS, Chapman L, Wells MV, Treadwell BL. Pituitary and adrenal hormones in rheumatoid arthritis. *Lancet* 1962;1:232–5.
- 202 Treadwell BL, Sever ED, Savage O, Copeman WS. Side-effects of long-term treatment with corticosteroids and corticotrophin. *Lancet* 1964;1:1121–3.
- 203 Thomas TP. The complications of systemic corticosteroid therapy in the elderly. A retrospective study. *Gerontology* 1984;30:60–5.
- 204 Hoes JN, Jacobs JW, Boers M *et al.* EULAR evidence-based recommendations on the management of systemic glucocorticoid therapy in rheumatic diseases. *Ann Rheum Dis* 2007;66:1560–7.
- 205 Kalbak K. Incidence of arteriosclerosis in patients with rheumatoid arthritis receiving long-term corticosteroid therapy. *Ann Rheum Dis* 1972;31:196–200.
- 206 Jackson SH, Beevers DG, Myers K. Does long-term low-dose corticosteroid therapy cause hypertension? *Clin Sci* 1981;61(Suppl 7):381s–3s.
- 207 Hafstrom I, Rohani M, Deneberg S, Wornert M, Jogestrand T, Frostegard J. Effects of low-dose prednisolone on endothelial function, atherosclerosis, and traditional risk factors for atherosclerosis in patients with rheumatoid arthritis—a randomized study. *J Rheumatol* 2007;34:1810–6.
- 208 Iversson LL, Salt PJ. Inhibition of catecholamine uptake 2 by steroids in the isolated rat heart. *Br J Pharmacol* 2006;40:528–30.
- 209 Kalsner S. Steroid potentiation of responses to sympathomimetic amines in aortic strips. *Br J Pharmacol* 1969;36:582–93.
- 210 Haigh RM, Jones CT. Effect of glucocorticoids on alpha 1-adrenergic receptor binding in rat vascular smooth muscle. *J Mol Endocrinol* 1990;5:41–8.
- 211 Aubert J, Darimont C, Safonova I, Ailhaud G, Negrel R. Regulation by glucocorticoids of angiotensinogen gene expression and secretion in adipose cells. *Biochem J* 1997;328:701–6.
- 212 Rhen T, Cidlowski JA. Antiinflammatory action of glucocorticoids—new mechanisms for old drugs. *N Engl J Med* 2005;353:1711–23.
- 213 Jonsson SW, Backman C, Johnson O *et al.* Increased prevalence of atherosclerosis in patients with medium term rheumatoid arthritis. *J Rheumatol* 2001;28:2597–602.
- 214 Arosio E, De Marchi S, Rigoni A, Prior M, Delva P, Lechi A. Forearm haemodynamics, arterial stiffness and microcirculatory reactivity in rheumatoid arthritis. *J Hypertens* 2007;25:1273–8.
- 215 Guidelines Committee. 2003 European Society of Hypertension-European Society of Cardiology guidelines for the management of arterial hypertension. *J Hypertens* 2003;21:1011–53.
- 216 JBS 2: Joint British Societies' guidelines on prevention of cardiovascular disease in clinical practice. *Heart* 2005;91 (Suppl 5):v1–52.
- 217 Appel LJ, Brands MW, Daniels SR, Karanja N, Elmer PJ, Sacks FM. Dietary approaches to prevent and treat hypertension: a scientific statement from the American Heart Association. *Hypertension* 2006;47:296–308.
- 218 Dickinson HO, Mason JM, Nicolson DJ *et al.* Lifestyle interventions to reduce raised blood pressure: a systematic review of randomized controlled trials. *J Hypertens* 2006;24:215–33.
- 219 Rosenberg L, Kaufman DW, Helmrich SP, Shapiro S. The risk of myocardial infarction after quitting smoking in men under 55 years of age. *N Engl J Med* 1985;313:1511–4.
- 220 Sacks FM, Appel LJ, Moore TJ *et al.* A dietary approach to prevent hypertension: a review of the Dietary Approaches to Stop Hypertension (DASH) Study. *Clin Cardiol* 1999;22:III6–10.
- 221 Geleijnse JM, Giltay EJ, Grobbee DE, Donders AR, Kok FJ. Blood pressure response to fish oil supplementation: metaregression analysis of randomized trials. *J Hypertens* 2002;20:1493–9.
- 222 Beyer FR, Dickinson HO, Nicolson DJ, Ford GA, Mason J. Combined calcium, magnesium and potassium supplementation for the management of primary hypertension in adults. *Cochrane Database Syst Rev* 2006;3: CD004805.
- 223 Dickinson HO, Nicolson DJ, Cook JV *et al.* Calcium supplementation for the management of primary hypertension in adults. *Cochrane Database Syst Rev* 2006; CD004639.
- 224 British Hypertension Society. http://www.bhsoc.org/Healthy_Eating.stm (18 April 2008, date last accessed).
- 225 British Heart Foundation. http://www.bhf.org.uk/keeping_your_heart_healthy/default.aspx (18 April 2008, date last accessed).
- 226 Metsios GS, Stavropoulos-Kalinoglou A, Nevill AM, Douglas KM, Koutedakis Y, Kitas GD. Smoking significantly increases basal metabolic rate in patients with rheumatoid arthritis. *Ann Rheum Dis* 2008;67:70–3.
- 227 Hypertension: management of hypertension in adults in primary care: partial update. <http://www.nice.org.uk/nicemedia/pdf/cg034quickrefguide.pdf> (18 April 2008, date last accessed).
- 228 Dekkers JC, Geenen R, Godaert GL, Bijlsma JW, van Doornen LJ. Elevated sympathetic nervous system activity in patients with recently diagnosed rheumatoid arthritis with active disease. *Clin Exp Rheumatol* 2004;22:63–70.
- 229 Evrengul H, Dursunoglu D, Cobankara V *et al.* Heart rate variability in patients with rheumatoid arthritis. *Rheumatol Int* 2004;24:198–202.
- 230 Kaul CL, Ramarao P. Renin release and the sympathetic nervous system. *Drugs Today* 2000;36:699–713.
- 231 van den Meiracker AH, Boomsma F. The angiotensin II-sympathetic nervous system connection. *J Hypertens* 2003;21:1453–4.
- 232 Dandona P, Dhindsa S, Ghanim H, Chaudhuri A. Angiotensin II and inflammation: the effect of angiotensin-converting enzyme inhibition and angiotensin II receptor blockade. *J Hum Hypertens* 2007;21:20–7.
- 233 Gulick T, Chung MK, Pieper SJ, Lange LG, Schreiner GF. Interleukin 1 and tumor necrosis factor inhibit cardiac myocyte beta-adrenergic responsiveness. *Proc Natl Acad Sci USA* 1989;86:6753–7.
- 234 Liu SJ, Zhou W, Kennedy RH. Suppression of beta-adrenergic responsiveness of L-type Ca²⁺ current by IL-1beta in rat ventricular myocytes. *Am J Physiol* 1999;276:H141–8.
- 235 Daneshtalab N, Lewanczuk RZ, Russell A, Jamali F. Rheumatoid arthritis does not reduce the pharmacodynamic response to valsartan. *J Clin Pharmacol* 2004;44:245–52.
- 236 Daneshtalab N, Lewanczuk RZ, Russell AS, Jamali F. Drug-disease interactions: losartan effect is not downregulated by rheumatoid arthritis. *J Clin Pharmacol* 2006;46:1344–55.
- 237 Dessein PH, Joffe BI. Insulin resistance and impaired beta cell function in rheumatoid arthritis. *Arthritis Rheum* 2006;54:2765–75.
- 238 Dessein PH, Joffe BI, Stanwin AE, Christian BF, Veller M. Glucocorticoids and insulin sensitivity in rheumatoid arthritis. *J Rheumatol* 2004;31:867–74.
- 239 Lindholm LH, Ibsen H, Borch-Johnsen K *et al.* Risk of new-onset diabetes in the losartan intervention for endpoint reduction in hypertension study. *J Hypertens* 2002;20:1879–86.
- 240 Mancia G, Grassi G, Zanchetti A. New-onset diabetes and antihypertensive drugs. *J Hypertens* 2006;24:3–10.
- 241 La Civita L, Pitaro N, Rossi M *et al.* Amlodipine in the treatment of Raynaud's phenomenon. *Br J Rheumatol* 1993;32:524–5.
- 242 Warren JB, Loi RK. Captopril increases skin microvascular blood flow secondary to bradykinin, nitric oxide, and prostaglandins. *FASEB J* 1995;9:411–8.
- 243 Dziadzio M, Denton CP, Smith R *et al.* Losartan therapy for Raynaud's phenomenon and scleroderma: clinical and biochemical findings in a fifteen-week, randomized, parallel-group, controlled trial. *Arthritis Rheum* 1999;42:2646–55.
- 244 Singh G, Mithal A, Mendelsohn A, Triadafilopoulos. Acute myocardial infarctions in rheumatoid arthritis and diabetes: a tale of two diseases and a call for action. *Ann Rheum Dis* 2004;63:68.
- 245 Dessein PH, Joffe BI, Veller MG *et al.* Traditional and nontraditional cardiovascular risk factors are associated with atherosclerosis in rheumatoid arthritis. *J Rheumatol* 2005;32:435–42.
- 246 Dessein PH, Tobias M, Veller MG. Metabolic syndrome and subclinical atherosclerosis in rheumatoid arthritis. *J Rheumatol* 2006;33:2425–32.
- 247 Karvounaris SA, Sidiropoulos PI, Papadakis JA *et al.* Metabolic syndrome is common among middle-to-older aged Mediterranean patients with rheumatoid arthritis and correlates with disease activity: a retrospective, cross-sectional, controlled, study. *Ann Rheum Dis* 2007;66:28–33.
- 248 Dessein PH, Joffe BI. Suppression of circulating interleukin-6 concentrations is associated with decreased endothelial activation in rheumatoid arthritis. *Clin Exp Rheumatol* 2006;24:161–7.
- 249 Maradit-Kremers H, Nicola PJ, Crowson CS, Ballman KV, Gabriel SE. Cardiovascular death in rheumatoid arthritis: a population-based study. *Arthritis Rheum* 2005;52:722–32.
- 250 Maradit-Kremers H, Crowson CS, Nicola PJ *et al.* Increased unrecognized coronary heart disease and sudden deaths in rheumatoid arthritis: a population-based cohort study. *Arthritis Rheum* 2005;52:402–11.
- 251 Kremers HM, Nicola PJ, Crowson CS, Ballman KV, Gabriel SE. Prognostic importance of low body mass index in relation to cardiovascular mortality in rheumatoid arthritis. *Arthritis Rheum* 2004;50:3450–7.
- 252 La Montagna G, Cacciapuoti F, Buono R *et al.* Insulin resistance is an independent risk factor for atherosclerosis in rheumatoid arthritis. *Diab Vasc Dis Res* 2007;4:130–5.
- 253 Pamuk ON, Unlu E, Cakir N. Role of insulin resistance in increased frequency of atherosclerosis detected by carotid ultrasonography in rheumatoid arthritis. *J Rheumatol* 2006;33:2447–52.
- 254 Panoulas VF, Douglas KM, Milionis HJ *et al.* Serum uric acid is independently associated with hypertension in patients with rheumatoid arthritis. *J Hum Hypertens* 2008;22:177–82.
- 255 Dessein PH, Joffe BI. Insulin resistance and impaired beta cell function in rheumatoid arthritis. *Arthritis Rheum* 2006;54:2765–75.
- 256 Dessein PH, Woodiwiss AJ, Joffe BI, Norton GR. Aminotransferases are associated with insulin resistance and atherosclerosis in rheumatoid arthritis. *BMC Cardiovasc Disord* 2007;7:31.
- 257 Wallberg-Jonsson S, Ohman ML, Dahlqvist SR. Cardiovascular morbidity and mortality in patients with seropositive rheumatoid arthritis in Northern Sweden. *J Rheumatol* 1997;24:445–51.
- 258 Roman MJ, Devereux RB, Schwartz JE *et al.* Arterial stiffness in chronic inflammatory diseases. *Hypertension* 2005;46:194–9.
- 259 Wolfe F, Michaud K. Heart failure in rheumatoid arthritis: rates, predictors, and the effect of anti-tumor necrosis factor therapy. *Am J Med* 2004;116:305–11.