

Ice velocity and climate variations for Baltoro Glacier, Pakistan

D.J. QUINCEY,¹ L. COPLAND,² C. MAYER,³ M. BISHOP,⁴ A. LUCKMAN,⁵ M. BELÒ⁶

¹*Institute of Geography and Earth Sciences, Penglais Campus, Aberystwyth University, Aberystwyth SY23 3DB, UK
E-mail: djq@aber.ac.uk*

²*Department of Geography, University of Ottawa, Ottawa, Ontario K1N 6N5, Canada*

³*Commission for Glaciology, Bavarian Academy of Sciences and Humanities, Alfons-Goppel Strasse 11,
D-80539 Munich, Germany*

⁴*Department of Geography and Geology, University of Nebraska at Omaha, Omaha, Nebraska 68182, USA*

⁵*School of the Environment and Society, Swansea University, Singleton Park, Swansea SA2 8PP, UK*

⁶*Trimble Navigation, Centro Torri Bianche, Palazzo Larice, I-20059 Vimercate, Milan, Italy*

ABSTRACT. The recent dynamic behaviour of Karakoram glaciers is expected to differ from that shown by glaciers in the central and eastern Himalaya because of regional variations in precipitation and temperature trends. However, there are insufficient quantitative data to support or confute such hypotheses. We present velocity data covering the period 1993–2008 for Baltoro Glacier, one of the longest glaciers in the Karakoram. Velocity measurements were made using cross-correlation feature tracking applied to European Remote-sensing Satellite (ERS-1 and -2) and Envisat advanced synthetic aperture radar (ASAR) data, supplemented by differential global positioning system (DGPS) measurements. We find a gradual acceleration of the glacier during the early 2000s, in particular during winter months. Multi-seasonal data reveal a large difference between summer and winter flow characteristics, but only in the upper ablation zone. Summer 2005 was a particularly dynamic period following from the heavy winter snowfall of 2004, indicating the importance of basal meltwater availability for glacier flow. Transverse velocity profiles indicate that Baltoro Glacier undergoes ‘block’ flow across much of the upper ablation zone during the summer, which we interpret as evidence of widespread basal sliding. The DGPS data confirm the rapid increase in flow detected during 2005. Modelled climatic data reveal decreasing summer temperatures and increasing precipitation over the study period, helping to explain the observed dynamic variations and their differences from glaciers elsewhere in the Himalaya.

INTRODUCTION

Reports of mountain glacier recession in response to sustained climatic warming are widespread, with some of the most extensive areas of stagnant, downwasting ice being observed in parts of the eastern Himalaya (Bolch and others, 2008; Quincey and others, 2009). However, patterns of climate warming are not uniform, and the western Himalaya appears to be a region not conforming to the global trend (Yadav and others, 2004). Indeed, climate data for the period 1961–2000 show a reduction in mean and minimum summer temperatures (Fowler and Archer, 2006), and an increase in winter mean and maximum temperature coincident with an increase in winter snowfall (Archer and Fowler, 2004). Oxygen isotope data derived from tree-rings, largely indicative of winter precipitation, also suggest that the 20th century was the wettest period in the mountains of northern Pakistan of the past 1000 years (Treydte and others, 2006). Consequently, some Karakoram glaciers have been observed to be thickening and advancing (Hewitt, 2005).

The Karakoram is home to several of the highest mountains in the world, including K2. Extreme topography such as this enables the formation of large, dynamic glaciers (Kerr, 1993); indeed four of the longest Asian glaciers (Biafo, Siachen, Sarpo Laggo and Baltoro) descend from the K2 and Gasherbrum groups (Mayer and others, 2006). The long-term dynamics (mass balance, horizontal and vertical fluctuations, ice flow) of these glaciers reflects regional climatic conditions (Dyurgerov and Meier, 2000), although the short-term relationships are unclear due to the paucity of climate station data, particularly at high elevations (Fowler

and Archer, 2006). Modelled climate data therefore provide the only realistic means of estimating past precipitation and temperature data for the Karakoram, and are particularly appropriate for use on a regional scale (Uppala and others, 2005).

The short-term dynamic response of Karakoram glaciers is often inhibited by their thick surface-debris cover, which has an insulating effect on ablation (Benn and Lehmkühl, 2000). However, rapid changes in flow characteristics, in particular glacier surges, have been widely noted in the region (Hewitt, 1969, 2005, 2007). It has been shown that the flow of valley glaciers can fluctuate on seasonal and annual timescales, and that motion is often linked to variations in water supply to the subglacial drainage system (Willis, 1995; Kavanaugh and Clarke, 2001; Purdie and others, 2008). However, there is a paucity of such information for the Karakoram. Useful measurements of glacier velocity have been derived from point-based GPS data and the manual tracking of surface features across sequential imagery (Mayer and others, 2006). Automated feature tracking has also been used (Copland and others, 2009), but without linkage to short- or long-term climatic variations.

Our aim is to build on this previous work and to examine the spatial and temporal variability in surface velocity exhibited by Baltoro Glacier, one of the longest non-polar glaciers in the world. Using robust cross-correlation feature-tracking algorithms (e.g. Strozzi and others, 2002), spatially comprehensive seasonal and annual velocity fields are presented from the mid- to late 1990s through to early 2008. These remote-sensing data are complemented and supported by two sets of field-based measurements from the



Fig. 1. Baltoro Glacier, with major peaks, tributary glaciers and profile locations (in yellow) highlighted.

summers of 2004 and 2005, periods of particularly interesting dynamical behaviour. Modelled climate data (European Centre for Medium-Range Weather Forecasts reanalysis, ERA-40) are presented for the period 1958–2001, showing temperature and precipitation trends on a seasonal scale. The controlling parameters on long-term glacier flow are discussed within the context of the modelled meteorological data, and the possible controls on short-term surface velocity fluctuations are explored.

STUDY AREA AND RELEVANT PREVIOUS WORK

Baltoro Glacier

Baltoro Glacier descends from the western face of K2 (8611 m a.s.l.) via Godwin Austen Glacier, and terminates approximately 63 km down-valley at an elevation of ~3500 m. It is fed by one other major tributary, Baltoro South Glacier, which itself descends from the Gasherbrum group of peaks (maximum elevation 8080 m) via Abruzzi Glacier (Fig. 1). A number of small debris-covered tributaries join the main glacier tongue along its northern and southern margins, some of which exhibit surge-type characteristics (Diolaiuti and others, 2003), although there is no evidence that Baltoro Glacier itself surges.

The wider Karakoram climate is strongly influenced by troughs of low pressure in the westerly circulation (Lüdecke and Kuhle, 1991), leading to maximum precipitation during winter and spring months (Archer and Fowler, 2004), and by monsoonal incursions, which bring a small amount of additional summer precipitation. In the specific case of Baltoro Glacier, climatic variables are strongly influenced by topography: precipitation rates increase with altitude such that almost all precipitation above the equilibrium line (around 5500 m) falls as snow (Mayer and others, 2006), and on a local scale, ablation rates are influenced by slope and aspect as well as elevation (Mihalcea and others, 2006).

The upper reaches of Baltoro Glacier are largely debris-free, but downstream of the equilibrium line the debris cover

becomes increasingly thick and extensive. At the confluence between the Godwin Austen and Baltoro South tributaries, known as Concordia (~4600 m a.s.l.; Fig. 2a), approximately 70% of the glacier surface is buried by a thin layer (5–15 cm) of debris. At around 4100 m (Urdukas), the debris cover is almost 100% and is 30–40 cm thick, whereas the debris is up to 1 m deep close to the glacier terminus (~3450 m a.s.l.; Mihalcea and others, 2008). Semi-permanent surface meltwater ponds punctuate the debris cover across the lower ablation area (Fig. 2b), occasionally reaching sizes of ~100 × 100 m or more. Water discharge from the glacier occurs primarily via a single outlet at the glacier terminus, where large volumes of very heavily debris-laden subglacial water are released in the summer (Fig. 2c). According to historical data, the glacier terminus position has gently fluctuated since 1913, with a 200 m advance in the early 20th century, followed by a similar recession since 1954. Repeat photographs suggest that there has been negligible surface thickening or downwasting over this same period (Mayer and others, 2006).

Previous velocity measurements in the Karakoram

There have been several previous measurements of seasonal ice velocity variability in the Karakoram, predominantly focusing on the largest glaciers in this region. For example, on Batura Glacier, the Batura Glacier Investigation Group (1979) used a combination of terrestrial stereophotography and control survey to measure the surface motion across the ablation area of the glacier at a total of 129 locations in 1974/75. They found average velocities of ~100 m a⁻¹, with a peak of 517.5 m a⁻¹. Velocities were higher from June to September than the mean annual rate, although typical summer increases were <20%. In contrast, the motion at the glacier terminus varied little between summer and winter.

On Biafo Glacier, Hewitt and others (1989) used a theodolite to measure the position of a network of nine stakes across the middle of the glacier's ablation area in 1985/86, with summer velocities of 128–226 m a⁻¹. Winter




Fig. 2. Baltoro Glacier showing (a) Concordia, the confluence of Baltoro South and Godwin Austen tributaries; (b) the debris-covered tongue viewed from Urdukas base camp, ~ 14 km up-glacier from the terminus (ice flow is from right to left); and (c) a view of the heavily debris-covered terminus from the proglacial outwash area.

velocities were only recorded at two stakes, although these measurements indicated that the glacier moved about half as fast in the winter as in the summer. Copland and others (2009) used satellite image feature tracking to determine velocities over the period 2006–07 at these same locations, and found that annual motion was consistent with the earlier measurements (within error limits).

Previous measurements on Baltoro Glacier

In summer 2004, a combined Italian/German expedition performed extensive measurements across the ablation zone of Baltoro Glacier, assessing the production of meltwater on debris-covered glacier tongues (Mayer and others, 2006). Ablation was measured at more than 50 stakes spread across the central part of the ablation area from 4000 m up to 5000 m. At 32 of these stakes, the surface velocity of the glacier was also determined by repeated differential global positioning system (DGPS) measurements (Mayer and others, 2006). Maximum velocities of 214 m a^{-1} were found close to Concordia, while typical summer velocities of $\sim 100 \text{ m a}^{-1}$ were recorded across most of the rest of the central ablation area. In contrast, annual velocities derived from manual feature tracking showed values approximately half of the measured summer velocities. This large seasonal velocity variation was attributed to strong basal sliding, at least during the summer months.

Copland and others (2009) measured annual velocities over the lowermost 13 km of Baltoro Glacier using feature tracking of 2006–07 Advanced Spaceborne Thermal Emission and Reflection Radiometer (ASTER) satellite scenes, with velocities ranging from $\sim 75 \text{ m a}^{-1}$ near the upper part of the study region (around Urdukas; Fig. 1) to $< 15 \text{ m a}^{-1}$ close to the glacier terminus. Transverse velocity profiles

showed strong lateral shearing indicative of basal sliding at a distance of ~ 12 km from the glacier terminus, but transverse profiles more indicative of internal deformation ~ 5 km from the terminus. Comparison with summer DGPS measurements indicated that summer speed-ups occurred close to the upper transverse profile, but there was little evidence for summer speed-ups close to the lower transverse profile. This lack of seasonal velocity variations in the near-terminus region is similar to that described by the Batura Glacier Investigation Group (1979; see above).

DATA AND METHODS

ERS-1 and -2 and Envisat ASAR

The European Remote-sensing Satellites have provided the glaciological community with fine-resolution radar data since the launch of ERS-1 in 1992. More recently, the launch of the advanced synthetic aperture radar (ASAR) sensor (on board the Envisat satellite) has continued the systematic collection of C-band data (wavelength 5–7 cm), giving a well-stocked archive for the Karakoram region for the previous 16 years. In the current study, six ERS-1 and -2 scenes were selected for the period 1992–2000, and eight ASAR scenes were selected for the period 2003–present. The resolutions of the raw data were approximately 7.9 m (range) \times 4.0 m (azimuth), equating to 20–30 m in ground range at scene centre. Images were selected to provide pairs covering annual (single year), summer (March/April to August/September), and winter (August/September to March/April) periods (Table 1). All data were selected from the same orbit, ensuring the minimum spatial baseline and consistent imaging conditions.

Table 1. Remote-sensing data used

	Date of acquisition	Track	Frame	Date of acquisition	Track	Frame
ERS-1/2	24 Dec. 1992	377	2887	8 Apr. 1993	377	2887
	20 Apr. 1996		2887	25 May 1996		2887
	7 Dec. 1997		2887	11 Apr. 1999		2888
ASAR	3 Aug. 2003	377	2886	4 Apr. 2004	377	2886
	22 Aug. 2004		2886	24 Apr. 2005		2887
	11 Sep. 2005		2887	5 Mar. 2006		2887
	16 Sep. 2007		2886	13 Apr. 2008		2886
Annual pairs		Summer pairs		Winter pairs		
8 Apr. 1993–20 Apr. 1996		25 May 1996–7 Dec. 1997		7 Dec. 1997–11 Apr. 1999		
3 Aug. 2003–22 Aug. 2004		4 Apr. 2004–22 Aug. 2004		3 Aug. 2003–4 Apr. 2004		
4 Apr. 2004–24 Apr. 2005		24 Apr. 2005–11 Sep. 2005		22 Aug. 2004–24 Apr. 2005		
22 Aug. 2004–11 Sep. 2005				11 Sep. 2005–5 Mar. 2006		
24 Apr. 2005–5 Mar. 2006						
11 Sep. 2005–16 Sep. 2007						
5 Mar. 2006–13 Apr. 2008						

Cross-correlation and feature tracking (including error estimations)

Surface-displacement measurements are made using cross-correlation feature tracking and are used here as a surrogate for glacier velocity. Cross-correlation feature tracking relies on the identification of displaced features across two time-separated synthetic aperture radar (SAR) images (Lucchitta and others, 1995). Image patches are defined at a size commensurate with the feature patterns to be tracked, and the search area is defined on the basis of the likely displacement over the imaged period. In the current study, the windows were sized at 30×150 pixels (pattern size) and 64×320 pixels (search area), equating to ~ 600 m and ~ 1280 m, respectively, on the ground. Sampling of the displacement field was made on a much finer grid (4×20 pixels, equating to ~ 50 m on the ground). The relatively large window size is necessary to capture sufficient ground features in the presence of image speckle, and may result in a degree of smoothing of the displacement fields. However, the spatial resolution of the technique can be considerably higher than the window size might imply. This is because the correlation field does not identify the average displacement but finds the displacement of the dominant features within the search window (in terms of area and contrast). Thus sharp gradients in velocity (e.g. between the glacier and its lateral moraines) are not problematic.

The cross-correlation method was implemented using a transformation of the SAR images to spatial frequencies using the fast Fourier transform (Strozzi and others, 2002). Feature matches are made based on the peak of the cross-correlation results and each match is assigned a signal-to-noise ratio (SNR) calculated from the height of the correlation peak relative to the average level of correlation. Matches below a specified SNR (7.0 in the present study) are rejected. Further noise is removed from the displacement map before projection in Universal Transverse Mercator (UTM) coordinates by masking areas of shadow and radar layover.

Error in the remaining data is caused by changes in surface features over time (thereby causing small-scale errors

in the matching) and image distortions introduced by the processing chain (e.g. projection from SAR to UTM coordinates). Although it is difficult to quantify the error exactly, previous studies have shown that by using well-separated image pairs (of the order of a year or more), the uncertainty in the data can be kept as low as $2\text{--}3 \text{ m a}^{-1}$ (Luckman and others, 2007). It is possible to validate these estimates by collecting measured displacement data over known stable areas; in the current study, the error in the data was consistently less than $\pm 7 \text{ m a}^{-1}$ for the annually separated data, and $\pm 11 \text{ m a}^{-1}$ for the seasonal ($\sim 4\text{--}6$ months separation) data. Such errors are small compared with the magnitude of displacement recorded over the glacier surface, and well within the observed trends in the seasonal and annual comparisons.

DGPS

DGPS data were collected during the summer months of 2004 and 2005 by two separate field teams.

During the 2004 field season, stake position measurements were performed with Trimble 5700 geodetic GPS receivers, and differentially corrected using a network of benchmarks installed at camps along the glacier margin. Ice velocities were measured at cross-sections at Urdukas, Gore I, Concordia and across the two main glacier tributaries: Godwin Austen and Baltoro South (Fig. 1). In addition, a longitudinal profile was collected along a central flowline from Urdukas to the highest stake on the Baltoro South tributary (~ 4640 m). Revisit periods varied from 1 day in the upper elevation areas to 13 days for the cross-profile at Urdukas (Table 2). Error in the horizontal displacement recorded over a single day was calculated to be $\sim 4 \text{ m a}^{-1}$. For the longest revisit period at Urdukas, the accuracy of the velocity data is calculated to be better than 0.3 m a^{-1} .

During summer 2005, ice-motion data were collected along the length of Baltoro Glacier from repeated measurement of poles drilled into the ice (upper ablation area), or of crosses painted on large, stable supraglacial boulders (lower ablation area). Positions were recorded with a Trimble R7

Table 2. Comparison of DGPS vs satellite-derived velocities. Points 1–14 surveyed by Copland; all other points surveyed by Mayer

Point	Field data						Feature-tracking data				
	GPS latitude	GPS longitude	GPS elevation	GPS survey 1	GPS survey 2	GPS velocity	GPS direction	Annual 2004/05 velocity	Annual 2005/06 velocity	Summer 2004 velocity	Summer 2005 velocity
	°	°	m			ma ⁻¹	°	ma ⁻¹	ma ⁻¹	ma ⁻¹	ma ⁻¹
1	35.69366	76.16118	3483.5	25-Jun-05	17-Jul-05	10.3	241.7	No data	No data	No data	No data
2	35.72106	76.23045	3816.4	28-Jun-05	16-Jul-05	38.3	242.1	36.1	33.3	51.0	38.1
3	35.72472	76.23064	3835.2	28-Jun-05	16-Jul-05	44.9	245.9	38.7	44.1	57.7	49.5
4	35.71891	76.23159	3810.5	28-Jun-05	16-Jul-05	19.7	234.9	15.3	24.2	43.1	32.9
5	35.73274	76.28443	3991.2	30-Jun-05	15-Jul-05	91.5	251.8	58.4	75.3	62.3	85.7
6	35.73558	76.28507	3995.8	01-Jul-05	15-Jul-05	117.1	255.5	63.9	75.6	79.1	96.4
7	35.73634	76.30848	4055.1	02-Jul-05	15-Jul-05	135.5	273.9	79.9	84.5	92.0	99.1
8	35.74663	76.40521	4280.5	06-Jul-05	15-Jul-05	245.0	249.9	112.4	132.1	142.6	163.3
9	35.74468	76.40401	4268.2	06-Jul-05	14-Jul-05	239.4	248.7	115.7	120.8	174.2	158.4
10	35.74147	76.51768	4565.3	10-Jul-05	19-Jul-05	168.4	277.7	134.7	149.4	175.9	172.6
11	35.74044	76.52116	4580.4	10-Jul-05	19-Jul-05	159.5	285.7	123.4	136.5	154.4	163.7
12	35.75117	76.52272	4590.5	11-Jul-05	18-Jul-05	125.7	215.8	85.0	97.7	94.5	No data
13	35.79587	76.51830	4774.1	12-Jul-05	18-Jul-05	154.6	177.5	No data	No data	No data	No data
14	35.81586	76.51389	4837.7	12-Jul-05	18-Jul-05	159.2	167.1	124.1	130.8	116.7	No data
c1	35.74326	76.50225	4538.2	08-Jul-04	11-Jul-04	205.0	264.6	155.5	172.2	185.1	209.8
c2	35.74254	76.5022	4538.6	08-Jul-04	11-Jul-04	195.0	264.7	156.2	174.9	185.0	210.5
c3	35.74182	76.50218	4542.3	08-Jul-04	11-Jul-04	191.1	264.5	155.1	179.8	186.6	210.8
c4	35.74116	76.50208	4545.0	08-Jul-04	11-Jul-04	186.0	264.5	144.9	180.2	181.0	206.5
c5	35.74050	76.50202	4522.5	08-Jul-04	11-Jul-04	182.9	264.8	152.9	179.2	183.3	233.2
g1	35.74787	76.36437	4184.1	04-Jul-04	13-Jul-04	72.9	280.5	58.0	76.8	71.5	103.8
g2	35.74350	76.36432	4189.7	04-Jul-04	13-Jul-04	121.7	277.5	75.2	87.5	103.1	115.6
g3	35.73899	76.36373	4178.6	04-Jul-04	13-Jul-04	126.8	274.6	82.5	88.1	102.9	121.1
g4	35.73460	76.36432	4181.7	04-Jul-04	13-Jul-04	124.8	277.5	71.9	98.6	105.7	134.0
g5	35.73039	76.36312	4181.6	04-Jul-04	13-Jul-04	83.9	283.6	54.4	57.8	78.5	94.1
bs1	35.71425	76.54687	4676.0	07-Jul-04	11-Jul-04	141.4	308.6	125.9	135.9	157.2	152.3
bs2	35.72302	76.53747	4638.7	07-Jul-04	11-Jul-04	132.6	326.5	137.6	125.8	138.2	144.0
bs4	35.72373	76.53839	4624.4	07-Jul-04	11-Jul-04	131.4	328.0	121.9	119.1	132.0	142.3
bs5	35.73490	76.52913	4606.2	07-Jul-04	11-Jul-04	131.7	322.6	129.6	131.5	146.9	160.6
u1	35.73075	76.28591	3983.9	02-Jul-04	15-Jul-04	49.9	255.0	61.3	53.1	50.7	59.8
u2	35.73367	76.28544	3984.8	02-Jul-04	15-Jul-04	82.9	255.0	60.6	72.1	71.2	87.9
u3	35.73653	76.28432	3988.7	02-Jul-04	15-Jul-04	95.7	256.6	65.3	74.9	83.9	98.7
u4	35.73984	76.28442	3997.3	02-Jul-04	15-Jul-04	100.3	257.6	79.2	75.1	83.4	96.5
u5	35.74434	76.28348	4007.6	02-Jul-04	15-Jul-04	94.5	257.1	65.7	66.4	77.7	93.6
bno1	35.75079	76.51999	4575.6	11-Jul-04	11-Jul-04	116.8	236.5	100.0	115.1	107.8	No data
bno2	35.76311	76.52181	4638.1	09-Jul-04	11-Jul-04	121.5	193.6	No data	No data	No data	No data
bno5	35.77192	76.51951	4665.3	09-Jul-04	11-Jul-04	120.3	201.6	No data	No data	No data	No data
bno3	35.77328	76.52267	4677.3	09-Jul-04	11-Jul-04	117.3	196.4	No data	No data	No data	No data
bno4	35.77153	76.51837	4670.4	09-Jul-04	11-Jul-04	127.7	197.6	No data	No data	No data	No data
blo1	35.74002	76.31347	4076.3	03-Jul-04	14-Jul-04	120.3	272.2	86.9	80.6	97.6	105.0
blo2	35.73953	76.34159	4261.1	03-Jul-04	14-Jul-04	132.7	279.2	82.1	98.5	114.3	133.6
blo4	35.74416	76.40943	4273.6	05-Jul-04	12-Jul-04	181.9	252.1	116.4	142.1	149.6	167.7
blo5	35.74342	76.42856	4337.1	05-Jul-04	12-Jul-04	185.8	286.0	129.6	148.6	163.5	191.3
blo6	35.73993	76.44192	4368.2	05-Jul-04	12-Jul-04	199.8	291.9	134.7	165.3	179.6	212.5
blo7	35.73699	76.48062	4444.6	08-Jul-04	11-Jul-04	194.8	250.9	142.0	160.0	171.7	208.6
blo8	35.74048	76.4917	4495.3	08-Jul-04	11-Jul-04	213.8	240.9	165.0	193.5	193.9	234.0
							Mean	99.9	111.7	121.7	138.5

DGPS unit, with site occupation of ≥ 30 min per visit and periods of 6–28 days between revisits (Table 2). Differential correction was undertaken using the Precise Point Positioning (PPP) method, or by comparison with base stations set up in nearby camps (located on bedrock) or in Skardu (~ 80 km southwest). Errors in positioning are estimated to be ~ 6 ma⁻¹ for the shortest time interval data and ~ 1.5 ma⁻¹ for the longest time interval (28 day) data.

ERA-40 climate re-analysis data

Climate re-analysis data refer to the processing of meteorological observations by use of a physical model, in which data are produced in a physically consistent way and interpolated to a three-dimensional (3-D) grid. The ERA-40 monthly time-series data represent a second-generation analysis produced by the European Centre for Medium-Range Weather Forecasts, Reading, UK. Although the ERA-40




Fig. 3. Average velocity field, 2003–08, derived from six annual pairs (cf. Table 1) and orthoprojected using Shuttle Radar Topography Mission (SRTM) digital elevation model (DEM) data. Inset: centre-line velocity profile; location as depicted by dashed yellow line in the main image.

model resolution is relatively fine from a global perspective, it is quite coarse when examining results on spatial scales of hundreds of kilometres. For this reason, spatial averaging over the K2 region was used to describe regional climate fluctuations since the averaging procedure produces statistically more significant and robust results, although it does not account for spatial differences in temperature and precipitation resulting from topographic variability within the region.

A monthly time series was generated for our region to examine the long-term trends in temperature and precipitation. For temperature, the summer trend was considered most important, as melting dominates during the summer ablation season, and despite an increase in winter temperatures observed over our region, they are still below freezing and have little impact on ablation. The summer mean was computed by averaging June, July and August temperatures, and a Gaussian filter was used to produce a low-frequency trend from 1958 to 2001.

For precipitation, the annual mean data were considered most important, as snowfall occurs during all seasons of the year. A seasonal analysis was also performed to determine which seasons contribute most to the annual precipitation trend. A Gaussian filter was used to produce a low-frequency trend from 1958 to 2001. The annual precipitation data best characterize total snowfall and potential accumulation.




Fig. 4. Transverse velocity profiles extracted from annually averaged data, 2003–08. Profile locations as depicted in Figures 1 and 3.

Tropical Rainfall Measuring Mission (TRMM) precipitation data

Regional observational precipitation data were acquired from the TRMM data visualization and analysis website (<http://lake.nascom.nasa.gov/tovas/>). Specifically, the 3B43 V6 rainfall data product was used. This product is considered the 'best' estimate of rainfall based upon satellite observational data as well as rain gauge data (Huffman and others, 2007). TRMM estimates of the monthly accumulated precipitation were employed to generate a spatially averaged time series using $0.25^\circ \times 0.25^\circ$ data. The annual mean precipitation data from 1998 to 2007 are most relevant to this study, although we also conducted a seasonal analysis to better elucidate climate/velocity variations.

RESULTS

Spatial patterns in flow

Derived velocity data demonstrate that Baltoro Glacier is currently active across its entire ablation and debris-covered area, with displacements of $\sim 15\text{--}20 \text{ m a}^{-1}$, well in excess of the estimated noise in the data ($\pm 7 \text{ m a}^{-1}$), detected immediately up-glacier of the terminus (Fig. 3). There is a general pattern of increasing motion with distance from the terminus, with the maximum velocity just below Concordia. This maximum of $\sim 180 \text{ m a}^{-1}$ is located at the peak of a steep flow gradient (Fig. 3 inset). Up-glacier, the velocity drops steeply as the profile extends up the Baltoro South tributary, reaching a maximum of $\sim 170 \text{ m a}^{-1}$, until the feature-tracking algorithm is no longer able to track features reliably over the clean-ice area.

Transverse velocity profiles extracted from annually averaged data are in agreement with the general trend of increasing flow with distance from the terminus (Fig. 4). The lowermost profile (t1) lacks displacement data towards the glacier margins, but shows uniform velocity across the glacier width and a gradual reduction in flow with distance from the centre line. Profiles t2–t5 are of a distinctly 'blocky' nature, with little variation in flow speed across almost the entire width of the glacier, and a rapid decrease




Fig. 5. Annual centre-line velocity profiles, 1993–2008. Date format is day/month/year.

in velocity at the glacier margins from rapidly flowing to almost stagnant ice.

Annual fluctuations (1992–2008)

Annual velocity profiles show a strong consistency in the spatial variability of flow across the entire glacier snout (Fig. 5). The greatest consistency is in the lowermost 16 km of the glacier, where multi-annual profiles largely replicate one another. Up-glacier, the profiles separate into two approximate groups: one group reflecting dynamics through the 1990s to 2004/05; and the second group reflecting conditions from 2004/05 to 2008. Within the latter group, velocities during 2005 were greatest, with an apparent speed-up of ~ 20 – 25% immediately down-glacier of Concordia compared with earlier measurements. In recent years, velocities appear to have decreased again, although locally they remain ~ 15 – 20% greater than displacements measured through the 1990s and early 2000s. Up-glacier of Concordia, all profiles display a more consistent pattern, with very little detectable interannual variability.

Seasonal fluctuations ('winter' vs 'summer' pairs)

Summer and winter velocity data are in agreement with previous reports of a strong seasonal velocity gradient on Baltoro Glacier (Mayer and others, 2006; Fig. 6). There is a strong consistency in flow profiles for the lowermost 10 km of the glacier snout, beyond which the summer motion significantly exceeds winter motion. The greatest differences are apparent in the area around Concordia, where the summer velocities exceed the winter velocities by up to 65%. Further up-glacier, the seasonal speed-up is still evident, with summer velocities consistently exceeding winter velocities by up to $\sim 35\%$.

A notable feature of the seasonal velocity data is the apparent speed-up of Baltoro Glacier during winter months in recent years. Centre-line profiles demonstrate that the data collected over winter 2003/04 by as much as 25%, with a gradual increase in the intervening years (Fig. 6). Again, the greatest differences are recorded around Concordia, with, significantly, little annual variability detectable up-glacier on the Baltoro South tributary.

Summer-only velocity data confirm a significant speed-up of the glacier during summer 2005, with velocities exceeding those during the similar period in 2004 by approximately 20% in the area immediately below Concordia.




Fig. 6. Seasonal centre-line velocity profiles, 1996–2006. Date format is day/month/year.

DGPS measurements (2004 and 2005)

Surface velocities measured by DGPS in summer 2004 are characterized by a pattern of increasing motion with distance from the terminus, up to a maximum of $>200 \text{ m a}^{-1}$ close to Concordia (Fig. 7; Table 2). Further up-glacier, measured surface displacements decrease markedly on both tributaries of the main Baltoro tongue, with a reduction in flow of $>50\%$ within 3.6 km upstream of Concordia. Local velocity increases are also observed, in particular where main tributaries (e.g. Biarchedi and Yermanendu Glaciers; Fig. 1) contribute to the ice flux.

Cross-profiles acquired at Concordia, on the Baltoro South tributary and on the Godwin Austen tributary, are restricted to the central part of the glacier and therefore exhibit little lateral variability. In contrast, cross-profiles at Gore I and Urdukas extend close to the glacier margin, where the surface velocities are shown to be 40–50% lower than ice velocities in the middle of the glacier ($\sim 70 \text{ m a}^{-1}$ compared with $\sim 125 \text{ m a}^{-1}$ at Gore 1, and $\sim 50 \text{ m a}^{-1}$ compared with $\sim 100 \text{ m a}^{-1}$ at Urdukas).

A comparison between DGPS measurements in both study years indicates that velocities were consistently higher in 2005 than in 2004 for stakes at nearby locations, despite the measurements having been made over nearly identical periods in each summer (Table 2; Fig. 7). For example, velocities at Gore II were 245.0 and 239.4 m a^{-1} at points 8 and 9 over the period 6–14 July 2005, but only 181.9 m a^{-1} at a nearby location (blo4) for 5–12 July 2004. Similarly, for the two overlapping points on the Urdukas transect, measured ice motion was (a) 117.1 m a^{-1} for 1–15 July 2005 (point 6) compared with 95.7 m a^{-1} for 2–15 July 2004 (point u3), and (b) 91.5 m a^{-1} for 30 June–15 July 2005 (point 5) compared with 82.9 m a^{-1} (point u2) for 2–15 July 2004. For all these data, ice-flow direction varied by $<3.5^\circ$ between stake pairs, indicating that velocity changes cannot be attributed to differences in flow direction.

Overall, the field measurements from both 2004 and 2005 compare well with the data extracted by feature tracking (Table 2; Fig. 7 inset), confirming the large seasonal velocity variations on the glacier as well as the increased velocities in summer 2005 compared with summer 2004. For all of the feature-tracking points provided in Table 2, average annual velocities were $\sim 12\%$ higher in 2005/06 than 2004/05, and $\sim 14\%$ higher in summer 2005 than in summer 2004.




Fig. 7. DGPS data collected during summer 2004 and summer 2005 field seasons. Inset: summer 2004/05 velocity profiles in grey, with field measurements for comparison.

ERA-40 temperature and precipitation

The ERA-40 summer mean temperature data of 1958–2001 depict a negative summer temperature trend from 1958 to 1990, and a positive trend from 1990 to 2001 (Fig. 8a). This finding is in general agreement with the work of Archer and Fowler (2004), although their analysis was based upon linear regression. The increase in temperatures during the 1990s suggests an increased influence of the southwest monsoon, possibly related to El Niño, which would have brought in warmer air currents, thereby increasing precipitation during the summer. Our seasonal analysis of ERA-40 summer precipitation trends indicates increasing precipitation during the 1990s.

The ERA-40 mean annual precipitation trend is positive from 1970 to 1990, and negative during the 1990s (Fig. 8b). Seasonal analysis indicates that the positive trend is primarily due to precipitation during the spring (westerlies) and summer. Other seasons show a negative trend during this time frame. Collectively, the precipitation and summer temperature trends support the interpretation of positive mass-balance conditions from 1970 to 1990. The negative trend in the 1990s is partially confirmed in the TRMM precipitation time series (Fig. 8d). Furthermore, a positive trend in summer precipitation from 1970 to 2001 demonstrates the potential influence of the summer monsoon given the temperature trend (Fig. 8c). Over this region, the




Fig. 8. (a) Mean summer temperature, (b) mean annual precipitation, and (c) mean summer precipitation, all derived from ERA-40 re-analysis data. (d) Mean annual precipitation, derived from TRMM estimates.

westerlies provide the majority of the precipitation, and summer precipitation can also come from the westerlies. It is expected, however, that an increase in summer precipitation from the westerlies is associated with a decrease in temperatures.

TRMM precipitation

The TRMM mean annual precipitation data reveal a decrease in precipitation in the late 1990s, and non-linear smoothing reveals a positive trend from 2000 to 2005, followed by a negative trend (Fig. 8d). The highest annual accumulation occurs from 2003 to 2006. Consequently, the region experiences increases in precipitation during different time frames, due to variations in the influence and magnitude of the westerlies and the summer monsoon. Collectively, increases in annual precipitation and decreasing summer temperatures indicate that Baltoro Glacier is accumulating more mass at higher altitudes. It is important to note, however, that the region lacks direct mass-balance data, and that the spatio-temporal variations in ablation and snow accumulation are highly governed by multi-scale topographic influences.

DISCUSSION

Several glaciers in the Karakoram are reported to have been thickening and advancing since the mid- to late 1990s (Hewitt, 2005), in contrast to the trends of stagnant ice and surface downwasting observed in most other areas of the Himalaya (Berthier and others, 2007; Bolch and others, 2008; Quincey and others, 2009). Modelled climate data presented here indicate that, in recent years, accumulation at high altitude has increased in the Karakoram, leading to positive mass balances and, ultimately, ice thickening and advance. In the specific case of Baltoro Glacier, field observations, centre-line velocity profiles presented here and by Copland and others (2009), surface topographic information, and observed glacier morphology all indicate that it is currently gaining mass. There is only modest evidence of frontal fluctuations (Mayer and others, 2006; Belò and others, 2008), but this is unsurprising given the estimated glacier response time, which is of the order of at least several decades for such a large ice mass (cf. equation (1b) of Jóhannesson and others (1989)).

In general terms, the pattern of linearly increasing flow with distance upstream from the terminus is typical of debris-covered glaciers observed elsewhere in the wider Himalaya (e.g. Scherler and others, 2008; Quincey and others, 2009). In several places, the velocity profile is complicated by the contributions of smaller tributary glaciers to the main Baltoro tongue, as well as the varying valley morphology. For example, the confluence of the two major glaciers at Concordia is one of the most interesting parts of the velocity profile, with the steep increase in flow clearly related to the convergence of the two masses. The decrease in flow immediately down-glacier is equally as steep, however, perhaps reflecting the wide valley morphology at this point. Similarly, another area of locally decreased flow, at around 19 km from the terminus, appears to reflect wide valley walls before the tributary of Yermanendu Glacier constricts the main glacier tongue again, increasing the velocity profile accordingly.

These general flow patterns are repeated year-on-year across the glacier surface, but with varying magnitude.

Medium-term (i.e. >5 years) patterns in flow are likely to reflect local climatic trends, but validating such linkages is difficult. There is a general paucity of climate station data available in the Karakoram, particularly at high elevations. Modelled climate data are therefore a useful source of precipitation and temperature information (Radiá and Hock, 2006). However, their accuracy is dependent upon the quality of observational data (coverage and volume), model parameterization schemes, and data assimilation methods. Several investigations have indicated concern associated with using ERA-40 re-analysis data (e.g. Allan and others, 2004; Bengtsson and others, 2004; Hagemann and others, 2005), particularly because regional biases in temperature and precipitation are known to occur. The regional bias for the Himalaya does not tend to vary seasonally, but positive precipitation biases have been found and appear to be more pronounced in the summer (Hagemann and others, 2005).

A number of studies have attempted to validate ERA-40 data specifically for assessing climate trends and glacier response to climate in high-altitude environments. Strong correlations between observed and modelled precipitation and temperature data have been found in alpine Switzerland (Kunz and others, 2007), the French Alps (Martin, 2004) and alpine Sweden (Radiá and Hock, 2006). There is no long-term climate station record for the high-altitude K2 region, other than that of the Skardu valley climate station at a much lower altitude. Consequently, there are no representative observational data to compare with ERA-40. Observational estimates of precipitation are available from TRMM data of the region from 1998 to 2001. There is good correlation between the two time series, indicating the reliability of our ERA-40 precipitation data. Furthermore, ERA-40 temperature trends correspond well with regional climate station temperature trends reported by Fowler and Archer (2006), indicating the ERA-40 temperature data are also of good quality.

Our seasonal analysis of ERA-40 data indicates that winter precipitation increased from 1958 to the early 1970s, and then decreased systematically. These data suggest that the medium-term speed-up of Baltoro Glacier is related to the increase in spring precipitation from the westerlies from 1970 to 1990, as precipitation rates are much higher than in winter, and the increase in precipitation during the summer can also add to the overall accumulation. For more recent years, seasonal analysis of TRMM data indicates an increase in winter precipitation from 1998 to 2006. An increase in precipitation also occurred during spring months from 2000 to 2006, with more accumulation observed in spring vs winter. In contrast, a decrease in summer precipitation was found for this time period. Therefore, increasing precipitation from the westerlies during the winter and spring may explain the non-summer speed-up of Baltoro Glacier since the mid-1990s.

Previous work has also shown statistically significant increases in winter and summer precipitation in the Karakoram since the early 1960s (Archer and Fowler, 2004), although widespread mass loss was observed in Karakoram glaciers for the majority of the 20th century, until the mid-1990s (Yao and others, 2006; Hewitt, 2007). The frequently heavy debris cover of these glaciers is likely to have provided a muted response to climatic effects (Benn and Lehmkuhl, 2000), and it is possible that recent changes are a symptom of the lag-time associated with the surface cover. As stated, simple response time theory (e.g. Jóhannesson,

1989) suggests that geometric changes on Baltoro Glacier may be possible from mass-balance perturbations on the timescale of decades, and modelling studies elsewhere have shown that rapid areal and volumetric changes (albeit recession) can take place under certain temperature–precipitation scenarios (e.g. Oerlemans and others, 1998). Nevertheless, glacier expansion in the Karakoram has occurred on a temporal scale less than the estimated response time of Baltoro Glacier. It has also occurred non-uniformly (Hewitt, 2005), indicating that factors other than, or in addition to, changing mass balance are influential in changing glacier velocities over the longer term.

The sudden speed-up of Baltoro Glacier observed during summer 2005 is an interesting departure from the gradual trend of increasing flow in recent years, and supplies evidence for meltwater playing an important role in glacier flow. Fluctuations in surface velocity have long been linked to the volume of meltwater available to the subglacial drainage system (Willis, 1995; Kavanaugh and Clarke, 2001; Purdie and others, 2008), particularly in cases where sliding is the predominant flow mechanism. Weather during the winter of 2004/05 was particularly anomalous in this region, with the international media reporting periods of record cold and prolonged snowfall that produced the deepest accumulations for decades, resulting in the burial of entire villages, emergency air drops of food supplies, >1000 deaths from hypothermia, house collapses and widespread avalanching (www.unicef.org/infobycountry/pakistan_25386.html; <http://edition.cnn.com/2005/WORLD/asiapcf/02/24/asia.cold.snap>). TRMM results presented here confirm that accumulation has been increasing in the region since 2000, and seasonal analyses show the precipitation increase has been particularly strong during winter months.

Our data indicate that, in the specific case of summer 2005, the rapid melting of the deep winter snowpack is likely to have introduced large volumes of meltwater to the glacier bed, thus increasing water pressure in subglacial channels and conduits that were not sized for such anomalously large water flows. In turn, this would have led to increased basal sliding and surface ice motion (cf. Kavanaugh and Clarke, 2001). At Concordia in particular, there are many large and deep supraglacial streams in the summer, especially where Baltoro South and Godwin Austen glaciers join. These streams are large enough to make navigation between these glaciers difficult, and field observations in summer 2005 indicate that they descend to the glacier bed via moulins a few kilometres downstream of Concordia. These moulins coincide with the start of the locations where the highest overall velocities are recorded (Fig. 3), and where the most marked speed-ups occurred in summer 2005 (Fig. 6). Overall, therefore, it is possible that the flow regime of the glacier is heavily influenced by both long-term mass variations and periodic summer meltwater availability.

Previous work on the seasonal variability in flow of Baltoro Glacier concluded that basal sliding plays a major part in ice transport (Mayer and others, 2006). Data presented here support this assertion, at least for the upper parts of the ablation area where the difference between summer and winter velocities is at its greatest. Up-glacier transverse velocity profiles are also indicative of a dominantly sliding flow regime, with their 'blocky' nature indicating the down-glacier movement of ice en masse, termed elsewhere as plug flow (Kick, 1962). The strong

consistency of summer and winter profiles in the lowermost 10 km of the glacier tongue indicates that deformation dominates there, rather than basal sliding. Similar conclusions were reached by Copland and others (2009).

CONCLUSIONS

The seasonal and temporal variability in surface ice motion exhibited by Baltoro Glacier over a 15 year period has been quantified by a combined field and remote-sensing based approach. The two datasets are highly complementary, with point-based field data revealing local variations in flow over short periods and image-based feature tracking providing glacier-wide flow fields over longer temporal windows. These data demonstrate that Baltoro Glacier is very active, even immediately up-glacier of its terminus. The highest surface velocities occurred during summer 2005, following a peak in precipitation the previous winter. These data indicate that the flow regime of the glacier is heavily influenced by meltwater availability as well as local-scale mass variability. Furthermore, large seasonal velocity gradients indicate a flow regime dominated by basal sliding, at least in parts. Winter velocities reveal a gradual speed-up of the glacier year-on-year since 2003, implying a glacier in positive mass balance, which is supported by our climate data showing increasing accumulation in the region since 2000.

In the wider Himalayan context, the Karakoram is certainly anomalous, at least because the glaciers here do not show evidence of stagnation over their lower reaches (Copland and others, 2009). However, the finer spatial and temporal variabilities in glacier dynamics in this region remain poorly understood, and their anticipated impact on water resources is largely unknown. Further work is required to establish the degree to which reported positive mass balances and glacier acceleration in the region have resulted in ice thickening (or possibly thinning, in response to extension) and terminal fluctuations. With information on both horizontal and vertical surface changes, the accurate modelling of glacier responses to continued climatic forcing should become possible, and the implications for water resources better understood.

ACKNOWLEDGEMENTS

Remote-sensing data were supplied under a European Space Agency (ESA) Category-1 agreement (5033-Quincey), and funding for D.J. Quincey was provided by Research Councils UK. Funding for L. Copland was provided by the US National Science Foundation, the US National Geographic Society, the Natural Sciences and Engineering Research Council of Canada, the Canadian Foundation for Innovation, the Ontario Research Fund and the University of Ottawa. Field assistance was provided by the K2 2005 medical and science teams, Nazir Sabir Expeditions and dedicated porters and guides. Funding for M. Bishop was provided by the US National Science Foundation (BCS-0242339), the US National Geographic Society, and the National Aeronautics and Space Administration (NNG04GL84G). The expedition 2004 was generously supported by the Ev-K2-CNR 'Scientific and Technological Research in Himalaya and Karakorum' project. We thank reviewers A. Käab and T. Bolch, whose comments helped to improve the quality of the paper.

REFERENCES

- Allan, R.P., M.A. Ringer, J.A. Pamment and A. Slingo. 2004. Simulation of the Earth's radiation budget by the European Centre for Medium-Range Weather Forecasts 40-year reanalysis (ERA40). *J. Geophys. Res.*, **109**(D18), D18107. (10.1029/2004JD004816.)
- Archer, D.R. and H.J. Fowler. 2004. Spatial and temporal variations in precipitation in the Upper Indus Basin: global teleconnections and hydrological implications. *Hydrol. Earth Syst. Sci.*, **8**(1), 47–61.
- Batura Glacier Investigation Group. 1979. The Batura Glacier in the Karakoram mountains and its variations. *Sci. Sin. B*, **22**(8), 958–974.
- Belò, M., G. Diolaiuti, C. Mayer, C. Mihalcea, C. Smiraglia and G. Vassena. 2008. Himalayan-Karakorum glaciers: results and problems in the study of recent variations of major non-polar glaciers. In Bonardi, L., ed. *Mountain glaciers and climate change in the last century*. Milan, Servizio Glaciologico Lombardo. (Terra Glacialis 10.)
- Bengtsson, L., S. Hagemann and K.I. Hodges. 2004. Can climate trends be calculated from reanalysis data? *J. Geophys. Res.*, **109**(D11), D11111. (10.1029/2004JD004536.)
- Benn, D.I. and F. Lehmkühl. 2000. Mass balance and equilibrium-line altitudes of glaciers in high mountain environments. *Quat. Int.*, **65/66**, 15–29.
- Berthier, E., Y. Arnaud, R. Kumar, S. Ahmad, P. Wagnon and P. Chevallier. 2007. Remote sensing estimates of glacier mass balances in the Himachal Pradesh (Western Himalaya, India). *Remote Sens. Environ.*, **108**(3), 327–338.
- Bolch, T., M. Buchroithner, T. Pieczonka and A. Kunert. 2008. Planimetric and volumetric glacier changes in the Khumbu Himal, Nepal, since 1962 using Corona, Landsat TM and ASTER data. *J. Glaciol.*, **54**(187), 592–600.
- Copland, L. and 8 others. 2009. Glacier velocities across the central Karakoram. *Ann. Glaciol.*, **50**(52), 41–49.
- Diolaiuti, G., M. Pecci and C. Smiraglia. 2003. Liligo Glacier, Karakoram, Pakistan: a reconstruction of the recent history of a surge-type glacier. *Ann. Glaciol.*, **36**, 168–172.
- Dyurgerov, M.B. and M.F. Meier. 2000. Twentieth century climate change: evidence from small glaciers. *Proc. Natl. Acad. Sci. USA (PNAS)*, **97**(4), 1406–1411.
- Fowler, H.J. and D.R. Archer. 2006. Conflicting signals of climatic change in the Upper Indus Basin. *J. Climate*, **19**(17), 4276–4293.
- Hagemann, S., K. Arpe and L. Bengtsson. 2005. Validation of the hydrological cycle of ERA-40. *ECMWF ERA-40 Proj. Rep. Ser.* 24.
- Hewitt, K. 1969. Glacier surges in the Karakoram Himalaya (Central Asia). *Can. J. Earth Sci.*, **6**(4, Part 2), 1009–1018.
- Hewitt, K. 2005. The Karakoram anomaly? Glacier expansion and the 'elevation effect', Karakoram Himalaya. *Mt. Res. Dev.*, **25**(4), 332–340.
- Hewitt, K. 2007. Tributary glacier surges: an exceptional concentration at Panmah Glacier, Karakoram Himalaya. *J. Glaciol.*, **53**(181), 181–188.
- Hewitt, K., C.P. Wake, G.J. Young and C. David. 1989. Hydrological investigations at Biafo Glacier, Karakorum Range, Himalaya; an important source of water for the Indus River. *Ann. Glaciol.*, **13**, 103–108.
- Huffman, G.J. and 8 others. 2007. The TRMM Multisatellite Precipitation Analysis (TMPA): quasi-global, multiyear, combined-sensor precipitation estimates at fine scales. *J. Hydromet.*, **8**(1), 38–55.
- Jóhannesson, T., C. Raymond and E. Waddington. 1989. Time-scale for adjustment of glaciers to changes in mass balance. *J. Glaciol.*, **35**(121), 355–369.
- Kavanaugh, J.L. and G.K.C. Clarke. 2001. Abrupt glacier motion and reorganization of basal shear stress following the establishment of a connected drainage system. *J. Glaciol.*, **47**(158), 472–480.
- Kerr, A. 1993. Topography, climate and ice masses: a review. *Terra Nova*, **5**(4), 332–342.
- Kick, W. 1962. Variations of some central Asiatic glaciers. *IASH Publ.* 58 (Symposium at Obergurgl 1962 – *Variations of the Regime of Existing Glaciers*), 223–229.
- Kunz, H., S.C. Scherrer, M.A. Liniger and C. Appenzeller. 2007. The evolution of ERA-40 surface temperatures and total ozone compared to observed Swiss time series. *Meteorol. Z.*, **16**(2), 171–181.
- Lucchitta, B.K., C.E. Rosanova and K.F. Mullins. 1995. Velocities of Pine Island Glacier, West Antarctica, from ERS-1 SAR images. *Ann. Glaciol.*, **21**, 277–283.
- Luckman, A., D.J. Quincey and S. Bevan. 2007. The potential of satellite radar interferometry and feature tracking for monitoring flow rates of Himalayan glaciers. *Remote Sens. Environ.*, **111**(2–3), 172–181.
- Lüdecke, C. and M. Kuhle. 1991. Comparison of meteorological observations at Mt. Everest and K2: examples of the 1984 and 1986 expedition. *Meteorol. Atmos. Phys.*, **47**(1), 55–60.
- Martin, E. 2004. Validation of Alpine snow in ERA-40. *ECMWF ERA-40 Proj. Rep. Ser.* 14.
- Mayer, C., A. Lambrecht, M. Belò, C. Smiraglia and G. Diolaiuti. 2006. Glaciological characteristics of the ablation zone of Baltoro glacier, Karakorum, Pakistan. *Ann. Glaciol.*, **43**, 123–131.
- Mihalcea, C., C. Mayer, G. Diolaiuti, A. Lambrecht, C. Smiraglia and G. Tartari. 2006. Ice ablation and meteorological conditions on the debris-covered area of Baltoro glacier, Karakoram, Pakistan. *Ann. Glaciol.*, **43**, 292–300.
- Mihalcea, C. and 7 others. 2008. Spatial distribution of debris thickness and melting from remote-sensing and meteorological data, at debris-covered Baltoro glacier, Karakoram, Pakistan. *Ann. Glaciol.*, **48**, 49–57.
- Oerlemans, J. and 10 others. 1998. Modelling the response of glaciers to climate warming. *Climate Dyn.*, **14**(4), 267–274.
- Purdie, H.L., M.S. Brook and I.C. Fuller. 2008. Seasonal variation in ablation and surface velocity on a temperate maritime glacier: Fox Glacier, New Zealand. *Arct. Antarct. Alp. Res.*, **40**(1), 140–147.
- Quincey, D.J., A. Luckman and D. Benn. 2009. Quantification of Everest region glacier velocities between 1992 and 2002, using satellite radar interferometry and feature tracking. *J. Glaciol.*, **55** (192), 596–606.
- Radiá, V. and R. Hock. 2006. Modeling future glacier mass balance and volume changes using ERA-40 reanalysis and climate models: sensitivity study at Storglaciären, Sweden. *J. Geophys. Res.*, **111**(F3), F03003. (10.1029/2005JF000440.)
- Scherler, D., S. Leprince and M.R. Strecker. 2008. Glacier-surface velocities in alpine terrain from optical satellite imagery – accuracy improvement and quality assessment. *Remote Sens. Environ.*, **112**(10), 3806–3819.
- Strozzi, T., A. Luckman, T. Murray, U. Wegmüller and C.L. Werner. 2002. Glacier motion estimation using satellite-radar offset-tracking procedures. *IEEE Trans. Geosci. Remote Sens.*, **40**(11), 2834–2391.
- Treydte, K.S. and 6 others. 2006. The twentieth century was the wettest period in northern Pakistan over the past millennium. *Nature*, **440**(7088), 1179–1182.
- Uppala, S.M. and 45 others. 2005. The ERA-40 re-analysis. *Q. J. R. Meteorol. Soc.*, **131**(612), 2961–3211.
- Willis, I.C. 1995. Intra-annual variations in glacier motion: a review. *Progr. Phys. Geogr.*, **19**(1), 61–106.
- Yadav, R.R., W.-K. Park, J. Singh and B. Dubey. 2004. Do the western Himalayas defy global warming? *Geophys. Res. Lett.*, **31**(17), L17201. (10.1029/2004GL020201.)
- Yao, T., J. Pu and S. Liu. 2006. Changing glaciers in High Asia. In Knight, P.G., ed. *Glacier science and environmental change*. Oxford, Blackwell, 275–282.