

 Open access • Journal Article • DOI:10.1017/S0004972715000751

Idempotent rank in the endomorphism monoid of a non-uniform partition

— [Source link](#) 

Igor Dolinka, James East, James D. Mitchell

Published on: 09 Apr 2015 - [arXiv: Group Theory](#)

Topics: [Idempotent matrix](#), [Semigroup](#), [Endomorphism](#) and [Idempotence](#)

Related papers:

- [Idempotent rank in the endomorphism monoid of a nonuniform partition](#)
- [Idempotent Elements of the Semigroups \$B_X\(D\)\$ Defined by Semilattices of the Class \$\Sigma_3\(X,8\)\$ When \$Z \neq \emptyset\$](#)
- [On the Singular Part of the Partition Monoid](#)
- [The rank of the endomorphism monoid of a uniform partition](#)
- [Idempotent Elements of the Semigroups \$B_X\(D\)\$ Defined by Semilattices of the Class \$\Sigma_3\(X,8\)\$ When \$Z = \emptyset\$](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/idempotent-rank-in-the-endomorphism-monoid-of-a-non-uniform-3rwwlk03sj>

Idempotent rank in the endomorphism monoid of a non-uniform partition

Igor Dolinka*

Department of Mathematics and Informatics

University of Novi Sad, Trg Dositeja Obradovića 4, 21101 Novi Sad, Serbia

`dockie@dmi.uns.ac.rs`

James East†

Centre for Research in Mathematics; School of Computing, Engineering and Mathematics

University of Western Sydney, Locked Bag 1797, Penrith NSW 2751, Australia

`J.East@uws.edu.au`

James D. Mitchell

Mathematical Institute, School of Mathematics and Statistics

University of St Andrews, St Andrews, Fife KY16 9SS, United Kingdom

`jdm3@st-and.ac.uk`

April 20, 2015

Abstract

We calculate the rank and idempotent rank of the semigroup $\mathcal{E}(X, \mathcal{P})$ generated by the idempotents of the semigroup $\mathcal{T}(X, \mathcal{P})$, which consists of all transformations of the finite set X preserving a non-uniform partition \mathcal{P} . We also classify and enumerate the idempotent generating sets of this minimal possible size. This extends results of the first two authors in the uniform case.

Keywords: Transformation semigroups, idempotents, generators, rank, idempotent rank.

MSC: 20M20; 20M17.

Dedicated to the memory of Prof. Gordon B. Preston.

1 Introduction

Let S be a monoid with identity 1, and $E(S) = \{s \in S : s^2 = s\}$ the set of all idempotents of S . For a subset $U \subseteq S$, we write $\langle U \rangle$ for the submonoid of S generated by U , which consists of all products $u_1 \cdots u_k$ with $u_1, \dots, u_k \in U \cup \{1\}$. The *rank* of S , denoted $\text{rank}(S)$, is the minimal cardinality of a subset $U \subseteq S$ such that $S = \langle U \rangle$. If S is idempotent generated, then the *idempotent rank* of S , denoted $\text{idrank}(S)$, is the minimal cardinality of a subset $U \subseteq E(S)$ such that $S = \langle U \rangle$.

The *full transformation semigroup* on a set X , denoted \mathcal{T}_X , is the set of all transformations of X (i.e., all functions $X \rightarrow X$), under the semigroup operation of composition. Let $\mathcal{P} = \{C_i : i \in I\}$ be a partition of X ; that is, the sets C_i are non-empty, pairwise disjoint, and their union is all of X . The set

$$\mathcal{T}(X, \mathcal{P}) = \{f \in \mathcal{T}_X : (\forall i \in I)(\exists j \in I) C_i f \subseteq C_j\},$$

consisting of all transformations of X preserving \mathcal{P} , is a subsemigroup of \mathcal{T}_X . A calculation of $\text{rank}(\mathcal{T}(X, \mathcal{P}))$ for finite X is given in [2] and [1] for the uniform and non-uniform cases, respectively. (Recall that \mathcal{P} is *uniform* if $|C_i| = |C_j|$ for all $i, j \in I$.) We write $\mathcal{E}(X, \mathcal{P}) = \langle E(\mathcal{T}(X, \mathcal{P})) \rangle$ for the idempotent generated subsemigroup of $\mathcal{T}(X, \mathcal{P})$. In [3], the first two authors calculated $\text{rank}(\mathcal{E}(X, \mathcal{P}))$ and $\text{idrank}(\mathcal{E}(X, \mathcal{P}))$ in the case of X being finite and \mathcal{P} uniform; among other things, it was shown that the rank and idempotent rank are equal, and the idempotent generating sets of this minimal possible size were also classified and enumerated. The purpose of the current work is to extend these results

*The first author gratefully acknowledges the support of Grant No. 174019 of the Ministry of Education, Science, and Technological Development of the Republic of Serbia, and Grant No. 1136/2014 of the Secretariat of Science and Technological Development of the Autonomous Province of Vojvodina.

†The second author gratefully acknowledges the support of the Glasgow Learning, Teaching, and Research Fund in partially funding his visit to the third author in July, 2014.

to the non-uniform case. Our main results include the classification and enumeration of the idempotents of $\mathcal{T}(X, \mathcal{P})$ (Propositions 3.1 and 3.2); the calculation of the rank and idempotent rank of $\mathcal{E}(X, \mathcal{P})$ (Theorem 3.16 — in particular, the rank and idempotent rank are equal unless \mathcal{P} has exactly two blocks of size 1 (and at least one other block)); and the classification and enumeration of all idempotent generating sets of the minimal possible size (Proposition 3.15 and Theorem 3.17).

2 Preliminaries

In this section, we state a number of results we will need concerning \mathcal{T}_X and $\mathcal{T}(X, \mathcal{P})$ for uniform \mathcal{P} . For the remainder of the article, we fix a finite set X . The group of units of \mathcal{T}_X is the *symmetric group* \mathcal{S}_X , which consists of all permutations of X (i.e., all bijections $X \rightarrow X$). Denote by $\mathcal{E}_X = \langle E(\mathcal{T}_X) \rangle$ the idempotent generated subsemigroup of \mathcal{T}_X . We generally denote the identity element of any monoid by 1; in particular, $1 \in \mathcal{T}_X$ denotes the identity map on X , which we also sometimes write as id_X . If $x, y \in X$ and $x \neq y$, then we write $e_{xy} \in \mathcal{T}_X$ for the transformation defined by

$$ze_{xy} = \begin{cases} x & \text{if } z = y \\ z & \text{if } z \in X \setminus \{y\}. \end{cases}$$

It is clear that $e_{xy} \in E(\mathcal{T}_X)$ for all x, y . We write $\mathcal{D}_X = \{e_{xy} : x, y \in X, x \neq y\}$. The next result collects several facts from [4–6]. We always interpret a binomial coefficient $\binom{m}{n}$ to be 0 if $m < n$.

Theorem 2.1. *Let X be a finite set with $|X| = n \geq 0$. Then*

$$\mathcal{E}_X = \langle \mathcal{D}_X \rangle = \{1\} \cup (\mathcal{T}_X \setminus \mathcal{S}_X).$$

Further, $\text{rank}(\mathcal{E}_X) = \text{idrank}(\mathcal{E}_X) = \rho_n$, where $\rho_2 = 2$ and $\rho_n = \binom{n}{2}$ if $n \neq 2$. □

The minimal idempotent generating sets of \mathcal{E}_X were characterised in [6] in terms of strongly connected tournaments. Such tournaments were enumerated in [7], and it was shown in [3] that any idempotent generating set for \mathcal{E}_X contains one of minimal size.

Theorem 2.2. *Let X be a finite set with $|X| = n \geq 0$. Then any idempotent generating set for $\mathcal{E}_X = \langle E(\mathcal{T}_X) \rangle$ contains an idempotent generating set of minimal possible size. The number of minimal idempotent generating sets for \mathcal{E}_X is equal to σ_n , where $\sigma_2 = 1$ and $\sigma_n = w_n$ for $n \neq 2$, and where the numbers w_n satisfy the recurrence*

$$w_0 = 1, \quad w_n = F_n - \sum_{s=1}^{n-1} \binom{n}{s} w_s F_{n-s} \quad \text{for } n \geq 1,$$

where $F_n = 2^{\binom{n}{2}} = 2^{n(n-1)/2}$. □

The following analogues of Theorems 2.1 and 2.2 were proved in [3].

Theorem 2.3. *Let $S = \mathcal{E}(X, \mathcal{P})$, where \mathcal{P} is a uniform partition of the finite set X into $m \geq 0$ blocks of size $n \geq 1$. Then $\text{rank}(S) = \text{idrank}(S) = \rho_{mn}$, where $\rho_{21} = 2$ and $\rho_{mn} = m\rho_n + n! \binom{m}{2}$ if $(m, n) \neq (2, 1)$. The numbers ρ_n are defined in Theorem 2.1. □*

Theorem 2.4. *Let $S = \mathcal{E}(X, \mathcal{P})$, where \mathcal{P} is a uniform partition of the finite set X into $m \geq 0$ blocks of size $n \geq 1$. Then any idempotent generating set of S contains an idempotent generating set of minimal possible size. The number of minimal idempotent generating sets for S is equal to σ_{mn} , where*

$$\sigma_{mn} = \begin{cases} 1 & \text{if } m = 0 \\ \sigma_n & \text{if } m = 1 \\ \sigma_m & \text{if } n = 1 \\ \sigma_n^m \times \sum_{k=0}^m \binom{m}{k} w_{nk} (2^{n!} - 2)^k & \text{if } m, n \geq 2. \end{cases}$$

The numbers σ_n are defined in Theorem 2.2, and the numbers w_{nk} satisfy the recurrence

$$w_{00} = 1, \quad w_{nk} = F_{nk} - \sum_{s=1}^{n-1} \binom{n}{s} \sum_{l=0}^k w_{sl} F_{n-s, k-l} \quad \text{for } n \geq 1,$$

where $F_{nk} = \binom{\binom{n}{2}}{k} \cdot 2^{\binom{n}{2}-k}$. □

Remark 2.5. It might seem odd to include the $m = 0$ case (when $X = \emptyset$) in the previous two results, but these will be useful for stating and proving later results such as Theorems 3.16 and 3.17.

3 The semigroup $\mathcal{E}(X, \mathcal{P})$

For a non-negative integer k , we write $[k] = \{1, \dots, k\}$, which we interpret to be empty if $k = 0$. We write $\mathcal{T}_k = \mathcal{T}_{[k]}$, and similarly for $\mathcal{S}_k, \mathcal{E}_k, \mathcal{D}_k$. Denote by $\mathcal{T}(k, l)$ the set of all functions $[k] \rightarrow [l]$, noting that $\mathcal{T}(k, k) = \mathcal{T}_k$. The *image*, *rank* and *kernel* of a function $f : A \rightarrow B$ are defined by

$$\text{im}(f) = \{af : a \in A\}, \quad \text{rank}(f) = |\text{im}(f)|, \quad \text{ker}(f) = \{(a, b) \in A \times A : af = bf\},$$

respectively. Obviously,

$$\text{im}(fg) \subseteq \text{im}(g), \quad \text{rank}(fg) \leq \min(\text{rank}(f), \text{rank}(g)), \quad \text{ker}(fg) \supseteq \text{ker}(f)$$

for all functions $f : A \rightarrow B$ and $g : B \rightarrow C$.

Recall that X is a fixed finite set. We also fix a *non-uniform* partition $\mathcal{P} = \{C_1, \dots, C_m\}$ of X . We will write $n_i = |C_i|$ for each i and assume that $n_1 \geq \dots \geq n_m$. We write $n = |X| = n_1 + \dots + n_m$. For convenience, we assume that $C_i = \{i\} \times [n_i]$ for each $i \in [m]$, so $X = \{(i, j) : i \in [m], j \in [n_i]\}$.

We now define some parameters associated to the partition \mathcal{P} that will make statements of results cleaner (see especially Theorems 3.16 and 3.17). For $i \in [n]$, define the sets

$$M_i = \{q \in [m] : n_q = i\} \quad \text{and} \quad N_i = \{j \in [i-1] : M_j \neq \emptyset\} = \{n_q : q \in [m], n_q < i\},$$

and put $\mu_i = |M_i|$ and $\nu_i = |N_i|$. In particular, μ_i is the number of blocks of \mathcal{P} of size i . As an example, if $n = 22$, $m = 8$ and $(n_1, \dots, n_8) = (5, 5, 3, 2, 2, 2, 2, 1)$, then the values of μ_i, ν_i are as follows:

i	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22
μ_i	1	4	1	0	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
ν_i	0	1	2	3	3	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4

Let $f \in \mathcal{T}(X, \mathcal{P})$. There is a transformation $\bar{f} \in \mathcal{T}_m$ such that, for all $i \in [m]$, $C_i f \subseteq C_{i\bar{f}}$. Also, for each $i \in [m]$, there is a function $f_i \in \mathcal{T}(n_i, n_{i\bar{f}})$ such that $(i, j)f = (i\bar{f}, jf_i)$ for all $j \in [n_i]$. The transformation $f \in \mathcal{T}(X, \mathcal{P})$ is uniquely determined by f_1, \dots, f_m, \bar{f} , and we will write $f = [f_1, \dots, f_m; \bar{f}]$. The product in $\mathcal{T}(X, \mathcal{P})$ may easily be described in terms of this notation. Indeed, if $f, g \in \mathcal{T}(X, \mathcal{P})$, then $fg = [f_1 g_{1\bar{f}}, \dots, f_m g_{m\bar{f}}; \bar{f}\bar{g}]$. Note that $\overline{fg} = \bar{f}\bar{g}$ and $(fg)_i = f_i g_{i\bar{f}} \in \mathcal{T}(n_i, n_{i\bar{f}\bar{g}})$ for all $f, g \in \mathcal{T}(X, \mathcal{P})$ and $i \in [m]$. When \mathcal{P} is uniform, each f_i belongs to $\mathcal{T}(n, n) = \mathcal{T}_n$ (where n is the common size of each block of \mathcal{P}), and $\mathcal{T}(X, \mathcal{P})$ is a wreath product $\mathcal{T}_n \wr \mathcal{T}_m$, as noted in [2].

There is a useful way to picture a transformation $f = [f_1, \dots, f_m; \bar{f}] \in \mathcal{T}(X, \mathcal{P})$. For example, with $m = 5$, and $\bar{f} = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 2 & 2 & 4 & 2 & 5 \end{pmatrix} \in \mathcal{T}_5$, the transformation $f = [f_1, f_2, f_3, f_4, f_5; \bar{f}]$ is pictured in Figure 1. (Note that these diagrams are not supposed to imply that the sets C_1, \dots, C_m have the same size.) This diagrammatic representation allows for easy




Figure 1: Diagrammatic representation of an element of $\mathcal{T}(X, \mathcal{P})$.

visualisation of the multiplication. For example, if f is as above, and if $g = [g_1, g_2, g_3, g_4, g_5; \bar{g}]$ where $\bar{g} = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 1 & 3 & 1 & 4 & 4 \end{pmatrix}$, then the product $fg = [f_1 g_{1\bar{g}}, f_2 g_{2\bar{g}}, f_3 g_{3\bar{g}}, f_4 g_{4\bar{g}}, f_5 g_{5\bar{g}}; \bar{f}\bar{g}]$ may be calculated as in Figure 2. Such diagrammatic methods may be used to verify various equations; an example is given in the proof of Proposition 3.3 (see Figure 4), but the rest are left to the reader.

The next result was proved in [3, Proposition 3.1] in the context of uniform partitions, but the argument works unmodified in the non-uniform case.

Proposition 3.1. *A transformation $f \in \mathcal{T}(X, \mathcal{P})$ is an idempotent if and only if*

- (i) $\bar{f} \in E(\mathcal{T}_m)$,
- (ii) $f_i \in E(\mathcal{T}_{n_i})$ for all $i \in \text{im}(\bar{f})$, and
- (iii) $\text{im}(f_i) \subseteq \text{im}(f_{i\bar{f}})$ for all $i \in [m] \setminus \text{im}(\bar{f})$.

□




Figure 2: Diagrammatic calculation of a product in $\mathcal{T}(X, \mathcal{P})$.

A formula for $|E(\mathcal{T}(X, \mathcal{P}))|$ was also given in [3, Proposition 3.1] in the uniform case. That formula seems impossible to extend to the non-uniform case, but we may give a recurrence analogous to that of [3, Proposition 3.2]. For a subset $A \subseteq [m]$, write $X_A = \bigcup_{a \in A} C_a$ and $\mathcal{P}_A = \{C_a : a \in A\}$. So \mathcal{P}_A is a partition of X_A (which is empty if A is empty).

Proposition 3.2. *Write $e(X, \mathcal{P}) = |E(\mathcal{T}(X, \mathcal{P}))|$. Then*

$$\begin{aligned}
 e(X, \mathcal{P}) &= 1 && \text{if } X \text{ is empty} \\
 e(X, \mathcal{P}) &= \sum_A e(X_{A^c}, \mathcal{P}_{A^c}) \sum_{a \in A} \sum_{l=1}^{n_a} \binom{n_a}{l} l^{n_a-l} && \text{if } X \text{ is non-empty,}
 \end{aligned}$$

where the outer sum is over all $A \subseteq [m]$ with $1 \in A$, and we write $A^c = [m] \setminus A$ and $n_A = |X_A| = \sum_{a \in A} n_a$.

Proof. The statement for X empty is clear, so suppose X is non-empty. An idempotent $f \in E(\mathcal{T}(X, \mathcal{P}))$ is uniquely determined by:

- (i) the set $A = \{i \in [m] : i\bar{f} = 1\bar{f}\}$,
- (ii) the element $a = 1\bar{f} \in A$ (note that $1\bar{f} \in A$ as \bar{f} is an idempotent),
- (iii) the image $\text{im}(f_a)$, say of size $l \in [n_a]$ — there are $\binom{n_a}{l}$ choices for these points, each of which are mapped identically by f_a ,
- (iv) the images under f of the elements of $(\bigcup_{b \in A} C_b) \setminus (\{a\} \times \text{im}(f_a))$, which must all be in $\{a\} \times \text{im}(f_a)$ — there are l^{n_a-l} choices for these images, and then finally
- (v) the restriction of f to $X_{A^c} = \bigcup_{i \in A^c} C_i$ — this restriction belongs to $E(\mathcal{T}(X_{A^c}, \mathcal{P}_{A^c}))$, which has size $e(X_{A^c}, \mathcal{P}_{A^c})$.

Multiplying these values and summing over relevant A, a, l gives the desired result. \square

We now move on to study the idempotent generated subsemigroup $\mathcal{E}(X, \mathcal{P}) = \langle E(\mathcal{T}(X, \mathcal{P})) \rangle$ of $\mathcal{T}(X, \mathcal{P})$. For simplicity, we will write $E = E(\mathcal{T}(X, \mathcal{P}))$ and $S = \mathcal{E}(X, \mathcal{P}) = \langle E \rangle$.

As in Section 2, for $k \geq 2$ and $i, j \in [k]$ with $i \neq j$, we write $e_{ij} \in \mathcal{T}_k$ for the idempotent transformation defined by

$$l e_{ij} = \begin{cases} i & \text{if } l = j \\ l & \text{if } l \in [k] \setminus \{j\}. \end{cases}$$

Note that k (the size of the set on which e_{ij} acts) depends on the context. For non-negative integers k, l , we write $\text{Inj}(k, l)$ (resp., $\text{Surj}(k, l)$) for the set of all injective (resp., surjective) functions $[k] \rightarrow [l]$. Note that $\text{Inj}(k, k) = \text{Surj}(k, k) = \mathcal{S}_k$, while if $k \neq l$, then (exactly) one of $\text{Inj}(k, l)$ or $\text{Surj}(k, l)$ is empty.

In what follows, certain special idempotents from E will play a crucial role. For $i, j \in [m]$ with $i \neq j$ and for any $f \in \text{Inj}(n_j, n_i)$ or $\text{Surj}(n_j, n_i)$, as appropriate, we write

$$e_{ij;f} = [1, \dots, 1, f, 1, \dots, 1; e_{ij}],$$

where f is in the j th position. Note that here $e_{ij} = \overline{e_{ij;f}}$ refers to the idempotent $e_{ij} \in \mathcal{T}_m$. The transformations $e_{ij;f}$ trivially satisfy conditions (i–iii) of Proposition 3.1, so $e_{ij;f} \in E$. If $k \in [m]$ and $g \in \mathcal{T}_{n_k}$, we will write $g^{(k)} = [1, \dots, 1, g, 1, \dots, 1; 1]$, where g is in the k th position. For example, with $m = 5$, the transformations $e_{42;f}$ and $g^{(2)}$ are pictured in Figure 3. For any $k \in [m]$, and for any subset $U \subseteq \mathcal{T}_{n_k}$, we write $U^{(k)} = \{g^{(k)} : g \in U\}$. If U is a subsemigroup of \mathcal{T}_{n_k} , then $U^{(k)}$ is a subsemigroup of $\mathcal{T}(X, \mathcal{P})$ and is isomorphic to U . Note that the k th coordinate of $e_{ij}^{(k)} = [1, \dots, 1, e_{ij}, 1, \dots, 1; 1]$ is $e_{ij} \in \mathcal{T}_{n_k}$.



Figure 3: Diagrammatic representation of $e_{42,f}$ (left) and $g^{(2)}$ (right) from $\mathcal{T}(X, \mathcal{P})$ with $m = 5$.

Proposition 3.3. *The semigroup $S = \mathcal{E}(X, \mathcal{P})$ is generated by $G_1 \cup G_2$, where*

$$G_1 = \{e_{ij}^{(k)} : k \in [m], i, j \in [n_k], i \neq j\} \quad \text{and} \quad G_2 = \{e_{ij;f} : i, j \in [m], i \neq j, f \in \text{Inj}(n_j, n_i) \cup \text{Surj}(n_j, n_i)\}.$$

Proof. Since the elements of $G_1 \cup G_2$ are idempotents, it suffices to show that $E \subseteq \langle G_1 \cup G_2 \rangle$. So let $f \in E$. Write A_1, \dots, A_r for the $\ker(\bar{f})$ -classes of $[m]$. Since f is an idempotent, we have $f = f_1 \cdots f_r$ where, for each $s \in [r]$, $f_s \in \mathcal{T}(X, \mathcal{P})$ is defined by

$$xf_s = \begin{cases} xf & \text{if } x \in X_{A_s} = \bigcup_{a \in A_s} C_a \\ x & \text{if } x \in X \setminus X_{A_s}. \end{cases}$$

So it suffices to show that $f_1, \dots, f_r \in \langle G_1 \cup G_2 \rangle$. Let $s \in [r]$, and write $A = A_s = \{a_1, \dots, a_k\}$. For simplicity, write $g = f_s = [g_1, \dots, g_m; \bar{g}]$. Without loss of generality, suppose $A\bar{g} = a_k$. By Proposition 3.1 and Theorem 2.1, we have $g_{a_k} \in E(\mathcal{T}_{n_{a_k}}) \subseteq \langle \mathcal{D}_{n_{a_k}} \rangle$, and it quickly follows that $g_{a_k}^{(a_k)} \in \langle G_1 \rangle$. In particular, if $k = |A| = 1$, then $g = g_{a_k}^{(a_k)} \in \langle G_1 \rangle$. So suppose $k \geq 2$. Now fix some $1 \leq j < k$. Let $e_j \in E(\mathcal{T}_{n_{a_j}})$ be such that $\ker(e_j) = \ker(g_{a_j})$, and let $h_j \in \text{Inj}(n_{a_j}, n_{a_k}) \cup \text{Surj}(n_{a_j}, n_{a_k})$ be any injective or surjective (as appropriate) map that extends the map $h'_j : \text{im}(e_j) \rightarrow \text{im}(g_{a_k})$ defined by $(xe_j)h'_j = xg_{a_j}$ for $x \in [n_{a_j}]$. In Figure 4, we show that

$$g = e_1^{(a_1)} \cdots e_{k-1}^{(a_{k-1})} \cdot g_{a_k}^{(a_k)} \cdot e_{a_k a_1; h_1} \cdots e_{a_k a_{k-1}; h_{k-1}}.$$

(In the diagram, we only picture the action of the transformations on $X_A = C_{a_1} \cup \cdots \cup C_{a_k}$, and the pictured ordering of the blocks is not meant to imply that $a_1 < \cdots < a_k$.) Again, each $e_j^{(a_j)}$ belongs to $\langle G_1 \rangle$, and clearly each $e_{a_k a_j; h_j}$ belongs to G_2 . This completes the proof. \square




Figure 4: Diagrammatic proof that $g = e_1^{(a_1)} \cdots e_{k-1}^{(a_{k-1})} \cdot g_{a_k}^{(a_k)} \cdot e_{a_k a_1; h_1} \cdots e_{a_k a_{k-1}; h_{k-1}}$; see the proof of Proposition 3.3 for more details.

Our next task is to calculate $\text{rank}(S)$ and $\text{idrank}(S)$. In order to do this, we will show that the generating set $G_1 \cup G_2$ from Proposition 3.3 may be significantly reduced in size. The next sequence of results (specifically, Lemmas 3.4, 3.6, 3.8 and 3.13) show what kind of transformations are essential in any (idempotent) generating set.

Recall that $\mathcal{E}_{n_i} = \langle E(\mathcal{T}_{n_i}) \rangle = \{1\} \cup (\mathcal{T}_{n_i} \setminus \mathcal{S}_{n_i})$. So $\mathcal{E}_{n_i}^{(i)}$, which consists of all maps $[1, \dots, 1, f, 1, \dots, 1; 1] \in \mathcal{T}(X, \mathcal{P})$ with $f \in \mathcal{E}_{n_i}$ in the i th position, is a subsemigroup of S isomorphic to \mathcal{E}_{n_i} . The proof of [3, Lemma 4.3] is easily adapted to show the following.

Lemma 3.4. *Let $i \in [m]$. Then $S \setminus \mathcal{E}_{n_i}^{(i)}$ is an ideal of S . Consequently, any generating set for S contains a generating set for $\mathcal{E}_{n_i}^{(i)}$. \square*

Since the map $\mathcal{T}(X, \mathcal{P}) \rightarrow \mathcal{T}_m : f \mapsto \bar{f}$ is a homomorphism, it follows from Proposition 3.1 that $\bar{f} \in \mathcal{E}_m = \{1\} \cup (\mathcal{T}_m \setminus \mathcal{S}_m)$ for all $f \in S$. We will frequently make use of this fact. The next simple result describes the preimage of $1 \in \mathcal{T}_m$ under the above map.

Lemma 3.5. *Let $f \in S$. If $\bar{f} = 1$, then $f_i \in \mathcal{E}_{n_i}$ for all $i \in [m]$.*

Proof. Let $f = h_1 \cdots h_k$, where $h_1, \dots, h_k \in E$, and write $h_j = [h_{j1}, \dots, h_{jm}; \bar{h}_j]$ for each j . Since $1 = \bar{f} = \bar{h}_1 \cdots \bar{h}_k$, we see that $\bar{h}_j = 1$ for all j . It follows that $h_{ji} \in E(\mathcal{T}_{n_i})$ for each i, j . So $f_i = h_{1i} \cdots h_{ki} \in \langle E(\mathcal{T}_{n_i}) \rangle = \mathcal{E}_{n_i}$ for each i . \square

For $1 \leq i < j \leq m$, we write $\varepsilon_{ij} = \varepsilon_{ji}$ for the equivalence relation on $[m]$ with unique non-trivial equivalence class $\{i, j\}$. Note that $\ker(e_{ij}) = \ker(e_{ji}) = \varepsilon_{ij}$. We also write $\Delta = \{(i, i) : i \in [m]\}$ for the trivial equivalence on $[m]$ (i.e., the equality relation on $[m]$).

Lemma 3.6. *Let $1 \leq i < j \leq m$ and $f \in \text{Inj}(n_j, n_i)$, and suppose $e_{ij;f} = gh$ where $g, h \in S$ and $g \neq 1$. Then*

- (i) $\ker(\bar{g}) = \varepsilon_{ij}$,
- (ii) g_1, \dots, g_m are injective,
- (iii) $g_i \in \mathcal{S}_{n_i}$ and $g_j g_i^{-1} = f$.

Consequently, any generating set for S contains such an element g for each such i, j, f . Further, if g is an idempotent, then

- (iv) if $n_i = n_j$, then either $g = e_{ij;f}$ or $g = e_{ji;f^{-1}}$,
- (v) if $n_i > n_j$, then $g = e_{ij;f}$.

Proof. Now, $[1, \dots, 1, f, 1, \dots, 1; e_{ij}] = e_{ij;f} = gh = [g_1 h_{1\bar{g}}, \dots, g_m h_{m\bar{g}}; \bar{g}\bar{h}]$. Since each $g_r h_{r\bar{g}}$ is injective (equal to either 1 or f), it follows that each g_r is injective, establishing (ii). If $\bar{g} = 1$, then $g_r \in \mathcal{E}_{n_r}$ for each r by Lemma 3.5; but the only injective element of \mathcal{E}_{n_r} is the identity element 1, so $g_r = 1$ for all r , giving $g = [1, \dots, 1; 1] = 1$, a contradiction. So $\bar{g} \neq 1$, whence $\bar{g} \in \mathcal{T}_m \setminus \mathcal{S}_m$. But then $\Delta \neq \ker(\bar{g}) \subseteq \ker(\bar{g}\bar{h}) = \ker(e_{ij}) = \varepsilon_{ij}$, so that $\ker(\bar{g}) = \varepsilon_{ij}$, giving (i). Put $k = i\bar{g} = j\bar{g}$.

Next we claim that $n_k = n_i$. Indeed, suppose this was not the case. Since $g_i \in \text{Inj}(n_i, n_k)$, we have $n_i \leq n_k$, so we must in fact have $n_i < n_k$. Let $L = \{r \in [m] : n_r > n_i\}$, noting that $k \in L$. Since $n_1 \geq \dots \geq n_m$, it follows that $L = [s]$ for some $s \geq 1$. Now, since \bar{g} maps $[m] \setminus \{i, j\}$ injectively into $[m] \setminus \{k\}$, and since $s < i < j$, it follows that there exists $r \in L = [s]$ such that $r\bar{g} > s$. But then $g_r \in \mathcal{T}(n_r, n_{r\bar{g}})$ with $n_{r\bar{g}} \leq n_i < n_r$, contradicting the fact that g_r is injective. This completes the proof of the claim.

In particular, it follows that $g_i \in \text{Inj}(n_i, n_k) = \text{Inj}(n_i, n_i) = \mathcal{S}_{n_i}$. Also, $h_k = h_{i\bar{g}} \in \mathcal{T}(n_k, n_i)$ since $i = ie_{ij} = i\bar{g}\bar{h} = k\bar{h}$. We also have $1 = g_i h_k$, so that $h_k = g_i^{-1}$, from which it follows that $f = g_j h_{j\bar{g}} = g_j h_k = g_j g_i^{-1}$, completing the proof of (iii).

Next, suppose G is an arbitrary generating set for S . By considering an expression $e_{ij;f} = h_1 \cdots h_k$, where $h_1, \dots, h_k \in G \setminus \{1\}$, we see that $h_1 \in G$ satisfies conditions (i–iii).

Finally, suppose g is an idempotent. Since $\bar{g} \in E(\mathcal{T}_m)$ by Proposition 3.1, and since $\ker(\bar{g}) = \varepsilon_{ij}$, it follows that $\bar{g} = e_{ij}$ or $\bar{g} = e_{ji}$. Suppose first that $\bar{g} = e_{ij}$. Proposition 3.1 also gives $g_r \in E(\mathcal{T}_{n_r})$ for each $r \in \text{im}(\bar{g}) = [m] \setminus \{j\}$; but each g_r is injective, so it follows that $g_r = 1$ if $r \neq j$. We also have $f = g_j g_i^{-1} = g_j$, giving $g = e_{ij;f}$. Next suppose $\bar{g} = e_{ji}$. In particular, $n_i = n_j$ as $g_i \in \text{Inj}(n_i, n_j)$ and $n_i \geq n_j$. Again, we have $g_r = 1$ for all $r \in [m] \setminus \{i\}$, and this time we have $f = g_j g_i^{-1} = g_i^{-1}$, which gives $g_i = f^{-1}$, and $g = e_{ji;f^{-1}}$. This completes the proof. \square

Let $i, j \in [m]$ with $n_i > n_j$. As a consequence of the previous result, we see that any idempotent generating set G for S contains all of $\{e_{ik;f} : k \in [m], n_k = n_j, f \in \text{Inj}(n_j, n_i)\}$. It might be tempting to guess that G must also contain all of $\{e_{ki;f} : k \in [m], n_k = n_j, f \in \text{Surj}(n_i, n_j)\}$. But this is far from the case. In fact, G need only contain a *single* element of the latter set, as we show in Lemma 3.8 and Proposition 3.15, the proof of which requires the next technical result (which will also be useful elsewhere).

Lemma 3.7. *Let $j \in [n]$ and put $L = \{q \in [m] : n_q > j\}$. Suppose $f \in S$ is such that $\bar{f}|_L$ is injective and $|C_q f| > j$ for all $q \in L$. Then $\bar{f}|_L = \text{id}_L$ and $f_q \in \mathcal{E}_{n_q}$ for all $q \in L$.*

Proof. If $L = \emptyset$, there is nothing to show, so suppose $L \neq \emptyset$. By Proposition 3.3, we may write $f = g_1 \cdots g_k$ where $g_1, \dots, g_k \in E$ and $\text{rank}(\bar{g}_l) \geq m-1$ for each $l \in [k]$. (We could be more specific by insisting that $g_1, \dots, g_k \in G_1 \cup G_2$, but it will be convenient later to argue more generally, as we do here.) We claim that $\bar{g}_l|_L = \text{id}_L$ for each l . Indeed, suppose this is not the case, and let $l \in [k]$ be minimal so that $\bar{g}_l|_L \neq \text{id}_L$. Since $\bar{g}_l \neq 1$, it follows that $\text{rank}(\bar{g}_l) = m-1$, so $\bar{g}_l = e_{ab}$ for some $a, b \in [m]$ with $a \neq b$. Note that at least one of a, b belongs to L since $\bar{g}_l|_L \neq \text{id}_L$. We could not have $a, b \in L$ or else then $(a\bar{g}_1 \cdots \bar{g}_{l-1})\bar{g}_l = a\bar{g}_l = b\bar{g}_l = (b\bar{g}_1 \cdots \bar{g}_{l-1})\bar{g}_l$, giving $a\bar{f} = b\bar{f}$, contradicting the assumption that $\bar{f}|_L$ is injective. We also could not have $a \in L$ or else then $b \in [m] \setminus L$ so that $\bar{g}_l|_L = \text{id}_L$, another contradiction. So $a \in [m] \setminus L$ and $b \in L$. But then $|C_b f| = |C_b g_1 \cdots g_k| \leq |(C_b g_1 \cdots g_{l-1})g_l| \leq |C_b g_l| \leq |C_a| \leq j$, contradicting the assumption that $|C_q f| > j$ for all $q \in L$. This establishes the claim that $\bar{g}_l|_L = \text{id}_L$ for each l . It obviously follows that $\bar{f}|_L = \text{id}_L$. For $l \in [k]$, write $g_l = [g_{l1}, \dots, g_{lm}; \bar{g}_l]$. Since $L \subseteq \text{im}(\bar{g}_l)$ and $g_l \in E$ for each $l \in [k]$, it follows from Proposition 3.1 that $g_{lq} \in E(\mathcal{T}_{n_q})$ for each $l \in [k]$ and $q \in L$. Since $\bar{g}_l|_L = \text{id}_L$ for all $l \in [k]$, it follows that $f_q = g_{1q} \cdots g_{kq} \in \mathcal{E}_{n_q}$ for each $q \in L$. \square

Lemma 3.8. *Let $i, j \in [m]$ be such that $n_i > n_j$, and let $f \in \text{Surj}(n_i, n_j)$. Suppose $e_{ji;f} = g_1 \cdots g_r$ where $g_1, \dots, g_r \in S \setminus \{1\}$. Put $L = \{q \in [m] : n_q > n_j\}$. Let $l \in [r]$ be minimal so that $\bar{g}_l|_L \neq \text{id}_L$. Let $h = g_l$ and write $h = [h_1, \dots, h_m; \bar{h}]$. Then*

- (i) $\text{rank}(\bar{h}) = m-1$, (ii) $|C_i h| = n_j$, and (iii) h_q is injective for all $q \in L \setminus \{i\}$.

Consequently, any generating set for S contains such an element h for each such i, j . Further, if h is an idempotent, then $h = e_{ki;f'}$ for some $k \in [m]$ with $n_k = n_j$ and some $f' \in \text{Surj}(n_i, n_j)$.

Proof. First note that $\bar{h} \neq 1$ since $\bar{h}|_L \neq \text{id}_L$, so $\bar{h} \in \mathcal{T}_m \setminus \mathcal{S}_m$ and $\text{rank}(\bar{h}) \leq m-1$. But also $\text{rank}(\bar{h}) \geq \text{rank}(e_{ji}) = m-1$, so (i) holds.

Next, suppose (ii) does not hold. First note that $n_j = |C_j| = |C_i e_{ji;f}| = |C_i g_1 \cdots g_r| \leq |(C_i g_1 \cdots g_{l-1})h| \leq |C_i h|$, since $\bar{g}_1 \cdots \bar{g}_{l-1}$ acts as the identity on L and $i \in L$. So, since we are assuming that $|C_i h| \neq n_j$, we must have $|C_i h| > n_j$. A similar calculation shows that $n_q \leq |C_q h|$ for any $q \in L \setminus \{i\}$. In particular, together with the assumption that $|C_i h| > n_j$, this gives $|C_q h| > n_j$ for all $q \in L$. Since $(\bar{g}_1 \cdots \bar{g}_{l-1} \bar{h} \bar{g}_{l+1} \cdots \bar{g}_r)|_L = e_{ji}|_L$ is injective, and since $(\bar{g}_1 \cdots \bar{g}_{l-1})|_L = \text{id}_L$, it follows that $\bar{h}|_L$ is injective. But then Lemma 3.7 says that $\bar{h}|_L = \text{id}_L$, a contradiction. This completes the proof of (ii).

Next, let $q \in L \setminus \{i\}$ be arbitrary. We have already seen that $|C_q h| \geq n_q$. But we also trivially have $|C_q h| \leq |C_q| = n_q$, whence $h|_{C_q}$ is injective, and (iii) holds. As in the proof of Lemma 3.6, the statement about the generating set G follows quickly.

Finally, suppose h is an idempotent. Since $\bar{h} \in E(\mathcal{T}_m)$ and $\text{rank}(\bar{h}) = m-1$, it follows that $\bar{h} = e_{ab}$ for some $a, b \in [m]$ with $a \neq b$. Since $(\bar{g}_1 \cdots \bar{g}_{l-1})|_L = \text{id}_L$ but $(\bar{g}_1 \cdots \bar{g}_{l-1} \bar{h})|_L \neq \text{id}_L$, we again conclude that $a \in [m] \setminus L$ and $b \in L$. We observe that $b = i$. Indeed, if this was not the case, then we would have $|C_b h| \leq |C_a| = n_a < n_b$, contradicting the fact that h_q is injective for all $q \in L \setminus \{i\}$. In particular, $\bar{h} = e_{ai}$. So h maps C_i into C_a , which gives $n_j = |C_i h| \leq |C_a| = n_a$. But also $n_a \leq n_j$ since $a \in [m] \setminus L$, so it follows that $n_a = n_j$. So far we know that

$$h = [h_1, \dots, h_i, \dots, h_m; e_{ai}].$$

Since we know that h maps C_i into C_a and $|C_i h| = n_j = |C_a|$, it follows that $h_i \in \text{Surj}(n_i, n_j)$. We wish to show that $h = e_{ai;h_i}$. This will complete the proof of the lemma (with $k = a$ and $f' = h_i$). It remains to show that $h_q = 1$ for all $q \in [m] \setminus \{i\}$. In fact, since h is an idempotent, and since $\text{im}(\bar{h}) = \text{im}(e_{ai}) = [m] \setminus \{i\}$, we already know that $h_q \in E(\mathcal{T}_{n_q})$ for all $q \in [m] \setminus \{i\}$, so it suffices to show that each such h_q is injective. We already know this is the case for $q \in L \setminus \{i\}$. We also know that $C_a = C_i h \subseteq C_a h$ by Proposition 3.1(iii), so it follows that h_a is surjective. But all surjective transformations of a finite set are injective, so it follows that h_a is injective. It remains to establish the injectivity of each h_q with $q \in [m] \setminus (L \cup \{a\})$. To do this, we must consider two separate cases. To simplify notation, put $u = g_1 \cdots g_{l-1}$ and $v = g_l \cdots g_r$, and write $u = [u_1, \dots, u_m; \bar{u}]$ and $v = [v_1, \dots, v_m; \bar{v}]$. Since $e_{ji;f} = uv = [u_1 v_{1\bar{u}}, \dots, u_m v_{m\bar{u}}; \bar{u}\bar{v}]$, it follows that u_q is injective for all $q \in [m] \setminus \{i\}$. Similarly, $u_q h_{q\bar{u}}$ is injective for all such q .

Case 1. Suppose first that $\bar{u} = 1$. Then $u_q \in \mathcal{E}_{n_q}$ for each $q \in [m]$. Since u_q is injective for each $q \in [m] \setminus \{i\}$, it follows that $u_q = 1$ for each such q . So

$$uh = [u_1 h_1, \dots, u_m h_m; e_{ai}] = [h_1, \dots, h_{i-1}, u_i h_i, h_{i+1}, \dots, h_m; e_{ai}].$$

In particular, for any $q \in [m] \setminus \{i\}$, $h_q = u_q h_q = u_q h_{q\bar{u}}$ is injective, as noted above. This completes the proof in this case.

Case 2. Finally, suppose $\bar{u} \neq 1$. So $\bar{u} \in \mathcal{T}_m \setminus \mathcal{S}_m$. Since $\varepsilon_{ij} = \ker(e_{ji}) = \ker(\bar{u}\bar{v}) \supseteq \ker(\bar{u}) \neq \Delta$, it follows that $\ker(\bar{u}) = \varepsilon_{ij}$. We claim that $a \notin \text{im}(\bar{u})$. Indeed, suppose this was not the case, so $a = c\bar{u}$ for some $c \in [m]$. Since $\bar{u}|_L = \text{id}_L$ and $j\bar{u} = i\bar{u}$, it follows that \bar{u} maps $L \cup \{j\}$ into L , so $c \in [m] \setminus (L \cup \{j\})$. But then

$$ce_{ji} = c\bar{g}_1 \cdots \bar{g}_r = c\bar{u}e_{ai}\bar{g}_{l+1} \cdots \bar{g}_r = ae_{ai}\bar{g}_{l+1} \cdots \bar{g}_r = ie_{ai}\bar{g}_{l+1} \cdots \bar{g}_r = i\bar{u}e_{ai}\bar{g}_{l+1} \cdots \bar{g}_r = ie_{ji} = j,$$

a contradiction, since $c \notin \{i, j\}$. This completes the proof of the claim that $a \notin \text{im}(\bar{u})$. Since $\text{rank}(\bar{u}) = m - 1$, it follows that $\text{im}(\bar{u}) = [m] \setminus \{a\}$. Let $Q = [m] \setminus L$, and put

$$Y = \bigcup_{q \in Q \setminus \{j\}} C_q \quad \text{and} \quad Z = \bigcup_{q \in Q \setminus \{a\}} C_q.$$

Since $|C_a| = n_j = |C_j|$, it follows that $|Y| = |Z|$. Since \bar{u} maps $Q \setminus \{j\}$ bijectively onto $Q \setminus \{a\}$, and since u_p is injective for all $p \in Q$, it follows that u_p is bijective for each $p \in Q \setminus \{j\}$. Now let $q \in Q \setminus \{a\}$ be arbitrary. Since $Q \setminus \{a\} = [m] \setminus (L \cup \{a\})$, the proof of the lemma will be complete if we can show that h_q is injective. Put $p = q\bar{u}^{-1} \in Q \setminus \{j\}$. Note that

$$uh = [u_1 h_{1\bar{u}}, \dots, u_p h_q, \dots, u_m h_{m\bar{u}}; \bar{u}e_{ai}].$$

Since $u_p h_q$ is injective, and since u_p is bijective, it follows that h_q is injective. As noted above, this completes the proof of the lemma. \square

We are now able to give a lower bound for $\text{rank}(S)$. The next result is stated in terms of the parameters μ_i, ν_i, ρ_i introduced at the beginning of this section and in Theorem 2.1.

Corollary 3.9. *We have $\text{rank}(S) \geq \rho$, where*

$$\rho = \sum_{i=1}^n \left(\mu_i \rho_i + i! \binom{\mu_i}{2} + \mu_i \nu_i \right) + \sum_{1 \leq i < j \leq n} \mu_i \mu_j \frac{j!}{(j-i)!}.$$

Proof. Let G be an arbitrary generating set for S . By Lemma 3.4, G contains a generating set for $\mathcal{E}_{n_r}^{(r)}$ for each $r \in [m]$. These are pairwise disjoint, and each has size at least $\text{rank}(\mathcal{E}_{n_r}) = \rho_{n_r}$, so G contains at least

$$\sum_{r=1}^m \rho_{n_r} = \sum_{i=1}^n \mu_i \rho_i \tag{3.9.1}$$

transformations coming from these generating sets of $\mathcal{E}_{n_1}^{(1)}, \dots, \mathcal{E}_{n_m}^{(m)}$.

Next, fix some $i \in [n]$ with $\mu_i \geq 2$. Lemma 3.6 tells us that for each $p, q \in M_i$ with $p < q$, and for each $f \in \mathcal{S}_i$, G contains some transformation g such that

$$(i) \ker(\bar{g}) = \varepsilon_{pq}, \quad (ii) g_1, \dots, g_m \text{ are injective}, \quad (iii) g_p \in \mathcal{S}_i \text{ and } g_q g_p^{-1} = f.$$

(For future reference, we note that if this g is an idempotent, then Lemma 3.6 gives $g = e_{ij;f}$ or $e_{ji;f^{-1}}$.) There are $\binom{\mu_i}{2}$ such p, q , and there are $i!$ such f . Summing over all appropriate i , and noting that $\binom{\mu_i}{2} = 0$ if $\mu_i \leq 1$, we see that G contains at least

$$\sum_{\substack{i \in [n] \\ \mu_i \geq 2}} i! \binom{\mu_i}{2} = \sum_{i=1}^n i! \binom{\mu_i}{2} \tag{3.9.2}$$

transformations of this type.

Next, suppose $1 \leq p < q \leq m$ are such that $n_p > n_q$. Let $f \in \text{Inj}(n_q, n_p)$ be arbitrary. Lemma 3.6 tells us that G must contain a transformation g such that

$$(i) \ker(\bar{g}) = \varepsilon_{pq}, \quad (ii) g_1, \dots, g_m \text{ are injective}, \quad (iii) g_p \in \mathcal{S}_{n_p} \text{ and } g_q g_p^{-1} = f.$$

There are $|\text{Inj}(n_q, n_p)| = n_p! / (n_p - n_q)!$ such transformations. For $1 \leq i < j \leq n$, there are $\mu_i \mu_j$ choices of $1 \leq p < q \leq m$ with $j = n_p$ and $i = n_q$, so G contains at least

$$\sum_{\substack{1 \leq p < q \leq m \\ n_p > n_q}} \frac{n_p!}{(n_p - n_q)!} = \sum_{1 \leq i < j \leq n} \mu_i \mu_j \frac{j!}{(j-i)!} \tag{3.9.3}$$

transformations of this type.

Finally, let $i \in [n]$ be such that $\mu_i \neq 0$, and suppose $p \in [m]$ is such that $n_p = i$. Let $L = \{q \in [m] : n_q \geq i\}$. If $1 \leq j < i$ is such that $\mu_j \neq 0$, then Lemma 3.8 says that G must contain some transformation g such that

$$(i) \text{rank}(\bar{g}) = m - 1, \quad (ii) |C_p g| = j, \text{ and} \quad (iii) g_q \text{ is injective for all } q \in L \setminus \{i\}.$$

There are μ_i such p , and ν_i such j . So G contains at least

$$\sum_{\substack{i \in [n] \\ \mu_i \neq 0}} \mu_i \nu_i = \sum_{i=1}^n \mu_i \nu_i \quad (3.9.4)$$

transformations of this type. Finally, adding equations (3.9.1–3.9.4) shows that $|G| \geq \rho$. Since G is an arbitrary generating set, the result follows. \square

We show below that this lower bound for $\text{rank}(S)$ is precise (see Theorem 3.16). In fact, we will also show that $\text{idrank}(S) = \text{rank}(S)$ apart from the special case in which $\mu_1 = 2$. In order to deal with that case, we need Lemma 3.13 below, which will also be useful when we later classify and enumerate the minimal idempotent generating sets for S (Theorem 3.17). But first we need a number of technical results.

Let $i \in [n]$. Recall that $M_i = \{q \in [m] : n_q = i\}$. Let $X_i = \bigcup_{q \in M_i} C_q$, and put

$$S_i = \{f \in S : f|_{X \setminus X_i} = \text{id}_{X \setminus X_i}, X_i f \subseteq X_i\}.$$

The reader should not confuse S_i with \mathcal{S}_i , the symmetric group on $[i]$. Let $\mathcal{P}_i = \{C_q : q \in M_i\}$. So \mathcal{P}_i is a uniform partition of X_i into μ_i blocks of size i . We aim to show that S_i is isomorphic to $\mathcal{E}(X_i, \mathcal{P}_i)$, the idempotent generated subsemigroup of $\mathcal{T}(X_i, \mathcal{P}_i)$. The following was proved in [3, Proposition 4.1].

Proposition 3.10. *Let $i \in [n]$ and write $M_i = \{q_1, \dots, q_{\mu_i}\}$. Then $f = [f_{q_1}, \dots, f_{q_{\mu_i}}; \bar{f}] \in \mathcal{T}(X_i, \mathcal{P}_i)$ belongs to $\mathcal{E}(X_i, \mathcal{P}_i)$ if and only if one of the following holds:*

- (i) $\bar{f} = 1$ and $f_{q_1}, \dots, f_{q_{\mu_i}} \in \mathcal{E}_i$, (ii) $\bar{f} \in \mathcal{T}_{X_i} \setminus \mathcal{S}_{X_i}$. \square

Lemma 3.11. *Let $i \in [n]$. Then S_i is isomorphic to $\mathcal{E}(X_i, \mathcal{P}_i)$.*

Proof. There is an obvious embedding $\phi : \mathcal{T}(X_i, \mathcal{P}_i) \rightarrow \mathcal{T}(X, \mathcal{P})$ defined, for $f \in \mathcal{T}(X_i, \mathcal{P}_i)$, by

$$x(f\phi) = \begin{cases} xf & \text{if } x \in X_i \\ x & \text{if } x \in X \setminus X_i. \end{cases}$$

So $\mathcal{E}(X_i, \mathcal{P}_i)$ is isomorphic to its image, $T = \mathcal{E}(X_i, \mathcal{P}_i)\phi$. It remains to show that $S_i = T$. Clearly, $T \subseteq S_i$. Conversely, suppose $f \in S_i$, and put $g = f|_{X_i} \in \mathcal{T}(X_i, \mathcal{P}_i)$. Obviously, $f = g\phi$, so it suffices to prove that $g \in \mathcal{E}(X_i, \mathcal{P}_i)$. But this follows quickly from Lemma 3.5 and Proposition 3.10. \square

Next, we aim to show that any idempotent generating set for S must contain a generating set for each S_i . To do this, we require the next technical result.

Lemma 3.12. *Let $r \in [n]$ with $\mu_r \neq 0$, and let $h \in S$ be such that $\text{rank}(\bar{h}) = m - 1$, $M_r \bar{h} \subseteq M_r$, $|M_r \bar{h}| = \mu_r - 1$, and h_q is injective for all $q \in [m]$. Suppose also that $g \in E \setminus S_r$ is such that $\text{rank}(\bar{h}\bar{g}) = m - 1$ and $(hg)_q$ is injective for all $q \in [m]$. Then $g|_Y = \text{id}_Y$ where $Y = X_r h$. In particular, $(hg)|_{X_r} = h|_{X_r}$.*

Proof. It is clear that $(hg)|_{X_r} = h|_{X_r}$ follows from $g|_Y = \text{id}_Y$, so we just prove the latter. By assumption, we have $M_r \bar{h} = M_r \setminus \{a\}$ for some $a \in M_r$. Note that $m - 1 = \text{rank}(\bar{h}\bar{g}) \leq \text{rank}(\bar{g}) \leq m$. We now break up the proof into cases, according to whether $\text{rank}(\bar{g}) = m$ or $m - 1$.

Case 1. Suppose first that $\text{rank}(\bar{g}) = m$. So $\bar{g} = 1$ and $g_q \in E(\mathcal{T}_{n_q})$ for all $q \in [m]$. Let $q \in M_r \setminus \{a\}$, and suppose $q = p\bar{h}$ where $p \in M_r$. Then $(hg)_p = h_p g_{p\bar{h}} = h_p g_q$. But h_p and $(hg)_p$ are injective, by assumption. Since $p\bar{h} = q \in M_r$, it follows that $h_p \in S_r$, so in fact $g_q = h_p^{-1}(hg)_p$ is injective. Since also $g_q \in E(\mathcal{T}_r)$, as noted above, we conclude that $g_q = 1$. Since this is true for all $q \in M_r \setminus \{a\}$, and since $\bar{g} = 1$, it follows that $g|_Y = \text{id}_Y$, as desired.

Case 2. Suppose now that $\text{rank}(\bar{g}) = m - 1$. Since $g \in E$, we must have $\bar{g} = e_{bc}$ for some $b, c \in [m]$ with $b \neq c$. Since $\text{rank}(\bar{h}\bar{g}) = \text{rank}(\bar{h}) = m - 1$, we cannot have both $b, c \in \text{im}(\bar{h})$. We also have $g_q \in E(\mathcal{T}_{n_q})$ for all $q \in \text{im}(\bar{g}) = [m] \setminus \{c\}$. We now consider two subcases, according to whether a belongs to $\text{im}(\bar{h})$ or not.

Subcase 2.1. Suppose first that $a \notin \text{im}(\bar{h})$. Since b, c do not both belong to $\text{im}(\bar{h})$, we must have either $b = a$ or $c = a$. Suppose first that $c = a$. Note that $\bar{g}|_{M_r \setminus \{a\}} = \text{id}_{M_r \setminus \{a\}}$ and $g_q \in E(\mathcal{T}_r)$ for all $q \in M_r \setminus \{a\}$ so, as in Case 1, we conclude that $g_q = 1$ for all such q , and therefore, $g|_Y = \text{id}_Y$. Now suppose $b = a$. If $c \notin M_r$, then again $\bar{g}|_{M_r \setminus \{a\}} = \text{id}_{M_r \setminus \{a\}}$ and $g_q \in E(\mathcal{T}_r)$ for all $q \in M_r \setminus \{a\}$, and the proof concludes as above. So suppose $c \in M_r$.

In fact, we will show that this case is not possible. Since $\text{im}(\bar{h}) = [m] \setminus \{a\}$ and $M_r \bar{h} \subseteq M_r$, it follows that \bar{h} maps $[m] \setminus M_r$ bijectively into $[m] \setminus M_r$. But h_q is injective for all $q \in [m] \setminus M_r$, so it follows that $h|_{X \setminus X_r} \in \mathcal{T}_{X \setminus X_r}$ is injective, and hence bijective. We deduce that h_q is bijective for all $q \in [m] \setminus M_r$. Let $q \in [m] \setminus M_r$ and put $p = q\bar{h}^{-1}$. Since $(gh)_p = h_p g_q$ is injective, it follows that g_q is injective. But also $g_q \in E(\mathcal{T}_{n_q})$ for all such q , giving $g_q = 1$ for $q \in [m] \setminus M_r$. Since also $\bar{g}|_{[m] \setminus M_r} = \text{id}_{[m] \setminus M_r}$, we would have $g \in S_r$, a contradiction. This completes the proof in Subcase 2.1.

Subcase 2.2. Finally, suppose $a \in \text{im}(\bar{h})$. As before, if $c \notin M_r \setminus \{a\}$, then $g|_Y = \text{id}_Y$ quickly follows. So suppose $c \in M_r \setminus \{a\}$. Actually, we will show that this is impossible. In particular, $c \in \text{im}(\bar{h})$, so $b \notin \text{im}(\bar{h})$. Next we claim that $n_b < r$. Indeed, suppose this is not the case. Note first that $b \notin \text{im}(\bar{h})$ implies $n_b \neq r$, since $M_r \subseteq \text{im}(\bar{h})$. So we must have $n_b > r$. Then some $q \in [m]$ with $n_q > r$ is mapped by \bar{h} to some $p \in [m]$ with $n_p \leq r < n_q$. But then $|C_q h| \leq |C_p| < |C_q|$, contradicting the fact that h_q is injective. This completes the proof of the claim that $n_b < r$. But now let $q \in M_r$ be such that $q\bar{h} = c$. Then $|C_q h g| \leq |C_c g| \leq |C_b| = n_b < r = |C_q|$, contradicting the assumption that $(hg)_q$ is injective. This completes the proof. \square

Lemma 3.13. *Let U be an arbitrary idempotent generating set for S and let $r \in [n]$. Then $U \cap S_r$ is a generating set for S_r .*

Proof. If $\mu_r = 0$, then $S_r = \{1\} = \langle \emptyset \rangle$, and the result is trivially true. So suppose $\mu_r \geq 1$, and write $U_r = U \cap S_r$. By Lemma 3.11 and [3, Corollary 4.2], S_r is generated by all elements of the form

$$(a) \ e_{ij}^{(k)} \text{ for } i, j \in [r] \text{ with } i \neq j \text{ and all } k \in M_r, \text{ and} \quad (b) \ e_{ij;f} \text{ for } i, j \in M_r \text{ with } i \neq j \text{ and all } f \in S_r.$$

So it suffices to show that $\langle U_r \rangle$ contains each element of type (a) and (b). Now, for each $k \in M_r$, U contains a generating set V for $\mathcal{E}_{n_k}^{(k)} = \mathcal{E}_r^{(k)}$ by Lemma 3.4, and we clearly have $V \subseteq U_r$. In particular, $\langle U_r \rangle$ contains all elements of type (a). So now fix $f \in S_r$ and $i, j \in M_r$ with $i \neq j$. Consider an expression $e_{ij;f} = g_1 \cdots g_k$, where $g_1, \dots, g_k \in U \setminus \{1\}$. Let $L = \{q \in [k] : g_q \in U_r\}$, and write $L = \{q_1, \dots, q_l\}$ where $q_1 < \dots < q_l$. We first aim to show that $(g_1 \cdots g_k)|_{X_r} = (g_{q_1} \cdots g_{q_l})|_{X_r}$.

For $p \in [k]$, let $h_p = g_1 \cdots g_p$, and write $h_p = [h_{p_1}, \dots, h_{p_m}; \bar{h}_p]$. We claim that for all $p \in [k]$,

$$\begin{aligned} (i) \ h_{pq} \text{ is injective for all } q \in [m], & \quad (iii) \ M_r \bar{h}_p \subseteq M_r, \\ (ii) \ \text{rank}(\bar{h}_p) = m - 1, & \quad (iv) \ |M_r \bar{h}_p| = \mu_r - 1. \end{aligned}$$

By Lemma 3.6, $h_1 = g_1 = e_{ij;f}$ or $e_{ji;f^{-1}}$, so the claim is true for $p = 1$. Now suppose (i–iv) all hold for some $1 \leq p < k$. Since $e_{ij;f} = h_{p+1} g_{p+2} \cdots g_k$, it is clear that each $h_{p+1,q}$ is injective, so (i) holds. Next, note that $m - 1 = \text{rank}(\bar{g}_1) \leq \text{rank}(\bar{h}_{p+1}) \leq \text{rank}(e_{ij}) = m - 1$, giving (ii). Next, we have $|M_r \bar{h}_{p+1}| \leq |M_r \bar{g}_1| = \mu_r - 1$. But $\text{rank}(\bar{h}_{p+1}) = m - 1$, so $|A \bar{h}_{p+1}| \geq |A| - 1$ for any subset $A \subseteq [m]$, and (iv) follows. For (iii), note that the induction hypothesis gives $M_r \bar{h}_p \subseteq M_r$. If $g_{p+1} \in S_r$, then $M_r \bar{g}_{p+1} \subseteq M_r$ by definition, so $M_r \bar{h}_{p+1} = M_r \bar{h}_p \bar{g}_{p+1} \subseteq M_r$. If $g_{p+1} \notin S_r$, then all the conditions of Lemma 3.12 are satisfied (with $h = h_p$ and $g = g_{p+1}$). We conclude then that $g_{p+1}|_Y = \text{id}_Y$, where $Y = X_r h_p$. In particular, $\bar{g}_{p+1}|_{M_r \bar{h}_p} = \text{id}_{M_r \bar{h}_p}$. But then $M_r \bar{h}_{p+1} = M_r \bar{h}_p \bar{g}_{p+1} = M_r \bar{h}_p \subseteq M_r$. This completes the proof of the inductive step and, hence, of the claim.

Now suppose $p \in [k]$ is such that $g_p \notin S_r$. In particular, $p \geq 2$ (since $g_1 \in S_r$, as noted above). By the above claim (and as noted in its proof), the conditions of Lemma 3.12 are satisfied (for $h = h_{p-1}$ and $g = g_p$), so we conclude that $h_p|_{X_r} = h_{p-1}|_{X_r}$. So, if $Q = [k] \setminus L = \{p \in [k] : g_p \notin U_r\}$, and $Q = \{p_1, \dots, p_s\}$ where $s = k - l$ and $p_1 < \dots < p_s$, then

$$\begin{aligned} e_{ij;f}|_{X_r} &= (g_1 \cdots g_k)|_{X_r} = h_{p_s}|_{X_r} (g_{p_s+1} \cdots g_k) \\ &= h_{p_s-1}|_{X_r} (g_{p_s+1} \cdots g_k) \\ &= h_{p_s-1}|_{X_r} (g_{p_{s-1}+1} \cdots g_{p_s-1}) (g_{p_s+1} \cdots g_k) \\ &= h_{p_s-1-1}|_{X_r} (g_{p_{s-1}+1} \cdots g_{p_s-1}) (g_{p_s+1} \cdots g_k) \\ &\quad \vdots \\ &= (g_1 \cdots g_{p_1-1})|_{X_r} (g_{p_1+1} \cdots g_{p_2-1}) \cdots (g_{p_s+1} \cdots g_k) \\ &= ((g_1 \cdots g_{p_1-1}) (g_{p_1+1} \cdots g_{p_2-1}) \cdots (g_{p_s+1} \cdots g_k))|_{X_r} \\ &= (g_{q_1} \cdots g_{q_l})|_{X_r}. \end{aligned}$$

But also $e_{ij;f}|_{X \setminus X_r} = \text{id}_{X \setminus X_r} = (g_{q_1} \cdots g_{q_l})|_{X \setminus X_r}$, so it follows that $e_{ij;f} = g_{q_1} \cdots g_{q_l} \in \langle U_r \rangle$, completing the proof. \square

Corollary 3.14. *If $\mu_1 = 2$, then $\text{idrank}(S) \geq \rho + 1$, where ρ is defined in Corollary 3.9.*

Proof. Let G be an arbitrary idempotent generating set for S . By Lemma 3.13, G contains a generating set U_i for $S_i \cong \mathcal{E}(X_i, \mathcal{P}_i)$ for each $i \in [n]$, and we have $|U_i| \geq \text{rank}(\mathcal{E}(X_i, \mathcal{P}_i)) = \rho_{\mu_i, i}$. By Theorem 2.3, $\rho_{\mu_i, i} = \mu_i \rho_i + i! \binom{\mu_i}{2}$, unless $i = 1$, in which case $\rho_{\mu_i, i} = \rho_{21} = 2 = 1 + (\mu_1 \rho_1 + 1! \binom{\mu_1}{2})$. So G contains at least

$$\sum_{i=1}^n \rho_{\mu_i, i} = 1 + \sum_{i=1}^n \left(\mu_i \rho_i + i! \binom{\mu_i}{2} \right)$$

elements from these generating sets of S_1, \dots, S_n . By the last two paragraphs of the proof of Corollary 3.9, G contains an additional

$$\sum_{i=1}^n \mu_i \nu_i + \sum_{1 \leq i < j \leq n} \mu_i \mu_j \frac{j!}{(j-i)!}$$

elements. Adding the two expressions above, we see that $|G| \geq \rho + 1$, and the proof is complete. \square

For the proof of the next result, we use the standard notation $f = \begin{pmatrix} A_1 & \dots & A_k \\ a_1 & \dots & a_k \end{pmatrix}$ to indicate that f is the function with domain $A_1 \cup \dots \cup A_k$ that maps each of the points in A_q to a_q for each $q \in [k]$.

Proposition 3.15. *For each $q \in [n]$, let U_q be an idempotent generating set for S_q . For each $i \in [m]$, choose sets $J_i \subseteq [m]$ such that $|J_i| = \nu_{n_i}$ and $\{n_j : j \in J_i\} = \{n_q : q \in [m], n_q < n_i\}$. For each $i \in [m]$ and $j \in J_i$, choose some $f_{ij} \in \text{Surj}(n_i, n_j)$. Then $U = U_1 \cup \dots \cup U_n \cup W_1 \cup W_2$ is a generating set for S , where*

$$W_1 = \{e_{ij;f} : 1 \leq i < j \leq m, n_i > n_j, f \in \text{Inj}(n_j, n_i)\} \quad \text{and} \quad W_2 = \{e_{ji;f_{ij}} : i \in [m], j \in J_i\}.$$

Proof. By Proposition 3.3, it suffices to prove that $G_1 \cup G_2 \subseteq \langle U \rangle$. Since $S_q = \langle U_q \rangle$ for each $q \in [n]$, it follows that $\langle U \rangle$ contains

- (i) each $e_{ij}^{(k)}$ with $k \in [m]$, $i, j \in [n_k]$ and $i \neq j$, and
- (ii) each $e_{ij;f}$ with $i, j \in [m]$, $i \neq j$, $n_i = n_j$ and $f \in \mathcal{S}_{n_i}$.

Since $W_1 \subseteq U$, it remains to show that $\langle U \rangle$ contains

- (iii) each $e_{ji;f}$ with $i, j \in [m]$, $n_i > n_j$ and $f \in \text{Surj}(n_i, n_j)$.

Let i, j, f be as in (iii). Let $k \in J_i$ be such that $n_k = n_j$, and for simplicity, put $g = f_{ik}$. So $e_{ki;g} \in U$. Write $f = \begin{pmatrix} A_1 & \dots & A_{n_j} \\ 1 & \dots & n_j \end{pmatrix}$ and $g = \begin{pmatrix} B_1 & \dots & B_{n_j} \\ 1 & \dots & n_j \end{pmatrix}$. Also, choose b_1, \dots, b_{n_j} such that $b_q \in B_q$ for each q . Put $h = \begin{pmatrix} A_1 & \dots & A_{n_j} \\ b_1 & \dots & b_{n_j} \end{pmatrix} \in \mathcal{T}_{n_i}$. Since $n_j < n_i$, note that $h \in \mathcal{E}_{n_i}$. So $h^{(i)} \in \mathcal{E}_{n_i}^{(i)} \subseteq \langle U \rangle$. It is clear that $e_{ki;f} = h^{(i)} e_{ki;g} \in \langle U \rangle$. In particular, if $k = j$, then we have shown that $e_{ji;f} = e_{ki;f} \in \langle U \rangle$. So suppose $k \neq j$. Now choose a_1, \dots, a_{n_j} so that $a_q \in A_q$ for each q . Put $d = \begin{pmatrix} 1 & \dots & n_j \\ a_1 & \dots & a_{n_j} \end{pmatrix} \in \text{Inj}(n_j, n_i)$. Note that $e_{ij;d} \in W_1 \subseteq U$, and that $e_{jk;1} \in \langle U \rangle$ as shown above, since $n_j = n_k$. It is clear that $df = 1 \in \mathcal{S}_{n_j}$, and one may then easily check that $e_{ji;f} = (e_{ij;d} e_{jk;1} e_{ki;f})^2 \in \langle U \rangle$, completing the proof. \square

We are now ready to prove the two main results of the paper. Again, these are stated in terms of the parameters μ_i, ν_i, ρ_i introduced at the beginning of this section and in Theorem 2.1.

Theorem 3.16. *We have $\text{rank}(S) = \text{idrank}(S) = \rho$, where*

$$\rho = \sum_{i=1}^n \left(\mu_i \rho_i + i! \binom{\mu_i}{2} \right) + \sum_{1 \leq i < j \leq n} \mu_i \mu_j \frac{j!}{(j-i)!},$$

unless $\mu_1 = 2$ in which case $\text{rank}(S) = \rho$ and $\text{idrank}(S) = \rho + 1$.

Proof. Let $U = U_1 \cup \dots \cup U_n \cup W_1 \cup W_2$ be an idempotent generating set as described in Proposition 3.15, with U_1, \dots, U_n of minimal size. As in the proof of Corollary 3.9,

$$|W_1| = \sum_{1 \leq i < j \leq n} \mu_i \mu_j \frac{j!}{(j-i)!} \quad \text{and} \quad |W_2| = \sum_{r=1}^m \nu_{n_r} = \sum_{i=1}^n \mu_i \nu_i.$$

As in the proof of Corollary 3.14,

$$|U_1| + \dots + |U_n| = \sum_{i=1}^n \left(\mu_i \rho_i + i! \binom{\mu_i}{2} \right),$$

unless $\mu_1 = 2$, in which case we must add 1 to the right hand side of this last expression. Adding these values, we conclude that

$$|U| = \begin{cases} \rho & \text{if } \mu_1 \neq 2 \\ \rho + 1 & \text{if } \mu_1 = 2. \end{cases}$$

Combined with Corollaries 3.9 and 3.14, and noting that $U \subseteq E$, this shows that $\text{rank}(S) = \text{idrank}(S) = \rho$ if $\mu_1 \neq 2$, and also that $\text{idrank}(S) = \rho + 1$ if $\mu_1 = 2$. To complete the proof, it suffices to prove that $S = \langle V \rangle$ for some $V \subseteq S$ with $|S| = \rho$ if $\mu_1 = 2$. For the remainder of the proof, we assume that $\mu_1 = 2$.

We have already seen that $S = \langle U \rangle$ and $|U| = \rho + 1$. Also, since there is a unique generating set of size 2 for $S_1 \cong \mathcal{E}(X_1, \mathcal{P}_1) \cong \mathcal{E}_2$, namely $U_1 = \{f, g\}$ where $f = e_{m-1, m; 1}$ and $g = e_{m, m-1; 1}$, we see that U must contain both f and g . Let $h \in \text{Inj}(1, n_1) = \text{Inj}(n_m, n_1)$ be arbitrary, and put $e = e_{1m; h}$. So $e \in W_1 \subseteq U$. It is easy to check that $e = (eg)f$ and $g = f(eg)$. It follows that $\langle e, f, g \rangle = \langle eg, f \rangle$ and so $S = \langle V \rangle$, where $V = (U \setminus \{e, g\}) \cup \{eg\}$. Since $|V| = \rho$, this completes the proof. \square

For the statement of the next result, by ‘‘minimal idempotent generating set’’ we mean an idempotent generating set that has the smallest possible size.

Theorem 3.17. (i) *Every idempotent generating set of S contains a minimal idempotent generating set.*

(ii) *Every minimal idempotent generating set of S is of the form described in Proposition 3.15 and with each U_q of minimal size.*

(iii) *The number of minimal idempotent generating sets of S is equal to*

$$\prod_{i=1}^n \sigma_{\mu_i, i} \times \prod_{\substack{1 \leq i < j \leq n \\ \mu_i \neq 0 \neq \mu_j}} \mu_i \mu_j S(j, i) i!,$$

where $S(j, i)$ is a Stirling number (of the second kind) and the numbers $\sigma_{\mu_i, i}$ are defined in Theorem 2.4.

Proof. Let U be an arbitrary idempotent generating set for S . By Lemma 3.13, U contains an idempotent generating set U_r of S_r for each $r \in [n]$. By Theorem 2.4, each U_r contains an idempotent generating set V_r of minimal size. As in the proof of Corollary 3.9, U must contain the sets

$$W_1 = \{e_{ij; f} : 1 \leq i < j \leq m, n_i > n_j, f \in \text{Inj}(n_j, n_i)\} \quad \text{and} \quad W_2 = \{e_{ji; f_{ij}} : i \in [m], j \in J_i\},$$

for some choice of sets J_i and functions $f_{ij} \in \text{Surj}(n_i, n_j)$. The set $V_1 \cup \dots \cup V_n \cup W_1 \cup W_2 \subseteq U$ has size $\text{idrank}(S)$, as stated in Theorem 3.16, and is a generating set for S by Proposition 3.15. This completes the proof of (i).

If U is an arbitrary idempotent generating set of minimal possible size, then we must in fact have $U = V_1 \cup \dots \cup V_n \cup W_1 \cup W_2$ (in the above notation), proving (ii). For each $i \in [n]$, we may choose V_i in $\sigma_{\mu_i, i}$ ways. To specify W_2 , for each $k \in M_j$ where $j \in [n]$ is such that $\mu_j \neq 0$, and for each $i \in [n]$ with $i < j$ and $\mu_i \neq 0$, we must choose some $e_{kl; f}$ where $l \in M_i$ and $f \in \text{Surj}(j, i)$. There are μ_j such k , μ_i such l , and $|\text{Surj}(j, i)| = S(j, i) i!$ such f . Multiplying these values as appropriate gives (iii). \square

References

- [1] Jo˜ao Ara˜ujo, Wolfram Bentz, J.D. Mitchell, and Csaba Schneider. The rank of the semigroup of transformations stabilising a partition of a finite set. *Preprint*, 2014, [arXiv:1404.1598](#).
- [2] Jo˜ao Ara˜ujo and Csaba Schneider. The rank of the endomorphism monoid of a uniform partition. *Semigroup Forum*, 78(3):498–510, 2009.
- [3] Igor Dolinka and James East. Idempotent generation in the endomorphism monoid of a uniform partition. *Comm. Algebra*, to appear, [arXiv:1407.3312](#).
- [4] Gracinda Gomes and John M. Howie. On the ranks of certain finite semigroups of transformations. *Math. Proc. Cambridge Philos. Soc.*, 101(3):395–403, 1987.
- [5] J. M. Howie. The subsemigroup generated by the idempotents of a full transformation semigroup. *J. London Math. Soc.*, 41:707–716, 1966.
- [6] J. M. Howie. Idempotent generators in finite full transformation semigroups. *Proc. Roy. Soc. Edinburgh Sect. A*, 81(3-4):317–323, 1978.
- [7] E. M. Wright. The number of irreducible tournaments. *Glasgow Math. J.*, 11:97–101, 1970.