

Identification and Mode of Action of a Plant Natural Product Targeting Human Fungal Pathogens

DORSAZ, Stéphane, *et al.*

Abstract

Candida albicans is a major cause of fungal diseases in humans and its resistance to available drugs is of concern. In an attempt to identify novel antifungal agents, we initiated a small scale screening of a 199 natural plant compounds (NPs) library. In vitro susceptibility profiling experiments identified 33 NPs with activity against *C. albicans* ($MIC_{50} \leq 32 \mu\text{g/ml}$). Among the selected NPs, the sterol alkaloid tomatidine was further investigated. Tomatidine originates from *Solanum lycopersicum* (tomato) and exhibited high fungistatic activity against *Candida* species ($MIC_{50} \leq 1 \mu\text{g/ml}$) but no cytotoxicity against mammalian cells. Genome-wide transcriptional analysis of *C. albicans* tomatidine-treated cells revealed a major alteration (upregulation) of ergosterol genes suggesting that the ergosterol pathway was targeted by this NP. Consistent with this transcriptional response, sterol content analysis of tomatidine-treated cells showed not only inhibition of Erg6 (C-24 sterol methyltransferase) but also of Erg4 (C-24 sterol reductase) activity. A forward genetic approach in *Saccharomyces cerevisiae* coupled with whole [...]

Reference

DORSAZ, Stéphane, *et al.* Identification and Mode of Action of a Plant Natural Product Targeting Human Fungal Pathogens. *Antimicrobial Agents and Chemotherapy*, 2017, vol. 61, no. 9, p. e00829-17

DOI : 10.1128/AAC.00829-17

PMID : 28674054

Available at:

<http://archive-ouverte.unige.ch/unige:103514>

Disclaimer: layout of this document may differ from the published version.



UNIVERSITÉ
DE GENÈVE

1 **Identification and mode of action of a plant natural product targeting human**
2 **fungal pathogens**

3 Stéphane Dorsaz,[§] Tiia Snäka,[§] Quentin Favre-Godal,[†] Pierre Maudens,[†] Nathalie Boulens,[†]
4 Pascal Furrer,[†] Samad Nejad Ebrahimi,[‡] Matthias Hamburger,[‡] Eric Allémann,[†] Katia Gindro,[°]
5 Emerson Ferreira Queiroz,[†] Howard Riezman,[⊥] Jean-Luc Wolfender,[†] Dominique Sanglard ^{*,§}

6

7 [§] Institute of Microbiology, University of Lausanne and University Hospital Center, Rue du
8 Bugnon 48, CH-1011 Lausanne, Switzerland.

9 [†] School of Pharmaceutical Sciences, EPGL, University of Geneva - University of Lausanne, Rue
10 Michel Servet 1, 1211 Geneva 4, Switzerland

11 [⊥] Department of Biochemistry, NCCR Chemical Biology, University of Geneva, 30 Quai Ernest-
12 Ansermet, 1211 Genève 4, Switzerland.

13 [°]Agroscope, Strategic Research Division Plant Protection, Mycology and Biotechnology, Route
14 de Duiller 50, P.O. Box 1012, 1260 Nyon, Switzerland

15 [‡]Division of Pharmaceutical Biology, Department of Pharmaceutical Sciences, University of
16 Basel, Klingelbergstrasse 50, CH-4056 Basel, Switzerland

17

18 **Running Title:** Natural products as antifungal agents

19 **Keywords:** Antifungal agents, natural products, sterol biosynthesis, antifungal therapy

20

21

22

23 *: corresponding author:
24 Institute of Microbiology
25 University of Lausanne and University Hospital Center
26 CH-1011 Lausanne, Switzerland
27 Tel: +41 21 3144083
28 e-mail: Dominique.Sanglard@chuv.ch

29

30 **Abstract**

31 *Candida albicans* is a major cause of fungal diseases in humans and its resistance to available
32 drugs is of concern. In an attempt to identify novel antifungal agents, we initiated a small
33 scale screening of a 199 natural plant compounds (NPs) library. *In vitro* susceptibility profiling
34 experiments identified 33 NPs with activity against *C. albicans* ($MIC_{50} \leq 32 \mu\text{g/ml}$). Among the
35 selected NPs, the sterol alkaloid tomatidine was further investigated. Tomatidine originates
36 from *Solanum lycopersicum* (tomato) and exhibited high fungistatic activity against *Candida*
37 species ($MIC_{50} \leq 1 \mu\text{g/ml}$) but no cytotoxicity against mammalian cells. Genome-wide
38 transcriptional analysis of *C. albicans* tomatidine-treated cells revealed a major alteration
39 (upregulation) of ergosterol genes suggesting that the ergosterol pathway was targeted by
40 this NP. Consistent with this transcriptional response, sterol content analysis of tomatidine-
41 treated cells showed not only inhibition of Erg6 (C-24 sterol methyltransferase) but also of
42 Erg4 (C-24 sterol reductase) activity. A forward genetic approach in *Saccharomyces cerevisiae*
43 coupled with whole genome sequencing identified 2 non-synonymous mutations in *ERG6*
44 (amino acids: D249G and G132D) responsible for tomatidine resistance. Our results therefore
45 identified unambiguously Erg6, a sterol C-24 methyltransferase absent in mammals, as the
46 main direct target of tomatidine. We tested the *in vivo* efficacy of tomatidine in a mouse
47 model of *C. albicans* systemic infection. Treatment with a nano-crystal pharmacological
48 formulation successfully decreased the fungal burden in infected kidneys as compared to
49 placebo and thus confirmed the potential of tomatidine as a therapeutic agent.

50 **Introduction**

51 Invasive fungal infections are an increasing threat to human health. In the developed
52 countries, these infections predominantly occur in the context of increasingly aggressive
53 immunosuppressive therapies. The overall mortality for invasive diseases caused by *Candida*
54 and *Aspergillus* spp. is 30–50%, despite the advent of new diagnostic and therapeutic
55 strategies (1). The fight against *C. albicans* infections necessitates the use of antifungal agents
56 and continued efforts are required to improve the therapeutic outcomes associated with
57 fungal infections.

58 Antifungal drugs that are currently available in the treatment of *Candida* infections belong to
59 four different chemical classes and include polyenes, azoles, pyrimidine analogues and
60 echinocandins (2). While polyenes and azoles target sterols and their biosynthesis, pyrimidine
61 analogues perturb nucleic acid biosynthesis and echinocandins interfere with cell wall
62 biosynthesis. The activity against common fungal pathogens and their detailed mode of action
63 is summarised in several available reviews (3, 4). The repeated or long-term use of antifungal
64 agents in medicine has facilitated the development of resistance in clinically relevant species
65 (5). When occurring, antifungal resistance can be a serious clinical problem due to the limited
66 number of available agents. In general, the incidence of antifungal resistance among human
67 fungal pathogens is low to moderate, especially when compared to the incidence of antibiotic
68 resistance among bacterial pathogens. Antifungal resistance occurrence has to be considered
69 independently for each antifungal class and for each fungal species. Moreover,
70 epidemiological data regarding incidence of resistance among fungal species is not identically
71 distributed worldwide (6, 7). Taken together, the small number of available antifungal agents
72 and the occurrence of resistance reveal the urgent need for novel active compounds.

73 Natural products (NPs) have already provided a vast resource for active ingredients in
74 medicines. The reason for this success can be explained by the high chemical diversity of NPs
75 and the effects of evolutionary pressure to create biologically active molecules and/or the
76 structural similarity of protein targets across many species (8). In the field of antimicrobials,
77 NPs met important successes. Starting with the discovery of penicillin, the pharmaceutical
78 industry has relied on this source extensively for antibiotic development. Nowadays, 80% of

79 all available clinically used antibiotics are directly (or indirectly) derived from NPs (9). Some
80 antifungals including polyenes and echinocandins derive directly from NPs.

81 The discovery of structurally novel NPs with suitable pharmacological properties as antibiotic
82 leads has weakly progressed in the recent decades (10). Innovative strategies provided
83 comprehensive antifungal profilings of given NPs and an understanding of their mode of
84 action for target identification and validation (11).

85 In a precedent study, we reported a strategy to identify antifungal NPs from plant crude
86 extracts (12). This strategy relied on the use of a *C. albicans* isolate highly susceptible to
87 growth inhibitors and in which traces of inhibitory NPs could be detected. NPs were identified
88 with a bioassay enabling a rapid detection tool of antifungal activity. With the determination
89 of chemical structures of the identified NPs, novel compounds could be readily processed for
90 further evaluation with *in vivo* approaches (13).

91 In this study, we report a small scale screening of selected NPs with an in-depth
92 characterization of their biological properties. The compounds were tested on the basis of
93 activity against different pathogenic and non-pathogenic yeasts and of their toxicity for
94 mammalian cells. One of promising compounds (tomatidine) showing high activity against *C.*
95 *albicans* was further investigated. Tomatidine mode of action was in-depth characterized for
96 the first time and its activity was confirmed *in vivo*.

97

98 Results

99 Screening of small scale library of plant NPs for antifungal activities

100 A library of 199 natural products (NPs) with potential antifungal activity was built. Compounds
101 were selected either according to previously reported activities or by structural analogy to
102 scaffolds that were known to be active. These compounds were obtained in two ways, either
103 by targeted isolation from plant extracts (29 different plants were investigated making 53%
104 of the investigated NPs) or by commercial acquisition after selection based on structural
105 similarity with documented antifungals (see Material and Methods; compounds are listed in
106 Supplementary Table S1). The 199 NPs were subjected to standard *in vitro* microdilution
107 susceptibility assays (EUCAST method) with *C. albicans* under acidic and neutral conditions
108 (pH of 4.6 and 7.0). These different values were chosen to reflect pH changes in the different
109 host niches of *C. albicans*. Results are summarized in Table 1. Considering a threshold for
110 antifungal activity of 32 µg/ml, our analysis identified 33 NPs exhibiting antifungal activities.
111 The activity threshold (32 µg/ml) was selected since we estimated that setting a high
112 threshold for a MIC value obtained *in vitro* would be problematic when testing activities *in*
113 *vivo* and to reach therapeutic concentration ranges in animals. With this threshold, while 2
114 compounds were active at neutral pH and 18 at low pH only, 13 were active at both pH
115 conditions (Table 1).

116 In order to further characterize their antifungal properties, active NPs were profiled for their
117 activity spectrum against several other clinically relevant *Candida* strains (*C. glabrata*, *C.*
118 *tropicalis*, *C. parapsilosis* and *C. krusei*) as well as another related non-pathogenic yeast
119 (*Saccharomyces cerevisiae*). As shown in Fig. 1, a variety of activity profiles was observed with
120 compounds active on all strains, while others only on a small species subset, suggesting a
121 diversity in their mode of action and in the target cell response. *C. glabrata* strains (azole
122 sensitive (AS) and azole resistant (AR)) exhibited the most resistant phenotypes. To identify
123 compounds with high antifungal potential, two major clinical antifungal agents (fluconazole
124 and caspofungin) were added to the selected NPs and all clustered according to their activity
125 profile. Four NPs (pyridoxatin, glc-3-medicagenic acid, medicagenic acid, plumbagin) were
126 grouped together with caspofungin (Fig. 1, purple highlighted) and exhibited a strong
127 inhibitory activity against all species tested. The nearest neighbouring cluster was containing

128 fluconazole and included three other NPs (formosanin C, tomatidine and taxodion; Fig. 1, red
129 highlighted). This cluster was characterized by strong overall activities but with reduced
130 susceptibility of specific yeast such as *C. glabrata* (AR) and *C. krusei* for fluconazole and *C.*
131 *glabrata* strains for tomatidine.

132 Biofilms are defined as complex cell populations with intrinsic resistance to many antifungal
133 drugs as compared to planktonic cells (14). We tested the selected NPs on *C. albicans* mature
134 biofilms formed *in vitro* using an activity cut-off value higher than the one used for planktonic
135 cells (MIC \leq 50 $\mu\text{g/ml}$). Most of the compounds (25 out of 33) were active against biofilms and
136 exhibited a MIC against biofilm higher than the MICs measured with planktonic cells, except
137 for NPs originating from *Waltheria indica*. Anti-biofilm activity of these NPs was previously
138 reported (15).

139 We next tested the activity of the selected NPs (33) on a Gram-negative bacterium (*E. coli*).
140 All but one compound (pyridoxatin, Table 1) were inactive against this bacterial species (MIC
141 >64 $\mu\text{g/ml}$), arguing for a fungal-specific inhibitory activity.

142 The last step in the screening process was to determine the potential acute toxicity of selected
143 NPs on mammalian cells. The 24 most active antifungal NPs (MIC \leq 16 $\mu\text{g/ml}$) were therefore
144 tested with standard cytotoxicity assays on Hela cells and the lethal dose (LD₅₀) was
145 determined. Our results indicated that only 3 NPs including tomatidine, medicagenic acid and
146 medicagenic acid 3-O-glucopyranoside, showed a selectivity index (SI) with an acceptable
147 range (>100). All the antifungal profiling assays are summarized in Table 1. Since the
148 antifungal activity of the two medicagenic acid-based NPs were already reported (16), we
149 focused our efforts in understanding the antifungal properties and mode of action of
150 tomatidine.

151

152 **Antifungal activity of tomatidine**

153 Tomatidine (Fig. 2A) is a sterol alkaloid produced by tomato (*Solanum lycopersicum*). It is the
154 precursor of the sterol glycol-alkaloids α -tomatine, which is a well-known antifungal saponin
155 against phytopathogens (17). Antifungal and antiparasitic activities of tomatidine have been
156 reported against *S. cerevisiae* and some parasites such as *Leishmania amazonensis* and

157 *Phytomonas serpens* (18-20). These studies revealed that tomatidine exposure induces a
158 perturbation of ergosterol biosynthesis and suggested the 24-sterol methyl transferase (24-
159 SMT-Erg6) as potential target within the sterol biosynthetic pathway. Erg6 is responsible for
160 the key structural difference between cholesterol and ergosterol (20, 21). This is consistent
161 with the fact that *ERG6* gene is present in fungi but absent in higher eukaryotes such as
162 mammals. Erg6 therefore represents an attractive target for the development of antifungal
163 agents.

164 Tomatidine susceptibility assays on several clinically-relevant *Candida* species are shown on
165 Fig. 2B. All strains, except *C. glabrata*, were highly susceptible to tomatidine. Similar results
166 were obtained with standard microdilution susceptibility assays with *C. krusei* and *C. albicans*
167 showing the highest sensitivity (Table 2, MIC = 0.625 μ M). We used a larger panel of *C.*
168 *albicans* isolates (n=9) and found that the tomatidine MIC50 (concentration needed to inhibit
169 at least 50% of the tested population) was at 0.3125 μ M. MIC50 values for *C. krusei* (n=8) and
170 *C. tropicalis* (n=9) were both at 1.25 μ M (Table S4). Tomatidine was not active against *C.*
171 *albicans* mature biofilms (Table 1), but the ability of *C. albicans* to form hyphae in
172 filamentation-inducing *in vitro* conditions was severely compromised in presence of
173 tomatidine. As shown in Fig. 2C, the addition of tomatidine to *C. albicans* cultures under
174 hyphal inducing conditions resulted in the formation of pseudohyphae and/or unseparated
175 budding yeast cells. Tomatidine exhibited fungistatic activity in *C. albicans* as revealed by
176 time-kill assays at different drug concentrations (Fig. 2D). Only weak fungicidal activity of
177 tomatidine (10 μ M) was observed in *C. krusei* with CFU counts decreased by about 80% as
178 compared to the starting inoculum after 24 h of incubation (Fig. 2D).

179

180 **Global transcriptional analysis of tomatidine-treated *C. albicans* cells**

181 Genome-wide transcriptional analysis in the presence of given drugs has been often used as
182 a mean to highlight their modes of action and to propose possible cellular targets (22). The
183 effect of tomatidine treatments on *C. albicans* transcriptome was investigated *in vitro* using
184 two different exposure times (1- and 3 h) at a fixed drug concentration (2.5 μ M). This drug
185 concentration was corresponding to the MIC obtained in YEPD (see Table 2) and led to a 30
186 % and 50 % growth inhibition after 1 h and 3 h incubation, respectively (data not shown).

187 Total RNAs was recovered from treated and untreated cells and genome-wide transcriptional
188 analysis was performed using RNA seq (see Materials and Methods). After 1- and 3 h of
189 tomatidine exposure, 129 and 1149 genes (2% and 21 % of the genes expressed) were
190 identified as differentially expressed between treated and untreated cells (false discovery
191 rate (FDR) ≤ 0.05 and fold-change ≥ 2), respectively. Hundred twelve (112) genes were found
192 to be affected at both 1- and 3 h exposure time. Seventy seven (77) genes were upregulated,
193 32 genes downregulated and 3 genes were inversely regulated as compared to untreated
194 conditions (Fig. 3A, supplementary File S1). GO term analysis on the 77 commonly
195 upregulated genes revealed a clear alteration of the ergosterol biosynthetic pathway
196 (corrected p-value < 0.0004) which included 6 genes (*ERG2*, *ERG3*, *ERG4*, *ERG6*, *ERG11*,
197 *ERG25*) (Fig. 3B). Interestingly, other components of the ergosterol pathway were found to
198 be upregulated by tomatidine at 1 h exposure time only including *ERG1*, *ERG5*, *ERG24* and
199 *UPC2* (the master sterol transcriptional regulator) and at 3 h exposure time only including
200 *ERG9* and *ERG28* (supplementary File S1). Interestingly, among all the *ERG* genes differentially
201 regulated by tomatidine treatment, the most affected was *ERG6* with 7.4- and 11.8 fold-
202 change of expression after 1- and 3 h treatment, respectively. This important fold-change was
203 confirmed by qPCR analysis (up to 40-fold increase at 3 h treatment) (Fig. 3C).

204 Response to steroid hormone stimulus was another overrepresented GO term among
205 upregulated genes (corrected p-value < 0.002). The three corresponding genes were part of
206 the *TAC1* regulon and included the major drug efflux system in *C. albicans* with the two ABC
207 transporters (*CDR1* and *CDR2*) and the transcriptional regulator itself (*TAC1*) (Fig. 3B). qPCR
208 analysis on *ERG4*, *ERG6*, *ERG11*, *CDR1* and *CDR2* expression on the same RNA samples
209 validated the RNA-seq data analysis (Fig. 3C, supplementary Fig. S1).

210 GO analysis identified only one enriched GO term (ion transport; corrected p-value < 0.003)
211 in commonly down-regulated gene (32 genes) with 2 genes involved in iron or calcium
212 transport (*FET34*, *FTR1*).

213 Exposing *C. albicans* cells to tomatidine for 3 h leads to large transcriptional changes (total of
214 1149 genes) with 1037 genes specifically affected after that exposure time. GO term analysis
215 identified processes that are characteristics of growth inhibition/arrest or drug-related stress

216 such as glucose transport (in upregulated genes) and the transcriptional and translational
217 machinery in downregulated genes (see supplementary File S1).

218 In order to better characterize the response to tomatidine, we used a Gene Set Enrichment
219 Analysis (GSEA) to identify transcriptional signatures from other drug-induced transcriptional
220 studies overlapping with the present tomatidine transcriptional data (Supplementary File S2).
221 As shown in Fig. 3D, tomatidine up- and downregulated genes (red and blue, respectively)
222 shared the highest number of regulated genes with the fluconazole_UP/DOWN gene sets,
223 followed by another azole-related gene set (ketoconazole_UP/DOWN) and gene set from cells
224 lacking the master regulator of ergosterol genes *UPC2*. Using another published set of
225 transcriptional data that includes several *in vitro* growth stress conditions (such as hypoxia,
226 (23), tomatidine gene set was sharing regulated genes with cells under hypoxic conditions,
227 which adaptive response mainly is mediated by *UPC2* (supplementary Fig. S2) (24). Taken
228 together, these analyses revealed that tomatidine had a transcriptional signature closely
229 related to azole drugs and strongly suggests that the ergosterol biosynthetic pathway is also
230 targeted by this NP.

231

232 **Tomatidine as an inhibitor of the sterol biosynthetic pathway**

233 As suggested by our transcriptional analysis, tomatidine is likely to interfere with ergosterol
234 biosynthesis given that the upregulation of *ERG* genes upon treatment has been shown to be
235 a cell response mechanism to drugs targeting the ergosterol pathway (25). In order to detect
236 more precisely the affected step(s) of this pathway, we performed an in-depth sterol content
237 analysis of *C. albicans*, *C. krusei* and *S. cerevisiae* cells treated with tomatidine. Total sterols
238 were extracted from yeast cells and next subjected to gas liquid chromatography coupled
239 with mass spectrometry (GC-MS) analysis.

240 As shown in Fig. 4A, untreated *C. albicans* cells contained, as expected, ergosterol (blue color)
241 as the most abundant sterol (>96% of total sterol). Zymosterol (the product known to
242 accumulate in the mutant for *ERG6*, a 24-C methyl transferase) (red color) was the major
243 sterol (84% of total sterols) in tomatidine-treated cells, thus suggesting a strong inhibition of
244 this enzymatic step by tomatidine. Furthermore, careful analysis of sterol content allowed the

245 identification of small amount of two other sterol intermediates including cholesta-trienol (6
246 %) and ergosta-5-7-24-(28)-tetraenol (8%). Enzymes involved in the late stage of ergosterol
247 biosynthesis are usually acting independently of each other which enables the pathway to
248 proceed even if one step is inhibited by a drug or is genetically suppressed leading to the
249 synthesis of alternative sterols. One of those is cholesta-trienol (green color) which was
250 detected in tomatidine-inhibited cells, where successive actions of Erg2, Erg3 and Erg5
251 occurred in absence of previous Erg6 activity. However, Erg4, the last enzyme involved in the
252 sterol pathway, requires first Erg6 action to produce its substrate. More specifically, the
253 methylene group added by Erg6 at position C-24 of zymosterol is the substrate for the C-24
254 reductase activity of Erg4. Therefore the detection of the Erg4 substrate (ergosta-5-7-24-(28)-
255 tetraenol (purple color)), was unanticipated and suggested that (i) tomatidine was also
256 inhibiting Erg4 and that (ii) Erg6 was not completely inhibited by tomatidine at the
257 concentration tested.

258 In order to be able to assess more accurately the inhibition of Erg4 activity, we next performed
259 a progressive decrease of Erg6 inhibition by titrating down tomatidine concentration. As
260 shown in Fig. 4A, decreasing tomatidine concentration from 2 μ M to 0.25 μ M resulted in the
261 release of Erg6 inhibition (zymosterol decreased from 84 % to 7 % of total sterol) and the
262 accumulation of ergosta-5-7-24-(28)-tetraenol (up to 72% of total sterol at 0.25 μ M
263 tomatidine), thus confirming the ability of tomatidine to inhibit Erg4. As the concentration of
264 tomatidine required to accumulate zymosterol and ergosta-5-7-24-(28)-tetraenol up to 84 %
265 and 72 % of total sterols is approximatively 2 μ M and 0.25 μ M, respectively, one can estimate
266 that tomatidine is about 10 times more effective as an inhibitor of Erg4 as compared to Erg6.
267 Interestingly, this Erg4/Erg6 dual inhibition was also found in *C. krusei* but not in *S. cerevisiae*
268 in which only Erg6 inhibition could be detected (supplementary Fig. S3).

269 Following these analysis, a *C. albicans* *ERG6* deletion strain was engineered in order to
270 validate that Erg6 is a target of tomatidine. We reasoned that, if the Erg6 target is absent,
271 then tomatidine will not be active against this mutant. As already described, the deletion of
272 *ERG6* results in increased susceptibility to a large variety of drugs, with the exception of azoles
273 and polyenes (26). The *erg6 Δ / Δ* mutant was indeed resistant to fluconazole with a 128-fold
274 higher MIC than the wild type (Fig. 4B). Interestingly, the *erg6 Δ / Δ* mutant constructed here
275 was as susceptible to tomatidine as the parental strain (Fig. 4B). This indicates a pleiotropic

276 susceptibility of the strain or potential secondary target(s) of tomatidine in addition to the
277 ergosterol pathway. Similar results were obtained using a *S. cerevisiae* *ERG6* mutant
278 (supplementary Fig. S4).

279

280

281

282 **Forward genetic approach to identify targets of tomatidine**

283 A forward genetic screen in *S. cerevisiae* was finally undertaken as an unbiased way to
284 discover the target(s) of tomatidine. The strain chosen in this approach lacked *PDR5* and
285 *MSH2* to avoid multidrug transporter-dependent resistance mechanisms and to increase the
286 rates at which resistance mutation may occur, respectively (27, 28). A *pdr5Δ msh2Δ* strain
287 (P1) was plated on YEPD media containing 10 μM tomatidine and, after a screening of more
288 than 2×10^8 cells, one resistant mutant (R1) was isolated. To obtain additional resistant
289 mutants, a second strategy inspired by Ojini *et al.* (28) was developed and consisted of
290 submitting cells to two drug-exposure periods in liquid media (72- and 48 h) interspersed by
291 a period of drug-free growth (48 h). Cultures with a robust growth at the end of the
292 experiment were further plated on solid YEPD media containing 10 μM tomatidine which
293 resulted in the isolation of four resistant mutants (R2, R3.1, R3.2 and R3.4) from two different
294 cultures. Tomatidine resistance of the five obtained mutants was confirmed by a serial
295 dilution assay (Fig. 5A). MIC results showed a 4-fold decrease in susceptibility for R1, R3.1 and
296 R3.2 as compared to the parental strain P1. Isolates R3.3 and R2 exhibited only a slight
297 decrease of susceptibility. Tomatidine resistance was specific since susceptibility assays with
298 different drug classes (fluconazole, caspofungin and amphotericin B) were in the range of
299 strain P1 (Table 3). Simplified sterol composition analysis using the unique
300 spectrophotometric absorbance signature of sterols confirmed that the mutants had normal
301 sterol composition with no detectable differences compared to the parental strain
302 (supplementary Fig. S5A).

303 The five mutant genomes were sequenced with that of parental strain P1. Alignment of
304 parental and mutant genomes to the reference genome (S288C) was followed by the

305 identification of non-synonymous polymorphisms uniquely present in the coding regions of
306 the resistant mutants (see Materials and Methods). All the non-synonymous mutations
307 inventoried in the five resistant mutants (between 26 to 60 according to the strain) are listed
308 in supplementary File S3. Interestingly, four out of the five resistant mutants contained a
309 missense mutation in *ERG6* with the following amino acid substitutions: D249G for R1, R3.1,
310 R3.2 and G132D for R2. These amino acid residues might play an important role in Erg6
311 function as they are conserved among fungi, plants or protozoa (29) (Fig. 5B), but seemed
312 dispensable for normal ergosterol synthesis. To determine if the two mutations were
313 sufficient to confer tomatidine resistance, the G132D and D249G substitutions were inserted
314 in a wild type *S. cerevisiae* strain (IMX585) using site-directed CRISPR-Cas9 genome editing
315 technology (30). Both strains (G132D and D249G) were 4-fold more resistant to tomatidine
316 than the wild type (Table 3), thus recapitulating (or increasing for G132D) the tomatidine
317 resistant phenotype of the original resistant mutants. Sterol profiles analysis showed that the
318 G132D strain, in contrast to the corresponding original resistant mutant and the D249G strain,
319 exhibited an altered sterol composition with intermediate profiles between an *ERG6* deletion
320 and a wild-type strain (supplementary Fig. S5B). This might explain the observed increase
321 fluconazole susceptibility of this strain (Table 3).

322 While the last resistant mutant (R3.3) retained a wild-type allele of *ERG6*, it exhibited a
323 frameshift mutation in *ACE2*. This gene encodes for a transcription factor required for septum
324 degradation after cytokinesis (31). Strain R3.3 exhibited a multicellular clumping phenotype,
325 which is identical to an *ACE2* deletion strain (supplementary Fig. S6). A decrease in tomatidine
326 susceptibility was observed in the *ACE2* deletion strain and thus recapitulated the resistance
327 phenotype of the R3.3 initial strain (Table 3).

328

329 **Tomatidine is targeted by efflux pumps**

330 The forward genetic approach was designed to exclude multidrug transporter-dependent
331 resistance mechanisms by using a *S. cerevisiae* strain lacking *PDR5*. However, the activation
332 of drug efflux in yeast is a common cell defence mechanism against toxic drugs, as observed
333 in our *C. albicans* genome-wide transcriptional analysis, in which the genes of the *TAC1*
334 regulon (including *CDR1*, *CDR2*) were among the tomatidine-upregulated genes (Fig. 3B). To

335 address whether tomatidine was a target of the efflux pump system in *C. albicans*, a set of
336 deletion mutants lacking each *CDR1*, *CDR2* or *MDR1* (another import efflux pump belonging
337 to the major facilitator superfamily of transporters) were used to evaluate their tomatidine
338 susceptibility. The *cdr1Δ/Δ* strain was the only one to exhibit increased susceptibility to
339 tomatidine in contrast to *CDR2* and *MDR1* mutants and the wild-type strain (Fig. 6A). This
340 data clearly indicated that, as many other drugs such as fluconazole (32), tomatidine is
341 targeted by the *CDR1* efflux pump.

342 A major problem arising during clinical treatment of candidiasis is the emergence of resistant
343 isolates. One important underlying mechanism consists of an upregulation of efflux pumps by
344 the acquisition of a hyperactive alleles of their regulator (33). To evaluate the potential cross-
345 resistance of hyperactive *TAC1* strains between fluconazole and tomatidine, we measured
346 the tomatidine susceptibility of two pairs of matched azole-susceptible (AS) and azole-
347 resistant (AR) *C. albicans* clinical isolates. The AR isolates carried *TAC1* hyperactive alleles.
348 While the difference in fluconazole MIC between DSY732 (AR) and DSY731 (AS) was of 64-fold
349 in magnitude, the tomatidine MIC between the two strains diverged only by 4-fold (Fig. 6B
350 left panel, Table 4). Similar results were obtained using *C. albicans* clinical isolates DSY1843
351 (AR) and DSY1841 (AS). Azole and tomatidine MICs increased by only 4- and 2-fold,
352 respectively (Fig. 6B right panel, Table 4). Taken together, these results suggest that the
353 common resistance mechanism against azole triggered by *TAC1* hyperactivity seems to have
354 limited effect on tomatidine susceptibility.

355 As *C. albicans* exhibited limited cross-resistance between fluconazole and tomatidine, we
356 next investigated the potential synergistic effect of their combination. Classical checkerboard
357 combination assays were performed and Fractional Inhibitory Concentration (FIC) values
358 determined (as described in Materials and Methods). As expected for drug targeting the same
359 pathway, fluconazole and tomatidine exhibited a strong additive effect in both *C. albicans* and
360 *C. krusei* with cell growth inhibition of more than 50% in the zone of additivity (Fig. 6C). We
361 then tested if the combination of the two drugs changed their fungistatic properties.
362 Interestingly, time-kill assay showed that fluconazole-tomatidine combination leads to a
363 fungicidal activity in *C. krusei* (>100-fold decrease in cells counts after 24 hours), but not in *C.*
364 *albicans* (Fig. 6D).

365

366 ***In vivo* activity of tomatidine**

367 To confirm the high potential of tomatidine as a therapeutic agent, the *in vivo* efficacy of the
368 compound was tested in a mouse model of *C. albicans* systemic infection. Due to its
369 hydrophobic nature, a nanoparticle-based formulation of tomatidine was developed to allow
370 its administration and to potentially enhance its bioavailability. Mice were infected through
371 the tail vein with *C. albicans* inoculum and were treated intraperitoneally (i.p.) with
372 tomatidine (50 mg/kg) or placebo at 6 hours, 24 hours and 31 hours post-infection (pi). Colony
373 forming units (CFU) were then determined in the kidneys 48h pi. As illustrated in Fig. 7, mice
374 treated with tomatidine exhibited statistically-significant reduced CFU compared to controls
375 (Mann-Whitney test, p-value = 0.031), thus highlighting the *in vivo* activity of tomatidine and
376 a therapeutic potential.

377

378 **Discussion**

379 **NPs as source of antifungal agents**

380 The aim of our study was to identify promising antifungal natural compounds starting with a
381 small scale screening of carefully selected compounds, proceeding with an extensive *in vitro*
382 characterisation of their antifungal properties and cytotoxicity, and then extending to the
383 identification of their cellular targets and the validation of their *in vivo* therapeutic potential.

384 A library of 199 natural products was built using both direct isolation from extracts of plants
385 with documented antifungal properties (29 different plants were investigated allowing to the
386 isolation of 53% of the investigated NPs) and commercial acquisition after selection based on
387 structural similarities with known antifungals. The efficiency of the preselection process
388 explains the high positive hit rate (17%; 33/199) of NPs with activity against human pathogens
389 using a cut-off MIC value (≤ 32 $\mu\text{g/ml}$). From extended bioactivity profiling procedure including
390 assays on different fungal strains, a bacterial strain and cytotoxicity assays for therapeutic
391 index evaluation, only three interesting leads were identified. Tomatidine stood out as being
392 a novel anti-*Candida* drug with a putative promising target.

393 Tomatidine is a sterol alkaloid from tomato plants with a cholesterol derived hydrophobic 27-
394 carbon skeleton and serves as precursor intermediate in the synthesis of a plant defence
395 metabolite, the glycol sterol alkaloid α -tomatine (34). The latter has been characterized as an
396 antifungal agent against a large variety of phytopathogens and possesses membrane
397 disruption properties (unspecific toxicity) caused by its ability to form complexes with
398 cholesterol and ergosterol (17, 35). Alpha-tomatine is present in most green parts of tomato
399 plants where it is subjected to degradation during maturation. In contrast, tomatidine, its
400 aglycone, has a different mode of action as it does not show any sterol binding activity (35).
401 It is poorly active on phytopathogens, it exhibits no toxicity *in vivo* and is present only in traces
402 in the tomato plant (17, 18, 36). An antifungal and antiparasitic activity of tomatidine has
403 been reported in *S. cerevisiae* and *Leishmania amazonensis* (18, 19) and a putative target, the
404 24-sterol methyl transferase (24-SMT-Erg6) was proposed. In the present study, the
405 antifungal activity of tomatidine was in-depth characterized for the first time against
406 important fungal pathogens. All *Candida* spp. except *C. glabrata* were susceptible to
407 tomatidine (MIC = 0.25 to 10 μM , Table 2).

408

409 **Tomatidine mode of action**

410 Multi-level investigations were then conducted in order to determine its mode of action and
411 to identify its molecular target. A first genomic strategy characterized tomatidine-specific
412 transcriptional signature in *C. albicans* and identified an important upregulation of the
413 ergosterol biosynthetic genes including *ERG6* as the most affected gene (Fig. 3). These results
414 confirmed studies published in *S. cerevisiae* (18) and pointed out the ergosterol pathway as
415 the target of tomatidine. Interestingly, the GSEA analysis identified fluconazole as mediating
416 the closest-related transcriptional signature to tomatidine (Fig. 3), which is consistent with
417 their inhibitory activity directed to the same pathway. Furthermore, this convergence of azole
418 and tomatidine activity was also showed by microscopic analysis of the cytological effect of
419 the drugs on the ultrastructure of *C. albicans* cells (supplementary Figure S7).

420 Considering the above findings, detailed sterol analysis using GC-MS was achieved in *C.*
421 *albicans* and *C. krusei* resulting in the indirect identification of (i) Erg6 inhibition through a
422 strong accumulation of the Erg6 substrate zymosterol in cells treated with high concentration
423 of tomatidine and (ii) Erg4 inhibition through a strong accumulation of ergosta-5-7-24-(28)-
424 tetraenol, its substrate, in cells treated with low concentration of tomatidine (Fig. 4). As Erg4
425 needs Erg6 activity to obtain its substrate, the identification of this dual Erg4-Erg6 inhibition
426 was made possible by the likely higher affinity of tomatidine for Erg4 as compared to Erg6.
427 Both enzymes are targeting the C-24 position in the ergosterol precursor and thereby
428 probably exhibit similar binding pocket conformation even if their protein sequence identity
429 is low (15 %). A dual effect on Erg4 and Erg6 had already been reported in *S. cerevisiae* for
430 azasterol, a sterol carrying a nitrogen in the side chain (37) and thus reinforces the idea of
431 structural similarities between the targets. Tomatidine dual inhibition was not detected in *S.*
432 *cerevisiae*, where no ergosta-5-7-24-(28)-tetraenol was identified even at low concentration.
433 This suggests that either Erg4 in *S. cerevisiae* is not inhibited by tomatidine or it has a lower
434 affinity for tomatidine than does Erg6. These intra-species differences could be attributed to
435 Erg4 independent genetic evolution as single point mutations in *ERG6* can affect sensitivity to
436 tomatidine.

437 It has been shown that deleting the target of fluconazole (*ERG11*) in *C. albicans* caused a
438 marked increase in its resistance to the drug (38). We applied the same paradigm to
439 tomatidine and tested the susceptibility of an *ERG6* deletion strain. Late-acting ergosterol
440 genes are not essential for cell viability and ergosterol is substituted by altered sterols in the
441 membrane of deletion strains (39). However, these substitutions impact the regulation of
442 membrane permeability and fluidity and are associated with diverse phenotypic alterations.
443 Yeast *ERG6* deletion mutant showed pleiotropic hypersensitivity to a broad range of
444 antifungal compounds and metabolic inhibitors reflecting an increased membrane
445 permeability and passive diffusion to small molecules (hydrophobic mostly) (26, 39, 40). The
446 fact that tomatidine susceptibility of *C. albicans* *ERG6* deletion strain was similar to the wild
447 type strain was intriguing. It is known that *ERG6* deletion can alter cell membrane
448 permeability to different drugs and this phenotype has been utilized by others in order to
449 improve the effect of different drugs (41). Given that *ERG6* deletion results in a slight
450 increased susceptibility to tomatidine as compared to wild type, it suggests that the
451 compound could still target other cellular components. Consistent with this hypothesis, we
452 identified *ACE2* as another putative target of tomatidine (see below). It is interesting to note
453 that the *C. albicans* *ERG6* deletion strain was resistant to fluconazole (Fig. 4B), which indicates
454 drug-dependent susceptibility phenotypes and thus excludes thereby an unspecific
455 hypersensitivity response to a given stressor.

456 Our forward genetic approach in *S. cerevisiae* was aimed to identify alternative targets of
457 tomatidine by avoiding the interference of efflux pump hyperactivity mechanisms. However,
458 our screen revealed Erg6 as major target and several explanations can be proposed. First, our
459 set of obtained resistant mutants was relatively small (5), thus limiting the discovery of
460 additional targets. Second, an alternative target could be accessible or present only in the
461 background of an *ERG6* deletion background, in which altered sterols are present with
462 compromised cell membrane functions. The hypothesis of a pleiotropic effect of the *ERG6*
463 deletion on the susceptibility to small hydrophobic molecules was tested using a sterol
464 alkaloid (solasodine) which possesses highly similar chemical structure to tomatidine (only
465 the planar configuration of the piperidine ring and an unsaturated alpha ring bond differs).
466 While this molecule was inactive against a *C. albicans* wild type strain (MIC >40 μ M), the
467 deletion of *ERG6* increased drug susceptibility (MIC = 20 μ M) and thus argued for an

468 unspecific drug hypersensitivity of this strain. Furthermore, *in vitro* filamentation studies with
469 *C. albicans* (Fig. 2D) attested of the similar pseudohyphae and/or unseparated budding yeast
470 cell phenotype between tomatidine-treated wild type cells and *ERG6* depleted cells arguing
471 for tomatidine has the major target. The forward genetic approach identified *ACE2* as a
472 possible tomatidine target. Interestingly, *ACE2* deletion in *C. albicans* results in upregulation
473 of several *ERG* genes of involved in sterol biosynthesis (42) and thus we reasoned that *ERG6*
474 could be upregulated in the resistant mutant R3.3, thus resulting in tomatidine resistance.
475 *ERG6* expression was however not significantly altered in R3.3 as compared to parent and
476 thus the basis of resistance by *ACE2* still remains unresolved.

477 In comparison to previous reports in *S. cerevisiae* that suggest Erg6 as a tomatidine target
478 (18), our forward genetic approach evidenced a direct interaction between Erg6 and
479 tomatidine, with a single substitution of well conserved amino acids (G132D or D249G)
480 sufficient to confer resistance. As a loss of function in Erg6 results in an increased
481 susceptibility to tomatidine, the two mutations did not affect drastically Erg6 function
482 (ergosterol was detected and amphotericin B susceptibility remained unaffected). In the
483 absence of X-ray-structure for sterol methyltransferase (Erg6), previous functional analysis
484 had identified functionally important residues in three regions using bioinformatics analysis,
485 mechanism-based inactivation and site-directed mutagenesis experiments (Fig. 5B) (29). The
486 two residues (G132D and D249G) were not included in these analyses and their effect on the
487 kinetic of C-methylation reaction remains to be determined. To address the mechanism of
488 Erg6 inhibition by tomatidine, studies of a known Erg6 inhibitor, 25-azalanosterol, have
489 suggested a non-competitive binding to a different site than the sterol binding site in the
490 active centre which leads to conformational changes deleterious to the catalytic reaction (29,
491 43). The two identified residues could mediate direct interactions to tomatidine and non-
492 conserved changes in these positions may result in a decreased affinity to the drug.

493 An important issue concerning the potential of tomatidine as an antifungal drug regard is its
494 resilience to efflux pump-mediated mechanisms, which are commonly acquired by azole
495 exposure. We showed here that tomatidine is the substrate of Cdr1 in *C. albicans*. By
496 extrapolation, we can hypothesize that the absence of activity against *C. glabrata*, which is
497 known to possess intrinsic resistance to antifungals, is related to its potent efflux machinery.
498 Nevertheless, we showed using matched azole-sensitive and azole-resistant clinical isolates

499 that the common resistance mechanism against azole through *TAC1* hyperactivity had limited
500 effect on tomatidine susceptibility. This reduced cross-resistance and the increased
501 fluconazole susceptibility of tomatidine-resistant strain (via G132D) raised the question about
502 the potential therapeutic advantage of the combined use of the two drugs. Our analysis
503 revealed an additive effect of their combination in a wild-type strain of *C. albicans*. In a recent
504 study (44), a screen for synergistic molecules in combination with fluconazole on a *C. albicans*
505 azole-resistant strain (with *ERG11* and *TAC1* mutations) identified a synergistic interaction
506 with tomatidine, thus confirming the high potential of tomatidine in drug combination
507 therapies.

508

509 ***In vivo* activity of tomatidine**

510 Our initial choice to screen a library of NPs was guided by the intrinsic properties of these
511 natural compounds as compared to synthetic products, namely their immense chemical
512 diversity, target specificity and intrinsic cell permeability. The critical step in developing new
513 antifungal agent is to validate the promising *in vitro* characteristics *in vivo* and, by choosing
514 NPs, we were hoping to bypass this bottle-neck. *In vivo* studies have validated that the target
515 of tomatidine (Erg6) is required for virulence in both mice and insect mini-host model of *C.*
516 *albicans* systemic infection ((45) and data not shown). Tomatidine is a highly hydrophobic
517 sterol-like molecule difficult to dissolve using common aqueous solvents. First attempts to
518 demonstrate *in vivo* activity of tomatidine in an animal model of *C. albicans* systemic infection
519 using (co)solvents (DMSO, ethanol) or cyclodextrin complexation (hydroxypropyl
520 betacyclodextrin) or surfactant (polyoxyl 35 castor oil) were not successful. We next reasoned
521 that other types of drug formulations could be more successful. Several studies have used
522 nanoparticles preparations to increase drug efficacy *in vivo*. For example, Moazeni et al (46)
523 have reverted *in vitro* azole resistance of *Candida* spp. with solid lipid nanoparticles prepared
524 with fluconazole. Inspired by this work, we prepared a nanoparticle-based tomatidine
525 formulation. This formulation, which did not affect the *in vitro* activity of tomatidine and did
526 not modify its chemical structure, was injected IP and decreased significantly fungal burden
527 in the kidney as compared to placebo. Further studies would be necessary to understand

528 tomatidine bioavailability and bio-distribution in mice, however our work clearly raised
529 tomatidine as a novel potential therapeutic antifungal agent.

530

531

532

533 **Acknowledgements**

534 The authors are thankful to the Swiss National Science Foundation for providing financial
535 support for this project, which aims to identify new antifungal compounds of natural origin
536 (Grant CR2313_143733 to J.L.W., E.F.Q., K.G. and D.S.). The authors are also thankful to
537 Françoise Ischer for excellent technical assistance, the Sanglard lab for helpful discussion and
538 Francine Voinesco for valuable technical assistance in transmission electron microscopy.

539

540 **Materials and Methods**

541 **Yeast strains culture and growth media**

542 Yeast and bacterial strains used in the study are listed in Table S1. NPs were dissolved in DMSO
543 to 10 µg/ml. Tomatidine (Tomatidine-HCL) was purchased from Phytolab (GmbH), dissolved
544 in DMSO at 2 mM concentration, heated for 20 min at 50°C for solubilization and stored at -
545 20°C upon usage.

546

547 **Antimicrobial susceptibility testing**

548 (i) Antifungal broth microdilution assays.

549 Antifungal susceptibility testing was carried out on the basis of EUCAST protocols with slight
550 modifications (47). Briefly, yeast strains were cultivated overnight at 30°C under constant
551 agitation in YEPD (1% yeast extract, 2% peptone, 2% glucose). Cultures were diluted to a
552 density of 2×10^5 cells per ml in RPMI (R8755- SIGMA) or Yeast Nitrogen Base (YNB) (MPbio)
553 buffered to adequate pH with HCL or NaOH and with a final concentration of 2% glucose.
554 Compounds from the NP library were dissolved in DMSO to 10 mg/ml as final concentration.
555 The final DMSO concentration was 1%. Two-fold serial dilutions were prepared from 32 to 1
556 µg/ml. Plates were incubated at 35°C for 24 h and then MICs were read with a
557 spectrophotometer plate reader set at 450 nm. The minimal inhibitory concentration (MIC)
558 was defined as the drug concentration at which the optical density was equal or decreased
559 more than 50% from that of the drug-free culture. For *S. cerevisiae* MICs, the media (YNB)
560 was complemented with Complete Supplement Mixture (CSM; Mpbio, Santa Ana, California,
561 USA) according to the supplier instructions. Assays were performed in triplicates.

562 For the detection of combinatory effect of tomatidine with fluconazole, a checkerboard assay
563 was set up in YNB pH 7 with 2-fold dilutions of fluconazole (0.008 to 0.5 µg/ml, 1 to 64 µg/ml)
564 and tomatidine (0.08 to 40 µM, 0.02 to 10 µM). Combinations of the different drug
565 concentrations enable to determine a Fractional Inhibitory Concentration Index (FIC index) as
566 described in (48). Drug combinations that give rise to a growth reduction of more than 50%
567 are then identified and the associate FIC index determined the properties of their

568 combination. A FIC index between 2 and 0.5 indicate an additive effect, while FIC < 0.5, a
569 synergetic effect. Average values are shown from three replicates.

570

571 (ii) Antifungal serial dilution susceptibility assay.

572 Drug susceptibility testing was also performed on solid YEPD agar plates containing specific
573 drug concentrations or vehicle (1% DMSO). Ten-fold serial dilutions of cells were spotted
574 starting with cell concentration of 10^6 cells/ml. Assays were performed at least in duplicates.

575

576 (iii) Antifungal biofilm susceptibility assay.

577 Antifungal susceptibility tests on *C. albicans* biofilms were conducted according to a published
578 protocol (49) with 48 h of biofilm formation and 48 h of antifungal treatment. Briefly, an
579 aliquot of a 100 μ l cell suspension (10^6 cells/ml) per well prepared in RPMI medium 0.2 %
580 glucose (pH 7) was deposited in each well of a 96-well plate and incubated at 37°C for 48 h to
581 allow biofilm formation. Wells were then washed twice with phosphate-buffered saline (PBS).
582 Two-fold serial dilutions of the compounds were prepared from 50 to 1.56 μ g/ml and added
583 to the wells containing the biofilms. Plates were incubated again for 48 h at 37°C and then
584 washed twice with PBS. A measurement of the metabolic activity of the sessile cells was
585 performed using a colorimetric assay with 2H-tetrazolium,2,3-bis(2-methoxy-4-nitro-5-
586 sulfophenyl)-5-[(phenylamino)carbonyl]-hydroxide salt (XTT) (X4626, Sigma Aldrich). Plates
587 were read with a spectrophotometer plate reader at 492 nm. The MIC was defined as the
588 drug concentration at which the optical density value was equal or less than 50% of the one
589 of the drug-free biofilm. Assays were performed in duplicates.

590

591 (iv) Antifungal drug time-kill assay.

592 Time-kill assay was performed as follows. Cells were cultured overnight in YEPD at 30°C,
593 adjusted to 2×10^5 cells/ml in YEPD and submitted to corresponding concentration of the
594 drugs, their combination or solvent. After 0-, 4-, 8- and 24 h incubation with the drug at 30°C
595 under agitation in a 3 ml liquid volume, cell viability was determined by plating cells on YEPD

596 agar plate for 16 h at 34°C and counting of colonies (colony forming unit, CFU). Fungicidal
597 effect was determined when at least a 2-fold log decrease of CFU/ml from the initial cell
598 density was measured.

599

600 (v) Antibacterial susceptibility assay.

601 Antibacterial susceptibility testing was carried out on the basis of CLSI Approved Standard
602 M7-A7 using microdilution method with CAMH (Cation-adjusted Müller-Hinton) broth.
603 Briefly, overnight *Escherichia coli* (ATCC® 25922) cell cultures were adjusted to McFarland 0.5
604 (10^8 Cells/ml) with NaCl. Final cell concentration was 3×10^5 CFU/ml. Drug two-fold serial
605 dilutions were prepared from 64 µg/ml to 2 µg/ml. Microplates were incubated at 37°C for
606 24 h and then MICs were read with a spectrophotometer plate reader at 450 nm. The MIC
607 was defined as the drug concentration at which the optical density was equal or decreased
608 more than 50% from that of the drug-free culture. Assays were performed in duplicates. Drugs
609 were tested at the pH at which an antifungal activity was detected during the microdilution
610 susceptibility screen (pH 7 if active at both pH values).

611

612 **Cell cytotoxicity assay**

613 Cytotoxicity assay were performed according to standard procedure with sulforhodamine B
614 (SRB) as a reporter (50). HeLa cells (ATCC CCL-2, Manassas, Virginia, USA) were cultured in
615 DMEM + 10% FBS, at 37°C with 5% CO₂. Ninety six-well plates were filled with a seeding
616 density of 10^4 cells/well. After 24 h of growth (day 1), cells were washed twice with PBS and
617 2-fold serial dilutions (starting at 100 µg/ml) of the compounds were added to the cells and
618 incubated for 48 h. The starting amount of cells was monitored by fixing the cells at day 1. At
619 day 3, cells were all washed twice with PBS and then fixed and labelled as described in the
620 standard procedure. Optical density was measured at 492 nm and the percentage of killed
621 cells could be determined using the following formula: $100 - (OD_{day3} / OD_{day1}) \times 100$. LD50 (lethal
622 dose 50) corresponded to the concentrations at which at least 50% of cells were killed. Assays
623 were performed in duplicates. The selectivity index (SI) was then calculated by dividing the
624 LD50 by the MIC against *C. albicans*.

625

626 **Hierarchical clustering of activity profiles**

627 MIC of the 40 active NPs against the 7 yeast strains was used to generate the heatmap and
628 pH of activity was indicated by a color scheme. NPs that showed activity at both pH were
629 labelled as active at neutral pH. Cluster analyses were performed by calculating the distance
630 matrix using the Euclidean method followed by Ward (Ward.D) hierarchal clustering using the
631 gplots package in R version 3.3.2.

632

633 **Calcofluor white staining**

634 Yeast cells were grown overnight and washed twice with PBS. Cells (10^5) were resuspended
635 in 200 μ l RPMI 0.2% glucose in 96-well plate and incubated for 3 h at 37°C. Ten μ l of calcofluor
636 white stain (Sigma) was added to the well. After 10 min at room temperature, 4 μ l of cell
637 suspension was mixed with 2 μ l of Mowiol (Sigma) and directly thereafter, fluorescence
638 microscopy was performed with a Zeiss Axioplan 2 microscope (Zeiss, Oberkochen, Germany).

639

640 **Sterol content analysis**

641 (i) Total sterol extraction

642 About 2×10^5 /ml cells were cultured for 16 h at 30°C under agitation in 15 ml YEPD
643 supplemented with the indicated tomatidine concentrations (with 1% DMSO final
644 concentration). Cells were treated with tri-chloro-acetic acid (TCA) to a final concentration of
645 5% to stop metabolism and the cells incubated for 10 minutes on ice. Harvested cells were
646 then washed twice, first in 5% TCA (distilled water) and then in distilled water to remove
647 traces of the YEPD medium. Cells were resuspended in 3 ml of distilled water and 10^9 cells
648 were used to perform total sterol extraction as described in (51) and to determine the
649 amounts of total cellular sterols (esterified and non-esterified). Cells were resuspended in 1
650 ml 60% KOH to which 1 ml of 0.5% pyrogallol-containing methanol and 1 ml of methanol were
651 added in a screw cap glass tube. Tubes were heated at 85°C for 2 h and returned to room
652 temperature. Sterols were extracted three times with 2 ml of petroleum ether (high boiling

653 point). The combined petroleum ether phases were dried under N₂ flow, resuspended in 1/1
654 methanol/chloroform (v/v) and sonicated for 5 min for further analysis by GC–MS.

655

656 (ii) GC-MS analysis

657 Sterols were analysed by gas liquid chromatography-mass spectrometry (GC- MS) as
658 described in (51). *ERG4* and *ERG6* deletion mutant extracts were used to determine the
659 positions of the known sterols.

660

661 **Tomatidine formulation**

662 The nanosuspension was obtained by adding 6 mg of tomatidine, 300 µL 2% (w/V) D- α -
663 tocopherol polyethylene glycol 1000 succinate (TPGS, Sigma-Aldrich) as stabilizer and 700 µL
664 of purified water as non-solvent in a 2 ml tube (tomatidine/TPGS (50/50)). A wet milling with
665 579 mg of zirconium bead (BeadBug TM, Sigma-Aldrich, prefilled tubes of 2.0 mL with 0.5 mm
666 Zirconium beads, triple-pure, high impact) was performed for 70 hours on a vortex Genie 2.
667 Then, the nanosuspensions were frozen by dipping the tubes in liquid nitrogen. Subsequently,
668 the nanosuspensions were lyophilized for 48 hours using a Christ Alpha 2-4 LD Plus freeze-
669 dryer. The particle size distribution was determined by dynamic light scattering (DLS) using a
670 Zetasizer 3000HSa. The samples were dispersed in 9/10 volume of filtered purified water and
671 stirred for 20 min with a vortex to ensure a uniform dispersion free of aggregates. 1/10
672 volume of PBS 10X was added prior injection.

673

674

675

676 **Transmission electron microscopy (TEM)**

677 The *C. albicans* strain CAF2-1 was grown in YNB liquid cultures for 2 h at 37°C (15 ml plastic
678 tubes). Next, miconazole (1 mg/ml in DMSO) was added at a concentration of 10 µg/ml and
679 cultures were grown during 18 h to evaluate the cytotoxic effect of this commercial product

680 on the yeast strain. The cytotoxic effect of tomatidine was evaluated according to the same
681 experiment, except that this compound (1 mg/ml in DMSO) was added at a concentration of
682 20 μ M and cultures were grown during 18 h. Cell preparation was performed as described in
683 (12). Thin sections were observed with a transmission electron microscope (Philips CM10)
684 with a Mega View II camera. Control cells were obtained in the same way without drug
685 treatment.

686

687 **Selection of NPs**

688 A smart chemical library containing 199 natural products (NPs) with potential antifungal
689 activity was constructed. Among these compounds, 53% were previously isolated from crude
690 plant extracts that presented an antifungal activity in our laboratory. The bioguided isolation
691 process of these compounds were performed by bioautography using wild and genetic
692 modified strains of *C. albicans* (12). In parallel, NPs with i) structures closely related to NPs
693 possessing antifungal activity from published sources and ii) with unknown antifungal activity,
694 were selected and acquired in commercial catalogues (compounds are listed in
695 Supplementary Table S1). The identity and the purity of the commercial compounds obtained
696 were systematically performed by nuclear magnetic resonance (NMR) and high resolution
697 mass spectrometry (HRMS) analysis.

698

699 **Genome-wide transcriptional analysis**

700 (i) RNA extraction and processing

701 Overnight YEPD culture of *C. albicans* SC5314 strain was diluted 1:200 in 5 ml YEPD media and
702 incubated under agitation at 30° C until early exponential growth phase ($OD_{540} = 0.3$). Fifty μ l
703 of solvent (DMSO) or 250 μ M tomatidine (diluted in DMSO) were added to the culture to
704 reach a concentration 1% DMSO and 2.5 μ M tomatidine. Total RNA was extracted after 1- or
705 3 h tomatidine/solvent exposure by mechanical disruption of the cells with glass beads as
706 previously described (52). Experiments were carried out in triplicates with 12 samples. Total
707 RNA extracts were treated with DNase using the DNA-free kit (Ambion-Life Technologies, Zug,

708 Switzerland) and RNA quality and integrity was verified with Fragment Analyzer™ Automated
709 CE System (Advanced Analytical). One µg of RNA was used to create sequencing libraries
710 through standard Illumina TruSeq stranded mRNA protocol. Each library (sample) received a
711 different index enabling several libraries to be multiplexed. Before RNA sequencing, libraries
712 were analyzed with a fragment analyzer to assess quality and fragment size and with a Qubit
713 fluorometer (Invitrogen) to determine cDNA concentration. Libraries were kept at -20°C until
714 sequencing.

715

716 (ii) RNA sequencing

717 The 12 libraries were run on Illumina HiSeq platform (HiSeq2500). Sequencing data were
718 processed using Illumina Pipeline software. Reads were filtered, trimmed, and counts align to
719 the SC5314 *C. albicans* reference genome using CLC workbench pipeline. The numbers of read
720 counts per gene locus was extracted. All reads were deposited at GEO under accession
721 number GSE96965.

722

723 (ii) RNA seq data analysis

724 Data normalization and gene expression analysis were performed in R (v3.2.3), using
725 Bioconductor packages (as described in (23)). The read count data were normalized using
726 TMM (trimmed mean of M-values) method available in the R package edgeR (53) and
727 transformed into log2 counts per million by Voom method from R package Limma (54). This
728 package was then used to apply a linear model with one factor per condition (4 conditions:
729 untreated 1h, tomatidine-treated 1h, untreated 3h, tomatidine-treated 3h (all in triplicates))
730 to the transformed data. Two contrasts representing the difference between tomatidine-
731 treated and untreated cells at each drug exposure time (1- and 3 h) were extracted from the
732 linear model to result in a moderated t statistic for all genes expressed.

733

734 **qPCR analysis**

735 Total RNA (same RNA samples as those used for RNA-seq experiments: 12 samples
736 representing 4 conditions in 3 biological replicates) were treated with DNase and 1 µg of
737 treated-RNA was used as template for cDNA synthesis using the high-fidelity cDNA synthesis
738 kit (Roche Diagnostics, Switzerland). Real-time quantitative PCR (qPCR) were performed to
739 determine relative gene expression of using primers and TaqMan probes (FAM-TAMRA
740 modified) with the iTaq Supermix with ROX (Bio-Rad AG, Switzerland) in a StepOnePlus real-
741 time PCR system (Applied Biosystems-Life Technologies, Switzerland). Each reaction was run
742 in duplicate. Primers and probes are listed in supplementary Table S3. Relative transcript
743 quantities were assessed using the $2^{(-\Delta\Delta CT)}$ method (55) to determine a normalized
744 expression ratio with *ACT1* as reference gene.

745

746 ***C. albicans* *ERG6* deletion strain constructions**

747 To delete the first allele of *ERG6*, two fragments of 571 bp and 569 bp of flanking 5'- and 3'-
748 UTR regions, respectively, were PCR amplified on SC5314 DNA with the following primer pairs
749 (*ERG6_5For_KPN1* and *ERG6_5Rev_Xho1*; *ERG6_3For_SacII* and *ERG6_3Rev_SacI*). These
750 primers contained restriction sites in order to insert the two amplicons sequentially in pSFS2A
751 (56). The plasmid obtained (pSD1) was then digested with *ScaI* and transformed in *C. albicans*
752 SC5314. Yeasts were transformed by a lithium-acetate procedure previously described (57).

753 Transformants were positively selected on YEPD plate containing 200 µg/ml nourseothricin
754 (Nour) (Werner Bioagents, Germany). The Nour selective cassette was then removed by
755 growing cells in YEPD media containing 2% maltose. Nour susceptible cells were used to
756 deleted the second allele. The same strategy was repeated for the second allele but, in order
757 to achieve it, a different 3'-end *ERG6* homologous region (at the end of the CDS and upstream
758 of the first allele, 343 bp) was amplified with the primers *ERG6_3CDSFor_SacII*,
759 *ERG6_3CDSRev_SacI*. The resulting construct was named pSD4.

760 Nour-resistant transformants were phenotypically screened using a simplified ergosterol
761 extraction and detection method (58). An alteration of the expected UV spectrophotometric
762 sterol profiles was detected in some transformants with an additional peak of absorbance at
763 230 nm indicating a perturbation in the ergosterol pathway and thus suggesting the deletion

764 of the second *ERG6* allele. GC-MS analysis confirmed the loss of *ERG6* function with
765 zymosterol being most abundant sterol (data not shown).

766

767 ***S. cerevisiae* forward genetic screen**

768 (i) Deletion of *MSH2* in *S. cerevisiae* DSY4743

769 The *MSH2 PDR5* deletion strain was constructed from a *PDR5* deletion strain (DSY4743) using
770 a PCR-based gene deletion approach as described previously [61]. Primers For_msh2_Sc and
771 Rev_msh2_Sc were used to amplify the *HIS3* selection marker. The *HIS3*-containing amplicons
772 was purified using the Nucleospin Gel and PCR Clean-Up kit (Macherey-Nagel, Düren,
773 Germany) according to the manufacturer's instructions and used to transform *S. cerevisiae*
774 *pdr5Δ* (DSY4743) using the standard lithium acetate protocol.

775

776 (ii) Selection of resistant mutants

777 Overnight cultures of *msh2Δ prd5Δ* strain (P1) cells were plated on solid media containing
778 several tomatidine concentration and incubated for two to seven days. One pop-out (R1) was
779 identified and resistance to tomatidine was confirmed using MIC broth dilution method. The
780 other tomatidine-resistant strains were identified as described previously by Ojini and
781 Gammie (28) with few modifications. The *msh2Δ prd5Δ* strain (P1) was grown to saturation
782 in 5 ml of YEPD medium at 30°C. Overnight cultures were diluted 1:200 in YEPD medium
783 containing 50 µg/ml of ampicillin and grown in the presence of 10 µM of tomatidine in 96-
784 well microtiter plates (Costar) in a shaking incubator at 30°C for 72 h. Cells were diluted 1:200
785 in the respective media and distributed into new 96-well plates and grown for 48 h at 30°C in
786 absence of the drug. The cultures were then diluted 1:200 in media containing tomatidine in
787 a new 96-well plate and optical density at 540 nm was recorded at six different time points
788 (T0, T16, T20, T24, T38 and T42) over a 42 h period. Several wells were selected based on
789 their growth profiles in presence of drug with high optical density after 40 h, and plated onto
790 YEPD containing 7.5 µM of tomatidine and incubated for four days at 30°C. Resistant pop-
791 outs were cultured overnight and resistance was retested with broth dilution method. Four

792 resistant strains arising from 2 different cultures (R2 and R3.1, R3.2, R3.3) were identified and
793 together with R1 submitted to whole genome sequencing.

794

795 (iii) Whole-genome sequencing

796 The five tomatidine-resistant cells and the parental strain were grown overnight in YEPD
797 medium at 30°C under constant agitation. Genomic DNA was extracted from yeast using the
798 Gentra Puregene Yeast/Bact. Kit (Qiagen, Hilden, Germany) with RNase treatment. DNA
799 concentration was verified by Qubit 2.0 Fluorometer (Thermo Fisher) and adjusted to 10
800 ng/ml for whole-genome sequencing. DNA quality was verified with Fragment Analyzer
801 (Advanced Analytical Technologies, Ankeny, IA, USA). Whole-genome sequencing was
802 performed at Fasteris SA (Plan-les-Ouates, Switzerland) using TrueSeq Nano DNA library
803 preparation and Illumina MiSeq. Paired-end read of 250 bp were performed, giving an
804 average of 118x coverage of each genome. The sequencing data was analysed using the CLC
805 Genomics Workbench (v.9.5.2) (Qiagen). The sequence reads were mapped to the reference
806 genome S288C. The average percentage of mapped reads was 93%. Mutations were
807 identified using the Variant Detector option by comparing P1 and tomatidine-resistant strains
808 to the reference genome. Functional consequences option was used to identify amino acid
809 changes. To identify mutations specific to the resistant strains, the variants of tomatidine-
810 resistant strains were compared to the P1 strain using the Compare Variants option. Only
811 non-synonymous mutations were taken into account with a frequency of $\geq 50\%$ of reads for
812 insertions and deletions (INDELS) and of $\geq 90\%$ of reads for single nucleotide variants (SNV).
813 Genome data are deposited at NCBI under BioProject PRJNA380059.

814

815 (iv) Construction of *S. cerevisiae* *ERG6* point mutated strains

816 To introduce specific point mutations in *ERG6* in *S. cerevisiae*, the clustered regularly
817 interspaced short palindromic repeat (CRISPR)-Cas9 genome editing system was used as
818 described previously (30). All primers are listed in Supplementary Table S3. A 20-nucleotide
819 guide sequence was selected using an online tool named CHOPCHOP
820 (<http://chopchop.cbu.uib.no/>). PAM 11 (position 251 849) and PAM 19 (position 252 596)

821 were chosen for the D249G (position 252 245) and G132D mutation (position 252 596),
822 respectively. The two repair fragments were constructed as described previously (59). All
823 fragments were purified using Nucleospin Gel and PCR Clean-up Kit (Macherey-Nagel).

824 Genome editing was performed by co-transformation of the guide, pMEL10 and repair
825 fragments into *S. cerevisiae* IMX581 and selection was carried out in YNB agar lacking uracil.
826 Verification of introduced mutations was performed by PCR amplification of *ERG6*
827 (*Erg6_verif_for* and *Erg6_verif_rev*, see supplementary Table S3) and by sequence analysis as
828 described above. The gRNA plasmid pMEL10 were removed by counter-selection pressure
829 with 5-fluoroorotic acid (5-FOA, Toronto Research Chemicals, TRC) and effect of mutations on
830 tomatidine susceptibility was verified by broth dilution method.

831

832 **Mice experiments and ethics statement**

833 All animal experiments were performed at the University Hospital Center of Lausanne with
834 approval through the Institutional Animal Use Committee, Affaires Vétérinaires du Canton de
835 Vaud, Switzerland (authorization n° 1734.3), according to decree 18 of the federal law on
836 animal protection. Female BALB/c mice (8 weeks-old; Charles River France) were housed in
837 ventilated cages with free access to food and water. SC5314 strain was grown overnight under
838 agitation at 30°C in YEPD medium, subsequently diluted 100-fold in YEPD medium and grown
839 again overnight under agitation at 30°C. Overnight cultures were washed twice with PBS and
840 resuspended in 5 ml PBS. The concentration of each culture was measured through optical
841 density, and each strain was diluted in PBS to the desired concentration (4×10^5 CFU/ml).
842 Mice were injected through the lateral tail vein with 250 μ l of a cell suspension containing 1.6
843 $\times 10^6$ cells/ml. At 7-, 24- and 31 h post-infection, tomatidine formulation or placebo was
844 administered through intraperitoneal (IP) injection in a volume of 200 μ l. At 48 h post-
845 infection, the kidneys were recovered, weighted and the CFU were determined as previously
846 described (60). The ratio CFU/g of kidney was determined. Outliers analysis was first
847 performed in GraphPad Prism using default parameters (ROUT, Q= 1%) and final number of
848 individuals per group were 10 for placebo and 9 for tomatidine-treated mice. Statistical
849 analyses of the differences between CFU values were performed using the Mann–Whitney
850 test. The weight and temperature of the animals were monitored daily.

852 **Figures Legends**

853

854 **Figure 1:** Heatmap and cluster analysis of the activity profiles for 33 active NPs.

855 MIC values of the NPs against seven yeast species are ordered by hierarchical clustering using
856 Euclidean distance method and represented with a heatmap intensity color code (Heatmap
857 legend, MIC ($\mu\text{g/ml}$): dark blue = 0.125, light blue = 32, white >32 (inactive)). The pH at which
858 activity was detected is indicated by a color code in the first column (orange: pH 4.6, green:
859 pH 7). Fluconazole and caspofungin were added to the list of compounds and their clusters
860 highlighted in red and purple colors, respectively.

861

862 **Figure 2:** Tomatidine susceptibility assays on *Candida* spp.

863 (A) Structure of tomatidine.

864 (B) Spotting susceptibility assays of *Candida* spp. in YEPD agar plates. Ten-fold serial dilutions
865 of indicated strains were spotted onto agar plates containing increasing tomatidine
866 concentrations (DMSO, 1, 5, 10 μM) and were incubated for 1 day at 34°C. Azole-resistant
867 strains are indicated with "AR".

868 (C) Calcofluor white staining of *C. albicans* cells exposed to tomatidine (magnification: 100X).
869 Cells were exposed to 2.5 μM tomatidine for 3 h or untreated (DMSO 1%) in RPMI (0.2 %
870 glucose) at 37°C and then labelled with calcofluor white to stain chitin. Depicted cells are
871 representative of the vast majority of cells from three independent experiments.

872 (D) Time-kill assay for tomatidine treated-cells. 2×10^5 *C. albicans* and *C. krusei* cells were
873 treated with different supra-MIC concentrations of tomatidine in YEPD and the CFU/ml were
874 determined after 4-, 8- and 24 h of exposure. Relative expressions to initial (T0) CFU/ml values
875 (100%) were calculated. Experiments were performed in duplicates with average and SEM
876 graphically represented. Y-axis is on log₁₀ scale.

877

878 **Figure 3:** Genome-wide transcriptional analysis of tomatidine-treated cells.

879 (A) Venn diagram showing the number of genes differentially regulated by tomatidine (2.5
880 μM) as compared to control cells after 1- and 3 h exposure (FDR >0.05 , fold-change ≥ 2).

881 (B) GO term analysis of the commonly up- and down- regulated genes upon 1h and 3h
882 tomatidine treatment. GO term and their FDR value are indicated and the expression of
883 corresponding genes represented with a heat-map for both drug exposure durations. The
884 scaled expression of each gene, denoted as the row Z-score, is plotted in red-green colour
885 scale with red indicating high expression and blue indicating low expression. A Z-score of 0
886 (black) correspond to the mean expression level of a particular gene and Z-score scale
887 indicates the numbers of standard deviation (positive or negative). Genes within each GO
888 term group are listed in an increasing adjusted p-value manner.

889 (C) qPCR analysis of *ERG11*, *ERG6* and *ERG4* expression in tomatidine-treated cells. Fold-
890 change increase in expression compared to untreated cells is represented in a barplot. T-test
891 analysis were performed between treated and untreated group for each gene relative
892 expression and significant differences detected (all p-value < 0.05).

893 (D) Gene set enrichment analysis (GSEA) of *C. albicans* genes regulated by tomatidine. The list
894 of drug-regulated genes was generated from published transcriptional data (see
895 supplementary File S2, *Candida_drug_treatment.gmt*). Tomatidine regulated genes (3h) were
896 ranked according to their fold-change. The list was then imported into the GSEA software.
897 Analysis parameters were as follows: norm, meandiv; scoring_scheme, weighted; set_min,
898 15; nperm, 1000; set_max, 500. GSEA results were uploaded into Cytoscape 3.0 with the
899 following parameters: p-value cut-off, 0.01; FDR q-value, 0.05. Red nodes represent enriched
900 gene lists in upregulated genes from the GSEA. Blue nodes represent enriched gene lists in
901 downregulated genes from the GSEA. Nodes are connected by edges when overlaps exist
902 between nodes. The size of nodes reflects the total number of genes that are connected by
903 edges to neighbouring nodes. The labels of the list (corresponding to list of genes up- and
904 down-regulated by drug treatments, gene deletions or stress conditions) are indicated next
905 to the nodes (details in supplementary File S2). XSA, XSB: oxidative shock, HU6H: hydroxyurea
906 treatment.

907

908 **Figure 4:** Total sterol analysis in tomatidine-treated cells with GC-MS.

909 (A) Sterol composition of *C. albicans* cells treated with increasing amount of tomatidine. GC-
910 MS analysis of cells exposed to tomatidine with the percentage of the different sterols are
911 shown in a bar plot. Bar colors correspond to sterol molecules illustrated on the left panel
912 where the last steps in the ergosterol biosynthetic pathway and an alternative pathway
913 following *ERG6* inhibition are shown. Experiment was repeated (duplicates) and gave similar
914 results.

915 (B) Susceptibility assay of the *C. albicans* *ERG6* deletion strain. Wild type (*ERG6/ERG6*),
916 heterozygous (*ERG6/erg6Δ*) or homozygous *ERG6* (*erg6Δ/erg6Δ*) deletion strains were
917 subjected to serial dilution susceptibility assay on YEPD plate containing indicated
918 concentration of tomatidine or fluconazole (right panel) and to standard MIC determination
919 assay in YNB pH 7 media (left panel).

920

921 **Figure 5:** Forward genetic approach in *S. cerevisiae*.

922 (A) Tomatidine susceptibility of resistant mutants. Fungal cells were spotted on YEPD agar
923 containing different concentrations of tomatidine (and a drug-free control) as indicated.
924 Plates were incubated at 30°C for 48 h.

925 (B) Alignment of the sterol methyltransferase protein sequences. Sequence alignment of
926 sterol methyltransferase (SMT) from distant eukaryote species (Fungi: *S. cerevisiae*, *C.*
927 *albicans*, *Pneumocystis carinii* (*P. carinii*), *Giberella zeae* (*G. zeae*); Green plants: *Glycine max*
928 (*G. max*), *Arabidopsis thaliana* (*A. thaliana*); Euglenozoa: *Trypanosoma brucei* (*T. brucei*).
929 Conserved residues are highlighted in grey. Black boxes indicated highly conserved region that
930 have a role in substrate binding and enzymatic activity (29). The two mutated residues found
931 in sterol C-24 methyltransferase (Erg6) of tomatidine-resistant strains (G132D and D249G) are
932 represented by red boxes and are conserved across all aligned sequences. Strains carrying the
933 *ERG6* mutation are indicated below the alignment. Protein sequences were retrieved from
934 the following Genbank sources: *S. cerevisiae* KZV08836, *C. albicans* AOW28252, *P. carinii*
935 KTW25893, *G. zeae* ESU10532, *G. max* NP_001238391, *A. thaliana* NP_173458, *T. brucei*

936 AAZ40214. A multiple alignment was performed using MUSCLE in the Geneious software
937 (version 9.1.4, default parameters).

938

939 **Figure 6:** Tomatidine targets efflux pumps and induces only partial cross-resistance with
940 fluconazole

941 (A) Serial dilution susceptibility assays of *C. albicans* strains carrying deletion in efflux pumps
942 (genotype are indicated) on YEPD agar plates. Ten-fold serial dilutions of indicated strains
943 were spotted onto agar plates containing increasing tomatidine concentrations (\emptyset , 0.5, 1, 5
944 μ M) and were incubated for 1 day at 34°C.

945 (B) Tomatidine MICs of two matched pairs of azole-susceptible (AS) and azole-resistant (AR)
946 strains. MICs were performed in YNB media for both tomatidine and fluconazole and are
947 indicated by the arrows. The “ Δ ” symbol indicates the MIC fold-change differences between
948 AS and AR matched strains.

949 (C) Two heatmaps representing color plots of checkerboard MIC tests. Each box corresponds
950 to the relative growth (compared to drug-free control) resulting from a specific combination
951 of tomatidine and fluconazole (RPMI, pH 7). Individual MICs of each drug are underlined by
952 black lines. Black zones correspond to additive interactions (FIC index between 2 and 0.5).
953 Yellow zones indicate a synergetic effect (FIC < 0.5).

954 (D) Time-kill assay of *C. albicans* and *C. krusei* cells exposed to a combination of tomatidine
955 and fluconazole. Cells were treated with indicated amount of drugs, individually or in
956 combination, in YEPD and CFU/ml determined after 4-, 8- and 24 h of exposure and relative
957 expression to initial (T0) CFU values (100%) calculated. Experiments were performed in
958 duplicates with average and SEM graphically represented. Y-axis is on log₁₀ scale. A dashed
959 line was plotted on *C. krusei* graph at Y = 1 to delimit the fungicidal threshold (2-times log₁₀
960 decrease).

961

962

963

964 **Figure 7:** *In vivo* efficacy of tomatidine in a mice model of systemic infection

965 (A) Schematic representation of the infection and treatment protocol. Mice were infected
966 with 4×10^5 CFU through the tail vein. Tomatidine treatment (50 mg/kg) (or placebo) was
967 administered intraperitoneally (IP) at 7-, 24- and 31 h post-infection (pi). At 48 h pi, animals
968 were sacrificed and kidneys collected for CFU determination.

969 (B) Fungal burden in tomatidine-treated mice. CFU per g of kidney are plotted for placebo and
970 tomatidine-treated mice. Significant differences in CFU distribution were assessed using
971 Mann Whitney test (n=9,10; p-value = 0.031).

972

973 **Table 1:** List of the 33 NPs with their biological activity profiles

Name	N°	Source	CAS number	<i>C. albicans</i> MIC (µg/ml)		<i>C. albicans</i> biofilm MIC (µg/ml)	<i>E. coli</i> MIC (µg/ml)	Cytotoxicity assay LD ₅₀	Selectivity Index LD ₅₀ /MIC <i>C. albicans</i>
				pH 7	pH 4.6				
morindone	1	<i>Morinda tomentosa</i>	478-29-5	32 ^{a)}	>32	6.25	>64	ND ^{c)}	ND
lucidine ω-methyl ether	2	<i>Morinda tomentosa</i>	NA ^{b)}	>32	32	NA	>64	ND	ND
morindoquinone	3	<i>Morinda tomentosa</i>	New NP	>32	16	25	>64	>100	6.25
avocadene	4	<i>Persea americana</i>	24607-08-7	16	8	25	>64	50	6.25
plumbagin	5	Sigma	481-42-5	2	4	50	>64	100	50
alpha-hederin	6	<i>Schefflera systila</i>	27013-91-8	16	4	50	>64	100	6.25
Glc-3 medicagenic acid	7	<i>Dolichos kilimandscharicus</i>	49792-23-6	>32	2	>50	>64	>100	≥100
2-Propen-1-one, 1-(2,4-dihydroxy-6-methoxy-3,5-dimethylphenyl)-3-phenyl	8	<i>Myrica serrata</i>	65349-31-7	>32	8	25	>64	25	3.125
O-methyllawsone	9	<i>Swertia calycina</i>	2348-82-5	16	8	12.5	>64	25	1.5625
dihydrochelerythrine	10	<i>Fagara zanthoxyloides</i>	6880-91-7	8	>32	25	>64	25	3.125

simplexene D	11	<i>Swartzia simplex</i>	New NP	>32	16	25	>64	>100	≥12.5
waltherione G	12	<i>Waltheria indica</i>	1632043-42-5	>32	32	25	>64	ND	ND
waltherione F	13	<i>Waltheria indica</i>	1632043-41-4	>32	8	12.5	>64	50	6.25
8-deoxoantidesmone	14	<i>Waltheria indica</i>	NA	>32	16	25	>64	>100	12.5
waltherione E	15	<i>Waltheria indica</i>	954367-81-8	>32	4	12.5	>64	>100	≥50
pterostilbene	16	Sigma	537-42-8	32	32	50	>64	ND	ND
(5S,10S)-11,15(S)- dihydroxy-12- methoxyswartziarboreol G	17	<i>Swartzia simplex</i>	1830306-56-3	32	16	50	>64	50	3.125
pulsatilla saponin D	18	<i>Odondatenia puncticulosa</i>	68027-15-6	>32	16	>50	>64	100	6.25
3β-O-[β-D-xylopyranosyl- (1→3)]-α-L- rhamnopyranosyl-(1→2)- [β-D- glucopyranosyl- (1→4)]-α-L- arabinopyranosyl]	19	<i>Odondatenia puncticulosa</i>	NA	>32	8	>50	>64	50	6.25
hederagenin									
garcinone C	20	Phytolab	76996-27-5	8	32	>50	>64	50	6.25

pennogenin tetraglycoside	21	Phytolab	68124-04-9	4	8	>50	>64	6.25	1.5625
tomatidine hydrochloride	22	Phytolab	6192-62-7	< 1	16	>50	>64	>100	≥ 200
formosanin C	23	Phytolab	50773-42-7	1	4	25	>64	12.5	12.5
medicagenic acid	24	Phytolab	599-07-5	>32	2	>50	>64	>100	≥100
pyridoxatin	25	Sigma	135529-30-5	4	32	25	16	100	25
Isograndifoliol	26	<i>Perovskia atriplicifolia</i>	1445475-53-5	16	32	>50	>64	100	25
taxodion	27	<i>Salvia leriifolia</i>	19026-31-4	16	8	12.5	>64	50	3.125
waltherione N	28	<i>Waltheria indica</i>	New NP	>32	32	12.5	>64	ND	ND
5(R)-vanessine	29	<i>Waltheria indica</i>	New NP	>32	32	12.5	>64	ND	ND
waltherione Q	30	<i>Waltheria indica</i>	New NP	>32	32	25	>64	ND	ND
antidesmone	31	<i>Waltheria indica</i>	222629-77-8	>32	32	12.5	>64	ND	ND
waltherione I	32	<i>Waltheria indica</i>	1632043-44-7	>32	32	12.5	>64	ND	ND
waltherione J	33	<i>Waltheria indica</i>	1632043-46-9	>32	16	12.5	>64	50	3.125

974

975 a): numbers in bold face indicate that the value met threshold requirements.

976 b): NA: not available

977 c): ND: not determined

978 **Table 2:** Tomatidine and fluconazole MICs for *Candida* spp.

Strain	RPMI pH 7		YNB pH 7		YPED pH 6.5	
	Tomatidine (μ M)	Fluconazole (μ g/ml)	Tomatidine (μ M)	Fluconazole (μ g/ml)	Tomatidine (μ M)	Fluconazole (μ g/ml)
<i>C. albicans</i> (SC5314)	0.625	0.125	0.625	0.5	1.25-2.5	0.5
<i>C. krusei</i> (DSY471) (AR) ^{a)}	0.625	32	0.625	32	0.3125	>128
<i>C. tropicalis</i> (DSY472)	1.25	0.25 - 0.5	5	>128	2.5	32
<i>C. parapsilosis</i> (DSY473)	10	2	2.5	8	5	16
<i>C. glabrata</i> (DSY562)	>40	2	>40	32	>40	64
<i>C. glabrata</i> (DSY562) (AR)	>40	128	>40	>128	>40	>128

979 ^{a)}: (AR): azole-resistant strains

980

981 **Table 3:** MICs of *S. cerevisiae* strains to different antifungals

Strain	MIC			
	Tomatidine (μM)	Fluconazole ($\mu\text{g/ml}$)	Caspofungin ($\mu\text{g/ml}$)	Amphotericin B ($\mu\text{g/ml}$)
P1	1.25	4	0.25	2
R2	1.25	4	0.25	2
R3.1	5	4	0.25	2
R3.2	5	4	0.25	2
R3.3	2.5	4	0.25	2
R1	5	4	0.25	2
IMX581	5	16	0.125	2
D249G	20	16	0.125	2
G132D	20	4	0.125	2

982

983

984

985 **Table 4:** Susceptibility of *C. albicans* clinical isolates to tomatidine and fluconazole

Strain	MIC (YNB pH 7)	
	Tomatidine (μM)	Fluconazole ($\mu\text{g/ml}$)
DSY731 (AS) ^{a)}	0.625	1
DSY732 (AR) ^{a)}	2.5	64
DSY1841 (AS)	2.5	8
DSY1843 (AR)	5	32

986 ^{a)} (AR): azole-resistant strains, (AS): azole-susceptible strains

987

988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038

References

1. **Denning DW, Hope WW.** 2010. Therapy for fungal diseases: opportunities and priorities. *Trends Microbiol* **18**:195-204.
2. **Thompson GR, 3rd, Cadena J, Patterson TF.** 2009. Overview of antifungal agents. *Clin Chest Med* **30**:203-215, v.
3. **Barker KS, Rogers PD.** 2006. Recent insights into the mechanisms of antifungal resistance. *Curr Infect Dis Rep* **8**:449-456.
4. **Sanglard D, Bille J, Calderone R.** 2002. Current understanding of the mode of action and of resistance mechanisms to conventional and emerging antifungal agents for treatment of *Candida* infections, p 349-383, *Candida and Candidiasis*.
5. **Ostrosky-Zeichner L, Casadevall A, Galgiani JN, Odds FC, Rex JH.** 2010. An insight into the antifungal pipeline: selected new molecules and beyond. *Nat Rev Drug Discov* **9**:719-727.
6. **Pfaller MA, Moet GJ, Messer SA, Jones RN, Castanheira M.** 2011. Geographic variations in species distribution and echinocandin and azole antifungal resistance rates among *Candida* bloodstream infection isolates: report from the SENTRY Antimicrobial Surveillance Program (2008 to 2009). *J Clin Microbiol* **49**:396-399.
7. **Pfaller MA, Diekema DJ, Castanheira M, Jones RN.** 2011. Definitions and Epidemiology of *Candida* Species not Susceptible to Echinocandins. *Current Fungal Infection Reports* doi:10.1007/s12281-011-0053-y.
8. **Larsson J, Gottfries J, Muresan S, Backlund A.** 2007. ChemGPS-NP: tuned for navigation in biologically relevant chemical space. *J Nat Prod* **70**:789-794.
9. **Newman DJ, Cragg GM, Snader KM.** 2003. Natural products as sources of new drugs over the period 1981-2002. *J Nat Prod* **66**:1022-1037.
10. **Roemer T, Xu D, Singh SB, Parish CA, Harris G, Wang H, Davies JE, Bills GF.** 2011. Confronting the challenges of natural product-based antifungal discovery. *Chem Biol* **18**:148-164.
11. **Znaidi S.** 2015. Strategies for the identification of the mode-of-action of antifungal drug candidates, p 183-210. *In* Coste AT, Vandeputte P (ed), *Antifungals: From Genomics to Resistance and the Development of Novel Agents*. Caister Academic Press, United Kingdom.
12. **Favre-Godal Q, Dorsaz S, Queiroz EF, Conan C, Marcourt L, Wardojo BP, Voinesco F, Buchwalder A, Gindro K, Sanglard D, Wolfender JL.** 2014. Comprehensive approach for the detection of antifungal compounds using a susceptible strain of *Candida albicans* and confirmation of in vivo activity with the *Galleria mellonella* model. *Phytochemistry* **105**:68-78.
13. **Coste A, Amorim-Vaz S.** 2014. Animal Models to study Fungal Virulence and Antifungal Drugs, p 288-315. *In* Coste A, Vandeputte P (ed), *Antifungals*. Casiter Academic Press, Norfolk, UK.
14. **Taff HT, Mitchell KF, Edward JA, Andes DR.** 2013. Mechanisms of *Candida* biofilm drug resistance. *Future Microbiol* **8**:1325-1337.
15. **Cretton S, Dorsaz S, Azzollini A, Favre-Godal Q, Marcourt L, Ebrahimi SN, Voinesco F, Michellod E, Sanglard D, Gindro K, Wolfender JL, Cuendet M, Christen P.** 2016. Antifungal Quinoline Alkaloids from *Waltheria indica*. *J Nat Prod* **79**:300-307.
16. **Polacheck I, Zehavi U, Naim M, Levy M, Evron R.** 1986. Activity of compound G2 isolated from alfalfa roots against medically important yeasts. *Antimicrob Agents Chemother* **30**:290-294.
17. **Sandrock RW, Vanetten HD.** 1998. Fungal Sensitivity to and Enzymatic Degradation of the Phytoanticipin alpha-Tomatine. *Phytopathology* **88**:137-143.
18. **Simons V, Morrissey JP, Latijnhouwers M, Csukai M, Cleaver A, Yarrow C, Osbourn A.** 2006. Dual effects of plant steroidal alkaloids on *Saccharomyces cerevisiae*. *Antimicrob Agents Chemother* **50**:2732-2740.

- 1039 19. **Medina JM, Rodrigues JC, Moreira OC, Atella G, Souza W, Barrabin H.** 2015. Mechanisms of
1040 growth inhibition of *Phytomonas serpens* by the alkaloids tomatine and tomatidine. *Mem*
1041 *Inst Oswaldo Cruz* **110**:48-55.
- 1042 20. **Medina JM, Rodrigues JC, De Souza W, Atella GC, Barrabin H.** 2012. Tomatidine promotes
1043 the inhibition of 24-alkylated sterol biosynthesis and mitochondrial dysfunction in
1044 *Leishmania amazonensis* promastigotes. *Parasitology* **139**:1253-1265.
- 1045 21. **Desmond E, Gribaldo S.** 2009. Phylogenomics of sterol synthesis: insights into the origin,
1046 evolution, and diversity of a key eukaryotic feature. *Genome Biol Evol* **1**:364-381.
- 1047 22. **Iorio F, Rittman T, Ge H, Menden M, Saez-Rodriguez J.** 2013. Transcriptional data: a new
1048 gateway to drug repositioning? *Drug Discov Today* **18**:350-357.
- 1049 23. **Amorim-Vaz S, Tran Vdu T, Pradervand S, Pagni M, Coste AT, Sanglard D.** 2015. RNA
1050 Enrichment Method for Quantitative Transcriptional Analysis of Pathogens In Vivo Applied to
1051 the Fungus *Candida albicans*. *MBio* **6**:e00942-00915.
- 1052 24. **Grahl N, Cramer RA, Jr.** 2010. Regulation of hypoxia adaptation: an overlooked virulence
1053 attribute of pathogenic fungi? *Med Mycol* **48**:1-15.
- 1054 25. **Henry KW, Nickels JT, Edlind TD.** 2000. Upregulation of ERG genes in *Candida* species by
1055 azoles and other sterol biosynthesis inhibitors. *Antimicrob Agents Chemother* **44**:2693-2700.
- 1056 26. **Jensen-Pergakes KL, Kennedy MA, Lees ND, Barbuch R, Koegel C, Bard M.** 1998.
1057 Sequencing, disruption, and characterization of the *Candida albicans* sterol
1058 methyltransferase (ERG6) gene: drug susceptibility studies in *erg6* mutants. *Antimicrob*
1059 *Agents Chemother* **42**:1160-1167.
- 1060 27. **Lang GI, Parsons L, Gammie AE.** 2013. Mutation rates, spectra, and genome-wide
1061 distribution of spontaneous mutations in mismatch repair deficient yeast. *G3 (Bethesda)*
1062 **3**:1453-1465.
- 1063 28. **Ojini I, Gammie A.** 2015. Rapid Identification of Chemoresistance Mechanisms Using Yeast
1064 DNA Mismatch Repair Mutants. *G3 (Bethesda)* **5**:1925-1935.
- 1065 29. **Nes WD.** 2005. Enzyme redesign and interactions of substrate analogues with sterol
1066 methyltransferase to understand phytosterol diversity, reaction mechanism and the nature
1067 of the active site. *Biochem Soc Trans* **33**:1189-1196.
- 1068 30. **Mans R, van Rossum HM, Wijsman M, Backx A, Kuijpers NG, van den Broek M, Daran-**
1069 **Lapujade P, Pronk JT, van Maris AJ, Daran JM.** 2015. CRISPR/Cas9: a molecular Swiss army
1070 knife for simultaneous introduction of multiple genetic modifications in *Saccharomyces*
1071 *cerevisiae*. *FEMS Yeast Res* **15**.
- 1072 31. **Laabs TL, Markwardt DD, Slattery MG, Newcomb LL, Stillman DJ, Heideman W.** 2003. ACE2
1073 is required for daughter cell-specific G1 delay in *Saccharomyces cerevisiae*. *Proc Natl Acad*
1074 *Sci U S A* **100**:10275-10280.
- 1075 32. **Sanglard D, Ischer F, Monod M, Bille J.** 1996. Susceptibilities of *Candida albicans* multidrug
1076 transporter mutants to various antifungal agents and other metabolic inhibitors. *Antimicrob*
1077 *Agents Chemother* **40**:2300-2305.
- 1078 33. **Coste A, Selmecki A, Forche A, Diogo D, Bougnoux ME, d'Enfert C, Berman J, Sanglard D.**
1079 2007. Genotypic evolution of azole resistance mechanisms in sequential *Candida albicans*
1080 isolates. *Eukaryot Cell* **6**:1889-1904.
- 1081 34. **Cardenas PD, Sonawane PD, Heinig U, Bocobza SE, Burdman S, Aharoni A.** 2015. The bitter
1082 side of the nightshades: Genomics drives discovery in Solanaceae steroidal alkaloid
1083 metabolism. *Phytochemistry* **113**:24-32.
- 1084 35. **Roddick JG.** 1979. Complex-Formation between Solanaceous Steroidal Glycoalkaloids and
1085 Free Sterols In vitro. *Phytochemistry* **18**:1467-1470.
- 1086 36. **Friedman M.** 2002. Tomato glycoalkaloids: role in the plant and in the diet. *J Agric Food*
1087 *Chem* **50**:5751-5780.

- 1088 37. **Pierce AM, Unrau AM, Oehlschlager AC, Woods RA.** 1979. Azasterol inhibitors in yeast.
1089 Inhibition of the delta 24-sterol methyltransferase and the 24-methylene sterol delta 24(28)-
1090 reductase in sterol mutants of *Saccharomyces cerevisiae*. *Can J Biochem* **57**:201-208.
- 1091 38. **Sanglard D, Ischer F, Parkinson T, Falconer D, Bille J.** 2003. *Candida albicans* mutations in
1092 the ergosterol biosynthetic pathway and resistance to several antifungal agents. *Antimicrob*
1093 *Agents Chemother* **47**:2404-2412.
- 1094 39. **Lees ND, Skaggs B, Kirsch DR, Bard M.** 1995. Cloning of the late genes in the ergosterol
1095 biosynthetic pathway of *Saccharomyces cerevisiae*--a review. *Lipids* **30**:221-226.
- 1096 40. **Gaber RF, Copple DM, Kennedy BK, Vidal M, Bard M.** 1989. The yeast gene ERG6 is required
1097 for normal membrane function but is not essential for biosynthesis of the cell-cycle-sparking
1098 sterol. *Mol Cell Biol* **9**:3447-3456.
- 1099 41. **Clark DD, Peterson BR.** 2003. Analysis of protein tyrosine kinase inhibitors in recombinant
1100 yeast lacking the ERG6 gene. *Chembiochem* **4**:101-107.
- 1101 42. **Kelly M, MacCallum D, Clancy S, Odds F, Brown A, Butler G.** 1975. [Chronic heart diseases.
1102 6th diagnostic-therapeutic discourse of the ZFA in Freudstadt]. *Z Allgemeinmed* **51**:1330-
1103 1331.
- 1104 43. **Nes WD, Jayasimha P, Zhou W, Kanagasabai R, Jin C, Jaradat TT, Shaw RW, Bujnicki JM.**
1105 2004. Sterol methyltransferase: functional analysis of highly conserved residues by site-
1106 directed mutagenesis. *Biochemistry* **43**:569-576.
- 1107 44. **Robbins N, Spitzer M, Yu T, Cerone RP, Averette AK, Bahn YS, Heitman J, Sheppard DC,**
1108 **Tyers M, Wright GD.** 2015. An Antifungal Combination Matrix Identifies a Rich Pool of
1109 Adjuvant Molecules that Enhance Drug Activity against Diverse Fungal Pathogens. *Cell Rep*
1110 **13**:1481-1492.
- 1111 45. **Becker JM, Kauffman SJ, Hauser M, Huang L, Lin M, Sillaots S, Jiang B, Xu D, Roemer T.**
1112 2010. Pathway analysis of *Candida albicans* survival and virulence determinants in a murine
1113 infection model. *Proc Natl Acad Sci U S A* **107**:22044-22049.
- 1114 46. **Moazeni M, Kelidari HR, Saedi M, Morteza-Semnani K, Nabili M, Gohar AA, Akbari J,**
1115 **Lotfali E, Nokhodchi A.** 2016. Time to overcome fluconazole resistant *Candida* isolates: Solid
1116 lipid nanoparticles as a novel antifungal drug delivery system. *Colloids Surf B Biointerfaces*
1117 **142**:400-407.
- 1118 47. **Rodriguez-Tudela JL, Arendrup MC, Barchiesi F, Bille J, Chryssanthou E, Cuenca-Estrella M,**
1119 **Dannaoui E, Denning DW, Donnelly JP, Dromer F, Fegeler W, Lass-Flörl C, Moore C,**
1120 **Richardson M, Sandven P, Velegriaki A, Verweij P.** 2008. EUCAST Definitive Document EDef
1121 7.1: method for the determination of broth dilution MICs of antifungal agents for
1122 fermentative yeasts: Subcommittee on Antifungal Susceptibility Testing (AFST) of the
1123 ESCMID European Committee for Antimicrobial Susceptibility Testing (EUCAST)*. *Clinical*
1124 *Microbiology and Infection* **14**:398-405.
- 1125 48. **Meletiadiis J, Pournaras S, Roilides E, Walsh TJ.** 2010. Defining fractional inhibitory
1126 concentration index cutoffs for additive interactions based on self-drug additive
1127 combinations, Monte Carlo simulation analysis, and in vitro-in vivo correlation data for
1128 antifungal drug combinations against *Aspergillus fumigatus*. *Antimicrob Agents Chemother*
1129 **54**:602-609.
- 1130 49. **Pierce CG, Uppuluri P, Tummala S, Lopez-Ribot JL.** 2010. A 96 well microtiter plate-based
1131 method for monitoring formation and antifungal susceptibility testing of *Candida albicans*
1132 biofilms. *J Vis Exp* doi:10.3791/2287.
- 1133 50. **Vichai V, Kirtikara K.** 2006. Sulforhodamine B colorimetric assay for cytotoxicity screening.
1134 *Nat Protoc* **1**:1112-1116.
- 1135 51. **Guan XL, Riezman I, Wenk MR, Riezman H.** 2010. Yeast lipid analysis and quantification by
1136 mass spectrometry. *Methods Enzymol* **470**:369-391.

1137 52. **Sanglard D, Ischer F, Calabrese D, Majcherczyk PA, Bille J.** 1999. The ATP binding cassette
1138 transporter gene CgCDR1 from *Candida glabrata* is involved in the resistance of clinical
1139 isolates to azole antifungal agents. *Antimicrob Agents Chemother* **43**:2753-2765.

1140 53. **Robinson MD, McCarthy DJ, Smyth GK.** 2010. edgeR: a Bioconductor package for
1141 differential expression analysis of digital gene expression data. *Bioinformatics* **26**:139-140.

1142 54. **Law CW, Chen Y, Shi W, Smyth GK.** 2014. voom: Precision weights unlock linear model
1143 analysis tools for RNA-seq read counts. *Genome Biol* **15**:R29.

1144 55. **Schmittgen TD, Livak KJ.** 2008. Analyzing real-time PCR data by the comparative CT method.
1145 *Nat Protocols* **3**:1101-1108.

1146 56. **Reuss O, Vik A, Kolter R, Morschhauser J.** 2004. The SAT1 flipper, an optimized tool for gene
1147 disruption in *Candida albicans*. *Gene* **341**:119-127.

1148 57. **Delarze E, Ischer F, Sanglard D, Coste AT.** 2015. Adaptation of a *Gaussia princeps* Luciferase
1149 reporter system in *Candida albicans* for in vivo detection in the *Galleria mellonella* infection
1150 model. *Virulence* **6**:684-693.

1151 58. **Arthington-Skaggs B, Jradi H, Desai T, Morrison C.** 1999. Quantitation of ergosterol content:
1152 novel method for determination of fluconazole susceptibility of *Candida albicans*. . *J*
1153 *ClinMicrobiol* **37**:3332-3337.

1154 59. **Ryan OW, Poddar S, Cate JH.** 2016. CRISPR-Cas9 Genome Engineering in *Saccharomyces*
1155 *cerevisiae* Cells. *Cold Spring Harb Protoc* **2016**:pdb prot086827.

1156 60. **Vandeputte P, Ischer F, Sanglard D, Coste AT.** 2011. In vivo systematic analysis of *Candida*
1157 *albicans* Zn2-Cys6 transcription factors mutants for mice organ colonization. *PLoS One*
1158 **6**:e26962.

1159