

Identification of Novel Hybrids Between *Cryptococcus neoformans* var. *grubii* VNI and *Cryptococcus gattii* VGII

Mojgan Aminnejad · Mara Diaz · Michael Arabatzis · Elizabeth Castañeda · Marcia Lazera · Aristeia Velegraki · Deborah Marriott · Tania C. Sorrell · Wieland Meyer

Received: 4 July 2011 / Accepted: 3 October 2011 / Published online: 13 November 2011
© Springer Science+Business Media B.V. 2011

Abstract *Cryptococcus neoformans* and *Cryptococcus gattii* are pathogenic yeasts causing meningoencephalitis in immunocompromised and immunocompetent hosts. The fungus is typically haploid, and sexual reproduction occurs normally between individuals with opposite mating types, α and **a**. *C. neoformans* var. *grubii* (serotype A) is comprised of molecular types VNI, VNII, and VNB, and *C. neoformans* var. *neoformans* (serotype D) contains the molecular type VNIV. Additionally, diploid or aneuploid AD hybrids (VNIII) have been reported. *C. gattii* contains the molecular types VGI, VGII, VGIII, and VGIV, which encompass both serotypes B and C. To identify possible hybrid strains, *URA5*-RFLP analysis was performed on 350 globally obtained clinical, environmental, and

veterinary isolates. Four clinical isolates from cerebrospinal fluid showed combination patterns of *C. neoformans* var. *grubii* and *C. gattii*: Brazil ($n = 2$), Colombia ($n = 1$), and India ($n = 1$). These strains were monokaryotic and diploid or aneuploid. M13 PCR fingerprinting showed that they contained fragments of both proposed parental groups. Luminex IGS genotyping identified these isolates as hybrids with two different molecular type combinations: three VNI/VGII and one VNI/VGI. Blue color development on CGB agar was delayed in three isolates and absent in one. *C. gattii*-specific PCR confirmed the presence of *C. gattii* in the hybrids. *CAP59* allele-specific PCR revealed that all the hybrids contained both serotype A and B alleles. Determination of mating-type allelic

M. Aminnejad
Molecular Mycology Research Laboratory, Westmead,
Australia

M. Aminnejad · T. C. Sorrell · W. Meyer
Centre for Infectious Diseases and Microbiology, Sydney
Medical School-Westmead, Westmead Millennium
Institute, Sydney Emerging Infections and Biosecurity
Institute, University of Sydney, Sydney, NSW, Australia

M. Diaz
Rosenstiel School of Marine and Atmospheric Sciences,
University of Miami, Miami, USA

M. Arabatzis · A. Velegraki
Mycology Laboratory, Department of Microbiology,
Medical School, National and Kapodistrian University
of Athens, Athens, Greece

E. Castañeda
Grupo de Microbiología, Instituto Nacional de Salud,
Bogotá, Colombia

M. Lazera
Mycology Laboratory, Instituto de Pesquisa Clínica
Evandro Chagas, Fundação Oswaldo Cruz, Rio de Janeiro,
Brazil

D. Marriott
Department of Microbiology and Clinical Infectious
Diseases, St. Vincent's Hospital, Sydney, NSW, Australia

W. Meyer (✉)
Molecular Mycology Research Laboratory, CIDM,
ICPMR, Level 3, Room 3114A, Westmead Hospital,
Darcy Road, Westmead, NSW 2145, Australia
e-mail: w.meyer@usyd.edu.au

patterns by PCR revealed that the isolates were α A **aB**. This is the first study discovering novel natural hybrids between *C. neoformans* molecular type VNI and *C. gattii* molecular type VGII.

Keywords *Cryptococcus neoformans* · *Cryptococcus gattii* · AB hybrids · Serotype · Mating type · Molecular type

Introduction

Cryptococcus neoformans and *Cryptococcus gattii* are encapsulated basidiomycetous yeasts that cause meningoencephalitis. *C. neoformans* is an opportunistic pathogen, causing disease in immunocompromised patients [1]. *C. gattii* is considered as a primary pathogen, as it typically affects immunocompetent hosts [2]. *C. neoformans* has a worldwide distribution, and its major natural reservoir is pigeon droppings [3], soil contaminated with avian guano [4], and decaying wood of many tree species [5]. *C. gattii* is associated with decaying wood, plant debris, and soil [6, 7]; its distribution is limited, but includes tropical or subtropical areas [8] and certain temperate zones in Australia [9], Canada [7, 10], Greece [11], Italy [12], South Africa [13], Spain [14], and the North-West of the USA [15].

Cryptococcus is typically a haploid fungus with a bipolar heterothallic mating system containing one mating-type locus, which occurs in either of the two alleles, **a** or α . Sexual reproduction normally occurs between opposite mating types [16]. *C. neoformans* is comprised of two varieties: *C. neoformans* var. *grubii* (serotype A, molecular types VNI & VNII) and *C. neoformans* var. *neoformans* (serotype D, molecular type VNIV). In addition, diploid and aneuploid AD hybrids have been reported in clinical and environmental samples [1, 17]. Recently, based on a multilocus sequence typing (MLST) method, a new molecular type (VNB) has been identified among African and South American isolates of serotype A [18, 19]. *C. gattii* is comprised of serotypes B and C, and contains molecular types VGI, VGII, VGIII, and VGIV [17, 20]. Overall, serotype A isolates are the most common pathogens, being responsible for over 90% of cases of HIV-associated cryptococcosis worldwide [21–23].

In addition to AD hybrids, several inter- and intra-varietal diploid or aneuploid hybrids have been reported in the *C. neoformans*/*C. gattii* species complex [24–27]. Recently, three BD *C. neoformans* var. *neoformans* VNIV \times *C. gattii* VGI hybrids from two HIV-negative patients in the Netherlands and one AB *C. neoformans* var. *grubii* VNI \times *C. gattii* VGI hybrid from an HIV-positive patient in Quebec, Canada, have been described [24, 25]. Intra-varietal AA hybrids between VNII and VNB have also been reported [26, 27].

In the present study, a random selection of global clinical, environmental, and veterinary isolates maintained in the culture collection of the Molecular Mycology Research Laboratory at Westmead Hospital, Australia, was investigated to detect the presence of possible hybrids between the two species *C. neoformans* and *C. gattii* and characterize them in order to better understand the overall population structure of this important human/animal pathogen.

Materials and Methods

Clinical and Reference Strains

A random selection of 350 global *C. neoformans* and *C. gattii* isolates of clinical, environmental, and veterinary origins was obtained from the culture collection of Molecular Mycology Research Laboratory at Westmead Hospital, Australia, and screened for potential hybrid strains. Eight laboratory standard reference strains representing each molecular type, WM 148 (VNI, serotype A), WM 626 (VNII, serotype A), WM 628 (VNIII, serotype AD), WM 629 (VNIV, serotype D), WM 179 (VGI, serotype B), WM 178 (VGII), WM 175 (VGIII), and WM 779 (VGIV, serotype C) [17], were used for the identification of the major molecular types. The strains H99 (*MAT* α , serotype A) [28], IUM 96-2828 (*MAT***a**, serotype A) [29], JEC21 (*MAT* α , serotype D) [30], JEC20 (*MAT***a**, serotype D) [32], and WM 04.79 = LA599 (VGII *MAT***a**) [31] were used as controls for mating, serotype identification, luminex, and flow cytometry experiments.

Genomic DNA Extraction

Isolates were grown on Sabouraud's dextrose agar at 30°C for 48–72 h. Genomic DNA was then extracted according to Ferrer et al. [32]. Prior to performing

PCRs, the DNA was diluted in sterile deionized water to a concentration of 10 ng/μl.

URA5-RFLP Analysis

The *URA5* gene of all 350 isolates was amplified with primers *URA5*, 5'-ATGTCCTCCCAAGCCCTCGACTCCG-3' and *SJ101*, 5'-TTAAGACCTCTGAA CACCGTACTC-3', as previously described by Meyer et al. [17]. PCR products were digested with *HhaI* and *Sau96I* enzymes at 37°C over night. Digestion products were separated by 3% agarose gel electrophoresis and molecular types were identified by comparison of each profile with those of the eight standard reference strains [17]. Isolates with mixed patterns were considered to be potential hybrids.

Ploidy Determination by Fluorescence Flow Cytometry

Flow cytometry was performed on 10,000 cells of each potential hybrid isolate as described previously [33] and analyzed with a Becton–Dickinson FACSCallibur (BD Biosciences, San Jose, CA, USA). Cells stained with propidium iodide were examined by fluorescent microscopy to check the number of nuclei per cell.

Luminex Suspension Array

Oligonucleotide probes targeting either *C. neoformans* or *C. gattii*, specific oligonucleotide probes for serotype A and D, plus four probes for each of the molecular types of *C. gattii* were used as described by Diaz and Fell [34]. In addition, to separate VNI and VNII molecular type, two new specific probes for each VNI and VNII molecular types were designed based on sequence data from the IGS1 region of the ribosomal DNA, CNN 1a/1b: 5'-GGAATAAGGGA TGGATAAA-3' and CNN 1c: 5'-GGAATAAGG AATGGATAAA-3'. Luminex analysis was performed on potential hybrid isolates as described by Diaz and Fell [34], Diaz et al. [35], Fulton et al. [36] and Diaz and Fell [37]. Detection analysis was carried out on the Luminex 100 analyzer (Luminex Corporation, Austin, TX, USA), a flow cytometer with a dual laser system that allows simultaneous bead set recognition and quantification of targets based on the fluorescence of the reporter molecule. One hundred microspheres of each set were analyzed, which

represents 100 replicate measurements. Median fluorescent intensity (MFI) values were calculated with a digital signal processor and the Luminex proprietary software. A blank (without sample DNA) and a negative control (unrelated DNA) were included in the assay to ensure that sample contamination during the reaction will not result in a false-positive interpretation of the data. A positive signal was defined by a signal intensity at least twice the background or blank MFI value after subtraction of the background. The experiment was run twice.

M13 Fingerprinting

The minisatellite-specific core sequence of the wild-type phage M13 (5'-GAGGGTGGCGGTTCT-3') [38] was used as a single primer in the PCR, previously described by Meyer et al. [17]. Amplification products were separated by electrophoresis on 1.4% agarose gels, containing 0.5 μg/ml ethidium bromide in 1× Tris–borate-EDTA (TBE) buffer (89.1 mM Tris base, 88.9 mM Boric acid, 50 mM EDTA, pH 8.0) at 70 V and visualized under UV light. The 1-Kb DNA Ladder (Invitrogen, Carlsbad, USA) was used as a molecular size marker. WM 148 (VNI), WM 626 (VNII), WM 179 (VGI), and WM 178 (VGII) were used as reference strains.

Growth on CGB Medium

Hybrid isolates were grown at 27°C on canavanine-glycine-bromothymol blue agar (CGB) medium, which distinguishes between *C. gattii* (cobalt blue) and *C. neoformans* (yellow) growth, due to a pH-dependent color change [39]. The color of CGB medium was assessed after 5 and 15 days incubation, since it was predicted that the blue color development would be delayed in hybrid strains. Strains WM 148 (VNI) and WM 179 (VGI) were used as controls.

C. gattii-Specific PCR

To confirm the presence of *C. gattii* in the putative hybrids, *C. gattii*-specific PCR was performed using the 660 U and 660 L *C. gattii*-specific primers, which are based on an anonymous DNA fragment (216 bp) that was amplified by randomly amplified polymorphic DNA (RAPD) analysis from a *C. gattii* strain [40]. 10 μl of each amplification product was electrophoresed at

80 V on 2% agarose gel in 1× TBE buffer containing 0.5 µg/ml ethidium bromide and visualized under UV light. The 1-Kb Plus DNA Ladder (Invitrogen, Carlsbad, USA) was used as a molecular size marker. WM 148 (VNI) and WM 179 (VGI) were used as control strains.

CAP59 Allele-Specific PCR

To identify the serotype-specific allele combination in the potential hybrids, the *CAP59* gene was amplified with primers CH-Cap 59F and CH-Cap 59R, as described previously [41]. Amplicons were subjected to analysis using three separate restriction enzymes—*BsmFI*, *HpaII*, and *AgeI*—according to the instructions of the manufacturer (New England Biolabs, Ipswich, MA, USA). Digested fragments were separated and visualized on a 3% agarose gel stained with ethidium bromide. Enzymatic restriction with *BsmFI* and *HpaII* distinguishes between the A, D, and B/C alleles. Serotype AD strains exhibit a mixture of A and D restriction profiles. The B and C alleles can be differentiated by *AgeI* digestion [41].

GPA1 and PAK1 Allele-Specific PCR

The presence or absence of serotype A- and D-specific alleles of the *GPA1* gene was determined by PCR using the primer sets JOHE2596/JOHE3241 specific for serotype A allele and JOHE2596/JOHE3240 specific for serotype D allele [42], and for *PAK1* gene it was determined using the primer sets JOHE3066/JOHE3236 specific for serotype A allele and JOHE3066/JOHE3065 specific for serotype D allele [43], as described previously. The following cryptococcal strains were used as control strains for different serotypes: H99 (serotype A), JEC21 (serotype D), WM 179 (serotype B), and WM 779 (serotype C).

Determination of Mating-Type Allelic Partners

To determine the mating-type allelic partners of the potential hybrid strains, the *MF α a* locus was amplified with the primer sets MF α F/MF α R and MFaF/MFaR [44], the *STE12 α a* locus was amplified with the primer sets MF α U/MF α L [40] and JOHE9779/JOHE9780 [10], the *STE20 α a* locus was amplified with the primer set JOHE9421/JOHE9422 [10], the *A α* allele of the *STE20 α a* locus was amplified with the

primer set JOHE7264/JOHE7266 [43], the *D α* allele of the *STE20 α a* locus was amplified with the primer set JOHE7267/JOHE7269 [43], the *Aa* allele of the *STE20 α a* locus was amplified with the primer set JOHE7270/JOHE7271 [43], the *Da* allele of the *STE20 α a* locus was amplified with the primer set JOHE7273/JOHE7274 [43], the *A α* allele of the *SXII α 2a* locus was amplified with the primer set JOHE15634/JOHE15635 [43], and the *D α* allele of the *SXII α 2a* locus was amplified with the primer set JOHE15634/JOHE15636 [43], as previously published. The following cryptococcal strains were used as control strains for the two mating types: H99 (*A α*), IUM 96-2828 (*Aa*), JEC21 (*D α*), JEC20 (*Da*), WM 179 (*B α*), and WM 04.79 (*VGII MATa*).

Results

Mixed patterns of both *C. neoformans* var. *grubii* and *C. gattii* were observed in cerebrospinal fluid isolates from four patients with cryptococcal meningitis. Comparison of the *URA5*-RFLP profiles of all 350 isolates with those obtained from the standard reference strains revealed mixed patterns of VNI/VGI for one isolate (WM 2617) from a patient in India and VNI/VGII for three isolates (WM 05.272, WM 05.459, and WM 05.532) from patients in Colombia and Brazil (Fig. 1a; Table 1).

Determination of ploidy using flow cytometry analysis showed that all potential hybrid isolates contained ~2× the amount of DNA in the haploid control strain, confirming that they were diploids or aneuploids (Fig. 2). All cells were uninucleate.

The putative hybrids were analyzed using Luminex xMAP technology, a flow cytometer that allows the simultaneous identification of cryptococcal varieties and their genotypes by mixing different sets of microspheres containing specific capture probes derived from IGS target sequences. The *C. neoformans* probe (CNN b), *C. neoformans* var. *grubii* probe (CNN 1b), *C. gattii*-specific probe (CNG), and the *C. neoformans* var. *grubii* VNI probe (CNN 1a/1b) gave a positive signal for all four *C. neoformans*/*C. gattii* hybrid isolates. Additionally, for WM 05.272, WM 05.459 and WM 05.532 positive signals were observed from VGII-specific probe (CNG 3), and for WM 2617, the probe specific for VGI (CNG 4c) gave positive signals (data not shown). Therefore, this technique




Fig. 1 **a** RFLP profiles of the putative hybrid isolates as well as standard strains for all molecular types obtained via double digest of the *URA5* gene with *HhaI* and *Sau96I*. Boxes indicate fragments occurring both in hybrid isolates and their parental groups. Lane M, 1-kb⁺ DNA ladder (Invitrogen, Carlsbad, USA). **b** M13 PCR fingerprint of the putative hybrid isolates and the standard strains for each of the putative parental groups. Boxes indicate fragments occurring both in hybrid isolates and their parental groups. Lane M', 1-kb DNA ladder (Invitrogen, Carlsbad, USA)

identified the isolates as hybrids with two different molecular type combinations, VNI/VGI for WM 2617 and VNI/VGII for WM 05.272, WM 05.459, and WM 05.532.

Comparisons of the M13 PCR fingerprints of the putative hybrids with the standard strains of the major molecular types showed that the putative hybrid

isolates contained fragments of both proposed parental groups (VNI and VGI for WM 2617 and VNI, and VGII for WM 05.272, WM 05.459, and WM 05.532) as determined by *URA5*-RFLP (Fig. 1 b; Table 1).

After 5 days of incubation, the positive control, *C. gattii* (WM 178), produced a cobalt blue color on CGB medium, whereas the negative control, *C. neoformans* (WM 148), remained yellow (no growth). It is of interest that the three AB hybrids of VNI and VGII produced an intermediate greenish-blue color at day 5, which became blue on day 15, whereas the VNI/VGI hybrid (WM 2617) remained yellow on CGB medium. All four putative hybrids containing serotype B were confirmed by *C. gattii*-specific PCR, giving a positive amplification (data not shown).



Restriction enzyme analysis of the *CAP59* gene with *BsmFI*, *HpaII*, and *AgeI* showed that all hybrids had both serotype A- and B-specific alleles (Fig. 3). Positive amplifications were obtained for all hybrid isolates with the *MATα* serotype A allele primer sets specific for *STE20α* (JOHE7264/LOHE7266) and *SXI1α* (JOHE15634/JOHE15635), and the serotype A allele primer sets specific for *GPA1* (JOHE2596/JOHE3241) and *PAK1* (JOHE3066/JOHE3236). No amplification was obtained with the *MATa* serotype A allele primer sets specific for *STE20a* (JOHE7270/JOHE7271). Similar negative results were obtained when using the primer sets for *SXI2a* (JOHE15629/JOHE15630), and serotype D allele primer sets for *GPA1* (JOHE2596/JOHE3240) and *PAK1* (JOHE3066/JOHE3065). In addition, the hybrid isolates yielded amplicons with the primer sets specific for *MFα* (MFαF/MFαR and MFαU/MFαL) and *STE12α* (STE12αF809/STE12αR1607) in *MATα* and *STE20a* (JOHE9421/JOHE9422) in *MATa*. When the STE12aF537/STE12aR1299 primers were used, amplicons were obtained for *MATa* in isolates WM 2617 and WM 05.272. The primer set JOHE9787/JOHE9788 amplified *MATa* of WM 05.272, whereas MFαF/MFαR primers failed to amplify the hybrid isolates. All the reference strains gave amplicons with the expected primer pairs (Table 2). These results indicate that the hybrid isolates had AαBα combination.

Discussion

Although *C. neoformans* and *C. gattii* are predominantly haploid organisms, naturally occurring hybrids

Table 1 Hybrid strains identified in this study

Strain #	Origin	Source	Date of isolation	Underlying disease	Site of sample	Molecular type	Mating/serotype
WM 2617	Vellore, Tamil Nadu, India	Clinical	8/09/1995	HIV+	CSF	VNI/VGI	α ABa
WM 05.272	Cúcuta, Colombia	Clinical	2003	HIV–	CSF	VNI/VGII	α ABa
WM 05.459	Castelo do Piauí, Piauí, Brazil	Clinical	12/1996	HIV–	CSF	VNI/VGII	α ABa
WM 05.532	Caxias, Maranhão, Brazil	Clinical	03/2001	HIV–	CSF	VNI/VGII	α ABa

Fig. 2 Flow cytometry profiles of haploid control strain (a) and a putative hybrid isolate (b) after staining with the fluorescent dye propidium iodide. *1n*, *2n*, and *4n* indicate nuclear content. The x-axis indicates fluorescence intensity reflecting the DNA content, and the y-axis indicates cell counts**Fig. 3** Restriction profiles of *CAP59* gene for hybrid isolates and reference strains of different serotypes of *C. neoformans* and *C. gattii* on 3% agarose gel with *Bsm*FI, *Hpa*II, and *Age*I restriction enzymes. Lane M, 1-kb⁺ DNA ladder (Invitrogen, Carlsbad, USA)

of AD (inter-varietal), AA (intra-varietal), and BD and AB (inter-species hybrids) that are virulent (i.e., cause mammalian disease) have been reported [1, 24, 25, 27].

In the current study, four AB hybrid strains from patients with cryptococcal meningitis were identified and characterized. By fluorescence-activated cell sorter (FACS) analysis, these strains were diploid or

aneuploid. PCR analysis revealed that these strains were heterozygous for serotype A- and B-specific alleles and the *MAT α* and *MATa* mating-type loci. Based on these observations, these hybrids could either be produced by somatic fusion followed by karyogamy of a *MAT α* serotype A cell with a *MATa* of a serotype B cell, or by mating of a *MAT α* serotype A

Table 2 Serotype- and mating-type-specific PCR analysis results for the identified hybrid isolates in comparison with reference strains

Trait	Gene	Isolate/strain									
		WM 2617	WM 05.272	WM 05.459	WM 05.532	H99 (A α)	IUM 96-2828 (Aa)	JEC21 (D α)	JEC20 (Da)	WM 179 (B α)	WM 04.79 (VG a)
Genes in the <i>MAT</i> locus	<i>STE20Aα</i>	+	+	+	+	+	–	–	–	ND	ND
	<i>SXIIAα</i>	+	+	+	+	+	–	–	–	ND	ND
	<i>STE20Dα</i>	–	–	–	–	–	–	+	–	ND	ND
	<i>SXII Dα</i>	–	–	–	–	–	–	+	–	ND	ND
	<i>STE20Aa</i>	–	–	–	–	–	+	–	–	ND	ND
	<i>SXI2Aa</i>	–	–	–	–	–	+	–	–	ND	ND
	<i>STE20Da</i>	–	–	–	–	–	–	–	+	ND	ND
	<i>SXI2Da</i>	–	–	–	–	–	–	–	+	ND	ND
	<i>MFα</i>	+	+	+	+	+	–	+	–	+	–
	<i>MFa</i>	–	+	–	–	–	+	–	+	–	+
	<i>STE12α</i>	+	+	+	+	+	–	ND	ND	ND	ND
	<i>STE12a</i>	+	+	–	–	–	+	ND	ND	ND	ND
	<i>STE20a</i>	+	+	+	+	ND	ND	ND	ND	–	+
	Other genes	<i>GPAI-A</i>	+	+	+	+	+	+	–	–	ND
<i>PAKI-A</i>		+	+	+	+	+	+	–	–	ND	ND
<i>GPAI-D</i>		–	–	–	–	–	–	+	+	ND	ND
<i>PAKI-D</i>		–	–	–	–	–	–	+	+	ND	ND

ND not done

strain with a *MATa* serotype B strain, to produce diploid basidiospores. Different analyses revealed that the strain WM 2617 from India portrayed genetic traits of *C. neoformans* var. *grubii* (molecular type VNI) and *C. gattii* (molecular type VGI), whereas the other three strains (WM 05.272, WM 05.459, and WM 05.532) from South America consisted of a novel genetic hybrid of *C. neoformans* var. *grubii* (molecular type VNI) and *C. gattii* (molecular type VGII). To our knowledge, this is the first report in the literature documenting the occurrence of novel natural hybrids between *C. neoformans* molecular type VNI and *C. gattii* molecular type VGII. It is notable that the strain WM 2617 reacts negative on CGB medium. Negative, weak, and delayed positive reactions on CGB medium have also been shown for other *C. neoformans* \times *C. gattii* hybrids [24, 25]. Negative reaction on CGB media may be due to inhibitory effect by canavanine, which is a dominant trait. As such routine screening of isolates on CGB medium may miss those hybrids.

The lack of amplification of the *MATa* locus with primer pairs: STE12aF537/STE12aR1299 and JOHE9787/JOHE9788 in some, and of MFaF/MFaR

primers in all reported hybrids could be due to a loss of the *MATa* allele in some genes of the *C. gattii* parent mating locus during the meiotic process. Alternatively, the primers may not have been specific, due to diverging sequences in the mating-type locus, which prevented annealing of the primers, and hence hindered the amplification, or these isolates were homozygous for one or more mating loci. Loss of genetic material has been observed in AD, BD, and other AB hybrids [24, 25, 45]. Even if hybrid strains possess two genome copies, it is likely that most of the duplicated genes would eventually be lost. However, the presence of two copies of one gene provides an opportunity to evolve new functions, as has been shown for *Saccharomyces cerevisiae* [46].

Development of polyploidy enables tolerance to genetic alteration and provides an additional mechanism for generating genetic diversity. It is a driving force in the evolution of many eukaryotes, and populations with both haploids and diploids may afford greater adaptability than those limited to only one, due to the shuttling between diploid and haploid

states [27]. Since *C. neoformans* and *C. gattii* are environmental pathogens that live in a wide range of ecological habitats, it has been suggested that they have evolved more diverse genotypes and phenotypes to survive in these different niches [47]. Formation of hybrids within *C. neoformans* and *C. gattii* populations might generate genotypic and phenotypic diversity and lineages that can occupy new ecological niches. In this case, hybridization would be advantageous for the fungus. However, it can also be a disadvantage, since some interspecies hybrids exhibit reduced fitness [48]. It has also been observed that diploid strains of *C. neoformans* can convert to a haploid (normal) state, suggesting that polyploidy may be associated with genetic instability [27].

The novelty of the work described herein is that the *C. gattii* parent of three of the four AB hybrids belongs to the VGII molecular type, which has not been reported previously. The *C. gattii* parent of previously reported BD and AB hybrids was of the molecular type VGI [24, 25], which is the same as the hybrid isolate WM 2617 reported in this study. The fact that we found similar hybrids in different countries suggests that either the inter-species mating process is not restricted to a specific geographic area and has independent, non-clonal origins, or that hybrid strains expanded clonally and were dispersed to diverse geographic regions.

It is notable that most hybrids to date have been clinical strains, though the implication of this observation with respect to disease pathogenesis, virulence, and the spread of infection, is not known. Further investigations to determine the relative virulence of these strains are underway.

Acknowledgments The authors thank Mary Matthews for providing the strain from India; Maggie Wang and Stephen Schibeci for technical assistance on flow cytometry analysis; and Fottes Al. Panetsos and Hariklia Moshoyiannis for technical assistance on Luminex analysis. This work was supported by a University of Sydney Postgraduate Award and a CIDM Public health scholarship to MAm. The work of the Sydney Institute for Emerging Infections and Biosecurity is supported by a grant from the Sydney Medical School Foundation. TCS is a Sydney Medical School Foundation Fellow. The Bodosakis Foundation and the Research fund SARG 70/3/6915 National and Kapodistrian University of Athens partially funded this work. Flow cytometry was performed in the Flow Cytometry Core Facility that is supported by Westmead Millennium Institute, National Health and Medical Research Council and Cancer Institute New South Wales, Australia.

References

1. Kwon-Chung KJ, Bennett JE. Epidemiologic differences between the two varieties of *Cryptococcus neoformans*. *Am J Epidemiol.* 1984;120:123–30.
2. Speed B, Dunt D. Clinical and host differences between infections with the two varieties of *Cryptococcus neoformans*. *Clin Infect Dis.* 1995;21:28–34.
3. Emmons CW. Prevalence of *Cryptococcus neoformans* in pigeon habitats. *Public Health Rep.* 1960;75:362–4.
4. Cafarchia C, Rommito D, Latta R, Camarda A, Montagna MT, Otranto D. Role of birds of prey as carriers and spreaders of *Cryptococcus neoformans* and other zoonotic yeasts. *Med Mycol.* 2006;44:458–92.
5. Randhawa HS, Kowshik T, Chowdhary A, Preeti SK, Khan ZU, Sun S, Xu J. The expanding host tree species spectrum of *Cryptococcus gattii* and *Cryptococcus neoformans* and their isolations from surrounding soil in India. *Med Mycol.* 2008;46:823–33.
6. Mitchell TG, Castañeda E, Nielson K, Wanke B, Lazera MS. Environmental niches for *Cryptococcus neoformans* and *Cryptococcus gattii*. In: Heitman J, Kozel TR, Kwon-Chung KJ, Perfect JR, Casadevall A, editors. *Cryptococcus: from human pathogen to model yeast*. Washington: ASM Press; 2011. p. 237–59.
7. Kidd SE, Chow Y, Mak S, Bach PJ, Chen H, Hingston AO, Kronstad JW, Bartlett KH. Characterization of environmental sources of the human and animal pathogen, *Cryptococcus gattii*, in British Columbia, Canada, and the Pacific Northwest of the United States. *Appl Environ Microbiol.* 2007;73:1433–43.
8. Kwon-Chung KJ, Bennett JE. High prevalence of *Cryptococcus neoformans* var. *gattii* in tropical and subtropical regions. *Zentralbl Bakteriol Mikrobiol Hyg.* 1984;257: 213–8.
9. Ellis DH. *Cryptococcus neoformans* var. *gattii* in Australia. *J Clin Microbiol.* 1987;25:430–1.
10. Fraser JA, Subaran RL, Nicholas CB, Heitman J. Recapitulation of the sexual cycle of the primary fungal pathogen *Cryptococcus neoformans* var. *gattii*: implications for an outbreak on Vancouver Island, Canada. *Eukaryot Cell.* 2003;2:1036–45.
11. Velegraki A, Kiosses VG, Pitsouni H, Toukas D, Daniilidis VD, Legakis NJ. First report of *Cryptococcus neoformans* var. *gattii* serotype B from Greece. *Med Mycol.* 2001; 39:419–22.
12. Romeo O, Scordino F, Criseo G. Environmental isolation of *Cryptococcus gattii* serotype B, VGI/MAT α strains in southern Italy. *Mycopathologia.* 2011;171:423–30.
13. Karstaedt AS, Crewe-Brown HH, Dromer F. Cryptococcal meningitis caused by *Cryptococcus neoformans* var. *gattii*, serotype C, in AIDS patients in Soweto, South Africa. *Med Mycol.* 2002;40:7–11.
14. Colom MF, Frasés S, Ferrer C, Jover A, Andreu M, Reus S, Sánchez M, Torres-Rodríguez JM. First case of human cryptococcosis due to *Cryptococcus neoformans* var. *gattii* in Spain. *J Clin Microbiol.* 2005;43:3548–50.
15. Byrnes EJ III, Li W, Lewit Y, Ma H, Voelz K, Ren P, Carter DA, Chaturvedi V, Bildfell RJ, May RC, Heitman J. Emergence and pathogenicity of highly virulent

- Cryptococcus gattii* genotypes in the Northwest United States. PLoS Pathog. 2010;6(4):e1000850. doi:10.1371/journal.ppat.1000850.
16. Lin X, Heitman J. The biology of the *Cryptococcus neoformans* species complex. Annu Rev Microbiol. 2006;60:69–105.
 17. Meyer W, Castaneda A, Jackson S, Huynh M, Castaneda E. The Ibero American Cryptococcal Study Group. Molecular typing of IberoAmerican *Cryptococcus neoformans* isolates. Emerg Infect Dis. 2003;9:189–95.
 18. Litvintseva AP, Thakur R, Vilgalys RJ, Mitchell T. Multilocus sequence typing reveals three genetic subpopulations of *Cryptococcus neoformans* var. *grubii* (serotype A), including a unique population in Botswana. Genetics. 2006;172:2223–38.
 19. Ngamskulrungrroj P, Gilgado F, Faganello J, Litvintseva AP, Leal AL, Tsui KM, Mitchell TG, Vainstein MH, Meyer W. Genetic diversity of the *Cryptococcus* species complex suggests that *Cryptococcus gattii* deserves to have varieties. PLoS One. 2009;4:e5862.
 20. Kwon-Chung KJ, Boekhout T, Fell JW, Diaz M. Proposal to conserve the name *Cryptococcus gattii* against *C. honduricus*, *C. bacillisporus* (Basidiomycota, Hymenozetes, Tremellomycetidae). Taxon. 2002;51:804–6.
 21. Currie BP, Casadevall A. Estimation of the prevalence of Cryptococcal infection among patients infected with the human immunodeficiency virus in New York City. Clin Infect Dis. 1994;19:1029–33.
 22. Franzot SP, Hamdan JS, Currie BP, Casadevall A. Molecular epidemiology of *Cryptococcus neoformans* in Brazil and the United States: evidence for both local genetic differences and a global clonal population structure. J Clin Microbiol. 1997;35:2243–51.
 23. Viviani MA, Cogliati M, Esposto MC, Lemmer K, Tintelnot K, Colom Valiente MF, Swinne D, Velegriaki A, Velho R. Molecular analysis of 311 *Cryptococcus neoformans* isolates from a 30-month ECMM survey of cryptococcosis in Europe. FEMS Yeast Res. 2006;6:614–9.
 24. Bovers M, Hagen F, Kuramae EE, Diaz MR, Spanjaard L, Dromer F, Hoogveld HL, Boekhout T. Unique hybrids between the fungal pathogens *Cryptococcus neoformans* and *Cryptococcus gattii*. FEMS Yeast Res. 2006;6:599–607.
 25. Bovers M, Hagen F, Kuramae EE, Hoogveld HL, Dromer F, St-Germain G, Boekhout T. AIDS patient death caused by novel *Cryptococcus neoformans* × *C. gattii* hybrid. Emerg Infect Dis. 2008;14:1105–8.
 26. Bovers M, Hagen F, Kuramae EE, Boekhout T. Six monophyletic lineages identified within *Cryptococcus neoformans*, *C. gattii* by multilocus sequence typing. Fungal Genet Biol. 2008;45:400–21.
 27. Lin X, Patel S, Litvintseva AP, Floyd A, Mitchell TG, Heitman J. Diploids in the *Cryptococcus neoformans* serotype A population homozygous for the α mating type originate via unisexual mating. PLoS Pathog. 2009;5:e1000283.
 28. Franzot SP, Salkin IR, Casadevall A. *Cryptococcus neoformans* var. *grubii*: separate varietal status for *Cryptococcus neoformans* serotype A isolates. J Clin Microbiol. 1999;37:838–40.
 29. Viviani MA, Esposto MC, Cogliati M, Montagna MT, Wickes BL. Isolation of a *Cryptococcus neoformans* serotype A MATa strain from the Italian environment. Med Mycol. 2001;39:383–6.
 30. Kwon-Chung KJ, Edman JC, Wickes BL. Genetic association of mating types and virulence in *Cryptococcus neoformans*. Infect Immun. 1992;60:602–5.
 31. Ngamskulrungrroj P, Sorrell TC, Chindamporn A, Chaiprasert A, Poonwan N, Meyer W. Association between fertility and molecular sub-type of global isolates of *Cryptococcus gattii* molecular type VGII. Med Mycol. 2008;46:665–73.
 32. Ferrer C, Colom F, Frasés S, Mulet E, Abad JL, Alió JL. Detection and identification of fungal pathogens by PCR and by ITS2 and 5.8S ribosomal DNA typing in ocular infections. J Clin Microbiol. 2001;39:2873–9.
 33. Dien BS, Peterson MS, Srien F. Cell cycle analysis in *Saccharomyces cerevisiae*. In: Darzynkiewicz Z, Robinson JP, Crissman HA, editors. Methods in cell biology, vol. 42. USA: Academic Press; 1994. p. 457–75.
 34. Diaz MR, Fell JW. Use of a suspension array for rapid identification of the varieties and genotypes of the *Cryptococcus neoformans* species complex. J Clin Microbiol. 2005;43:3662–72.
 35. Diaz MR, Dunbar SA, Jacobson JW. Multiplexed detection of fungal nucleic acid signatures. In: Robinson JP, Darzynkiewicz Z, Hoffman R, Nolan JP, Rabinovitch PS, Watkins S, editors. Current protocols in cytometry. New York: Wiley; 2008. 44:13.9.1–13.9.21. doi:10.1002/0471142956.cy1309s44.
 36. Fulton R, McDade R, Smith P, Kienker L, Kettman J. Advanced multiplexed analysis with the Flowmetrix system. Clin Chem. 1997;43:1749–56.
 37. Diaz MR, Fell JW. High-throughput detection of pathogenic yeasts of the genus *Trichosporon*. J Clin Microbiol. 2004;42:3696–706.
 38. Vassart G, Georges M, Monsieur R, Brocas H, Lequarre AS, Christophe D. A sequence in M13 phage detects hyper-variable minisatellites in human and animal DNA. Science. 1987;235:683–4.
 39. Kwon-Chung KJ, Polacheck I, Bennett JE. Improved diagnostic medium for separation of *Cryptococcus neoformans* var. *neoformans* (serotypes A and D) and *Cryptococcus neoformans* var. *gattii* (serotypes B and C). J Clin Microbiol. 1982;15:535–7.
 40. Halliday CL, Bui T, Krockenberger M, Malik R, Ellis DH, Carter DA. Presence of α and β mating type in environmental and clinical collections of *Cryptococcus neoformans* var. *gattii* strains from Australia. J Clin Microbiol. 1999;37:2920–6.
 41. Enache-Angoulvant A, Chandener J, Symoens F, Lacube P, Bolognini J, Douchet C, Poirot JL, Hennequin C. Molecular identification of *Cryptococcus neoformans* serotypes. J Clin Microbiol. 2007;45:1261–5.
 42. Lengeler KB, Cox GM, Heitman J. Serotype AD strains of *Cryptococcus neoformans* are diploid or aneuploid and are heterozygous at the mating-type locus. Infect Immun. 2001;69(1):115–22.
 43. Lin X, Litvintseva AP, Nielsen K, Patel S, Floyd A, Mitchell TG, Heitman J. α AD α hybrids of *Cryptococcus neoformans*: evidence of same-sex mating in nature and hybrid fitness. PLoS Genet. 2007;3:e186.
 44. Chaturvedi S, Rodeghier B, Fan J, McClelland CM, Wickes BL, Chaturvedi V. Direct PCR of *Cryptococcus neoformans*

- MAT α* and *MATa* pheromones to determine mating type, ploidy, and variety: a tool for epidemiological and molecular pathogenesis studies. *J Clin Microbiol.* 2000;38:2007–9.
45. Kwon-Chung KJ, Varma A. Do major species concepts support one, two or more species within *Cryptococcus neoformans*? *FEMS Yeast Res.* 2006;6:574–87. doi: [10.1111/j.1567-1364.2006.00088.x](https://doi.org/10.1111/j.1567-1364.2006.00088.x).
46. Kellis M, Birren BW, Lander ES. Proof and evolutionary analysis of ancient genome duplication in the yeast *Saccharomyces cerevisiae*. *Nature.* 2004;428:617–24.
47. Hunter PR, Fraser CA. Application of the theory of adaptive polymorphism to the ecology and epidemiology of pathogenic yeasts. *Appl Environ Microbiol.* 1990;56:2219–22.
48. Seehausen O. Hybridization and adaptive radiation. *Trends Ecol Evol.* 2004;19:198–207.