

# Identification of phenolic compounds that suppress the virulence of *Xanthomonas oryzae* on rice via the type III secretion system

SUSU FAN<sup>1</sup>, FANG TIAN<sup>1,\*</sup>, JIANYU LI<sup>1</sup>, WILLIAM HUTCHINS<sup>2</sup>, HUAMIN CHEN<sup>1</sup>, FENGHUAN YANG<sup>1</sup>, XIAOCHEN YUAN<sup>2</sup>, ZINING CUI<sup>3</sup>, CHING-HONG YANG<sup>2,\*</sup> AND CHENYANG HE<sup>1</sup>

<sup>1</sup>State Key Laboratory for Biology of Plant Diseases and Insect Pests, Institute of Plant Protection, Chinese Academy of Agricultural Sciences, Beijing 100193, China

<sup>2</sup>Department of Biological Sciences, University of Wisconsin-Milwaukee, Milwaukee, WI 53211, USA

<sup>3</sup>Guangdong Province Key Laboratory of Microbial Signals and Disease Control, Department of Plant Pathology, South China Agricultural University, Guangzhou 510642, China

## SUMMARY

The targeting of bacterial type III secretion systems (T3SSs), which are critical virulence factors in most Gram-negative pathogens, is regarded as an alternative strategy for the development of novel anti-microbial drugs. *Xanthomonas oryzae* pv. *oryzae* (*Xoo*) and *X. oryzae* pv. *oryzicola* (*Xoc*) are two of the most important bacterial pathogens on rice, which cause leaf blight and leaf streak diseases, respectively. To identify potential anti-virulence drugs against these two pathogens, we screened a library of plant phenolic compounds and derivatives for their effects on the *Xoo* T3SS. Ten of 56 compounds significantly inhibited the promoter activity of a harpin gene, *hpa1*. These inhibitors were further tested for their impact on the hypersensitive response (HR) caused by *Xoo* on non-host tobacco plants. The results showed that pretreatment of *Xoo* with TS006 (*o*-coumaric acid, OCA), TS010, TS015 and TS018 resulted in significantly attenuated HR without affecting bacterial growth or survival. In addition, Cya translocation assays demonstrated that the translocation of two T3 effectors was suppressed by the four inhibitors. Quantitative reverse transcription-polymerase chain reaction (qRT-PCR) analysis showed that mRNA levels of representative genes in the *hrp* (*hypersensitive response and pathogenicity*) cluster, as well as the regulatory genes *hrpG* and *hrpX*, were reduced by treatment with the four inhibitors, suggesting that expression of the *Xoo* T3SS was suppressed. The expression of other virulence factors was not suppressed, which indicated possible T3SS-specific inhibition. Finally, we demonstrated that these inhibitors reduced the disease symptoms of *Xoo* and *Xoc* on the rice cultivar (*Oryza sativa*) IR24 to varying extents.

**Keywords:** bacterial leaf blight, bacterial leaf streak, harpin, HrpG, HrpX, hypersensitive response, *o*-coumaric acid.

## INTRODUCTION

Traditional antibiotic therapy, which aims to affect the processes essential for bacterial growth and survival, has led to strong selective pressure to develop antibiotic resistance in pathogenic bacteria (Rasko and Sperandio, 2010). An alternative approach is to identify new agents that target bacterial virulence factors, rather than their growth (Barczak and Hung, 2009; Rasko and Sperandio, 2010). The type III secretion system (T3SS) is a highly conserved virulence factor in many Gram-negative pathogenic bacteria, and is not necessary for bacterial survival *in vitro* (Büttner, 2012; Cornelis, 2006). Thus, it is regarded as an ideal target for the development of novel antimicrobial drugs (Charro and Mota, 2015; Marshall and Finlay, 2014). So far, several different classes of small-molecule compounds have been identified as active T3SS inhibitors in a wide range of pathogenic bacteria, including *Escherichia*, *Yersinia*, *Salmonella* and *Erwinia* species (Felise *et al.*, 2008; Jessen *et al.*, 2014; Wang *et al.*, 2011; Yang *et al.*, 2014). These inhibitors function by directly targeting the components of the T3SS apparatus (Bowlin *et al.*, 2014; Jessen *et al.*, 2014), or regulating T3SS gene expression (Garrity-Ryan *et al.*, 2010; Yang *et al.*, 2014), or through more indirect interactions (Bowlin *et al.*, 2014; Wang *et al.*, 2011).

Phenolic compounds are ubiquitous secondary metabolites in plants, which are crucial in many aspects of plant life, especially during their interactions with the environment (Lattanzio, 2013). Some phenolics play key roles in plant defence responses to pathogen or insect attacks. For example, salicylic acid (SA), a simple phenolic compound, is the signalling molecule required for the induction of systemic acquired resistance in plants (Durrant and Dong, 2004). Recently, it has been discovered that the novel phenolic compounds *o*-coumaric acid (OCA), *trans*-cinnamic acid (TCA) and *p*-coumaric acid (PCA) modulate the expression of T3SS genes in *Dickeya dadantii*, a broad-host-range phytopathogen (Li *et al.*, 2009; Yang *et al.*, 2008). Some of their derivatives have been found to suppress T3SS gene expression in *Pseudomonas aeruginosa* and *Erwinia amylovora* (Khokhani *et al.*, 2013; Yamazaki *et al.*, 2012).

\*Correspondence: Email: ftian@ippcaas.cn; chyang@uwm.edu

*Xanthomonas oryzae* pv. *oryzae* (*Xoo*) is the causal agent of bacterial blight (BB), one of the most serious diseases of rice worldwide (Nino-Liu *et al.*, 2006). *Xoo* typically invades rice vascular tissue through wounds or natural openings, such as the hydathode (Nino-Liu *et al.*, 2006). *Xanthomonas oryzae* pv. *oryzicola* (*Xoc*) is another bacterial pathogen of rice, which colonizes the mesophyll parenchyma tissue to cause bacterial leaf streak disease. *Xoc* enters the leaf primarily through the stomata (Nino-Liu *et al.*, 2006). To date, multiple virulence factors have been identified that contribute to bacterial invasion at different stages, such as adhesion-like proteins, exopolysaccharide (EPS) and lipopolysaccharide (LPS), biofilm formation and the type II secretion system (T2SS), which is used for the secretion of extracellular enzymes, including xylanase and cellulase (Büttner and Bonas, 2010; Das *et al.*, 2009; Kim *et al.*, 2009; Rai *et al.*, 2012; Rajeshwari *et al.*, 2005). Like many other Gram-negative plant-pathogenic bacteria, *Xoo* and *Xoc* also possess a T3SS to inject and deliver effector proteins into host cells. It is encoded by the *hypersensitive response and pathogenicity* (*hrp*) gene locus (Cho *et al.*, 2008; Zou *et al.*, 2006). The T3SS and its secreted components play a critical role in conferring pathogenicity on the host and triggering the hypersensitive response (HR) on resistant or non-host plants (Alfano and Collmer, 1997; Büttner and Bonas, 2003). The core operon consists of more than 20 genes on several transcriptional units, which contain *hrp*, *hrc* (*hrp-conserved*) and *hpa* (*hrp-associated*) genes (Cho *et al.*, 2008; Zou *et al.*, 2006). Two types of effector have been found in *X. oryzae*: TAL (transcription activator-like) and non-TAL effectors (Salzberg *et al.*, 2008; White and Yang, 2009). These effectors often determine the consequence of interactions between the bacterium and different hosts.

*hrp* gene expression is tightly regulated, and is induced *in planta* or in specially prepared minimal medium, which is designed to mimic *in planta* conditions. Expression is suppressed in nutrient-rich medium (Tang *et al.*, 2006; Tsuge *et al.*, 2002). The *hrp* genes in *Xanthomonas* spp. and *Ralstonia solanacearum* have been classified as *hrp* group II, which differ from group I of *E. amylovora* and *Pseudomonas syringae* (Alfano and Collmer, 1997; Tang *et al.*, 2006). The expression of *hrp* genes in group I is activated by alternative sigma factor HrpL (Chatterjee *et al.*, 2002; Xiao *et al.*, 1994). In group II, *hrpG* and *hrpX*, which are spatially located away from the *hrp* gene cluster, are two key known regulatory genes of *hrp* gene expression (Wengelnik and Bonas, 1996; Wengelnik *et al.*, 1996). HrpG, a response regulator belonging to the OmpR family of two-component signal transduction systems (TCS), positively regulates the expression of *hrpX* (Wengelnik *et al.*, 1996). HrpX is an AraC family regulator which activates the transcription of other *hrp* genes (*hrpB* to *hrpF*), together with the genes encoding T3 effectors (Wengelnik and Bonas, 1996). Most of the genes in the HrpG regulon are regulated by HrpX. HrpX interacts with a cis-element within the promoter region of *hrp*

genes, known as the plant-inducible promoter (PIP)-box, which is also present in the promoter of many T3 effectors (Noel *et al.*, 2001; Wengelnik and Bonas, 1996). In addition to HrpG and HrpX, HrpD6 has been reported recently as a *hrp* regulator in *Xoc*, responsible for the control of a subgroup of *hrp* genes (Li *et al.*, 2011).

In this article, a small library of phenolic compounds was screened for their effectiveness on T3SS expression of *Xoo*. Four inhibitors, which can suppress the HR in tobacco without killing bacterial cells, were identified and selected for further analysis. Their effects on effector translocation and expression of representative *hrp* genes were also examined. Virulence assays indicated that the symptoms in rice caused by both *Xoo* and *Xoc* were weakened by these four inhibitors to varying degrees.

## RESULTS

### Screening of inhibitors which affect the T3SS of *Xoo*

In order to identify potential inhibitors of the *Xoo* T3SS, a library of 56 plant phenolic compounds and their derivatives (Tables 1 and S1, see Supporting Information) was screened for their effects on the promoter activity of the *hpa1* gene, which encodes a harpin protein in *Xoo* (Furutani *et al.*, 2003; Wang *et al.*, 2008). Expression of *hpa1* is induced in the *hrp*-inducing medium XOM2, and is controlled by the regulatory protein HrpX (Furutani *et al.*, 2003). The promoter region of *hpa1* (Fig. S1, see Supporting Information) was inserted into the promoter-probe vector pPROBE-AT (Li *et al.*, 2009), which contains a promoterless green fluorescence protein (GFP) reporter gene, resulting in pPhpa1. *Xoo* PXO99<sup>A</sup> strain carrying pPhpa1 was grown in XOM2 supplemented with each of the compounds at a final concentration of 200  $\mu$ M for 15 h before the promoter activity of *hpa1* was measured. Eleven of the 56 compounds were found to be insoluble in XOM2 medium (Table S1). Thus, their effects could not be determined. The other 45 compounds were screened for alterations in *hpa1* promoter activity through the highly efficient fluorescence-activated cell sorting (FACS) system. The mean fluorescence intensity (MFI), representing the promoter activity of *hpa1* in each treatment, was recorded (Table 1). We also calculated the ratio of MFI after treatment by each compound to that of the solvent control, resulting in a number indicated by %DMSO (DMSO, dimethylsulfoxide). The results showed that 25 compounds altered *hpa1* promoter activity significantly in comparison with the DMSO control and, of these, 10 compounds inhibited the *hpa1* promoter activity by at least 60% (Table 1).

### Five compounds suppress the HR caused by *Xoo* in tobacco


To determine whether the 10 inhibitors interfered with the function of T3SS in *Xoo*, we performed a secondary screening

**Table 1** Screening for inhibitors of the *Xanthomonas oryzae* pv. *oryzae* (*Xoo*) type III secretion system (T3SS) by fluorescence-activated cell sorting assays.

Phenolic compound	Average MFI $\pm$ SD <sup>a</sup>	%DMSO <sup>b</sup>
XOM2	15.63 $\pm$ 1.96	
Dimethylsulfoxide (DMSO)	14.97 $\pm$ 2.33	
TS001, <i>trans</i> -cinnamic acid	9.41 $\pm$ 0.10*	62.86
TS002, 2,4-dihydroxycinnamic acid	12.69 $\pm$ 0.10	84.77
TS006, <i>o</i> -coumaric acid	3.58 $\pm$ 0.08**	23.91
TS007, 3-(4-hydroxyphenyl)propionic acid	19.85 $\pm$ 0.74*	132.60
TS013, <i>trans</i> -3-(3-pyridyl)acrylic acid	7.66 $\pm$ 1.06*	51.17
TS015, <i>trans</i> -2-methoxycinnamic acid	4.43 $\pm$ 0.01*	29.59
TS027, <i>trans</i> -4-aminocinnamic acid	13.59 $\pm$ 1.28	90.78
TS028, <i>trans</i> -4-nitrocinnamic acid	10.26 $\pm$ 1.17	68.54
TS103, <i>trans</i> -4-hydroxycinnamohydroxamic acid	7.92 $\pm$ 0.26*	52.91
TS114, diethyl <i>trans</i> -2-(4-hydroxyphenyl)-vinylphosphonate	22.2 $\pm$ 3.11	148.30
TS115, <i>trans</i> -2-(4-hydroxyphenyl)vinylphosphonic acid	17.74 $\pm$ 1.31	118.50
TS117, <i>p</i> -coumarylamine	14.45 $\pm$ 0.23	96.53
TS118, <i>N</i> -(4-methoxycinnamyl)phthalimide	16.89 $\pm$ 0.34	112.83
TS130, <i>trans</i> -3-hydroxycinnamohydroxamic acid	5.37 $\pm$ 0.53*	35.87
TS132, <i>trans</i> -cinnamohydroxamic acid	9.10 $\pm$ 0.52*	60.79
TS139, 3-phenylpropionohydroxamic acid	4.85 $\pm$ 0.03*	32.40
TS141, <i>trans</i> -4-fluorocinnamohydroxamic acid	5.34 $\pm$ 0.44*	35.67
TS160, <i>trans</i> -2-methylcinnamohydroxamic acid	3.97 $\pm$ 0.04**	26.52
XOM2	20.26 $\pm$ 0.50	
DMSO	19.67 $\pm$ 1.45	
TS003, 3,4-dihydroxycinnamic acid	12.54 $\pm$ 1.09*	63.75
TS004, <i>p</i> -coumaric acid	22.08 $\pm$ 2.78	112.25
TS008, hydrocinnamic acid	18.98 $\pm$ 0.90	96.49
TS010, <i>trans</i> -2-phenylcyclopropane-1-carboxylic acid	6.68 $\pm$ 1.69**	33.96
TS011, <i>trans</i> -3-(2-thienyl)acrylic acid	18.46 $\pm$ 1.66	93.85
TS012, <i>trans</i> -3-indoleacrylic acid	4.41 $\pm$ 0.06**	22.42
TS014, <i>trans</i> -3-(4-imidazolyl)acrylic acid	18.81 $\pm$ 1.28	95.63
TS016, <i>trans</i> -3-methoxycinnamic acid	8.72 $\pm$ 0.29**	44.33
TS018, <i>trans</i> -2-methylcinnamic acid	7.26 $\pm$ 0.16**	36.91
TS019, <i>trans</i> -3-methylcinnamic acid	10.79 $\pm$ 0.46**	54.86
TS022, <i>trans</i> -3-chlorocinnamic acid	9.23 $\pm$ 0.18**	46.92
TS024, <i>trans</i> -2-carboxycinnamic acid	21.07 $\pm$ 4.16	107.12
TS025, <i>trans</i> -cinnamamide	17.96 $\pm$ 1.87	91.31
TS026, <i>trans</i> -4-mercaptocinnamic acid	4.48 $\pm$ 0.06**	22.78
TS029, <i>trans</i> -4-formylcinnamic acid	16.72 $\pm$ 0.72**	85.00
TS030, methyl <i>trans</i> -cinnamate	28.43 $\pm$ 5.36	144.53
TS031, <i>trans</i> -4-carboxycinnamic acid	19.61 $\pm$ 1.09	99.69
TS032, cinnamyl alcohol	29.88 $\pm$ 1.49	151.91
TS033, salicylic acid	12.51 $\pm$ 1.36*	63.60
TS034, benzoic acid	21.65 $\pm$ 2.306	110.07
TS124, <i>trans</i> -4-hydroxymethylcinnamic acid	20.67 $\pm$ 0.06	105.08
TS125, <i>trans</i> -4-methoxycinnamohydroxamic acid	20.71 $\pm$ 0.97	105.29
TS131, <i>trans</i> -3,4-dihydroxycinnamohydroxamic acid	9.06 $\pm$ 0.61**	46.06
TS133, <i>trans</i> -3-(4-hydroxyphenyl)acryloylhydrazide	11.75 $\pm$ 0.42**	59.74
TS134, benzhydroxamic acid	12.01 $\pm$ 1.88*	61.06
TS135, salicylhydroxamic acid	10.21 $\pm$ 0.34**	51.91
TS136, phenylpropionic acid	14.80 $\pm$ 2.473	75.24

<sup>a</sup>Green fluorescent protein (GFP) mean fluorescence intensity (MFI) was determined for gated populations of bacterial cells by flow cytometry. Values are representative of at least two independent experiments, and three replicates were used for each experiment. Asterisks indicate statistically significant differences in MFI between bacterial cells grown in XOM2 with DMSO and XOM2 supplemented with 200  $\mu$ M of each compound (Student's *t*-test). \**P* < 0.05; \*\**P* < 0.01.


<sup>b</sup>%DMSO was used to represent the relative promoter activity of *hpa1* in *Xoo* cells grown in XOM2 supplemented with 200  $\mu$ M of each compound in comparison with that in XOM2 with DMSO only, which was calculated by the formula: %DMSO = 100  $\times$  MFI(XOM2 with phenolic compounds)/MFI(XOM2 with DMSO).



**Fig. 1** Effects of various compounds on the hypersensitive response (HR) induced by *Xanthomonas oryzae* pv. *oryzae* (*Xoo*) on *Nicotiana benthamiana*. Cell suspensions of wild-type *Xoo* PXO99<sup>A</sup> (wild-type, WT) at an optical density at 600 nm (OD<sub>600</sub>) of 0.3 were incubated with dimethylsulfoxide (DMSO) or each compound at a concentration of 200  $\mu$ M for 2 h before infiltration into tobacco leaves.  $\Delta hrpX$  was used as a type III secretion system (T3SS) deficiency control. Photographs were taken 24 h after infiltration. At least four independent experiments were performed with similar results.

by examining their effects on the HR-inducing ability of *Xoo* on tobacco. Infiltration of *Xoo* cells into non-host tobacco leaves can induce HR, whereas deletion of the major T3SS regulatory gene *hrpX* completely abolishes the HR, suggesting that a functional T3SS is required for this phenotype (Fig. 1). Therefore, the HR-inducing ability can indicate whether the T3SS is active in the bacteria. To determine the appropriate concentration of compounds for HR assays, we first examined the dose-dependent effects of TS006, one of the strongest inhibitors of *hpa1* promoter activity. TS006 began to show an inhibitory effect at a concentration of 50  $\mu$ M. The impact was significantly enhanced at 100  $\mu$ M and reached a plateau at 200  $\mu$ M (Fig. S2, see Supporting Information). Therefore, we chose to test the effects at concentrations of 100 and 200  $\mu$ M. Bacterial cell suspensions were incubated with DMSO or each compound for 2 h before they were infiltrated into tobacco leaves. Meanwhile, the cell suspensions were plated onto peptone sucrose agar

(PSA) to check whether bacterial survival was affected by these compounds. At a concentration of 100  $\mu$ M, none of them showed inhibition of HR (data not shown). When the concentration was increased to 200  $\mu$ M, TS006, TS010, TS012, TS015 and TS018 caused a complete loss of HR (Fig. 1), suggesting that these compounds might effectively suppress the function of the *Xoo* T3SS. However, the numbers of bacterial cells were significantly decreased after treatment with TS012 (Fig. S3A, see Supporting Information). Cell counting revealed that bacterial populations were reduced by at least 50% (Fig. S3B), suggesting that TS012 has strong bactericidal activity. Comparatively, the other five compounds, TS026, TS130, TS139, TS141 and TS160, did not show inhibition of the HR phenotype at 200  $\mu$ M (Fig. 1). Even when the concentration was increased to 400  $\mu$ M, they did not demonstrate any inhibitory effect at all (data not shown), suggesting that they might not be functional *in planta*.



**Fig. 2** Effects of various compounds on bacterial growth rates. (A) The growth rate of *Xanthomonas oryzae* pv. *oryzae* (*Xoo*) PXO99<sup>A</sup> in rich medium (M210) supplemented with dimethylsulfoxide (DMSO) or 200  $\mu$ M of TS006, TS010, TS012, TS015 or TS018. (B) The growth rate of *Xoo* PXO99<sup>A</sup> in *hrp*-inducing medium (XOM2) (plus 0.5% sucrose) supplemented with DMSO or 200  $\mu$ M of TS006, TS010, TS012, TS015 or TS018. The optical density at 600 nm (OD<sub>600</sub>) of the culture suspensions was measured every 6 h during the 72-h period. Two independent tests were performed with similar results.

### TS012 severely affects the growth rate of *Xoo*

As shown in the previous section, except for TS012, the other four compounds did not demonstrate any apparent bactericidal effects on *Xoo*. In line with this, we investigated their effect on bacterial growth at different stages. The growth rate of *Xoo* was measured in both rich medium (M210) and *hrp*-inducing medium (XOM2) for a period of 72 h. The nutrient-scarce medium XOM2 was supplemented with 0.5% sucrose to support bacterial growth (Tian *et al.*, 2015). TS006, TS010, TS012, TS015 and TS018 were added to the media at a concentration of 200  $\mu$ M. In comparison with the non-treatment control, bacterial growth rates with the addition of either DMSO or compounds TS006, TS010, TS015 or TS018 did not show statistically significant changes at different time points, indicating that these compounds did not affect bacterial growth (Fig. 2). In contrast, treatment with TS012 significantly reduced the growth rate of *Xoo*. As we were looking for compounds that suppressed bacterial virulence, but did not affect bacterial growth or survival, we focused on TS006, TS010, TS015 and

TS018 for the remainder of the study. To be consistent, we used a final concentration of 200  $\mu$ M of each compound in all other assays.

### TS006, TS010, TS015 and TS018 suppress the translocation of PXO\_04172 and PXO\_03702 in rice

Translocation of T3 effectors into the host cell is the primary function of the T3SS. To evaluate directly whether this process was affected by the four inhibitors, we examined their effect on the translocation of two non-TAL effectors, which have been confirmed previously to be translocated using a Cya-based reporter system (Furutani *et al.*, 2009). Two plasmids, pHM04172 and pHM03702, carrying T3 effectors XOO3803 and XOO4042 of *Xoo* strain MAFF31101 (Furutani *et al.*, 2009), corresponding to effectors PXO\_04172 and PXO\_03702 in PXO99<sup>A</sup>, were transformed into PXO99<sup>A</sup> wild-type and *hrpX* mutant strains for translocation assays. With only DMSO treatment, a considerable amount of cyclic adenosine monophosphate (cAMP) was detected in rice leaves infected with wild-type strains carrying PXO\_04172 or PXO\_03702, whereas it was barely detectable after infection with the *hrpX* mutant strains, suggesting that these two effectors were properly translocated by the T3SS of PXO99<sup>A</sup> (Fig. 3). Compared with DMSO treatment, there was a significant reduction in cAMP accumulation in the presence of TS006, TS010, TS015 and TS018 (Fig. 3). These results indicate that translocation of PXO\_04172 and PXO\_03702 was suppressed by these four inhibitors, suggesting that the translocation process through the T3SS in *Xoo* might be repressed under these conditions.

### Expression of representative *hrp/hrc* genes is suppressed by inhibitors TS006, TS010, TS015 and TS018

The data outlined above demonstrate that TS006, TS010, TS015 and TS018 suppress the *Xoo* T3SS without affecting bacterial growth, implying that they might specifically interfere with the regulatory pathway for T3SS gene expression. Therefore, we carried out quantitative reverse transcription-polymerase chain reaction (qRT-PCR) experiments to examine the effect of these inhibitors on the expression levels of T3SS genes. The first screening showed that TS006, TS010, TS015 and TS018 reduced the promoter activity of *hpa1* by at least 60% (Table 1). Here, the qRT-PCR assays demonstrated that the mRNA level of *hpa1* was reduced by over 80% in *Xoo* cells treated with these inhibitors, which is consistent with the results obtained from the promoter assay (Fig. 4A). Next, we examined the expression levels of other *hrp/hrc* genes, including *hrpE* (encoding the *hrp* pilus protein), *hrpF* (encoding a putative translocon protein), *hrcC* (encoding the outer-membrane secretin) and export apparatus genes *hrcT* and *hrcU* (Gürlebeck *et al.*, 2006). As expected, the mRNA levels of the tested *hrp/hrc* genes were reduced significantly when the inhibitors were present, in


comparison with the DMSO control (Fig. 4A,B). It is worth noting that TS006, among the four inhibitors, consistently showed the strongest inhibitory effect on all genes tested.

### The four inhibitors affect the expression of regulatory genes *hrpG* and *hrpX*

It is known that the expression of the *Xanthomonas* T3SS is controlled by the regulatory cascade consisting of HrpG and HrpX (Büttner and Bonas, 2010). As we observed a reduction in mRNA levels in representative *hpa1/hrp/hrpC* genes in *Xoo* after treatment with the four inhibitors, it was imperative to investigate whether the expression of *hrpG* and *hrpX* was affected. The mRNA levels of *hrpG* and *hrpX* were examined in *Xoo* cells grown in the presence of each inhibitor. In comparison with the DMSO control, the mRNA level of *hrpX* was reduced by approximately 50% in the presence of each inhibitor (Fig. 4C). In contrast, the mRNA level of *hrpG* was reduced by about 30% by TS006, TS015 and TS018, and even less by TS010 (Fig. 4C). These results suggest that the inhibitory effect of TS006, TS010, TS015 and TS018 on T3SS gene expression of *Xoo* exists through the HrpG–HrpX regulatory cascade, but with varying degrees of intensity.

### The promoters of *hrcC* and *hrpB* respond differentially to inhibitors TS006, TS010, TS015 and TS018

To further investigate whether HrpG or HrpX played a role in relaying the impact of the inhibitors on the transcription of T3SS genes in *Xoo*, we examined their effect on two different promoters within




**Fig. 3** The effect of TS006, TS010, TS015 and TS018 on the translocation of the type III effectors PXO\_04172 and PXO\_03702 from *Xanthomonas oryzae* pv. *oryzae* (*Xoo*) PXO99<sup>A</sup> into rice cells, examined by Cya translocation assays. The concentration of each compound was 200  $\mu$ M. An equal volume of dimethylsulfoxide (DMSO) was used as a control. The level of cyclic adenosine monophosphate (cAMP) indicates the level of the translocated T3 effectors. Three replicates were used in each experiment. At least three independent tests were performed with similar results.

their regulon. As mentioned above, most of the genes in the *hrp* cluster were activated by the transcriptional regulator HrpX, which recognizes the PIP-box (TTCGC-N<sub>15</sub>-TTCGC) in the promoter region, such as that of *hpa1* (Fig. S4A, see Supporting Information). The *hrpB* operon, spanning from *hrpB1* to *hrcT*, is located within the core *hrp* gene cluster of *Xoo* (Fig. 5A) (Cho *et al.*, 2008). *hrcC*, which is next to *hrcT* has been confirmed to be transcribed together with the *hrpB* operon (Cho *et al.*, 2008). Interestingly, it has been shown recently that *hrcC* in *Xoc* also uses an internal promoter for transcription (Li *et al.*, 2011). The internal promoter in front of *hrcC*, designated as  $P_{hrcC}$  does not contain a PIP-box, and was regulated by HrpG, but not HrpX (Li *et al.*, 2011). Sequence alignment of the promoter regions of *hrcC* in *Xoo* PXO99<sup>A</sup> and *Xoc* RS105 showed that they share over 90% identity (Fig. S4B). Therefore, we amplified this region, named  $P_{hrcC}$  in *Xoo* (Fig. 5A), to test whether its activity was affected by the four inhibitors. In addition, we amplified the promoter for the *hrpB* operon, designated as  $P_{hrpB}$ , which contains a perfect PIP-box for comparison (Fig. S4C). FACS assays were performed to examine the impact of TS006, TS010, TS015 and TS018 on the activity of  $P_{hrcC}$  and  $P_{hrpB}$ . After calculating the ratio of compound treatment to the DMSO control (DMSO%), we found that the activity of  $P_{hrcC}$  was not affected by these four compounds at all (Fig. 5B). In contrast, the activity of  $P_{hrpB}$  decreased by at least 50% after treatment with these inhibitors in comparison with the DMSO control. This effect was similar to their effect on the *hpa1* promoter (Fig. 5B). These results suggest that promoters containing a PIP-box might be the main targets of these inhibitors, which also supports the statement that HrpX plays an important role in the mediation of the inhibitory effect. Although the activity of  $P_{hrcC}$  was not affected by these inhibitors, its mRNA level was reduced significantly, as shown above (Fig. 4B). A plausible hypothesis might be that part of the mRNA transcribed from  $P_{hrpB}$  is reduced, leading to a decrease in the total mRNA level of *hrcC*.

### The expression of other virulence factors of *Xoo* is not suppressed by T3SS inhibitors

Previously, it has been shown that several compounds related to salicylidene acylhydrazides suppress both T3SS and the EPS amylovoran in *E. amylovora* (Yang *et al.*, 2014). Therefore, we wanted to investigate whether the effects of the *Xoo* T3SS inhibitors identified above were specific or effective on multiple virulence factors. The disruption of the T2SS structural genes *xpsE* and *xpsF* led to reduced secretion of extracellular enzymes and impaired virulence of *Xoo* (Ray *et al.*, 2000; Sun *et al.*, 2005). The mRNA levels of *xpsE* and *xpsF* were measured by qRT-PCR in *Xoo* cells treated with the T3SS inhibitors TS006, TS010, TS015 and TS018. In comparison with the DMSO control, the four compounds did not reduce the mRNA levels of *xpsE* and *xpsF* (Fig. 6A). The expression of representative *gum* genes in *Xoo* cells treated with the four inhibitors was also examined. The results showed that



**Fig. 4** Relative mRNA levels of *hrp* genes in *Xanthomonas oryzae* pv. *oryzae* (*Xoo*) PXO99<sup>A</sup> incubated with compounds TS006, TS010, TS015 and TS018, measured by quantitative reverse transcription-polymerase chain reaction (qRT-PCR). (A) The mRNA levels of *hrpA*, *hrpE* and *hrpF* were reduced significantly after treatment with these inhibitors in comparison with the dimethylsulfoxide (DMSO) control. (B) The mRNA levels of three *hrc* genes, i.e. *hrcC*, *hrcT* and *hrcU*, were also reduced significantly after treatment with these inhibitors in comparison with the DMSO control. (C) The mRNA levels of *hrpG* and *hrpX* genes were reduced to varying extents after treatment with different inhibitors in comparison with the DMSO control. The concentration of each compound used was 200  $\mu\text{M}$ . An equal volume of DMSO was used as a control. *gyrB* was used as an endogenous control for data analysis. Three replicates were used in each experiment. Three independent tests were performed with similar results. Asterisks indicate statistically significant differences (Student's *t*-test). \*\* $P < 0.01$ .

the mRNA levels of *gumB*, *gumD*, *gumK* and *gumM* were not reduced by these T3SS inhibitors (Fig. 6B).

The impact of the four inhibitors on the mRNA levels of *wxoE*, *smtA* and *wxoA* from cluster I, and *xanB* from cluster II, was similarly examined. Except for *smtA*, which was slightly down-regulated by TS010 and TS015, the other three genes did not show reduced expression after treatment by the four inhibitors. Overall, these results clearly demonstrate that the four T3SS inhibitors do not suppress major gene expression of T2SS, EPS and


LPS. Instead, their expression is enhanced in some cases (Fig. 6A–C), which is probably a result of negative cross-talk between different virulence factors. Therefore, we conclude that the inhibitory effect of these inhibitors is specific to T3SS, and not expanded to other virulence factors.

#### TS006, TS010, TS015 and TS018 suppress the water-soaking and disease symptoms of *Xoo* on rice

Our focus in this work was to demonstrate that these inhibitors can suppress the virulence phenotypes of *Xoo* on rice. On the seedlings of susceptible rice cultivar IR24, *Xoo* PXO99<sup>A</sup> induced water-soaked lesions after infiltration of bacterial cells into the leaves (Fig. 7A). The water-soaking symptoms were reduced to various levels by treatment of the bacterial cells with TS006, TS010, TS015 and TS018 (Fig. 7A). Moreover, the yellowish disease symptoms on adult IR24 plants were also weakened by these inhibitors (Fig. 7B), and the lesion lengths were significantly reduced in comparison with DMSO treatment (Fig. 7C). In addition, we also measured the bacterial population in the diseased leaf tissue. After treatment with each inhibitor, the bacterial numbers isolated from the leaves were significantly reduced (Fig. S5, see Supporting Information). These results indicate that TS006, TS010, TS015 and TS018 do indeed suppress the virulence of *Xoo* cells, and reflect their propagation *in planta*.

#### TS006, TS010, TS015 and TS018 suppress the virulence of *Xoc* on rice

As the *hrp* clusters among *Xanthomonas* spp. are highly conserved (Büttner and Bonas, 2010; Nino-Liu *et al.*, 2006), and the major regulatory proteins HrpG and HrpX are also functional in *Xoc* (Zou *et al.*, 2006), we speculated that it might be possible for these inhibitors to also suppress the T3SS of *Xoc*. Therefore, it was reasonable to test whether they suppressed the virulence of *Xoc* on rice. As shown in Fig. 8A, the disease symptoms caused by *Xoc* RS105 on rice seedlings were significantly reduced by TS006 and TS010 (Fig. 8A). On adult plants with only DMSO treatment, the lesion length can reach 4–5 cm, whereas, with the addition of the four compounds, lesion lengths were below 1 cm (Fig. 8B,C), suggesting that the virulence of *Xoc* was significantly suppressed by these inhibitors. As the four compounds exhibited stronger inhibitory effects on the virulence of *Xoc* than *Xoo* at the same concentration (200  $\mu\text{M}$ ), it is necessary to include the possibility that they might affect the growth or survival of *Xoc*. Therefore, we measured the growth rate of *Xoc* in both rich medium (nutrient broth, NB) and *hrp*-inducing medium (XOM2) (plus 0.5% sucrose) during a period of 72 h. TS006, TS010, TS015 and TS018 were added to the media at a final concentration of 200  $\mu\text{M}$ . In comparison with the non-treatment control, bacterial growth rates with the addition of either DMSO or each compound did not show significant alterations at different time points, indicating that these compounds did



**Fig. 5** Impact of TS006, TS010, TS015 and TS018 on the promoter activity of *hrpB* and *hrcC*. (A) Schematic diagram of  $P_{hrpB}$  and  $P_{hrcC}$ . (B) Promoter activities of *hrpB* and *hrcC* in *Xanthomonas oryzae* pv. *oryzae* (Xoo) PXO99<sup>A</sup> grown in XOM2 supplemented with 200  $\mu$ M of TS006, TS010, TS015 and TS018 for 15 h, measured by fluorescence-activated cell sorting (FACS). Dimethylsulfoxide (DMSO) was used as a negative control. Green fluorescent protein (GFP) mean fluorescence intensity (MFI) was determined by flow cytometry. %DMSO =  $100 \times \text{MFI}(\text{XOM2 with each compound})/\text{MFI}(\text{XOM2 with equal volume of DMSO})$ . Three independent tests were performed with similar results.

not affect the bacterial growth of *Xoc* (Fig. S6, see Supporting Information). Thus, we conclude that TS006, TS010, TS015 and TS018 suppress the virulence of *Xoc*, but do not affect its growth.

## DISCUSSION

Rice is the staple food of the world, feeding more than one-half of its population. However, rice is vulnerable to pathogen infection, which often leads to devastating diseases, resulting in severe yield losses. Bacterial leaf blight and leaf streak diseases caused by the two pathovars of *X. oryzae* are the most important bacterial diseases around the world, especially in Asia and Africa (Sundaram *et al.*, 2014). Here, we report, for the first time, that phenolic compounds TS006, TS010, TS015 and TS018 are able to suppress the disease symptoms of *Xoo* and *Xoc* on rice by specifically inhibiting the function of the T3SS. The compounds were identified by screening a small library of natural phenolics and their derivatives, some of which have been shown previously to affect the T3SS gene expression of several bacterial pathogens, including *D. dadantii* (Li *et al.*, 2009; Yang *et al.*, 2008), *P. aeruginosa* (Yamazaki *et al.*, 2012) and *E. amylovora* (Khokhani *et al.*, 2013; Li *et al.*, 2015).


A whole-cell-based high-throughput screening (HTS) approach has often been used to identify T3SS inhibitors from compound libraries (Marshall and Finlay, 2014). Here, the *Xoo* strain PXO99<sup>A</sup>, harbouring a reporter plasmid with a *gfp* gene transcriptionally fused to the *hpa1* promoter, was constructed for screen-

ing. Purposely, the *hpa1* promoter was used because Hpa1 has been confirmed to be a T3SS-secreted harpin protein in both *Xoo* and *Xoc* (Furutanin *et al.*, 2003; Zhu *et al.*, 2000; Zou *et al.*, 2006). Twenty-five of 45 compounds showed significant inhibitory effects on *hpa1* promoter activity (Table 1). This efficiency is much higher than that of HTS using large libraries containing thousands of natural or synthetic small molecules (Charro and Mota, 2015; Felise *et al.*, 2008). A secondary screening was performed to identify HR inhibitors from the 10 strongest compounds showing inhibition of promoter activity. At a concentration of 200  $\mu$ M, TS006, TS010, TS012, TS015 and TS018 showed suppression of HR of *Xoo* in tobacco (Fig. 1). Interestingly, the other five compounds (TS026, TS130, TS139, TS141 and TS160) did not demonstrate any inhibitory effects at all, even at higher concentrations (Fig. 1). There could be many reasons for their inability to suppress HR *in planta*. One possible explanation is that they might not be stable *in planta* to perform their functions. Of course, more work needs to be performed in the future to fully elucidate the exact cause.

As a result of their possible positive impact on agriculture, it is important to understand the functional mechanism of these T3SS inhibitors. Our results indicated that TS006, TS010, TS015 and TS018 suppressed *hrp/hrc* gene expression, probably through the major regulatory proteins HrpG and HrpX (Fig. 4). In addition, we examined the activity of  $P_{hrcC}$  which does not contain a PIP-box like  $P_{hpa1}$  (Fig. 5). In *Xoc*,  $P_{hrcCP}$  was positively regulated by HrpG, but not HrpX (Li *et al.*, 2011). The activity of  $P_{hrcC}$  was not suppressed by the four T3SS inhibitors, suggesting that the regulatory role of HrpX might be indispensable for the function of these inhibitors. We speculated that these inhibitors may exert their effects upstream of HrpG, as the expression of *hrpG* and *hrpX* was affected (Fig. 4). Presumably, phosphorylated HrpG activates the expression and production of HrpX, and HrpX regulates the downstream genes (Büttner and Bonas, 2010). However, the signalling transduction upstream of HrpG remains elusive. Recently, a putative histidine kinase of HrpG has been reported in *X. campestris* pv. *campestris* (Li *et al.*, 2014). In addition, the RNA-binding protein RsmA has been shown to positively regulate the T3SS by stabilizing HrpG mRNA in *X. citri* ssp. *citri* (Andrade *et al.*, 2014). It will be of great interest to study whether these components play similar roles in regulating the T3SS in *X. oryzae*, and whether they mediate the inhibitory effects of these compounds on the T3SS.

By comparison with previous screening results of these phenolic compounds in other bacteria, we observed some differences in their activities. For example, TS006 (OCA), the inhibitor of the *Xoo* T3SS, which has also been shown to inhibit the T3SS of *E. amylovora* (Khokhani *et al.*, 2013), was initially identified to induce T3SS gene expression in *D. dadantii* (Li *et al.*, 2009; Yang *et al.*, 2008). In contrast, TS004 (PCA), a strong inhibitor of the T3SS in *D. dadantii* and *E. amylovora*, did not show significant alteration of *hpa1* promoter activity in *Xoo* (Table 1). In *D. dadantii*, PCA





**Fig. 6** Impact of 200  $\mu\text{M}$  of TS006, TS010, TS015 and TS018 on the mRNA levels of genes encoding the type II secretion system (T2SS) (A), exopolysaccharide (EPS) (B) and lipopolysaccharide (LPS) (C) in *Xanthomonas oryzae* pv. *oryzae* (*Xoo*) PXO99<sup>A</sup>, measured by quantitative reverse transcription-polymerase chain reaction (qRT-PCR). The *gyrB* gene was used as an endogenous control for data analysis. Three replicates were used in each experiment. Three independent tests were performed with similar results. Asterisks indicate statistically significant differences (Student's *t*-test). \* $P < 0.05$ ; \*\* $P < 0.01$ . DMSO, dimethylsulfoxide.

inhibited the T3SS gene expression through the HrpX/HrpY TCS (Li *et al.*, 2009), which is a pathway that does not exist in *Xoo*. Therefore, different T3SS regulatory pathways might be a major reason for the distinctive activities detected for these compounds.


Given the possibility that the T3SS inhibitors might also affect the expression of other virulence factors, it was important to observe the behaviour of these types of T3SS inhibitor. Here, we analyzed the effects of TS006, TS010, TS015 and TS018 on the expression of other important virulence factors in *Xoo*, including the T2SS, EPS and LPS (Büttner and Bonas, 2010; Das *et al.*, 2009; Kim *et al.*, 2009; Rai *et al.*, 2012; Rajeshwari *et al.*, 2005). We found that most genes encoding their products were not affected by the four T3SS inhibitors (Fig. 6). These results are consistent with the proposed mechanism that the T3SS inhibitors act through

the HrpG–HrpX cascade, as the transcription of these genes is most probably not controlled by HrpX. Meanwhile, it was intriguing that some genes were up-regulated by the addition of the inhibitors. A similar phenomenon has also been reported previously. For example, in *E. amylovora*, transcriptome analysis demonstrated that a large number of genes were up-regulated when treated with T3SS inhibitors (Yang *et al.*, 2014). In *Salmonella*, when a compound was used to inhibit the secretion of the T3 substrate SipA, a significantly increased amount of the flagellin protein FliC was detected (Felise *et al.*, 2008). This might be a result of cross-talk between virulence factors in bacteria, which would warrant further investigation.

Water soaking is a symptom specifically related to the function of the AvrBs3/PthA family of effectors in Xanthomonads, which were more recently renamed as 'TAL (transcription activator-like)' effectors (Verdier *et al.*, 2012; White and Yang, 2009). In *Xoo* strain PXO99<sup>A</sup>, there are multiple TAL effectors, which confer gene-specific resistance in host plants (White and Yang, 2009). After syringe infiltration in the susceptible rice cultivar IR24, where no corresponding *R* genes exist, the PXO99<sup>A</sup> strain caused strong water-soaked lesions (Verdier *et al.*, 2012; Fig. 7A). We found that, after performing a treatment protocol with TS006 and TS010, the water-soaking phenotypes on rice were almost completely abolished (Fig. 7A). TS015 and TS018 also suppressed the water-soaking phenotype to varying levels (Fig. 7A). Using the translocation assays including two non-TAL effectors as representatives (Fig. 3), these results demonstrated that the primary function of the T3SS, which is the delivery of effectors into plant cells, was suppressed by TS006, TS010, TS015 and TS018 with different intensities.

Finally, virulence assays were performed to evaluate the effect of the four inhibitors in preventing the disease symptoms of *Xoo* and *Xoc* on rice. Unexpectedly, TS006 and TS010, which almost completely abolished the water-soaking symptoms, only reduced the disease lesion lengths of *Xoo* by 20%–30% (Fig. 7C). This might be because other virulence factors were still functional or even enhanced by these inhibitors (Fig. 6). In comparison, the reduction in disease lesion lengths of *Xoc* on rice was more evident (Fig. 8B,C). Although these T3SS inhibitors were screened using *Xoo* as the reporter strain, it was not surprising to observe their functions in *Xoc*, as the major T3SS regulatory pathways are very similar between *Xoo* and *Xoc* (Cho *et al.*, 2008; Zou *et al.*, 2006). With regard to why the inhibitors were more efficient in suppressing the virulence of *Xoc*, different infection methods between the two pathovars might be a possible reason.

In summary, we have demonstrated the inhibitory effect of phenolic compounds TS006, TS010, TS015 and TS018 on the T3SS of *Xoo* both *in vitro* and *in planta*. Furthermore, these compounds have been proven to be effective in suppressing the disease symptoms caused by *Xoo* and *Xoc* on rice leaves. These findings are important because they may provide potential anti-virulence



**Fig. 7** Compounds TS006, TS010, TS015 and TS018 suppress the virulence of *Xanthomonas oryzae* pv. *oryzae* (Xoo) PXO99<sup>A</sup> on rice cultivar IR24. (A) The effect of TS006, TS010, TS015 and TS018 on the water-soaking symptoms caused by Xoo PXO99<sup>A</sup> (wild-type, WT) on IR24 seedling.  $\Delta hrpX$  was used as a type III secretion system (T3SS) deficiency control. Photographs were taken 3 days after infiltration. The disease symptoms (B) and lesion lengths (C) of Xoo PXO99<sup>A</sup> on adult plants of rice cultivar IR24 were reduced after pretreatment with TS006, TS010, TS015 and TS018. Photographs were taken 14 days after infiltration. At least three independent tests were performed with similar results. Asterisks indicate statistically significant differences (Student's *t*-test). \*\**P* < 0.01. DMSO, dimethylsulfoxide.

drugs, which might be used to prevent infection in agricultural production in the future.

## EXPERIMENTAL PROCEDURES

### Bacterial strains, plasmids and growth conditions

The bacterial strains and plasmids used in this study are listed in Table 2. *Escherichia coli* was grown in Luria–Bertani (LB) medium at 37°C. Xoo wild-type strain PXO99<sup>A</sup> and the derived strains were grown in M210 medium (0.8% casein enzymatic hydrolysate, 0.5% sucrose, 0.4% yeast extract, 17.2 mM K<sub>2</sub>HPO<sub>4</sub>, 1.2 mM MgSO<sub>4</sub>·7H<sub>2</sub>O) or on PSA plates. XOM2 medium (0.18% D-(+)-xylose, 670 μM L-methionine, 10 mM sodium L-(+)-glutamate, 14.7 mM KH<sub>2</sub>PO<sub>4</sub>, 40 μM MnSO<sub>4</sub>, 240 μM Fe(III)-EDTA and 5 mM MgCl<sub>2</sub>, the pH was adjusted to 6.5 with KOH) was used for *hrp*-inducing conditions (Tsuje *et al.*, 2002). Xoc wild-type strain RS105 was grown on NA (0.5% peptone, 0.1% yeast, 1% sucrose, 0.3% beef extract and 1.5% agar) or NB (NA without agar) medium at 28°C. Antibiotics were used at the following final concentrations (μg/mL) when required: ampicillin (Ap), 100; spectinomycin (Sp), 50; kanamycin (Kan), 50; cephalixin (Cp), 25.


### Sources of the screened compounds

Compounds TS001 to TS035, TS108 to TS113, and TS134 to TS136 were purchased from commercial sources Aldrich (St. Louis, MO, USA), Alfa


Aesar (Ward Hill, MA, USA), and TCI (Tokyo, Japan). The remaining compounds were synthesized via the routes described in our previously published papers (Khokhani *et al.*, 2013; Yamazaki *et al.*, 2012). All compounds were dissolved in DMSO.

### Construction of the *hrpX* gene deletion mutant

An in-frame deletion mutation of the *hrpX* gene was constructed in PXO99<sup>A</sup> through homologous recombination using the suicide vector pKMS1, as described previously (Li *et al.*, 2011). The *sacB* gene (sucrose sensitivity counter-selectable marker) on pKMS1 confers suicide ability to the host bacterium during growth on high-concentration sucrose-containing medium. Briefly, approximately 600 bp of the upstream and 900 bp of the downstream region of the *hrpX* gene were amplified from PXO99<sup>A</sup> genomic DNA using the primer pairs *hrpXUSF/R* and *hrpXDSF/R*, respectively. The primers used in this study are listed in Table S2 (see Supporting Information). The two fragments were ligated into the suicide vector pKMS1, and introduced into PXO99<sup>A</sup> by electroporation. The transformants were first selected on NAN medium (consisting of kanamycin, 1% tryptone, 0.1% yeast extract, 0.3% peptone, 1.5% agar) and after continuous transfer culture in NBN broth (1% tryptone, 0.1% yeast extract, 1% sucrose, 0.3% peptone) four times. The potential mutants were selected on NAS medium (1% tryptone, 0.1% yeast extract, 10% sucrose, 0.3% peptone and 1.5% agar). The mutant candidates that grew on NAS, but were sensitive to kanamycin, were further confirmed by PCR.



**Fig. 8** Compounds TS006, TS010, TS015 and TS018 suppress the virulence of *Xanthomonas oryzae* pv. *oryzicola* (Xoc) RS105 on rice cultivar IR24. (A) The impact of TS006, TS010, TS015 and TS018 on disease symptoms caused by infiltration of Xoc RS105 (wild-type, WT) in IR24 seedlings. Photographs were taken 3 days after infiltration. (B) The impact of TS006, TS010, TS015 and TS018 on disease symptoms cause by needling infiltration of Xoc RS105 in IR24 adult plants. Photographs were taken 14 days after infiltration. (C) Lesion lengths on adult plants. At least three independent tests were performed with similar results. DMSO, dimethylsulfoxide.



10

### Construction of reporter strains and flow cytometry analysis

To screen compounds that induce or inhibit the expression of the PX099<sup>A</sup> T3SS, a 246-bp fragment containing the promoter region of *hpa1* was PCR amplified using the primers Phpa1-F and Phpa1-R. The amplified fragment was digested with *Bam*HI and *Eco*RI, and ligated to pPROBE-AT, a broad-host-range vector carrying a promoter-less *gfp* gene (Miller *et al.*, 2000), resulting in pPhpa1. This plasmid was then transferred to PX099<sup>A</sup> by electroporation. PX099<sup>A</sup> carrying the pPhpa1 or promoterless pPROBE-AT was grown in M210 overnight and transferred to XOM2 or XOM2 supplemented with 200  $\mu$ M of each compound. The promoter activity of *hpa1* was analysed using a FACS-Caliber flow cytometer (BD Bioscience, San Jose, CA, USA) as described previously (Yamazaki *et al.*, 2012). An equivalent volume of DMSO was added as a negative control. Three independent experiments were performed, and three replicates were used in each experiment. The promoter activities of *hrcC* and *hrcB* were analysed by a similar method. PhrcC-F/R and PhrcT-F/R were used to amplify the promoter regions of *hrcC* and *hrcB*, respectively (Li *et al.*, 2011).

### RNA extraction and qPCR analysis

Xoo cells were cultured in M210 medium overnight at 28°C and subcultured to XOM2 at an optical density at 600 nm ( $OD_{600}$ ) of 0.3, supplemented with DMSO or 200  $\mu$ M of each compound for 15 h. Total RNA

was isolated using an RNAprep Pure Bacteria Kit (Tiangen, Beijing, China). cDNA was synthesized using an HiScriptII Q RT SuperMix Kit (Vazyme, Nanjing, China). The cDNA levels of different samples were quantified by real-time PCR using a SYBR Green Master Mix (Vazyme). The relative levels of gene expression were determined using the  $2^{-\Delta\Delta CT}$  method (Livak and Schmittgen, 2001), with the DNA gyrase subunit B (*gyrB*) gene as the internal control (Tsuge *et al.*, 2002). Three technical replicates were used each time.

### HR assay

Xoo cells were cultured in M210 medium overnight at 28°C and resuspended in sterile distilled water. The bacterial suspensions were adjusted to  $OD_{600} = 0.3$ . *Nicotiana benthamiana* plants were used for HR assays. Cell suspensions were mixed with 200  $\mu$ M of each compound or DMSO, and incubated at 28°C for 2 h, before infiltration into tobacco using a needleless syringe. The HR symptoms were observed and photographed at 24 h after inoculation. Bacterial cell numbers were counted after serial dilutions and plating on PSA plates.

### Measurement of the growth rate

Xoo or Xoc cells were grown overnight in M210 or NB at 28°C. The cells were then resuspended in M210, NB or XOM2 (plus 0.5% sucrose) medium supplemented with DMSO or 200  $\mu$ M of each compound, starting

**Table 2** Strains and plasmids used in this study.

Strains and plasmids	Relevant characteristics	Reference or source
Strains		
<i>Xanthomonas oryzae</i> pv. <i>oryzae</i> PXO99 <sup>A</sup>	Wild-type strain, Philippine race 6, Cp <sup>r</sup>	Laboratory collection
$\Delta$ <i>hrpX</i>	<i>hrpX</i> gene deletion mutant of PXO99 <sup>A</sup> , Cp <sup>r</sup>	This study
<i>Xanthomonas oryzae</i> pv. <i>oryzicola</i> RS105	Wild-type, Chinese race 2	Dr Fengquan Liu
Plasmids		
pPROBE-AT	Promoter-probe vector, Ap <sup>r</sup>	
pKMS1	Suicidal vector carrying <i>sacB</i> gene for non-marker mutagenesis, Km <sup>r</sup>	Dr Gong-you Chen (Li <i>et al.</i> , 2011)
pPhpa1	pProbe-AT derivative with PCR fragment containing <i>hpa1</i> promoter region, Ap <sup>r</sup>	This study
pPhrcC	pProbe-AT derivative with PCR fragment containing <i>hrcC</i> promoter region, Ap <sup>r</sup>	This study
pPhrpB	pProbe-AT derivative with PCR fragment containing <i>hrpB</i> promoter region, Ap <sup>r</sup>	This study
pKHrpX	pKMS1 derivative carrying an in-frame <i>hrpX</i> mutation, Km <sup>r</sup>	This study
pHM04172	The putative promoter region and the 5'-coding region of PXO_04172 fused with a <i>cya</i> -tag, Sp <sup>r</sup>	Dr Seiji Tsuge (Furutani <i>et al.</i> , 2009)
pHM03702	The putative promoter region and the 5'-coding region of PXO_03702 fused with a <i>cya</i> -tag, Sp <sup>r</sup>	Dr Seiji Tsuge (Furutani <i>et al.</i> , 2009)

Ap<sup>r</sup>, ampicillin resistance; Cp<sup>r</sup>, cephalixin resistance; Km<sup>r</sup>, kanamycin resistance; Sp<sup>r</sup>, spectinomycin resistance.

at an OD<sub>600</sub> of 0.05. The growth rates were monitored every 6 h during the 72-h period using a Synergy 4 multimode microplate reader (BioTek, Winooski, VT, USA). Two independent experiments were performed, and three replicates were used in each experiment.

### Adenylate cyclase translocation assays

Plasmids expressing C-terminal Cya fusions of type III effectors PXO\_04172 and PXO\_03702, which were named as pHM04172 and pHM03702, were kindly provided by Dr Seiji Tsuge (Furutani *et al.*, 2009). PXO99<sup>A</sup> strain carrying pHM04172 or pHM03702 was grown in M210 medium overnight at 28°C, and resuspended in sterile distilled water to an OD<sub>600</sub> of 0.6. The bacterial cell suspensions were incubated with DMSO or 200 μM of each compound for 2 h before infiltration into the leaves of rice (*Oryza sativa* ssp. *indica* cultivar IR24) for translocation assays. Rice leaf sections (length, 1 cm) that included the inoculation sites were collected 3 days after inoculation, and homogenized in 300 μL of 0.1 M HCl with stainless beads using a vibration-ball mill (GRINOER, Beijing, China) (1000 rpm for 1 min). Then, cell debris was removed by centrifugation (13 400 g) for 2 min. The supernatant was collected in new tubes and stored at -80°C until use. The level of cAMP was quantified using the cAMP ELISA kit (Enzo Life Sciences, Lausen, Switzerland) system according to the manufacturer's instructions. The protein concentration of each sample was determined with a BCA Protein Assay kit (CWBI, Beijing, China).

### Pathogenicity assays

*Xoo* and *Xoc* cells were cultured in M210 or NB medium overnight at 28°C and resuspended in sterile distilled water. Cell suspensions at an OD<sub>600</sub> of 0.8 were mixed with 200 μM of the compounds or DMSO and incubated at 28°C for 2 h. *Oryza sativa* ssp. *indica* rice cultivar IR24 was

used for pathogenicity assays. On 2-week-old seedlings, bacterial cells of *Xoo* or *Xoc* were inoculated using a needleless syringe. On 2-month-old adult plants, bacterial cells of *Xoo* or *Xoc* were inoculated by the leaf clipping or leaf needling method, respectively. Plants were scored at 3 days post-inoculation (dpi) for symptoms in seedlings, and at 14 dpi for lesion lengths in adult rice plants. Plants were maintained in a glasshouse at 25°C (16 h of light and 8 h of darkness) during the experiments.

For bacterial population assay, the top 20 cm of each leaf was ground in sterilized distilled water and the cell suspensions were plated onto PSA after serial dilutions, and then incubated at 28°C. Bacterial colonies were counted after 72 h. At least 10 leaves were used for each compound treatment, and all experiments were repeated at least three times.

### ACKNOWLEDGEMENTS

We would like to thank Dr Seiji Tsuge for providing us with plasmids pHM04172 and pHM03702. We also thank Dr Fengquan Liu for providing the *Xoc* RS105 strain. This work was supported by grants from the Special Fund for Agro-Scientific Research in the Public Interest of China (201303015), the National Key Project for Basic Research (973 Project, 2015CB150600), National Science Foundation of China (31370160) for CH, Beijing Natural Science Foundation (5142017) for FT, and the Research Growth Initiative of the University of Wisconsin-Milwaukee and the Catalyst Grant in Advanced Automation of UWM Research Foundation for C-HY.

### REFERENCES

- Alfano, J.R. and Collmer, A. (1997) The type III (Hrp) secretion pathway of plant pathogenic bacteria: trafficking harpins, Avr proteins, and death. *J. Bacteriol.* **179**, 5655–5662.
- Andrade, M.O., Farah, C.S. and Wang, N. (2014) The post-transcriptional regulator *rsmA/csrA* activates T3SS by stabilizing the 5' UTR of *hrpG*, the master regulator of *hrp* genes, in *Xanthomonas*. *PLoS Pathog.* **10**, e1003945.
- Barczak, A.K. and Hung, D.T. (2009) Productive steps toward an antimicrobial targeting virulence. *Curr. Opin. Microbiol.* **12**, 490–496.

- Bowlin, N.O., Williams, J.D., Knoten, C.A., Torhan, M.C., Tashjian, T.F., Li, B., Aiello, D., Mecsas, J., Hauser, A.R., Peet, N.P., Bowlin, T.L. and Moir, D.T. (2014) Mutations in the *Pseudomonas aeruginosa* needle protein gene *pscF* confer resistance to phenoxycetamide inhibitors of the type III secretion system. *Antimicrob. Agents Chemother.* **58**, 2211–2220.
- Büttner, D. (2012) Protein export according to schedule: architecture, assembly, and regulation of type III secretion systems from plant- and animal-pathogenic bacteria. *Microbiol. Mol. Biol. Rep.* **76**, 262–310.
- Büttner, D. and Bonas, U. (2003) Common infection strategies of plant and animal pathogenic bacteria. *Curr. Opin. Microbiol.* **6**, 312–319.
- Büttner, D. and Bonas, U. (2010) Regulation and secretion of *Xanthomonas* virulence factors. *FEMS Microbiol. Rev.* **34**, 107–133.
- Charro, N. and Mota, L.J. (2015) Approaches targeting the type III secretion system to treat or prevent bacterial infections. *Expert Opin. Drug Discov.* **10**, 373–387.
- Chatterjee, A., Cui, Y. and Chatterjee, A.K. (2002) Regulation of *Erwinia carotovora* hrpL(Ecc) (sigma-L(Ecc)), which encodes an extracytoplasmic function subfamily of sigma factor required for expression of the HRP regulon. *Mol. Plant–Microbe Interact.* **15**, 971–980.
- Cho, H.J., Park, Y.J., Noh, T.H., Kim, Y.T., Kim, J.G., Song, E.S., Lee, D.H. and Lee, B.M. (2008) Molecular analysis of the *hrp* gene cluster in *Xanthomonas oryzae* pathovar *oryzae* KACC10859. *Microb. Pathog.* **44**, 473–483.
- Cornelis, G.R. (2006) The type III secretion injectisome. *Nat. Rev.* **4**, 811–825.
- Durrant, W.E., and Dong, X. (2004) Systemic acquired resistance. *Annu. Rev. Phytopathol.* **42**, 185–209.
- Das, A., Rangaraj, N. and Sonti, R.V. (2009) Multiple adhesin-like functions of *Xanthomonas oryzae* pv. *oryzae* are involved in promoting leaf attachment, entry, and virulence on rice. *Mol. Plant–Microbe Interact.* **22**, 73–85.
- Felise, H.B., Nguyen, H.V., Pfueter, R.A., Barry, K.C., Jackson, S.R., Blanc, M.P., Bronstein, P.A., Kline, T. and Miller, S.I. (2008) An inhibitor of gram-negative bacterial virulence protein secretion. *Cell Host Microbe*, **4**, 325–336.
- Furutani, A., Tsuge, S., Oku, T., Tsuno, K., Inoue, Y., Ochiai, H., Kaku, H. and Kubo, Y. (2003) Hpa1 secretion via type III secretion system in *Xanthomonas oryzae* pv. *oryzae*. *J. Gen. Plant Pathol.* **69**, 271–275.
- Furutani, A., Takaoka, M., Sanada, H., Noguchi, Y., Oku, T., Tsuno, K., Ochiai, H. and Tsuge, S. (2009) Identification of novel type III secretion effectors in *Xanthomonas oryzae* pv. *oryzae*. *Mol. Plant–Microbe Interact.* **22**, 96–106.
- Garrity-Ryan, L.K., Kim, O.K., Balada-Llasat, J.M., Bartlett, V.J., Verma, A.K., Fisher, M.L., Castillo, C., Songsungthong, W., Tanaka, S.K., Levy, S.B., Mecsas, J. and Aleksun, M.N. (2010) Small molecule inhibitors of LcrF, a *Yersinia pseudotuberculosis* transcription factor, attenuate virulence and limit infection in a murine pneumonia model. *Infect. Immun.* **78**, 4683–4690.
- Gürlebeck, D., Thieme, F. and Bonas, U. (2006) Type III effector proteins from the plant pathogen *Xanthomonas* and their role in the interaction with the host plant. *J. Plant Physiol.* **163**, 233–255.
- Jessen, D.L., Bradley, D.S. and Nilles, M.L. (2014) A type III secretion system inhibitor targets YopD while revealing differential regulation of secretion in calcium-blind mutants of *Yersinia pestis*. *Antimicrob. Agents Chemother.* **58**, 839–850.
- Khokhani, D., Zhang, C., Li, Y., Wang, Q., Zeng, Q., Yamazaki, A., Hutchins, W., Zhou, S.S., Chen, X. and Yang, C.H. (2013) Discovery of plant phenolic compounds that act as type III secretion system inhibitors or inducers of the fire blight pathogen, *Erwinia amylovora*. *Appl. Environ. Microb.* **79**, 5424–5436.
- Kim, S.Y., Kim, J.G., Lee, B.M. and Cho, J.Y. (2009) Mutational analysis of the gum gene cluster required for xanthan biosynthesis in *Xanthomonas oryzae* pv. *oryzae*. *Biotechnol. Lett.* **31**, 265–270.
- Lattanzio, V. (2013) Phenolic compounds: introduction. In: *Handbook of Natural Products* (Ramawat, K.G. and Mérillon, J.-M., eds), pp. 1543–1580. Berlin, Heidelberg: Springer.
- Livak, K. J., and Schmittgen, T. D. (2001) Analysis of relative gene expression data using real-time quantitative PCR and the 2<sup>-</sup>(Delta Delta C(T)) Methods, **25**, 402–408.
- Li, R.F., Lu, G.T., Li, L., Su, H.Z., Feng, G.F., Chen, Y., He, Y.Q., Jiang, B.L., Tang, D.J. and Tang, J.L. (2014) Identification of a putative cognate sensor kinase for the two-component response regulator HrpG, a key regulator controlling the expression of the *hrp* genes in *Xanthomonas campestris* pv. *campestris*. *Environ. Microbiol.* **16**, 2053–2071.
- Li, Y., Peng, Q., Selimi, D., Wang, Q., Charkowski, A.O., Chen, X. and Yang, C.H. (2009) The plant phenolic compound p-coumaric acid represses gene expression in the *Dickeya dadantii* type III secretion system. *Appl. Environ. Microb.* **75**, 1223–1228.
- Li, Y., Hutchins, W., Wu, X., Liang, C., Zhang, C., Yuan, X., Khokhani, D., Chen, X., Che, Y., Wang, Q. and Yang, C.H. (2015) Derivative of plant phenolic compound inhibits the type III secretion system of *Dickeya dadantii* via HrpX/HrpY two-component signal transduction and Rsm systems. *Mol. Plant Pathol.* **16**, 150–163.
- Li, Y.R., Zou, H.S., Che, Y.Z., Cui, Y.P., Guo, W., Zou, L.F., Chatterjee, S., Biddle, E.M., Yang, C.H. and Chen, G.Y. (2011) A novel regulatory role of HrpD6 in regulating *hrp-hrc-hpa* genes in *Xanthomonas oryzae* pv. *oryzicola*. *Mol. Plant–Microbe Interact.* **24**, 1086–1101.
- Marshall, N.C. and Finlay, B.B. (2014) Targeting the type III secretion system to treat bacterial infections. *Expert Opin. Ther. Targets*, **18**, 137–152.
- Miller, W.G., Leveau, J.H. and Lindow, S.E. (2000) Improved *gfp* and *inaZ* broad-host-range promoter-probe vectors. *Mol. Plant–Microbe Interact.* **13**, 1243–1250.
- Nino-Liu, D.O., Ronald, P.C. and Bogdanove, A.J. (2006) *Xanthomonas oryzae* pathovars: model pathogens of a model crop. *Mol. Plant Pathol.* **7**, 303–324.
- Noel, L., Thieme, F., Nennstiel, D. and Bonas, U. (2001) cDNA-AFLP analysis unravels a genome-wide *hrpG*-regulon in the plant pathogen *Xanthomonas campestris* pv. *vesicatoria*. *Mol. Microbiol.* **41**, 1271–1281.
- Rai, R., Ranjan, M., Pradhan, B.B. and Chatterjee, S. (2012) Atypical regulation of virulence-associated functions by a diffusible signal factor in *Xanthomonas oryzae* pv. *oryzae*. *Mol. Plant–Microbe Interact.* **25**, 789–801.
- Rajeshwari, R., Jha, G. and Sonti, R.V. (2005) Role of an in planta-expressed xylanase of *Xanthomonas oryzae* pv. *oryzae* in promoting virulence on rice. *Mol. Plant–Microbe Interact.* **18**, 830–837.
- Rasko, D.A. and Sperandio, V. (2010) Anti-virulence strategies to combat bacteria-mediated disease. *Nat. Rev. Drug Discov.* **9**, 117–128.
- Ray, S.K., Rajeshwari, R. and Sonti, R.V. (2000) Mutants of *Xanthomonas oryzae* pv. *oryzae* deficient in general secretory pathway are virulence deficient and unable to secrete xylanase. *Mol. Plant–Microbe Interact.* **13**, 394–401.
- Salzberg, S.L., Sommer, D.D., Schatz, M.C., Phillippy, A.M., Rabinowicz, P.D., Tsuge, S., Furutani, A., Ochiai, H., Delcher, A.L., Kelley, D., Madupu, R., Puiu, D., Radune, D., Shumway, M., Trapnell, C., Aparna, G., Jha, G., Pandey, A., Patil, P.B., Ishihara, H., Meyer, D.F., Szurek, B., Verdier, V., Koebnik, R., Dow, J.M., Ryan, R.P., Hirata, H., Tsuyumu, S., Won Lee, S., Seo, Y.S., Sriariyanum, M., Ronald, P.C., Sonti, R.V., Van Sluys, M.A., Leach, J.E., White, F.F. and Bogdanove, A.J. (2008) Genome sequence and rapid evolution of the rice pathogen *Xanthomonas oryzae* pv. *oryzae* PXO99<sup>A</sup>. *BMC Genomics*, **9**, 204.
- Sun, Q.H., Hu, J., Huang, G.X., Ge, C., Fang, R.X. and He, C.Z. (2005) Type-II secretion pathway structural gene *xpsE*, xylanase- and cellulase secretion and virulence in *Xanthomonas oryzae* pv. *oryzae*. *Plant Pathol.* **54**, 15–21.
- Sundaram, R.M., Chatterjee, S., Oliva, R., Laha, G.S., Cruz, C.V., Leach, J.E. and Sonti, R.V. (2014) Update on Bacterial Blight of Rice: Fourth International Conference on Bacterial Blight of Rice. *Rice*, **7**, 12.
- Tang, X., Xiao, Y. and Zhou, J.M. (2006) Regulation of the type III secretion system in phytopathogenic bacteria. *Mol. Plant–Microbe Interact.* **19**, 1159–1166.
- Tian, F., Yu, C., Li, H., Wu, X., Li, B., Chen, H., Wu, M. and He, C. (2015) Alternative sigma factor RpoN2 is required for flagellar motility and full virulence of *Xanthomonas oryzae* pv. *oryzae*. *Microbiol. Res.* **170**, 177–183.
- Tsuge, S., Furutani, A., Fukunaka, R., Oku, T., Tsuno, K., Ochiai, H., Inoue, Y., Kaku, H. and Kubo, Y. (2002) Expression of *Xanthomonas oryzae* pv. *oryzae* *hrp* genes in XOM2, a novel synthetic medium. *J. Gen. Plant Pathol.* **68**, 363–371.
- Verdier, V., Triplett, L.R., Hummel, A.W., Corral, R., Cernadas, R.A., Schmidt, C.L., Bogdanove, A.J. and Leach, J.E. (2012) Transcription activator-like (TAL) effectors targeting OsSWEET genes enhance virulence on diverse rice (*Oryza sativa*) varieties when expressed individually in a TAL effector-deficient strain of *Xanthomonas oryzae*. *New Phytol.* **196**, 1197–1207.
- Wang, D., Zetterström, C.E., Gabrielsen, M., Beckham, K.S.H., Tree, J.J., Macdonald, S.E., Byron, O., Mitchell, T.J., Gally, D.L., Herzyk, P., Mahajan, A., Uvell, H., Burchmore, R., Smith, B.O., Elofsson, M. and Roe, A.J. (2011) Identification of bacterial target proteins for the salicylidene acylhydrazide class of virulence-blocking compounds. *J. Biol. Chem.* **286**, 29 922–29 931.
- Wang, X.Y., Song, C.F., Miao, W.G., Ji, Z.L., Wang, X., Zhang, Y., Zhang, J.H., Hu, J.S., Borth, W. and Wang, J.S. (2008) Mutations in the N-terminal coding region of the harpin protein Hpa1 from *Xanthomonas oryzae* cause loss of hypersensitive reaction induction in tobacco. *Appl. Microbiol. Biotechnol.* **81**, 359–369.
- Wengelnik, K. and Bonas, U. (1996) HrpXv, an AraC-type regulator, activates expression of five of the six loci in the *hrp* cluster of *Xanthomonas campestris* pv. *vesicatoria*. *J. Bacteriol.* **178**, 3462–3469.
- Wengelnik, K., Van den Ackerveken, G. and Bonas, U. (1996) HrpG, a key hrp regulatory protein of *Xanthomonas campestris* pv. *vesicatoria* is homologous to two-component response regulators. *Mol. Plant–Microbe Interact.* **9**, 704–712.
- White, F.F. and Yang, B. (2009) Host and pathogen factors controlling the rice–*Xanthomonas oryzae* interaction. *Plant Physiol.* **150**, 1677–1686.

- Xiao, Y., Heu, S., Yi, J., Lu, Y. and Hutcheson, S.W. (1994) Identification of a putative alternate sigma factor and characterization of a multicomponent regulatory cascade controlling the expression of *Pseudomonas syringae* pv. *syringae* Pss61 *hrp* and *hmrA* genes. *J. Bacteriol.* **176**, 1025–1036.
- Yamazaki, A., Li, J., Zeng, Q., Khokhani, D., Hutchins, W.C., Yost, A.C., Biddle, E., Toone, E.J., Chen, X. and Yang, C.H. (2012) Derivatives of plant phenolic compound affect the type III secretion system of *Pseudomonas aeruginosa* via a GacS–GacA two-component signal transduction system. *Antimicrob. Agents Chemother.* **56**, 36–43.
- Yang, F., Korban, S.S., Pusey, P.L., Elofsson, M., Sundin, G.W. and Zhao, Y. (2014) Small-molecule inhibitors suppress the expression of both type III secretion and amylovoran biosynthesis genes in *Erwinia amylovora*. *Mol. Plant Pathol.* **15**, 44–57.
- Yang, S., Peng, Q., San Francisco, M., Wang, Y., Zeng, Q. and Yang, C.H. (2008) Type III secretion system genes of *Dickeya dadantii* 3937 are induced by plant phenolic acids. *PLoS One*, **3**, e2973.
- Zhu, W., MaGbanua, M.M. and White, F.F. (2000) Identification of two novel *hrp*-associated genes in the *hrp* gene cluster of *Xanthomonas oryzae* pv. *oryzae*. *J. Bacteriol.* **182**, 1844–1853.
- Zou, L.F., Wang, X.P., Xiang, Y., Zhang, B., Li, Y.R., Xiao, Y.L., Wang, J.S., Walmsley, A.R. and Chen, G.Y. (2006) Elucidation of the *hrp* clusters of *Xanthomonas oryzae* pv. *oryzicola* that control the hypersensitive response in nonhost tobacco and pathogenicity in susceptible host rice. *Appl. Environ. Microbiol.* **72**, 6212–6224.

## SUPPORTING INFORMATION

Additional Supporting Information may be found in the online version of this article at the publisher's website:

**Fig. S1** DNA sequence of *hpa1* gene and its promoter region. Sequences for a plant-inducible promoter (PIP)-box, start codon, stop codon and the primers used for amplification of the promoter region are indicated in bold.

**Fig. S2** Effects of TS006 on promoter activity of *hpa1* at different concentrations. The growth of *Xanthomonas oryzae* pv. *oryzae* (*Xoo*) in XOM2 supplemented with different concentrations of TS006 was recorded by measuring the absorbance of the

bacterial suspension at 600 nm. DMSO, dimethylsulfoxide; OD, optical density.

**Fig. S3** Effects of various compounds on bacterial survival. Bacterial cells were incubated with dimethylsulfoxide (DMSO) or various compounds at a concentration of 200  $\mu\text{M}$  for 2 h at 28°C before being plated on a peptone sucrose agar (PSA) plate after serial dilutions. Three independent tests were performed with similar results. CFU, colony-forming unit.

**Fig. S4** Sequence alignment of the promoter regions of *hpa1* (A), *hrcC* (B) and *hrpB* (C) between *Xanthomonas oryzae* pv. *oryzae* (*Xoo*) and *Xanthomonas oryzae* pv. *oryzicola* (*Xoc*).

**Fig. S5** Effects of TS006, TS010, TS015 and TS018 on the bacterial population of *Xanthomonas oryzae* pv. *oryzae* (*Xoo*) PXO99<sup>A</sup> in leaves of IR24. Bacterial populations were measured in 20-cm leaf segments at 14 days after inoculation. CFU, colony-forming unit; DMSO, dimethylsulfoxide.

**Fig. S6** Effects of various compounds on bacterial growth rates. (A) Growth rate of *Xanthomonas oryzae* pv. *oryzicola* (*Xoc*) RS105 in rich medium (NB, nutrient broth) supplemented with dimethylsulfoxide (DMSO) or 200  $\mu\text{M}$  of TS006, TS010, TS015 or TS018. (B) Growth rate of *Xoc* RS105 in *hrp*-inducing medium XOM2 (plus 0.5% sucrose) supplemented with DMSO or 200  $\mu\text{M}$  of TS006, TS010, TS015 or TS018. The optical density at 600 nm ( $\text{OD}_{600}$ ) of the culture suspensions was measured every 6 h during the 72-h period. Two independent tests were performed with similar results.

**Table S1** Phenolic compounds precipitated when added to XOM2.

**Table S2** Primers used in this study.