

Identification of Potential Prostate Cancer Preventive Agents through Induction of Quinone Reductase *in Vitro*¹

James D. Brooks, Michael F. Goldberg, Linda A. Nelson, David Wu, and William G. Nelson²

Department of Urology, Stanford University School of Medicine [J. D. B.], and the Departments of Urology [M. G., D. W., W. G. N.], Oncology [W. G. N.], Pharmacology [W. G. N.], and Medicine [W. G. N.], The Johns Hopkins University School of Medicine, Baltimore, Maryland 21321-1000

Abstract

Human prostate cancer is characterized by an early and near-universal loss of expression of the phase 2 enzyme glutathione *S*-transferase- π (GSTP1). We hypothesize that a mechanism-based prostate cancer preventive strategy could involve induction of phase 2 enzymes within the prostate to compensate for the loss of GSTP1 expression. NAD[P]H:(quinone-acceptor) oxidoreductase (quinone reductase or QR) enzymatic activity, a surrogate of phase 2 enzyme response, was measured after treating the human prostate cancer cell line LNCaP with known phase 2 enzyme-inducing agents from 10 distinct chemical classes. QR enzymatic activity was assayed in microtiter plates using the menadione-coupled reduction of tetrazolium dye. Degree of induction was expressed as fold-increase over control and corrected for toxicity. Compounds were also tested in LNCaP-5-aza-C, an LNCaP subline selected in 5-aza-cytidine that expresses GSTP1, and in the human liver cell line HepG2. LNCaP showed robust induction of QR enzymatic activity after treatment with a subset of the phase 2 enzyme-inducing agents. All Michael acceptors were effective at inducing QR activity in LNCaP. Some phenolic antioxidants, heavy metal salts, and quinones also significantly increased QR activity, although inducer potency varied widely within these classes of compounds. Some of the isothiocyanates, mercaptans, bifunctional inducers, and trivalent arsenicals also produced modest QR induction, but peroxides and dithiolethiones were inactive. LNCaP-5-aza-C and LNCaP responded similarly to all compounds, but the pattern of response for HepG2 differed significantly. The differences in QR responsiveness between the prostate cell lines and HepG2 suggest that prostate tissues may have a unique pattern

of response to phase 2-inducing agents distinct from other tissue types. Our data suggest that measurement of QR induction in prostate cancer cell lines may help identify potential cancer chemopreventive agents effective in the prostate.

Introduction

Prostate cancer is the most commonly diagnosed noncutaneous malignancy and second leading cause of cancer death in American men (1). One striking feature of this disease is the tremendous disparity in incidence and mortality rates worldwide. In contrast to Western industrialized nations, prostate cancer is rarely diagnosed and contributes little to cancer mortality in Asia (2, 3). Migration studies suggest that lifestyle and/or the environment are important determinants of prostate cancer pathogenesis. Men who emigrate from Asia to the United States acquire higher rates of prostate cancer, and subsequent generations of American-born Asian men retain this elevated risk (4–7). Although the environmental factors responsible for this change in risk are unknown, this observation suggests that lifestyle changes may prevent the development of prostate cancer or slow the progression of the disease. The development of preventive intervention strategies has become particularly pressing because large cohorts of men are identified who are at increased risk for prostate cancer, including African-Americans, those with a family history of prostate cancer, and men carrying genetic markers associated with prostate cancer risk (8–12).

The ideal prostate cancer preventive strategy has not been defined. Antiproliferative agents, compounds that induce differentiation, and drugs that alter the androgen milieu of the prostate have all been proposed as potential preventive approaches and are currently being evaluated in clinical trials (13). Another possible strategy, yet untested in prostate cancer, involves induction of enzymes of carcinogen defense (phase 2 enzymes), thereby buttressing the innate defenses of the prostate cell to slow accumulation of genetic alterations responsible for the development and progression of the disease. We have collected provocative evidence that such a strategy may be particularly relevant to prostatic carcinogenesis. Virtually all human prostate cancer cancers, regardless of grade or stage, lack expression of the phase 2 enzyme GSTP³ (14–16). This loss of expression is associated with extensive methylation of deoxycytidine residues in the 5'-regulatory regions of the *GSTP1* gene. Intriguingly, this alteration appears to be an early event in prostatic carcinogenesis in that it can be found in prostatic intraepithelial neoplasia, a purported prostate cancer

Received 11/18/02; revised 5/8/02; accepted 5/16/02.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

¹ Supported by NIH Grants AT00437, CA58236, and CA70196 (to W. G. N.), the Doris Duke Foundation (to J. D. B.), Department of Defense Grant DAMD17-98-1-8555 (to J. D. B.), and the American Foundation for Urologic Disease (to J. D. B.).

² To whom requests for reprints should be addressed, at The Johns Hopkins University School of Medicine, Room 151, Bunting-Blaustein Cancer Research Building, 1650 Orleans Street, Baltimore, MD 21321-1000. Phone (410) 614-1661; Fax (410) 502-9817.

³ The abbreviations used are: GSTP, glutathione *S*-transferase π ; QR, quinone reductase; I_{max} , maximum fold-induction for each of the compounds compared with DMSO-treated controls; EC_{100} , concentration of each compound at maximum induction; IC_{50} , concentration toxic to 50% of cells.

precursor lesion (17). Mice genetically lacking *GSTP1* have increased susceptibility to DMBA/TPA-induced skin cancer, suggesting that loss of this enzyme in itself can contribute to carcinogenesis (18).

A large body of evidence suggests that induction of phase 2 enzymes, and in particular the glutathione transferases, will prevent carcinogen-induced tumors in a number of species (19, 20). NAD[P]H:(quinone-acceptor) oxidoreductase or QR, a cytosolic FAD-dependent flavoprotein, is induced coordinately with the glutathione transferases and has served as a surrogate marker of phase 2 enzyme responsiveness *in vivo* and *in vitro* (21–25). QR protects cells against quinones and highly reactive semiquinones by catalyzing an obligate two-electron reduction of quinones to hydroquinones (26). In the prostate, QR has been shown to protect against formation of mutagenic 4-catechol-estrogen DNA adducts in Noble rats (27). *In vitro* methods have been devised to rapidly screen agents for QR induction and have been used to identify synthetic and diet-derived candidate chemopreventive agents (21–23). Several of these compounds have later been shown to prevent carcinogenesis in animal models (28, 29).

Previous *in vitro* screens of phase 2 enzyme-inducing compounds have usually been carried out using the Hepa1c1c7 murine hepatoma cell line. Although this cell line has documented utility in the identification of novel agents, it is unknown whether the responses observed in this cell line can be extrapolated to other tissue or cell types, to responses *in vivo*, or to other species. Because human prostate cancer selectively lacks *GSTP1* expression, we hypothesize that compounds able to induce phase 2 enzyme activity within prostate epithelial cells may hold promise as prostate cancer preventive agents. To evaluate the possibility of phase 2 enzyme induction in human prostatic cells, we screened a diverse set of 55 compounds for their ability to induce QR enzymatic activity in the human prostate cancer cell line LNCaP. Compounds evaluated include monofunctional inducers (Michael reaction acceptors, diphenols, quinones, isothiocyanates, peroxides, azo dyes, and heavy metals), bifunctional inducers, as well as other putative cancer preventive agents. Induction of QR activity was also assayed in an LNCaP subline (LNCaP-5-aza-C) that expresses *GSTP1* and in the human hepatoblastoma cell line HepG2. Measurement of toxicity of agents for each of the cell lines was carried out in parallel plates treated identically.

Materials and Methods

Reagents. Vinylene trithiocarbonate, 1,2-dithiol[4,3-*c*]-1,2-dithiole-3,6-dithione, dimethyl fumarate, dimethyl maleate, 1-nitro-1-cyclohexene, phenyl isothiocyanate, phenethyl isothiocyanate, benzyl isothiocyanate, chalcone, perillyl alcohol, and selenium were obtained from Aldrich Chemical Co. (Milwaukee, WI). Linomide and a related compound, 2,4-quinolinediol, were a gift of Dr. John T. Isaacs (Johns Hopkins Oncology Center). All other compounds were purchased from Sigma Chemical Co. (St. Louis, MO).

Cell Culture. The human prostate cancer cell line LNCaP and human hepatoblastoma cell line HepG2 were obtained from American Type Culture Collection. The LNCaPazaC cell line was derived from selection of the LNCaP cell line in 5 μ M 5-aza-cytidine, a noncompetitive inhibitor of DNA methyltransferase, and stably expresses the *GSTP* enzyme (16). LNCaP and LNCaP-5-aza-C cell lines were cultured in 96-well plates at a density of 10,000 cells/well in 200 μ l of RPMI 1640 and grown in a humidified incubator in 5% CO₂ at 37°C. HepG2 cells were plated at a density of 4000 cells/well and

grown similarly. The following day, the medium was aspirated and replaced with RPMI 1640 supplemented with 10% charcoal-stripped FCS, 100 units/ml penicillin G, 100 μ g/ml streptomycin, and 0.1% DMSO. Test compounds were dissolved in DMSO and diluted in the medium such that the concentration of DMSO did not exceed 0.1%. Two-fold serial dilutions of each compound were made in the microtiter plates so that an entire row (eight wells) represented a single concentration of the compound. One row treated with DMSO alone served as a control, and another row containing only medium was used as a blank in absorbance determinations. After 48 h of exposure to each compound, plates were assayed for quinone reductase activity.

Quinone Reductase Assays. Quinone reductase activity was assessed by the menadione-coupled reduction of tetrazolium dye as modified from Prochaska and Santamaria (30). Medium was gently aspirated, and the cells were lysed by incubation at 37°C with 50 μ l of 0.08% digitonin and 2 mM EDTA (pH 7.8) with gentle agitation for 30 min. During this incubation, a stock solution was prepared by combining 16.7 mg of BSA, 7.5 mg of 3-(4,5-dimethylthiazo-2-yl)-2,5-diphenyltetrazolium bromide, 0.6 mg of NADP, 1.25 ml of 0.5 M Tris HCl (pH 7.4), 166.7 μ l of 1.5% Tween 20, 166.7 μ l of 150 mM glucose 6-phosphate, 16.7 μ l of 7.5 mM FAD, 50 units of yeast glucose 6-phosphate dehydrogenase, and distilled water to a final volume of 25 ml for each plate to be assayed. Immediately before use, 25 μ l of 50 mM menadione dissolved in acetonitrile were added to this stock solution. Two hundred μ l of the complete stock solution was added simultaneously to the cell lysate in all 96 wells of the plate. Plates were immediately placed in a Tecan 96-well plate automated optical scanner, and readings at 610 nm were taken every 30 s. In virtually all instances, a change in absorbance attributable to the formation of blue-brown reduced tetrazolium dye was linear for well over 5 min; therefore, a single reading at 5 min was used for all compounds as described by Prochaska *et al.* (21).

Toxicity Assessment. Toxicity of the compounds was assessed in parallel plates treated identically to those used in assays for quinone reductase activity (30). After 48 h of exposure to each compound, cells were fixed with methanol and stained with 0.5% crystal violet for 5 min. Plates were then washed with distilled water and allowed to air dry overnight. Bound dye was dissolved with 200 μ l/well of 1% SDS, and the plates were scanned at 610 nm.

Inducer Potency. QR activity, in arbitrary units, was calculated automatically from the mean activity for all eight wells at each concentration for each compound using software developed in our laboratory. Activity was corrected for toxicity at each concentration as described (30). Inducer potency (fold-induction of QR activity) was expressed as the ratio of corrected QR activity for treated cells to corrected QR activity for the vehicle controls.

Results

QR Induction in LNCaP. Phase 2 enzyme-inducing agents comprise a chemically diverse set of compounds and have been demonstrated to prevent carcinogen-induced tumors in a variety of model systems. To characterize the phase 2 enzyme responsiveness of the human prostate cancer cell line LNCaP, we measured QR activity after treatment with 34 different phase 2-inducing agents from 10 distinct chemical classes. Compounds were selected because of their ability to induce phase 2 enzyme activity in Hepa1c1c7 murine hepatoma cells or another model system.

Table 1 Quinone reductase inducer potency and toxicity of compounds of diverse chemical classes in LNCaP cells

| Compound | Dose range (μM) | I_{Max} Mean \pm SD | P^a | EC_{100} (μM) | IC_{50} (μM) |
|--|------------------------------|--------------------------------|------------|-------------------------------------|------------------------------------|
| Bifunctional inducers (PAHs) | | | | | |
| Benzo(a)pyrene | 0.040–200 | NI ^b | | NI | >200 |
| 20-Methylcholanthrene | 0.78–200 | NI | | NI | >200 |
| β -naphthoflavone | 0.008–500 | NI | | NI | 500 \pm 150 |
| Sudan 1 | 0.020–500 | 1.59 \pm 0.169 | .00008 | 25 | 260 \pm 62 |
| Sudan 2 | 0.20–50 | NI | | NI | >50 |
| Sudan 3 | 0.20–500 | 1.28 \pm 1.89 | .0008 | 50 | >500 |
| 1-[2-Thiazolylazo]-2-naphthol | 0.20–100 | NI | | NI | 13 \pm 2 |
| 1-[2-Pyridoylazo]-2-naphthol | 0.20–50 | NI | | NI | 7.5 \pm 2 |
| Isothiocyanates | | | | | |
| Benzyl isothiocyanate | 0.20–50 | 1.10 \pm 0.096 | .001 | 1.56 | 3.1 \pm 0.5 |
| Phenyl isothiocyanate | 2.0–500 | NI | | NI | 170 \pm 24 |
| Phenethyl isothiocyanate | 0.20–50 | 1.19 \pm 0.143 | .005 | 3.125 | 6 \pm 0.9 |
| Phenolic antioxidants | | | | | |
| Butylated hydroxyanisole | 0.78–200 | 1.17 \pm 0.072 | .009 | 25 | 120 \pm 60 |
| Butylated hydroxytoluene | 2.0–500 | NI | | NI | 300 \pm 100 |
| Catechol | 0.78–200 | 2.14 \pm 0.279 | .000001 | 12.5 | 75 \pm 6 |
| Resorcinol | 0.20–5000 | NI | | NI | 5000 \pm 900 |
| Heavy metal salts | | | | | |
| CdCl ₂ | 0.20–50 | 1.52 \pm 0.213 | .002 | 3.25 | 10 \pm 2.5 |
| HgCl ₂ | 0.20–50 | 5.54 \pm 0.235 | | 50 | 50 \pm 12 |
| ZnCl ₂ | 2.0–500 | NI | .00000001 | NI | 150 \pm 16 |
| Peroxides | | | | | |
| Cumene hydroperoxide | 0.39–100 | NI | | NI | 25 \pm 5 |
| Hydrogen peroxide | 2.0–500 | NI | | NI | 70 \pm 17 |
| <i>tert</i> -Butyl hydroperoxide | 0.39–100 | NI | | NI | 13.5 \pm 5 |
| Mercaptans | | | | | |
| 1,2-Ethanedithiol | 2.0–500 | 1.27 \pm 0.109 | .0006 | 31.25 | 70 \pm 14 |
| Michael acceptors | | | | | |
| Coumarin | 0.20–5000 | 1.22 \pm 0.152 | .003 | 50 | 2500 \pm 1100 |
| Dimethyl maleate | 2.0–500 | 2.46 \pm 0.312 | .000005 | 62.5 | 100 \pm 16 |
| Dimethyl fumarate | 0.78–200 | 2.05 \pm 0.151 | .000000007 | 80 | 100 \pm 23 |
| α -Methylene- γ -butyrolactone | 0.20–50 | 2.20 \pm 0.203 | .0000003 | 25 | 30 \pm 1.4 |
| 1-Nitro-1-cyclohexene | 0.20–50 | 1.47 \pm 0.196 | .002 | 3.125 | 7 \pm 1.1 |
| Quinones | | | | | |
| Hydroquinone | 0.20–50 | 3.54 \pm 0.431 | .0000009 | 12.5 | >50 |
| Ethoxyquin | 0.20–500 | NI | | NI | 70 \pm 7 |
| Dithiolethiones | | | | | |
| [1,2]Dithiolo-dithiole-dithione | 2.0–500 | NI | | NI | 125 \pm 11 |
| Vinylene trithiocarbonate | 2.0–500 | NI | | NI | >500 |
| Trivalent arsenicals | | | | | |
| Phenylarsine oxide | 0.008–50 | 1.54 \pm 0.256 | .0008 | 0.125 | 0.13 \pm 0.03 |
| Sodium <i>m</i> -arsenite | 0.040–10 | NI | | NI | 3.5 \pm 0.5 |

^a QR activity of treated cells compared with vehicle-treated controls by a two-tailed Student's *t* test.

^b NI, not induced.

LNCaP readily responded to several phase 2 enzyme-inducing agents. Table 1 shows I_{Max} , EC_{100} , and IC_{50} . A typical induction profile is shown in Fig. 1. All Michael acceptors, particularly dimethyl maleate, dimethyl fumarate, and methylene butyrolactone, reliably produced significant induction of QR activity in LNCaP. No other chemical class universally produced robust QR induction in LNCaP, and responses to individual members of each class varied widely. For instance, catechol was the only phenolic antioxidant to robustly increase QR activity; hydroquinone and HgCl₂ were also the only compounds in their classes to induce QR. Two of the bifunctional inducers (planar aromatic hydrocarbons, known to induce both phase 1 and phase 2 enzymes) produced modest induction in LNCaP as did the isothiocyanates, peroxides, mercaptans and trivalent arsenicals. Somewhat surprisingly, the dithiolethiones failed to induce quinone reductase altogether.

QR Induction in LNCaP by Other Cancer Chemopreventive Agents. We evaluated whether 21 compounds implicated as potential chemopreventive agents could influence phase 2

enzymatic activity in LNCaP (Table 2). Epigallocatechin was the only tea catechin to produce slight QR induction in LNCaP at near-toxic doses. Two selenium compounds, sodium selenite and selenium dioxide, produced modest elevation of QR activity at concentrations approaching their IC_{50} for LNCaP. Of the remaining diverse set of compounds, quercetin (1.66-fold) and chalcone (1.44-fold) produced modest increases in QR activity, whereas curcumin and *para*-coumaric acid produced more significant levels of induction at micromolar doses. Curcumin-treated cells showed QR induction over baseline starting at 6.25 μM that peaked at 2.01-fold at 25 μM . Curcumin was toxic at slightly higher doses (IC_{50} , 50 μM). *para*-Coumaric acid produced QR induction in LNCaP that began at 62.5 μM and increased linearly with dose. *para*-Coumaric acid was not toxic to LNCaP, even at high concentrations (IC_{50} , >1000 μM).

Induction Patterns in LNCaP Cell Lines Differ from HepG2. Phase 2 enzyme response has been reported to vary significantly between different species and between different tissue types in single animals (28, 31–35). Because LNCaP

Fig. 1. QR induction over a range of concentrations for LNCaP, LNCaP-5-aza-C, and HepG2 after treatment with HgCl₂. Similar induction profiles were generated for the 55 compounds screened in the three cell lines. Toxicity profiles were plotted similarly. Bars, SD.




Table 2 Inducer potency and toxicity of putative preventive agents in LNCaP cells

| Compound | Dose range (μM) | I _{Max} Mean ± SD | P | EC ₁₀₀ (μM) | IC ₅₀ (μM) |
|---------------------------|-----------------|----------------------------|-----------|------------------------|-----------------------|
| Catechins | | | | | |
| Catechin | 2.0–500 | NI | | NI | >500 |
| Epicatechin | 2.0–500 | NI | | NI | >500 |
| Epicatechin gallate | 2.0–500 | NI | | NI | 230 ± 74 |
| Epigallocatechin | 2.0–500 | 1.26 ± 0.287 | .0005 | 62.5 | 80 ± 16 |
| Epigallocatechin gallate | 2.0–500 | NI | | NI | 95 ± 40 |
| Selenium compounds | | | | | |
| Selenium dioxide | 0.20–50 | 1.59 ± 0.132 | .0002 | 6.25 | 7.5 ± 0.8 |
| Selenium | 0.20–50 | NI | | NI | 40 ± 13 |
| Selenocystamine | 0.20–50 | NI | | NI | 11 ± 3 |
| Selenocysteine | 0.20–50 | NI | | NI | 14 ± 4 |
| Selenomethionine | 0.20–50 | NI | | NI | >50 |
| Selenium sulfate | 0.20–50 | NI | | NI | >50 |
| Sodium selenate | 0.20–50 | NI | | NI | 35 ± 13 |
| Sodium selenite | 0.20–50 | 1.48 ± 0.339 | .005 | 3.125 | 5.5 ± 0.8 |
| Others | | | | | |
| Perillyl alcohol | 2.0–500 | NI | | NI | >500 |
| Quercetin | 2.0–500 | 1.66 ± 0.171 | .0005 | 15.63 | 62.5 ± 13 |
| Chalcone | 2.0–500 | 1.44 ± 0.118 | .000002 | 31.25 | 62 ± 12 |
| Lycopene | 1.0–250 | NI | | NI | >250 |
| Curcumin | 0.20–50 | 2.01 ± 0.239 | .00002 | 25 | 50 ± 9 |
| Limonene | 2.0–500 | NI | | NI | 400 ± 180 |
| Linomide | 0.20–50 | NI | | NI | 3.9 ± 1.8 |
| 2,4-Quinolinediol | 2.0–500 | NI | | NI | >500 |
| para-Coumaric acid | 3.91–1000 | 2.28 ± 0.171 | .00000008 | 1000 | >1000 |

^a Abbreviations given in Table 1.

displayed a spectrum of induction to the 34 compounds that differed from that reported for Hepalcl7, we wondered whether those differences were attributable to their species of origin, to their tissue of origin, or to both. In addition, because normal prostatic epithelial cells express GSTP1, we were curious whether reexpression of GSTP1 in LNCaP would affect QR induction (14). Therefore, we evaluated the degree and pattern of QR response to all 55 compounds we had tested in LNCaP in the human hepatoblastoma cell line HepG2 and in an LNCaP cell line selected in 5-aza-cytidine (LNCaP-5-aza-C)

which, unlike the parent cell line, expresses the phase 2 enzyme GSTP (GSTP1; Ref. 16). In both HepG2 and LNCaP-5-aza-C, all compounds were tested over the range of concentrations listed for LNCaP in Table 1. Toxicity measurements were carried out in parallel plates handled identically.

Depicted in Fig. 2 are the 35 compounds that produced a QR response in at least one of the three cell lines. The remaining 20 compounds failed to generate significant QR induction in any of the cell lines and are not shown. Although there were some minor quantitative differences in response between

| | LNCaP | LNCaPazaC | HepG2 |
|--------------------------|------------------|------------------|------------------|
| 20-Methylcholanthrene | NI | NI | 1.32±0.20 |
| Butylated hydroxyanisole | 1.17±0.072 | 1.55±0.13 | 1.87±0.21 |
| Butylated hydroxytoluene | NI | NI | 1.22±0.09 |
| Benzyl isothiocyanate | 1.10±0.096 | NI | 1.37±0.10 |
| Beta-naphthoflavone | NI | NI | 4.98±1.98 |
| Benzo[a]pyrene | NI | NI | 1.62±0.13 |
| Catechol | 2.14±0.28 | 2.47±0.38 | 1.30±0.13 |
| CdCl ₂ | 1.52±0.21 | 1.40±0.18 | 1.21±0.10 |
| Chalcone | 1.44±0.12 | 1.60±0.43 | 1.41±0.19 |
| Coumarin | 1.22±0.15 | 1.31±0.10 | 1.42±0.13 |
| Curcumin | 2.01±0.24 | 2.48±0.12 | 1.74±0.24 |
| Dimethyl fumarate | 2.05±0.15 | 3.76±0.63 | 1.78±0.18 |
| Dimethyl maleate | 2.46±0.31 | 3.24±0.39 | 2.25±0.07 |
| Epicatechin | NI | 1.36±0.16 | NI |
| 1,2-Ethanedithiole | 1.27±0.11 | 1.18±0.13 | NI |
| Epigallocatechin | 1.26±0.29 | NI | NI |
| Ethoxyquin | NI | 1.53±0.15 | 1.38±0.09 |
| HgCl ₂ | 5.54±0.24 | 3.69±0.24 | 1.9±0.22 |
| Hydroquinone | 3.54±0.43 | 5.66±0.93 | 2.12±0.17 |
| Methylene butyrolactone | 2.20±0.20 | 3.61±0.51 | 1.75±0.14 |
| Nitrocyclohexene | 1.47±0.20 | 1.80±0.19 | 1.26±0.12 |
| Pyridoylazo-2-naphthol | NI | NI | 1.70±0.37 |
| Phenylarsine oxide | 1.54±0.26 | 2.00±0.28 | 1.23±0.11 |
| Para-coumaric acid | 2.26±0.17 | 2.02±0.15 | 1.14±0.11 |
| Phenethyl isothiocyanate | 1.19±0.14 | NI | 1.32±0.23 |
| Quercetin | 1.66±0.17 | 1.59±0.22 | 1.69±0.13 |
| Sodium m-arsenite | NI | NI | 1.43±0.10 |
| Selenium dioxide | 1.89±0.13 | 1.54±0.19 | NI |
| Sodium selenite | 1.48±0.34 | NI | NI |
| Sudan 1 | 1.59±0.17 | 1.90±0.11 | 2.63±0.19 |
| Sudan 2 | NI | NI | 2.55±0.28 |
| Sudan 3 | 1.28±0.19 | 1.29±0.11 | 1.87±0.12 |
| Thiazolylazo-2-naphthol | NI | NI | 1.47±0.15 |
| ZnCl ₂ | NI | 1.69±0.31 | 1.46±0.12 |

Fig. 2. I_{Max} s for LNCaP, LNCaP-5-aza-C, and HepG2 for 35 compounds effective in at least one of the cell lines. All values listed show significant induction of QR in treated cells compared with vehicle-treated control at $P < 0.05$ by a two-tail Student's t test. An additional 20 compounds that had no effect in any of the three cell lines are not shown. Differences in the patterns of response are highlighted with darker grays representing greater QR induction. The pattern of response is similar between the prostate cell lines and contrasts sharply with HepG2.

LNCaP and LNCaP-5-aza-C (e.g., HgCl₂, methylene butyrolactone), in most cases QR responsiveness in the GSTP1-expressing LNCaPazaC cell line was virtually identical to the parental cell line. QR induction in HepG2 differed significantly from the prostate cell lines for most of the compounds tested. HepG2 responded robustly to bifunctional inducing agents including β -naphthoflavone, benzo(a)pyrene, and the azo dyes, whereas the prostate cell lines usually responded meagerly to these agents or failed to respond altogether. Response to most monofunctional inducers, although present, was often blunted somewhat in HepG2 compared with the prostate cell lines. Toxicity profiles and IC₅₀ levels differed little between the prostate and liver cell lines (not shown). Thus, there appear to be significant qualitative and quantitative differences in the pattern of QR response between cells derived from different tissues.

Discussion

The human prostate cancer cell line LNCaP appears to be an excellent model for identifying potential prostate cancer preventive agents that act through induction of phase 2 enzymes. LNCaP expresses QR, possesses QR enzymatic activity, and has the capacity to respond to phase 2 enzyme-inducing agents. Because sulforaphane induces several phase 2 enzymes and glutathione synthetic pathways in LNCaP, QR appears to be a valid surrogate of phase 2 enzyme activity in this cell line (36).

Reexpression of GSTP1 by selection with 5-azacytidine did little to alter the pattern or degree of QR responsiveness to chemically diverse compounds. Phase 2 enzyme response in LNCaP and LNCaP-5-aza-C did differ significantly from that of HepG2 and that reported for the murine hepatoma cell line Hepa1c1c7. In part, these differences may be attributable to their tissue of origin, or, for Hepa1c1c7, their species of origin, particularly because rodent cells are more labile in their phase 2 enzyme response than human cells (28, 31–35). Furthermore, the carcinogen *N*-OH-2-amino-1-methyl-6-phenylimidazo[4,5-*b*]pyridine can be activated directly in prostate cancer cell lines, and reexpression of GSTP1 will prevent this activation (37). Therefore, preventive agents that act through induction of phase 2 enzymes may be particularly relevant to human prostate cancer prevention and should be tested for efficacy in human prostatic cell lines. It should be noted, however, that toxic compounds, such as HgCl₂ and some azo dyes, can also induce QR activity. Additional work will be necessary to test whether agents effective in prostate cells *in vitro* are safe and will also produce phase 2 enzyme induction *in vivo*.

The LNCaP cell lines showed a distinct pattern of QR response to monofunctional inducers of several chemical classes. LNCaP and LNCaP-5-aza-C showed robust QR induction when treated with classic Michael reaction acceptors including dimethyl fumarate, dimethyl maleate, and methylene butyrolactone, suggesting this class of compounds may hold promise as prostate cancer preventive agents. Within other chemical classes, QR induction in the LNCaP cell lines was more varied. Both cell lines displayed significant QR induction when treated with catechol, HgCl₂, and hydroquinone but little or no induction when treated with other members of these chemical classes. Similarly, sulforaphane will produce vigorous QR induction in LNCaP; yet we observed very little response to other isothiocyanates in this study (36). We were surprised that dithiolethiones failed to induce QR in LNCaP, particularly because they are effective in other *in vivo* and *in vitro* model systems and ongoing clinical trials in liver cancer with these agents (38, 39). Our findings raise questions whether dithiolethiones would be effective as prostate cancer chemopreventive agents.

The factors underlying the unique patterns of response in LNCaP and the other cell lines are unknown. Phase 2 enzyme response is regulated transcriptionally by Nrf2 binding at antioxidant response enhancer elements (40). Treatment with phase 2 enzyme-inducing agents activates mitogen-activated protein kinase, protein kinase C, and phosphatidylinositol 3-kinase pathways that lead to release of Nrf2 from Keap1 in the cytoplasm, translocation of Nrf2 to the nucleus, and binding together with Maf to antioxidant response enhancers (41). The induction capacity of any compound will be influenced by its ability to stimulate thiol-dependent sensors in the cytoplasm, a process that depends on the biochemical milieu of the cell. Spencer *et al.* (42) and Talalay *et al.* (43) have proposed that the capacity of a compound to induce phase 2 enzyme expression is directly related to its ability to act as a Michael acceptor. For heavy metal salts, inductive capacity parallels their affinity for sulfhydryl groups (44). Therefore, compounds active in LNCaP may be more prone to exist as Michael acceptors or other chemical species with high affinity for thiol groups (45). Zhang and colleagues (46–48) have reported that phase 2 enzymatic induction by isothiocyanates parallels the accumulation of glutathione conjugates intracellularly, and that this accumulation can be affected by GSH concentration and glutathione transferase activity. We did not observe a direct relationship between GSTP1 expression and inducer potency between the

LNCaP and LNCaP-5-aza-C cell lines, suggesting that factors beyond glutathione transferase activity may account for phase 2 enzyme responsiveness in these prostatic cell lines.

The ability of a compound to act as a phase 2 enzyme-inducing agent may also depend on the unique profile of gene expression in each cell line. Large-scale gene expression profiling has demonstrated that cell lines possess unique gene expression patterns that retain many features of their tissue of origin (49). These findings suggest that the response to chemopreventive agents observed *in vitro* may parallel their effects *in vivo*. The expression data also highlight that the response to any compound will depend upon genes expressed in the cell line in which it is tested. Indeed, the differences in QR response we observed in LNCaP and HepG2 could be attributable to the differences in the genes they express. For instance, LNCaP cells showed little QR induction after treatment with bifunctional inducing agents, whereas these agents were the most potent QR inducers in HepG2. Bifunctional inducers require conversion by phase 1 enzymes into oxidized metabolites that then induce phase 2 enzymatic activity (43). We have observed previously that LNCaP cells are unable to activate the heterocyclic amine PhIP (2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine) by *N*-hydroxylation into carcinogenic *N*-OH-PhIP, suggesting low or absent phase 1 enzymatic activity (37). Thus, one possible explanation for the meager QR induction in LNCaP cells in response to bifunctional inducers is that they do not express the enzymes necessary to metabolize the compounds into QR-inducing agents. In addition to this difference between LNCaP and HepG2, we suspect that the unique patterns of response to diverse phase 2 enzyme-inducing agents in different cell lines may be attributable to other poorly characterized differences in gene expression, such as differences in the pattern of expression of thiol-dependent sensing proteins and cell line-specific expression of metabolic enzymes and signaling pathways. Gene expression profiling and proteomics will help define the molecular underpinnings of the phase 2 enzyme response in different tissues (50).

Our limited survey of candidate chemopreventive agents for QR induction in LNCaP demonstrates the potential for this model system in identifying novel agents for use in prostate cancer. Curcumin reliably produced robust induction of QR at micromolar doses in the prostate cells. The potency of curcumin in LNCaP undoubtedly relates to its ability to act as a classic Michael acceptor (51). Curcumin is also intriguing as a prostate cancer preventive agent because it possesses anti-inflammatory effects and inflammation, and free radical generation has been implicated in prostatic carcinogenesis (52–54). Curcumin can also inhibit cyclooxygenase-2 and inducible nitric oxide synthase, and clinical trials are under way to evaluate cyclooxygenase-2 inhibitors as prostate cancer preventive agents (55). Although there is some debate about curcumin, its ability to quench free radicals as well as induce phase 2 enzymes make it attractive as a prostate cancer preventive agent (56, 57).

The flavonoids quercetin and chalcone both produced modest induction in QR at micromolar doses. Both are distributed widely in plants and have been shown to act as phase 2 enzyme-inducing agents in other systems (58, 59). The ability of these compounds to induce QR in prostatic cells may help explain the observed inverse correlation between vegetable consumption and prostate cancer risk (60–63). Other epidemiological studies have noted an inverse correlation between serum selenium levels and prostate cancer risk, and we observed modest induction of QR by selenium dioxide and sodium selenite (64, 65). Our results suggest that one of the ways

that selenium may act to prevent prostate cancer is by inducing phase 2 enzyme activity.

para-Coumaric acid readily induced QR activity with little toxicity at high doses. Tomatoes possess relatively high levels of *para*-coumaric acid, and tomato consumption has been associated with a decreased risk of prostate cancer (66, 67). Previous work has ascribed this preventive effect to lycopene, the most potent quencher of singlet oxygen of all carotenoids but, in our hands, lacking QR inducing activity (68). Our findings raise the intriguing possibility that lycopene and *para*-coumaric acid in tomatoes may act in concert to protect against prostate cancer by quenching free radicals and inducing carcinogen defenses in prostate cells.

In summary, the human prostate cancer cell line LNCaP could serve as a model for future screens to identify phase 2 enzyme-inducing chemopreventive agents with activity in human prostate tissues. Although Michael acceptors appear most promising as prostate phase 2 enzyme-inducing agents, several other classes of compounds also show robust activation not easily predicted by their chemical class. Future efforts will focus on identifying additional phase 2 enzyme-inducing agents effective in the prostate *in vitro* and *in vivo* and in defining the factors responsible for the unique pattern of response to phase 2 enzyme-inducing agents in prostate cells.

References

- Greenlee, R. T., Hill-Harmon, M. B., Murray, T., and Thun, M. Cancer statistics, 2001. *CA Cancer J. Clin.*, 51: 15–36, 2001.
- Carter, B. S., Carter, H. B., and Isaacs, J. T. Epidemiologic evidence regarding predisposing factors to prostate cancer. *Prostate*, 16: 187–197, 1990.
- Yu, H., Harris, R. E., Gao, Y. T., Gao, R., and Wynder, E. L. Comparative epidemiology of cancers of the colon, rectum, prostate and breast in Shanghai, China versus the United States. *Int. J. Epidemiol.*, 20: 76–81, 1991.
- Shimizu, H., Ross, R. K., Bernstein, L., Yatani, R., Henderson, B. E., and Mack, T. M. Cancers of the prostate and breast among Japanese and white immigrants in Los Angeles County. *Br. J. Cancer*, 63: 963–966, 1991.
- Whittemore, A. S., Kolonel, L. N., Wu, A. H., John, E. M., Gallagher, R. P., Howe, G. R., Burch, J. D., Hankin, J., Dreon, D. M., West, D. W., *et al.* Prostate cancer in relation to diet, physical activity, and body size in blacks, whites, and Asians in the United States and Canada. *J. Natl. Cancer Inst.*, 87: 652–661, 1995.
- Haenzel, W., and Kurihara, M. Studies of Japanese migrants. I. Mortality from cancer and other diseases among Japanese men in the United States. *J. Natl. Cancer Inst.*, 40: 43–68, 1968.
- Danley, K. L., Richardson, J. L., Bernstein, L., Langholz, B., and Ross, R. K. Prostate cancer: trends in mortality and stage-specific incidence rates by racial/ethnic group in Los Angeles County, California (United States). *Cancer Causes Control*, 6: 492–498, 1995.
- Smith, J. R., Freije, D., Carpten, J. D., Gronberg, H., Xu, J., Isaacs, S. D., Brownstein, M. J., Bova, G. S., Guo, H., Bujnovszky, P., Nusskern, D. R., Damber, J. E., Bergh, A., Emanuelsson, M., Kallioniemi, O. P., Walker-Daniels, J., Bailey-Wilson, J. E., Beaty, T. H., Meyers, D. A., Walsh, P. C., Collins, F. S., Trent, J. M., and Isaacs, W. B. Major susceptibility locus for prostate cancer on chromosome 1 suggested by a genome-wide search [see comments]. *Science (Wash. DC)*, 274: 1371–1374, 1996.
- Xu, J., Meyers, D., Freije, D., Isaacs, S., Wiley, K., Nusskern, D., Ewing, C., Wilkens, E., Bujnovszky, P., Bova, G. S., Walsh, P., Isaacs, W., Schleutker, J., Matikainen, M., Tammela, T., Visakorpi, T., Kallioniemi, O. P., Berry, R., Schaid, D., French, A., McDonnell, S., Schroeder, J., Blute, M., Thibodeau, S., Trent, J., *et al.* Evidence for a prostate cancer susceptibility locus on the X chromosome. *Nat. Genet.*, 20: 175–179, 1998.
- Berry, R., Schroeder, J. J., French, A. J., McDonnell, S. K., Peterson, B. J., Cunningham, J. M., Thibodeau, S. N., and Schaid, D. J. Evidence for a prostate cancer-susceptibility locus on chromosome 20. *Am. J. Hum. Genet.*, 67: 82–91, 2000.
- Giovannucci, E., Stampfer, M. J., Krithivas, K., Brown, M., Brufsky, A., Talcott, J., Hennekens, C. H., and Kantoff, P. W. The CAG repeat within the androgen receptor gene and its relationship to prostate cancer. *Proc. Natl. Acad. Sci. USA*, 94: 3320–3323, 1997.
- Tavtigian, S. V., Simard, J., Teng, D. H., Abtin, V., Baumgard, M., Beck, A., Camp, N. J., Carillo, A. R., Chen, Y., Dayanar, P., Desrochers, M., Dumont, M., Farnham, J. M., Frank, D., Frye, C., Ghaffari, S., Gupte, J. S., Hu, R., Liev,

- D., Janecki, T., Kort, E. N., Laity, K. E., Leavitt, A., Leblanc, G., McArthur-Morrison, J., Pederson, A., Penn, B., Peterson, K. T., Reid, J. E., Richards, S., Schroeder, M., Smith, R., Snyder, S. C., Swedlund, B., Swensen, J., Thomas, A., Tranchant, M., Woodland, A. M., Labrie, F., Skolnick, M. H., Neuhausen, S., Rommens, J., and Cannon-Albright, L. A. A candidate prostate cancer susceptibility gene at chromosome 17p. *Nat. Genet.*, 27: 172–180, 2001.
13. Thompson, I. M., Coltman, C. A., Jr., and Crowley, J. Chemoprevention of prostate cancer: the Prostate Cancer Prevention Trial. *Prostate*, 33: 217–221, 1997.
14. Lee, W. H., Morton, R. A., Epstein, J. I., Brooks, J. D., Campbell, P. A., Bova, G. S., Hsieh, W. S., Isaacs, W. B., and Nelson, W. G. Cytidine methylation of regulatory sequences near the pi-class glutathione S-transferase gene accompanies human prostatic carcinogenesis. *Proc. Natl. Acad. Sci. USA*, 91: 11733–11737, 1994.
15. Lee, W.-H., Isaacs, W. B., Bova, G. S., and Nelson, W. G. CG island methylation changes near the *GSTP1* gene in prostatic carcinoma cells detected using the polymerase chain reaction: a new prostatic biomarker. *Cancer Epidemiol. Biomark. Prev.*, 6: 443–450, 1997.
16. Lin, X., Tascilar, M., Lee, W. H., Vles, W. J., Lee, B. H., Veeraswamy, R., Asgari, K., Freije, D., van Rees, B., Gage, W. R., Bova, G. S., Isaacs, W. B., Brooks, J. D., DeWeese, T. L., De Marzo, A. M., and Nelson, W. G. *GSTP1* CpG island hypermethylation is responsible for the absence of *GSTP1* expression in human prostate cancer cells. *Am. J. Pathol.*, 159: 1815–1826, 2001.
17. Brooks, J. D., Weinstein, M., Lin, X., Sun, Y., Pin, S. S., Bova, G. S., Epstein, J. I., Isaacs, W. B., and Nelson, W. G. CG island methylation changes near the *GSTP1* gene in prostatic intraepithelial neoplasia. *Cancer Epidemiol. Biomark. Prev.*, 7: 531–536, 1998.
18. Henderson, C. J., Smith, A. G., Ure, J., Brown, K., Bacon, E. J., and Wolf, C. R. Increased skin tumorigenesis in mice lacking pi class glutathione S-transferases. *Proc. Natl. Acad. Sci. USA*, 95: 5275–5280, 1998.
19. Wattenberg, L. W. Inhibition of carcinogenesis by minor dietary constituents. *Cancer Res.*, 52: 2085s–2091s, 1992.
20. Talalay, P., Fahey, J. W., Holtzclaw, W. D., Prestera, T., and Zhang, Y. Chemoprotection against cancer by phase 2 enzyme induction. *Toxicol. Lett.*, 82–83: 173–179, 1995.
21. Prochaska, H., Santamaria, A., and Talalay, P. Rapid detection of inducers of enzymes that protect against carcinogens. *Proc. Natl. Acad. Sci. USA*, 89: 2394–2398, 1992.
22. Zhang, Y., Talalay, P., Cho, C. G., and Posner, G. H. A major inducer of anticarcinogenic protective enzymes from broccoli: isolation and elucidation of structure. *Proc. Natl. Acad. Sci. USA*, 89: 2399–2403, 1992.
23. Prestera, T., Holtzclaw, W. D., Zhang, Y., and Talalay, P. Chemical and molecular regulation of enzymes that detoxify carcinogens. *Proc. Natl. Acad. Sci. USA*, 90: 2965–2969, 1993.
24. De Long, M. J., Prochaska, H. J., and Talalay, P. Induction of NAD(P)H:quinone reductase in murine hepatoma cells by phenolic antioxidants, azo dyes, and other chemoprotectors: a model system for the study of anticarcinogens. *Proc. Natl. Acad. Sci. USA*, 83: 787–791, 1986.
25. Fahey, J. W., Zhang, Y., and Talalay, P. Broccoli sprouts: an exceptionally rich source of inducers of enzymes that protect against chemical carcinogens. *Proc. Natl. Acad. Sci. USA*, 94: 10367–10372, 1997.
26. Chesis, P. L., Levin, D. E., Smith, M. T., Ernster, L., and Ames, B. N. Mutagenicity of quinones: pathways of metabolic activation and detoxification. *Proc. Natl. Acad. Sci. USA*, 81: 1696–1700, 1984.
27. Cavalieri, E. L., Devanesan, P., Bosland, M. C., Badawi, A. F., and Rogan, E. G. Catechol estrogen metabolites and conjugates in different regions of the prostate of Noble rats treated with 4-hydroxyestradiol: implications for estrogen-induced initiation of prostate cancer. *Carcinogenesis (Lond.)*, 23: 329–333, 2002.
28. Spencer, S. R., Wilczak, C. A., and Talalay, P. Induction of glutathione transferases and NAD(P)H:quinone reductase by fumaric acid derivatives in rodent cells and tissues. *Cancer Res.*, 50: 7871–7875, 1990.
29. Zhang, Y., Kensler, T. W., Cho, C. G., Posner, G. H., and Talalay, P. Anticarcinogenic activities of sulforaphane and structurally related synthetic norbornyl isothiocyanates. *Proc. Natl. Acad. Sci. USA*, 91: 3147–3150, 1994.
30. Prochaska, H. J., and Santamaria, A. B. Direct measurement of NAD(P)H:quinone reductase from cells cultured in microtiter wells: a screening assay for anticarcinogenic enzyme inducers. *Anal. Biochem.*, 169: 328–336, 1988.
31. Prochaska, H. J., and Fernandes, C. L. Elevation of serum phase II enzymes by anticarcinogenic enzyme inducers: markers for a chemoprotected state? *Carcinogenesis (Lond.)*, 14: 2441–2445, 1993.
32. De Long, M. J., Prochaska, H. J., and Talalay, P. Tissue-specific induction patterns of cancer-protective enzymes in mice by *tert*-butyl-4-hydroxyanisole and related substituted phenols. *Cancer Res.*, 45: 546–551, 1985.
33. van Lieshout, E. M., Peters, W. H., and Jansen, J. B. Effect of oltipraz, α -tocopherol, β -carotene and phenethylisothiocyanate on rat oesophageal, gastric, colonic and hepatic glutathione, glutathione S-transferase and peroxidase. *Carcinogenesis (Lond.)*, 17: 1439–1445, 1996.
34. Meyer, D. J., Harris, J. M., Gilmore, K. S., Coles, B., Kensler, T. W., and Ketterer, B. Quantitation of tissue- and sex-specific induction of rat GSH transferase subunits by dietary 1,2-dithiole-3-thiones. *Carcinogenesis (Lond.)*, 14: 567–572, 1993.
35. Hayes, J. D., and Pulford, D. J. The glutathione S-transferase supergene family: regulation of GST and the contribution of the isoenzymes to cancer chemoprotection and drug resistance. *Crit. Rev. Biochem. Mol. Biol.*, 30: 445–600, 1995.
36. Brooks, J. D., Paton, V. G., and Vidanes, G. Potent induction of phase 2 enzymes in human prostate cells by sulforaphane. *Cancer Epidemiol. Biomark. Prev.*, 10: 949–954, 2001.
37. Nelson, C. P., Kidd, L. C., Sauvageot, J., Isaacs, W. B., De Marzo, A. M., Groopman, J. D., Nelson, W. G., and Kensler, T. W. Protection against 2-hydroxyamino-1-methyl-6-phenylimidazo[4,5-*b*]pyridine cytotoxicity and DNA adduct formation in human prostate by glutathione S-transferase P1. *Cancer Res.*, 61: 103–109, 2001.
38. Wang, J. S., Shen, X., He, X., Zhu, Y. R., Zhang, B. C., Wang, J. B., Qian, G. S., Kuang, S. Y., Zarba, A., Egner, P. A., Jacobson, L. P., Munoz, A., Helzlsouer, K. J., Groopman, J. D., and Kensler, T. W. Protective alterations in phase I and 2 metabolism of aflatoxin B1 by oltipraz in residents of Qidong, People's Republic of China. *J. Natl. Cancer Inst.*, 91: 347–354, 1999.
39. Kensler, T. W., Curphey, T. J., Maxiutenko, Y., and Roebuck, B. D. Chemoprotection by organosulfur inducers of phase 2 enzymes: dithiolethiones and dithiols. *Drug Metab. Drug Interact.*, 17: 3–22, 2000.
40. Ramos-Gomez, M., Kwak, M. K., Dolan, P. M., Itoh, K., Yamamoto, M., Talalay, P., and Kensler, T. W. Sensitivity to carcinogenesis is increased and chemoprotective efficacy of enzyme inducers is lost in *nrf2* transcription factor-deficient mice. *Proc. Natl. Acad. Sci. USA*, 98: 3410–3415, 2001.
41. Hayes, J. D., and McMahon, M. Molecular basis for the contribution of the antioxidant responsive element to cancer chemoprotection. *Cancer Lett.*, 174: 103–113, 2001.
42. Spencer, S. R., Xue, L. A., Klenz, E. M., and Talalay, P. The potency of inducers of NAD(P)H:quinone-acceptor oxidoreductase parallels their efficiency as substrates for glutathione transferases. Structural and electronic correlations. *Biochem. J.*, 273: 711–717, 1991.
43. Talalay, P., De Long, M. J., and Prochaska, H. J. Identification of a common chemical signal regulating the induction of enzymes that protect against chemical carcinogenesis. *Proc. Natl. Acad. Sci. USA*, 85: 8261–8265, 1988.
44. Prestera, T., Zhang, Y., Spencer, S. R., Wilczak, C. A., and Talalay, P. The electrophile counterattack response: protection against neoplasia and toxicity. *Adv. Enzyme Regul.*, 33: 281–296, 1993.
45. Dinkova-Kostova, A. T., Massiah, M. A., Bozak, R. E., Hicks, R. J., and Talalay, P. Potency of Michael reaction acceptors as inducers of enzymes that protect against carcinogenesis depends on their reactivity with sulfhydryl groups. *Proc. Natl. Acad. Sci. USA*, 98: 3404–3409, 2001.
46. Zhang, Y. Role of glutathione in the accumulation of anticarcinogenic isothiocyanates and their glutathione conjugates by murine hepatoma cells. *Carcinogenesis (Lond.)*, 21: 1175–1182, 2000.
47. Zhang, Y. Molecular mechanism of rapid cellular accumulation of anticarcinogenic isothiocyanates. *Carcinogenesis (Lond.)*, 22: 425–431, 2001.
48. Ye, L., and Zhang, Y. Total intracellular accumulation levels of dietary isothiocyanates determine their activity in elevation of cellular glutathione and induction of phase 2 detoxification enzymes. *Carcinogenesis (Lond.)*, 22: 1987–1992, 2001.
49. Ross, D. T., Scherf, U., Eisen, M. B., Perou, C. M., Rees, C., Spellman, P., Iyer, V., Jeffrey, S. S., Van de Rijn, M., Waltham, M., Pergamenschikov, A., Lee, J. C., Lashkari, D., Shalon, D., Myers, T. G., Weinstein, J. N., Botstein, D., and Brown, P. O. Systematic variation in gene expression patterns in human cancer cell lines. *Nat. Genet.*, 24: 227–235, 2000.
50. Williams, E. D., and Brooks, J. D. New molecular approaches for identifying novel targets, mechanisms, and biomarkers for prostate cancer chemopreventive agents. *Urology*, 57: 100–102, 2001.
51. Dinkova-Kostova, A. T., and Talalay, P. Relation of structure of curcumin analogs to their potencies as inducers of phase 2 detoxification enzymes. *Carcinogenesis (Lond.)*, 20: 911–914, 1999.
52. Kuo, M. L., Huang, T. S., and Lin, J. K. Curcumin, an antioxidant and anti-tumor promoter, induces apoptosis in human leukemia cells. *Biochim. Biophys. Acta*, 1317: 95–100, 1996.
53. Sreejayan, and Rao, M. N. Curcuminoids as potent inhibitors of lipid peroxidation. *J. Pharm. Pharmacol.*, 46: 1013–1016, 1994.
54. De Marzo, A. M., Marchi, V. L., Epstein, J. I., and Nelson, W. G. Proliferative inflammatory atrophy of the prostate: implications for prostatic carcinogenesis. *Am. J. Pathol.*, 155: 1985–1992, 1999.

55. Surh, Y. J., Chun, K. S., Cha, H. H., Han, S. S., Keum, Y. S., Park, K. K., and Lee, S. S. Molecular mechanisms underlying chemopreventive activities of anti-inflammatory phytochemicals: down-regulation of COX-2 and iNOS through suppression of NF- κ B activation. *Mutat. Res.*, 480–481: 243–268, 2001.
56. Imaida, K., Tamano, S., Kato, K., Ikeda, Y., Asamoto, M., Takahashi, S., Nir, Z., Murakoshi, M., Nishino, H., and Shirai, T. Lack of chemopreventive effects of lycopene and curcumin on experimental rat prostate carcinogenesis. *Carcinogenesis (Lond.)*, 22: 467–472, 2001.
57. Dorai, T., Cao, Y. C., Dorai, B., Buttyan, R., and Katz, A. E. Therapeutic potential of curcumin in human prostate cancer. III. Curcumin inhibits proliferation, induces apoptosis, and inhibits angiogenesis of LNCaP prostate cancer cells *in vivo*. *Prostate*, 47: 293–303, 2001.
58. Dinkova-Kostova, A. T., Abeygunawardana, C., and Talalay, P. Chemoprotective properties of phenylpropenoids, bis(benzylidene)cycloalkanones, and related Michael reaction acceptors: correlation of potencies as phase 2 enzyme inducers and radical scavengers. *J. Med. Chem.*, 41: 5287–5296, 1998.
59. Miranda, C. L., Aponso, G. L., Stevens, J. F., Deinzer, M. L., and Buhler, D. R. Prenylated chalcones and flavanones as inducers of quinone reductase in mouse Hepa1c1c7 cells. *Cancer Lett.*, 149: 21–29, 2000.
60. Cohen, J. H., Kristal, A. R., and Stanford, J. L. Fruit and vegetable intakes and prostate cancer risk. *J. Natl. Cancer Inst.*, 92: 61–68, 2000.
61. Kolonel, L. N., Hankin, J. H., Whittemore, A. S., Wu, A. H., Gallagher, R. P., Wilkens, L. R., John, E. M., Howe, G. R., Dreon, D. M., West, D. W., and Paffenbarger, R. S., Jr. Vegetables, fruits, legumes and prostate cancer: a multiethnic case-control study. *Cancer Epidemiol. Biomark. Prev.*, 9: 795–804, 2000.
62. Le Marchand, L., Hankin, J. H., Kolonel, L. N., and Wilkens, L. R. Vegetable and fruit consumption in relation to prostate cancer risk in Hawaii: a reevaluation of the effect of dietary β -carotene. *Am. J. Epidemiol.*, 133: 215–219, 1991.
63. Spitz, M. R., Duphorne, C. M., Detry, M. A., Pillow, P. C., Amos, C. I., Lei, L., de Andrade, M., Gu, X., Hong, W. K., and Wu, X. Dietary intake of isothiocyanates: evidence of a joint effect with glutathione S-transferase polymorphisms in lung cancer risk. *Cancer Epidemiol. Biomark. Prev.*, 9: 1017–1020, 2000.
64. Nomura, A. M., Lee, J., Stemmermann, G. N., and Combs, G. F., Jr. Serum selenium and subsequent risk of prostate cancer. *Cancer Epidemiol. Biomark. Prev.*, 9: 883–887, 2000.
65. Brooks, J. D., Metter, E. J., Chan, D. W., Sokoll, L. J., Landis, P., Nelson, W. G., Muller, D., Andres, R., and Carter, H. B. Plasma selenium level before diagnosis and the risk of prostate cancer development. *J. Urol.*, 166: 2034–2038, 2001.
66. Giovannucci, E., Ascherio, A., Rimm, E. B., Stampfer, M. J., Colditz, G. A., and Willett, W. C. Intake of carotenoids and retinol in relation to risk of prostate cancer. *J. Natl. Cancer Inst.*, 87: 1767–1776, 1995.
67. Giovannucci, E. Tomatoes, tomato-based products, lycopene, and cancer: review of the epidemiologic literature. *J. Natl. Cancer Inst.*, 91: 317–331, 1999.
68. Di Mascio, P., Kaiser, S., and Sies, H. Lycopene as the most efficient biological carotenoid singlet oxygen quencher. *Arch. Biochem. Biophys.*, 274: 532–538, 1989.