

Identification of Prolactin-Sensitive GABA and Kisspeptin Neurons in Regions of the Rat Hypothalamus Involved in the Control of Fertility

Ilona C. Kokay, Sandra L. Petersen, and David R. Grattan

Centre for Neuroendocrinology and Department of Anatomy and Structural Biology (I.C.K., D.R.G.), University of Otago, Dunedin 9016, New Zealand; and Neuroscience and Behavior Program and Centre for Neuroendocrine Studies (S.L.P.), University of Massachusetts, Amherst, Massachusetts 01003

High levels of circulating prolactin are known to cause infertility, but the precise mechanisms by which prolactin influences the neuroendocrine axis are yet to be determined. We used dual-label *in situ* hybridization to investigate whether prolactin-receptor (PRLR) mRNA is expressed in GnRH neurons. In addition, because γ -aminobutyric acidergic and kisspeptin neurons in the rostral hypothalamus are known to regulate GnRH neurons and, hence, might mediate the actions of prolactin, we investigated whether these neurons coexpress PRLR mRNA. ³⁵S-labeled RNA probes to detect PRLR mRNA were hybridized together with digoxigenin-labeled probes to detect either GnRH, *Gad1/Gad2*, or *Kiss1* mRNA in the rostral hypothalamus of ovariectomized (OVX), estradiol-treated rats. Additional sets of serial sections were cut through the arcuate nucleus of OVX rats, without estradiol replacement, to examine coexpression of PRLR mRNA in the arcuate population of kisspeptin neurons. PRLR mRNA was highly expressed throughout the rostral preoptic area, particularly in periventricular regions surrounding the third ventricle, and there was a high degree of colocalization of PRLR mRNA in both *Gad1/Gad2* and *Kiss1* mRNA-containing cells (86 and 85.5%, respectively). In contrast, only a small number of GnRH neurons (<5%) was found to coexpress PRLR mRNA. In the arcuate nucleus of OVX rats, the majority of *Kiss1* mRNA-containing cells also coexpressed PRLR mRNA. These data are consistent with the hypothesis that, in addition to a direct action on a small subpopulation of GnRH neurons, prolactin actions on GnRH neurons are predominantly mediated indirectly, through known afferent pathways. (*Endocrinology* 152: 526–535, 2011)

Hyperprolactinaemia is the most common disorder of the hypothalamic-pituitary axis (1) and is well recognized as a major cause of reproductive dysfunction in both males and females. Approximately 40% of women presenting with secondary amenorrhoea have increased serum prolactin levels (2, 3). Men may similarly present with infertility and decreased libido, with hyperprolactinemia present in approximately 16% of patients with erectile dysfunction and 11% of men with oligospermia (4). Infertility and loss of libido are serious side effects caused by the hyperprolactinemia induced by neuroleptic drugs, affecting up to 70% of patients and representing

one of the major causes of noncompliance with antipsychotic medication (3). Further understanding of the mechanism of prolactin in suppressing reproduction may allow better therapy or drug design.

Precisely how hyperprolactinaemia disrupts the reproductive axis is unclear. In humans, elevated circulating levels of prolactin are associated with a marked reduction in both the frequency and amplitude of LH pulses (5), a direct result of GnRH pulses, and the suppression of LH pulsatility can be reversed by reducing serum prolactin concentrations to normal (6). Similarly, prolactin suppresses both the frequency and the amplitude of LH pulses

ISSN Print 0013-7227 ISSN Online 1945-7170

Printed in U.S.A.

Copyright © 2011 by The Endocrine Society

doi: 10.1210/en.2010-0668 Received June 14, 2010. Accepted October 29, 2010.

First Published Online December 22, 2010

Abbreviations: E, Estradiol; GABA, γ -aminobutyric acid; GAD, glutamic acid decarboxylase; OVX, ovariectomized; PRLR, prolactin receptor; RP3V, rostral periventricular area of the third ventricle; TIDA, tuberoinfundibular dopamine.

in rats (7–10), and direct measurements of GnRH secretion into the portal blood have revealed a prolactin-induced suppression of GnRH release (11, 12). Furthermore, during lactation, a physiological state characterized by high levels of circulating prolactin, pulsatile LH release is suppressed (13, 14), and the activity of GnRH neurons is reduced (15). Although there is evidence that prolactin can also act in the pituitary gland to suppress LH and FSH secretion (16–18), pulsatile GnRH replacement can reverse the infertility induced by hyperprolactinaemia (19, 20), suggesting that prolactin-induced suppression of GnRH release is the principal cause of infertility.

We recently reported that a small proportion of GnRH neurons in mice express the prolactin receptor (PRLR) and show prolactin-induced changes in cAMP response element-binding protein phosphorylation (21). These data suggested that prolactin might act directly on some GnRH neurons to regulate fertility. The infrequent expression of PRLRs in GnRH neurons, however, contrasts markedly with the relatively high levels of PRLR expression in regions of the rostral hypothalamus known to be involved in the control of GnRH secretion (22, 23). The majority of GnRH neuronal cell bodies reside in the preoptic area/anterior hypothalamus and the hypothalamic region extending from the anteroventral periventricular nucleus to the periventricular preoptic region [collectively designated the rostral periventricular area of the third ventricle (RP3V) (24)] is well established to be involved in the regulation of GnRH neurons (24–27). The RP3V also contains abundant PRLR mRNA (22, 23), suggesting that prolactin actions on GnRH neurons could be mediated indirectly through actions of RP3V neuronal populations. The neurochemical phenotype of the PRLR-expressing neurons in the RP3V, however, is not known.

Based on the current literature, the two most likely populations in the RP3V that could be involved in regulating GnRH neuronal activity are γ -aminobutyric acid (GABA)ergic and kisspeptin neurons. There is convincing evidence that the inhibitory neurotransmitter GABA is an important player in the regulation of GnRH secretion (28; also see reviews in Refs. 29, 30). GABAergic neurons form synapses onto GnRH neurons (31), and studies using *in situ* hybridization (32, 33) or single cell RT-PCR methodology (34) have shown that GnRH neurons express mRNA for GABA receptor subunits. In addition, a wealth of electrophysiological data has shown that GABA affects the activity of GnRH neurons (35–38). Neurons expressing the neuropeptide kisspeptin are also found prominently in the RP3V (39). GnRH neurons express G protein-coupled receptor 54, the kisspeptin receptor (40–42), and kisspeptin potently activates GnRH activity (40, 42, 43). In female rodents, kisspeptin is one of the essential

components mediating the onset of GnRH activity at puberty [reviewed by Clarkson *et al.* (44)] and plays a role in inducing the preovulatory GnRH surge (43, 45, 46).

We hypothesize that prolactin may influence GnRH neurons indirectly, through actions on afferent GABA and kisspeptin neuronal populations located in the RP3V. The aim of this study was to use dual-label *in situ* hybridization to determine whether these neuronal populations express PRLR mRNA. In addition, because there is a second population of kisspeptin neurons present in the arcuate nucleus of the rodent hypothalamus (43, 47–49), and this nucleus contains many PRLR-expressing neurons (22, 23, 50–52), we also investigated PRLR expression in the arcuate kisspeptin neurons. Finally, to extend data from mice showing expression of PRLRs in a subpopulation of GnRH neurons, we examined whether GnRH neurons in the rat expressed PRLR (long form) mRNA.

Materials and Methods

Animals and tissue preparation

All animal manipulations were approved by the University of Otago Animal Ethics committee. Virgin, adult female Sprague Dawley rats (10–12 wk old) were group housed under standard, controlled laboratory conditions (14-h light, 10-h dark cycle; temperature, 21 ± 1 C). Rats were bilaterally ovariectomized (OVX) under halothane anesthesia using sterile surgical procedures. Seven days later, silastic implants (length, 30 mm; inside diameter, 1.57 mm; outside diameter, 3.18 mm) containing either 150 μ g/ml of 17 β -estradiol (E) to restore low physiological levels (equivalent to diestrus) (OVX+E) or vehicle (sesame oil) were implanted sc. This level of E is not sufficient to produce a daily LH or prolactin surge. One week later, between 0930 and 1100 h, all rats were anesthetized with an overdose of sodium pentobarbitone, then transcardially perfused with buffered 2% paraformaldehyde. Brains were removed, postfixed overnight, then cryoprotected in 30% sucrose in phosphate buffer for 72 h before being frozen on powdered dry ice and stored at -80 C. Sets of serial, coronal sections of 16- μ m thickness were cut through the rostral hypothalamus from the preoptic region to the periventricular area (bregma 0.00 to -0.84 mm) and through the arcuate nucleus (bregma -2.10 to -3.30 mm) (53) and mounted onto aminopropyltriethoxysilane-coated slides. Sections were stored at -80 C until used for *in situ* hybridization.

Preparation of probes for *in situ* hybridization

To prepare probes to detect *Kiss1* and *Gad1/Gad2*, genes encoding kisspeptin-54 and glutamic acid decarboxylase (GAD)1 and GAD2, respectively, and PRLR mRNA, cDNA templates were first generated by RT-PCR. Primer pairs were designed using sequences from GenBank for the long form of the PRLR (accession no. M57668), for *Kiss1* (accession no. NM181692), and for *Gad2* and *Gad1* (accession nos. NM012563 and NM1077007, respectively), and these primer pairs used to generate cDNA templates (290–313 bp) that incorporated T7 and SP6 RNA polymerase sequence sites. The specificity of all the cDNA templates was confirmed by sequencing. ³⁵S-labeled

cRNA probes targeted to the PRLR were generated using T7 polymerase (antisense probe) or SP6 polymerase (sense probe) in accordance with the Riboprobe In Vitro kit (Promega, Madison, WI) protocol as described previously (51). Antisense and sense cRNA probes for *Kiss1*, *Gad1*, and *Gad2*, labeled with digoxigenin, were transcribed from the cDNA templates using a digoxigenin RNA-Labeling kit (Roche Diagnostics GmbH, Mannheim, Germany) and the appropriate polymerases. A cRNA probe targeted to GnRH mRNA was transcribed from a 330 bp *BamH1-Hind111* cDNA construct subcloned into a p65T7 vector. The cDNA was linearized with *BamH1*, then transcribed with T7 polymerase in the presence of digoxigenin uridine triphosphate, using a commercial digoxigenin RNA-Labeling kit (Roche). Unincorporated nucleotides were removed from all labeled probes with mini Quick Spin RNA purification columns (Roche).

Dual-label *in situ* hybridization histochemistry

Sections were pretreated with a 5 min, 2% paraformaldehyde fixation step, permeabilized with proteinase K, then acetylated and prehybridized in the hybridization solution [100 mM dithiothreitol, 0.3 M NaCl, 20 mM Tris (pH 8), 5 mM EDTA (pH 8), 1× Denhardt's solution, 10% dextran sulfate, and 50% formamide] without the labeled cRNA probes for 1–3 h. The probes were denatured at 95 C for 3 min, then diluted into hybridization solution, pipetted onto the sections, and the slides were coverslipped (Hybri-slips; Sigma-Aldrich, St. Louis, MO). Sections were hybridized for 16 h at 55 C with an ³⁵S-labeled RNA probe (1.2 × 10⁶ cpm per 120 μl of hybridization solution) complementary to the long form of the PRLR together with digoxigenin-labeled probes (0.5 ng/μl·Kb per section) specific for either GnRH, *Gad1* and *Gad2*, or *Kiss1* mRNA. After hybridization, sections were washed in sodium saline citrate solution (150 mM NaCl and 15 mM sodium citrate) containing β-mercaptoethanol, subjected to ribonuclease treatment, followed by a series of washes (most stringent, 0.1 M sodium saline citrate at 55 C with shaking for 2 h). To visualize the digoxigenin-labeled mRNA hybrids, sections were incubated in sheep antidigoxigenin antibody conjugated to alkaline phosphatase (1:2000 dilution) for 24 h at 4 C, then detected using nitroblue tetrazolium/5-bromo-4-chloro-3-indolyl phosphate as a substrate. After air-drying, slides were dipped in undiluted LM-1 Hypercoat emulsion (Amersham Pharmacia Biotech, Piscataway, NJ), sealed in lightproof boxes with desiccant and stored at 4 C for 4–5 wk. Slides were developed with D-19 developer (Kodak, New York, NY), fixed with Ilford Hypan fixer, dehydrated through alcohols, cleared in xylene, then mounted with VectaMount mounting medium, and coverslipped. To generate autoradiograms, the majority of slides were also apposed to film (Kodak BioMax MR) for 3–5 d before emulsion coating.

Data analysis

Sections were photographed under brightfield illumination with ×40 objective, using a SPOT RT digital camera and associated SPOT software (Diagnostic Instruments, Inc., Sterling Heights, MI), attached to an Olympus BX51 microscope. Digoxigenin-labeled cells were identified by the presence of a distinct blue/purple cytoplasmic precipitate. For GnRH neurons, all digoxigenin-labeled GnRH mRNA-positive cells were counted in five to seven sections throughout the preoptic area and rostral hypothalamus of OVX, E-treated rats. Data are presented as the total number counted, per brain (n = 4, mean ±

SEM). For *Kiss1* and *Gad1/Gad2* digoxigenin-labeled mRNA-positive cells, a single field of view was photographed at ×400 magnification on each side of the brain, using the third ventricle to define one edge of the field. All labeled cells within these fields were counted in each of four sections through the RP3V region of the hypothalamus of OVX+E rats (n = 4). In the arcuate nucleus, all *Kiss1* mRNA-positive cells were counted in four to six slides from both OVX and OVX+E rats (n = 4 in each group). For *Kiss1* and *Gad1/Gad2* mRNA, data are expressed as the mean number of identified cells per section in each brain (mean ± SEM).

To assess the number of PRLR-expressing cells, the number of silver grains over each cell (proportional to the number of PRLR mRNA hybrids) in these same sections was quantified using ImageJ software (National Institutes of Health). Silver grains were identified and analyzed under brightfield illumination and confirmed by darkfield inspection of the sections. Mean background levels of silver grains on each slide were sampled from five random areas, which were approximately equivalent in size to that of the silver grain clusters present over neuronal cell bodies. PRLR-positive cells were defined as those in which the density of clusters of silver grains overlying neuronal cell bodies was greater than three times background silver grain levels. Cells were counted as double labeled if a digoxigenin-labeled cell was also positive for PRLR mRNA. Differences between groups were calculated by ANOVA followed by Newman-Keuls *post hoc* tests using GraphPad Prism software for Macintosh (GraphPad, San Diego, CA). The level of significance was set at *P* < 0.05.

Results

Double-label *in situ* hybridization for PRLR mRNA and GnRH mRNA

As shown in the autoradiograph and associated dark-field image in Fig. 1, A and B, high levels of PRLR-expressing cells were found throughout the RP3V region, particularly in the area close to the third ventricle. A representative image of cells subjected to double-label *in situ* hybridization experiments to detect colocalization of PRLR and GnRH mRNA expression is shown in Fig. 1C. Neuronal cell bodies expressing GnRH mRNA hybrids, visible as a colored precipitate in the cytoplasm of the cell, were detected in a scattered pattern throughout the rostral preoptic area. This pattern of expression closely matches the well-established distribution of GnRH neurons (54), thus confirming the specificity of the digoxigenin-labeled probe. Although PRLR mRNA transcripts were abundant in the RP3V region of the hypothalamus and in the adjacent medial preoptic area (particularly in the ventromedial preoptic nucleus, which is sparsely populated with GnRH neurons), only a few GnRH neurons (≈5%) were found to coexpress PRLR mRNA, evidenced by the lack of visible silver grain clusters over GnRH neuronal cell bodies (Fig. 1, C and D).




FIG. 1. Dual labeling of GnRH and PRLR mRNA by *in situ* hybridization in the rostral preoptic area. A, Representative autoradiogram showing PRLR mRNA (black areas) in the rostral preoptic area (note also strong labeling in the choroid plexus). rPOA, Rostral preoptic area; CP, choroid plexus. B, Darkfield image of the box shown in A with PRLR mRNA revealed as white clusters of silver grains. C, Micrograph showing localization of GnRH neurons (blue/purple cytoplasmic precipitate) and PRLR mRNA (clusters of black grains). Section has been counterstained with Gills hematoxylin to identify cell nuclei. Insets, Brightfield (upper) and darkfield (lower); high-power views of the area in the box of C. White arrows indicate PRLR mRNA-positive cells, visualized as clusters of silver grains. Yellow arrow indicates a GnRH neuron. Note absence of silver grains over this cell, illustrating lack of colocalization of GnRH and PRLR mRNA. D, Histogram comparing the mean number of GnRH neurons counted per rat with the mean number of GnRH neurons expressing PRLR mRNA. Scale bar, 100 μ m (B and C) and 10 μ m (insets).

Colocalization of PRLR mRNA and *Gad1/Gad2* mRNA

GABAergic neurons were detected using a mixture of two probes that were targeted to the two genes (*Gad1* and *Gad2*), which encode the GABAergic synthesizing enzyme, glutamic acid decarboxylase. In the rat brain, almost all GABAergic neurons express both *Gad1* and *Gad2* mRNA (55, 56.) Results from dual-label *in situ* hybrid-

ization experiments for PRLR mRNA labeled with 35 S together with GABAergic neurons visualized using a mixture of the two digoxigenin-labeled cRNA probes are illustrated in Fig. 2. As has been characterized previously (57, 58), there was a large number of GABAergic neurons present throughout the RP3V. A high degree of colocalization of PRLR mRNA in *Gad1/Gad2* mRNA-containing cells was observed as shown by the presence of dense clusters of silver grains over individual *Gad1/Gad2*-digoxigenin-labeled cell soma (Fig. 2C). Over 85% of PRLR mRNA-positive cells expressed *Gad1/Gad2* mRNA, suggesting that the vast majority of these cells are GABAergic (Fig. 2D). This high level of coexpression was observed throughout the RP3V region as well as in the




FIG. 2. Coexpression of *Gad1/Gad2* (GAD) and PRLR mRNA in the RP3V region of the hypothalamus. A, Representative autoradiogram showing localization of PRLR mRNA (black areas) in the periventricular region. B, Most GAD-positive cells outside the RP3V do not colocalize with PRLR mRNA, illustrated in this example from the horizontal limb of the diagonal band of Broca. C, Low-power view showing localization of GAD mRNA and PRLR mRNA in the RP3V region. Insets c¹ and c² are high-power photomicrographs of boxes in C, showing examples of GAD mRNA-containing cells coexpressing PRLR mRNA in the RP3V region. Black arrows indicate double-labeled cells, whereas the white arrow indicates a non-GABAergic cell single labeled for PRLR mRNA. D, Histogram comparing the mean number of GAD neurons counted (average per section) with the mean number that coexpressed prolactin mRNA. Note that relatively few neurons that expressed PRLR were not also GAD positive. Scale bar, 100 μ m (C) and 10 μ m (B, c¹, and c²).

medial preoptic area, but was specific to these periventricular regions. In other brain regions present in the same sections, where GABA cells were also numerous, for example, the horizontal limb of the diagonal band of Broca, *Gad1/Gad2* mRNA-containing cells did not coexpress PRLR mRNA (Fig. 2B).

Colocalization of PRLR mRNA and *Kiss1* mRNA

1) RP3V *Kiss1* population

In the OVX+E rats, the pattern of *Kiss1* mRNA-expressing neurons in the RP3V was consistent with the expression patterns in rat brain reported elsewhere (59), with the highest number of *Kiss1*-expressing cells present more caudally in the periventricular area of the RP3V region (Fig. 3A). Throughout the RP3V, a high degree of colocalization of PRLR mRNA with *Kiss1* mRNA was detected with PRLR mRNA coexpressed by 86% of kisspeptin neurons (Fig. 3B). Consistent with the pattern of PRLR mRNA seen in the dual-label *Gad1/Gad2* *in situ* hybridization experiments, PRLR mRNA was not restricted to the kisspeptin cells but was abundantly ex-

pressed on cells throughout the RP3V. Levels of *Kiss1* mRNA in the group that were OVX but not given E implants were virtually undetectable.

2) Arcuate nucleus *Kiss1* population

Because *Kiss1* mRNA expression in the arcuate nucleus is markedly increased when levels of circulating E are low, PRLR expression in identified kisspeptin neurons was evaluated in sections from both the OVX rat group that received oil implants as well as those from OVX+E rats. Figure 4, A and B, compares the relative distribution of *Kiss1* mRNA-positive cells (brightfield) with PRLR mRNA (darkfield) in the same section. Note that there was extensive expression of PRLR in the arcuate nucleus in the dorsomedial region, distinct from the distribution of kisspeptin cells. Analysis of the dual-label experiment shows that, although significantly more *Kiss1*-containing cells were detected in the OVX group, numerous *Kiss1* cells were also clearly detectable in the steroid-replaced group. In the former group, the majority (79%) of kisspeptin neurons was found to coexpress prolactin mRNA (Fig. 4, D and E). In OVX rats given E replacement, many fewer kisspeptin cells were detected, and a smaller proportion of those cells was double labeled (45%) (Fig. 4, C and E).




FIG. 3. Coexpression of *Kiss1* and PRLR mRNA in the RP3V region of the hypothalamus. A, Low-power view showing distribution of *Kiss1* mRNA and PRLR mRNA in the periventricular region. a¹ and a², High-power photomicrographs of boxes in A showing examples of *Kiss1* mRNA-containing cells coexpressing PRLR mRNA. Black arrows indicate double-labeled cells, whereas white arrows indicate a cell single labeled for PRLR mRNA. B, Histogram comparing the mean number of *Kiss1*-positive neurons counted (average per section) with the mean number that coexpressed prolactin mRNA. In this region, there was also a large number of PRLR-containing cells that did not express *Kiss1* mRNA. Scale bar, 100 μ m (A) and 10 μ m (a¹ and a²).

Discussion

In the present study, we have documented high levels of PRLR expression in regions of the hypothalamus known to be involved in the regulation of GnRH neurons. Consistent with our previous data in mice (21), we found that a relatively small proportion of GnRH neurons express the long form of the PRLR, but that levels of PRLR expression were high in regions of the RP3V surrounding GnRH neurons. Within the RP3V, almost all GABAergic neurons coexpressed PRLR mRNA. Similarly, many kisspeptin neurons in both the RP3V and arcuate nucleus were found to contain PRLR mRNA. These data provide new insights into potential mechanisms by which prolactin might influence GnRH neurons, suggesting an indirect pathway mediated by both stimulatory and inhibitory afferent neurons.

Our data documenting PRLR mRNA expression in the rostral hypothalamus show a relatively restricted distribution of the receptor in the RP3V region of the rostral forebrain. These data are consistent with a range of previous *in situ* hybridization studies in both rats (22, 60) and mice (23). Our autoradiographs were exposed for a relatively short period; and hence, we did not see expression in regions such as the bed nucleus of the stria terminalis and medial septum, in which PRLR mRNA could be de-




FIG. 4. Coexpression of *Kiss1* and PRLR mRNA in the arcuate nucleus of the hypothalamus. **A**, A coronal section at the level of the arcuate nucleus photographed under both brightfield and (**B**) darkfield illumination. The brightfield micrograph shows the distribution of *Kiss1* mRNA detected using a digoxigenin-labeled probe (blue/purple cytoplasmic precipitate) and PRLR mRNA (clusters of black grains). The darkfield view illustrates the distribution of PRLR mRNA as clusters of black grains overlying cells. Note that much of the PRLR mRNA in the arcuate nucleus is not associated with kisspeptin neurons. **C**, Low-power micrograph showing distribution of *Kiss1* mRNA and PRLR mRNA in an OVX rat given E replacement (OVX+E). **c**¹ and **c**², High-power photomicrographs of boxes in **C** showing examples of *Kiss1* mRNA-containing cells coexpressing PRLR mRNA. *Black arrow* indicates a double-labeled cell, whereas *white arrows* indicate cells single labeled for PRLR mRNA. **D**, Low-power micrograph showing distribution of *Kiss1* mRNA and PRLR mRNA in an OVX rat. **d**¹, High-power photomicrograph of box in **D** showing examples of *Kiss1* mRNA-containing cells coexpressing PRLR mRNA. The *black arrows* indicate double-labeled cells, whereas *white arrows* indicate cells single labeled for PRLR mRNA. **E**, The mean number of *Kiss1*-positive neurons that coexpressed prolactin mRNA in OVX and OVX+E-replaced rats. *, $P < 0.05$. Scale bar, 100 μm (A–D) and 10 μm (**c**¹, **c**², and **d**¹).

tected in mice (23). This observation suggests that levels of PRLR mRNA are higher in the periventricular areas than in these other regions and that nuclei in these regions are highly prolactin sensitive.

By far the majority of neurons in the prolactin-sensitive regions of the RP3V area is GABAergic and/or glutamatergic (61, 62). There is a large body of evidence suggesting that GABA regulates GnRH neuronal activity. GABA neurons synapse directly on GnRH neurons (31), and GnRH neurons express mRNA for several subunits of the GABA_A (32) and GABA_B receptor (63). Because over 85% of the GABAergic neurons in this region express PRLR mRNA, it seems highly likely that at least some of these prolactin-responsive neurons will be directly innervating GnRH neurons. Pharmacological evidence consistently demonstrates an inhibitory influence over GnRH mRNA and GnRH and LH secretion (64, 65). Although there is some controversy (66, 67), electrophysiological evidence suggests that GABAergic neurons maintain a tonic inhibitory regulation of excitability of adult GnRH neurons (36, 65). GABA also alters intracellular calcium specifically in GnRH neurons (68, 69). Hyperprolactinaemia has been reported to increase the activity of hypothalamic GABAergic neurons (70–73). We have also described increased

activity of GABA neurons during lactation (74), when prolactin levels are elevated, and GABA levels in the cerebrospinal fluid are increased during lactation (75). These observations, together with the present data showing specific expression of PRLRs in GABAergic neurons in the RP3V, support the hypothesis that GABAergic neurons may mediate the suppressive effects of prolactin on GnRH neurons.

In recent years, there has been an explosion of interest in the role of kisspeptin in regulating GnRH neurons. Consistent with earlier reports (49, 59, 76), we identified two populations of kisspeptin neurons in the rodent that are reciprocally regulated by E: one population that is present in the RP3V and a second population in the arcuate nucleus. In OVX rats, kisspeptin mRNA was prominent in the arcuate nucleus, whereas there was a marked decrease in kisspeptin mRNA levels detected after E treatment. In contrast, in the RP3V, kisspeptin mRNA was undetectable in OVX rats but, after E treatment, was readily detectable. In both populations of kisspeptin neurons, a high proportion of neurons expressed PRLR mRNA, suggesting that they may be regulated by prolactin.

Kisspeptin neurons in the RP3V region project directly to the cell bodies of GnRH neurons (76) are activated by

elevated E and are essential for the generation of the pre-ovulatory GnRH surge (43, 45, 46, 77). Thus, the presence of PRLR mRNA in 85% of these cells suggests that prolactin-responsive neurons will directly innervate GnRH neurons. These data are consistent with the hypothesis that PRLR-mediated signaling might play a role in decreasing kisspeptin levels, contributing to infertility. Interestingly, in the RP3V, there were very few PRLR-positive cells that were not also labeled with *Gad1/Gad2* mRNA, suggesting that the vast majority of these cells is GABAergic. The colocalization of some of these cells with kisspeptin, therefore, raises the interesting possibility that at least some RP3V kisspeptin neurons might coexpress GAD. Further work will be required to confirm this.

Although evidence linking RP3V kisspeptin neurons to GnRH secretion is strong, the role of the arcuate kisspeptin neurons is less clear. It has been suggested that this population of kisspeptin cells may be involved in the negative feedback regulation of GnRH secretion (43, 49), although the anatomical evidence linking these cells to GnRH neurons in rodents is still a little thin. In the monkey, kisspeptin fibers have been identified in the median eminence, closely associated with the GnRH axons and axon terminals (78), and there are some data in rodents suggesting that kisspeptin-expressing neuronal fibers, identified by coexpression of prodynorphin and proneurokinin B, terminate near GnRH fibers in the median eminence (79). It has been proposed that the arcuate population of kisspeptin neurons stimulate tonic pulsatile secretion of GnRH and are a potential site of E negative-feedback suppression of GnRH secretion (80, 81). Prolactin actions on either population of kisspeptin neurons, therefore, might influence fertility by indirectly regulating GnRH neurons. During lactation, when prolactin levels are markedly elevated, levels of *Kiss1* mRNA in the arcuate nucleus are markedly suppressed (82, 83). This is particularly interesting, because E levels are extremely low at this time, which would normally favor elevated kisspeptin expression.

It is possible that the arcuate kisspeptin neurons play other modulatory roles, rather than the control of reproduction. There is now good evidence that kisspeptin cells in the arcuate nucleus coexpress both dynorphin and neurokinin B (80, 84), and earlier studies have shown that dynorphin cells project to the tuberoinfundibular dopamine (TIDA) neurons within the arcuate nucleus (85). Changes in TIDA neuronal activity are the most important factors maintaining homeostasis of prolactin levels in the blood. There is good evidence that endogenous opioid peptides modulate TIDA neuronal activity (86–88), particularly through both κ and μ receptors (87–89), and dynorphin may play a critical role here. There is also recent

evidence that kisspeptin might suppress TIDA neuronal activity (90). Hence, it is possible that prolactin-mediated signaling in the arcuate population of kisspeptin neurons might be involved in regulation of its own secretion.

The present study has shown that in hypothalamic nuclei involved in the control of fertility, both GABA and kisspeptin neurons express PRLR mRNA. In contrast, very few GnRH neurons express PRLR mRNA. These data are consistent with the hypothesis that prolactin actions on GnRH neurons are predominantly mediated indirectly through these known afferent pathways. Although hyperprolactinemia-induced infertility can be thought of as a pathological condition, arising, for example, from a pituitary adenoma, the mechanism for prolactin to suppress GnRH neurons has likely evolved as an adaptive response to pregnancy and lactation (91). Hyperprolactinemia during these physiological states is likely to contribute to the suppression of GnRH neurons that occurs in these states (13–15). This lactational infertility plays a critical role in birth spacing, ensuring optimal survival of mammalian young by minimizing maternal investment in a subsequent pregnancy (92).

Acknowledgments

Address all correspondence and requests for reprints to: Prof. Dave Grattan, Department of Anatomy and Structural Biology, University of Otago, P.O. Box 913, Dunedin 9054, New Zealand. E-mail: dave.grattan@otago.ac.nz.

This work was supported by the Health Research Council of New Zealand and by a Foundation for Research, Science, and Technology fellowship (I.C.K.).

Disclosure Summary: The authors have nothing to disclose.

References

1. Mah PM, Webster J 2002 Hyperprolactinemia: etiology, diagnosis, and management. *Semin Reprod Med* 20:365–374
2. Evans WS, Cronin MJ, Thorner MO 1982 Hypogonadism in hyperprolactinemia: proposed mechanisms. In: Ganong WF, Martini L, ed. *Frontiers in neuroendocrinology*. Vol 7, New York: Raven Press; 77–122
3. Meaney AM, O'Keane V 2002 Prolactin and schizophrenia: clinical consequences of hyperprolactinaemia. *Life Sci* 71:979–992
4. De Rosa M, Zarrilli S, Di Sarno A, Milano N, Gaccione M, Boggia B, Lombardi G, Colao A 2003 Hyperprolactinemia in men: clinical and biochemical features and response to treatment. *Endocrine* 20: 75–82
5. Matsuzaki T, Azuma K, Irahara M, Yasui T, Aono T 1994 Mechanism of anovulation in hyperprolactinemic amenorrhea determined by pulsatile gonadotropin-releasing hormone injection combined with human chorionic gonadotropin. *Fertil Steril* 62:1143–1149
6. Moul PJ, Rees LH, Besser GM 1982 Pulsatile gonadotropin secretion in hyperprolactinaemic amenorrhoea on the response to bromocriptine therapy. *Clin Endocrinol (Oxf)* 16:153–162

7. Cohen-Becker IR, Selmanoff M, Wise PM 1986 Hyperprolactinemia alters the frequency and amplitude of pulsatile luteinizing hormone secretion in the ovariectomized rat. *Neuroendocrinology* 42:328–333
8. Park SK, Keenan MW, Selmanoff M 1993 Graded hyperprolactinemia first suppresses LH pulse frequency and then pulse amplitude in castrated male rats. *Neuroendocrinology* 58:448–453
9. Park SK, Selmanoff M 1991 Dose-dependent suppression of post-castration luteinizing hormone secretion exerted by exogenous prolactin administration in male rats: a model for studying hyperprolactinemic hypogonadism. *Neuroendocrinology* 53:404–410
10. Fox SR, Hoefer MT, Bartke A, Smith MS 1987 Suppression of pulsatile LH secretion, pituitary GnRH receptor content and pituitary responsiveness to GnRH by hyperprolactinemia in the male rat. *Neuroendocrinology* 46:350–359
11. Koike K, Aono T, Miyake A, Tasaka K, Chatani F, Kurachi K 1984 Effect of pituitary transplants on the LH-RH concentrations in the medial basal hypothalamus and hypophyseal portal blood. *Brain Res* 301:253–258
12. Sarkar DK, Yen SS 1985 Hyperprolactinemia decreases the luteinizing hormone-releasing hormone concentration in pituitary portal plasma: a possible role for β -endorphin as a mediator. *Endocrinology* 116:2080–2084
13. Fox SR, Smith MS 1984 The suppression of pulsatile luteinizing hormone secretion during lactation in the rat. *Endocrinology* 115:2045–2051
14. Smith MS, Grove KL 2002 Integration of the regulation of reproductive function and energy balance: lactation as a model. *Front Neuroendocrinol* 23:225–256
15. Xu J, Kirigiti MA, Cowley MA, Grove KL, Smith MS 2009 Suppression of basal spontaneous gonadotropin-releasing hormone neuronal activity during lactation: role of inhibitory effects of neuropeptide Y. *Endocrinology* 150:333–340
16. Smith MS 1982 Effect of pulsatile gonadotropin-releasing hormone on the release of luteinizing hormone and follicle-stimulating hormone *in vitro* by anterior pituitaries from lactating and cycling rats. *Endocrinology* 110:882–891
17. Tortonese DJ, Brooks J, Ingleton PM, McNeilly AS 1998 Detection of prolactin receptor gene expression in the sheep pituitary gland and visualization of the specific translation of the signal in gonadotrophs. *Endocrinology* 139:5215–5223
18. Henderson HL, Townsend J, Tortonese DJ 2008 Direct effects of prolactin and dopamine on the gonadotroph response to GnRH. *J Endocrinol* 197:343–350
19. Polson DW, Sagle M, Mason HD, Adams J, Jacobs HS, Franks S 1986 Ovulation and normal luteal function during LHRH treatment of women with hyperprolactinaemic amenorrhoea. *Clin Endocrinol (Oxf)* 24:531–537
20. Lecomte P, Lecomte C, Lansac J, Gallier J, Sonier CB, Simonetta C 1997 Pregnancy after intravenous pulsatile gonadotropin-releasing hormone in a hyperprolactinaemic woman resistant to treatment with dopamine agonists. *Eur J Obstet Gynecol Reprod Biol* 74:219–221
21. Grattan DR, Jasoni CL, Liu X, Anderson GM, Herbison AE 2007 Prolactin regulation of gonadotropin-releasing hormone neurons to suppress luteinizing hormone secretion in mice. *Endocrinology* 148:4344–4351
22. Bakowska JC, Morrell JI 1997 Atlas of the neurons that express mRNA for the long form of the prolactin receptor in the forebrain of the female rat. *J Comp Neurol* 386:161–177
23. Brown RS, Kokay IC, Herbison AE, Grattan DR 2010 Distribution of prolactin-responsive neurons in the mouse forebrain. *J Comp Neurol* 518:92–102
24. Herbison AE 2008 Estrogen positive feedback to gonadotropin-releasing hormone (GnRH) neurons in the rodent: the case for the rostral periventricular area of the third ventricle (RP3V). *Brain Res Rev* 57:277–287
25. Lee WS, Smith MS, Hoffman GE 1990 Luteinizing hormone-releasing hormone neurons express Fos protein during the proestrous surge of luteinizing hormone. *Proc Natl Acad Sci USA [Erratum (1990) 87:8185]* 87:5163–5167
26. Wintermantel TM, Campbell RE, Porteous R, Bock D, Gröne HJ, Todman MG, Korach KS, Greiner E, Pérez CA, Schütz G, Herbison AE 2006 Definition of estrogen receptor pathway critical for estrogen positive feedback to gonadotropin-releasing hormone neurons and fertility. *Neuron* 52:271–280
27. Le WW, Berghorn KA, Rassnick S, Hoffman GE 1999 Periventricular preoptic area neurons coactivated with luteinizing hormone (LH)-releasing hormone (LHRH) neurons at the time of the LH surge are LHRH afferents. *Endocrinology* 140:510–519
28. Jarry H, Leonhardt S, Wuttke W 1991 γ -Aminobutyric acid neurons in the preoptic/anterior hypothalamic area synchronize the phasic activity of the gonadotropin-releasing hormone pulse generator in ovariectomized rats. *Neuroendocrinology* 53:261–267
29. Herbison AE 1998 Multimodal influence of estrogen upon gonadotropin-releasing hormone neurons. *Endocr Rev* 19:302–330
30. Smith MJ, Jennes L 2001 Neural signals that regulate GnRH neurons directly during the oestrous cycle. *Reproduction* 122:1–10
31. Leranath C, MacLusky NJ, Sakamoto H, Shanabrough M, Naftolin F 1985 Glutamic acid decarboxylase-containing axons synapse on LHRH neurons in the rat medial preoptic area. *Neuroendocrinology* 40:536–539
32. Petersen SL, McCrone S, Coy D, Adelman JP, Mahan LC 1993 GABA_A receptor subunit mRNAs in cells of the preoptic area: colocalization with LHRH mRNA using dual label *in situ* hybridization histochemistry. *Endocr J* 1:29–34
33. Jung H, Shannon EM, Fritschy JM, Ojeda SR 1998 Several GABAA receptor subunits are expressed in LHRH neurons of juvenile female rats. *Brain Res* 780:218–229
34. Herbison AE, Pape JR 2001 New evidence for estrogen receptors in gonadotropin-releasing hormone neurons. *Front Neuroendocrinol* 22:292–308
35. DeFazio RA, Heger S, Ojeda SR, Moenter SM 2002 Activation of A-type γ -aminobutyric acid receptors excites gonadotropin-releasing hormone neurons. *Mol Endocrinol* 16:2872–2891
36. Han SK, Abraham IM, Herbison AE 2002 Effect of GABA on GnRH neurons switches from depolarization to hyperpolarization at puberty in the female mouse. *Endocrinology* 143:1459–1466
37. Sullivan SD, Moenter SM 2004 γ -Aminobutyric acid neurons integrate and rapidly transmit permissive and inhibitory metabolic cues to gonadotropin-releasing hormone neurons. *Endocrinology* 145:1194–1202
38. Chen P, Moenter SM 2009 GABAergic transmission to gonadotropin-releasing hormone (GnRH) neurons is regulated by GnRH in a concentration-dependent manner engaging multiple signaling pathways. *J Neurosci* 29:9809–9818
39. Seminara SB, Crowley Jr WF 2008 Kisspeptin and GPR54: discovery of a novel pathway in reproduction. *J Neuroendocrinol* 20:727–731
40. Irwig MS, Fraley GS, Smith JT, Acohido BV, Popa SM, Cunningham MJ, Gottsch ML, Clifton DK, Steiner RA 2004 Kisspeptin activation of gonadotropin releasing hormone neurons and regulation of KISS-1 mRNA in the male rat. *Neuroendocrinology* 80:264–272
41. Han SK, Gottsch ML, Lee KJ, Popa SM, Smith JT, Jakawich SK, Clifton DK, Steiner RA, Herbison AE 2005 Activation of gonadotropin-releasing hormone neurons by kisspeptin as a neuroendocrine switch for the onset of puberty. *J Neurosci* 25:11349–11356
42. Messenger S, Chatzidakis EE, Ma D, Hendrick AG, Zahn D, Dixon J, Thresher RR, Malinge I, Lomet D, Carlton MB, Colledge WH, Caraty A, Aparicio SA 2005 Kisspeptin directly stimulates gonadotropin-releasing hormone release via G protein-coupled receptor 54. *Proc Natl Acad Sci USA* 102:1761–1766
43. Gottsch ML, Cunningham MJ, Smith JT, Popa SM, Acohido BV, Crowley WF, Seminara S, Clifton DK, Steiner RA 2004 A role for kisspeptins in the regulation of gonadotropin secretion in the mouse. *Endocrinology* 145:4073–4077

44. Clarkson J, Han SK, Liu X, Lee K, Herbison AE 2010 Neurobiological mechanisms underlying kisspeptin activation of gonadotropin-releasing hormone (GnRH) neurons at puberty. *Mol Cell Endocrinol* 324:45–50
45. Smith JT, Popa SM, Clifton DK, Hoffman GE, Steiner RA 2006 Kiss1 neurons in the forebrain as central processors for generating the preovulatory luteinizing hormone surge. *J Neurosci* 26:6687–6694
46. Clarkson J, d'Anglemont de Tassigny X, Moreno AS, Colledge WH, Herbison AE 2008 Kisspeptin-GPR54 signaling is essential for preovulatory gonadotropin-releasing hormone neuron activation and the luteinizing hormone surge. *J Neurosci* 28:8691–8697
47. Castellano JM, Navarro VM, Roa J, Pineda R, Sánchez-Garrido MA, García-Galiano D, Vigo E, Dieguez C, Aguilar E, Pinilla L, Tena-Sempere M 2009 Alterations in hypothalamic Kiss-1 system in experimental diabetes: early changes and functional consequences. *Endocrinology* 150:784–794
48. Clarkson J, d'Anglemont de Tassigny X, Colledge WH, Caraty A, Herbison AE 2009 Distribution of kisspeptin neurones in the adult female mouse brain. *J Neuroendocrinol* 21:673–682
49. Smith JT, Cunningham MJ, Rissman EF, Clifton DK, Steiner RA 2005 Regulation of Kiss1 gene expression in the brain of the female mouse. *Endocrinology* 146:3686–3692
50. Lerant A, Freeman ME 1998 Ovarian steroids differentially regulate the expression of prolactin receptors in neuroendocrine dopaminergic neuron populations—a double-label confocal microscopic study. *Brain Research* 802:141–154
51. Kokay IC, Grattan DR 2005 Expression of mRNA for prolactin receptor (long form) in dopamine and pro-opiomelanocortin neurones in the arcuate nucleus of non-pregnant and lactating rats. *J Neuroendocrinol* 17:827–835
52. Pi XJ, Grattan DR 1998 Distribution of prolactin receptor immunoreactivity in the brain of estrogen-treated, ovariectomized rats. *J Comp Neurol* 394:462–474
53. Paxinos GWC 2005 *The rat brain in stereotaxic coordinates*. 5th ed. Burlington, MA: Elsevier Academic Press
54. King JC, Tobet SA, Snavely FL, Arimura AA 1982 LHRH immunopositive cells and their projections to the median eminence and organum vasculosum of the lamina terminalis. *J Comp Neurol* 209:287–300
55. Feldblum S, Erlander MG, Tobin AJ 1993 Different distributions of GAD65 and GAD67 mRNAs suggest that the two glutamate decarboxylases play distinctive functional roles. *J Neurosci Res* 34:689–706
56. Esclapez M, Tillakaratne NJ, Tobin AJ, Houser CR 1993 Comparative localization of mRNAs encoding two forms of glutamic acid decarboxylase with nonradioactive in situ hybridization methods. *J Comp Neurol* 331:339–362
57. Okamura H, Abitbol M, Julien JF, Dumas S, Béroed A, Geffard M, Kitahama K, Bobillier P, Mallet J, Wiklund L 1990 Neurons containing messenger RNA encoding glutamate decarboxylase in rat hypothalamus demonstrated by in situ hybridization, with special emphasis on cell groups in medial preoptic area, anterior hypothalamic area and dorsomedial hypothalamic nucleus. *Neuroscience* 39:675–699
58. Curran-Rauhut MA, Petersen SL 2002 Regulation of glutamic acid decarboxylase 65 and 67 gene expression by ovarian steroids: identification of two functionally distinct populations of GABA neurones in the preoptic area. *J Neuroendocrinol* 14:310–317
59. Kauffman AS, Gottsch ML, Roa J, Byquist AC, Crown A, Clifton DK, Hoffman GE, Steiner RA, Tena-Sempere M 2007 Sexual differentiation of Kiss1 gene expression in the brain of the rat. *Endocrinology* 148:1774–1783
60. Chiu S, Wise PM 1994 Prolactin receptor mRNA localization in the hypothalamus by in situ hybridization. *J Neuroendocrinol* 6:191–199
61. Ottem EN, Godwin JG, Krishnan S, Petersen SL 2004 Dual-phenotype GABA/glutamate neurons in adult preoptic area: sexual dimorphism and function. *J Neurosci* 24:8097–8105
62. Maffucci JA, Gore AC 2009 Chapter 2: hypothalamic neural systems controlling the female reproductive life cycle gonadotropin-releasing hormone, glutamate, and GABA. *Int Rev Cell Mol Biol* 274:69–127
63. Sliwowska JH, Billings HJ, Goodman RL, Lehman MN 2006 Immunocytochemical colocalization of GABA-B receptor subunits in gonadotropin-releasing hormone neurons of the sheep. *Neuroscience* 141:311–319
64. Sagrillo CA, Grattan DR, McCarthy MM, Selmanoff M 1996 Hormonal and neurotransmitter regulation of GnRH gene expression and related reproductive behaviors. *Behav Genet* 26:241–277
65. Han SK, Todman MG, Herbison AE 2004 Endogenous GABA release inhibits the firing of adult gonadotropin-releasing hormone neurons. *Endocrinology* 145:495–499
66. Moenter SM, DeFazio RA 2005 Endogenous γ -aminobutyric acid can excite gonadotropin-releasing hormone neurons. *Endocrinology* 146:5374–5379
67. Watanabe M, Sakuma Y, Kato M 2009 GABAA receptors mediate excitation in adult Rat GnRH neurons. *Biol Reprod* 81:327–332
68. Romanò N, Lee K, Abrahám IM, Jasoni CL, Herbison AE 2008 Nonclassical estrogen modulation of presynaptic GABA terminals modulates calcium dynamics in gonadotropin-releasing hormone neurons. *Endocrinology* 149:5335–5344
69. Constantin S, Caraty A, Wray S, Duittoz AH 2009 Development of gonadotropin-releasing hormone-1 secretion in mouse nasal explants. *Endocrinology* 150:3221–3227
70. Locatelli V, Apud JA, Gudelsky GA, Cocchi D, Masotto C, Casanueva F, Racagni G, Müller EE 1985 Prolactin in cerebrospinal fluid increases the synthesis and release of hypothalamic γ -aminobutyric acid. *J Endocrinol* 106:323–328
71. Felman K, Tappaz M 1989 GABAergic biochemical parameters of the tuberoinfundibular neurons following chronic hyperprolactinemia. *Neuroendocrinology* 49:580–585
72. Kolbinger W, Beyer C, Föhr K, Reisert I, Pilgrim C 1992 Diencephalic GABAergic neurons in vitro respond to prolactin with a rapid increase in intracellular free calcium. *Neuroendocrinology* 56:148–152
73. Grattan DR, Selmanoff M 1994 Prolactin- and testosterone-induced inhibition of LH secretion after orchidectomy: role of preoptic and tuberoinfundibular γ -aminobutyric acidergic neurones. *J Endocrinol* 143:165–174
74. Kornblatt JJ, Grattan DR 2001 Lactation alters γ -aminobutyric acid neuronal activity in the hypothalamus and cerebral cortex in the rat. *Neuroendocrinology* 73:175–184
75. Qureshi GA, Hansen S, Södersten P 1987 Offspring control of cerebrospinal fluid GABA concentrations in lactating rats. *Neurosci Lett* 75:85–88
76. Clarkson J, Herbison AE 2006 Postnatal development of kisspeptin neurons in mouse hypothalamus; sexual dimorphism and projections to gonadotropin-releasing hormone neurons. *Endocrinology* 147:5817–5825
77. Adachi S, Yamada S, Takatsu Y, Matsui H, Kinoshita M, Takase K, Sugiura H, Ohtaki T, Matsumoto H, Uenoyama Y, Tsukamura H, Inoue K, Maeda K 2007 Involvement of anteroventral periventricular metastin/kisspeptin neurons in estrogen positive feedback action on luteinizing hormone release in female rats. *J Reprod Dev* 53:367–378
78. Ramaswamy S, Guerriero KA, Gibbs RB, Plant TM 2008 Structural interactions between kisspeptin and GnRH neurons in the mediobasal hypothalamus of the male rhesus monkey (*Macaca mulatta*) as revealed by double immunofluorescence and confocal microscopy. *Endocrinology* 149:4387–4395
79. Burke MC, Letts PA, Krajewski SJ, Rance NE 2006 Coexpression of dynorphin and neurokinin B immunoreactivity in the rat hypothalamus: morphologic evidence of interrelated function within the arcuate nucleus. *J Comp Neurol* 498:712–726

80. Navarro VM, Gottsch ML, Chavkin C, Okamura H, Clifton DK, Steiner RA 2009 Regulation of gonadotropin-releasing hormone secretion by kisspeptin/dynorphin/neurokinin B neurons in the arcuate nucleus of the mouse. *J Neurosci* 29:11859–11866
81. Krajewski SJ, Burke MC, Anderson MJ, McMullen NT, Rance NE 2010 Forebrain projections of arcuate neurokinin B neurons demonstrated by anterograde tract-tracing and monosodium glutamate lesions in the rat. *Neuroscience* 166:680–697
82. Yamada S, Uenoyama Y, Kinoshita M, Iwata K, Takase K, Matsui H, Adachi S, Inoue K, Maeda KI, Tsukamura H 2007 Inhibition of metastin (kisspeptin-54)-GPR54 signaling in the arcuate nucleus-median eminence region during lactation in rats. *Endocrinology* 148:2226–2232
83. Xu J, Kirigiti MA, Grove KL, Smith MS 2009 Regulation of food intake and gonadotropin-releasing hormone/luteinizing hormone during lactation: role of insulin and leptin. *Endocrinology* 150:4231–4240
84. Bogusz AL, Hardy SL, Lehman MN, Connors JM, Hileman SM, Sliwowska JH, Billings HJ, McManus CJ, Valent M, Singh SR, Nestor CC, Coolen LM, Goodman RL 2008 Evidence that γ -aminobutyric acid is part of the neural circuit mediating estradiol negative feedback in anestrous ewes. *Endocrinology* 149:2762–2772
85. Fitzsimmons MD, Olschowka JA, Wiegand SJ, Hoffman GE 1992 Interaction of opioid peptide-containing terminals with dopaminergic perikarya in the rat hypothalamus. *Brain Research* 581:10–18
86. Arbogast LA, Voegt JL 1998 Endogenous opioid peptides contribute to suckling-induced prolactin release by suppressing tyrosine hydroxylase activity and messenger ribonucleic acid levels in tuberoinfundibular dopaminergic neurons. *Endocrinology* 139:2857–2862
87. Callahan P, Baumann MH, Rabii J 1996 Inhibition of tuberoinfundibular dopaminergic neural activity during suckling: involvement of mu and κ opiate receptor subtypes. *J Neuroendocrinol* 8:771–776
88. Manzanares J, Wagner EJ, Lookingland KJ, Moore KE 1992 Effects of immunoneutralization of dynorphin1–17 and dynorphin1–8 on the activity of central dopaminergic neurons in the male rat. *Brain Res* 587:301–305
89. Andrews ZB, Grattan DR 2003 Opioid receptor subtypes involved in the regulation of prolactin secretion during pregnancy and lactation. *J Neuroendocrinol* 15:227–236
90. Szawka RE, Ribeiro AB, Leite CM, Helena CV, Franci CR, Anderson GM, Hoffman GE, Anselmo-Franci JA 2010 Kisspeptin regulates prolactin release through hypothalamic dopaminergic neurons. *Endocrinology* 151:3247–3257
91. Grattan DR, Kokay IC 2008 Prolactin: a pleiotropic neuroendocrine hormone. *J Neuroendocrinol* 20:752–763
92. Thapa S, Short RV, Potts M 1988 Breast feeding, birth spacing and their effects on child survival. *Nature* 335:679–682



Go to the *Translational Research in Endocrinology & Metabolism* site for a collection of articles from The Endocrine Society journals

www.endojournals.org/trem