

Identification of siderophore producing and cynogenic fluorescent *Pseudomonas* and a simple confrontation assay to identify potential bio-control agent for collar rot of chickpea

Anil S. Kotasthane¹ · Toshy Agrawal¹ · Najam Waris Zaidi² · U. S. Singh²

Received: 1 August 2016 / Accepted: 1 February 2017 / Published online: 8 June 2017
© Springer-Verlag Berlin Heidelberg 2017

Abstract In soil, plant roots coexist with bacteria and fungi that produce siderophores capable of sequestering the available iron. Microbial cyanogenesis has been demonstrated in many species of fungi and in a few species of bacteria (e.g., *Chromobacterium* and *Pseudomonas*). Fluorescent *Pseudomonas* isolates P29, P59, P144, P166, P174, P187, P191 and P192 were cyanogenic and produced siderophores in the presence of a strong chelator 8-Hydroxyquinoline (50 mg/l). A simple confrontation assay for identifying potential antagonists was developed. Fluorescent *Pseudomonas* isolates P66, P141, P144, P166 and P174 were antagonistic against both *Rhizoctonia solani* and *Sclerotium rolfsii*. Vigorous plant growth was observed following seed bacterization with P141, P200 and P240. In field experiments, seed bacterization with selected bacterial isolates resulted in reduced collar rot (*S. rolfsii*) incidence.

Keywords Fluorescent Pseudomonads · Collar rot · HCN · Siderophores · Confrontation assays

Introduction

Rhizosphere inhabiting fluorescent Pseudomonads are one of the most dominant and potentially most promising group of plant growth promoting rhizobacteria involved in the bio-control of plant diseases (Haas and Défago 2005; Glick

2014). They maintain soil health by employing a wide variety of mechanisms, including nitrogen fixation, enhanced solubilization of phosphate and phytohormone production that positively impact plant health (such as auxins and cytokinins) (Penrose and Glick 2003; Mirza et al. 2006). *Pseudomonas* spp. produce an arsenal of antimicrobials (including hydrogen cyanide (HCN), pyoluteorin, phenazines, pyrrolnitrin, siderophores, cyclic lipopeptides and 2,4-diacetylphloroglucinol (DAPG) (Thomashow and Weller 1996; Weller 2007). They also are able to promote plant growth and induce systemic resistance (ISR) in plants (Raaijmakers et al. 2009; Glick 2014). In the present study we evaluate fluorescent *Pseudomonas* isolates for siderophore (CAS assay-plate screening, CAS assay-spectrophotometric analysis, hydroxyquinoline test, tetrazolium test, FeCl₃ test and Arnow's assay) and HCN production. *Pseudomonas* spp. have been employed efficiently as commercial biocontrol agents (Loper and Lindow 1987; Walsh et al. 2001). However, there is always a scope for isolating better, locally adapted strains for deployment as biocontrol agents. Hence, we have screened local isolates of fluorescent Pseudomonads for developing formulation and possible commercialization for the management of collar rot of chickpea (*S. rolfsii* Sacc), one of the major biotic factors contributing towards low production (55–95% mortality of chickpea seedlings).

Materials and methods

Microorganisms and culture conditions

The experimental material consisted of purified 29 isolates of fluorescent *Pseudomonas* spp. from soils (rhizospheric and non-rhizospheric) of different geographical locations

✉ Anil S. Kotasthane
kotasthaneaj@yahoo.com

¹ Department of Plant Molecular Biology and Biotechnology, Indira Gandhi Krishi Vishwavidyalaya, Krishak Nagar, Raipur, CG 492006, India

² IRRRI India Office, New Delhi, India

of Chhattisgarh. Isolation of fluorescent pseudomonads was done by adopting serial dilution method on King's B (KB) medium. After incubation at 28 °C for 2 days, fluorescent pseudomonad colonies from plates were identified under UV light (366 nm). Isolates were characterized on the basis of biochemical tests as per the procedures outlined in Bergey's Manual of Systematic Bacteriology (Sneath et al. 1986). Isolated colonies of fluorescent *Pseudomonas* were further streaked onto KB agar plates to obtain pure cultures. The isolates were maintained (at -80 °C on King's B broth (Himedia) containing 30% (v/v) glycerol) in the culture collections of the Department of Plant Molecular Biology and Biotechnology, Indira Gandhi Krishi Vishwavidyalaya, Raipur, Chhattisgarh, India, and revived on King's B slants when required.

Siderophore production

Siderophore production (qualitative and quantitative) was determined by (CAS assay (Schwyn and Neilands 1987). Specific tests were carried out for identification of hydroxamate and Catecholate types of siderophores following the standard methods (Arnou 1937). Chrome azurol S solution was prepared and added to melted King's B agar medium in the ratio 1:15. Spot inoculation at the centre of the CAS plate was done from actively growing cultures of *Pseudomonas*. Colonies exhibiting an orange halo after 3 days of incubation (28 ± 2 °C) were considered positive for siderophore production and the diameter of the orange halo was measured.

Hydroxyquinoline mediated siderophore test

For selection of *Pseudomonas* isolates with high ability to siderophores, isolates were inoculated on King's B medium supplemented with a strong chelator 8-Hydroxyquinoline (50 mg/l) (De Brito et al. 1995). Inoculated isolates were incubated at 28 ± 2 °C for 48–72 h; only those bacteria that produce a more avid iron chelator will grow.

Arnou's assay

Arnou's assay was used for quantification of catechol type siderophore. For qualitative estimation of siderophores, actively growing cultures of *Pseudomonas* were inoculated to 20 ml King's B medium in 50 ml tubes and incubated for 3 days at 28 ± 2 °C. The bacterial cells were removed by centrifugation at 3000 rpm for 5 min. Three ml of the culture supernatant was then mixed with 0.3 ml of 5 N HCl solution, 1.5 ml of Arnou's reagent (10 g NaNO₂, 10 g Na₂MoO₄·2H₂O dissolved in 50 ml distilled water) and 0.3 ml of 10 N NaOH. After 10 min the presence or absence of pink colour was observed and noted.

Tetrazolium test

This test is based on the capacity of hydroxamic acid to reduce tetrazolium salt by hydrolysis of hydroxamate groups using a strong alkali. The reduction and release of alkali shows red colour to a pinch of tetrazolium salt when 1–2 drops of 2 N NaOH and 0.1 ml of test sample are added. Instant appearance of a deep red colour indicated the presence of hydroxamate siderophore.

FeCl₃ test

One ml of the culture supernatant was mixed with freshly prepared 0.5 ml of 2% aqueous FeCl₃ and observed for the presence and absence of deep red colour.

HCN production

The production of HCN was estimated by the method of Wei et al. (1991). The cultures were grown on KM plates supplemented with 4.4 g/l glycine as a precursor and the filter paper strips soaked in saturated picric acid solution were exposed to the growing *Pseudomonas* isolates. The plates were incubated for 7 days at 28 ± 2 °C and observations were recorded as change in the colour of filter paper to brown as positive indicator for HCN production.

Confrontation assay

Fluorescent *Pseudomonas* isolates were multiplied on King's B broth and incubated for 2 days at 28 °C till the fluorescent pigment appeared in the broth. Petri-plates containing pre-sterilized potato dextrose agar (PDA) medium were inoculated with plant pathogenic fungi *Sclerotium rolfsii* or *Rhizoctonia solani* (in the centre) and incubated at 25 °C for 3 days till the fungus completely covered the entire plate. Bipartite interactions were performed following a simple confrontation assay which was developed during the course of investigation. To identify prospective bio-agent, a simple confrontation assay was developed wherein edge of glass funnel was deployed for bio-agent inoculum deposition surrounding pre-inoculated fungal pathogen. The edge of a glass funnel was sterilized by dipping in alcohol followed by flaming. Broth containing young growing cell (3-day-old) of fluorescent *Pseudomonas* was dispensed in sterile petri dish and picked at the edge of the funnel by dipping. Care was taken to remove the excess inoculum by gently shaking the funnel. Inoculation was done by gently touching the edge of the funnel (containing fluorescent *Pseudomonas*) which encircled the pre-inoculated plant pathogenic fungi on agar plug




Fig. 1 Confrontation assay

equidistantly (Fig. 1). Inhibition zone was measured after 72 h of incubation at 28 ± 2 °C. Percent inhibition of pathogens by *Pseudomonas* isolates over control was calculated using the formula of (Vincent 1947): $[(\text{Growth of pathogen in control} - \text{growth of pathogen with } Pseudomonas \text{ isolate}) / \text{growth of pathogen in control}] \times 100$. Proposed technique has the following advantages: (1) uniform inoculum deposition during all combinations of bipartite interactions. (2) Replica-plating can be done of the inoculum picked on the edge of the funnel. (3) Ability to evaluate the antagonistic potential of a sporulating bio-inoculant (e.g. *Trichoderma* sp.). The only disadvantage with this technique is that for each bio-inoculant in the broth a separate plate is required to dispense the inoculums.

Field trial for testing selected fluorescent *Pseudomonas* isolates against chickpea collar rot

For field trials seed bacterization was done with stock cultures (used for confrontation assays) of selected fluorescent *Pseudomonas* isolate. Slurry for seed bacterization was prepared @ 5 ml of bacterial culture + 3 g of talcum powder/kg of chickpea seeds. Care was taken for uniform coating of all the seeds, which were dried in shade and then sown in a field naturally infested with *S. rolf sii*. Sowing was done in 2 M \times 7 M plots. Care was taken to carry out all normal agricultural practices. Observations were recorded on germination, plant vigour, mortality and bundle and grain weight.

Results

Qualitative and quantitative assay for siderophore production

In the present investigation, 29 isolates of *Pseudomonas* were screened by six different siderophore assays viz., CAS assay-plate screening, CAS assay-spectrophotometric analysis, hydroxyquinoline test, tetrazolium test, FeCl₃ test and Arnow's assay. All isolates exhibited an orange halo after 3 days of incubation (28 ± 2 °C) on CAS agar plate and, therefore, were considered positive for siderophore production. Intensity of orange halo and diameter showed wide variation among the isolates, which ranged from 11.50 (isolate P56) to 64.50 (isolate P191) mm. Isolates P29, P59, P66, P141, P144, P166, P174, P187, P191, P192, P200, P207 P229 and P260 were identified as producer of more avid iron chelator as they were tested +ve in King's B medium supplemented with a strong chelator 8-Hydroxyquinoline (50 mg/l). Three isolates (P7, P43 and P45) tested positive in Arnow's assay (detects catechol type of siderophores). We observed that isolates P43 and P45 tested -ve in HQ test but were +ve for other siderophore tests. All isolates produced deep red colour on addition of tetrazolium salt and NaOH indicating the production of hydroxamate type of siderophores (reduce tetrazolium salt by hydrolysis of hydroxamate group in presence of strong alkali). FeCl₃ test was positive for isolates P7, P23, P29, P43, P59, P66 P74, P80, P123, P141, P144, P174, P180, P187, P192, P200, P207 and P260. Only one isolate (P7)

Table 1 Quantitative and qualitative and estimation of siderophores by different tests, HCN production and antagonistic activity of fluorescent *Pseudomonas* isolates against *Rhizoctonia solani* and *Sclerotium rolfsii*

Isolates	Quantitative % siderophore units	Qualitative siderophore test					HCN production	% inhibition	
		CAS	Arnow's	FeCl ₃	Tetrazolium	HQ		<i>R. solani</i>	<i>S. rolfsii</i>
Fluorescent <i>Pseudomonas</i> isolates with high ACC deaminase activity (data not shown)									
P66	78.555 ± 0.39 ^a	63.5 ± 1.5 ^a		+ve	+ve	+++	-	68.89 ± 1.11 ^a	82.78 ± 2.78 ^b
P141	80.15 ± 0.15 ^a	55 ± 1 ^b		+ve	+ve	+	-	52.78 ± 2.78 ^c	43.335 ± 2.22 ^g
P200	49.475 ± 2.11 ^f	45.5 ± 1.5 ^{de}		+ve	+ve	+	-	60 ± 5.56 ^b	28.89 ± 1.11 ⁱ
P229	22.365 ± 1.32 ^l	36 ± 1 ^h				+	-	19.445 ± 2.78 ^j	28.885 ± 4.45 ⁱ
P260	71.05 ± 2.63 ^b	61.5 ± 0.5 ^a		+ve	+ve	++	-	37.775 ± 5.56 ^{gh}	32.775 ± 3.33 ^{hi}
Fluorescent <i>Pseudomonas</i> isolates ineffective against <i>R. solani</i> and <i>S. rolfsii</i>									
P2	37.25 ± 1.06 ⁱ	35 ± 1 ^h			+ve	+	-	40.5 ± 0.5 ^{fg}	17.25 ± 0.55 ^j
P3	30.2 ± 0.28 ^j	52.5 ± 2.5 ^{bc}			+ve	+	-	32.95 ± 0.45 ^{hi}	17.1 ± 0.4 ^j
P43	70.6 ± 0.85 ^b	41 ± 1 ^{efg}	+ve	+ve	+ve	-	-	26.2 ± 1.2 ^{ij}	33.95 ± 0.65 ^{hi}
P45	24.15 ± 0.21 ^l	16.5 ± 1.5 ⁱ	+ve		+ve	-	+	13.05 ± 0.55 ^{kl}	25.1 ± 0.7 ⁱ
P130	47.5 ± 0.32 ^f	31.5 ± 1.5 ^{hi}			+ve	+	-	25.15 ± 0.15 ^{ij}	31.85 ± 0.75 ^{hi}
Fluorescent <i>Pseudomonas</i> isolates moderately effective against <i>R. solani</i> and <i>S. rolfsii</i>									
P163	47.5 ± 0.71 ^f	40 ± 2 ^{fgh}			+ve	-	-	21.825 ± 0.58 ^j	45.15 ± 0.75 ^g
P123	43.78 ± 0.31 ^g	39 ± 2 ^{gh}		+ve	+ve	++	-	33.15 ± 0.65 ^{hi}	34 ± 0.7 ^h
P150	43.78 ± 0.31 ^g	42.5 ± 2.5 ^{efg}			+ve	+	+	38.2 ± 0.7 ^{gh}	44.95 ± 0.55 ^g
P23	48.25 ± 0.35 ^f	42 ± 2 ^{efg}		+ve	+ve	-	-	39.375 ± 0.63 ^{fg}	45.05 ± 0.65 ^g
Fluorescent <i>Pseudomonas</i> isolates effective against <i>S. rolfsii</i> (non HCN producing)									
P56	27.5 ± 0.71 ^k	11.5 ± 1.5 ⁱ			+ve	-	-	10.8 ± 0.8 ^l	50.65 ± 0.65 ^f
P184	40.615 ± 0.87 ^h	40.5 ± 0.5 ^{fgh}		+ve	+ve	-	-	18 ± 0.5 ^{jk}	50.75 ± 0.75 ^f
P207	36.16 ± 0.23 ⁱ	51 ± 1 ^{bc}		+ve	+ve	+++	-	46.875 ± 0.63 ^{de}	72.8 ± 0.6 ^c
Fluorescent <i>Pseudomonas</i> isolates effective against <i>S. rolfsii</i> (HCN producing)									
P132	23.835 ± 0.23 ^l	41 ± 1 ^{efg}			+ve	+	+	13.15 ± 0.65 ^{kl}	61.95 ± 0.85 ^d
P7	52.9 ± 0.14 ^e	41.5 ± 1.5 ^{efg}	+ve	+ve	+ve	++	+	21.625 ± 0.38 ^j	61.8 ± 0.7 ^d
P192	36.115 ± 0.16 ⁱ	54.5 ± 1.5 ^b		+ve	+ve	+++	+	28.2 ± 0.7 ⁱ	56.25 ± 0.65 ^c
P191	47.945 ± 0.08 ^f	64.5 ± 1.5 ^a			+ve	+++	+	30.8 ± 0.8 ⁱ	72.6 ± 0.4 ^c
P59	60.5 ± 0.71 ^c	39.5 ± 1.5 ^{gh}		+ve	+ve	+++	+	36.625 ± 0.38 ^{gh}	47.3 ± 0.6 ^{fg}
P29	60.5 ± 0.71 ^c	54.5 ± 1.5 ^b		+ve	+ve	+++	+	43.85 ± 1.35 ^{ef}	87.3 ± 0.6 ^a
P187	60.115 ± 0.16 ^c	51.5 ± 1.5 ^{bc}		+ve	+ve	+++	+	44.325 ± 0.57 ^{ef}	74.05 ± 0.75 ^c
Fluorescent <i>Pseudomonas</i> isolates effective against <i>R. solani</i> (non HCN producing)									
P74	27.75 ± 0.35 ^k	40.5 ± 2.5 ^{fgh}		+ve	+ve	+	-	50.7 ± 0.7 ^{cd}	35.4 ± 0.3 ^h
P80	32.5 ± 0.71 ^j	s ± 1.5 ^{def}		+ve	+ve	++	-	63.1 ± 0.6 ^b	63.85 ± 0.55 ^d
Fluorescent <i>Pseudomonas</i> isolates effective against <i>R. solani</i> and <i>S. rolfsii</i> (HCN producing)									
P144	43.5 ± 0.71 ^g	48.5 ± 1.5 ^{cd}		+ve	+ve	+++	+	50.7 ± 0.7 ^{cd}	76 ± 0.4 ^c
P166	60.115 ± 0.16 ^c	40 ± 3 ^{fgh}			+ve	+++	+	73.1 ± 0.6 ^a	73 ± 0.8 ^c
P174	55.5 ± 0.71 ^d	51 ± 1 ^{bc}		+ve	+ve	+++	+	51 ± 1 ^{cd}	82.95 ± 0.75 ^b
Max.	80.15	64.5						73.1	87.3
Min.	22.365	11.5						10.8	17.1
CV	2.352	5.22						6.625	3.645
CD (0.01)	2.962	6.33						6.876	5.12
CD (0.05)	2.198	4.697						5.102	3.799

Table 1 continued

Isolates	Quantitative % siderophore units	Qualitative siderophore test					HCN production	% inhibition	
		CAS	Arnow's	FeCl ₃	Tetrazolium	HQ		<i>R. solani</i>	<i>S. rolfsii</i>
F cal	527.375**	53.39**					90.948**	247.998**	

Values are average of three replications; values after \pm represent standard deviation

As per Duncan's grouping means with the same letter are not significantly different

CV coefficient of variance, CD critical difference, HQ hydroxyquinoline test

+++ Luxuriant/high growth

++ Medium growth

+ Low growth

– No growth

** Values are significant at 1 and 5% levels

Table 2 Efficacy of selected fluorescent *Pseudomonas* isolates against collar rot and yield of chickpea

Isolate no.	No. of plants	Plant vigour		Collar rot incidence	Bundle weight (kg)	Yield (gm)
		% poor vigour	% high vigour	% wilted plants		
Control	256	92.820 ^a \pm 0.63	7.965 ^c \pm 0.155	35.015 ^a \pm 0.635	2.070 ^{cd} \pm 0.25	598.000 ^d \pm 25
P66	289	17.425 ^d \pm 0.465	83.585 ^b \pm 0.545	16.130 ^b \pm 0.21	3.670 ^{ab} \pm 0.45	1361.500 ^b \pm 59.5
P141	303	5.130 ^c \pm 0.51	96.070 ^a \pm 0.69	11.105 ^c \pm 0.545	4.890 ^a \pm 0.43	1732.500 ^a \pm 10.5
P200	350	21.625 ^c \pm 0.485	79.535 ^c \pm 0.675	7.275 ^d \pm 0.705	3.785 ^{ab} \pm 0.425	1085.000 ^c \pm 36
P229	305	92.955 ^a \pm 0.165	7.820 ^e \pm 0.61	2.930 ^e \pm 0.31	2.735 ^{bc} \pm 0.355	574.500 ^d \pm 14.5
P260	367	77.620 ^b \pm 0.51	14.000 ^d \pm 0.65	2.645 ^e \pm 0.465	1.425 ^d \pm 0.225	277.000 ^e \pm 34
CV		1.331	1.715	5.750	16.755	5.126
Fcal		7145.771**	5242.546**	570.035**	11.890**	262.364**
CD (0.01)		5.529	3.062	2.668	1.923	178.267
CD (0.05)		1.669	2.021	1.761	1.269	117.675

Values after \pm represent standard deviation

Superscript values indicate Duncan's grouping means with the same letter are not significantly different

** Values are significant at 1 and 5% levels

tested positive for all siderophore tests. Carboxylate type of siderophore was determined by spectrophotometric method at 630 nm and the percentage of siderophore unit ranged from 12.23 (isolate P130) to 70.60% (isolate P43) (Table 1).

Screening for hydrogen cyanide production


To identify cynogenic fluorescent *Pseudomonas*, isolates were inoculated on KMB plates supplemented with 4.4 g/l glycine (precursor molecule of HCN) and incubated for 7 days at 28 ± 2 °C. Development of brown colour on filter paper strips soaked in saturated picric acid solution indicated +ve for HCN production. Out of the 29 fluorescent *Pseudomonas* tested, 12 isolates (P7, P29, P45, P59, P132, P144, P150, P166, P174, P187, P191 and P192) tested positive for HCN producing ability. We observed a correlation between inhibitory effects observed following confrontation assays and the ability to produce HCN by

fluorescent *Pseudomonas*. Cynogenic (HCN producers) isolates P132, P7, P192, P191, P59, P29, P187 exerted strong antagonism against *S. rolfsii*, where as a non-cynogenic isolate P74 exerted strong inhibitory effects against *R. solani*. On the contrary, a noncynogenic (P80) and three cyanogenic (P144, P166, P174) fluorescent *Pseudomonas* exerted strong inhibitory effects against both the soilborne fungal pathogens *R. solani* and *S. rolfsii* (Table 1).

In vitro antagonistic activity of *Pseudomonas* isolates against *R. solani* and *S. rolfsii*

There were differences in the antagonistic abilities of fluorescent *Pseudomonas* isolates against both (*R. solani* and *S. rolfsii*) pathogens (Fig. 1; Table 1). All of the 29 isolates of fluorescent *Pseudomonas* showed different degree of growth inhibitions of *R. solani* and *S. rolfsii*, ranging from 10.8 to 73.1% and 17.1 to 87.3%, respectively.

Fig. 2 Efficacy of selected fluorescent *Pseudomonas* isolates against collar rot and yield of chickpea



Confrontation assays revealed P166 and P29 as potential antagonists against *R. solani* and *S. rolfsii*, respectively, while isolates P56 and P3 were ineffective. Fluorescent

Pseudomonas isolates P29, P187, P191, P192 and P207 exerted strong inhibitory effects on the mycelia growth of *S. rolfsii* whereas isolates P66, P141, P144, P166 and P174

expressed strong inhibitory effects on both *R. solani* and *S. rolf sii*.

Efficacy of selected fluorescent *Pseudomonas* isolates against root rot and yield of chickpea

Through confrontation assay, isolates P66 and P141 was identified to be having very strong antagonistic activities against *R. solani* and *S. rolf sii*, whereas P200 was antagonistic only against *R. solani*; other two isolates P229 and P260 were ineffective against both the pathogens. For field trials these five fluorescent *Pseudomonas* isolates were selected for their efficacy against collar rot disease and yield of chickpea. Frequency of infected plants was very high in untreated (control) plot, whereas seed treatment with bacterial isolates had lower incidence of collar rot (Table 2; Fig. 2). Vigorous plant growth was observed following seed bacterization with P141, P200, P240 (95.38, 83.04 and 78.86%, respectively), whereas poor vigour was observed with P229 (7.21%) and P260 (13.35%) (Table 2; Fig. 2). Bundle weight and yield indicated significant differences between control and treated plots. In the order of increasing bundle weight and yield per plot, the bioefficacy of the isolates were P260 < P229 < P66 < P200 < P141 (Table 2).

Discussion

Understanding of the mechanisms involved in the antagonist interactions between bacteria, pathogen and host plant is important for efficient utilization of these natural resources in crop health management. (Thomashow and Weller 1991). Siderophore production by strains of *Pseudomonas* spp., for plant disease control, is of great interest because of its possibilities in the substitution of chemical pesticides. Similarly, microbial cyanogenesis has been demonstrated in a few bacterial species (belonging to the genera *Pseudomonas*, *Chromobacterium*, *Rhizobium* and several cyanobacteria (Blumer and Haas 2000). Glycine has generally been used as a precursor of cyanide in fungi and bacteria (Brysk et al. 1969; Wissing 1974) and cyanogenesis is one of the mechanisms of antagonism and biocontrol properties (Haas and Défago 2005; Lanteigne et al. 2012). In the present study, we have compared the ability of several fluorescent *Pseudomonads* to produce siderophores, cyanogenesis and antagonism in plate assay. Our study revealed that the isolates vary in the mechanisms and ability to inhibit pathogens. During the study a simple confrontation assay technique was developed which was advantageous as compared to earlier reported techniques (Dennis and Webster 1971; Fokkema 1978; Santoyo et al. 2010), wherein bipartite interactions were performed on

media plates by streaking bacterial bio-agents (forming quadrant) and placing mycelial plug of 4 mm in the centre. Our combined in vitro and field data show the potential of isolates P66, P141 and P200 to be developed as a commercial bioagent for the control of chickpea collar rot, a perennial problem in chickpea production compounded by the lack of host resistance against the pathogen *S. rolf sii*.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest in the publication.

References

- Arnou LE (1937) Colorimetric determination of the components of 3, 4-dihydroxyphenylalanine-tyrosine mixtures. *J Biol Chem* 118:531–537
- Blumer C, Haas D (2000) Mechanism, regulation, and ecological role of bacterial cyanide biosynthesis. *Arch Microbiol* 173:170–177
- Brysk MM, Lauinger C, Ressler C (1969) Biosynthesis of cyanide from [2-¹⁴C¹⁵N] glycine in *chromobacterium violaceum*. *Biochim Biophys Acta (BBA) Gen Subj* 184:583–588
- De Brito AM, Gagne S, Antoun H (1995) Effect of compost on rhizosphere microflora of the tomato and on the incidence of plant growth-promoting rhizobacteria. *Appl Environ Microbiol* 61:194–199
- Dennis C, Webster J (1971) Antagonistic properties of species-groups of *Trichoderma*: I. Production of non-volatile antibiotics. *Trans Br Mycol Soc* 57:25–IN3
- Fokkema NJ (1978) Fungal antagonisms in the phyllosphere. *Ann Appl Biol* 89:115–119
- Glick BR (2014) Bacteria with ACC deaminase can promote plant growth and help to feed the world. *Microbiol Res* 169:30–39
- Haas D, Défago G (2005) Biological control of soil-borne pathogens by fluorescent *pseudomonads*. *Nat Rev Microbiol* 3:307–319. doi:10.1038/nrmicro1129
- Lanteigne C, Gadkar VJ, Wallon T et al (2012) production of DAPG and HCN by *Pseudomonas* sp. LBUM300 contributes to the biological control of bacterial canker of tomato. *Phytopathology* 102:967–973. doi:10.1094/PHYTO-11-11-0312
- Loper JE, Lindow SE (1987) Lack of evidence for the in situ fluorescent pigment production by *Pseudomonas syringae* pv. *syringae* on bean leaf surfaces. *Phytopathology* 77:1449–1454
- Mirza MS, Mehnaz S, Normand P et al (2006) Molecular characterization and PCR detection of a nitrogen-fixing *Pseudomonas* strain promoting rice growth. *Biol Fertil Soils* 43:163–170
- Penrose DM, Glick BR (2003) Methods for isolating and characterizing ACC deaminase-containing plant growth-promoting rhizobacteria. *Physiol Plant* 118:10–15
- Raaijmakers JM, Paulitz TC, Steinberg C et al (2009) The rhizosphere: a playground and battlefield for soilborne pathogens and beneficial microorganisms. *Plant Soil* 321:341–361
- Santoyo G, Valencia-Cantero E, Orozco-Mosqueda M del C et al (2010) Papel de los sideróforos en la actividad antagonista de *Pseudomonas fluorescens* zum80 hacia hongos fitopatógenos (Role of siderophores in antagonistic activity of *Pseudomonas fluorescens* zum80 against phytopatogens). *Terra Latinoam* 28:53–60
- Schwyn B, Neilands JB (1987) Universal chemical assay for the detection and determination of siderophores. *Anal Biochem* 160:47–56

- Sneath PHA, Mair NS, Sharpe ME, Holt JG (1986) Bergey's manual of systematic bacteriology, vol 2. Williams and Wilkins, Baltimore, MD, p 964
- Thomashow LS, Weller DM (1991) Role of antibiotics and siderophores in biocontrol of take-all disease of wheat. In: The rhizosphere and plant growth. Springer, Netherlands, pp 245–251
- Thomashow LS, Weller DM (1996) Current concepts in the use of introduced bacteria for biological disease control: mechanisms and antifungal metabolites. In: Plant-microbe interactions. Springer, US, pp 187–235
- Vincent JM (1947) Distortion of fungal hyphae in the presence of certain inhibitors. Nature 159:850
- Walsh UF, Morrissey JP, O'Gara F (2001) *Pseudomonas* for biocontrol of phytopathogens: from functional genomics to commercial exploitation. Curr Opin Biotechnol 12:289–295
- Wei G, Kloepper JW, Tuzun S (1991) Induction of systemic resistance of cucumber to *Colletotrichum orbiculare* by select strains of plant growth-promoting rhizobacteria. Phytopathology 81:1508–1512
- Weller DM (2007) *Pseudomonas* biocontrol agents of soilborne pathogens: looking back over 30 years. Phytopathology 97:250–256
- Wissing F (1974) Cyanide formation from oxidation of glycine by a *Pseudomonas* species. J Bacteriol 117:1289–1294