

ARTICLE

Received 29 Jan 2015 | Accepted 27 Mar 2015 | Published 11 May 2015

DOI: 10.1038/ncomms8055

IL-10-producing intestinal macrophages prevent excessive antibacterial innate immunity by limiting IL-23 synthesis

Petra Krause¹, Venetia Morris¹, Jason A. Greenbaum², Yoon Park³, Unni Bjoerheden¹, Zbigniew Mikulski⁴, Tracy Muffley¹, Jr-Wen Shui¹, Gisen Kim¹, Hilde Cheroutre¹, Yun-Cai Liu³, Bjoern Peters⁵, Mitchell Kronenberg¹ & Masako Murai¹

Innate immune responses are regulated in the intestine to prevent excessive inflammation. Here we show that a subset of mouse colonic macrophages constitutively produce the anti-inflammatory cytokine IL-10. In mice infected with *Citrobacter rodentium*, a model for enteropathogenic *Escherichia coli* infection in humans, these macrophages are required to prevent intestinal pathology. IL-23 is significantly increased in infected mice with a myeloid cell-specific deletion of IL-10, and the addition of IL-10 reduces IL-23 production by intestinal macrophages. Furthermore, blockade of IL-23 leads to reduced mortality in the context of macrophage IL-10 deficiency. Transcriptome and other analyses indicate that IL-10-expressing macrophages receive an autocrine IL-10 signal. Interestingly, only transfer of the IL-10 positive macrophages could rescue IL-10-deficient infected mice. Therefore, these data indicate a pivotal role for intestinal macrophages that constitutively produce IL-10, in controlling excessive innate immune activation and preventing tissue damage after an acute bacterial infection.

¹Division of Developmental Immunology, La Jolla Institute for Allergy and Immunology, 9420 Athena Circle, La Jolla, California 92037, USA. ²Bioinformatics Core Facility, La Jolla Institute for Allergy and Immunology, 9420 Athena Circle, La Jolla, California 92037, USA. ³Division of Cell Biology, La Jolla Institute for Allergy and Immunology, 9420 Athena Circle, La Jolla, California 92037, USA. ⁴Microscopy Core Facility, La Jolla Institute for Allergy and Immunology, 9420 Athena Circle, La Jolla, California 92037, USA. ⁵Division of Vaccine Discovery, La Jolla Institute for Allergy and Immunology, 9420 Athena Circle, La Jolla, California 92037, USA. Correspondence and requests for materials should be addressed to M.K. (email: mitch@liai.org) or M.M. (email: murai@liai.org).

Interleukin 10 (IL-10) is an immunoregulatory cytokine that limits mucosal immune responses and minimizes immunopathology. Indeed, mutations in the interleukin 10 (IL-10) receptor gene (*Il10r*) were found in some patients with early-onset colitis^{1,2}. Similarly, mice deficient for IL-10 (*Il10*^{-/-}) or IL-10R (*Il10rb*^{-/-}) developed spontaneous inflammation of the large intestine, a process that was T-cell dependent and dominated by a pathogenic T helper type 1 and type 17 (Th1 and Th17) immune responses³⁻⁵.

A number of cell types can produce IL-10, including lymphocytes, monocytes, macrophages, mast cells, keratinocytes and intestinal epithelial cells. In several colitis models, the role of T-cell-derived IL-10 has proven to be central⁶⁻⁸, and in fact, mice with a conditional deletion of IL-10 in the CD4⁺ T-cell subset develop spontaneous inflammation of the intestine, as do those with a deletion in regulatory T cells (Treg) mediated by a Foxp3 driven Cre (refs 9,10). However, we found IL-10 from macrophages plays an indispensable role in the maintenance of Foxp3 expression by Treg and their function in mice with colitis¹¹. Despite this, mice with a conditional deletion of IL-10 or IL-10R in myeloid cell subsets^{12,13}, or of IL-10 specifically in macrophages¹⁴, did not spontaneously develop aggressive colitis, although they were susceptible to excessive inflammation following systemic LPS exposure^{12,13}. These data suggest that myeloid cell derived IL-10 might have a pivotal role in controlling mucosal immune responses after bacterial infection.

In the context of acute mucosal infections, the lack of IL-10 could be protective because of an enhanced inflammatory response with increased IL-12, tumour necrosis factor (TNF) and other cytokines¹⁵⁻²⁰. However, the absence of IL-10 also could lead to excessive inflammation^{17,21-23}. Therefore, in the context of acute bacterial infection of the intestine, it remains to be determined which cell type(s) produce IL-10, if IL-10 is protective or harmful, and which pathways are activated or inhibited by IL-10 secretion. To explore these issues, we have analysed mice with cell type-specific deletion of *Il10* that were infected with *Citrobacter rodentium* (*C. rodentium*). This is a Gram-negative bacterium that is considered a model for enteropathogenic and enterohemorrhagic *Escherichia coli* infections in humans. *C. rodentium* causes attaching and effacing mechanism of epithelial infection, leading to intestinal inflammation and diarrhoea. The bacteria normally are cleared in wild-type mice due to the actions of innate and adaptive immunity and the intestinal inflammation ultimately resolves²⁴⁻²⁶. Here we show that a unique subset of macrophages in the colonic lamina propria that constitutively produces IL-10 plays a critical role in preventing excessive inflammation following acute bacterial infection by limiting innate immunity, and that a major pathway by which IL-10 acts is through controlling IL-23 production.

Results

Myeloid cell IL-10 is important for survival from *C. rodentium*.

To assess the function of IL-10 in regulating mucosal immune responses after intestinal bacterial challenge, we used *Il10*^{-/-}, *Il10rb*^{-/-} and wild-type (wt) recipient mice that were infected with a sublethal dose of *C. rodentium* by oral gavage. The onset and severity of colitis in *Il10*^{-/-} mice is strongly influenced by the genetic background³, and we used mice on the C57BL6/J background that are more protected from inflammation under steady-state conditions. Indeed, none of the groups developed spontaneous colitis in our mouse colony (Supplementary Fig. 1). However, we found that all *Il10*^{-/-} and *Il10rb*^{-/-} mice died 7–12 days after infection, whereas all wt mice survived (Fig. 1a). At day 6 after infection, *Il10*^{-/-} and *Il10rb*^{-/-} mice showed

severe colonic inflammation, characterized by epithelial cell destruction, infiltration of mononuclear cells (Supplementary Fig. 1). These indicate that the role of IL-10 in modulating the initial mucosal immune response against *C. rodentium* is indispensable for survival and mucosal damage.

To identify which cell type(s) produce the IL-10 that is essential for host protection, we analysed mice with cell type-specific deletions of the *Il10* gene. These included *Il10*^{fllox/fllox}*LysM-Cre*, with deletion targeted primarily to CD11b⁺ myeloid cells, including macrophages, monocytes and neutrophils, *Il10*^{fllox/fllox}*Cd4-Cre* targeting mostly CD4⁺ T cells, and *Il10*^{fllox/fllox}*Cd11c-Cre* mice, which acts predominantly in dendritic cells (DCs). We used *Il10*^{fllox/fllox}*Cre*⁻ (*Il10*^{fllox/fllox}) littermates for controls. We found that *Il10*^{fllox/fllox}*LysM-Cre* mice started to die early after *C. rodentium* infection, with less than half surviving beyond day 7. In contrast, most of the *Il10*^{fllox/fllox}*Cd11c-Cre* and all of the *Il10*^{fllox/fllox}*Cd4-Cre* recipients survived to 21 days after infection (Fig. 1b). The average histology scores of each group were similar before *C. rodentium* infection. With the dose of bacteria we used, at day 6 we observed mild inflammation in *Il10*^{fllox/fllox}*Cd4-Cre* and *Il10*^{fllox/fllox} recipients, manifested by epithelial cell hyperplasia and some cellular infiltration. However, *Il10*^{fllox/fllox}*LysM-Cre* recipients, most of which survived to day 6, developed significantly more severe colonic inflammation, characterized by epithelial cell destruction, infiltration of mononuclear cells and submucosal oedema in the large intestine (Fig. 1c,d). In parallel with this, we found increased colonic epithelial permeability and bacterial dissemination to the spleen after infection in *Il10*^{fllox/fllox}*LysM-Cre* recipients (Fig. 1e,f). Despite this, bacterial numbers in the faeces of *Il10*^{fllox/fllox}*LysM-Cre* recipients at day 6 after bacterial challenge were similar compared with *Il10*^{fllox/fllox}*Cd4-Cre* and *Il10*^{fllox/fllox} mice (Fig. 1g). These data suggest intestinal epithelial cell destruction by excessive mucosal immune responses after infection in *Il10*^{fllox/fllox}*LysM-Cre* recipients probably contributed to bacterial dissemination and lethality.

Macrophages are the principal IL-10-producing cells.

We assessed the effectiveness and specificity of the deletion of *Il10* in the various cell populations by measuring the decrease in *Il10* mRNA at steady state. Indeed, we detected a decrease of ~90% in *Il10* mRNA transcripts in the CD11c^{int}CD11b⁺ macrophage population from the colon of *Il10*^{fllox/fllox}*LysM-Cre* mice compared with *Il10*^{fllox/fllox} controls, with little decrease in CD11c⁺CD11b⁻ cells or CD4⁺ T cells (Supplementary Fig. 2). Similarly, we detected a specific and nearly complete loss of *Il10* transcripts in CD4⁺ T cells in *Il10*^{fllox/fllox}*Cd4-Cre* mice and a similar decrease in the CD11c⁺CD11b⁻ population in *Il10*^{fllox/fllox}*Cd11c-Cre* mice. There also was a 15–20% decrease in *Il10* mRNA in the CD11c^{int}CD11b⁺ population in *Il10*^{fllox/fllox}*Cd11c-Cre* mice (Supplementary Fig. 2). The intermediate level of CD11c expression by macrophages in the CD11c^{int}CD11b⁺ population likely is responsible for the incomplete disruption of IL-10 expression and the preservation of protective function in the *Il10*^{fllox/fllox}*Cd11c-Cre* strain. Therefore, we conclude that the absence of a strong effect on the survival and inflammation in the *Il10*^{fllox/fllox}*Cd4-Cre* and *Il10*^{fllox/fllox}*Cd11c-Cre* strains is not due to the inefficient removal of the *Il10* gene.

Because the *LysM* promoter drives Cre recombinase expression in macrophages, monocytes and neutrophils, we determined which of these cell types produced IL-10 after infection. Consistent with a recent report²⁰, IL-10 mRNA was most highly expressed in CD11b⁺CD11c^{int} macrophages, which also express F4/80 and MHC class II, but was undetectable in granulocytes (Fig. 1h) in the large intestine. Therefore, the




Figure 1 | Myeloid cell-derived IL-10 is crucial for survival after *Citrobacter rodentium* infection. (a) *Il10*^{-/-} (black circle), *Il10rb*^{-/-} (grey triangle) and C57BL/6J (black square) mice or (b) *Il10*^{flox/flox}*LysM*-Cre (black circle), *Il10*^{flox/flox}*Cd4*-Cre (open circle), *Il10*^{flox/flox}*Cd11c*-Cre (black triangle) and *Il10*^{flox/flox} (black square) mice were infected orally with *C. rodentium* and survival was monitored. (a) Two independent experiments with 6 mice per group and (b) 3 independent experiments with 12-17 mice per group are presented. (a,b) Log-rank test, ***P* < 0.01, ****P* < 0.001 (c,d) inflammation of middle colon was assessed by histology before and 6 days after infection. Scale bars, 100 μm. (e) Epithelial barrier permeability was determined before and 6 days after infection. (f,g) Titres of *C. rodentium* in f spleen and g faeces were quantified 6 days of infection. (e-g) Each symbol represents a measurement from a single mouse. Student's *t*-test, ***P* < 0.01, ****P* < 0.001 (h) Analysis of *Il10* mRNA in macrophages or granulocytes at day 3 after infection. Results were representative of two independent experiments. NS, non significant.

reduced survival in *Il10*^{flox/flox}*LysM*-Cre mice is very likely due to the loss of IL-10 production by macrophages.

A significant proportion of macrophages were IL-10 competent, GFP⁺ cells in the large intestine before infection. By flow cytometry, these cells were more frequent in the colon

compared with the small intestine (Supplementary Fig. 3a). When analysed by immunohistology, both the GFP⁺ (IL-10⁺) and GFP⁻ (IL-10⁻) macrophages were scattered from the top of the villi to the crypt in both the small and the large intestine (Supplementary Fig. 3b). This population is maintained through

day 6 of infection (Supplementary Fig. 4), although the frequency tended to decrease.

To confirm that IL-10-producing macrophages regulate mucosal innate immune responses independently of adaptive immunity, we analysed intestinal inflammation triggered by *C. rodentium* using *Rag*^{-/-} and *Il10*^{-/-}*Rag*^{-/-} mice. *Il10*^{-/-}*Rag*^{-/-} mice had more severe mortality and severe colonic inflammation at day 6 after infection compared with *Rag*^{-/-} mice (Supplementary Fig. 5). Together, these data suggest that macrophage-derived IL-10 plays a nonredundant role in limiting the innate mucosal immune responses against intestinal infection with *C. rodentium*.

Macrophage IL-10 regulates myeloid cell recruitment to the colon. To determine if macrophage-derived IL-10 regulates inflammatory cell recruitment to the large intestine after *C. rodentium* infection, we examined the kinetics of cell accumulation in the colon lamina propria in *Il10*^{fllox/fllox}*LysM-Cre*,

Il10^{fllox/fllox}*Cd4-Cre* and *Il10*^{fllox/fllox} recipients during *C. rodentium* infection, and we characterized the infiltrating cell types by flow cytometric analysis. The total cell number in the colon lamina propria was comparable in all the groups before *C. rodentium* injection, and CD11b⁺Ly6G⁻Ly6C^{high} monocytes were undetectable (Fig. 2a). Six days after infection, we detected a significant increase in total cell numbers in *Il10*^{fllox/fllox}*LysM-Cre* mice compared with *Il10*^{fllox/fllox}*Cd4-Cre* and *Il10*^{fllox/fllox} recipients, and the population in the large intestine lamina propria consisted mainly of CD11b⁺Ly6G⁻Ly6C^{high} monocytes and CD11b⁺Ly6G^{high}Ly6C^{low} granulocytes (Fig. 2a). There was no increase in CD4⁺ T cells in any of these groups at this early time point (Fig. 2a).

Because myeloid cell infiltrates were increased in infected *Il10*^{fllox/fllox}*LysM-Cre* mice, we tested if chemokines that would attract these cells were also increased. Because *Il10*^{fllox/fllox}*Cd4-Cre* and *Il10*^{fllox/fllox} recipients were similar in survival and degree of inflammation, we compared *Il10*^{fllox/fllox}*LysM-Cre* mice with *Il10*^{fllox/fllox}*Cd4-Cre* mice. Although CCL2 and CXCL1, known to




Figure 2 | Macrophage IL-10 regulates myeloid cell recruitment to the colon. *Il10*^{fllox/fllox}*LysM-Cre* (black), *Il10*^{fllox/fllox}*Cd4-Cre* (white) and *Il10*^{fllox/fllox} (grey) mice were analysed before and at day 6 after *C. rodentium* infection. (a) Lamina propria cells were isolated at the indicated times and total cell numbers were counted. Frequencies of CD4⁺ T cells (CD4⁺TCR β ⁺), monocytes (Ly6G⁻Ly6C^{high}) and granulocytes (Ly6G^{high}Ly6C^{low}) were measured by flow cytometry and cell numbers were calculated according to the respective total cell number. Cell numbers are presented as averages and s.d. from nine mice per group from three independent experiments. (b,c) At the indicated time points, 3-mm fragments of the middle colon of *Il10*^{fllox/fllox}*LysM-Cre* (black), *Il10*^{fllox/fllox}*Cd4-Cre* (white) mice were cultured for 24 h. Chemokine production was measured in the supernatants by Multiplex cytokine assay and normalized to the weight of the corresponding tissue fragments. Data are presented as averages with s.d. from two independent experiments with six mice in each group. Student's *t*-test, **P* < 0.05, ***P* < 0.01, ****P* < 0.001.

recruit monocytes and granulocytes, respectively, were similar between $Il10^{lox/lox}LysM-Cre$ mice and $Il10^{lox/lox}Cd4-Cre$ mice (Fig. 2b), other chemokines that recruit monocytes and neutrophils, including CCL3, CCL4 and CCL5, were significantly increased in colon fragment culture supernatants from $Il10^{lox/lox}LysM-Cre$ mice (Fig. 2c). This was true even before *C. rodentium* infection (Fig. 2c), although this increased level before infection apparently was not sufficiently high to recruit monocytes and granulocytes and cause detectable inflammation. These data suggest that macrophage-derived IL-10 regulates the chemokine-dependent recruitment of monocytes and granulocytes to the large intestine during intestinal bacterial infection.

Increased pro-inflammatory cytokines in $Il10^{lox/lox}LysM-Cre$ mice. We examined if the increased colitis in *C. rodentium*-infected $Il10^{lox/lox}LysM-Cre$ mice was associated with the excessive production of pro-inflammatory cytokines using a Bio-Plex multicytokine assay. As illustrated (Fig. 3), IL-10 production was reduced before infection and early after infection in $Il10^{lox/lox}LysM-Cre$ compared with $Il10^{lox/lox}Cd4-Cre$ mice, suggesting that the majority of IL-10 in the colon under these conditions is produced by macrophages. Conversely, the production of TNF and IL-12p40 was increased in $Il10^{lox/lox}LysM-Cre$ recipients in colon fragment cultures before and more strongly during infection, while IL-6 and interferon γ levels were comparable. We used a different method, enzyme-linked immunosorbent assay (ELISA), to explore the expression of IL-12 family cytokines further. We detected a striking increase in IL-23p19 by day 6 after infection in $Il10^{lox/lox}LysM-Cre$ mice. However neither IL-23 nor IL-12p70 were significantly increased in $Il10^{lox/lox}LysM-Cre$ before infection, suggesting that IL-12p40 monomers or dimers, known to be produced in other contexts^{27,28}, likely were produced in the intestine under the lack of macrophage-derived IL-10, although the function of IL-12p40 homodimers is uncertain.

It has been reported that IL-23 induces IL-22 production from innate lymphoid cells^{29,30}. Indeed, consistent with the significant

increase in IL-23 by day 6 after infection, in colonic fragment cultures from infected $Il10^{lox/lox}LysM-Cre$ mice, we detected a profound increase in IL-22 production as well (Fig. 3).

Macrophage IL-10 influences mortality through the regulation of IL-23p19. We assessed if IL-10 acts on the IL-23p19-producing cells activated by *C. rodentium*. It has been reported IL-23 is produced by macrophages and DCs^{29,31,32}. To determine the dominant source of IL-23-producing cells, we sorted $CD11b^+CD11c^{int}F4/80^+$ colonic macrophages and $CD11b^-CD11c^+F4/80^-$ DCs from two strains of IL-10-deficient mice at day 6 after infection, and compared the *Il23p19*, *Il12p40* and *Il12p35* mRNA transcripts. Macrophages tended to have higher amounts of each of these transcripts, but importantly, mice with a *LysM Cre*-mediated deletion of IL-10 had increased amounts of *Il23p19* and *Il12p40* mRNA, while *Il12p35* mRNA transcripts were similar to mice with a CD4 *Cre*-mediated deletion (Fig. 4a). To determine if IL-10 by macrophages regulates IL-23 production, we sorted GFP⁺ (IL-10⁺) and GFP⁻ (IL-10⁻) macrophages from $Il10^{sfp}$ mice before and after *C. rodentium* infection. We detected significantly increased amounts of *Il23p19* mRNA in GFP⁻ macrophages compared with those in GFP⁺ cells after infection, suggesting an autocrine role for IL-10 in the reporter positive cells. There were very few *Il23p19* mRNA transcripts in either type of macrophages before infection, but even in the GFP⁺ macrophages there was a small increase in *Il23p19* mRNA following the exposure to *C. rodentium* (Fig. 4b). We also sorted macrophages from the large intestine of mice with a germline *Il10* deficiency at day 3 after infection, cultured them with or without the addition of recombinant IL-10, and analysed the supernatants by ELISA. Exogenous IL-10 completely abrogated IL-23p19 production by colonic macrophages from infected mice; but had little effect on the IL-6 production (Fig. 4c). IL-12p70 was undetectable at this time, with or without the addition of IL-10 (data not shown). These data suggest that IL-10 might inhibit IL-23 production




Figure 3 | Increased pro-inflammatory cytokines in the large intestine in $Il10^{lox/lox}LysM-Cre$ mice. $Il10^{lox/lox}LysM-Cre$ (black) and $Il10^{lox/lox}Cd4-Cre$ (white) mice were analysed at day 0 or day 6 after *C. rodentium* infection. At the indicated time points, 3-mm fragments of the middle colon were cultured for 24 h. Cytokine production was measured in the supernatants by Multiplex or ELISA and normalized to the weight of the corresponding tissue fragments. Averages and s.d. from two independent experiments with six mice in each group are presented. Student's *t*-test, * $P < 0.05$, *** $P < 0.001$. IFN- γ , interferon γ .




Figure 4 | Macrophage-derived IL-10 influences mortality through regulation of IL-23p19. (a) Analysis of gene expression by reverse transcription PCR (RT-PCR) in sorted CD11b⁻CD11c⁺F4/80⁻ DC or CD11b⁺CD11c^{int}F4/80⁺ macrophages from large intestines in *Il10*^{flox/flox}*LysM-Cre* or *Il10*^{flox/flox}*Cd4-Cre* mice at day 6 after *C. rodentium* infection. Averages and s.d. from two independent experiments with six mice in each group are shown. (b) Analysis of *Il23p19* mRNA transcripts by RT-PCR in large intestine lamina propria macrophages, including GFP⁺ (IL-10⁺) and GFP⁻ (IL-10⁻), sorted from five to eight *Il10*^{gfp} mice before and after 6 days after *C. rodentium* infection. These data are from two independent experiments. (c) Large intestine lamina propria macrophages were sorted from five to eight infected *Il10*^{-/-}*Rag*^{-/-} mice and cultured for 24 h in the absence (black bars) or presence (white bars) of 100 ng ml⁻¹ recombinant mouse IL-10. Supernatants were analysed for IL23p19 and IL-6 by ELISA. These data are from 2–3 independent experiments. (d,e) *Il10*^{-/-}*Rag*^{-/-} and *Rag*^{-/-} mice or (f,g) *Il10*^{flox/flox}*LysM-Cre* and *Il10*^{flox/flox}*Cd4-Cre* mice were infected with *C. rodentium*. (d–g) At the time of infection, 100 μg anti-IL-23 or rat IgG were injected per mouse intravenously. (d,f) Survival curve and (e,g) histology score from middle colon at day 6 after infection are shown. These are from two independent experiments with 3–4 mice per group. Student's *t*-test, **P* < 0.05, ***P* < 0.01, ****P* < 0.001.

in vivo following infection, in part by acting directly on the IL-23-producing macrophages.

To elucidate if increased IL-23 production in the absence of IL-10 is involved in intestinal damage and mortality, independent of adaptive immunity, we analysed *C. rodentium*-infected *Rag*^{-/-} and *Il10*^{-/-}*Rag*^{-/-} recipients that were treated with

IL-23p19 blocking or control antibodies. *Il10*^{-/-}*Rag*^{-/-} recipients treated with control antibodies died between 5 and 10 days after infection; however, a neutralizing IL-23p19 antibody treatment rescued the survival of the infected mice (Fig. 4d). Furthermore, IL-23p19 antibody treatment prevented tissue damage at day 6 after infection compared with control antibodies

(Fig. 4e). These data show that in the absence of IL-10 the innate immune response is unchecked, and excessive IL-23 contributes to early morbidity and death.

To verify that blocking IL-23 also can reduce mortality in fully immune competent mice lacking only macrophage-derived IL-10, we analysed *C. rodentium*-infected $Il10^{lox/lox}LysM-Cre$ mice, and as a control, $Il10^{lox/lox}Cd4-Cre$ mice. $Il10^{lox/lox}LysM-Cre$ mice treated with the isotype control antibody began to die at day 3 after infection, and none survived beyond day 11. However IL-23p19 antibody treatment had a significant effect on promoting the survival and preventing the epithelial cell damage of the infected $Il10^{lox/lox}LysM-Cre$ mice (Fig. 4f,g). Altogether, these data indicate that macrophage-derived IL-10 plays a dominant role in controlling the amount of IL-23 production by macrophages in the large intestine during the early phases of *C. rodentium* infection, and IL-23 is responsible, at least in part, for morbidity and mortality and is a marker for the excessive response by innate immune cells.

Gene signature of IL-10 competent large intestine macrophages. Macrophages are heterogeneous, and therefore we determined which subset was responsible for preventing the excessive intestinal damage after *C. rodentium* infection. We reported previously that IL-10 competent GFP⁺ cells in the large intestine from $Il10^{gfp}$ reporter mice were mainly CD11b⁺, CD11c^{int} and F4/80⁺ (ref. 11). Therefore we first gated GFP⁺ or GFP⁻ cells, then sorted IL-10 competent, GFP⁺ CD11b⁺CD11c^{int}F4/80⁺ macrophages (IL-10⁺) or their GFP⁻ counterparts (IL-10⁻) from the large intestine of $Il10^{gfp}$ reporter mice, and performed microarray analyses to obtain insight into their degree of relatedness (Fig. 5a). F4/80, CD11b and MHC class II expression levels in post sorted GFP⁺ (IL-10⁺) and GFP⁻ (IL-10⁻) populations were similar. While CD11c expression in the GFP⁻ population had a broad distribution, we sorted those cells that had an amount of CD11c comparable to the GFP⁺ population. Heatmap and MA plots showed that these two populations were closely related, with 242 mRNA transcripts that were differentially expressed (Fig. 5b,c). Two hundred and eleven genes were upregulated in the GFP⁺ cells, including the cell surface markers, *Vcam1*, *Cd209f* and *Cd163*. In addition, 31 genes, including *Il12b*, were downregulated (Fig. 5c).

An annotation search in DAVID (database for Annotation, Visualization and Integrated Discovery functional classification tool) yielded unexpectedly enriched functional categories among upregulated genes in GFP⁺ macrophages ($P < 0.05$), including: extracellular structure organization, extracellular matrix organization, biological adhesion, regulation of peptidyl-tyrosine, vasculature development and behaviour. However, no pathways were identified for the downregulated genes, likely reflecting the small number of genes with decreased expression (Fig. 5d and Supplementary Fig. 6).

We validated increased expression of *Il10* and *Cd163* in GFP⁺ colonic macrophages by real-time reverse transcription PCR. In contrast, both the macrophage populations produced *Tgfb1* mRNA (Fig. 5e). We also confirmed increased expression of CD163 in GFP⁺ macrophages by flow cytometry (Fig. 5f).

It has been reported CD163 is an IL-10 target gene *in vitro*³³. Therefore we compared IL-10 receptor expression in IL-10⁺ and IL-10⁻ colonic macrophages. However there was no difference in receptor expression. To determine if IL-10 provides an autocrine signal in macrophages *in vivo*, we crossed $Il10^{gfp}$ mice with $Il10rb^{-/-}$ mice. Interestingly, we found a similar frequency of GFP⁺ colonic macrophages in mice that had not received any IL-10R signals *in vivo* (Fig. 5f). However, intestinal macrophages

from $Il10rb^{-/-}$ mice showed reduced CD163 cell surface expression (Fig. 5f). To determine if an autocrine IL-10 signal could be responsible for the increased expression of CD163 by intestinal macrophages, we analysed cells from $Il10^{lox/lox}LysM-Cre$ and control mice. Intestinal macrophages expressed increased CD163 in $Il10^{lox/lox}Cd4-Cre$ and $Il10^{lox/lox}$ mice compared with $Il10^{lox/lox}LysM-Cre$ (Fig. 5g). These data suggest that autocrine IL-10 may play a role in the induction of the phenotype of the IL-10-producing colonic macrophage subset, but IL-10 signals, either autocrine or paracrine, are not required for the induction of IL-10 competence in these cells.

IL-10 competent macrophages are protective. To determine if the reporter positive subset of large intestinal macrophages is required for preventing the excessive inflammation after *C. rodentium* infection, we transferred sorted, reporter expressing GFP⁺ (IL-10⁺) or GFP⁻ (IL-10⁻) colonic macrophages from $Il10^{gfp}$ mice into *C. rodentium*-infected $Il10^{-/-}Rag^{-/-}$ recipients (Fig. 6a). When GFP⁺ (IL-10⁺) macrophages were transferred, we detected a marked increase in the survival and protection of tissue damage in the intestine compared with recipients of GFP⁻ cells (Fig. 6b). Indeed we detected a significant reduction of inflammatory cell recruitment, including monocytes and granulocytes, in the large intestine of recipients of GFP⁺ colonic macrophages after *C. rodentium* infection (Fig. 6c). Furthermore, chemokines potentially involved in the recruitment of monocytes and neutrophils to the large intestine, including CCL3, CCL4 and CCL5, were significantly decreased in the recipients of GFP⁺ cells (Fig. 6c). These data show that IL-10 from macrophages in the large intestine is indispensable for limiting the excessive inflammation by regulating the inflammatory cell recruitment during the innate response.

Discussion

A balanced innate immune response is required for the protection from infection, and preventing excessive inflammation and tissue damage. Here we address what mediators control excessive inflammation and mortality during innate immune responses against mucosal bacteria, and the cell type and the mechanism responsible for the resolution of inflammatory response early during infection.

IL-10 is a potent inhibitor of the production of TNF, IL-1, IL-6 and IL-12 *in vitro*^{34,35}, and is therefore a good candidate for limiting the excessive inflammation in the intestinal mucosae. In the context of mucosal infectious disease, studies using *Salmonella choleraesuis*¹⁵, *Klebsiella pneumoniae*¹⁶, *Listeria monocytogenes*¹⁸ and *Candida albicans*¹⁹ have reported an increase in pathogen clearance, because the lack of IL-10 enhanced pro-inflammatory cytokines. In these situations the lack of IL-10 was beneficial to the host. In contrast, during infection with *Toxoplasma gondii*^{17,21,22} and *Trypanosoma cruzi*²³, infection in the absence of IL-10 caused excessive and often lethal inflammatory responses. These data highlight the importance of IL-10 in influencing the proper balance of innate immunity following infection with a variety of pathogens.

We observed that *C. rodentium* was cleared slightly faster in mice deficient for IL-10, but these animals suffered from high morbidity and mortality, indicative of increased inflammation. A recent study, however, reported only a beneficial effect of the lack of IL-10 following *C. rodentium* infection²⁰. The discordant findings could reflect the fourfold increased dose of bacteria we used, but more likely reflect aspects of the endogenous microbiota in different mouse colonies, as the virulence of *C. rodentium* is dependent on other components of the microbial flora³⁶. By analysing mouse strains with T cell- or myeloid cell-specific




Figure 5 | Comparison of IL-10-producing and non-producing large intestinal macrophages. (a) Gating strategy for the microarray analysis of GFP⁺ (IL10⁺) and GFP⁻ (IL10⁻) colonic lamina propria macrophages. Cells that were CD45⁺, MHC class II⁺, CD11b⁺, CD11c^{int} and F4/80⁺ were analysed. As depicted, the GFP⁻ cells (gate #1) used to provide mRNA were the subset falling in gates #2 and #3, CD11b⁺, CD11c^{int} and which were F4/80⁺ (gate #3). The corresponding GFP⁺ cells (gate #4) were less heterogeneous as they were almost entirely, CD11b⁺CD11c^{int} (gate #5) and F4/80⁺ (gate #6). After the sort, F4/80, CD11c, CD11b and MHC class II expressions of cells in gate 3 (blue) and gate 6 (red) were examined by flow cytometry. (b) Heatmap of differentially expressed probe sets ($n = 242$) between IL-10⁺ and IL-10⁻ macrophages from the large intestine lamina propria (LI: macrophages) in *Il10^{gfp}* reporter mice, ordered by average difference in intensity. LI, Large Intestine. (c) MA plot of all probe sets on the array. Red points significantly above zero represent higher expression in IL-10⁺ macrophages compared to IL-10⁻ macrophages ($n = 211$), while points significantly below zero represent the inverse relationship ($n = 31$). Grey points indicate probe sets that did not exhibit a significant difference in expression levels, according to criteria described in Methods. (d) Enriched Gene Ontology biological processes among upregulated genes, as determined by DAVID. (e) IL-10⁺ and IL-10⁻ macrophages (CD11b⁺CD11c^{int}F4/80⁺) were sorted from large intestine lamina propria of *Il10^{gfp}* reporter mice and analysed for expression of *Il10*, *Cd163* and *Tgfb1* by real-time PCR. Data are presented relative to L32 expression and are presented as average and s.d. of two independent experiments. (f) (Left) GFP (IL-10) expression by colonic macrophages in *Il10rb*^{-/-}, *Il10^{gfp}il10rb*^{-/-} and *Il10^{gfp}* mice. Gated on CD45⁺ MHC class II⁺ CD11b⁺ CD11c^{int} F4/80⁺ cells. (Right) CD163 expression by IL-10 positive macrophages in *Il10^{gfp}* mice (red) and in *Il10^{gfp}il10rb*^{-/-} mice (blue). (g) CD163 expression in large intestinal macrophages (CD11b⁺CD11c^{int}F4/80⁺ cells) from *Il10^{fllox/fllox}LysM-Cre* (red), *Il10^{fllox/fllox}Cd4-Cre* (blue) and *Il10^{fllox/fllox}* (grey) mice. Data are representative of one of the two independent experiments.

deletion of IL-10, we could implicate macrophage IL-10 in limiting the destructive immune response. Consistent with this, intestinal tissue from *Il10^{fllox/fllox}Cd4-Cre* mice produced more IL-10 protein than tissue from *Il10^{fllox/fllox}LysM-Cre* mice, both at steady state and at early time points after infection, suggesting that macrophages are major IL-10-producing cells in the colon, both before and after bacterial infection. Indeed, colonic macrophages were GFP⁺ before and after infection in *Il10^{gfp}* reporter mice.

Macrophage IL-10 is indispensable for maintaining a balance in adaptive immune responses in the mucosae^{11,37–40}, and here

we demonstrate its importance during the innate immune response. It is possible, however, that IL-10-producing intestinal macrophages acquire inflammatory features during bacterial infection, or alternatively, that many intestinal macrophages can rapidly acquire the ability to produce IL-10. We showed that adoptive transfer of IL-10-producing intestinal macrophages from reporter mice, but not macrophage populations that did not express IL-10, rescued the survival of infected, IL-10-deficient mice. Therefore the GFP⁻ colonic macrophages could not induce sufficient IL-10 production to attain a protective function during this acute infection. Although we have shown that IL-10 from




Figure 6 | IL-10 competent macrophages can rescue survival after infection. *Il10*^{-/-} *Rag*^{-/-} mice were infected with *C. rodentium*. On day 1 after infection, macrophages (CD11b⁺ CD11c^{int}F4/80⁺ cells) were sorted from large intestine of *Il10*^{gfp} reporter mice and 5 × 10⁵ cells per mouse of GFP⁺ (IL-10⁺) or GFP⁻ (IL-10⁻) populations were injected intravenously. **(a)** Gating strategy for obtaining GFP⁺ (IL-10⁺) and GFP⁻ (IL-10⁻) colonic macrophages for adoptive cell transfer. CD45⁺ colonic lamina propria cells were selected on the basis of CD11b⁺, CD11c^{int} expression (gate #1) and F4/80 expression (gate #2) and further separated into GFP⁺ (IL-10⁺) (gate #4) or GFP⁻ (IL-10⁻; gate #3) populations. **(b)** Survival and histology score at day 6 after infection were monitored and is presented as a summary of two independent experiments with 3–5 mice per group. **(c)** (Top) total cell numbers were counted from large intestine at day 6 after infection. Frequencies of monocytes (Ly6G⁻Ly6C^{high}) and granulocytes (Ly6G^{high}Ly6C^{low}) were measured by flow cytometry and cell numbers were calculated according to the respective total cell number. Cell numbers are presented as averages and s.d. from three mice per group from two independent experiments. (Bottom) analysis of *Ccl3*, *Ccl4* and *Ccl5* mRNA transcripts in the large intestine at day 6 after infection. Data are presented as averages with s.d. from three mice per group from two independent experiments. Student's *t*-test, ***P* < 0.01, ****P* < 0.001.

colonic macrophages is protective, we cannot formally exclude the possibility that the GFP⁻ cells were not protective because they could not home to the intestine or survive well after transfer.

We consider this unlikely, because we did not see an increase in GFP⁺ cells after infection. Therefore, while we suggest that the IL-10-producing intestinal macrophages could be a relatively

stable subset; further experiments will be required to prove that this is the case. Interestingly, while the transcriptomes of the two colonic macrophage subsets are similar, the GFP⁺ cells have increased expression of genes involved in extracellular matrix organization, vascular development, and others involved in tissue regeneration, such as the epidermal growth factor receptor and the platelet derived growth factor receptor. These data suggest that the GFP⁺ macrophages may have a more prominent role in tissue homeostasis and repair.

Here we have demonstrated that after a bacterial challenge the absence of IL-10 production by macrophages led to increased IL-23 synthesis, predominantly by macrophages, an enhanced innate immune response and fatal colitis. It is curious that IL-12 synthesis was not greatly affected, but indeed, it has been reported that IL-23 is associated with innate immune pathology in the intestine^{41,42}. Consistent with this, IL-23 transgenic mice developed systemic inflammation that included the terminal ileum and colon, with infiltration of neutrophils and macrophages⁴³. A few other studies have demonstrated that IL-10 negatively regulates IL-23 production, and therefore the absence of IL-10 caused an increase in IL-17-producing T cells and spontaneous intestinal inflammation^{5,44}. We note, however, that the *Il10^{flox/flox}LysM-Cre* did not have increased IL-23 in the intestine at steady state nor did they spontaneously develop colitis.

Our data demonstrate that exogenous IL-10 reduced *in vitro* IL-23 production by intestinal macrophages from *C. rodentium*-infected mice. *In vivo*, there is evidence that IL-10 acts directly on the IL-10-producing macrophage subset, although IL-10 competence did not depend on any IL-10 signals. IL-10-producing macrophages expressed increased CD163, however, which was dependent on IL-10R expression by these cells. Furthermore, GFP⁻ macrophages from infected mice produced more IL-23p19 mRNA more than GFP⁺ macrophages. However, in our analyses <40% of colonic macrophages were GFP⁺, and therefore the reduced inflammation and improved survival in wt mice suggests a paracrine role for IL-10 is also possible.

Recent studies showed that IL-10 signals drive macrophages to express tolerogenic functions that prevent colitis^{14,45}. However, those investigations focused on either steady-state or adaptive mucosal immune responses, not on innate mucosal immune responses, and they did not provide a mechanism or tie the protective role of IL-10 to macrophage IL-23 production.

In humans, there is a study showing myeloid DCs from Crohn's disease patients produced a higher amount of IL-23 and a lower amount of IL-10 after a bacterial stimulation when compared with ulcerative colitis patients⁴⁶, leading us to propose that intestinal IL-10-producing myeloid cells might regulate IL-23 production not only in Crohn's disease, but also in bacterial infection in humans.

In summary, our data show that IL-10 synthesis by intestinal macrophages has a marked effect on regulating the innate immune response in mice through regulating macrophage IL-23 production after mucosal bacterial infection. Furthermore, intestinal macrophages are heterogeneous, and although IL-10 competent and IL-10⁻ macrophages are similar, we suggest that they may not rapidly interconvert *in vivo*.

Material and methods

Animals. Mice were bred and housed under specific pathogen free conditions in the vivarium of the La Jolla Institute for Allergy & Immunology (LJI, La Jolla, CA). Eight weeks old female *Il10^{-/-}*, *Il10rb^{-/-}* and *Rag^{-/-}* mice were purchased from Jackson laboratory. Eight weeks old female *Il10^{flox/flox}* (*VertX*) mice were obtained from Christopher Karp (Cincinnati Children's Hospital, OH) and also purchased from Jackson Laboratory. *Il10^{-/-}Rag^{-/-}* mice were described previously⁴⁷. *Il10^{flox/flox}* mice were obtained from Alexander Rudensky (Memorial Sloan-Kettering Cancer Center, NY) with the permission of Axel Roers (Dresden University of Technology,

Germany). *Il10^{flox/flox}* mice were bred to *Cd4-Cre*, *LysM-Cre* and *Cd11c-Cre* mice, which were all purchased from Jackson laboratory. All mouse strains were generated on the C57BL6/J background or backcrossed at least 10 generations. All procedures were approved by the LJI Animal Care and Use Committee.

C. rodentium infection. A chloramphenicol-resistant variant of the wt *C. rodentium* strain DBS100 was cultured as previously described³⁰. In brief, bacteria were cultured for 15–16 h in Luria-Bertani broth at 37 °C. Mice were infected with 2–3 × 10⁹ CFU *C. rodentium* by oral gavage. In some experiments functional grade IL23p19 (G23-8) antibody or control rat IgG were injected intravenously at 100 µg per mouse at the time of infection. Antibodies were purchased from eBiosciences (San Diego, CA). GFP⁺ (IL-10⁺) and GFP⁻ (IL-10⁻) macrophages were sorted from colon tissue of uninfected *Il10^{flox/flox}* (*VertX*) mice as CD45⁺, MHC class II⁺, CD11b⁺, F4/80⁺, CD11c^{int} cells. One day after infection, 5 × 10⁵ macrophages were injected intravenously per mouse. For CFU determination, the faecal pellets and the spleen were weighed, homogenized and plated on chloramphenicol-containing MacConkey agar plates in serial dilutions.

FITC-dextran permeability assay. *In vivo* permeability assays to assess intestinal barrier function were performed using fluorescein isothiocyanate (FITC)-labelled dextran (Sigma) as described previously³⁰. In brief, mice were infected with *C. rodentium*. At day 0 or 6 after infection, mice were gavaged with FITC-dextran (60 mg per 100 g body weight) and serum was collected retro-orbitally 4 h later. Blood cells in serum samples were removed and the fluorescence intensity of FITC-dextran was measured. FITC-dextran concentrations were determined from the standard curves generated by serial dilution.

Reagents and antibodies. For flow cytometry and cell sorting the following antibodies were used: CD11b (M1/70), CD11c (N418), F4/80 (BM8), CD45 (30-F11), IA/IE (M5/114.15.2), Ly6C (HK1.4), Ly6G (1A8), CD4 (RM4-5), TCRβ (H57-597), CD163 (polyclonal). Antibodies were purchased from BD Biosciences (San Diego, CA), eBiosciences (San Diego, CA), BioLegend (San Diego, CA) or Antibodies-online.com. Cell sorting was performed on an Aria II instrument (BD Biosciences). Cytokines and chemokines were quantified using either the Bio-plex multicytokine assay (Bio-Rad, Hercules, CA), as described previously⁴⁸, IL-23p19, IL-22 or IL-12p70 ELISA kits (eBiosciences, San Diego, CA) or a IL-6 ELISA kit (BD, San Diego, CA).

Colon fragment cultures. Large intestines were harvested and 3-mm colon fragments were cultured for 24 h in DMEM with 10% FBS and antibiotics. Supernatants were analysed for cytokines and chemokines in ratio to the weight of the corresponding tissue fragment.

Isolation of lamina propria cells. Large intestines were harvested, opened longitudinally and washed to remove faecal content. Intestines were cut into small pieces and incubated in a horizontal shaker at 37 °C in the presence of 2.5 mM EDTA three times for 20 min to remove epithelial cells. Colon pieces were minced and digested for 20 min with 1 mg ml⁻¹ Collagenase type VIII (Sigma, St Louis, MO) at 37 °C. Lamina propria cells were filtered and stained for flow cytometric analysis or cell sorting.

Histology. Tissue sections were taken from the middle colon and fixed in 10% formalin. After embedding in paraffin, 5-µm sections were cut and stained with hematoxylin-eosin. Histological scores were assigned in a blinded manner according to the following criteria: (i) mucosal architecture (0, normal; 1, focally abnormal; 2, diffusely abnormal; 3, severely abnormal); (ii) inflammatory cell infiltration of mucosa (0, normal; 1, mild infiltration; 2, moderate infiltration; 3, severe infiltration), submucosa (0, normal; 1, mild infiltration; 2 moderate infiltration; 3, severe infiltration), muscle (0, normal; 1 moderate to severe) and serosa (0, absent; 1, present); (iii) epithelial erosions and ulcerations (0, absent; 1, present); (iv) crypt abscesses (0, absent; 1, present); and (v) goblet cell loss (0, absent; 1, present). The maximal total score for colitis severity is 14.

Relative mRNA quantification. Total RNA was extracted from sorted cells or 2–3-mm long colon tissue fragments using the RNeasy Mini kit (Qiagen, Hilden, Germany) according to the manufacturer's instructions. Genomic DNA was digested with DNase I (Qiagen), cDNA was synthesized using iScript (Bio-Rad, Hercules, CA) and real-time PCR was performed using SYBRgreen (Roche, Indianapolis, IN) on a LightCycler instrument (Roche). The primers were as follows: L32, 5'-GAA ACT GGC GGA AAC CCA-3' (sense), 5'-GGA TCT GGC CCT TGA ACC TT-3' (antisense), CD163, 5'-GGT GGA CAC AGA ATG GTT CTT C-3' (sense), 5'-CCA GGA GCG TTA GTG ACA GC-3' (antisense), TGFβ1, 5'-ACC ATG CCA ACT TCT GTC-3' (sense), 5'-CGG GTT GTG TTG GTT GTA GA-3' (antisense), *Il10*, 5'-GCC ACA TGC TCC TAG AGC TG-3' (sense), 5'-CAG CTG GTC CTT TGT TTG AAA-3' (antisense), *Il23p19*, 5'-AAT AAT GTG CCC CGT ATC CA-3' (sense), 5'-GGA TCC TTT GCA AGC AGA AC-3' (antisense), *Il12p35*, 5'-GTG AAG ACG GCC AGA GAA A-3' (sense), 5'-GGT CCC GTG

TGA TGT CTT C-3' (antisense), *Il12p40*, 5'-AGC AGT AGC AGT TCC CCT GA-3' (sense), 5'-AGT CCC TTT GGT CCA GTG TG-3' (antisense).

Microarray data processing and analysis. All microarray data sets were processed in R, using customized scripts and Bioconductor modules⁴⁹. Raw intensity data were normalized with the 'rma' function, with default parameters. Differential expression analysis was performed using the 'limma' package. Probe sets were deemed significantly differentially expressed only if their corrected *P* values were <0.05 and the difference was at least threefold. Heatmaps containing the differentially expressed genes in each set were built using the 'regHeatmap' function of the 'Heatplus' module and sorted by average fold change. The 242 probe sets that were significantly differentially expressed were mined for enriched Gene Ontology biological process terms using DAVID (<http://david.abcc.ncifcrf.gov>).

Immunofluorescence. Colons and ilea were harvested, flushed with PBS (Life Technologies, Carlsbad, CA) and fixed with 4% paraformaldehyde (Electron Microscopy Sciences, Hatfield, PA) in PBS overnight at 4 °C and then equilibrated in 30% w/v sucrose solution in PBS for another 24 h. Tissues were briefly washed in PBS, frozen in OCT and stored at -80 °C. Cryostat sections were cut at 16-µm thickness and air-dried for 30 min. Sections were rehydrated for 10 min in PBS and nonspecific binding sites were saturated with 10% normal goat serum, 0.5% bovine serum albumin (both from Gemini Bio-Products, West Sacramento, CA) in the presence of 0.1% Triton X-100 in PBS for 1 h. Tissues were incubated with anti-F4/80 rat monoclonal antibodies (50 µg ml⁻¹, BM8, eBiosciences) and anti-GFP rabbit antiserum (5 µg ml⁻¹, ab290, Abcam, Boston, MA) overnight at 4 °C. Following washing, sections were reacted with donkey anti-rat DyLight 549-labelled (1.4 µg ml⁻¹, Jackson ImmunoResearch) and goat anti-rabbit F(ab')₂ fragment Alexa Fluor 647-labelled secondary antibodies (4 µg ml⁻¹, Invitrogen) for 1 h at room temperature. Slides were washed, mounted in Prolong Gold antifade reagent (Invitrogen) with a cover slip and examined with FluoView FV10i confocal microscope (Olympus). Imaging was performed at room temperature, using 60 × /1.35 NA oil objective. To improve feature visibility images were processed by contrast stretching using identical procedures.

Statistical analysis. The Student's *t*-test was used for statistical analysis except for the survival and histological scores, for which the log-rank test and the Mann-Whitney test were used, respectively. Differences were considered significant at *P* < 0.05.

References

- Glocker, E. O. *et al.* Inflammatory bowel disease and mutations affecting the interleukin-10 receptor. *N. Engl. J. Med.* **361**, 2033–2045 (2009).
- Moran, C. J. *et al.* IL-10R polymorphisms are associated with very-early-onset ulcerative colitis. *Inflamm. Bowel. Dis.* **19**, 115–123 (2013).
- Kuhn, R., Lohler, J., Rennick, D., Rajewsky, K. & Muller, W. Interleukin-10-deficient mice develop chronic enterocolitis. *Cell* **75**, 263–274 (1993).
- Spencer, S. D. *et al.* The orphan receptor CRF2-4 is an essential subunit of the interleukin 10 receptor. *J. Exp. Med.* **187**, 571–578 (1998).
- Yen, D. *et al.* IL-23 is essential for T cell-mediated colitis and promotes inflammation via IL-17 and IL-6. *J. Clin. Invest.* **116**, 1310–1316 (2006).
- Groux, H. *et al.* A CD4 T-cell subset inhibits antigen-specific T-cell responses and prevents colitis. *Nature* **389**, 737–742 (1997).
- Van Montfrans, C. *et al.* Prevention of colitis by interleukin 10-transduced T lymphocytes in the SCID mice transfer model. *Gastroenterology* **123**, 1865–1876 (2002).
- Asseman, C., Read, S. & Powrie, F. Colitogenic Th1 cells are present in the antigen-experienced T cell pool in normal mice: control by CD4+ regulatory T cells and IL-10. *J. Immunol.* **171**, 971–978 (2003).
- Roers, A. *et al.* T cell-specific inactivation of the interleukin 10 gene in mice results in enhanced T cell responses but normal innate responses to lipopolysaccharide or skin irritation. *J. Exp. Med.* **200**, 1289–1297 (2004).
- Rubtsov, Y. P. *et al.* Regulatory T cell-derived interleukin-10 limits inflammation at environmental interfaces. *Immunity* **28**, 546–558 (2008).
- Murai, M. *et al.* Interleukin 10 acts on regulatory T cells to maintain expression of the transcription factor Foxp3 and suppressive function in mice with colitis. *Nat. Immunol.* **10**, 1178–1184 (2009).
- Siewe, L. *et al.* Interleukin-10 derived from macrophages and/or neutrophils regulates the inflammatory response to LPS but not the response to CpG DNA. *Eur. J. Immunol.* **36**, 3248–3255 (2006).
- Pils, M. C. *et al.* Monocytes/macrophages and/or neutrophils are the target of IL-10 in the LPS endotoxemia model. *Eur. J. Immunol.* **40**, 443–448 (2010).
- Zigmond, E. *et al.* Macrophage-restricted interleukin-10 receptor deficiency, but not IL-10 deficiency, causes severe spontaneous colitis. *Immunity* **40**, 720–733 (2014).
- Arai, T. *et al.* Effects of *in vivo* administration of anti-IL-10 monoclonal antibody on the host defence mechanism against murine Salmonella infection. *Immunology* **85**, 381–388 (1995).
- Greenberger, M. J. *et al.* Neutralization of IL-10 increases survival in a murine model of *Klebsiella pneumoniae*. *J. Immunol.* **155**, 722–729 (1995).
- Gazzinelli, R. T. *et al.* In the absence of endogenous IL-10, mice acutely infected with *Toxoplasma gondii* succumb to a lethal immune response dependent on CD4+ T cells and accompanied by overproduction of IL-12, IFN-γ and TNF-α. *J. Immunol.* **157**, 798–805 (1996).
- Dai, W. J., Kohler, G. & Brombacher, F. Both innate and acquired immunity to *Listeria monocytogenes* infection are increased in IL-10-deficient mice. *J. Immunol.* **158**, 2259–2267 (1997).
- Vazquez-Torres, A., Jones-Carson, J., Wagner, R. D., Warner, T. & Balish, E. Early resistance of interleukin-10 knockout mice to acute systemic candidiasis. *Infect. Immun.* **67**, 670–674 (1999).
- Dann, S. M. *et al.* Attenuation of intestinal inflammation in IL-10 deficient mice infected with *Citrobacter rodentium*. *Infect. Immun.* **82**, 1949–1958 (2014).
- Reis e Sousa, C. *et al.* Paralysis of dendritic cell IL-12 production by microbial products prevents infection-induced immunopathology. *Immunity* **11**, 637–647 (1999).
- Suzuki, Y. *et al.* IL-10 is required for prevention of necrosis in the small intestine and mortality in both genetically resistant BALB/c and susceptible C57BL/6 mice following peroral infection with *Toxoplasma gondii*. *J. Immunol.* **164**, 5375–5382 (2000).
- Hunter, C. A. *et al.* IL-10 is required to prevent immune hyperactivity during infection with *Trypanosoma cruzi*. *J. Immunol.* **158**, 3311–3316 (1997).
- Mundy, R., MacDonald, T. T., Dougan, G., Frankel, G. & Wiles, S. *Citrobacter rodentium* of mice and man. *Cell. Microbiol.* **7**, 1697–1706 (2005).
- Eckmann, L. Animal models of inflammatory bowel disease: lessons from enteric infections. *Ann. NY Acad. Sci.* **1072**, 28–38 (2006).
- Bhinder, G. *et al.* The *Citrobacter rodentium* mouse model: studying pathogen and host contributions to infectious colitis. *J. Vis. Exp.* e50222 (2013).
- Gately, M. K. *et al.* The interleukin-12/interleukin-12-receptor system: role in normal and pathologic immune responses. *Annu. Rev. Immunol.* **16**, 495–521 (1998).
- Cooper, A. M. & Khader, S. A. IL-12p40: an inherently agonistic cytokine. *Trends. Immunol.* **28**, 33–38 (2007).
- Satpathy, A. T. *et al.* Notch2-dependent classical dendritic cells orchestrate intestinal immunity to attaching-and-effacing bacterial pathogens. *Nat. Immunol.* **14**, 937–948 (2013).
- Shui, J. W. *et al.* HVEM signalling at mucosal barriers provides host defence against pathogenic bacteria. *Nature* **488**, 222–225 (2012).
- Hunter, C. A. New IL-12-family members: IL-23 and IL-27, cytokines with divergent functions. *Nat. Rev. Immunol.* **5**, 521–531 (2005).
- Manta, C. *et al.* CX3CR1(+) macrophages support IL-22 production by innate lymphoid cells during infection with *Citrobacter rodentium*. *Mucosal Immunol.* **6**, 177–188 (2012).
- Sulhian, T. H. *et al.* Human monocytes express CD163, which is upregulated by IL-10 and identical to p155. *Cytokine* **12**, 1312–1321 (2000).
- de Waal Malefyt, R., Abrams, J., Bennett, B., Figdor, C. G. & de Vries, J. E. Interleukin 10(IL-10) inhibits cytokine synthesis by human monocytes: an autoregulatory role of IL-10 produced by monocytes. *J. Exp. Med.* **174**, 1209–1220 (1991).
- Saraiva, M. & O'Garra, A. The regulation of IL-10 production by immune cells. *Nat. Rev. Immunol.* **10**, 170–181 (2010).
- Kamada, N. *et al.* Regulated virulence controls the ability of a pathogen to compete with the gut microbiota. *Science New York, NY* **336**, 1325–1329 (2012).
- Denning, T. L., Wang, Y. C., Patel, S. R., Williams, I. R. & Pulendran, B. Lamina propria macrophages and dendritic cells differentially induce regulatory and interleukin 17-producing T cell responses. *Nat. Immunol.* **8**, 1086–1094 (2007).
- Ueda, Y. *et al.* Commensal microbiota induce LPS hyporesponsiveness in colonic macrophages via the production of IL-10. *Int. Immunol.* **22**, 953–962 (2010).
- Medina-Conterras, O. *et al.* CX3CR1 regulates intestinal macrophage homeostasis, bacterial translocation, and colitogenic Th17 responses in mice. *J. Clin. Invest.* **121**, 4787–4795 (2011).
- Rivollier, A., He, J., Kole, A., Valatas, V. & Kelsall, B. L. Inflammation switches the differentiation program of Ly6Chi monocytes from anti-inflammatory macrophages to inflammatory dendritic cells in the colon. *J. Exp. Med.* **209**, 139–155 (2012).
- Uhlir, H. H. *et al.* Differential activity of IL-12 and IL-23 in mucosal and systemic innate immune pathology. *Immunity* **25**, 309–318 (2006).
- Hue, S. *et al.* Interleukin-23 drives innate and T cell-mediated intestinal inflammation. *J. Exp. Med.* **203**, 2473–2483 (2006).
- Wiekowski, M. T. *et al.* Ubiquitous transgenic expression of the IL-23 subunit p19 induces multiorgan inflammation, runting, infertility, and premature death. *J. Immunol.* **166**, 7563–7570 (2001).
- Hoshi, N. *et al.* MyD88 signalling in colonic mononuclear phagocytes drives colitis in IL-10-deficient mice. *Nat. Commun.* **3**, 1120 (2012).

45. Shouval, D. S. *et al.* Interleukin-10 receptor signaling in innate immune cells regulates mucosal immune tolerance and anti-inflammatory macrophage function. *Immunity* **40**, 706–719 (2014).
46. Sakuraba, A. *et al.* Th1/Th17 immune response is induced by mesenteric lymph node dendritic cells in Crohn's disease. *Gastroenterology* **137**, 1736–1745 (2009).
47. Murai, M. *et al.* Peyer's patch is the essential site in initiating murine acute and lethal graft-versus-host reaction. *Nat. Immunol.* **4**, 154–160 (2003).
48. Park, Y. *et al.* TSC1 regulates the balance between effector and regulatory T cells. *J. Clin. Invest.* **123**, 5165–5178 (2013).
49. Gentleman, R. C. *et al.* Bioconductor: open software development for computational biology and bioinformatics. *Genome. Biol.* **5**, R80 (2004).

Acknowledgements

We thank Drs K. Ley and R. Hanna for critical reading of this manuscript, Dr C. Hedrick for discussions, C. Kim and K. Van Gunst for assistance with cell sorting. This work was supported by the National Institutes of Health Grants R21 AI091911 to M.M., PO1 AI089624 and Crohn's and the Colitis Foundation of America, Senior research award to M.K.

Author contributions

M.M. and M.K. designed experiments. P.K., V.M., Y.P., U.B., Z.M., T.M. and M.M. performed the experiments. M.M., P.K., J.A.G. and B.P. performed the bioinformatics analysis. J.-W.S., G.K., H.C., Y.-C.L. and B.P. helped with critical advice and discussion throughout. P.K., M.K. and M.M. wrote the manuscript.

Additional information

Accession codes. Microarray data has been deposited in the Gene Expression Omnibus (GEO) database under accession code GSE58677.

Supplementary Information accompanies this paper at <http://www.nature.com/naturecommunications>

Competing financial interests: The authors declare no competing financial interest.

Reprints and permission information is available at <http://npg.nature.com/reprintsandpermissions/>

How to cite this article: Krause, P. *et al.* IL-10-producing intestinal macrophages prevent excessive antibacterial innate immunity by limiting IL-23 synthesis. *Nat. Commun.* **6**:7055 doi: 10.1038/ncomms8055 (2015).