

Published in final edited form as:

Nature. 2008 July 17; 454(7202): 350–352. doi:10.1038/nature07021.

IL-21 and TGF- β are required for differentiation of human T_H17 cells

Li Yang^{1,*}, David E. Anderson^{1,*}, Clare Baecher-Allan¹, William D. Hastings¹, Estelle Bettelli¹, Mohamed Oukka¹, Vijay K. Kuchroo^{1,*}, and David A. Hafler¹

¹Division of Molecular Immunology, Center for Neurologic Diseases, Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts 02115, USA.

Abstract

The recent discovery of CD4⁺ T cells characterized by secretion of interleukin (IL)-17 (T_H17 cells) and the naturally occurring regulatory FOXP3⁺ CD4 T cell (nT_{reg}) has had a major impact on our understanding of immune processes not readily explained by the T_H1/T_H2 paradigm. T_H17 and nT_{reg} cells have been implicated in the pathogenesis of human autoimmune diseases, including multiple sclerosis, rheumatoid arthritis, inflammatory bowel disease and psoriasis^{1,2}. Our recent data and the work of others demonstrated that transforming growth factor- β (TGF- β) and IL-6 are responsible for the differentiation of naive mouse T cells into T_H17 cells, and it has been proposed that IL-23 may have a critical role in stabilization of the T_H17 phenotype³⁻⁵. A second pathway has been discovered in which a combination of TGF- β and IL-21 is capable of inducing differentiation of mouse T_H17 cells in the absence of IL-6 (refs⁶⁻⁸). However, TGF- β and IL-6 are not capable of differentiating human T_H17 cells^{2,9} and it has been suggested that TGF- β may in fact suppress the generation of human T_H17 cells¹⁰. Instead, it has been recently shown that the cytokines IL-1 β , IL-6 and IL-23 are capable of driving IL-17 secretion in short-term CD4⁺ T cell lines isolated from human peripheral blood¹¹, although the factors required for differentiation of naive human CD4 to T_H17 cells are still unknown. Here we confirm that whereas IL-1 β and IL-6 induce IL-17A secretion from human central memory CD4⁺ T cells, TGF- β and IL-21 uniquely promote the differentiation of human naive CD4⁺ T cells into T_H17 cells accompanied by expression of the transcription factor RORC2. These data will allow the investigation of this new population of T_H17 cells in human inflammatory disease.

To better understand regulation of IL-17A secretion from human CD4⁺ T cells, we used a strategy that would allow us to evaluate the effects of various combinations of cytokine on expansion of T_H17 cells from memory T cells versus differentiation of naive CD4⁺ lymphocytes into T_H17 cells. Specifically, we used high-speed flow cytometry for sorting these two distinct populations of CD4⁺ T cells from the peripheral blood of healthy subjects: CD4⁺CD25⁻CD62L⁺CD45RA^{hi} cells highly enriched for naive T cells and CD4⁺CD25⁻CD62L⁺CD45RA⁻ cells enriched for central memory T cells (T_{CM}; Fig. 1a). All cells enriched for a naive or a central memory phenotype expressed the chemokine receptor CCR7 (data not shown). These two T cell populations were then stimulated with plate-bound

©2008 Macmillan Publishers Limited. All rights reserved

Correspondence and requests for materials should be addressed to D.E.A. (danderson@rics.bwh.harvard.edu)..

*These authors contributed equally to this work.

Reprints and permissions information is available at www.nature.com/reprints.

Full Methods and any associated references are available in the online version of the paper at www.nature.com/nature.

Supplementary Information is linked to the online version of the paper at www.nature.com/nature.

anti-CD3 and soluble anti-CD28 monoclonal antibodies for 7 days in serum-free medium containing different combinations of cytokines implicated in CD4⁺ T_H17 cell differentiation.

As reported previously, the cytokine IL-1 β induced the greatest amount of IL-17A secretion from T_{CM} (Fig. 1b). The addition of IL-6 alone had little effect on induction of IL-17A, and when added with IL-1 β had no additive or synergistic effect on IL-17A production. Addition of IL-23 was also able to modestly enhance IL-17A secretion from T_{CM}. However, IL-1 β alone or together with IL-6 failed to induce IL-17A secretion from naive CD4⁺ T cells. In marked contrast, a combination of TGF- β and IL-21 was uniquely able to induce T_H17 differentiation. Whereas IL-21, IL-1 β or IL-6 induced significant amounts of interferon- γ (IFN- γ) secretion from naive T cells, the addition of TGF- β with IL-21 suppressed IFN- γ secretion and induced differentiation of T_H17 cells. Whereas we did observe variability in the extent of T_H17 differentiation among unrelated healthy donors (Fig. 1c), we always observed induction of IL-17A after differentiation in the presence of TGF- β with IL-21. Intracytoplasmic staining demonstrated, in agreement with enzyme-linked immunosorbent assay (ELISA) results, that the combination of TGF- β and IL-21 differentiated CD4⁺ T cells that secreted only IL-17A and no IFN- γ (Fig. 1d). A recent study¹² has demonstrated that the pathogenicity of mouse IL-17-secreting T cells is influenced by whether they also secrete IL-10. Using intracytoplasmic staining, we failed to observe any IL-10 when naive CD4⁺ T cells were differentiated in the presence of TGF- β and IL-21 (data not shown).

RORC2 is the human homologue of mouse ROR γ t—a transcription factor critical for the differentiation of mouse IL-17-secreting T cells. Thus, to understand the molecular mechanisms involved in differentiation of human T_H17 cells, we used quantitative PCR with reverse transcription (RT-PCR) to evaluate messenger RNA levels of *RORC2* and other molecules implicated in T_H17 differentiation. The combination of TGF- β and IL-21 induced high levels of *RORC2* (Fig. 2a), consistent with their ability to induce IL-17A secretion from naive human CD4⁺ T cells. It was of particular interest that the combination of TGF- β and IL-6 that induces T_H17 differentiation in mouse T cells also induced expression of *RORC2* in naive human CD4 cells. Because this combination of cytokines did not, however, induce IL-17A secretion, these data indicate that expression of *RORC2* in humans is not in itself sufficient to induce IL-17 production, and that another as yet unidentified transcription factor in combination with *RORC2*, perhaps the human homologue of mouse ROR α (ref. ¹³), may be required to induce IL-17A-secreting T_H17 cells.

T-bet (also known as TBX21) is the master regulator for T_H1 cells secreting IFN- γ , whereas GATA3 induces T_H2, IL-4 secreting CD4 cells. Messenger RNA expression levels of *Tbet* were highly concordant with amounts of IFN- γ secretion and were consistent with our findings that whereas TGF- β and IL-21 induce T_H17 cell differentiation with *RORC2* expression, TGF- β suppresses the induction of T-bet by IL-21. Similarly, there was no induction of *GATA3* with TGF- β and IL-21. The cytokines IL-6, IL-21 and IL-1 β but not TGF- β induced IL-23 receptor upregulation in stimulated naive CD4⁺ T cells. We also examined the expression of the T_{reg} transcription factor FOXP3. As has been previously reported in both mouse and human systems, *FOXP3* was induced by TGF- β 1. Induction of *FOXP3* was inhibited by IL-6 and to a greater extent by IL-21—transcription factors that induce *RORC2*. Thus, although the induction of *RORC2* and FOXP3 transcription factors was highly similar between mouse and human naive CD4 cells, the induction of IL-17A by IL-6 in combination with TGF- β is discordant between the species.

We and others have shown previously that IL-21 secreted by mouse CD4⁺ T cells can induce the secretion of IL-21 in an autocrine amplification loop^{6-8,14}. Thus, we examined whether human IL-21 similarly induced IL-21 secretion from naive CD4⁺ T cells; we also evaluated the effects of a combination of TGF- β with IL-21 and IL-1 β , given the ability of these cytokines

to induce IL-17 from naive and central memory CD4⁺ T cells. Consistent with results in mice, IL-21 upregulated IL-21, although IL-1 β induced even greater amounts of *IL21* mRNA (Fig. 2b). In contrast to what has been observed in mice, IL-21 also increased *IL22* mRNA levels in naive CD4⁺ T cells in the absence of any exogenous IL-23. However, TGF- β suppressed the expression of *IL21* and *IL22* mRNA induced by IL-21 (Fig. 2b). These data further highlight similar yet subtle differences between human and mouse CD4⁺ T cells, because whereas IL-21 induces *IL21* and *IL22*, differentiation to T_H17 cells with TGF- β inhibits the expression of these cytokines.

To confirm the unique function of TGF- β and IL-21 in the differentiation of T_H17 cells from naive human CD4⁺ T cells, we sorted CD4⁺CD25⁻CD62L⁺CD45RA^{hi} cells from human cord blood. As expected, a higher proportion of CD4 cells in the cord blood exhibited this naive phenotype relative to peripheral blood obtained from healthy adult subjects (data not shown). TGF- β and IL-21 induced the upregulation of *IL17A* and *RORC2* mRNA (Fig. 2c). Although IL-21 alone modestly induced *RORC2*, only TGF- β and IL-21 were able to induce *IL17A* mRNA. When given a very strong *in vitro* stimulus, naive CD4⁺ T cells sorted from cord blood secreted IL-17A protein (Fig. 2d). These data further indicate that TGF- β and IL-21 are critical in the differentiation of both human and mouse T_H17 cells.

Collectively, our data refine and extend our understanding of the regulation of IL-17A secretion from human CD4⁺ T cells and define the conditions required for human T_H17 cell differentiation. We confirm recent reports that IL-1 β together with IL-6 (ref. ⁹) or IL-23 (ref. ²) can induce IL-17A secretion, which is most apparent in the human memory CD4⁺ T cell subset. A combination of TGF- β plus IL-21 is required for the differentiation of T_H17 cells from naive T cells. It is important to note that to observe T_H17 differentiation from naive human CD4⁺ T cells, appropriate amounts of both IL-21 and TGF- β are needed: addition of IL-21 alone must be sufficient to upregulate IFN- γ secretion, and the amount of TGF- β added must inhibit IL-21-induced IFN- γ secretion (see Methods for details). T_H17 cells differentiated under these conditions are also notable for secretion of IL-17A in the absence of IFN- γ . In summary, our data together with previous reports in humans suggest that IL-1 β and IL-6 induced during the early stages of an inflammatory response may act on memory T cells to promote IL-17 and IL-21 secretion, with induced IL-21 able to synergize with TGF- β to promote differentiation of T_H17 cells from naive CD4⁺ T cells. IL-23 may serve to expand or stabilize the phenotype of previously differentiated T_H17 cells. These experiments allow for the characterization of human inflammatory T_H17 responses associated with infection and autoimmune diseases.

METHODS SUMMARY

Cell sorting

PBMCs were obtained from the peripheral blood of healthy subjects or from cord blood (AllCells) in compliance with institutional review board (IRB) protocols. CD4⁺ T cells were subsequently isolated by negative selection using magnetic beads (Miltenyi Biotec). Naive (CD25⁻CD62L⁺CD45RA^{hi}) and central memory (CD25⁻CD62L⁺CD45RA⁻) CD4⁺ T cells were obtained by staining with the following antibodies: CD62L-FITC, CD4-PerCP, CD45RA-PE-Cy7, CD25-APC-Cy7 (BD Pharmingen) and were sorted on a FACS Aria (BD Biosciences).

Differentiation assays

Naive or central memory CD4⁺ T cells were stimulated with plate-bound anti-CD3 and soluble CD28 monoclonal antibodies in serum-free X-VIVO15 medium (Biowhittaker) and cytokines (IL-6, 25 ng ml⁻¹; TGF- β 1, 5 ng ml⁻¹; IL-1 β , 12.5 ng ml⁻¹; IL-21, 25 ng ml⁻¹; IL-23, 25 ng ml⁻¹) for a period of 7 days, at which point supernatants were collected and tested by ELISA

for IFN- γ (BD Biosciences) or IL-17A (eBioscience) using paired antibodies. We have observed that concentrations of TGF- β ranging from 0.1 ng ml⁻¹ to 10 ng ml⁻¹ in the presence of IL-21 promote T_H17 differentiation, whereas 50 ng ml⁻¹ TGF- β suppresses differentiation. Intracytoplasmic staining was performed using standard methodologies and anti-IL-17-APC (R&D Systems) and anti-IFN- γ -PE or anti-IL-10-PE (BD Biosciences) antibodies.

Real-time PCR

All primers and probes were obtained from Applied Biosystems and used according to standard methodologies.

METHODS

Differentiation assays

Plates (96-well U-bottom) were coated with 1.5 μ g ml⁻¹ anti-CD3 monoclonal antibody (eBioscience, clone OKT3) in a volume of 50 μ l of PBS and were incubated overnight (16 h) at 4 °C. For T cells isolated from cord blood, a 96-well plate was pre-coated overnight at 4 °C with 3 μ g ml⁻¹ anti-CD3 monoclonal antibody (UCHT1, BD Biosciences). Antibody solution was then removed, plates were rinsed once with serum-free X-VIVO 15 medium (Biowhittaker), and naive or central memory CD4⁺ T cells (5×10^4 per well) were stimulated in serum-free X-VIVO15 medium with soluble CD28 (BD Biosciences, clone 28.2) monoclonal antibody (1.0 μ g ml⁻¹) and cytokines (IL-6, 25 ng ml⁻¹; TGF- β 1, 5 ng ml⁻¹; IL-1 β , 12.5 ng ml⁻¹; IL-21, 25 ng ml⁻¹; IL-23, 25 ng ml⁻¹) for a period of 7 days in the absence of IL-2. The cytokines IL-6, IL-1 β , IL-23 and TGF β 1 (catalogue number 240-B-002) were all obtained from R&D systems. The TGF β 1 was not acid-treated before addition. IL-21 was purchased from Cell Sciences (catalogue number CRI 172A). Supernatants were collected and tested by ELISA for IFN- γ (Endogen) or IL-17A (human IL-17A ELISA kit from eBioscience or human IL-17 duoset from R&D systems) using paired antibodies. We have observed that concentrations of TGF- β 1 ranging from 0.1 ng ml⁻¹ to 10 ng ml⁻¹ in the presence of IL-21 promote T_H17 differentiation, with lower doses of TGF β 1 associated with induction of a higher proportion of IL-17-producing cells than higher doses; use of 50 ng ml⁻¹ TGF- β 1 suppresses differentiation and IL-17 secretion. Intracytoplasmic staining was performed using standard methodologies and anti-IL-17-APC (R&D Systems) and anti-IFN- γ -PE or anti-IL-10-PE (BD Biosciences) antibodies. Cells were stimulated with phorbol 12-myristate 13-acetate (PMA; 10 ng ml⁻¹), iono-mycin (0.5 μ g ml⁻¹) and Golgistop for 5 h at 37 °C before intracellular staining.

Real-time PCR

After removing supernatants from wells in differentiation assays, 250 μ l per well of lysis buffer was added, at which point RNA was isolated using a RNeasy mini kit (Qiagen). Total RNA was converted to complementary DNA using Taqman Reverse transcription reagents (Applied Biosystems). Quantitative PCR was performed using a 7500 Fast Real-time PCR system (Applied Biosystems). All primers and probes were obtained from Applied Biosystems and used according to standard methodologies.

Acknowledgements

We thank D. Kozoriz for assistance with flow cytometric cell sorting. Supported by grants from the NIH (D.A.H., V.K.K.), the National Multiple Sclerosis Society (D.A.H., V.K.K.), the Juvenile Diabetes Research Foundation (D.A.H., V.K.K.), and the American Cancer Society (D.E.A.). D.A.H. and V.K.K. are recipients of a Javitz Investigator award from the NIH.

References

1. Kikly K, Liu L, Na S, Sedgwick JD. The IL-23/Th(17) axis: therapeutic targets for autoimmune inflammation. *Curr. Opin. Immunol* 2006;18:670–675. [PubMed: 17010592]
2. Wilson NJ, et al. Development, cytokine profile and function of human interleukin 17-producing helper T cells. *Nature Immunol* 2007;8:950–957. [PubMed: 17676044]
3. Bettelli E, et al. Reciprocal developmental pathways for the generation of pathogenic effector T_H17 and regulatory T cells. *Nature* 2006;441:235–238. [PubMed: 16648838]
4. Mangan PR, et al. Transforming growth factor- β induces development of the T_H17 lineage. *Nature* 2006;441:231–234. [PubMed: 16648837]
5. Veldhoen M, Hocking RJ, Atkins CJ, Locksley RM, Stockinger B. TGF β in the context of an inflammatory cytokine milieu supports de novo differentiation of IL-17-producing T cells. *Immunity* 2006;24:179–189. [PubMed: 16473830]
6. Korn T, et al. IL-21 initiates an alternative pathway to induce proinflammatory T_H17 cells. *Nature* 2007;448:484–487. [PubMed: 17581588]
7. Nurieva R, et al. Essential autocrine regulation by IL-21 in the generation of inflammatory T cells. *Nature* 2007;448:480–483. [PubMed: 17581589]
8. Zhou L, et al. IL-6 programs T_H-17 cell differentiation by promoting sequential engagement of the IL-21 and IL-23 pathways. *Nature Immunol* 2007;8:967–974. [PubMed: 17581537]
9. Acosta-Rodriguez EV, Napolitani G, Lanzavecchia A, Sallusto F. Interleukins 1 β and 6 but not transforming growth factor- β are essential for the differentiation of interleukin 17-producing human T helper cells. *Nature Immunol* 2007;8:942–949. [PubMed: 17676045]
10. Evans HG, Suddason T, Jackson I, Taams LS, Lord GM. Optimal induction of T helper 17 cells in humans requires T cell receptor ligation in the context of Toll-like receptor-activated monocytes. *Proc. Natl Acad. Sci. USA* 2007;104:17034–17039. [PubMed: 17942669]
11. Laurence A, O'Shea JJ. T_H-17 differentiation: of mice and men. *Nature Immunol* 2007;8:903–905. [PubMed: 17712339]
12. McGeachy MJ, et al. TGF- β and IL-6 drive the production of IL-17 and IL-10 by T cells and restrain T_H-17 cell-mediated pathology. *Nature Immunol* 2007;8:1390–1397. [PubMed: 17994024]
13. Yang XO, et al. T helper 17 lineage differentiation is programmed by orphan nuclear receptors ROR α and ROR γ . *Immunity* 2008;28:29–39. [PubMed: 18164222]
14. Wei L, Laurence A, Elias KM, O'Shea JJ. IL-21 is produced by Th17 cells and drives IL-17 production in a STAT3-dependent manner. *J. Biol. Chem* 2007;282:34605–34610. [PubMed: 17884812]




Figure 1. TGF- β and IL-21 promote T_H17 differentiation from naive CD4⁺ T cells

a, CD4⁺ T cells were sorted into populations enriched for naive or central memory T helper cells. **b**, IFN- γ and IL-17A secretion is shown from T cells stimulated for 7 days in the presence of the indicated cytokines. Standard deviation using T cells from three unrelated subjects is represented. TGF- β - and IL-21-induced IL-17 secretion is highly significant ($P < 0.01$). Tx, treatment. **c**, IL-17 secretion from naive T cells from six different donors is represented (mean \pm s.e.m.). **d**, Intracellular expression of IL-17 and IFN- γ from one of five experiments is shown.




Figure 2. TGF-β and IL-21 induce RORC2 in naive CD4⁺ T cells

a, mRNA expression levels (fold-induction relative to T cells without exogenous cytokines) of *RORC2*, *Tbet*, *GATA3*, *IL23R* and *FOXP3* are shown after naive T cells were differentiated as indicated. **b**, Fold-induction \pm s.e.m. ($n = 3$) of *IL21* and *IL22* are represented. **c**, Mean expression \pm s.e.m. of *IL17A* and *RORC2* are shown ($n = 3$) for naive T cells obtained from cord blood. **d**, Shown is intracytoplasmic staining of IL-17 and IFN-γ from cord blood naive T cells after 7 days of differentiation. Similar results were seen in another independent assay.