

Note: Snapshot PDF is the proof copy of corrections marked in EditGenie, the layout would be different from typeset PDF and EditGenie editing view.

Author Queries & Comments:

Q1 : The distinction between surnames can be ambiguous, therefore to ensure accurate tagging for indexing purposes online (e.g. for PubMed entries), please check that the highlighted surnames have been correctly identified, that all names are in the correct order and spelt correctly.

Response: Please replace B.S. Dezfuli with B. Sayyaf Dezfuli

Q2 : Please check the affiliation.

Response: Affiliation is correct

Q3 : Please check table 2 alignment

Response: Please align the words "Musculature" "Liver" "Rest of viscera" "Overall" in the centre of each column

Q4 : fig. 3b, c and see further: is there something missing after 'further'?

Response: Please delete "see further" and insert fig. 5a, e

Q5 : "Fernandez *et al.*, 2005" has been changed to "Fernández *et al.*, 2005" with respect to the reference list. Please check.

Response: Ok

Q6 : "Sitja-Bobadilla *et al.* (2016)" has been changed to "Sitjà-Bobadilla *et al.* (2016)" with respect to the reference list. Please check.

Response: Ok

Q7 : Please provide publisher name for reference "Karasev (1990)."

Response: "Please add as Publisher "Institute of Marine Research"

CM1 : Please as Verso running head put "B. Sayyaf Dezfuli et al."

Color coding for Comments and editing changes:

1. Author	(a) Text insertion; (b) Text Deletion; (c) Text Formatting; (d) Text Comments
2. Editor	All track changes and comments.
3. Assitant Editor	All track changes and comments.
4. Associate Editor	All track changes and comments.
5. Forum Editor	All track changes and comments.
6. Production editor	All track changes and comments.
7. Copyeditor	All track changes and comments.
8. Proofreader	All track changes and comments.
9. Managing editor	All track changes and comments.
10. Digital review editor	All track changes and comments.
11. Master Copy	All track changes and comments.

anisakid nematode larvae and a coccidian in *Micromesistius poutassou* from NE Atlantic waters

Verso running head : B.S. Dezfuli *et al.*

Recto running head : *Journal of Helminthology*

 B.S. Dezfuli^{1*}, E. Simoni², G. Bosi³, M. Palomba^{4,5}, S. Mattiucci^{4,5}, L. Giulietti⁶, M. Bao⁶, A. Levsen⁶, P. Cipriani^[Q1]⁶

¹ Department of Life Sciences & Biotechnology, University of Ferrara, St. Borsari 46, 44121 Ferrara, Italy

² Bioacoustics Research Laboratory, Department of Neurosciences, University of Padua, via G. Orus, 2b, 35129 Padua, Italy

³ Department of Health, Animal Science and Food Safety, Università degli Studi di Milano, St. Trentacoste 2, 20134 Milan, Italy

⁴ Department of Public Health and Infectious Diseases, 'Sapienza-University of Rome', P.le Aldo Moro, 5, 00185 Rome, Italy

⁵ Laboratory Affiliated to 'Istituto Pasteur Italia-Fondazione Cenci Bolognietti', Viale Regina Elena, 291, 00161 Roma, Italy^[Q2]

⁶ Section of contaminants and biohazards, Institute of Marine Research (IMR), Nordnes, Bergen, Norway

Author for correspondence: B.S. Dezfuli, Fax: 0039-0532 455715. E-mail: dzb@unife.it

History : received : 2020-09-13 revised : 2020-11-2 accepted : 2020-11-02

Copyright Line: Copyright © The Author(s), 2021. Published by Cambridge University Press

Abstract

A survey on *Anisakis simplex* (*sensu stricto* (s.s.)) from blue whiting, *Micromesistius poutassou*, in the north-eastern Atlantic Ocean revealed the occurrence of high infection levels of third larval stages in visceral organs and flesh. Larvae were genetically identified with a multilocus approach as *A. simplex* (s.s.). Histochemical, immunohistochemical and ultrastructural observations were conducted on 30 *M. poutassou* specimens. Gonads, pyloric caeca and flesh harboured encapsulated larvae of *A. simplex* (s.s.) but no intense host reaction was encountered around the parasite in the above organs. In the liver, the most infected organ, the larvae co-occurred with the coccidian *Goussia* sp. Within the granuloma around the *A. simplex* (s.s.) larvae, two concentric layers were recognized, an inner mostly comprising electron-dense epithelioid cells and an outer layer made of less electron-dense epithelioid cells. Macrophages and macrophage aggregates (MAs) were abundant out of the granulomas, scattered in parenchyma, and inside the MAs, the presence of engulfed *Goussia* sp. was frequent. In liver tissue co-infected with *Goussia* sp. and *A. simplex* (s.s.), hepatocytes showed cytoplasmic rarefaction and acute cell swelling. Results suggest that the host-induced encapsulation of *A. simplex* (s.s.) larvae is a strategic compromise to minimize collateral tissue damage around the larval infection sites, to facilitate the survival of both parasite and host.

Key words

- Blue whiting
- *Anisakis simplex* (s.s.)
- *Goussia*
- granuloma
- immunohistochemistry
- ultrastructure

Introduction

The blue whiting (*Micromesistius poutassou*) (Risso, 1827) is a small oceanic and benthopelagic gadoid fish, broadly distributed in North Atlantic waters and the Mediterranean Sea. The species is of considerable economic importance with total catches reaching 1538 kt in 2017, of which Norway was the largest producer with close to 400 kt (FAO, 2020). The parasite fauna of blue whiting seems to have a core of parasite species present throughout its north-east (NE) Atlantic distribution. The dominating parasite species appear to be the anisakid nematode *Anisakis simplex* (s.s.) and the coccidian *Goussia* sp., with prevalence reaching 100% for both parasites in catches from the NE Atlantic Ocean and the North Sea (Berland, 2006; Levsen *et al.*, 2018).

Many studies have focused on the adverse health effects of larval nematodes on marine fish species (reviewed by Buchmann, 2012; Mackenzie & Hemmingsen, 2014; Horbowy *et al.*, 2016; Mladineo *et al.*, 2016; Debenedetti *et al.*, 2019). Formation of granulomas on the surface of internal organs (Buchmann & Mehrdana, 2016; Dezfuli *et al.*, 2017) or in the skeletal musculature is a common response to larval nematodes in fish (Umberger *et al.*, 2013; Cipriani *et al.*, 2016). Granulomas are focal chronic inflammatory lesions that appear as nodules in tissues and organs of the host (Ferguson, 2006; Dezfuli *et al.*, 2016). Cellular immune responses in fish-parasite systems generally involve the following cell components: neutrophils (Jørgensen *et al.*, 2018;

Furtado *et al.*, 2019), mast cells (MCs) (Reite & Evensen, 2006; Buchmann, 2012; Dezfuli *et al.*, 2018), macrophage aggregates (MAs) (Agius & Roberts, 2003; Dezfuli *et al.*, 2017) and epithelioid cells (Ferguson, 2006; Dezfuli *et al.*, 2017).

The present study aimed to perform a histochemical, immunohistochemical and ultrastructural analysis of different organs of blue whiting from the NE Atlantic Ocean parasitized by *Anisakis simplex* (s.s.) and *Goussia* sp. to document the immunohistopathological responses to the parasites. Our results are the first to provide data on the cellular component of the granuloma around *Anisakis simplex* (s.s.) larvae in the liver of a fish. Indeed, scanning electron microscopy on histological sections of fish liver with any helminth has not been reported previously.

Materials and methods

Fish sampling and epidemiological data

Samples of blue whiting ($n = 150 + 30$) were obtained from a commercial fishing catch in April 2018 off St Kilda (N58°04' W09°40') in the NE Atlantic Ocean (FAO 27 area, Division VI a), on board the fishery and research vessel MS *Kings Bay* (Institute of Marine Research, cruise no. 'Kings Bay' 2018843).

Tissue samples of $n = 30$ fish were collected and processed for immunohistochemistry analysis within 1 hour post-catch. Thus, the fish stomach, pyloric caeca, intestine, liver, gonads and skeletal muscles were inspected for the presence of *Anisakis* sp. larvae. Pieces of organ tissue (approx. $15 \times 15 \times 10$ mm) containing *Anisakis* larvae were excised, photographed and then fixed in 10% neutral buffered formalin for 24 h. For negative control, tissue pieces of a single uninfected liver were processed the same way as infected organ samples. After 24 h, formalin was substituted with ethanol 70% and the samples were stored and prepared for transport to the laboratory.

Furthermore, 150 specimens were randomly selected from the same fishing catch and inspected for detection of anisakid nematodes and other parasites, to be used as a reference for overall infection data. Fish were weighed, measured and disposed for parasitological survey, conducted by the UV press method as previously described (Levsen *et al.*, 2018). The larvae localized in the fish host were counted and morphologically assigned to the genus level using bright-field microscopy following the diagnostic keys of Berland (1961). Subsequently, a subsample of 80 *Anisakis* spp. Larvae, randomly selected from morphologically confirmed *Anisakis* type I larvae, were washed in saline solution and stored at -70°C for further genetic identification.

Molecular identification

A total of $n = 100$ *Anisakis* type I larvae were identified to the species level based on the direct sequence analysis of mitochondrial (mtDNA *cox2*) and nuclear (elongation factor *EF1* α -1 of nDNA) gene loci. Total DNA was extracted from a tissue portion (≈ 2 mg) of each *Anisakis* larva. The Quick-gDNA™ Miniprep Kit (ZYMO RESEARCH) was used as extraction method. DNA obtained was quantified using a NanoDrop®TC1-E20 spectrophotometer (BioTek Synergy HT).

For the sequencing of the mtDNA *cox2* gene locus, amplification was performed using the primers 211 F (5'-TTT TCT AGT TAT ATA GAT TGR TTT YAT- 3') and 210R (5'-CAC CAA CTC TTA AAA TTA TC- 3') (Mattiucci *et al.*, 2014). Polymerase chain reaction (PCR) was carried out according to the procedures provided by Mattiucci *et al.* (2014). The sequences obtained were compared with those already deposited for the same gene in GenBank for other *Anisakis* species. *Anisakis* spp. larvae, identified by the mtDNA *cox2* gene, were sequenced at the elongation factor (*EF1* α -1 nDNA) nuclear gene. The *EF1* α -1 nDNA was amplified using the primers EF-F (5'-TCCTCAAGCGTTGTATCTGTT-3') and EF-R (5'-AGTTTTGCCACTAGCGGTTC-3') (Mattiucci *et al.*, 2016). The PCR conditions and procedures followed those previously reported (Mattiucci *et al.*, 2016). The sequences obtained for the *EF1* α -1 nDNA gene for the larval specimens were compared with those previously deposited in GenBank, at the diagnostic positions (i.e. 186 and 286) as previously detailed (Mattiucci *et al.*, 2016).

Quantitative infection assessment focused on nematode prevalence and abundance values, separately for liver, rest of viscera and skeletal muscle of fish. The epidemiological parameters considered were prevalence (P, %), mean abundance (mA), and mean intensity (mI) with standard deviation (\pm SD) and range of infection (min–max).

Histopathology

For histological purpose, fixed pieces of visceral organs and skeletal muscles of 30 *M. poutassou* were sent to the University of Ferrara (Italy). Each organ sample was embedded in paraffin wax and cut with routine histological techniques. Multiple 5- μm sections were taken from each tissue block, stained with either Alcian Blue–haematoxylin and eosin (AB/HE), Alcian Blue 8 GX pH 2.5 and periodic acid Schiff's (AB/PAS) and Giemsa, examined and photographed using a Nikon Microscope ECLIPSE 80i (Nikon, Tokyo, Japan).

Histochemistry and immunohistochemistry

Immunohistochemistry tests were carried out on sections (4–6 μm) of liver, gonads, pyloric caeca and skeletal muscles using the

antibodies and dilution indicated in [table 1](#). The sections were rehydrated and treated for 20 min with 3% H₂O₂ in Tris- buffered saline (TBS) pH 7.6 (TBS: 0.05 M Tris- HCl, 0.15 M NaCl) to block endogenous peroxidase. Slides used for the detection of the rabbit polyclonal anti-tumour necrosis factor (TNF)- α and mouse monoclonal Mast cell tryptase antibodies were heated 2 \times 5 min in microwave at 500 W in 0.01 M citrate buffer pH 6.0 for antigen retrieval. Sections were rinsed 2 \times 5 min in TBS, then incubated for 30 min with 1:20 goat normal serum or with 1:20 rabbit normal serum to avoid tissue-unspecific reaction. The use of one or the other normal serum depends on the host species in which the used primary antibody was produced (see [table 1](#)). After a washing step of 2 \times 5 min in TBS, sections were incubated at room temperature for 24 h with the primary antibody. The optimum dilution for each antibody was determined by several trials in our laboratories. Slides treated with polyclonal anti-rabbit primary antibodies were washed twice for 5 min in TBS and incubated for 1 h with 1:200 biotinylated anti-rabbit immunoglobulins (Vector Labs, Burlingame, CA, USA). The sections treated with monoclonal anti-mouse primary antibodies were incubated for 1 h with 1:200 biotinylated anti-mouse immunoglobulins (Vector Labs). Finally, slides were treated for 1 h at room temperature with the streptavidin-biotin/horseradish peroxidase complex (Vectastain® ABC kit, Vector Labs) and the reaction was developed with a freshly prepared solution of 4 mg of 3,3'-diaminobenzidine tetrahydrochloride (Sigma-Aldrich, St. Louis, MO, USA) in 10 ml of 0.05 M TBS containing 0.1 ml of 3% H₂O₂. After 15 min washing steps in running tap water, sections were quickly washed in TBS, counterstained for 15 min with Mayer's haematoxylin, de-hydrated and mounted with Eukitt.

Table 1. Antibodies used in the present study, their specificity, source and dilution.

Antibody anti-	Specificity	Source and code	Dilution
Histamine	Rabbit, polyclonal	Sigma-Aldrich, Saint Louis, MO, USA, H7403	1:50
Mast cell chymase	Mouse, monoclonal	Santa Cruz Biotechnology Inc., Santa Cruz, CA, USA, sc-59586	1:50
Mast cell tryptase	Mouse, monoclonal	Dako Glostrup, Denmark (now part of Agilent, Santa Clara, CA, USA), M7052	1:50
Serotonin	Rabbit, polyclonal	Chemicon (now part of Merck Millipore, Darmstadt, Germany), AB938	1:100
Tumor necrosis factor (TNF- α)	Rabbit, polyclonal	Abcam, Cambridge, UK, ab6671	1:100

In liver, the immunohistochemical results could be invalidated by the presence of melanin in the MAs. A melanin bleaching protocol was used to remove melanin (Chung *et al.*, 2016), treating liver sections with 0.5% H₂O₂ in 10 mM Tris/HCl pH 10.0 at 80°C for 15 min prior of the primary antibody incubation. Further control protocols were performed by (1) incubation step with phosphate-buffered saline (PBS) pH 7.4 instead of the primary antibody; (2) treatment with PBS instead of the secondary antibody; (3) melanin bleaching treatment prior of the incubation step with PBS instead of the primary antibody; (4) melanin bleaching protocol prior of the treatment with PBS instead of the secondary antibody. All these procedures gave the expected results.

Transmission electron microscopy

For transmission electron microscopy, several pieces mostly up to 7 mm in size of all ten infected livers and one single liver with no *Anisakis* sp. larva were fixed in 2.5% glutaraldehyde in 0.1 M sodium cacodylate buffer for 3 h at 4°C and embedded in Durcupan ACM resins. Stained ultrathin sections were examined in a Hitachi H-800 at an acceleration of 100 kV as detailed in Dezfuli *et al.* (2018).

Scanning electron microscopy

For scanning electron microscopy (SEM), pieces of ten livers with visible encapsulated *Anisakis* sp. larvae were prepared with routine techniques for SEM. Furthermore, 7- μ m histological sections from the above infected livers and sections from a single liver free of *Anisakis* sp. larvae were placed on coverslips, then dehydrated through a graded ethanol series and coated with gold in an Edward S 150 sputter coater. Samples were then examined with a Cambridge Stereoscan 360 (London, UK) at an acceleration voltage of 20 kV.

Results

Parasite molecular identification and epidemiological data

Out of the $n = 150$ *M. poutassou* specimens analysed, 69 were males and 81 females (total length, mean \pm SD = 273.5 ± 27.4 mm; mean weight \pm SD = 104.1 ± 38.4 g) (table 2).

Table 2. *Anisakis simplex* (s.s.) infection in *Micromesistius poutassou* [Q3].

Fish reference sample		Musculature	Liver	Rest of viscera	Overall
$n = 150$ TL = 273.5 ± 27.4 mm TW = 104.1 ± 38.4 g	Infected	136	148	149	149
	P (%)	90.7	98.7	97.3	99.3
	A	6.5	24.3	8.0	38.8
	$mI (\pm SD)(\text{min-max})$	$7.2 \pm 9.3(1-65)$	$24.6 \pm 37.7(1-264)$	$8.2 \pm 9.1(1-66)$	$39.0 \pm 50.8(3-377)$
	N_{LTot} (%)	976 (16.8%)	3647 (62.7%)	1193 (20.5%)	5816 (100%)

n , number of fish; TL, mean total length \pm SD; TW, mean total weight \pm SD; Infected, number of infected fish; P (%), prevalence; A, mean abundance; $mI (\pm SD)$, mean intensity \pm SD with range (min-max); N_{LTot} (%), number of total larvae and percentage in the indicated site of infection.

A total of 5816 nematode larvae were recovered from the viscera and flesh of the examined fish. Based on basic diagnostic morphological characters, all the third-stage larvae (L3) nematodes were recognized as *Anisakis* spp. showing larval Type I characters (*sensu* Berland, 1961).

According to the sequences of 629 bp in length from the mtDNA *cox2* gene locus, all the *Anisakis* spp. larvae ($n = 100$) were assigned to the species *A. simplex* (s.s.). The sequences obtained matched 99–100%, with the sequences previously deposited in GenBank for the species. In addition, the identity of the specimens recognized by mtDNA *cox2* sequences analysis as belonging to *A. simplex* (s.s.) was confirmed by sequence analysis of the locus *EF1 α -1* nDNA, showing of the two diagnostic nucleotide positions characteristic for the species *A. simplex* (s.s.), as reported in Mattiucci *et al.* (2016).

A total of ten mtDNA *cox2* sequences were deposited in GenBank (accession numbers: from MT989556 to MT989565).

Anisakis simplex (s.s.) larvae were detected in the liver (fig. 1a, b, c), the viscera, muscle and gonad of the fish (fig. 2a, b), in the muscle the larvae were encapsulated in tissue (fig. 2a). Data of prevalence (P), mean abundance (mA), and mean intensity (mI) of *A. simplex* (s.s.) larvae at different sites of infection (liver, rest of viscera and muscle) are given in table 2.

Fig. 1. Photos after necropsy of blue whiting, *Micromesistius poutassou*. (a) numerous *Anisakis simplex* (s.s.) (L3) larvae (arrows) encapsulated on surface of liver are visible; (b) portion of infected liver with some *A. simplex* (s.s.) larvae (arrows); (c) heavily infected liver, all the surface of the organ is covered with encapsulated anisakid larvae.





Fig. 2. Photos after necropsy of blue whiting, *Micromesistius poutassou*. (a) larva of *Anisakis simplex* (s.s.) (arrow) is encapsulated in host skeletal muscle; (b) ovary of *M. poutassou* with *A. simplex* (s.s.) larvae (arrows) encysted on organ surface.



Only one fish out of the 150 was not infected with *A. simplex* (s.s.). The overall prevalence recorded for *A. simplex* (s.s.) was $P = 99.3\%$ (149 infected out of 150), with mean intensity 39 (table 2). Prevalence and mean intensities in liver and rest viscera were $P = 98.7\%$, $mI = 24.6$; and $P = 97.3\%$, $mI = 8.2$, respectively (table 2). The prevalence in the skeletal muscle was $P = 90.7\%$, with $mI = 7.2$. The relative proportion of *A. simplex* (s.s.) larvae by infection site was 62.7% in liver, 20.5% of larvae in the rest of viscera and 16.8% in the skeletal muscle (table 2). In a few *M. poutassou*, larvae of *A. simplex* were observed on the surface of the testes and ovaries (fig. 2b).

Histopathology

During dissection and then observations of histological sections of visceral organs and flesh, it was noted that liver, gonads, pyloric caeca and hypaxial flesh surrounding visceral cavity were the organs infected with encapsulated *A. simplex* (s.s.) larvae.

Among them, liver was the most parasitized organ, with the presence of numerous larvae on its surface (fig. 3a) and, frequently, also inside the hepatic tissue. Infected liver showed varying degrees of alterations ranging from hydropic degeneration (cellular oedema) to acute cell swellings (fig. 3b, c and fig. 5a, e see further [Q4]). Additionally, the liver always harboured oocysts of the coccidian *Goussia* sp. (100% prevalence was observed also in the 30 samples used for immunohistochemistry analyses) (figs. 3b, c and 4a, b, c). Moreover, single macrophages, and MAs, were found scattered in the parenchyma and, inside the MAs, engulfed *Goussia* sp. oocysts were often noticed (fig. 3c).

Fig. 3. Histological sections of infected organs of *Micromesistius poutassou*. (a) periphery of liver with encysted *A. simplex* (s.s.) larvae (arrows)(alcian blue and periodic acid Schiff's, AB/PAS stain), scale bar = 200 µm; (b) micrograph shows liver with co-occurrence of *A. simplex* (arrow) and coccidian *Goussia* sp. (curved arrows)(AB/PAS stain), scale bar = 20 µm; (c) high magnification of macrophage aggregates (MAs) in liver, several *Goussia* sp. (arrows) are engulfed by MAs, note acute swelling of hepatocytes (alcian blue and haematoxylin & Eosin, AB/H&E stain), scale bar = 20 µm; (d) longitudinal section of a pyloric caecum with *A. simplex* (s.s.) larva (arrow) externally encysted, lack of cellular response around nematode is evident (AB/H&E stain), scale bar = 200 µm.





Fig. 4. Micrographs of immunohistochemical reactions on infected liver of *Micromesistius poutassou*. (a) periphery of liver, co-occurrence of *Goussia* sp. (curved arrows) and *A. simplex* (s.s.) larva (asterisk) is evident, note positive immunoreactivity of macrophage aggregates (MAs) to anti-mast cell tryptase (arrows), scale bar = 100 µm; (b) high magnification of MAs positive to anti-mast cell tryptase, some *Goussia* sp. (arrows) are engulfed by MAs, several hepatocytes are infected with *Goussia* sp. (curved arrows), acute swelling of hepatocytes is visible, scale bar = 50 µm; (c) single macrophage (curved arrows) shows more intense reactivity to anti-tumour necrosis factor (TNF-α) in comparison to MAs (arrows), hepatocytes infected with *Goussia* sp. show acute swelling, scale bar = 50 µm.



The skeletal muscle of many fish was infected with *A. simplex* (s.s.), mostly in the hypaxial part of the flesh, but no host cellular reaction was encountered around the encapsulated larvae. Few blue whiting harboured nematode larvae on visceral peritoneum of the gonads, nevertheless, no immune cells encircled *A. simplex* (s.s.) larvae. In several blue whiting, *A. simplex* (s.s.) larvae were encapsulated on the external surface of the pyloric caeca, observations of the histological sections did not reveal appreciable host cellular reactions (fig. 3d).

Histochemistry and immunohistochemistry


In the liver of blue whiting, macrophages and MAs were immunoreactive to the antibody anti-mast cell tryptase, mainly in peripheral zone of the organ and near the nematode encapsulated larva (fig. 4a). Indeed, positive MAs engulfed coccidian *Goussia* sp. (fig. 4b). MAs also showed a weak reactivity to the antibody anti-TNF- α (fig. 4c). A more intense reactivity to anti-TNF- α was observed in single macrophage scattered in the liver parenchyma (fig. 4c). No results were obtained with the antibodies anti-histamine, -mast cell chymase and -serotonin. In sections of infected gonads, muscle and pyloric caeca no immunoreactivity to antibody anti-mast cell tryptase and anti-TNF- α was recorded.

Transmission electron microscopy

Ultrastructurally, in the infected liver, especially hepatocytes adjacent to each nematode larva had amorphous cytoplasmic vacuolation or rarefaction, liver cell with high level of lipid tended to possess numerous osmiophilic droplets (fig. 5a). Hepatocytes nuclei were enlarged and rich in euchromatin, with a thin rim of heterochromatin in the periphery of the nucleus attached to the nuclear membrane (fig. 5b). Each *A. simplex* (s.s.) larva was surrounded by a granuloma in which two distinct layers were recognized. An inner electron-dense epithelioid cells layer (fig. 5c) was formed of flattened cells with slight vacuolated cytoplasm. Ultrastructurally, the epithelioid cells were similar to macrophages and appeared to be compressed and cytoplasmic borders were indistinct (fig. 5c, d). The outer layer was less electron dense than the inner one and formed by epithelioid cells and collagen bundles scattered among them. In the outer layer, the epithelioid cells had elongated nuclei (fig. 5c) and presented numerous desmosomes (fig. 5d).

Fig. 5. Transmission electron microscopy of infected liver of *Micromesistius poutassou*. (a) Micrograph shows lipid droplets (arrows) within the cytoplasm of hepatocytes, bad condition of tissue, namely swollen of hepatocytes and their

cytoplasmic vacuolation are notable, scale bar = 3.3 μm ; (b) high magnification of hepatocyte nucleus rich in euchromatin (arrow), thin rim of heterochromatin (curved arrows) is also evident, scale bar = 0.8 μm ; (c) interface region between *A. simplex* (s. s.) larva (arrow) and hepatic tissue, granuloma around each nematode larva is formed by two distinct layers, an inner layer of electron-dense epithelioid cells (curved arrows) and an outer layer formed by epithelioid cells with elongated nuclei (arrow heads), scale bar = 2.5 μm ; (d) higher magnification of interface region, thick arrow shows *A. simplex* (s. s.) larva, occurrence of several desmosomes (arrows) between epithelioid cells are evident, scale bar = 0.4 μm ; (e) two *Goussia* sp. (arrows) inside the cytoplasm of a hepatocyte, lipid droplet within the cytoplasm is visible, occurrence of coccidians induced displacement of the hepatocyte nuclei (curved arrows) to a marginal position, scale bar = 2.5 μm .




As mentioned earlier, the prevalence of *Goussia* sp. in the liver was 100%. Additionally, the hepatocyte cytoplasm frequently hosted more than one coccidian, consequently displacing the nucleus into a marginal position and narrowing the cytoplasm (fig. 5e).

Scanning electron microscopy

SEM on pieces of liver with encapsulated *A. simplex* (s.s.) larvae revealed that the larvae were often covered with amorphous material (fig. 6a). SEM of histological sections of infected livers showed also that co-infections with *A. simplex* (s.s.) and the coccidian *Goussia* sp. commonly occurred (fig. 6b, c). All livers of blue whiting were infected with *Goussia* sp., and, frequently, more than one *Goussia* sp. oocyte infected single hepatocytes (fig. 6c).

Fig. 6. Scanning electron microscopy (SEM) of infected liver of *Micromesistius poutassou*. (a) *Anisakis simplex* (s.s.) larva (arrow) encapsulated on liver surface and covered with mucus-like material, scale bar = 400 μm ; (b) SEM of liver histological section, typical appearance of preventricular portion of the oesophagus of *A. simplex* (s.s.) larva (arrow) encapsulated in

hepatic tissue is evident, scale bar = 100 μm ; (c) high magnification reveals co-occurrence of *A. simplex* (s.s.) larva (arrow) and *Goussia* sp. (curved arrows) inside the hepatocytes, note acute swelling and degeneration of liver cells, scale bar = 20 μm .



Discussion

Anisakis simplex (s.s.) seems to be able to utilize several pelagic and demersal teleost fish species as vectors to the definitive cetacean host level. In this respect, blue whiting appears to be one of the most important transport hosts in NE Atlantic oceanic waters (Levsen *et al.*, 2018). On the other hand, the fish species represents a suitable host in the life cycle of the species of the *A. simplex* (s.s.), being found hosting also the sibling species *A. pegreffii* in the Mediterranean Sea waters (Levsen *et al.*, 2018; Mattiucci *et al.*, 2018). Fishes acquire larval stages of the *Anisakis* spp. by ingesting infected planktonic or semi-planktonic crustacean or prey fish. Once in the new fish host, the larva penetrates the intestinal wall of fish and enters the visceral cavity, crawling over organs, encysting on their surface or migrating into the host muscle (Mattiucci *et al.*, 2018). Migration of larvae in fish musculature can occur when fish is still alive, *intra vitam* (Larsen *et al.*, 2002), or after host death, post mortem (Hauck, 1977; Cipriani *et al.*, 2016). The success of helminth infection largely depends on their capacity to manipulate and or evade the general efficiency of hosts immune system (Franke *et al.*, 2014).

According to the mtDNA *cox2* and *EF1 α -1* nDNA sequences analyses, all the *Anisakis* spp. larvae (n = 100) were identified as *Anisakis simplex* (s.s.). This finding is congruent with previous recordings on *M. poutassou* from that fishing ground of the NE Atlantic Ocean (Levsen *et al.*, 2018), and also with the geographic distribution of this parasite species (Mattiucci *et al.*, 2018). The reference sample of 150 specimens of *M. poutassou* inspected for detection of anisakid nematodes showed high infection levels, with 149 fish out of 150 infected by *A. simplex* (s.s.), and a mean intensity $I_m = 39$. Similar infection levels were already reported, from the same area, in previous studies (Karasev, 1990; Fernández *et al.*, 2005 [Q5]; Berland, 2006; Levsen *et al.*, 2018), demonstrating that this gadoid species seems to tolerate well the high intensity of *A. simplex* (s.s.) larvae.

It is well known that extraintestinal helminths frequently become encapsulated, as an evolved strategy to reduce parasite immune reactivity (Dezfuli *et al.*, 2017) and, thus, the extent of collateral damage to the infected host tissue. The granuloma is a chronic inflammatory focal response, most often organized in concentric cellular layers in/on the host tissue (Ferguson, 2006). Encapsulation is considered a mutual adaptation between parasite and the host's immune response, representing a trade-off to ensure viability of both parasite and host (Bruschi & Chiumiento, 2011; Dezfuli *et al.*, 2017). In infected gonads, pyloric caeca and fillets of blue whiting, no appreciable host cellular response was noticed around *A. simplex* (s.s.) larvae. In a similar study, no

cellular host reaction was seen in the muscles of *Paralichthys lethostigma* Jordan & Gilbert, 1884, parasitized by two species of philometrid nematodes (de Buron & Roumillat, 2010). Conversely, in skeletal muscle of *Perca fluviatilis* Linnaeus, 1758 infected with larvae of the nematode *Eustrongylides* sp. Jägerskiöld, 1909, numerous MAs and MCs were seen throughout the thick fibro-connectival layer of the capsule enclosing the larvae (Dezfuli *et al.*, 2015).

The reasons for the lack or presence of immune cells in granulomas in visceral organs and skeletal muscle of fish is unknown and open to conjecture. It has been also hypothesized that this concomitant immunity may represent a trade-off between greater pathology that may be caused by the presence of dead larvae in host organs compared to encapsulation and isolation of live anisakids (reviewed by Buchmann, 2012). Immune responses in fish against parasites depend on the eliciting agents and host species (Gauthier *et al.*, 2004) since different parasites may differentially affect the various parts of the host physiology and immunological pathways (Ferguson, 2006).

In *M. poutassou*, the liver was the most heavily infected organ, showing 100% prevalence of *Goussia* sp. and 99.3% of *A. simplex* (s.s.) larvae. In liver, recruitment of epithelioid cells, macrophages and macrophages aggregates were noticed. One of the well-conserved innate defence mechanisms is phagocytosis (Rieger *et al.*, 2012; Grayfer *et al.*, 2014). In fish, generally, two populations of professional phagocytes were described, mononuclear phagocytes (circulating monocytes and tissue macrophages) (Secombes & Ellis, 2012; Esteban *et al.*, 2015) and neutrophils (Secombes & Ellis, 2012). In infected liver of *M. poutassou* very few neutrophils were noticed. With regard to macrophages, in most fish species they exist in clusters (MAs) in the posterior kidney, spleen and less frequently in intestine and liver (Agius & Roberts, 2003) and contain pigments like hemosiderin, chromolipoids and melanin (Agius & Roberts, 2003; Secombes & Ellis, 2012; Dezfuli *et al.*, 2017).

The size and distribution of MAs might be an indicator of the general health status in fish (Latoszek *et al.*, 2019), and a biomarker of their immune capacity and activation (Stosik *et al.*, 2019). Indeed, in fish liver, MAs may develop in association with chronic inflammatory lesions (Agius & Roberts, 2003; Estensoro *et al.*, 2014; Dezfuli *et al.*, 2017). In the liver of blue whiting, single macrophages as well as numerous MAs were encountered in close proximity to *A. simplex* (s.s.) larvae and *Goussia* sp. oocysts. Data on involvement of macrophages and MAs against nematode infections appeared in (Dezfuli *et al.*, 2016; Ventura *et al.*, 2016) and relationship between MAs and protozoans was reviewed by Sitjà-Bobadilla *et al.* (2016). Indeed, those authors provided an image of carp macrophages which ingested merozoites of *Goussia carpelli*. It was reported that an increase in number and size of MAs in the spleen is the onset of the inflammatory response (Agius & Roberts, 2003) and in some instances it is also due to occurrence of engulfed parasites in MAs (Estensoro *et al.*, 2014; Sitjà-Bobadilla *et al.*, 2016 [Q6]). Livers of all 150 blue whiting harboured high numbers of coccidians, and MAs engulfed variable numbers of *Goussia* sp. Thus, our assumption is most likely there could be a direct relationship between number of coccidian and that of MAs of blue whiting. Detoxification, recycling of exogenous and endogenous materials, and focal destruction have been attributed to MAs (Agius & Roberts, 2003). Frequently, numerous single macrophages, and MAs, were noticed in close proximity to the *A. simplex* (s.s.) larvae, but we failed to find where MAs had engulfed portions of the nematode, presumably because of large size of parasite larva.

In the infected liver of blue whiting, the layers of the capsule surrounding each larva were made by epithelioid cells. Epithelioid cells show morphological similarity to epithelial cells, and it is generally accepted that they are transformed macrophages that form upon persistent inflammatory stimulation (Gauthier *et al.*, 2004). The origin of epithelioid cells from macrophages was documented from *in vitro* investigation (Secombes, 1996). It is interesting to note that the liver of fish is able to produce granuloma/capsule formed by metabolically active cells like macrophages and less active cells like epithelioid cells. Similar granuloma/capsules were encountered in the liver of another fish species naturally infected with the nematode *Brevimulticaecum* sp. Shikhobalova and Mozgovoi, 1952 (Dezfuli *et al.*, 2016).

With regard to *Anisakis* spp. larvae in fish, it was mentioned that host cellular reactions are likely one of the stimuli inducing larval coiling and production of a defence wall made by an inner layer of dead host cells, and outer layer of leucocytes (Larsen *et al.*, 2002). Even at the high prevalence and intensity of infection of *A. simplex* (s.s.) in blue whiting liver, it seems unlikely that the parasite is able to induce severe damage in the fish, with a consequent host population decline. Most probably a balance between nematode damage and fish defence response occurs in liver thanks to potential liver compensatory properties which is well known (Jia *et al.*, 2019). In terms of fish host - *Anisakis* interaction, recent studies have shown that a significant up-regulation of the gene coding for a Kunitz-type trypsin inhibitor protein occurs in *A. simplex* (s.s.) larvae infecting the liver of blue whiting (Palomba *et al.*, 2020). The Kunitz-type trypsin inhibitor protein of *Anisakis* seems to have a role in the inhibition of the host's trypsinases, as previously observed in other nematode species (Bernard & Peanasky, 1993). It could be thus hypothesized that an invading larva of *A. simplex* (s.s.) increases transcripts of an inhibitor protein against the fish trypsinases, abundant in the liver tissue, as a possible result of the parasite's mechanism of evasion of the host immune response (Palomba *et al.*, 2020).

In fish, TNF- α is a cytokine involved in modulation of different immune cells in response to the presence of pathogens and tissue inflammation (Seno *et al.*, 2002; Hong *et al.*, 2013; Williams *et al.*, 2015). TNF- α was mainly produced by monocytes and macrophages (Hehlhans & Pfeffer, 2005), and was observed in cells of several tissues and organs of the turbot (Ronza *et al.*, 2015).

The presence of immunoreactive macrophages to the antibody anti-TNF- α in the liver of the blue whiting is clearly associated with the hepatic degeneration due to the *A. simplex* (s.s.) larvae and *Goussia* sp. In the liver of rats, an increase of TNF- α level was observed after the degeneration induced by a traumatic event (Gao *et al.*, 2014).

Tryptase is a serine protease produced and stored in mast cell granules (McNeil *et al.*, 2007). Tryptase has been reported in gill mast cells of zebrafish (Dobson *et al.*, 2008), and in mucosal mast cells of the intestine of the fat snook, *Centropomus parallelus* Poey, 1860 (da Silva *et al.*, 2017), suggesting that the occurrence of tryptase is supportive in host tissue repair processes (Duchesne *et al.*, 2011). In the present parasitized liver samples of blue whiting, numerous macrophages and MAs were noticed, and they were positive to the antibody anti-mast cell tryptase. The presence of tryptase may suggest that liver tissue repairing processes were activated and employed as a response to the parasite infections.

In conclusion, the liver of blue whiting was the most infected organ in the present study. However, *A. simplex* (s.s.) larvae were efficiently sequestered inside the granulomas, since the surrounding parenchyma did not show any signs of fibrosis. Most likely, this focal encapsulation of the nematode permits uninfected portions of the liver to maintain their functions and provide adequate fish health condition and, thus, proper recruitment of the NE Atlantic blue whiting stock. Above assumption is based also on overall acceptable health condition of fish during inspection, indeed, it seems that *Goussia* sp. has a low virulence on fish liver with 100% prevalence in this organ and several hepatocytes with coccidians inside.

Acknowledgements

We would like to G. Servadei, C. Maestri for technical helps in laboratory and P. Boldrini from Centre of Electron Microscopy of University of Ferrara for technical assistance.

Financial support

This study was supported by grants from the University of Ferrara to B.S.D (FAR, year 2019). S. Mattiucci has contributed to the study with a grant from 'Istituto Pasteur Italia-Fondazione Cenci Bolognetti'.

Conflict of interest

None.

Ethical standards

The authors assert that all procedures contributing to this study comply with the ethical standards of the relevant national and institutional guides.

References

- Agius C and Roberts RJ** (2003) Melano-macrophage centers and their role in fish pathology. *Journal of Fish Diseases* **26**, 499–509.
- Berland B** (1961) Nematodes from some Norwegian marine fishes. *Sarsia* **2**, 1–50.
- Berland B** (2006) Musings on nematode parasites. *Fisken Go Havent* **11**, 1–30.
- Bernard VD and Peanasky RJ** (1993) The serine protease inhibitor family from *Ascaris suum*: Chemical determination of the five disulphide bridges. *Archives of Biochemistry Biophysics* **303**, 367–376.
- Bruschi F and Chiumiento L** (2011) *Trichinella* inflammatory myopathy: Host or parasite strategy? *Veterinary Parasitology* **23**, 42.
- Buchmann K** (2012) Fish immune responses against endoparasitic nematodes – experimental models. *Journal of Fish Diseases* **35**, 623–635.
- Buchmann K and Mehrdana F** (2016) Effects of anisakid nematodes *Anisakis simplex* (s.l.), *Pseudoterranova decipiens* (s.l.) and *Contracaecum osculatatum* (s.l.) on fish and consumer health. *Food and Waterborne Parasitology* **4**, 13–22.
- Chung JY, Choi J, Sears JD, et al.** (2016) A melanin-bleaching methodology for molecular and histopathological analysis of formalin-fixed paraffin-embedded tissue. *Laboratory Investigation* **96**, 1116–1127.
- Cipriani P, Acerra V, Bellisario B, Sbaraglia GL, Cheleschi R, Nascetti G and Mattiucci S** (2016) Larval migration of the zoonotic parasite *Anisakis pegreffii* (Nematoda: Anisakidae) in European anchovy, *Engraulis encrasicolus*: Implications to seafood safety. *Food Control* **59**, 148–157.
- da Silva WF, Simões, MJ, Gutierrez RC, Egami MI, Santos AA, Antoniazzi MM and Ranzani-Paiva MJT** (2017) Special dyeing, histochemistry, immunohistochemistry and ultrastructure: A study of mast cells/eosinophilic granules cells (MCs/EGC) from *Centropomus parallelus* intestine. *Fish & Shellfish Immunology* **60**, 502–508.

- Debenedetti ÁL, Madrid E, Trelis M, Codes FJ, Gil-Gómez F, Sáez-Durán S and Fuentes MV** (2019) Prevalence and risk of anisakid Larvae in fresh fish frequently consumed in Spain: An overview. *Fishes* **4**, 1–16.
- de Buron I and Roumillat WA** (2010) Histopathology of two philometrid parasites of the southern flounder, *Paralichthys lethostigma*. *Journal of Wildlife Diseases* **46**, 277–282.
- Dezfuli BS, Manera M, Lorenzoni M, Pironi F, Shinn AP and Giari L** (2015) Histopathology and the inflammatory response of European perch, *Perca fluviatilis* muscle infected with *Eustrongylides* sp. (Nematoda). *Parasites & Vectors* **8**, 227–220.
- Dezfuli BS, Fernandes CE, Galindo GM, Castaldelli G, Manera M, DePasquale JA and Giari L** (2016) Nematode infection in liver of the fish *Gymnotus inaequilabiatus* (gymnotiformes: Gymnotidae) from the pantanal region in Brazil: Pathobiology and inflammatory response. *Parasites & Vectors* **9**, 473.
- Dezfuli BS, Manera M, DePasquale JA, Pironi F and Giari L** (2017) Liver of the fish *Gymnotus inaequilabiatus* and nematode larvae infection: Histochemical features and expression of proliferative cell nuclear antigen. *Journal of Fish Diseases* **40**, 1765–1774.
- Dezfuli BS, Giari L, Lorenzoni M, Carosi A, Manera M and Bosi G** (2018) Pike intestinal reaction to *Acanthocephalus lucii* (acanthocephala): Immunohistochemical and ultrastructural surveys. *Parasites & Vectors* **11**, 424.
- Dobson JT, Seibert J, The EM, Da'as S, Fraser RB, Paw BH and Berman JN** (2008) Carboxypeptidase5 identifies a novel mast cell lineage in the zebrafish providing new insight into mast cell fate determination. *Blood* **112**, 2969–2972.
- Duchesne E, Tremblay MH and Côté CH** (2011) Mast cell tryptase stimulates myoblast proliferation; a mechanism relying on protease activated receptor-2 and cyclooxygenase-2. *BMC Musculoskeletal Disorders* **12**, 235.
- Esteban MÁ, Cuesta A, Chaves-Pozo E and Meseguer J** (2015) Phagocytosis in teleosts. Implications of the new cells involved. *Biology* **4**, 907–922.
- Estensoro I, Mulero I, Redondo M, Alvarez-Pellitero P, Mulero V and Sitjà-Bobdella A** (2014) Modulation of leukocytic populations of gilthead sea bream (*Sparus aurata*) by the intestinal parasite *Enteromyxum leei* (Myxozoa: Myxosporidia). *Parasitology* **141**, 425–440.
- FAO** (2020, April 22) *Fishery Statistical Collections - Global Capture Production* [Document]. URL Retrieved from <http://www.fao.org/fishery/statistics/global-capture-production/en>
- Ferguson HW** (2006) *Systemic pathology of fish: A text and atlas of normal tissues in teleosts and their responses in disease*. 2nd edn. London, UK, Scotian Press.
- Fernández M, Aznar FJ, Montero FE and Raga JA** (2005) Endoparasites of the blue whiting, *Mcromesistius poutassou* from north-west Spain. *Journal of Helminthology* **79**, 15–21.
- Franke F, Rahn AK, Dittmar J, Erin N, Rieger JK, Haase D and Scharsack JP** (2014) *In vitro* leukocyte response of three-spined sticklebacks (*Gasterosteus aculeatus*) to helminth parasite antigens. *Fish & Shellfish Immunology* **36**, 130–140.
- Furtado WE, Cardoso L, Figueredo AB, Marchiori NC and Martins ML** (2019) Histological and hematological alterations of silver catfish *Rhamdia quelen* highly parasitized by *Lernaea cyprinacea*. *Diseases of Aquatic Organisms* **135**, 157–168.
- Gao M, Xie B, Gu C, Li H, Zhang F and Yu Y** (2014) Targeting the proinflammatory cytokine tumor necrosis factor- α to alleviate cardiopulmonary bypass-induced lung injury (review). *Molecular Medicine Reports* **11**, 2373–2378.
- Gauthier DT, Vogelbein WK and Ottinger CA** (2004) Ultrastructure of *Mycobacterium marinum* granuloma in striped bass *Morone saxatilis*. *Diseases of Aquatic Organisms* **62**, 121–132.
- Grayfer L, Hodgkinson JW and Belosevic M** (2014) Antimicrobial responses of teleost phagocytes and innate immune evasion strategies of intracellular bacteria. *Developmental & Comparative Immunology* **43**, 223–242.
- Hauck AK** (1977) Occurrence and survival of the larval nematode *Anisakis* sp. in the flesh of fresh, frozen, brined, and smoked pacific herring, *Clupea harengus pallasi*. *Journal of Parasitology* **63**, 515–519.
- Hehlgans T and Pfeffer K** (2005) The intriguing biology of the tumour necrosis factor/tumour necrosis factor receptor superfamily: Players, rules and the games. *Immunology* **115**, 1–20.
- Hong S, Li R, Xu Q, Secombes CJ and Wang T** (2013) Two types of TNF- α exist in teleost fish: Phylogeny, expression, and bioactivity analysis of type-II TNF- α 3 in rainbow trout *Oncorhynchus mykiss*. *Journal of Immunology* **191**, 5959–5972.
- Horbowy J, Podolska M and Nadolna-Altyń K** (2016) Increasing occurrence of anisakid nematodes in the liver of cod (*Gadus morhua*) from the Baltic Sea: Does infection affect the condition and mortality of fish. *Fisheries Research* **179**, 98–103.
- Jia J, Qin J, Yuan X, Liao Z, Huang J, Wang B, Sun C and Li W** (2019) Microarray and metabolome analysis of hepatic response to fasting and subsequent refeeding in zebrafish (*Danio rerio*). *BMC Genomics* **20**, 919.

- Jørgensen LG, Korbust R, Jeberg F, Kania PW and Buchmann K** (2018) Association between adaptive immunity and neutrophil dynamics in zebrafish (*Danio rerio*) infected by a parasitic ciliate. *PLoS One* **11**, 1–18.
- Karasev AB** (1990) Ecological and geographical analysis of the northeast Atlantic blue whiting parasitic fauna. pp. 307–318 in **Monstad T** (Ed) *Biology and fisheries of Norwegian spring spawning herring and blue whiting in the northeast Atlantic*. Bergen. [Q71]
- Larsen AH, Bresciani J and Buchmann K** (2002) Interactions between ecto- and endoparasites in trout *Salmo trutta*. *Veterinary Parasitology* **103**, 167–173.
- Latoszek E, Kamaszewski M, Milczarek K, Puppel K, Szudrowicz H, Adamski A and Ostaszewska T** (2019) Histochemical characteristics of macrophages of butterfly splitfin *Ameioba splendens*. *Folia Biologica (Kraków)* **67**, 53–60.
- Levsen A, Svanevik CS, Cipriani P, Mattiucci S, Gay M, Hastie LC and Pierce GJ** (2018) A survey of zoonotic nematodes of commercial key fish species from major European fishing ground-introducing the FP7 PARASITE exposure assessment study. *Fisheries Research* **202**, 4–21.
- Mackenzie K and Hemmingsen W** (2014) Parasites as biological tags in marine fisheries research: European Atlantic waters. *Parasitology* **142**, 1–14.
- Mattiucci S, Cipriani P, Webb SC, Paoletti M, Marcer F, Bellisario B and Nascetti G** (2014) Genetic and morphological approaches distinguish the three sibling species of the *Anisakis simplex* species complex, with a species designation as *Anisakis berlandi* n. sp. for *A. simplex* sp. C (Nematoda: Anisakidae). *Journal of Parasitology* **100**, 199–214.
- Mattiucci S, Acerra V, Paoletti M, Cipriani P, Levsen A, Webb SC and Nascetti G** (2016) No more time to stay “single” in the detection of *Anisakis pegreffii*, *A. simplex* (s. s.) and hybridization events between them: A multi-marker nuclear genotyping approach. *Parasitology* **143**, 998–1011.
- Mattiucci S, Cipriani P, Levsen A, Paoletti M and Nascetti G** (2018) Molecular epidemiology of anisakis and anisakiasis: An ecological and evolutionary road Map. *Advances in Parasitology* **99**, 93–263.
- McNeil HP, Adachi R and Stevens RL** (2007) Mast cell-restricted tryptases: Structure and function in inflammation and pathogen defense. *Journal of Biological Chemistry* **282**, 20785–20789.
- Mladineo I, Popovic M, Drmic-Hofman I and Poljak V** (2016) A case report of *Anisakis pegreffii* (Nematoda, anisakidae) identified from archival paraffin sections of a Croatian patient. *BMC Infectious Diseases* **16**, 42.
- Palomba M, Cipriani P, Giulietti L, Levsen A, Nascetti G and Mattiucci M** (2020) Differences in gene expression profiles of seven proteins in third-stage larvae of *Anisakis simplex* (*sensu stricto*) by sites of infection in blue whiting (*Micromesistius poutassou*). *Genes* **11**, 559.
- Reite OB and Evensen Ø** (2006) Inflammatory cells of teleostean fish: A review focusing on mast cells/eosinophilic granule cells and rodlet cells. *Fish Shellfish Immunology* **20**, 192–208.
- Rieger AM, Konowalchuk JD, Grayfer L, Katzenback BA, Havixbeck JJ, Kiemele MD and Barreda DR** (2012) Fish and mammalian phagocytes differentially regulate pro-inflammatory and homeostatic responses *in vivo*. *PLoS One* **7**, e47070.
- Ronza P, Losada AP, Villamarín A, Bermúdez R and Quiroga MI** (2015) Immunolocalization of tumor necrosis factor alpha in turbot (*Scophthalmus maximus*, L.) tissues. *Fish & Shellfish Immunology* **45**, 470–476.
- Secombes CJ** (1996) The nonspecific immune system: Cellular defenses. pp. 63–103 in **Iwama G, Nakanishi T** (Eds) *The fish immune system*. London, Academic Press.
- Secombes CJ and Ellis AE** (2012) The immunology of teleosts. pp. 144–664 in **Roberts RJ** (Ed) *Fish pathology*. Chichester, Blackwell Publishing.
- Seno H, Sawada M, Fukuzawa H, Morita-Fujisawa Y, Takaishi S, Hiai H and Chiba T** (2002) Involvement of tumor necrosis factor alpha in intestinal epithelial cell proliferation following paneth cell destruction. *Scandinavian Journal of Gastroenterology* **37**, 154–160.
- Sitjà-Bobadilla A, Estensoro I and Pérez-Sánchez J** (2016) Immunity to gastrointestinal microparasites of fish. *Developmental and Comparative Immunology* **64**, 187–201.
- Stosik MP, Tokarz-Deptuła B and Deptuła W** (2019) Melanomacrophages and melanomacrophage centres in osteichthyes. *Central European Journal of Immunology* **44**, 201–205.
- Umberger CM, de Buron I, Roumillat WA and Mcelroy E** (2013) Effects of a muscle-infecting parasitic nematode on the locomotor performance of their fish host. *Journal of Fish Biology* **82**, 1250–1258.
- Ventura AS, Ishikawa MM, de Araujo Gabriel AM, Silbiger HLN, Cavichiolo F and Takemoto RM** (2016) Histopathology from liver of tuiuba (*Gymnotus* spp.) parasitized by larvae of nematodes. *Ciência Rural* **46**, 1233–1239.

Williams JM, Duckworth CA, Burkitt MD, Watson AJM, Campbell BJ and Pritchard DM (2015) Epithelial cell shedding and barrier function: A matter of life and death at the small intestinal villus tip. *Veterinary Pathology* **52**, 445–455.

Loading https://sonoisa.github.io/xyjax_ext/xypic.js