

Impact of Extended-Duration Shifts on Medical Errors, Adverse Events, and Attentional Failures

Laura K. Barger^{1,2}, Najib T. Ayas^{3,4,5}, Brian E. Cade¹, John W. Cronin^{1,2}, Bernard Rosner⁶, Frank E. Speizer⁶, Charles A. Czeisler^{1,2*}

1 Division of Sleep Medicine, Department of Medicine, Brigham and Women's Hospital, Boston, Massachusetts, United States of America, **2** Division of Sleep Medicine, Harvard Medical School, Boston, Massachusetts, United States of America, **3** Sleep Disorders Program, University of British Columbia, Vancouver, British Columbia, Canada, **4** Department of Medicine, Vancouver General Hospital, Vancouver, British Columbia, Canada, **5** Center for Clinical Epidemiology and Evaluation, Vancouver Coastal Health Research Institute, Vancouver, British Columbia, Canada, **6** Channing Laboratory, Department of Medicine, Brigham and Women's Hospital, Harvard Medical School, Massachusetts, United States of America

Funding: See section at end of manuscript.

Competing Interests: See section at end of manuscript.

Academic Editor: Emmanuel Mignot, Stanford University, United States of America

Citation: Barger LK, Ayas NT, Cade BE, Cronin JW, Rosner B, et al. (2006) Impact of extended-duration shifts on medical errors, adverse events, and attentional failures. *PLoS Med* 3(12): e487. doi:10.1371/journal.pmed.0030487

Received: January 5, 2006

Accepted: September 29, 2006

Published: December 12, 2006

Copyright: © 2006 Barger et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abbreviations: ACGME, Accreditation Council for Graduate Medical Education; AHRQ, Agency for Healthcare Research and Quality; CI, confidence interval; NRMP, National Residency Matching Program

* To whom correspondence should be addressed. E-mail: caczeisler@hms.harvard.edu

ABSTRACT

Background

A recent randomized controlled trial in critical-care units revealed that the elimination of extended-duration work shifts (≥ 24 h) reduces the rates of significant medical errors and polysomnographically recorded attentional failures. This raised the concern that the extended-duration shifts commonly worked by interns may contribute to the risk of medical errors being made, and perhaps to the risk of adverse events more generally. Our current study assessed whether extended-duration shifts worked by interns are associated with significant medical errors, adverse events, and attentional failures in a diverse population of interns across the United States.

Methods and Findings

We conducted a Web-based survey, across the United States, in which 2,737 residents in their first postgraduate year (interns) completed 17,003 monthly reports. The association between the number of extended-duration shifts worked in the month and the reporting of significant medical errors, preventable adverse events, and attentional failures was assessed using a case-crossover analysis in which each intern acted as his/her own control. Compared to months in which no extended-duration shifts were worked, during months in which between one and four extended-duration shifts and five or more extended-duration shifts were worked, the odds ratios of reporting at least one fatigue-related significant medical error were 3.5 (95% confidence interval [CI], 3.3–3.7) and 7.5 (95% CI, 7.2–7.8), respectively. The respective odds ratios for fatigue-related preventable adverse events, 8.7 (95% CI, 3.4–22) and 7.0 (95% CI, 4.3–11), were also increased. Interns working five or more extended-duration shifts per month reported more attentional failures during lectures, rounds, and clinical activities, including surgery and reported 300% more fatigue-related preventable adverse events resulting in a fatality.

Conclusions

In our survey, extended-duration work shifts were associated with an increased risk of significant medical errors, adverse events, and attentional failures in interns across the United States. These results have important public policy implications for postgraduate medical education.

The Editors' Summary of this article follows the references.



Introduction

Although the Accreditation Council for Graduate Medical Education (ACGME) has recently placed limitations on resident work hours in an attempt to reduce fatigue-related medical errors, the practice of working for more than 24 h consecutively remains the cornerstone of American post-graduate medical education. Moreover, a 1999 report from the Institute of Medicine revealed that between 48,000 and 98,000 deaths each year occur due to a medical error [1].

A recent randomized trial reported that interns working extended-duration shifts (defined as at least 24 h continuously at work) had significantly more polysomnographically recorded attentional failures and made significantly more serious medical errors than those scheduled to work shifts 16 h or longer [2,3]. However, that trial was not large enough to determine whether extended-duration work shifts increased the risk of preventable adverse events, for example injury due to a non-intercepted serious error in medical management, although a trend in that direction was reported [3]. To address the impact of extended-duration work shifts on patient safety in a large and more diverse population of interns, we conducted a nationwide study of US interns in which we collected monthly data on self-reported attentional failures (defined as “nodding off or falling asleep” during patient-care or educational activities), significant medical errors, and preventable adverse events (including fatalities).

Methods

Details regarding participant recruitment have been described in detail elsewhere [4]. Briefly, in the spring of 2002, we sent email announcements to all individuals who successfully matched to a residency program in the National Residency Matching Program (NRMP) and to all known email addresses of graduating fourth-year medical students of US programs in an effort to reach as many of the 18,447 individuals who matched to residencies via the NRMP. These emails directed potential participants to a secure Web site which provided detailed information about the study and enabled participants to provide electronic informed consent.

In June 2002, an email was sent to the 3,429 interns who had volunteered to participate in the study, and they were directed to a password-coded secure Web site to complete a baseline survey that solicited background data. From July 2002 to May 2003, on the 28th day of each month, emails were sent to those who had completed the baseline survey (the study cohort) to direct participants to a secure Web site to complete a monthly survey. Monthly surveys were available on the Web site until the 27th day of the next month, and we ensured that each participant answered each monthly survey only once.

The monthly survey contained detailed questions regarding work hours, sleep, activities during the month, number of days off, and the number of extended-duration work shifts. Among the 60 questions they were asked each month, interns were asked to report whether they had made a significant medical error (“Do you believe sleep deprivation or fatigue caused you to make a significant medical error?” [henceforth referred to here as “fatigue-related errors”] and “Do you believe you made any significant medical errors other than due to sleep deprivation or fatigue?” [henceforth referred to

here as “non-fatigue-related errors”]). If they answered affirmatively, they were directed to a supplementary survey that elicited further information about the error(s), including whether they had resulted in an adverse patient outcome (i.e., a preventable adverse event) or a patient fatality (i.e., a preventable adverse event resulting in a fatality). Moreover, interns were also asked to report how many times they had nodded off or had fallen asleep (attentional failure) during specific patient-care activities (during surgery and while talking to, or examining, patients) and educational activities (during rounds with the attending physicians and during lectures, seminars, or grand rounds). The remaining questions addressed secondary outcomes such as caffeine usage, health, and mood, and served as distracters for the main hypothesis.

The Web sites were hosted and maintained by Pearson Assessments (<http://www.pearsonncs.com>). Data were transmitted electronically on a weekly basis through secure means from Pearson Assessments to the Brigham and Women’s Hospital. All demographic and potentially identifiable data were then stored separately from the main database. A certificate of confidentiality was issued by the Centers for Disease Control; the data were also protected by federal statute (Public Health Service Act 42 USC) as a consequence of funding from the Agency for Healthcare Research and Quality. The Brigham and Women’s Hospital/Partners HealthCare System Human Research Committee approved the procedures for the protocol, and electronic informed consent was obtained from all participants.

Work-Hour Validation

A random subset of participants (7%) completed daily work/sleep diaries, and these diaries were validated in a separate study using continuous work-hour monitoring by direct observation and polysomnographic recordings [2]. Those completing the work/sleep diaries recorded daily work hours for at least 21 out of 28 d and completed the corresponding monthly survey. Pearson product-moment correlation was used to determine the association between daily average work hours and number of extended-duration work shifts reported in the diary and through the monthly survey.

Statistical Analysis

Our analysis to determine whether the number of extended-duration shifts worked per month was associated with significant medical errors and with attentional failures during surgery, while examining patients, while on rounds with attending physicians, and during educational events employed a case-crossover component of self-matching [5,6]. Specifically, months were classified according to the number of reported extended-duration shifts worked (i.e., zero, between one and four, and five or more shifts) per month and whether or not a particular outcome occurred at least once during the month. The denominator for our events was thus the intern-month. That is, each participant was considered as a separate stratum in the analysis, and therefore interns acted as their own controls. This case-crossover analysis thus eliminated the need to account for potential between-participant confounders such as age, gender, or medical specialty. A Mantel-Haenszel test was then used to calculate a pooled odds ratio of at least one

outcome occurring during months with between one and four, or five or more extended-duration shifts worked (using months with no extended-duration shifts as the comparison group) [7].

To address potential reporting bias, we conducted a sub-analysis of the data from the 682 interns who completed all monthly surveys. In addition, given that the ACGME established new resident work-hour guidelines in 2003, we conducted a sub-analysis in which we included only those intern-months that were in compliance with current ACGME guidelines based on the frequency of extended-duration work shifts (i.e., those intern-months with nine or fewer extended-duration work shifts).

To limit this analysis to months in which interns worked full time, months reported to have fewer than 150 h worked were excluded. Odds ratios are reported with 95% confidence intervals [CIs]. SAS 8.2 (SAS Institute [http://www.sas.com]) was used for statistical analysis, and $p < 0.05$ defined statistical significance.

Results

As we have reported elsewhere [4], 18.6% of the interns who were sent an advertisement announcing the study volunteered to participate and enrolled electronically in the study (3,429 interns). Of those volunteers, 2,737 completed the baseline survey and were considered to be the study cohort (80% response rate of volunteers). The mean number of surveys completed per participant was 7.2 ± 4.0 (mean \pm standard deviation); 682 participants (25%) completed all 12 surveys. The number of surveys received per month ranged from 1,167 to 2,118 ($1,548 \pm 376$). A total of 19,740 surveys (2,737 baseline surveys and 17,003 monthly surveys) were collected. Of the intern-months, 81% were in compliance with current ACGME guidelines with respect to the frequency of extended-duration work shifts.

Our survey cohort was 53% female and 47% male with a mean age of 28.0 ± 3.9 y. Compared with data obtained from the Electronic Residency Application Service (41% female, 56% male, mean age = 30.2 y [P. Jolly, Association of American Medical Colleges, personal communication]), our cohort was slightly younger and had a relatively greater proportion of women. Of our participants, 85% had graduated from US medical schools in 43 states, and 15% had graduated from foreign medical schools. Thus, we had fewer foreign medical graduates compared with the overall statistics from the NRMP (74% US medical graduates, 26% foreign medical graduates) [8]. The distribution of medical/surgical specialties of the survey participants was similar to the distribution of specialties matched through the NRMP [4]. The mean number of extended-duration work shifts reported per month was 3.9 ± 3.4 (median = 4; range = 0–18), and their average duration was 32.0 ± 3.7 h [4]. The mean weekly work hours reported in each of the three categories of extended-duration shifts per month (zero, between one and four, and five or more extended-duration shifts) were 55.5 ± 13.8 (29% of intern-months), 64.8 ± 15.5 (24% of intern-months), and 79.3 ± 14.8 h (47% of intern-months), respectively.

Work-Hour Validation

As reported elsewhere [4], the Pearson correlation coefficients for the work hours and number of extended-duration

work shifts reported in the monthly survey versus actual hours and extended-duration shifts worked, as indicated by the daily work diaries completed by a subset of our cohort, was $r = 0.76$ ($n = 192$, $p < 0.001$), and $r = 0.94$ ($n = 40$, $p < 0.001$), respectively.

Significant Medical Errors and Preventable Adverse Events

The frequency of extended-duration work shifts had a substantial impact on reported rates of fatigue-related significant medical errors. That is, the reported rates of fatigue-related significant medical errors were increased substantially as the number of extended-duration work shifts increased (Table 1). Interns reported at least one fatigue-related significant medical error in 3.8% of months with no extended-duration shifts. However, at least one fatigue-related significant medical error was reported in 9.8% of months with between one and four extended-duration work shifts and 16% of months with five or more extended-duration work shifts. Reported rates of fatigue-related significant medical errors that resulted in a preventable adverse patient outcome also increased as the number of reported extended-duration shifts worked per month increased (0.2%, 1.1%, and 1.6%, respectively, in months with zero, between one and four, and five or more extended-duration shifts per month). A similar significant relationship was seen with fatigue-related errors associated with a fatality.

The odds ratios of reporting at least one significant medical error due to fatigue were 3.5 (95% CI, 3.3–3.7) and 7.5 (95% CI, 7.2–7.8), respectively, in months with between one and four reported extended-duration shifts and with five or more reported extended-duration shifts compared to months with no extended-duration shifts (reference group) (Table 1). A similar relationship was seen with fatigue-related preventable adverse events, with odds ratios of 8.7 (95% CI, 3.4–22) and 7.0 (95% CI, 4.3–11), respectively. Although the total number of medical errors resulting in a fatality was low, this rate also significantly increased in months with five or more extended-duration shifts per month (odds ratio = 4.1, 95% CI, 1.4–12). The number of reported fatigue-related medical errors increased as the number of extended-duration shifts per month increased regardless of gender or age. Males: odds ratio = 4.7 (95% CI, 4.0–5.4) and odds ratio = 8.8 (95% CI, 8.0–9.6) for between one and four extended-duration shifts and five or more extended-duration shifts per month, respectively. Females: odds ratio = 2.9 (95% CI, 2.6–3.2) and odds ratio = 6.7 (95% CI, 6.2–7.1) for between one and four extended-duration shifts and five or more extended-duration shifts per month, respectively. For participants below age 28 years, odds ratio = 3.8 (95% CI, 3.4–4.3) and odds ratio = 6.6 (95% CI, 6.1–7.2) for between one and four extended-duration shifts and five or more extended-duration shifts per month, respectively. For participants over age 28 years, odds ratio = 3.2 (95% CI, 2.8–3.7) and odds ratio = 8.6 (95% CI, 7.8–9.4) for between one and four extended-duration shifts and five or more extended-duration shifts per month, respectively (not shown in Table 1).

Significant medical errors made for reasons other than fatigue or sleep deprivation were not as strongly associated with the number of extended-duration shifts worked per month. Although rates of non-fatigue-related significant medical errors were greater in months with more frequent extended-duration work shifts, the relationship was modest

Table 2. Self-Reported Attentional Failures per Person-Month with Corresponding Odds Ratios (95% CIs)

Attentional Failure	Extended-Duration Shifts (0)			Extended-Duration Shifts (1–4)			Extended-Duration Shifts (≥5)		
	Number of Person-Months with Positive Response	Rate Of Positive Response/ Person-Month	Number of Person-Months with Positive Response	Rate of Positive Response/ Person-Month	Number of Person-Months with Positive Response	Odds Ratio (95% CI)	Number of Person-Months with Positive Response	Rate of Positive Response/ Person-Month	Odds Ratio (95% CI)
Nodding off or falling asleep during surgery	45	0.028	1,933	0.044	85	2.1 (1.7–2.7)	4,298	0.073	1.4 (1.3–1.6)
Nodding off or falling asleep while talking to or examining patients	85	0.028	3,275	0.039	128	1.5 (1.3–1.7)	7,245	0.054	2.1 (2.0–2.2)
Nodding off or falling asleep during rounds with the attending physician	192	0.081	2,990	0.15	455	2.3 (2.2–2.4)	6,817	0.22	5.5 (5.4–5.7)
Nodding off or falling asleep during lectures, seminars, or grand rounds	1,392	0.46	3,175	0.59	1,873	1.99 (1.96–2.03)	7,075	0.70	4.3 (4.3–4.4)

Number of person months vary from outcome as missing values for the outcome (i.e., did not answer yes or no) were eliminated from the analysis. Also, for surgery, rounds with attending physicians, examining patients, or attending lectures, participants were also asked if they did not do any of these activities that month. If they did not do any of these activities that month, these months were also excluded from the analysis. Rates represent the proportion of months when the subject reported one or more of the outcomes (regardless of how many were reported). As such, it is not the actual rate of outcomes that were reported, but rather the proportion of months that are positive (i.e., have at least one of the outcomes indicated). Odds ratios (95% CIs) are reported in months with extended-duration work shifts as compared to months without extended-duration work shifts.
doi:10.1371/journal.pmed.0030487.t002

Increases were also found for attentional failures during patient examinations (odds ratio = 1.5 [95% CI, 1.3–1.7] for between one and four extended-duration shifts and 2.1 [95% CI, 2.0–2.2] for five or more extended-duration shifts) compared to the reference group (no extended-duration shifts).

For educational activities (i.e., rounds with attending physicians or lectures), the results were more marked. The odds ratios of reporting at least one episode of falling asleep during rounds with attending physicians were significantly increased in months with between one and four extended-duration shifts (2.3; 95% CI, 2.2–2.4) and with five or more extended-duration shifts (5.5; 95% CI, 5.4–5.7). Odds ratios of falling asleep during lectures were similarly increased in months with between one and four extended-duration shifts (1.99; 95% CI, 1.96–2.03) and in months with five or more extended-duration shifts (4.3; 95% CI, 4.3–4.4), with 70% of intern-months with five or more extended-duration shifts having reports of falling asleep at least once during lectures. Interestingly, interns reported falling asleep during lectures at least once in 46% of months with no extended-duration work shifts.

Sleep

Interns reported sleeping 179.3 ± 54.3 h (mean ± standard deviation) in months with no extended-duration work shifts, 176.6 ± 45.1 h in months with between one and four extended-duration work shifts, and 165.3 ± 39.2 h in months with five or more extended-duration work shifts.

Discussion

We found that extended-duration work shifts were associated with adverse effects on patient safety (i.e., fatigue-related significant medical errors and preventable adverse events, including those resulting in fatalities). The risk of self-reported attentional failures during patient-care and educational activities was also significantly affected by extended-duration work shifts. That is, during months with frequent extended-duration work shifts (i.e., five or more extended-duration shifts in the month), interns were significantly more likely to fall asleep during surgery, while talking to or examining patients, during rounds, and during lectures or seminars, potentially affecting their ability to deliver patient care or to learn.

The hours of reported sleep per month decreased as the number of extended-duration work shifts, the number of reported medical errors, and the number of reported attentional failures increased—and this is consistent with laboratory studies that have unequivocally demonstrated the profound effects of sleep deprivation on alertness and performance [9–15]. Indeed, the adverse effects after 24 h of continuous wakefulness on neurobehavioral performance are comparable to that of a blood alcohol concentration of 0.10% [16,17]. Our results confirm the recently published findings by Landrigan et al. [3] and by Lockley et al. [2]. During a traditional schedule of extended-duration work shifts every other shift, as compared to a timetable where the scheduled length of work shifts was limited to less than 16 h, Landrigan et al. reported that serious medical errors were 36% more frequent [3], while Lockley et al. reported that polysomnographically defined attentional failures were ap-



proximately twice as frequent [2]. Moreover, our findings extend these results by confirming the associations between extended-duration work shifts, medical errors, preventable adverse events (including those resulting in fatalities), and attentional failures in interns in multiple specialties and hospitals across the United States.

In addition to the impact of these fatigue-related serious medical errors on patient safety, the perception of having made a medical error that causes an adverse patient outcome creates significant emotional distress for physicians that can last days or years and may include feelings of fear, guilt, anger, embarrassment, and humiliation [18]. The personal distress and reduced empathy associated with perceived medical errors increase the odds of future medical errors [19]. Without coping mechanisms, such as accepting responsibility, discussion with colleagues, and disclosure to patients, physicians may use dysfunctional methods of dealing with errors, such as alcohol and drug abuse [20,21].

We also demonstrated that extended-duration work shifts adversely affected medical education, as the odds of falling asleep in lectures and while on rounds with attending physicians increased significantly when extended-duration shifts were worked. Not only is adequate sleep crucial for maintaining vigilance [22], but sleep also plays an important role in memory consolidation and learning [23,24]. The amount of sleep reported by most interns is less than that shown to be necessary to perform satisfactorily on cognitive performance tasks that are implicitly necessary for learning [25]. Additionally, the high level of chronic partial sleep deprivation, consistent with prior reports [26,27], may account for the high base rate of attentional failures in these participants.

We acknowledge that, despite the large amount of data on intern-months collected from interns across specialties from programs throughout the United States, there are a number of limitations to our study. First, the individuals who volunteered for our study represent a small proportion of interns in the United States. Therefore, it is possible that the participants in our study may not be representative of the entire population of interns. For instance, our sample could be biased towards interns with a specific interest in the issue of resident work hours, thus raising the potential of a reporting bias. Additionally, those individuals who responded to our survey may have been more responsible and therefore more likely to report medical errors (although the responsibility of the participant could not account for the relationship between the numbers of extended-duration shifts and reported medical errors that were identified by the within-person analysis). However, potential participants were not informed of the specific study hypotheses, and the questions addressing our primary exposure and outcome variables were distributed among questions regarding secondary outcome measures such as caffeine usage, mood, and stress. Moreover, because each intern serves as his/her own control in our case-crossover analysis, it would have taken considerable time over a number of months—and a conscious effort to mislead—for intentionally biased respondents to affect our results deliberately. However, it remains a possibility that those who were more attentive to the topic of medical errors may have volunteered to participate in the study.

Second, the data for medical errors relied on self-reports and were not independently validated. Additionally, because

“fatigue or sleep deprivation” was part of the question regarding medical-error outcomes, it may have inflated the association between this suggested “cause” and the medical error. Third, the definition of what incidents interns believe constituted a “significant error” or “adverse patient outcome” may have varied from person to person. However, the consistency of the results when the outcome of the error resulted in a fatality provides a common metric of severity. In addition, because each intern was compared only to him- or herself, there is no reason to believe that he or she would have changed the criteria for what was significant by the nature of the rotation schedule.

Fourth, in our case-crossover analysis, it is possible that during months without extended-duration work shifts, interns may be on rotations with fewer opportunities to make errors (e.g., radiology rotation, dermatology, ambulatory elective), than on more clinically-oriented rotations when extended-duration work-shift frequency was greater (e.g., intensive-care unit, hospital ward) resulting in more opportunities to make errors. Additionally, there may be other additional organizational features (e.g., staffing, supervision) that differ between rotations. To address partially this issue of monthly activity, we conducted a secondary analysis in which we limited the data to only those months on hospital wards. When we did so, we found that the association between the frequency of extended-duration work shifts and medical errors and adverse events was comparable to the results of the overall analysis. While the number of hours at work increases as the number of shifts increases and may partially account for the increased rate of events, it should be noted that the percentage increase in weekly work hours (a 16.7% increase in the category of between one and four extended-duration shifts per month and a 42.9% increase in the number of five or more extended-duration shifts per month, compared to months with no extended-duration shifts) cannot account for the observed increase in the rate of events. For months with between one and four extended-duration shifts and for months with five or more extended-duration shifts, fatigue-related adverse events increased by ~450% and ~700%, respectively. For months with five or more extended-duration shifts, fatigue-related adverse events resulting in a fatality increased by ~300%.

We also recorded the frequency of both fatigue-related and non-fatigue-related errors. We found that even though the rate of non-fatigue-related errors increased as extended-duration work-shift frequency increased (presumably for the two reasons outlined above), the magnitude of the increase was much less than the increased rate of fatigue-related errors. This finding suggests that the increased rate of fatigue-related errors was not predominately due to a confounding effect of monthly activity. However, it remains possible that interns were more tired when working more frequent extended-duration shifts and that they may have been more likely to under-report or to over-report medical errors, misperceive errors in their work, or to attribute errors to fatigue. Landrigan and colleagues found that physicians perceived fewer medical errors than those assessed by an independent observer [3].

Fifth, for many of our analyses, we collected information about the events (i.e., errors) and exposures (i.e., number of extended-duration shifts worked per month) simultaneously from the monthly surveys. This could lead to reporting bias as

interns may be more inclined to send back surveys during months in which incidents occurred. However, when analyses were limited to participants returning all surveys throughout the year, the relationships between errors and the frequency of extended-duration shifts did not change substantially. Sixth, there may be a carry-over effect from one month to the next. For example, an intern may have worked five extended-duration shifts in the first month and no extended-duration shifts in the second month. If that intern reported a medical error at the beginning of the second month, our data analysis would not have been able to detect the potential carry-over effect from the high number of extended-duration shifts in the first month.

Finally, although self-reporting is a well-established method of eliciting information regarding medical errors, prior work has demonstrated that even very robust stimulated self-reporting systems detect only a fraction of all medical errors [28]. Consequently, it is likely that the interns committed a greater number of serious medical errors, including those resulting in preventable adverse events, than we have reported here. Thus, although there are a number of potential biases and confounders in our study, given the consistency and magnitude of our findings, it is unlikely that they could account for the striking association between extended-duration work shifts and fatigue-related significant medical errors, preventable adverse events, and attentional failures.

According to the 1999 report from the Institute of Medicine, between 48,000 and 98,000 deaths each year occur due to a medical error [1]. Recently, the ACGME has placed limitations on resident work hours in an attempt to reduce fatigue-related medical errors. Historically, when such attempts have been made to reduce resident work hours in the United States, the frequency of extended-duration work shifts (>24 h) has been reduced (e.g., from a frequency of Q2 to Q3 to Q4 [where Q2 is an extended duration shift occurring every other night, Q3 is an extended duration shift occurring every third night, and Q4 is an extended duration shift occurring every fourth night]). However, the practice of working for more than 24 h consecutively has remained the cornerstone of American postgraduate medical education. In fact, the recent (2003) ACGME work-hour guidelines for postgraduate medical education programs effectively continue to sanction up to nine extended-duration shifts (of up to 30 h consecutively) per month, since every other shift can be an extended-duration work shift under the new ACGME guidelines [29]. Still, interns working extended-duration shifts within these ACGME guidelines reported significant numbers of medical errors, including those that resulted in adverse patient outcomes and fatalities. Furthermore, 83.6% of interns reported working more hours than allowed by ACGME standards in the year following their introduction [30].

These data, collected from interns in all specialties across the United States, are not consistent with the recent suggestions by a member of the ACGME Residency Review Committee in Surgery [31] that safety hazards associated with resident fatigue are limited to a small subset of trainees. Even interns who worked well below the current 80-h ACGME weekly work-hour limits (averaging 64.8 h of work per week), but who continued to work up to one extended-duration shift per week (half the weekly frequency allowed under current ACGME standards), had 8-fold greater odds of reporting an

adverse event than those who did not work extended-duration work shifts. This finding is consistent with data from numerous studies documenting that 24 h consecutively of wakefulness impairs short-term memory, degrades neuro-behavioral performance, and greatly increases the risk of both errors of commission and omission and attentional failures [23,32,33]. Additionally, Ayas and colleagues recently reported that the odds of an intern having a percutaneous injury increased by 61% after ≥ 20 h at work [34]. These findings are also consistent with the recent demonstration that elimination of extended-duration work shifts reduces attentional failures and serious medical errors among interns working in intensive-care units [2,3].

Our results thus reveal that the practice of scheduling 24-h or greater extended-duration work shifts, as currently sanctioned by the ACGME, may pose a significant increased risk of safety hazards to patients, contribute to the occurrence of medical errors that are attributable to fatigue or sleep deprivation and to consequent preventable fatal and nonfatal adverse events, and may also interfere with the primary educational purpose of residency training. These results have important public policy implications in terms of postgraduate medical education and suggest that directors of training programs should consider alternative coverage schedules for trainees with the objective of eliminating extended-duration shifts. In Europe, where the tradition of extended-duration “on call” shifts originated more than a century ago, work shifts of all physicians (including those in training) have recently been limited to 13 h consecutively [35], thereby eliminating extended-duration work shifts altogether. Fletcher et al. recently published a review of interventions aimed at reducing US resident work hours, including strategies such as day and night float teams and the use of physician extenders [36]. Future studies should explore the applicability of our findings regarding the association between medical errors and extended-duration work shifts to all practicing physicians in the United States.

Supporting Information

Alternative Language Abstract S1. Translation of the Abstract into Hungarian by S. Kantor

Found at doi:10.1371/journal.pmed.0030487.sd001 (102 KB PDF).

Alternative Language Abstract S2. Translation of the Abstract into Polish by L. Kubin

Found at doi:10.1371/journal.pmed.0030487.sd002 (27 KB PDF).

Alternative Language Abstract S3. Translation of the Abstract into Portuguese by F. Louzada

Found at doi:10.1371/journal.pmed.0030487.sd003 (22 KB PDF).

Alternative Language Abstract S4. Translation of the Abstract into French by C. Gronfier

Found at doi:10.1371/journal.pmed.0030487.sd004 (32 KB PDF).

Alternative Language Abstract S5. Translation of the Abstract into Chinese by L. Ling

Found at doi:10.1371/journal.pmed.0030487.sd005 (188 KB PDF).

Alternative Language Abstract S6. Translation of the Abstract into Spanish by C. Robles

Found at doi:10.1371/journal.pmed.0030487.sd006 (28 KB PDF).

Alternative Language Abstract S7. Translation of the Abstract into Japanese by T. Tanigawa

Found at doi:10.1371/journal.pmed.0030487.sd007 (16.5 KB PDF).

Acknowledgments

The authors would especially like to thank the interns who took time from their busy work schedules to participate in this study. We would particularly like to thank the NRMP and the Association of American Medical Colleges, especially Jordan J. Cohen, MD, Paul Jolly, PhD, and the Division of Medical School Services and Studies, for their assistance with recruitment; DeWitt C. Baldwin, MD and Steven R. Daugherty, PhD for their help in designing the questionnaires; Tim Ayas, MD and Sharlene Hudson, MD for questionnaire review; Michael Schulzer, MD, PhD for his assistance with data analysis; Steven W. Lockley, PhD for his assistance with the study design; Christopher P. Landrigan, MD for his assistance with data interpretation; and Joseph B. Martin, PhD, MD for his support and encouragement of this work. We are also appreciative of Cheryl W. Were and Mohammed Rasheed at Pearson Assessments for their commitment to this project.

Author contributions. LKB, NTA, BEC, FES, and CAC designed the study and survey instruments. LKB and JWC enrolled participants. BEC provided technical design of study database and data management. LKB, NTA, BEC, and BR analyzed the data. LKB, NTA, BEC, JWC, BR, FES, and CAC contributed to writing the paper.

Funding: This study was supported by grants from the National Institute for Occupational Safety and Health within the US Centers for Disease Control (Grant R01 OH07567) and the Agency for Healthcare Research and Quality (AHRQ) (Grant R01 HS12032). JWC was the recipient of an AHRQ National Research Service Award (F32 HS14130), and JWC and LKB were the recipients of National Heart, Lung and Blood Institute fellowships (T32 HL079010) in the program of training in Sleep, Circadian and Respiratory Neurobiology at Brigham and Women's Hospital. NTA is supported by a New Investigator Award from the Canadian Institutes of Health Research and British Columbia Lung Association, a Michael Smith Foundation Scholar Award, and a Departmental Scholar Award from the University of British Columbia. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: BEC is a past consultant for Vanda Pharmaceuticals. CAC, LKB, BR, and FES are paid employees of the Brigham and Women's Hospital, a subsidiary of Partners HealthCare System which employs thousands of interns and residents. CAC has received consulting fees from, or has served as a paid member of, scientific advisory boards for Accelerator, Actelion, the American Physiological Society, Aventis, Avera Pharmaceuticals, Cephalon, Coca-Cola, Hypnion, the NASA Jet Propulsion Laboratory (managed by the California Institute of Technology), the National Center for Sleep Disorders Research (which is part of the National Heart, Lung and Blood Institute), Oxford Biosignals, Pfizer, Morgan Stanley, Sleep Multimedia, the Sleep Research Society (for which CAC has served as president), Respiroics, Takeda Pharmaceuticals, Unilever, Vanda Pharmaceuticals, and Warburg-Pincus. CAC owns an equity interest in Hypnion and Vanda Pharmaceuticals. CAC has received lecture fees from the Accreditation Council of Graduate Medical Education (which established the current work-hour standards for interns), the American Academy of Allergy, Asthma and Immunology Program Directors, the Association of University Anesthesiologists, Beth-Israel Deaconess Medical Center, Brown Medical School and affiliated Rhode Island Hospital, Cephalon, Dalhousie University, Harvard University, London Deanery, MPM Capital, Partners HealthCare, Philips Lighting, Sanofi-Aventis, the Smithsonian Institution, Takeda, the Society of Neurological Surgeons, the University of Michigan, the University of Pennsylvania, the University of Pittsburgh, the University of Virginia Medical School, and the University of Wisconsin Medical School. CAC holds a number of process patents in the field of sleep/circadian rhythms (e.g., photic resetting of the human circadian pacemaker), all of which are assigned to the Brigham and Women's Hospital per institutional policy. CAC has also received research prizes with monetary awards from the American Academy of Sleep Medicine, the American Clinical and Climatological Association, the Association for Patient-Oriented Research, the National Institute for Occupational Safety and Health, and the National Sleep Foundation, clinical trial research contracts from Cephalon and Pfizer, an investigator-initiated research grant from Cephalon, and unrestricted research and education funds from Cephalon, Philips Lighting, and the Brigham and Women's Hospital. CAC is the incumbent of an endowed professorship provided to Harvard University by Cephalon. Since 1985, CAC has served as an expert witness on various legal cases related to sleep and/or circadian rhythms, but has never served as an expert witness for a commercial research sponsor or on a matter related to interns or residents.

References

- Kohn LT, Corrigan JM, Donaldson MS, editors, Committee on Quality of Healthcare in America, Institute of Medicine (1999) To err is human: Building a safer health system. Washington (DC): National Academy Press. 312 p.
- Lockley SW, Cronin JW, Evans EE, Cade BE, Lee CJ, et al. (2004) Effect of reducing interns' weekly work hours on sleep and attentional failures. *N Engl J Med* 351: 1829–1837.
- Landrigan CP, Rothschild JM, Cronin JW, Kaushal R, Burdick E, et al. (2004) Effect of reducing interns' work hours on serious medical errors in intensive-care units. *N Engl J Med* 351: 1838–1848.
- Barger LK, Cade BE, Ayas NT, Cronin JW, Rosner B, et al. (2005) Extended work shifts and the risk of motor vehicle crashes among interns. *N Engl J Med* 352: 125–134.
- Redelmeier DA, Tibshirani RJ (1997) Interpretation and bias in case-crossover studies. *J Clin Epidemiol* 50: 1281–1287.
- Maclure M (1991) The case-crossover design: A method of studying transient effects on the risk of acute events. *Am J Epidemiol* 133: 144–153.
- Rosner B (2000) Fundamentals of biostatistics. Pacific Grove (California): Duxbury Press. 792 p.
- National Resident Matching Program (2006). Available: <http://www.nrmp.org>. Accessed: 30 October 2006.
- Dinges DF (1989) The nature of sleepiness: Causes, contexts, and consequences. In: Stunkard AJ, Baum A, editors. Perspectives in behavioral medicine: Eating, sleeping, and sex. Hillsdale, New Jersey: Lawrence Erlbaum Associates. pp. 147–179.
- Carskadon MA, Dement WC (1992) Multiple sleep latency tests during the constant routine. *Sleep* 15: 396–399.
- Pilcher JJ, Huffcutt AI (1996) Effects of sleep deprivation on performance: A meta-analysis. *Sleep* 19: 318–326.
- Lorenzo I, Ramos J, Arce C, Guevara MA, Corsi-Cabrera M (1995) Effect of total sleep deprivation on reaction time and waking EEG activity in man. *Sleep* 18: 346–354.
- Koslowsky M, Babkoff H (1992) Meta-analysis of the relationship between total sleep deprivation and performance. *Chronobiol Int* 9: 132–136.
- Johnson LC (1982) Sleep deprivation and performance. In: Webb WB, editor. Biological rhythms, sleep, and performance. New York: John Wiley. pp. 111–141.
- Wilkinson RT (1972) Effects of up to 60 hours' sleep deprivation on different types of work. *Ergonomics* 7: 175–186.
- Dawson D, Reid K (1997) Fatigue, alcohol and performance impairment. *Nature* 388: 235.
- Arnedt JT, Owens J, Crouch M, Stahl J, Carskadon MA (2005) Neuro-behavioral performance of residents after heavy night call vs after alcohol ingestion. *JAMA* 294: 1025–1033.
- Christensen JF, Levinson W, Dunn PM (1992) The heart of darkness: The impact of perceived mistakes on physicians. *J Gen Intern Med* 7: 424–431.
- West CP, Huschka MM, Novotny PJ, Sloan JA, Kolars JC, et al. (2006) Association of perceived medical errors with resident distress and empathy. A prospective longitudinal study. *JAMA* 296: 1071–1078.
- Goldberg RM, Kuhn G, Andrew LB, Thomas HA (2004) Coping with medical mistakes and errors in judgment. *Ann Emerg Med* 39: 287–292.
- Newman MC (1996) The emotional impact of mistakes on family physicians. *Arch Fam Med* 5: 71–75.
- van Dongen HPA, Maislin G, Mullington JM, Dinges DF (2003) The cumulative cost of additional wakefulness: Dose-response effects on neurobehavioral functions and sleep physiology from chronic sleep restriction and total sleep deprivation. *Sleep* 26: 117–126.
- Stickgold R, James L, Hobson JA (2000) Visual discrimination learning requires sleep after training. *Nat Neurosci* 3: 1237–1238.
- Walker MP, Brakefield T, Morgan A, Hobson JA, Stickgold R (2002) Practice with sleep makes perfect: Sleep-dependent motor skill learning. *Neuron* 35: 205–211.
- Baldwin DC Jr, Daugherty SR (2004) Sleep deprivation and fatigue in residency training: Results of a national survey of first- and second-year residents. *Sleep* 27: 217–223.
- Howard SK, Gaba DM, Rosekind MR, Zarcone VP (2002) The risks and implications of excessive daytime sleepiness in resident physicians. *Acad Med* 77: 1019–1025.
- Richardson GS, Wyatt JK, Sullivan JP, Orav E, Ward A, et al. (1996) Objective assessment of sleep and alertness in medical housestaff and the impact of protected time for sleep. *Sleep* 19: 718–726.
- O'Neil AC, Petersen LA, Cook EF, Bates DW, Lee TH, et al. (1993) Physician reporting compared with medical-record review to identify adverse medical events. *Ann Intern Med* 119: 370–376.
- Association of American Medical Colleges (October 2001) AAMC policy guidance on graduate medical education. Assuring quality patient care and quality education. Available: <http://www.aamc.org/patientcare/gmepolicy/gmepolicy.pdf>. Accessed 30 October 2006.
- Landrigan CP, Barger LK, Cade BE, Ayas NT, Czeisler CA (2006) Interns' compliance with Accreditation Council for Graduate Medical Education work-hour limits. *JAMA* 296: 1063–1070.
- Fischer JE (2005) Surgeons: Employees or professionals? *Am J Surg* 190: 1–3.

32. Cajochen C, Khalsa SBS, Wyatt JK, Czeisler CA, Dijk DJ (1999) EEG and ocular correlates of circadian melatonin phase and human performance decrements during sleep loss. *Am J Physiol* 277: R640–R649.
33. Horowitz TS, Cade BE, Wolfe JM, Czeisler CA (2003) Searching night and day: A dissociation of effects of circadian phase and time awake on visual selective attention and vigilance. *Psychol Sci* 14: 549–557.
34. Ayas NT, Barger LK, Cade BE, Hashimoto DM, Rosner B, et al. (2006) Extended work duration and the risk of self-reported percutaneous injuries in interns. *JAMA* 296: 1055–1062.
35. Incomes Data Services (2006). European Working Time Directive. Available: <http://www.incomesdata.co.uk/information/worktimedirective.htm#Article3>. Accessed 30 October 2006.
36. Fletcher KE, Underwood W III, Davis SQ, Mangrulkar RS, McMahon LF Jr, et al. (2005) Effects of work hour reduction on residents' lives: A systematic review. *JAMA* 294: 1088–1100.

Editors' Summary

Background. In the United States, medical students who are doing their internship (first year of postgraduate clinical training) regularly work in the clinic for longer than 24 h at a time. It is already known that doctors or students who work for long shifts make more medical errors and are less able to pay attention to what they are doing. Many thousands of adverse medical events per year including, in the extreme, deaths of patients, are thought to result from medical errors, but it is not clear whether doctors or students working long shifts—as opposed to, for example, an increase in total number of hours worked—are the cause of many or any of these errors.

Why Was This Study Done? This research group wanted to find out whether long shifts worked by interns had an effect on reported medical errors, and hence patient safety, and specifically whether any harm that happened to patients might otherwise have been preventable.

What Did the Researchers Do and Find? The researchers contacted all US medical school graduates beginning their internships from one particular year-group by email, and asked each person whether they wanted to take part in a confidential survey. Individuals who agreed to participate were directed to a secure website to enter basic information about themselves and then to complete a form each month. On that form the interns gave information about their working hours, hours of sleep, and number of extended-duration shifts worked, and completed questions about medical errors in the past month. Then, for each intern in the study, researchers compared month by month the number of medical errors and the number of extended-duration shifts that had been worked. A total of 2,737 interns took part in the survey.

Compared to months in which no extended-duration shifts were worked, in those months in which between one and four, and more than five extended-duration shifts were worked, the doctors were, respectively, three and seven times more likely to report at least one fatigue-related significant medical error. Similarly, fatigue-related adverse events increased by around seven and eight times, respectively, compared with months in which no extended-duration shifts were worked. Fatigue-related preventable adverse events associated with the death of the patient increased by ~300% in interns working more than five extended-duration shifts per month; they were also more likely to fall asleep during lectures, rounds, and clinical activities, including surgery.

What Do These Findings Mean? Guidelines for graduate medical education in the United States still allow up to nine marathon shifts (30 h at a stretch) per month, even though the total number of hours worked is capped. This study shows that the long shifts worked by interns are bad for patient safety, as they are more likely to cause harm that would not otherwise happen.

Additional Information. Please access these Web sites via the online version of this summary at <http://dx.doi.org/doi:10.1371/journal.pmed.0030487>.

- The US Food and Drug Administration has resources on its website about medication errors
- The National Sleep Foundation aims to improve public health and safety by achieving understanding of sleep and sleep disorders
- Wikipedia (an internet encyclopedia anyone can edit) has a page about residency training in the United States