

Northumbria Research Link

Citation: Ritti-Dias, Raphael Mendes, Correia, Marilia Almeida, Carvalho, Juliana Ferreira, Braghieri, Heloisa Amaral, Wolosker, Nelson, Cucato, Gabriel and Kanegusuku, Hécio (2022) Impact of the COVID-19 pandemic on health lifestyle in patients with peripheral artery disease: A cross-sectional study. *Journal of Vascular Nursing*, 40 (1). pp. 54-58. ISSN 1062-0303

Published by: Elsevier

URL: <https://doi.org/10.1016/j.jvn.2022.01.001>
<<https://doi.org/10.1016/j.jvn.2022.01.001>>

This version was downloaded from Northumbria Research Link:
<https://nrl.northumbria.ac.uk/id/eprint/48656/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)

Journal of Vascular Nursing

Impact of the COVID-19 Pandemic on Health Lifestyle in Patients with Peripheral Artery Disease: A Cross-sectional Study

--Manuscript Draft--

Manuscript Number:	JVN-D-20-00067R7
Article Type:	Research Paper
Section/Category:	Regular Submission
Keywords:	Coronavirus 2019, social isolation, peripheral artery disease, physical activity, mental health.
Corresponding Author:	Hélcio kanegusuku Hospital Israelita Albert Einstein BRAZIL
First Author:	Raphael Mendes Ritti-Dias, PhD
Order of Authors:	Raphael Mendes Ritti-Dias, PhD Marília Almeida Correia, PhD Juliana Ferreira Carvalho, BSc Heloisa Amaral Braghieri, MSc Nelson Wolosker, PhD Gabriel Grizzo Cucato, PhD Hélcio kanegusuku, PhD
Response to Reviewers:	<p>Reviewer:</p> <p>MATERIAL AND METHODS Please indicate year. Answer: As suggested, we added this information: "...on March 16, 2020 with quarantine, and non-essential services closed on March 24, 2020. Data collection was performed through a phone interview, between May 15, 2020 and August 22, 2020...".</p> <p>Understanding there is expertise in research with PAD there remains a concern for the validity of the tool used. Suggest commenting on the potential bias and lack of rigor and validity secondary to a non validated questionnaire Answer: We completely agree with the reviewer. Thus, as suggested, we rewrote the sentence: "...The questionnaire used in this study was a non validated questionnaire developed by our group, based on questionnaires and previous studies. 11-15 Despite the group has experience in research with PAD, the use of non validated questionnaire has less methodological rigor and is also more susceptible to information bias...".</p>

Reviewer:

MATERIAL AND METHODS

Please indicate year.

Answer: As suggested, we added this information: "...on March 16, 2020 with quarantine, and non-essential services closed on March 24, 2020. Data collection was performed through a phone interview, between May 15, 2020 and August 22, 2020...".

Understanding there is expertise in research with PAD there remains a concern for the validity of the tool used. Suggest commenting on the potential bias and lack of rigor and validity secondary to a non validated questionnaire

Answer: We completely agree with the reviewer. Thus, as suggested, we rewrote the sentence: "...The questionnaire used in this study was a non validated questionnaire developed by our group, based on questionnaires and previous studies.¹¹⁻¹⁵ Despite the group has experience in research with PAD, the use of non validated questionnaire has less methodological rigor and is also more susceptible to information bias...".

The majority of patients self-reported spending more time watching TV and sitting.

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12
- 13
- 14
- 15
- 16
- 17
- 18
- 19
- 20
- 21
- 22
- 23
- 24
- 25
- 26
- 27
- 28
- 29
- 30
- 31
- 32
- 33
- 34
- 35
- 36
- 37
- 38
- 39
- 40
- 41
- 42
- 43
- 44
- 45
- 46
- 47
- 48
- 49
- 50
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60
- 61
- 62
- 63
- 64
- 65

Only 28.7% of patients self-reported practicing physical exercise.

Most patients self-reported worse physical and mental health.

Original Article

Impact of the COVID-19 Pandemic on Health Lifestyle in Patients with Peripheral Artery Disease: A Cross-sectional Study

Running head: COVID-19 and Peripheral Artery Disease

Authors: Raphael Mendes Ritti-Dias¹, Marilia de Almeida Correia², Juliana Ferreira de Carvalho³, Heloisa Amaral Braghieri¹, Nelson Wolosker³, Gabriel Grizzo Cucato⁴, Hécio Kanegusuku³

Affiliations: ¹Graduated Program in Rehabilitation Sciences, Universidade Nove de Julho, São Paulo, Brazil;

² Graduated Program in Medicine, Universidade Nove de Julho, São Paulo, Brazil;

³Hospital Israelita Albert Einstein, São Paulo, Brazil;

⁴Department of Sport, Exercise & Rehabilitation, Northumbria University, Newcastle Upon Tyne, UK.

***Corresponding Author:**

Raphael Mendes Ritti-Dias, PhD

Universidade Nove de Julho, Rua Vergueiro 235, São Paulo – SP – Brazil. ZIP-code:

01504-000. Phone: (+5519) 99940 6878; e-mail: raphaelritti@gmail.com

Funding: RR and NW receive a research productivity fellowship granted by Brazilian National Council for Scientific and Technological Development.

Conflict of interest: The authors have no conflict of interest.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Original Article

**Impact of the COVID-19 pandemic on health lifestyle in patients with peripheral artery dis-
ease**

Running head: COVID-19 and Peripheral Artery Disease

Conflict of interest

The authors have no conflict of interest.

1
2
3
4 **ABSTRACT**
5

6 **Background:** Social isolation has been one of the main strategies to prevent the spread of Coro-
7 navirus 2019 (COVID-19). However, the impact of social isolation on the lifestyle of patients with
8 peripheral artery disease (PAD) and claudication symptoms remains unclear.
9

10 Objectives: To analyze the perceptions of patients with PAD of the impact of social isolation pro-
11 voked by COVID-19 pandemic on health lifestyle.
12
13

14 **Design:** Cross-sectional.
15

16 **Setting:** The database of studies developed by our group involving patients with PAD from public
17 hospitals in São Paulo.
18

19 **Methods:** In this cross-sectional survey study, 136 patients with PAD (61% men, 68±9 years old,
20 0.55±0.17 ankle-brachial index, 82.4% with a PAD diagnosis ≥5 years old) were included. Health
21 lifestyle factors were assessed through a telephone interview using a questionnaire containing
22 questions related to: (a) COVID-19 personal care; (b) mental health; (c) health risk habits; (d)
23 eating behavior; (e) lifestyle; (f) physical activity; (g) overall health; and (h) peripheral artery dis-
24 ease health care.
25

26 **Results:** The majority of patients self-reported spending more time watching TV and sitting during
27 the COVID-19 pandemic and only 28.7% were practicing physical exercise. Anxiety and unhap-
28 piness were the most prevalent feelings self-reported among patients and 43.4% reported a decline
29 in walking capacity.
30

31 **Conclusion:** Most patients with PAD self-reported increased sedentary behavior, lower physical
32 activity level, and worse physical and mental health during the COVID-19 pandemic. Thus, it is
33 necessary to adopt strategies to improve the quality of life of these patients during this period.
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 **Keywords:** Coronavirus 2019, social isolation, peripheral artery disease, physical activity, mental
5
6 health.
7
8
9

10 11 **INTRODUCTION**

12
13
14 Patients with peripheral artery disease (PAD) and claudication symptoms present with
15 walking impairment ¹ and several cardiovascular risk factors, such as smoking, hypertension, and
16 dyslipidemia. ¹⁻⁵ The first line therapy for these patients includes stimulus to physical activity prac-
17 tice, improvements in healthy eating, the treatment of comorbid conditions, such as. hypertension,
18 cardiac disease and more recently, management of mental health. ^{6,7}
19
20
21
22
23
24
25

26 In December 2019, a novel severe acute respiratory syndrome coronavirus (SARS-Cov-2)
27 called Coronavirus 2019 (COVID-19) was first identified in the city of Wuhan, China and subse-
28 quently declared a global pandemic, with more than 200.174.880 cases and 4.255.890 deaths reg-
29 istered worldwide by August 5, 2021. ⁸ Social isolation has been one of the main strategies used
30 to prevent the spread of COVID-19, however, this strategy has resulted in unintended negative
31 consequences on the lifestyle of the population. ⁹⁻¹⁰ Amnar et al. (2020), through an international
32 online survey involving people from different continents, observed a reduction in physical activity
33 level, an increase in daily sitting time, and unhealthy eating habits during the COVID-19 pan-
34 demic. Hossain et al. (2020) in a review study also observed worsening mental health of the pop-
35 ulation due to social isolation. Whether similar results occur in patients with PAD, who frequently
36 report low physical activity level, poor eating habits, and mental health, is unclear. Thus, the cur-
37 rent study investigated the perceptions of patients with PAD and the impact of COVID-19 on
38 health lifestyle. The hypothesis of the study was that social isolation negatively affects lifestyle,
39 aggravating the physical and mental health, and unhealthy eating habits of these patients.
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

2. MATERIAL AND METHODS

Study design and patients

This observational, descriptive, cross-sectional survey study, included patients with PAD and claudication symptoms (i.e. residents of metropolitan cities) recruited from the database of studies developed and published before the COVID-19 pandemic by our group.^{1,3-5,11-13} This study was approved by the Universidade Nove de Julho' Ethics Committee before data collection (CAAE #31529220.8.0000.5511). All participants verbally gave informed consent by phone prior to participation. Participants did not identify themselves and their answers were only included in the sample if they gave authorization before the protocol started. All procedures follow the national legislation and Declaration of Helsinki.

Patients were included if they met the following criteria: a) agreed to participate and respond to all questions of the survey; b) previous diagnosis of PAD; c) age > 45 years old; d) had ankle-brachial index (ABI) ≤ 0.90 in one or both legs, and; e) absence of non-compressible vessels, amputated limbs and/or ulcers. Patients were only excluded if: a) presented disabilities such as cognitive, hearing, or speech during phone call that compromises the answer to the questionnaire.

Data collection

Social isolation in São Paulo was recommended on March 16, 2020 with quarantine, and non-essential services closed on March 24, 2020. Data collection was performed through a phone interview, between May 15, 2020 and August 22, 2020. The evaluation of the impact of COVID-19 on health lifestyle was assessed through a questionnaire developed by the researchers of the study, based on questionnaires and previous studies.¹¹⁻¹⁵

1
2
3
4 The questionnaire was composed of questions divided into 8 domains: (a) COVID-19 per-
5 sonal care; (b) mental health; (c) health risk habits; (d) eating behavior; (e) lifestyle; (f) physical
6 activity; (g) overall health; and (h) peripheral artery disease health care. The questions used in the
7 present analysis are presented below.
8
9

10
11
12
13
14 Personal information was accessed from our database including information on sex (“fe-
15 male” or “male”), date of birth (DD/MM/YYYY), time of PAD diagnosis (in years), and PAD
16 severity (ankle-brachial index).
17
18
19

20
21 Covid-19 personal care: involved questions on the recommendations for personal care dur-
22 ing the Covid-19 pandemic. The patient was required to report all that were applicable from the
23 possible answers: “washing hands”, “alcohol gel”, “avoid leaving home (shelter in place)”, “avoid
24 crowds”, “avoid other family members (outside household)”, “avoid shaking hands”, “avoid eating
25 out”, and “social isolation”. In order to gain direct information about a COVID-19 diagnosis, pa-
26 tients were asked: 1 – Have you had contact with someone who was diagnosed with COVID-19?,
27 2 – Have you been diagnosed with COVID-19? If yes, 3 – Have you recovered? Answers to all
28 questions were “Yes” or “No”.
29
30
31
32
33
34
35
36
37
38
39
40

41 Mental health: This domain is composed of 3-self-reported items aiming to identify fre-
42 quent feelings related to the COVID-19 pandemic. The following questions were used: 1 – Due to
43 the COVID-19, are you feeling more anxious?; 2 – Due to the COVID-19, are you feeling over-
44 whelmingly unhappy?; 3 - Due to the COVID-19, are you feeling more stressed?; 4- Due to the
45 COVID-19, are you feeling depressed? Answers to all questions were “Yes” or “No”.
46
47
48
49
50
51
52

53 Health risk behavior: This domain aimed to identify frequent social habits. The selected
54 questions were as follows: 1 – Do you smoke?; 2 – Due to social isolation, do you spend more
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 time sitting?; 3 – With the COVID-19 pandemic, do you feel that your television use has increased?

5
6
7 Answers to all items were “No”, or “Yes”.

8
9 Eating behavior: To explore the possible impacts of COVID-19 on the frequent eating hab-
10 its, the following questions were asked: 1 – Due to the COVID-19 pandemic, are your eating hab-
11 its: “unchanged”, “worsened” or “improved”?; 2 –Due to the COVID-19 pandemic, do you feel
12 that your fruit intake has increased?; 3 – Due to the COVID-19 pandemic, do you feel that your
13 intake of sweets has increased? 4 – Due to the COVID-19 pandemic, do you feel that your intake
14 of vegetables has increased? 5 – Due to the COVID-19 pandemic, do you feel that your fried food
15 intake has increased? Possible answers to questions 2 to 5 of this topic were “No”, or “Yes”.

16
17
18
19
20
21
22
23
24
25
26 Physical activity: In order to assess frequent physical activity habits, participants were
27 asked the following questions: 1-Did you regularly practice physical activity before the COVID-
28 19 pandemic?; 2- Have you been practicing physical activity regularly, at least once a week, during
29 the COVID-19 pandemic?; 3 – How many times do you exercise a week at the moment?; 4 –Do
30 you usually exercise less than 30 min, between 30 and 60 minutes or more than 60 minutes?; 5 –
31 What type of exercise do you do?

32
33
34
35
36
37
38
39
40
41 Overall health: From the list of diseases, the participant is required to report the presence
42 of all diagnosed diseases, such as hypertension, diabetes, high cholesterol, high triglycerides, car-
43 diopathy, or other.

44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
Peripheral artery disease health care: from this domain patients were asked: “Do you feel
that your ability to walk has decreased in recent weeks?” Possible answers were “yes” or “no”.

1
2
3
4 **Statistical analyses**
5

6 All statistical analysis were performed using the software SPSS (version 20). The data are
7 presented as mean and standard deviations for continuous variables and relative frequencies for
8 categorical variables.
9
10
11
12
13
14

15
16 **3. RESULTS**
17

18 A flowchart of the study is provided in Figure 1. One hundred and sixty-seven patients
19 were contacted. Of these, 18 patients died before the COVID-19 pandemic, 3 did not agree to
20 participate, and 10 did not meet the inclusion criteria. Thus, 136 patients with PAD and claudica-
21 tion symptoms (61% men, 68±9 years old, 0.55±0.17 ankle-brachial index, 82.4% had the PAD
22 diagnosis ≥5 years old) participated in the study. Of these, two patients had been diagnosed with
23 and recovered from COVID-19. The majority of patients followed preventive attitudes to avoid
24 COVID-19 contagion, such as social isolation (88.2%) and hand sanitization (98.5%) (Table 1).
25
26
27
28
29
30
31
32
33
34

35
36 ***Table 1***
37

38 Table 2 shows the sedentary and physical activity habits of patients with PAD during the
39 COVID-19 pandemic. Most patients self-reported spending more time watching TV and sitting,
40 58.8% and 76.5%, respectively. Before the COVID-19 pandemic, 54.4% of patients were physi-
41 cally active, decreasing to 28.7% during the COVID-19 pandemic.
42
43
44
45
46
47

48 ***Table 2***
49

50 Table 3 shows the eating habits, physical and mental health of patients with PAD during
51 the COVID-19 pandemic. Approximately, seventy-nine percent of patients had not modified their
52 eating habits and only 10.3% reported worsened eating habits. In addition, approximately, forty-
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 three percent of patients reported a reduction in walking capacity, anxiety, and unhappiness had
5
6 increased in more than 30.1% of patients.
7
8

9 ***Table 3***
10

11 12 13 14 **4. DISCUSSION**

15
16 This study revealed the majority of patients self report spending more time watching TV
17
18 and sitting during the COVID-19 pandemic, with only 28.7% physically exercising. In addition,
19
20 forty-three percent of patient reported the most prevalent self reported feeling of anxiety, unhap-
21
22 piness, and noted decline in walking capacity.
23
24

25
26 The majority of the patients presented with different co-morbidities, such as hypertension,
27
28 dyslipidemia, and diabetes. The study results corroborate with other studies ¹⁻⁵ that also demon-
29
30 strated a high prevalence of cardiovascular risk factors in patients with PAD, classifying them as
31
32 a risk group for COVID-19. ¹⁶ Jin et al (2020) state the number of co-morbidities has been associ-
33
34 ated with higher severity and mortality among patients with a COVID-19 diagnosis. Among our
35
36 sample, two patients had been diagnosed with and recovered from COVID-19. This result is inter-
37
38 esting because São Paulo had the highest number of cases and deaths of all cities in Brazil. ¹⁷ The
39
40 low number of cases of COVID-19 among our sample may be related to their high adherence to
41
42 preventive measures. Most of our patients were in social isolation (88.2%), avoiding contact with
43
44 other people (97.1%) or leaving their homes (83.8%), and using hygienic strategies, such as hand
45
46 sanitization (98.5%). These precautions suggest that preventive attitudes could be useful to avoid
47
48 contagion in this group.
49
50
51
52

53
54
55 In accordance with other studies performed with subjects without PAD, ^{18,19} patients self-
56
57 reported spending more time watching TV and sitting compared with the period before COVID-
58
59
60
61
62
63
64
65

1
2
3
4 19. This is alarming for these patients who already presented high sedentary behavior¹³ and also,
5
6 especially, among the patients with a higher body mass index and lower walking capacity.¹² A
7
8 small number of patients with PAD also self-reported (28.7%) the continuation of regular physical
9
10 exercise practice. Recent studies^{9,20,21} also observed a negative impact of the COVID-19 pandemic
11
12 on physical activity level of the population in different countries. Studies^{3,22} performed prior to
13
14 the COVID-19 pandemic highlighted that the majority of PAD patients were physically inactive.
15
16 This result corroborates the impairment in walking capacity reported by 43.4% of patients com-
17
18 pared with before COVID-19. Lower walking capacity has been associated with worse prognosis
19
20 in this population.^{23,24} Thus, exacerbation of these harmful behaviors due to social isolation has
21
22 aggravated the worsening in health of these patients.
23
24
25
26
27

28
29 Many patients reported being more anxious, unhappy, stressed, and depressed than before
30
31 the pandemic, which is particularly worrying since studies before the COVID-19 pandemic
32
33 showed that patients with PAD already presented deleterious alterations in mental health compared
34
35 to subjects without PAD.^{25,26} Factors such as social distancing from family and friends, for being
36
37 included in a main risk group, and also reduced physical activity level might have contributed to
38
39 these factors. Future studies should investigate the factors associated with mental health during
40
41 social isolation in patients with PAD.
42
43
44

45
46 Since physical activity, management of cardiovascular risk factors, and mental health are
47
48 the cornerstones of clinical treatment of PAD, these results could be useful for health authorities
49
50 to develop future strategies to avoid the clinical impairment caused by the COVID-19 pandemic
51
52 in PAD. In this context, our data support the importance of studies analyzing the feasibility and
53
54 health effects of behavior change programs using telemedicine consultations to improve overall
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 health (sedentary time, physical and mental health, healthy eating habits, and physical activity),
5
6 unsupervised home-based exercise programs, and tele-rehabilitation programs.
7
8
9

10 11 **5. LIMITATIONS** 12

13
14 The questionnaire used in this study was a non validated questionnaire developed by our
15
16 group, based on questionnaires and previous studies.¹¹⁻¹⁵ Despite the group has experience in re-
17
18 search with PAD, the use of non validated questionnaire has less methodological rigor and is also
19
20 more susceptible to information bias. The PAD severity was not assessed due to the social isolation
21
22 adopted to contain of COVID-19, however, as all patients were recruited from a previously com-
23
24 piled database. We can attest all of them have a confirmed PAD diagnosis. In addition, a group of
25
26 individuals without PAD was not included in this study.
27
28
29
30
31
32

33 34 **6. CONCLUSION** 35

36 Most patients with PAD self-reported increased sedentary behavior, lower physical activity
37
38 level, and worse physical and mental health during the COVID-19 pandemic. Behavior change
39
40 programs involving nutritional, psychological, physical activity monitoring among others in order
41
42 to improve quality of life these patients should be adopted to counteract the consequences of
43
44 COVID-19 pandemic on lifestyle.
45
46
47
48
49

50 51 **7. References** 52

- 53 1. Farah BQ, Souza Barbosa JP, Cucato GG, Chehuen Mda R, Gobbo LA, Wolosker N, et al.
54
55 Predictors of walking capacity in peripheral arterial disease patients. Clinics. 2013; 68: 537-41.
56
57
58
59
60
61
62
63
64
65

- 1
2
3
4 2. Brevetti G, Oliva G, Di Giacomo S, Bucur R, Annecchini R, Di Iorio A. Intermittent claudi-
5
6 cation in older patients: risk factors, cardiovascular comorbidity, and severity of peripheral arterial
7
8 disease. *J Am Geriatr Soc.* 2001; 49: 1261-2.
9
- 10
11 3. Gerage AM, Correia MA, Oliveira PML, Palmeira AC, Domingues WJR, Zeratti AE, et al.
12
13 Physical Activity Levels in Peripheral Artery Disease Patients. *Arq Bras Cardiol.* 2019; 113: 410-
14
15
16 6.
17
- 18
19 4. Cavalcante BR, Germano-Soares AH, Gerage AM, Leicht A, Tassitano RM, Bortolotti H, et
20
21 al. Association between physical activity and walking capacity with cognitive function in periph-
22
23 eral artery disease patients. *Eur J Vasc Endovasc Surg.* 2018; 55: 672-8.
24
- 25
26 5. Kanegusuku H, Cucato GG, Domiciano RM, Longano P, Puech-Leao P, Wolosker N, et al.
27
28 Impact of obesity on walking capacity and cardiovascular parameters in patients with peripheral
29
30 artery disease: A cross-sectional study. *J Vasc Nurs.* 2020; 38: 66-71.
31
- 32
33 6. Aboyans V, Ricco JB, Bartelink MEL, Bjorck M, Brodmann M, Cohnert T, et al. 2017 ESC
34
35 Guidelines on the Diagnosis and Treatment of Peripheral Arterial Diseases, in collaboration with
36
37 the European Society for Vascular Surgery (ESVS): Document covering atherosclerotic disease of
38
39 extracranial carotid and vertebral, mesenteric, renal, upper and lower extremity arteries Endorsed
40
41 by: the European Stroke Organization (ESO). The Task Force for the Diagnosis and Treatment of
42
43 Peripheral Arterial Diseases of the European Society of Cardiology (ESC) and of the European
44
45 Society for Vascular Surgery (ESVS). *Eur Heart J.* 2018; 39: 763-816.
46
47
- 48
49 7. Norgren L, Hiatt WR, Dormandy JA, Nehler MR, Harris KA, Fowkes FG. Inter-Society Con-
50
51 sensus for the Management of Peripheral Arterial Disease (TASC II). *J Vasc Surg.* 2007;45 Suppl
52
53 S:S5-67.
54
55
56
57
58
59
60
61
62
63
64
65

- 1
2
3
4 8. World Health Organization. Coronavirus disease 2019 (COVID-19) pandemic.
5
6 <https://www.who.int/emergencies/diseases/novel-coronavirus-2019> (Accessed August 5, 2021).
7
8
- 9 9. Ammar A, Brach M, Trabelsi K, Chtourou H, Boukhris O, Masmoudi L, et al. Effects of
10 COVID-19 Home Confinement on Eating Behaviour and Physical Activity: Results of the ECLB-
11 COVID19 International Online Survey. *Nutrients*. (2020) (In press).
12
13
- 14 10. Hossain MM, Sultana A, Purohit N. Mental health outcomes of quarantine and isolation for
15 infection prevention: A systematic umbrella review of the global evidence. *Epidemiol Health*.
16 (2020) (In press).
17
18
- 19 11. Cucato GG, Correia MdA, Farah BQ, Saes GF, Lima AHdA, Ritti-Dias RM, et al. Validation
20 of a brazilian portuguese version of the walking estimated-limitation calculated by history
21 (WELCH). *Arq Bras Cardiol*. 2016;106(1):49-55.
22
23
- 24 12. Farah BQ, Ritti-Dias RM, Cucato GG, Montgomery PS, Gardner AW. Factors Associated
25 with Sedentary Behavior in Patients with Intermittent Claudication. *Eur J Vasc Endovasc Surg*.
26 2016; 52: 809-14.
27
28
- 29 13. Farah BQ, Ritti-Dias RM, Montgomery PS, Casanegra AI, Silva-Palacios F, Gardner AW.
30 Sedentary behavior is associated with impaired biomarkers in claudicants. *J Vasc Surg*. 2016; 63:
31 657-63.
32
33
- 34 14. Kaiser MJ, Bauer JM, Ramsch C, Uter W, Guigoz Y, Cderholm T, et al. Validation of the
35 Mini Nutritional Assessment short-form (MNA®-SF): A practical tool for identification of nutri-
36 tional status. *J Nutr Health Aging*. 2009; 13:782–88.
37
38
- 39 15. The World Health Organization quality of life assessment (WHOQOL): Position paper from
40 the World Health Organization. *Soc Sci Med*. 1995;41(10):1403-1409.
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

16. Jin JM, Bai P, He W, Wu F, Liu XF, Han DM, et al. Gender Differences in Patients With COVID-19: Focus on Severity and Mortality. *Front Public Health*. (2020) (In press).
17. COVID-19. Painel Coronavírus. <https://covid.saude.gov.br> (assessed 12 May 2021).
18. Cheval B, Sivaramakrishnan H, Maltagliati S, Fessler L, Forestier C, Sarrazin P, et al. Relationships between changes in self-reported physical activity, sedentary behaviour and health during the coronavirus (COVID-19) pandemic in France and Switzerland. *J Sports Sci*. 2021; 9:699-704.
19. Luciano F, Cenacchi V, Vegro V, Pavei G. COVID-19 lockdown: physical activity, sedentary behaviour and sleep in Italian medicine students. *Eur J Sport Sci*. (2020) (In press).
20. Caputo EL, Reichert FF. Studies of Physical Activity and COVID-19 During the Pandemic: A Scoping Review. *J Phys Act Health*. 2020; 17:1275-84.
21. Tison GH, Avram R, Kuhar P, Abreau S, Marcus GM, Pletcher MJ, et al. Worldwide Effect of COVID-19 on Physical Activity: A Descriptive Study. *Ann Intern Med* 2020; 173:767-70.
22. Heikkila K, Coughlin PA, Pentti J, Kivimaki M, Halonen JI. Physical activity and peripheral artery disease: Two prospective cohort studies and a systematic review. *Atherosclerosis* 2019; 286: 114-20.
23. Gengo e Silva Rde C, de Melo VF, Wolosker N, Consolim-Colombo FM. Lower functional capacity is associated with higher cardiovascular risk in Brazilian patients with intermittent claudication. *J Vasc Nurs*. 2015; 33: 21-5.
24. Nead KT, Zhou M, Diaz Caceres R, Olin JW, Cooke JP, Leeper NJ. Walking impairment questionnaire improves mortality risk prediction models in a high-risk cohort independent of peripheral arterial disease status. *Cir Cardiovasc Qual Outcomes*. 2013; 6: 255-61.

1
2
3
4 25. Thomas M, Patel KK, Gosch K, Labrosciano C, Mena-Hurtado C, Fitridge R, et al. Mental
5 health concerns in patients with symptomatic peripheral artery disease: Insights from the POR-
6 TRAIT registry. J Psychosom Res. (2020) (In press).
7
8

9
10
11 26. Sliwka A, Furgal M, Maga P, Drelicharz L, Mika P, Wloch T, et al. The role of psycho-
12 pathology in perceiving, reporting and treating intermittent claudication: a systematic review. Int
13 Angiol. 2018; 37: 335-45.
14
15
16
17
18
19
20
21
22

23
24 **Figure legends**

25
26 Fig. 1 Participants' flowchart
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65




Table 1. Factors related with COVID-19 in patients with peripheral artery disease.

	N=136
Variables, n (% yes)	
COVID-19 diagnosis	2 (1.5)
Recovered COVID-19*	2 (100)
<i>Preventive attitudes</i>	
Social isolation	120 (88.2)
Hands sanitization	134 (98.5)
Avoid leaving home	114 (83.8)
Avoid contact people	132 (97.1)
Contact with people diagnosed with COVID-19	7 (5.1)
<i>Comorbid conditions</i>	
Smoking	21 (15.4)
Ex-smoker	84 (61.8)
Diabetes mellitus	61 (44.9)
Hypertension	114 (83.8)
Dyslipidemia	105 (77.2)
Respiratory disease	22 (16.2)

Data presented as absolute and relative frequency. *: n=2.

Table 2. Impact of COVID-19 on sedentary and physical activity habits in patients with peripheral artery disease (n=136).

Patients with peripheral artery disease	
Variables, n (% yes)	
<i>Sedentary habits</i> (n=136)	
Spend more time watching TV	80 (58.8)
Spend more time sitting	104 (76.5)
<i>Physical activity habits</i> (n=136)	
Physical exercise before COVID-19	74 (54.4)
Physical exercise currently	39 (28.7)
<i>Modalities</i> (n=39)	
Aerobic exercise	24 (61.5)
Functional exercise	14 (35.9)
<i>Frequency</i> (n=39)	
1–2 x/week	8 (20.5)
3–4 x/week	11 (28.2)
5–7 x/week	20 (51.3)
<i>Duration</i> (n=39)	
≤ 30 min	10 (25.6)
31-60 min	23 (59.0)
≥ 61 min	6 (15.4)

Data presented as absolute and relative frequency.

Table 3. Impact of COVID-19 on eating habits and physical and mental health in patients with peripheral artery disease.

		N=136
Variables, n (% yes)		
<i>Eating habits</i>		
	healthier	15 (11.0)
	equal	107 (78.7)
	less healthy	14 (10.3)
	Increase in consumption of fried foods	14 (10.3)
	Increase in consumption of sweets	24 (17.6)
	Increase in consumption of fruits	46 (33.8)
	Increase in consumption of vegetables/legumes	33 (24.3)
<i>Physical health</i>		
	Decline in walking capacity	59 (43.4)
<i>Mental health</i>		
	More anxious	69 (50.7)
	Unhappier	41 (30.1)
	More stressed	34 (25.0)
	More depressed	27 (19.9)

Data presented as absolute and relative frequency.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Conflict of interest

The authors have no conflict of interest.