

Impacts of Modified Atmosphere Packaging and Controlled Atmospheres on Aroma, Flavor, and Quality of Horticultural Commodities

James Mattheis¹ and John K. Fellman²

ADDITIONAL INDEX WORDS. anaerobiosis, volatiles, ethanol, acetaldehyde, esters

SUMMARY. The commercial use of modified atmosphere packaging (MAP) technology provides a means to slow the processes of ripening and senescence during storage, transport, and marketing of many fresh fruit and vegetables. The benefits of MAP and controlled atmosphere (CA) technologies for extending postharvest life of many fruit and vegetables have been recognized for many years. Although both technologies have been and continue to be extensively researched, more examples of the impacts of CA on produce quality are available in the literature and many of these reports were used in development of this review. Storage using MAP, similar to the use of CA storage, impacts most aspects of produce quality although the extent to which each quality attribute responds to CA or modified atmosphere (MA) conditions varies among commodities. Impacts of MAP and CA on flavor and aroma are dependent on the composition of the storage atmosphere, avoidance of anaerobic conditions, storage duration, and the use of fresh-cut technologies before storage.

A number of factors modify the impact of MAP on produce quality. Package gas composition determines the effectiveness of any packaging system. Reduced O₂ and elevated CO₂ concentrations must be sufficiently stringent to slow metabolism and provide shelf-life extension while also being within the tolerance range of the stored commodity to avoid induction of anaerobic stress. Temperature management is critical as gas composition within packages changes with temperature (see Kader et al., 1989). Duration in the package after sealing is also important, as commodity tolerance to atmospheres may change over time (Mattheis et al., 1997). Finally, various processes of produce ripening and senescence do not have the same O₂ and CO₂ optima for maximizing beneficial responses.

Impacts of MAP and CA on quality

Most aspects of produce quality are impacted by MA (Kader et al., 1989; Weichmann, 1986).

The cost of publishing this paper was defrayed in part by the payment of page charges. Under postal regulations, this paper therefore must be hereby marked *advertisements* solely to indicate this fact.

¹USDA, ARS Tree Fruit Research Laboratory, 1104 N. Western Avenue, Wenatchee, WA 98801.

²Department of Horticulture and Landscape Architecture, Washington State University, Pullman, WA 99164-6414.

Color changes during produce ripening and senescence are highly impacted by storage atmosphere conditions, particularly the change from green to yellow. Storage in high CO₂ and/or low O₂ results in reduced loss of chlorophyll as well as reduced accumulation of other pigments including anthocyanin, lycopene, xanthophylls and carotenoids (Barth et al., 1993; Barth and Hong, 1996; Salunkhe and Wu, 1973; Wang et al., 1971; Zhuang et al., 1994). Within the tolerance range of O₂ and CO₂, reduced development of browning may also occur (Brecht et al., 1973; Smyth et al., 1998). Outside the tolerance range, particularly for CO₂, browning may be intensified (Brecht et al., 1973; Chen et al., 1981). Undesirable changes in texture are often reduced by high CO₂ and/or low O₂ within the tolerance range (see Weichmann, 1986). This includes reduced softening as well as reduced development of toughness after processing (Lougheed and Dewey, 1966). Outside of the tolerance range, processes of softening may be accelerated (Nanos and Mitchell, 1991; Patterson, 1982). CA and MAP can reduce the use of carbohydrates and titratable acids resulting in slower acid and sugar loss during storage (Salunkhe and Wu, 1973). Alternatively, loss of sugar and titratable acid can be enhanced if anaerobic conditions develop within the package.

MAP and CA conditions also impact a number of compounds that contribute to produce nutritive value. Ascorbate losses decrease when many commodities are stored under low O₂ conditions at low temperatures (Bangerth, 1977; Delaporte, 1971; Platenius and Jones, 1944). For many fruit and vegetables, as CO₂ partial pressure is increased under low or ambient O₂ partial pressure conditions, ascorbate losses increase but may recover to prestress amounts after storage in air (Weichmann, 1986). Loss of b-carotene in carrots (*Daucus carota* L.) decreases at low O₂ partial pressure but losses increase with the addition of CO₂ (Weichmann, 1986). The response to CO₂ is concentration dependent, below 5% (1% = 1.013 kPa at sea level) losses increase, at 10% losses decrease relative to carrots stored in air.

Impacts of MAP and CA on flavor and aroma

Flavor can also be altered by storage in MA. When CO₂ and O₂ are

maintained in the tolerance range, flavor deterioration is slowed by the combination of reduced loss of sugar, acid, and changes in other compounds that contribute to flavor. Flavor may deteriorate at a faster rate if anaerobic conditions develop due to both increased sugar and/or acid loss, and development of off-flavors. Off-flavors result from the accumulation of ethanol and acetaldehyde as well as altered produc-

tion of other compounds that can contribute to flavor and aroma. Ethanol is used in ester production by fruit (Berger and Drawert, 1984), and increased ethanol present after anaerobiosis can result in increased synthesis of ethyl esters. The increased amount of ethyl esters can alter aroma directly and also due to reduced production of other esters that are produced under normal conditions of fruit ripening (Mattheis et al., 1991).




Fig. 1. Production of straight C-chain esters by 'Gala' apple fruit following storage at 0 °C in air or controlled atmosphere. Fruit were removed from storage then held at 20 °C for 7 d before analysis by gas chromatography-mass spectrometry.




Fig. 2. Production of branched C-chain esters by 'Gala' apples following storage at 0 °C in air or controlled atmosphere. Fruit were removed from storage then held at 20 °C for 7 d before analysis by gas chromatography-mass spectrometry.

The excessive production of sulfur compounds following storage under anaerobic conditions contributes to off-flavor development in broccoli (*Brassica oleracea* L. Italica Group) (Forney et al., 1991). Reduced production of compounds that contribute to aroma in the absence of anaerobic conditions can occur following storage in low O₂/high CO₂ atmospheres (Guadagni et al., 1971; Patterson et al., 1974).

Responses to low O₂ and/or high CO₂ partial pressures that impact fruit aroma vary with commodity, cultivars and storage conditions. Aroma of minimally processed honeydew melons (*Cucumis melo* L. var. *inodorus* Naud.) deteriorates when melon pieces are stored in air at 5 °C (41 °F) (Portella and Cantwell, 1998). Storage of these pieces in air plus 15% CO₂ reduces aroma deterioration as indicated by higher taste panel ratings after 12 d storage. Although storage in 15% CO₂ reduced firmness loss in only one of the 4 cultivars evaluated, aroma scores for all 4 cultivars were higher compared to pieces stored in air. Strawberries (*Fragaria × ananassa* Duch. 'Pajaro') stored in 20% CO₂ contain higher amounts of acetaldehyde, ethanol and other ethyl esters compared to fruit stored in air over a 12 d period (Larsen and Watkins, 1995). Accumulation of these compounds occurs gradually during storage, and off-flavor development occurs coincident with this accumulation. Storage in air after removal from high CO₂ reduces the detection of off-flavors depending on the duration of anaerobiosis. Off-flavors detected in fruit stored 6 d in 20% CO₂ were no longer present after an additional 6 d storage in air. Ethyl acetate and ethyl butanoate are typically produced by 'Pajaro' strawberries in small amounts. Production of both increased following storage at 20% CO₂. Of these two compounds, ethyl butanoate has a lower odor threshold, therefore production of ethyl butanoate may have contributed to the loss of off-flavors by masking the presence of other compounds, particularly ethyl acetate, that contribute to off-flavor development. Concentrations of other volatiles that contribute to strawberry flavor and aroma were impacted less, if at all, by high CO₂. For example, concentrations of *E*-2-hexenal, *g*-decalactone, 2,5-dimethyl-4-methoxy-3(2*H*)-furanone and 2,5-dimethyl-4-hydroxy-3(2*H*)-furanone, all of which contribute to the flavor of strawberries (Larsen et al., 1992)

were similar in fruit stored in air or high CO₂.

Storage atmosphere and storage duration are important factors determining volatile production by sweet cherries (*Prunus avium* L. 'Bing') (Mattheis et al., 1997). Storage in nonanaerobic conditions (5% O₂ with up to 10% CO₂) for 12 weeks at 0 °C (32 °F) did not impact production of benzaldehyde and *E*-2-hexenal, two compounds that contribute to sweet cherry aroma and flavor (Schmid and Grosch, 1986). Storage in 5% O₂ with 0.1% CO₂ resulted in an accumulation of 2-propanol although no off-flavor developed. Sweet cherries produce only trace amounts of ethanol and ethyl esters under normal storage conditions, however, these compounds accumulated in fruit stored 12 weeks in 5% O₂ with 15 or 20% CO₂, indicating tolerance of 'Bing' sweet cherries to high CO₂ environments changes with storage duration.

Production of volatiles that contribute to apple flavor and aroma can be negatively impacted by storage atmospheres that do not induce anaerobiosis. As O₂ and CO₂ partial pressures are reduced and increased, respectively, fruit volatile production following removal from storage is reduced (Streif and Bangerth, 1988). The degree to which volatile production is reduced increases with the duration of CA storage. Volatile production can increase after a period of air storage following removal from a controlled atmosphere, however, the amount of recovery decreases with increased duration of CA storage.

Apple volatile concentration is also reduced in the package in a low oxygen MAP system (Song et al., 1997).

A differential impact of O₂ and CO₂ partial pressures on the types of esters produced by apples (*Malus sylvestris* Mill., *M. pumila* Mill., *M. domestica* Borkh.) has been reported (Brachmann et al., 1993). At 1% O₂ with 1% CO₂, production of straight C chain esters is reduced while storage in 1% O₂ with 3% CO₂ also impacts production of esters with branched C chains. Results from a storage experiment (J. Mattheis, unpublished) using 'Gala' apples further illustrate the importance of atmosphere composition and storage duration on poststorage volatile production. Apples were stored for up to 120 d at 0 °C in air or CA at 1 or 3% O₂ with 0.1% to 5% CO₂. Fruit were held in air at 20 °C (68 °F) for 7 d following removal from CA, then volatile emissions were characterized. A reduction in straight C chain ester production was observed after 60 d storage only in fruit stored in 1% O₂ (Fig. 1). Production of branch chain esters was not altered by any of the CA treatments after 60 d (Fig. 2). After 90 d, straight chain ester production decreased with the addition of CO₂ to either O₂ treatment, while CO₂ addition reduced branched chain ester production only for fruit stored at 1% O₂. The impact of storage CO₂ partial pressure on straight and branched chain ester production after 120 d storage is evident only for fruit stored at 3% CO₂. As CO₂ partial pressure increases, ester




Fig. 3. Production of 4-allylanisole by 'Gala' apples following storage at 0 °C in air or controlled atmosphere. Fruit were removed from storage then held at 20 °C for 7 d before analysis by gas chromatography-mass spectrometry.

production is reduced. At 1% O₂ ester production is low regardless of CO₂ concentration. Impacts of CA conditions on production of 4-methoxyallylbenzene, a product of the shikimic acid pathway (Manitto et al., 1974) that imparts a spicy character to apple aroma (Williams et al., 1977), were also related to O₂ and CO₂ partial pressures and duration of storage (Fig. 3). Production of 4-methoxyallylbenzene decreased with increased CO₂ partial pressure after 90 d storage, however after 120 d this CO₂ response was evident only for fruit stored at 3% O₂.

The use of MAP may also impact flavor and aroma directly by the nature of interactions between aroma compounds and packaging materials. Films slow movement of volatile aroma compounds but the solubility coefficients and diffusivity of volatile aroma compounds varies for different polymers (Charara et al., 1992). This variability may result in differential movement of aroma compounds through packaging films and the potential for change in aroma and flavor. Aroma transfer through methylcellulose-based films increases with the water vapor transfer rate (Debeaufort and Voilley, 1994), therefore packaging that is optimized for a high rate of water movement also provides for higher transmission of aroma compounds. Restricted movement of volatile compounds through packaging materials can also be used to increase concentrations of exogenous volatile materials to enhance flavor and/or reduce development of decay (Song et al., 1997).

Conclusions

The impact of MAP and CA on flavor and aroma is dependent on a number of factors. Temperature management is critical to avoid induction of anaerobic metabolism that leads to off-flavor development and loss of quality. Atmospheres that reduce the rate of quality deterioration while avoiding induction of anaerobiosis may still result in reduced production of other compounds that contribute to flavor and aroma. Reduced production of aroma compounds develops over long storage periods typical of CA applications, however, impacts on flavor development after short-term storage have not been well characterized. The potential for flavor enhancement and decay reduction by introduction of exogenous vola-

tile materials into packages provides another tool to manipulate fruit and vegetable quality by the use of MAP.

Literature cited

- Bangerth, F. 1977. The effect of different partial pressures of CO₂, C₂H₄ and O₂ in the storage atmosphere on the ascorbic acid content of fruits and vegetables. *Qualitas Plant.* 27:125-133.
- Barth, M.M. and Z.A. Hong. 1996. Packaging design affects antioxidant vitamin retention and quality of broccoli florets during postharvest storage. *Postharvest Biol. Technol.* 9 :141-150.
- Barth, M.M., E.L. Kerbel, A.K. Perry, and S.J. Schmidt. 1993. Modified atmosphere packaging affects ascorbic acid, enzyme activity and market quality of broccoli. *J. Food Sci.* 58 :140-143.
- Berger, R.G. and F. Drawert. 1984. Changes in the composition of volatiles by post-harvest application of alcohols to Red Delicious apples. *J. Sci. Food Agr.* 35:1318-1325.
- Brachmann, A., J. Streif, and F. Bangerth. 1993. Relationship between a reduced aroma production and lipid metabolism of apples after long-term controlled atmosphere storage. *J. Amer. Soc. Hort. Sci.* 118:243-247.
- Brecht, P.E., A.A. Kader, and L.L. Morris. 1973. The effect of composition of the atmosphere on duration of exposure on brown stain of lettuce. *J. Amer. Soc. Hort. Sci.* 98:536-538.
- Charara, Z.N., J.W. Williams, R.H. Schmidt, and M.R. Marshall. 1992. Orange flavor absorption into various polymeric packaging materials. *J. Food Sci.* 57:963-966.
- Chen, P.M., W.M. Mellenthin, S.B. Kelly, and T.J. Facticeau. 1981. Effects of low oxygen and temperature on quality retention of 'Bing' cherries during prolonged storage. *J. Amer. Soc. Hort. Sci.* 106: 533-535.
- Debeaufort, F. and A. Voilley. 1994. Aroma compound and water vapor permeability of edible films and polymeric packagings. *J. Agr. Food Chem.* 42: 2871-2875.
- Delaporte, N. 1971. Influence of oxygen atmospheres on the loss of ascorbic acid during storage of apples. *Food Sci. Tech.* 4:106-112.
- Forney, C.F., J.P. Mattheis, and R.K. Austin. 1991. Volatile compounds produced by broccoli under anaerobic conditions. *J. Agr. Food Chem.* 39:2257-2259.
- Guadagni, D.G., J.L. Bomben, and J.S. Hudson. 1971. Factors influencing the development of aroma in apple peels. *J. Sci. Food Agr.* 22:111-115.
- Kader, A.A., D. Zagory, and E.L. Kerbel. 1989. Modified atmosphere packaging of fruits and vegetables. *Crit. Rev. Food Sci. Nutr.* 28:1-30.
- Larsen, M., L. Poll, and C.E. Olsen. 1992. Evaluation of the aroma composition of some strawberry (*Fragaria ananassa* Duch) cultivars by use of odour threshold values. *Zeitschrift Fuer Lebensmittel-Untersuchung und Forschung* 195:536-539.
- Larsen, M. and C.B. Watkins. 1995. Firmness and aroma composition of strawberries following short-term high carbon dioxide treatments. *HortScience* 30:303-305.
- Lougheed, E.C. and D.H. Dewey. 1966. Factors affecting the tenderizing effect of modified atmospheres on asparagus spears during storage. *Proc. Amer. Soc. Hort. Sci.* 89:336-345.
- Manitto, P., D. Monti, and P. Gramitica. 1974. Biosynthesis of phenylpropanoid compounds. Part I. Biosynthesis of eugenol in *Ocimum basilicum* L. *J. Chem. Soc. Perkin Trans* 1:1727-1731.
- Mattheis, J.P., D.A. Buchanan, and J.K. Fellman. 1991. Change in apple fruit volatiles after storage in atmospheres inducing anaerobic metabolism. *J. Agr. Food Chem.* 39:1602-1605.
- Mattheis, J.P., D.A. Buchanan, and J.K. Fellman. 1997. Volatile constituents of 'Bing' sweet cherry fruit following controlled atmosphere storage. *J. Agr. Food Chem.* 45:212-216.
- Nanos, G.D. and F.G. Mitchell. 1991. Carbon dioxide injury and flesh softening following high-temperature conditioning in peaches. *HortScience* 26:562-563.
- Patterson, B.D., S.G.S. Hatfield, and M. Knee. 1974. Residual effects of controlled atmosphere storage on the production of volatile compounds by two varieties of apples. *J. Sci. Food Agr.* 25:843-849.
- Patterson, M.E. 1982. CA storage of cherries, p. 149-154. In: D.G. Richardson and M. Meheriuk (eds.). *Controlled atmospheres for storage and transport of perishable agricultural commodities.* Timber Press, Beaverton, Ore.
- Platenius, H. and J.B. Jones. 1944. Effect of modified atmosphere storage on ascorbic acid content of some vegetables. *Food Res.* 9:378-385.
- Portella, S.I. and M.I. Cantwell. 1998. Quality changes of minimally processed honeydew melons stored in air or controlled atmosphere. *Postharvest Biol. Technol.* 14:351-357.
- Salunkhe, D.K. and M.T. Wu. 1973. Effects of subatmospheric pressure storage on ripening and associated chemical changes of certain deciduous fruits. *J. Amer. Soc. Hort. Sci.* 98:12-14.
- Schmid, W. and W. Grosch. 1986. Quantitative analysis of the volatile flavour compounds having high aroma values from sour (*Prunus cerasus* L.) and sweet (*Prunus avium* L.) cherry juices and jams. *Z. Lebensm. Unters. Forsch* 193:39-44.
- Smyth, A.B. J. Song, and A.C. Cameron. 1998. Modified atmosphere packaged cut iceberg lettuce: effect of temperature and O₂ partial pressure on respiration and quality. *J. Agr. Food Chem.* 46: 4556-4562.
- Song, J., W. Deng, L. Fan, J. Verschoor, and R. Beaudry. 1997. Aroma volatiles and quality changes in modified atmosphere packaging, p. 89-95. In: J. Gorny (ed.). *Proc. 7th Intl. Controlled Atmosphere Res. Conf. vol. 5. Fresh-cut fruits and vegetables and MAP.* Univ. Calif. Postharvest Hort. Ser. 19.
- Streif, J. and F. Bangerth. 1988. Production of volatile aroma substances by 'Golden Delicious' apple fruit after storage for various times in different CO₂ and O₂ concentrations. *J. Hort. Sci.* 63:193-199.
- Wang, S.S., N.F. Haard, and G.R. DiMarco. 1971. Chlorophyll degradation during controlled atmosphere storage of asparagus. *J. Food Sci.* 36:657-661.
- Weichmann, J. 1986. The effect of controlled atmosphere storage on the sensory and nutritional quality of fruits and vegetables. *Hort Rev.* 8:101-125.
- Williams, A.A., O.G. Tucknott, and M.J. Lewis. 1977. 4-methoxyallylbenzene: An important aroma component of apples. *J. Sci. Food Agr.* 28: 185-190.
- Zhuang, H. K.Y., M.M. Barth, and D.F. Hildebrand. 1994. Packaging influenced total chlorophyll, soluble protein, fatty acid composition and lipoygenase activity in broccoli florets. *J. Food Sci.* 59 :1171-1174.