

## Review Article

# Implication of Gut Microbiota in Cardiovascular Diseases

Wenyi Zhou <sup>1,2</sup>, Yiyu Cheng,<sup>1</sup> Ping Zhu <sup>2</sup>, M. I. Nasser <sup>2</sup>, Xueyan Zhang <sup>1</sup>,  
and Mingyi Zhao <sup>1</sup>

<sup>1</sup>Department of Pediatrics, The Third Xiangya Hospital, Central South University, Changsha, Hunan 410013, China

<sup>2</sup>Guangdong Cardiovascular Institute, Guangdong Provincial People's Hospital, Guangdong Academy of Medical Sciences, Guangzhou, Guangdong 510100, China

Correspondence should be addressed to Mingyi Zhao; 36163773@qq.com

Received 3 May 2020; Revised 13 July 2020; Accepted 16 July 2020; Published 26 September 2020

Guest Editor: Wai Lydia Tai

Copyright © 2020 Wenyi Zhou et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Emerging evidence has identified the association between gut microbiota and various diseases, including cardiovascular diseases (CVDs). Altered intestinal flora composition has been described in detail in CVDs, such as hypertension, atherosclerosis, myocardial infarction, heart failure, and arrhythmia. In contrast, the importance of fermentation metabolites, such as trimethylamine N-oxide (TMAO), short-chain fatty acids (SCFAs), and secondary bile acid (BA), has also been implicated in CVD development, prevention, treatment, and prognosis. The potential mechanisms are conventionally thought to involve immune regulation, host energy metabolism, and oxidative stress. However, numerous types of programmed cell death, including apoptosis, autophagy, pyroptosis, ferroptosis, and clockophagy, also serve as a key link in microbiome-host cross talk. In this review, we introduced and summarized the results from recent studies dealing with the relationship between gut microbiota and cardiac disorders, highlighting the role of programmed cell death. We hope to shed light on microbiota-targeted therapeutic strategies in CVD management.

## 1. Introduction

Cardiovascular disease (CVD), with its rising prevalence rate and mortality, entails both health threats and economic burdens to our society. As a chronic progressive condition, the development of CVDs often begins with risk factors like obesity, type 2 diabetes, and hypertension, most of which would irreversibly damage vascular structure and eventually lead to detrimental clinical outcomes like arterial thrombosis and ischemic stroke. While heredity can only be blamed for less than 20% occurrence of CVDs, dietary and nutritional statuses are two stimuli with more profound and lasting impacts [1]. Therefore, increasing evidence has suggested a close relationship between gut microbiota and CVD development [2].

The gut microbiota refers to trillions of commensal microorganisms located in the intestine in a certain proportion, whose balance is easily disturbed by food intake, lifestyle, and environment [3]. Considered a complex organ, the microbial community is required in the committed step

through which food would be converted into small compounds and metabolites, thus modulating intestine structure, gut barrier integrity, inflammatory status, and host metabolism both directly and indirectly [4]. Since Hippocrates claimed that “all diseases begin in the gut” centuries ago, a great body of research has demonstrated the interplay between intestinal microbiota and diseases, including colorectal cancer [5], cerebral ischemia-reperfusion injury [6], liver fibrosis [7], and CVDs [8]. The gut microbiota accounts for 0.2–2.0 kg of the weight of an adult and approximately 50% of the dry weight of adult feces. The enormous genome of microbial genes and their functions are described as the microbiome, which outnumbers the human genome tremendously [3, 9]. Although the characteristics of the gut community may be inherited in early life, the composition could also be altered by external conditions [10, 11]. Appropriate gut microbiota structure and metabolite functions are essential in homeostasis maintenance, whereas gut dysbiosis contributes to atherosclerosis, hypertension, heart failure, arrhythmia,

cardiac tumours, and others [12]. However, its underlying mechanisms are multifactorial and yet to be determined.

In this report, we introduce the role of gut microbiota in CVDs and summarize possible mechanisms, which may provide a theoretical basis and shed light on novel therapeutic strategies in the prevention and treatment of CVDs.

## 2. Mechanisms Underlying the Interaction between Gut Microbiota and the Host

The community of gut microbiota consists mostly of bacteria, fungi, and viruses in which the primary component is bacteria. There are 5 major families in the intestinal flora: *Bacteroidetes*, *Firmicutes*, *Actinobacteria*, *Proteobacteria*, and *Verrucomicrobia* [13]. Although the variety of species is abundant, the architecture of gut microbiota is comparatively fixed in different sites. However, the differences in gut microorganism quantities between locations are significant, with the ascending colon containing the largest number [13]. Under physiological conditions, more than 90% of the bacteria comprise *Bacteroidetes* and *Firmicutes*, while an elevated *Firmicutes/Bacteroidetes* (F/B) proportion is associated with CVDs [14]. Koliada et al. found that with the body mass index (BMI) in Ukraine adult population increasing, their F/B ratio raised likewise after removing other confounders such as age or smoking [15]. Subsequently, evaluation of children's gut microbiota composition and BMI had confirmed F/B ratio as a key risk indicator for childhood obesity [16]. Additionally, the F/B ratio is related to low-grade inflammation leading diabetes mellitus [17]. These diseases serve as both risk factors and stimulatives for CVDs. In addition to intestinal integrity maintenance, gut metabolites serve as essential messengers in the communication between gut microbiota and the host. Here, we review the mechanisms underlying the interaction between gut microbiota and the host, especially in CVDs.

**2.1. Immunoregulation.** Generated by fiber fermentation in the colon, short-chain fatty acids (SCFAs) include three major products, namely, acetate, propionate, and butyrate, all of which contain less than six carbons [18]. Apart from being nutrients and energy sources for intestinal epithelial cells, these small-molecule metabolites could enter the blood circulation, participate in immune regulation and inflammation modulation either by binding to G protein-coupled receptors (GPCRs) or by inhibiting histone deacetylases (HDACs) [18], and thereby influence gut homeostasis and host diseases. Laurence et al. found that SCFAs induce NLRP3 inflammasome activation and subsequent abundant IL-18 secretion in a GPR43- and GPR109A-dependent manner, thus eliciting favourable effects on intestinal integrity maintenance [19]. Of note, GPR43 and GPR109A are two receptors that are expressed on intestinal epithelial cells and some immune cells, where GPR43 mainly binds to acetate and propionate, while GPR109A is specifically activated by butyrate [20]. Studies have demonstrated that SCFAs beneficially upregulate not only the proliferation and differentiation of regulatory T cells (Tregs) but also the anti-inflammatory IL-10 secreted from Foxp3+ Tregs, which are mediated through GPR43 (also

known as *Ffar2*) activation and HDAC inhibition [21]. Additionally, butyrate was shown to suppress proinflammatory factors, including IL-6, IL-12, and NO, from intestinal macrophages by HDAC inhibition [18]. Likewise, Bartolomeus et al. recently proved that the anti-inflammatory role of SCFAs such as propionate significantly reduced the number of effector memory T cells and T helper 17 cells, thus mitigating cardiovascular damage [22]. However, the proinflammatory functions mediated by GPR41 (also known as *Ffar3*) and GPR43 were reported elsewhere [23], indicating that SCFA-induced immunoregulatory effects are dependent on the distinct cell types.

Additionally, trimethylamine N-oxide (TMAO) is generally investigated as a risk indicator for cardiovascular diseases, diabetes mellitus, nonalcoholic fatty liver disease, and other metabolic events [24–26]. As the end-product of dietary choline and L-carnitine, TMAO is converted from trimethylamine (TMA) in the liver by flavin-containing monooxygenases (FMOs), especially FMO3 [24]. However, how exactly TMAO functions to regulate homeostasis is seldom discussed. According to Sun et al., TMAO induces inflammation by activating the ROS-TXNIP-NLRP3 inflammasome, thereby contributing to endothelial dysfunction in human umbilical vein endothelial cells [27]. Similarly, Yue et al. showed that TMAO promotes the release of the inflammatory cytokines IL-1 $\beta$  and IL-18 via activation of the NLRP3 inflammasome from foetal human colon cells in a time- and dose-dependent manner [28]. Moreover, injection of TMAO was shown to significantly increase inflammatory markers, including cyclooxygenase 2, IL-6, E-selectin, and ICAM1, through the MAPK and NF- $\kappa$ B signalling pathways, which then recruit leukocytes and induce vascular inflammation [29]. In these fine experiments in which treatments against TMAO were adopted, inflammatory damage was prevented. Taken together, the proinflammatory role of TMAO is established.

Plasma cholesterol, the key cellular membranes constituent and precursor of steroid hormones, vitamin D, and bile acids, is positively correlative with cardiovascular diseases. There are two main sources of cholesterol, with one-third being exogenous from daily dietary and the other two-third synthesized inside the body [30]. Confirmed with various models, microbial regulation is believed to be critically involved in cholesterol balance modulation [31]. To begin with, gut microbiome is reported to convert cholesterol into poorly absorbed coprostanol, reducing the risk of cardiovascular diseases [30, 32]. Further elucidation reveals that the presence of intestinal sterol metabolism A genes is responsible for such metabolism mediation [32]. Another key aspect the gut microbiota enrolled is bile acids metabolism. Bile acids deconjugation yields free bile acids as well as free glycine or taurine residues, which requires the participation of bile salt hydrolase enzymes (BSHs) [30]. The presence of BSHs was found within *Clostridium*, *Bifidobacterium*, *Lactobacillus*, and others. With higher degree of bile salts deconjugation, more free BAs were excreted into feces [30]. Primary bile acids refer to steroid molecules that result from the decomposition of cholesterol in the liver. Most of them are recycled back to the liver, while the rest enter the intestine,

where they are converted into secondary bile acids by gut microbiota [33]. The most well-studied secondary bile acids are deoxycholic acid (DCA), lithocholic acid (LCA), and ursodeoxycholic acid (UDCA), which often function through their receptors, including G protein-coupled BA receptor 1 (TGR5), farnesoid X receptor (FXR), and vitamin D receptor (VDR) [33]. When bound to the TGR5 receptor, secondary bile acids cause the activation of macrophages and then the production of inflammatory cytokines [34]. Interestingly, researchers found that low concentrations of secondary bile acids bring anti-inflammatory effects, while high concentrations would instead cause damage. For example, Wang et al. demonstrated that low-dose DCA mitigates the inflammatory response in birds [35].

Additionally, these products from commensal microbiota would trigger innate immune signalling, thereby communicating with the host. Microbial-associated molecular patterns (MAMPs) including LPS or peptidoglycan are recognized by receptors like Toll-like receptors (TLRs), NOD-like receptors (NLRs), and others [4]. The strong connection between TLRs and atherosclerosis was confirmed in genetic mice researches. In the TLR4<sup>-/-</sup> apoE<sup>-/-</sup> mice model fed with cholesterol-rich diet, the size of aortic plaque was significantly reduced [36]. Interestingly, deficiency of TLR2 in myeloid cells had no influence in the development of atherosclerosis, suggesting the role of endothelial TLR2 in atherogenesis [37]. Furthermore, the development of arterial thrombosis was relative to NOD2, TLR2, and TLR9 signalling in platelets as well as TLR2 and TLR4 pathways in endothelial cells [4].

**2.2. Energy Metabolism and Homeostasis.** Among the numerous risk factors contributing to CVD, abnormal immune regulation and metabolic disorders represent two major elements. Metabolic syndromes such as obesity, dyslipidosis, hyperglycaemia, and insulin resistance are closely related to the occurrence and development of CVD. In recent years, the link between gut microbiota, metabolism, and CVD has gained much attention. For instance, Den and his coworkers considered SCFAs to carry metabolic benefits for those with a high-fat diet through inhibition of peroxisome proliferator-activated receptor gamma (PPAR $\gamma$ ), converting lipid synthesis to lipid oxidation [38]. Moreover, a fiber-rich diet upregulates the levels of SCFAs in the gut, which then promotes intestinal gluconeogenesis [39]. SCFAs accelerate the production of GLP-1 by binding to GPR41 and GPR43, therefore facilitating insulin secretion [39]. In contrast, TMAO aggravates triglyceride accumulation and lipogenesis in the livers of high-fat diet-fed mice [40]. Propionate was found to induce glycogenolysis and hyperglycaemia via the upregulation of glucagon and fatty acid-binding protein 4 (FABP4), thereby hindering the effects of insulin [41]. In mice with obesity, bile acid promotes GLP-1 secretion via the TGR5 pathway, thereby modulating blood sugar [42]. Notably, there is multiplicity in the associations between gut microbiota and their microbiome. For instance, TMAO could alter the bile acid profile and metabolism, thus contributing to liver steatosis and atherosclerosis [40, 43], whereas bile acid stimulates FMO3 expression via FXR, eventually resulting in TMAO production (Bennett et al., 2013).

Moreover, butyrate was found to restore bile acid dysregulation and counteract hepatic inflammation [44].

To sum up, the gut microbiota communicates with the host through diverse manners. To begin with, SCFAs and secondary bile acids are two of the main products by gut microbiota. They play their immune-regulatory role either by directly affecting the proliferation of immune cells or by stimulating the production of cytokines. Moreover, SCFAs are involved in both lipid and sugar metabolism. Second, TMAO that primarily comes from L-carnitine and choline consumption participates in inflammatory modulation by promoting IL-18 and IL-1 $\beta$  release or activating MAPK/NF- $\kappa$ B signalling pathway, thus upregulating the levels of COX2, IL-6, and ICAM1. Moreover, MAMPs including LPS and peptidoglycan serve as another vital contributor in the development of atherosclerosis and arterial thrombosis, mainly through TLRs and NLRs (Figure 1).

**2.3. Programmed Cell Death.** Apart from the well-known immune and inflammation modulation properties of gut microbiota, accumulating evidence has revealed its potential in the determination of diverse manners of cell death (Figure 2).

**2.3.1. Apoptosis.** Characterized by the formation of a distinctive apoptotic body, apoptosis is one of the most widely investigated programmed cell deaths. It is often observed in myocardial infarction, heart failure, and other vascular damage. Saito et al. found that *Bacteroides fragilis* (*B. fragilis*) is able to protect HT29 cells from apoptosis resulting from Shiga toxin [45]. Butyrate promotes vascular smooth muscle cell growth via proliferation arrest as well as apoptosis inhibition [46]. Notably, there are proapoptotic effects as well. Sodium propionate was reported to induce apoptosis in H1299 and H1703 lung cancer cells, as evidenced by increased protein expression of p21, Bad, and Bax as well as apoptosis markers, including cleaved PARP and cleaved caspase 3 [47]. According to Nie et al., *Bifidobacterium* (BIF) ameliorates TNF- $\alpha$ -induced cell apoptosis in Caco-2 cells [48]. Likewise, butyrate causes apoptosis and cell cycle arrest in kidney epithelial cells [49].

**2.3.2. Autophagy.** Nie et al. discovered that BIF ameliorates TNF- $\alpha$ -induced autophagy in Caco-2 cells by suppressing the level of p62 and inhibiting the expression of autophagy-related markers such as Beclin1 and LC3II [48]. According to their research, BIF may provide a therapeutic target aimed at the Kawasaki disease, which is highly related to acquired heart disease in children. Lannucci and his coworkers proved that SCFAs induce autophagy in hepatic cells by uncoupling protein 2 (UCP2) [50]. Accordingly, Qiao et al. demonstrated that sodium butyrate contributes to the reduction in  $\alpha$ -synuclein both via the inhibition of the PI3K/Akt/mTOR autophagy pathway and enhancement of Atg5-mediated autophagy, manifested as elevated LC3II and reduced p62 expression [51].

**2.3.3. Pyroptosis.** As a type of proinflammatory cell death, pyroptosis is characterized by swollen cells, subcellular organelle damage, and the release of cytokines, including the NLRP3 inflammasome, NLRP6, an apoptosis-associated speck-like protein containing CARD (ASC), cysteinyl-




FIGURE 1: Mechanisms involved in gut microbiota-host communication. Short-chain fatty acids (SCFAs), mainly propionate, acetate, and butyrate, stimulate Foxp3<sup>+</sup> Tregs and macrophages via GPR43 activation and HDAC inhibition. Foxp3<sup>+</sup> Tregs subsequently produce the anti-inflammatory cytokine IL-10, while proinflammatory cytokines such as IL-6 and IL-12 are secreted by macrophages. Moreover, Th17 cells and effector memory T cells were downregulated by SCFAs. By suppressing PPAR $\gamma$ , SCFAs promote lipid oxidation. Although insulin production was enhanced by SCFAs, glycogenolysis and gluconeogenesis were both observed to occur even with SCFA treatment. L-carnitine and choline consumption contribute to the release of trimethylamine (TMA), which is then converted by FMO into trimethylamine N-oxide (TMAO). Both SCFAs and TMAO activate the NLRP3 inflammasome, leading to IL-18 and IL-1 $\beta$  release. Through the MAPK/NF- $\kappa$ B signalling pathway, TMAO increases the levels of COX2, IL-6, and ICAM1. Secondary bile acids such as deoxycholic acid (DCA), lithocholic acid (LCA), and ursodeoxycholic acid (UDCA) are produced in the intestine by gut microbiota and then participate in inflammatory modulation and blood sugar regulation.

aspartate-specific proteinase 1 (caspase-1), and gasdermin D. Data have shown that sodium butyrate is capable of breaking down the gingival epithelial barrier by inducing pyroptosis [52]. Similarly, TMAO promotes vascular endothelial cell pyroptosis via ROS production, thus resulting in the development of atherosclerosis [53]. However, Gu et al. proved the antipyroptosis effects of sodium butyrate on renal glomerular endothelial cells, protecting them from damage caused by high glucose [54]. From the perspective of the mechanism, the classic caspase-1-gasdermin D pathway and NF- $\kappa$ B/I $\kappa$ B- $\alpha$  signalling may both be involved [54]. Moreover, Cohen et al. confirmed that *Vibrio proteolyticus* (VPRH), a Gram-negative bacterium from the gut of a wood borer, induces pyroptosis by activating the NLRP3 inflammasome and caspase-1, thereby resulting in IL-1 $\beta$  secretion, suggesting that the NLRP3 inflammasome pyroptotic pathway can benefit the host during infection [55].

**2.3.4. Ferroptosis.** Induced by lipid reactive oxygen species accumulation, ferroptosis refers to another distinct kind of cell death mediated by mitochondria. Studies concerning whether gut microbiota are implicated in ferroptosis are rather rare. Until recently, Robert et al. proposed that supplementation of omega-3 polyunsaturated fatty acids (n-3

PUFAs) and butyrate may both facilitate mitochondrial Ca<sup>2+</sup>- and Gpx4-dependent ferroptosis [56]. Hopefully, this hypothesis may shed light on the link between gut microbiota and ferroptosis as well as accelerate related research.

**2.3.5. Clockophagy.** The circadian rhythm, namely, clockophagy, is controlled by a complex circadian clock gene network including the ARNTL, CLOCK, CRY2, and PER2 genes [57]. The interaction between circadian rhythms and diverse gut microbiota has been well studied, where the acute sleep-wake cycle shift alters the functional profiles of gut microbes. Together, the clock-microbial communities affect host homeostasis [58]. The circadian rhythm of SCFA production was observed by Segers et al. to cause rhythmicity in intestinal movement [59]. However, such effects were abolished by the deletion of Bmal1 [59]. Besides, Marques et al. found that in hypertensive mice, a high-fiber diet changes the composition of the gut microbiota and restores gut dysbiosis, which may be partially due to increased levels of clock genes in the heart and kidney [60]. Additionally, a negative correlation between the phylum *Firmicutes* and Bmal1 as well as a positive correlation between *Bacteroidetes* and Bmal1 was observed in mice [61].




FIGURE 2: Manners of cell death induced by gut microbiota. A variety of gut flora have been demonstrated to be effective in regulating cell death. (a) Muciniphila and (b) fragilis were shown to counteract apoptosis. In contrast, sodium propionate has the ability to induce apoptosis. Interestingly, the effects of butyrate on apoptosis are controversial, manifesting elevated biomarkers such as P21, Bad, Bax, and caspase-3. In addition, SCFAs stimulate autophagy, while *Bifidobacterium* is autophagy-protective, with decreased expression of P62, Beclin1, and LC3II. Sodium butyrate promotes autophagy by inhibiting the PI3K/Akt/mTOR pathway. Additionally, it is involved in pyroptosis via regulation of the caspase-1/gasdermin D pathway. In addition, TMAO stimulates ROS activation and thus induces pyroptosis. Along with pyroptosis, the NLRP3/NLRP6 inflammasome and IL-1 $\beta$  are produced. Moreover, clockophagy can reverse gut dysbiosis. For instance, SCFAs are capable of controlling rhythmicity via clock genes such as Bmal1.

### 3. Implications of Gut Microbiota in CVDs

To concisely describe the role of gut microbiota in cardiovascular disease, the positive or negative effects of gut microbiota on CVDs are listed in Table 1.

**3.1. Hypertension.** Hypertension (HTN) has been a key link in the occurrence and development of cardiovascular diseases. Although HTN is currently beyond cure, it is preventable and controllable. According to the mosaic theory advanced by Irvin Page, HTN is induced by multiple factors, including inheritance, diet, and environment [62]. HTN also has extensive impacts on various tissues and organs, such as endothelial cells, the kidneys, and brain. Moreover, in recent years, the value of gut microbiota in HTN has been widely investigated.

In the work conducted by Li et al., fecal transplantation was performed from hypertensive individuals to germ-free mice. Along with microbiota shift, blood pressure was also elevated in those mice, indicating the contributing role of gut microbiota in hypertension [63]. It has been demonstrated that butyrate-producing bacteria and butyrate levels are relatively low in patients with HTN, indicating that imbalanced host-microbiome cross talk is relevant to systolic blood pressure [64]. Accordingly, in mice pretreated with angiotensin II, supplementation with butyrate effectively lowered blood pressure [65]. Interestingly, the same team found that gut barrier dysfunction is another contributor to HTN, as evidenced by elevated levels of zonulin, a gut epithelial

tight junction protein regulator [65]. However, the same metabolite may yield contradictory biological effects through different receptors. For instance, Jennifer et al. found that propionate may upregulate blood pressure via olfactory receptor 78 (Olf78) while exerting hypotensive effects through activation of Gpr41 [66]. In-depth knowledge reveals that vascular inflammation and endothelial dysfunction are two key processes in the development of hypertension [67]. In mice fed with Western diet, endothelial dysfunction was associated with decreased proportion of *Bifidobacterium* spp., whereas antibiotic administration helped mitigate such vascular damage [68]. As compared with germ-free mice, the conventionally raised mice pretreated with Ang II presented with a higher level of IL-4 and IL-10, indicating a vascular inflammation-prone role of enteric flora [68]. In a meta-analysis of 8 studies, a higher circulating TMAO level was positively associated with hypertension risk, which was dose-dependent [69]. Liu and coworkers identified that administration of the *Lactobacillus rhamnosus* GG strain is an effective approach to prevent exacerbation of HTN, which is in part mediated by reducing TMAO levels [70]. However, it is worth noting that the application of TMAO alone would not alter blood pressure in normotensive rats but prolonged the hypertensive-prone effects of angiotensin II [71]. More recently, a novel mechanism different from inflammation or immunity regulation has been presented. In high salt-induced hypertensive mice, elevated blood pressure is closely related to increased levels of intestinal-derived corticosterone [72].

TABLE 1: The exact role of different gut microbiota in CVDs.

CVDs Species	Atherosclerosis	Myocardial infarction	Heart failure	Arrhythmia
<i>Enterobacteriaceae</i>	Negative			
<i>Ruminococcus gnavus</i>	Negative			
<i>Eggerthella lenta</i>	Negative			
<i>Roseburia intestinalis</i>	Positive			
<i>Faecalibacterium cf. prausnitzii</i>	Positive			
<i>Synergistetes phylum</i>		Negative		
<i>Lachnospiraceae</i> family		Negative		
<i>Spirochaetes</i> phylum		Negative		
<i>Syntrophomonadaceae</i> family		Negative		
<i>Tissierella</i> and <i>Soehngenia</i> genera		Negative		
<i>Lactobacillus plantarum</i> 299v		Positive		
<i>Faecalibacterium prausnitzii</i>			Positive	
<i>Bacteroides fragilis</i>			Positive	
<i>Ruminococcus</i>				Negative
<i>Streptococcus</i>				Negative
<i>Enterococcus</i>				Negative
<i>Faecalibacterium</i>				Positive
<i>Alistipes</i>				Positive
<i>Oscillibacter</i>				Positive
<i>Bilophila</i>				Positive

Taken together, these results established that the gut microbiota is involved in blood pressure regulation. However, the underlying mechanisms still await further validation.

**3.2. Atherosclerosis and Arterial Thrombosis.** Initially related to dyslipidaemia, abnormal accumulation of macrophages, and massive production of inflammatory cytokines, atherosclerosis is considered a chronic inflammatory disease that underlies end-stage CVDs such as myocardial infarction or heart failure. In recent years, people have started to consider gut microbiota potent regulators during the development of atherosclerotic lesions. Koren et al. first identified bacterial DNA in atherosclerotic plaques, and the amount of DNA was associated with the infiltration of leukocytes in the plaques [73]. Moreover, the altered composition of the gut microbiome was confirmed in a metagenome-wide association study encompassing 218 individuals with atherosclerosis and 187 healthy controls. Specifically, the abundances of *Enterobacteriaceae*, *Ruminococcus gnavus*, and *Eggerthella lenta* were significantly increased in those with atherosclerosis, whereas *Roseburia intestinalis* and *Faecalibacterium cf. prausnitzii*, both butyrate-yielding bacteria, were reduced [74]. The above findings strongly suggest correlations between gut microbiota and atherosclerosis.

With the use of atherosclerosis-prone germ-free mice and antibiotic treatments, the role of gut microbiota in atherosclerosis development was further elucidated (Table 2). First people suggested that bacterial or viral infection is necessary for the initiation of atherosclerosis. However, such hypothesis was overturned by Samuel and his colleagues' work [75].

Apolipoprotein (apo) E<sup>-/-</sup> murine model was often adopted for atherosclerosis research given the self-driven ability of atherosclerotic plaque formation. Samuel et al. compared the atherosclerosis lesion in germ-free apoE<sup>-/-</sup> animals with those raised in conventional environment, and they found no evident difference [75]. Alternatively, with the help of antibiotics to suppress gut microflora, choline-enhanced atherosclerosis in aorta was off-set along with reduced macrophage and scavenger receptor CD36 [76]. However, given the complexity of enteric flora, the pro- or antiatherosclerosis role of gut microbiota depends. Kasahara and his colleagues demonstrated that *Roseburia intestinalis* is capable of ameliorating atherosclerosis by shaping gene expression, enhancing fatty acid metabolism, and reducing the inflammatory response [77]. However, treatment with butyrate markedly mitigates the formation of atherosclerotic plaques via the upregulation of ABCA1 and subsequent cholesterol efflux [78]. In contrast, the production of TMAO by gut microbiota yields negative effects on atherosclerosis [79].

Rupture of the atherosclerotic plaque would likely cause arterial thrombus elsewhere, resulting in detrimental consequences. For one, the LPS-TLR pathway is a major contributor in thrombosis formation. Both TLR2 and TLR4 were found expressed on endothelial cells and platelets. Activation of TLR2 and TLR4 pathway would facilitate the release of VWF and factor VIII expression, contributing to platelet-proinflammatory cell aggregation [80]. For another, gut microbiota metabolites take part in arterial thrombosis as well. Feces transplantation of TMAO-rich gut microbiota into germ-free mice would promote platelet function and

TABLE 2: Researches of gut microbiota in CVDs.

Diseases	Sample	Observations	Mechanism	Ref.
Hypertension	HTN patients	Decreased butyrate-producing bacteria and butyrate level	SCFA-dependent	[62]
	Ang-II pretreated mice	Reduced BP after butyrate administration; increased zonulin level	SCFA-dependent; gut barrier dysfunction	[65]
	Mice	Increased BP after propionate treatment	Olf78-dependent	[66]
	Mice	Decreased BP after propionate treatment	Gpr41-dependent	[66]
		<i>Lactobacillus rhamnosus</i> GG prevents HTN development	Reduced TMAO levels	[70]
	Mice	High salt-induced HTN	Increased intestinal-derived corticosterone	[72]
	Patients	Bacterial DNA observed in atherosclerotic plaques	/	[73]
Atherosclerosis		<i>Roseburia intestinalis</i> ameliorates atherosclerosis	Alter gene expression, induce fatty acid metabolism, and reduce inflammation response	[77]
	apoE <sup>-/-</sup> mice	Comparable atherosclerosis lesion in germ-free apoE <sup>-/-</sup> animals and their conventionally raised counterparts	/	[75]
		Choline-enhanced atherosclerosis in aorta was off-set by antibiotics	Reduced macrophage and scavenger receptor CD36	[76]
	apoE <sup>-/-</sup> mice with HFD	Butyrate mitigates atherosclerotic plaque formation	Upregulation of ABCA1 and subsequent cholesterol efflux	[78]
	AMI rat model	Increased <i>Synergistetes phylum</i> , <i>Lachnospiraceae family</i> , <i>Spirochaetes phylum</i> , <i>Syntrophomonadaceae family</i> , and <i>Tissierella</i> and <i>Soehngenia</i> genera	In parallel with gut barrier impairment	[83]
Myocardial infarction	STEMI patients	Over 12% plasma bacteria originated from the gut	Partially associated with an inflammatory response	[84]
	Patients presenting with chest pain	Predictive value of plasma TMAO levels for incident cardiovascular events	TMAO-related proinflammatory monocytes augment	[85]
	Mice	Improve cardiac repair and post-MI outcome through modulation of immune composition	Gut microbiota-derived SCFAs modulate immune composition	[86]
		<i>Lactobacillus plantarum</i> 299v improved ischemia tolerance and acute cardiac injury after MI	Reduce leptin level	[87]
	Mice	<i>Bacteroides fragilis</i> reduces ventricular remodelling	Increased Foxp3+ Treg cells and anti-inflammatory cytokine	[92]
Heart failure		Depletion of SCFAs finally leads to HF	Intestinal barrier destruction, with endotoxin translocation	[93, 94]
	Mice	TMAO alters cardiac muscle cells contractility	Promotion of calcium ions release	[95, 96]
		TMAO confers detrimental effects on adult cardiomyocytes	T-tubule network damage; Ca handling dysfunction	[97]
	Mice	Pulmonary edema, cardiac enlargement, and decreased ejection fraction	TMAO-dependent	[98]
	Patients	TMAO increases susceptibility to HF	Induction of myocardial fibrosis	[99]
	Overload-induced HF mice	DMB ameliorates adverse cardiac structural remodelling	Downregulating TMAO levels	[100]
	Patients	Shared common features of gut microbiota dysbiosis	Alike ratio of <i>Firmicutes</i> and <i>Bacteroidetes</i>	[104, 105]
Arrhythmia	Patients	Thrombus formation; platelet hyperreactivity	Elevated TMAO level	[107]
		TMAO stimulates ischemia-induced VA	Release of proinflammatory markers such as IL-1 $\beta$ and TNF- $\alpha$	[109]
	Canine AF model	Gut microbes counteracts AF progression	TMAO production and CANS activation	[110]
	Mice	Reduced susceptibility to cardiac ventricular arrhythmias	SCFA-dependent	[22]

arterial thrombosis [81]. Recently, another gut microbial metabolite, Phenylacetylglutamine (PAGln), was shown to induce hyperreactivity of platelet via adrenergic receptors [82].

**3.3. Myocardial Infarction.** The connection between intestinal flora and myocardial infarction (MI) has been supported by a growing body of literature. In a rat model of acute myocardial infarction (AMI), enrichment of the *Synergistetes* phylum, *Lachnospiraceae* family, *Spirochaetes* phylum, *Syntrophomonadaceae* family, and *Tissierella and Soehngenia* genera was observed compared with the sham group, which is in parallel with gut barrier impairment [83]. In patients with ST-elevation myocardial infarction (STEMI), systemic microbiome alteration was also observed. Over 12% of plasma bacteria were identified to originate from the gut after STEMI, which is partially associated with the inflammatory response [84]. Accordingly, reduced cardiac damage and decreased inflammation were noticed following the abrogation of bacterial translocation [84]. Of clinical value, plasma TMAO levels may be potential markers to predict the risks of incident cardiovascular events in patients presenting with chest pain [85]. Such potency may in part be explained by TMAO-related proinflammatory monocyte augmentation [85]. Moreover, Tang et al. demonstrated that gut microbiota-derived SCFAs would benefit cardiac repair and improve post-MI outcome through modulation of immune composition [86]. With the administration of the probiotic *Lactobacillus plantarum* 299v, the leptin level in blood was reduced, leading to enhancement of ischemic tolerance in the myocardium and alleviation of acute cardiac injury after MI [87].

**3.4. Heart Failure.** As an irreversible end-stage disease, heart failure (HF) is characterized by oedema and dyspnoea, with a five-year mortality rate of over 50% [88]. At present, a growing body of research has confirmed the “gut hypothesis of heart failure” [89, 90]. That is, decreased cardiac output in HF leads to intestinal mucosa barrier damage and dysbacteriosis, with elevated levels of pathogenic bacteria such as *Candida* [91] and reduced levels of anti-inflammatory bacteria such as *Faecalibacterium prausnitzii* [3]. Reciprocally, intestinal flora promotes HF development by participating in mucosal immunity modulation [3]. Segmented filamentous bacteria can stimulate the secretion of IL-6 and IL-23 and then promote the differentiation of Th17 cells. *Bacteroides fragilis* increases the abundance of Foxp3+ Treg cells and induces the secretion of anti-inflammatory cytokines, which have been found to reduce ventricular remodelling in MI mice [92].

Not surprisingly, metabolites of intestinal flora are also important for HF. Although studies concerning SCFAs and HF are limited, it has been proven that SCFAs are beneficial for the intestinal mucosa [3]. The depletion of SCFAs would result in intestinal barrier destruction, which then facilitates the translocation of endotoxin into blood circulation and finally leads to HF [93, 94].

However, the level of TMAO has long been recognized as a risk factor. Savi et al. found that TMAO promotes the release of calcium ions in cardiac muscle cells of healthy mice and thus alters their contractility [95, 96]. Recently, the in-

depth work carried out by Jin et al. showed that TMAO confers detrimental effects on adult cardiomyocytes by inducing T-tubule network damage and Ca handling dysfunction [97]. When TMAO was administered to HF mice, Organ et al. found that mouse cardiac function deteriorated significantly, characterized by pulmonary oedema, cardiac enlargement, and decreased ejection fraction [98]. Schuett et al. proved that TMAO could enhance patient susceptibility to HF by increasing myocardial fibrosis [99]. Likewise, Wang and his team proved that 3,3-dimethyl-1-butanol (DMB) ameliorates adverse cardiac structural remodelling in overload-induced HF mice by downregulating TMAO levels [100]. Given the critical role of TMAO in HF, it may serve as a potential therapeutic target.

**3.5. Arrhythmia.** Arrhythmia, including atrial fibrillation (AF), ventricular arrhythmia (VA), and atrioventricular block, is emerging as intractable CVD that contributes to heart failure or sudden cardiac death. Up-to-date studies have shown that anticancer therapies may induce cardiotoxicities, such as corrected QT interval prolongation and arrhythmia [101]. Additionally, Vahdatpour et al. found that atrial arrhythmia can be secondary to chronic lung disease-associated pulmonary hypertension [102]. Due to its prevalence and accompanying adverse events, investigation about arrhythmia has deepened, and we are now looking at the implications between gut microbiota and arrhythmia.

Zuo et al. previously identified variable metabolic patterns as well as imbalanced gut microbiota composition in patients with AF in which *Ruminococcus*, *Streptococcus*, and *Enterococcus* significantly increased while *Faecalibacterium*, *Alistipes*, *Oscillibacter*, and *Bilophila* obviously reduced [103]. Later, they found that patients with persistent AF (psAF) shared a great proportion of common features of gut microbiota dysbiosis [104]. In their latest study, the fecal microbiota from patients with psAF and those with paroxysmal AF were investigated, verifying a similar pattern of gut microbiota, with similar ratios of *Firmicutes* to *Bacteroidetes* [105].

Svingen et al. conducted a study in thousands of patients with suspected stable angina and proposed that plasma TMAO levels are definitely related to AF [106]. It is well known that thrombi can easily take place in the left atrial appendage of patients with AF, which then leads to embolism. Gong et al. found that in patients with AF, elevated TMAO levels are related to thrombus formation, manifested as platelet hyperreactivity [107]. It has been confirmed that the cardiac autonomic nervous system (CANS) can regulate the pathophysiology of AF or VA [108]. Meng et al. first proposed that preserving dysbacteriosis or modulating metabolites such as TMAO may be a target to treat arrhythmia due to the ability of TMAO to stimulate CANS and deteriorate ischaemia-induced VA by releasing proinflammatory markers such as IL-1 $\beta$  and TNF- $\alpha$  [109]. Similarly, according to the experiment of Yu et al., gut microbes have the ability to counteract AF progression by producing TMAO and can thus activate CANS in a rapid atrial pacing-induced canine AF model [110]. Likewise, in a propionate-treated hypertensive mouse model, the susceptibility to cardiac ventricular arrhythmias was significantly reduced, indicating



possible links between SCFAs and arrhythmia development [22]. Although the connection between gut microbiota and arrhythmia has been established, the precise underlying mechanisms still await further investigation (Table 2).

#### 4. Microorganism-Targeted Therapies

**4.1. Fecal Microbiota Transplantation.** As an effective approach to directly introduce intestinal flora, fecal microbiota transplantation (FMT) has gained much attention. The therapeutic value of FMT in gastrointestinal diseases, neurological and psychiatric disorders, and immunology regulation has been extensively examined [22, 111, 112]. However, studies concerning its application in CVDs are limited. Although oral supplementation of resveratrol has been proven to improve glucose homeostasis by altering gut microbiota, in the work of Kim and his colleagues [113], FMT from resveratrol-fed mice to obese mice was found to yield better results than oral administration of resveratrol alone, indicating that FMT is more straightforward and direct. Moreover, Hu et al. showed that FMT could abolish the increased proportion of *Firmicutes/Bacteroidetes*, diminish inflammatory infiltration in cardiomyocytes, and thereby attenuate myocarditis in mice [5]. However, in a double-blind trial involving 20 patients, the composition of intestinal flora was altered in the recipients after FMT from vegetarians, whereas the vasculitis indicators presented no improvement [114]. There are also disadvantages to FMT. For instance, endotoxins are transferred along with the donor microbiome. How to weigh the pros and cons of actual practice is still an issue to be addressed. To guarantee the reliable and smooth application of FMT in clinical use, the establishment of stool banks is on its way.

**4.2. Probiotic Administration.** Among the numerous bacteria residing in the host intestine, some are beneficial. An extra boost of these bacteria would probably bring positive results, thus leading to the application of probiotics. In a meta-analysis involving 846 individuals with hypertension, mild reductions in blood pressure, body mass index (BMI), and blood glucose levels were observed after probiotic administration, supporting the beneficial role of probiotics in blood pressure control [115]. Similarly, in other studies with spontaneously hypertensive rats, the probiotics *Bifidobacterium breve* and *Lactobacillus fermentum* were found to elicit antihypertensive effects by restoring gut microbiota balance and preventing endothelial dysfunction [116], whereas long-term supplementation with kefir ameliorated high blood pressure via improvement in intestinal integrity [117]. Moreover, in apoE<sup>-/-</sup> mice fed with HFD, supplementation with *Lactobacillus rhamnosus* GR-1 markedly reduced atherosclerotic lesion size by alleviating oxidative stress and inflammation [118]. Likewise, *Lactobacillus plantarum* ZDY04 has been shown to downregulate serum TMAO levels, which is a critical factor contributing to atherosclerosis development [119].

**4.3. Herbal Medicine.** Traditional Chinese medicine (TCM), which mainly utilizes herbs and their extracts, has recently been demonstrated to treat CVDs via intestinal microbial modulation. Ou et al. reviewed and summarized the mecha-

nisms of gut flora in TCM's theory of "stasis of intermingled phlegm and blood stasis" [120]. For example, the fact that TMAO promotes thrombosis might be one of the major causes of CVDs [121]. Anlu et al. showed that berberine originating from the Chinese herb *Coptis chinensis* has the ability to regulate the "microbiota-metabolism-immunity" axis [122]. Moreover, resveratrol derived from *Polygonum cuspidatum* was demonstrated to attenuate TMAO-induced atherosclerosis in apoE<sup>-/-</sup> mice by remodelling microbiota as well as decreasing TMAO and BA levels [123]. In addition, Ghosh et al. found that curcumin, a phytochemical component of *Curcuma longa*, attenuates atherosclerosis in LDLR<sup>-/-</sup> mice by regulating intestinal barrier function [124]. Anwar et al. showed that Trigonelline, which is purified from the seeds of *Trigonella foenum-graecum*, can inhibit the growth of *Citrobacter freundii* and subsequently decrease the production of TMAO in mice [125].

#### 5. Conclusion

Evidence from a compilation of studies of animals and humans indicates that the implications of gut microbiota and their metabolites in CVDs are well established. With high-throughput technologies, verification of the intestinal flora composition and in-depth mechanistic exploration are accessible. However, the links between gut microbiota and disease development are so complex that they involve immune regulation, the inflammatory response, gut barrier integrity, metabolic homeostasis, etc. Further investigations into the specific mechanisms are needed, which then share the possibility of being transferred into clinical practice.

#### Conflicts of Interest

The authors declare no conflicts of interest, financial or otherwise.

#### Authors' Contributions

MZ conceived of and designed the study and revised the manuscript for important intellectual content; XZ performed the literature search. YC generated the figures and tables; MN performed the background research. MZ and PZ edited the manuscript. WZ and YC drafted the manuscript. All authors have read and approved the content of the manuscript. Wenyi Zhou, Yiyu Cheng, and Ping Zhu contributed equally to this work.

#### Acknowledgments

This research was funded by the National Key Research and Development Program of China (2018YFA0108700), the NSFC Projects of International Cooperation and Exchanges (81720102004), the National Natural Science Foundation of China (81974019, 81970248), and the National Training Program of Innovation and Entrepreneurship for Undergraduates (2020105330125). AJE edited the manuscript for grammar, punctuation and spelling.

## References

- [1] J. M. Brown and S. L. Hazen, "Microbial modulation of cardiovascular disease," *Nature Reviews Microbiology*, vol. 16, no. 3, pp. 171–181, 2018.
- [2] W. H. Tang, T. Kitai, and S. L. Hazen, "Gut microbiota in cardiovascular health and disease," *Circulation Research*, vol. 120, no. 7, pp. 1183–1196, 2017.
- [3] W. Tang, D. Y. Li, and S. L. Hazen, "Dietary metabolism, the gut microbiome, and heart failure," *Nature Reviews Cardiology*, vol. 16, no. 3, pp. 137–154, 2019.
- [4] K. Kiouptsi and C. Reinhardt, "Contribution of the commensal microbiota to atherosclerosis and arterial thrombosis," *British journal of pharmacology*, vol. 175, no. 24, pp. 4439–4449, 2018.
- [5] T. T. Kim, N. Parajuli, M. M. Sung et al., "Fecal transplant from resveratrol-fed donors improves glycaemia and cardiovascular features of the metabolic syndrome in mice," *American Journal of Physiology-Endocrinology and Metabolism*, vol. 315, no. 4, pp. E511–E519, 2018.
- [6] J. Liu, T. Zhang, Y. Wang et al., "Baicalin ameliorates neuropathology in repeated cerebral ischemia-reperfusion injury model mice by remodeling the gut microbiota," *Aging (Albany NY)*, vol. 12, no. 4, pp. 3791–3806, 2020.
- [7] S. Wan, Y. Nie, Y. Zhang, C. Huang, and X. Zhu, "Gut microbial dysbiosis is associated with profibrotic factors in liver fibrosis mice," *Frontiers in cellular and infection microbiology*, vol. 10, p. 18, 2020.
- [8] M. Jin, Z. Qian, J. Yin, W. Xu, and X. Zhou, "The role of intestinal microbiota in cardiovascular disease," *Journal of Cellular and Molecular Medicine*, vol. 23, no. 4, pp. 2343–2350, 2019.
- [9] F. Z. Marques, C. R. Mackay, and D. M. Kaye, "Beyond gut feelings: how the gut microbiota regulates blood pressure," *Nature Reviews Cardiology*, vol. 15, no. 1, pp. 20–32, 2018.
- [10] Y. Li, H. S. Faden, and L. Zhu, "The response of the gut microbiota to dietary changes in the first two years of life," *Frontiers in Pharmacology*, vol. 11, p. 334, 2020.
- [11] F. Fava, L. Rizzetto, and K. M. Tuohy, "Gut microbiota and health: connecting actors across the metabolic system," *Proceedings of the Nutrition Society*, vol. 78, no. 2, pp. 177–188, 2019.
- [12] W. H. W. Tang, F. Bäckhed, U. Landmesser, and S. L. Hazen, "Intestinal microbiota in cardiovascular health and disease: JACC State-of-the-Art Review," *Journal of the American College of Cardiology*, vol. 73, no. 16, pp. 2089–2105, 2019.
- [13] P. B. Eckburg, E. M. Bik, C. N. Bernstein et al., "Diversity of the human intestinal microbial flora," *Science*, vol. 308, no. 5728, pp. 1635–1638, 2005.
- [14] S. R. Gill, M. Pop, R. T. DeBoy et al., "Metagenomic analysis of the human distal gut microbiome," *Science*, vol. 312, no. 5778, pp. 1355–1359, 2006.
- [15] A. Koliada, G. Syzenko, V. Moseiko et al., "Association between body mass index and Firmicutes/Bacteroidetes ratio in an adult Ukrainian population," *BMC Microbiology*, vol. 17, no. 1, pp. 1–6, 2017.
- [16] C. Indiani, K. F. Rizzardi, P. M. Castelo, L. F. C. Ferraz, M. Darrieux, and T. M. Parisotto, "Childhood obesity and Firmicutes/Bacteroidetes ratio in the gut microbiota: a systematic review," *Childhood Obesity*, vol. 14, no. 8, pp. 501–509, 2018.
- [17] A. Pascale, N. Marchesi, S. Govoni, A. Coppola, and C. Gazzaruso, "The role of gut microbiota in obesity, diabetes mellitus, and effect of metformin: new insights into old diseases," *Current Opinion in Pharmacology*, vol. 49, pp. 1–5, 2019.
- [18] P. V. Chang, L. Hao, S. Offermanns, and R. Medzhitov, "The microbial metabolite butyrate regulates intestinal macrophage function via histone deacetylase inhibition," *Proceedings of the National Academy of Sciences*, vol. 111, no. 6, pp. 2247–2252, 2014.
- [19] L. Macia, J. Tan, A. T. Vieira et al., "Metabolite-sensing receptors GPR43 and GPR109A facilitate dietary fibre-induced gut homeostasis through regulation of the inflammasome," *Nature communications*, vol. 6, no. 1, p. 6734, 2015.
- [20] J. Tan, C. McKenzie, M. Potamitis, A. N. Thorburn, C. R. Mackay, and L. Macia, "The role of short-chain fatty acids in health and disease," *Advances in Immunology*, vol. 121, pp. 91–119, 2014.
- [21] P. M. Smith, M. R. Howitt, N. Panikov et al., "The microbial metabolites, short-chain fatty acids, regulate colonic Treg cell homeostasis," *Science*, vol. 341, no. 6145, pp. 569–573, 2013.
- [22] H. Bartolomeaus, A. Balogh, M. Yakoub et al., "Short-chain fatty acid propionate protects from hypertensive cardiovascular damage," *Circulation*, vol. 139, no. 11, pp. 1407–1421, 2019.
- [23] M. H. Kim, S. G. Kang, J. H. Park, M. Yanagisawa, and C. H. Kim, "Short-chain fatty acids activate GPR41 and GPR43 on intestinal epithelial cells to promote inflammatory responses in mice," *Gastroenterology*, vol. 145, no. 2, pp. 396–406.e10, 2013.
- [24] G. G. Schiattarella, A. Sannino, E. Toscano et al., "Gut microbe-generated metabolite trimethylamine-N-oxide as cardiovascular risk biomarker: a systematic review and dose-response meta-analysis," *European Heart Journal*, vol. 38, no. 39, pp. 2948–2956, 2017.
- [25] R. Zhuang, X. Ge, L. Han et al., "Gut microbe-generated metabolite trimethylamine-N-oxide and the risk of diabetes: a systematic review and dose-response meta-analysis," *Obesity Reviews*, vol. 20, no. 6, pp. 883–894, 2019.
- [26] O. Manor, N. Zubair, M. P. Conomos et al., "A multi-omic association study of trimethylamine N-oxide," *Cell Reports*, vol. 24, no. 4, pp. 935–946, 2018.
- [27] X. Sun, X. Jiao, Y. Ma et al., "Trimethylamine N-oxide induces inflammation and endothelial dysfunction in human umbilical vein endothelial cells via activating ROS-TXNIP-NLRP3 inflammasome," *Biochemical and Biophysical Research Communications*, vol. 481, no. 1-2, pp. 63–70, 2016.
- [28] C. Yue, X. Yang, J. Li et al., "Trimethylamine N-oxide prime NLRP3 inflammasome via inhibiting ATG16L1-induced autophagy in colonic epithelial cells," *Biochemical and Biophysical Research Communications*, vol. 490, no. 2, pp. 541–551, 2017.
- [29] M. M. Seldin, Y. Meng, H. Qi et al., "Trimethylamine N-oxide promotes vascular inflammation through signaling of mitogen-activated protein kinase and nuclear Factor- $\kappa$ B," *Journal of the American Heart Association*, vol. 5, no. 2, 2016.
- [30] A. Kriaa, M. Bourgin, A. Potiron et al., "Microbial impact on cholesterol and bile acid metabolism: current status and future prospects," *Journal of Lipid Research*, vol. 60, no. 2, pp. 323–332, 2019.
- [31] R. Villette, P. Kc, S. Beliard et al., "Unraveling host-gut microbiota dialogue and its impact on cholesterol levels," *Frontiers in Pharmacology*, vol. 11, 2020.

- [32] D. J. Kenny, D. R. Plichta, D. Shungin et al., "Cholesterol metabolism by uncultured human gut bacteria influences host cholesterol level," *Cell Host & Microbe*, vol. 28, no. 2, pp. 245–257.e6, 2020.
- [33] J. M. Ridlon, S. C. Harris, S. Bhowmik, D. J. Kang, and P. B. Hylemon, "Consequences of bile salt biotransformations by intestinal bacteria," *Gut Microbes*, vol. 7, no. 1, pp. 22–39, 2016.
- [34] S. A. Joyce and C. G. Gahan, "Disease-associated changes in bile acid profiles and links to altered gut microbiota," *Digestive Diseases*, vol. 35, no. 3, pp. 169–177, 2017.
- [35] H. Wang, J. D. Latorre, M. Bansal et al., "Microbial metabolite deoxycholic acid controls *Clostridium perfringens*-induced chicken necrotic enteritis through attenuating inflammatory cyclooxygenase signaling," *Scientific Reports*, vol. 9, no. 1, article 14541, 2019.
- [36] K. S. Michelsen, M. H. Wong, P. K. Shah et al., "Lack of Toll-like receptor 4 or myeloid differentiation factor 88 reduces atherosclerosis and alters plaque phenotype in mice deficient in apolipoprotein E," *Proceedings of the National Academy of Sciences*, vol. 101, no. 29, pp. 10679–10684, 2004.
- [37] A. E. Mullick, P. S. Tobias, and L. K. Curtiss, "Modulation of atherosclerosis in mice by Toll-like receptor 2," *The Journal of Clinical Investigation*, vol. 115, no. 11, pp. 3149–3156, 2005.
- [38] G. den Besten, A. Bleeker, A. Gerding et al., "Short-chain fatty acids protect against high-fat diet-induced obesity via a PPAR $\gamma$ -Dependent switch from lipogenesis to fat oxidation," *Diabetes*, vol. 64, no. 7, pp. 2398–2408, 2015.
- [39] M. Hernandez, E. E. Canfora, J. Jocken, and E. E. Blaak, "The short-chain fatty acid acetate in body weight control and insulin sensitivity," *Nutrients*, vol. 11, no. 8, p. 1943, 2019.
- [40] X. Tan, Y. Liu, J. Long et al., "Trimethylamine N-oxide aggravates liver steatosis through modulation of bile acid metabolism and inhibition of farnesoid X receptor signaling in nonalcoholic fatty liver disease," *Molecular Nutrition & Food Research*, vol. 63, no. 17, article e1900257, 2019.
- [41] A. Tirosh, E. S. Calay, G. Tuncman et al., "The short-chain fatty acid propionate increases glucagon and FABP4 production, impairing insulin action in mice and humans," *Science Translational Medicine*, vol. 11, no. 489, p. eaav0120, 2019.
- [42] A. Bronden and F. K. Knop, "Glucose-metabolic effects of pharmacotherapy-induced modulation of bile acid physiology," *The Journal of Clinical Endocrinology & Metabolism*, vol. 105, no. 1, pp. 362–373, 2020.
- [43] L. Ding, M. Chang, Y. Guo et al., "Trimethylamine-N-oxide (TMAO)-induced atherosclerosis is associated with bile acid metabolism," *Lipids in Health and Disease*, vol. 17, no. 1, p. 286, 2018.
- [44] L. Sheng, P. K. Jena, Y. Hu et al., "Hepatic inflammation caused by dysregulated bile acid synthesis is reversible by butyrate supplementation," *The Journal of pathology*, vol. 243, no. 4, pp. 431–441, 2017.
- [45] K. Saito, R. Suzuki, Y. Koyanagi, H. Isogai, H. Yoneyama, and E. Isogai, "Inhibition of enterohemorrhagic *Escherichia coli* O157:H7 infection in a gnotobiotic mouse model with pre-colonization by *Bacteroides* strains," *Biomedical Reports*, vol. 10, no. 3, pp. 175–182, 2019.
- [46] O. P. Mathew, K. Ranganna, J. Mathew et al., "Cellular effects of butyrate on vascular smooth muscle cells are mediated through disparate actions on dual targets, histone deacetylase (HDAC) Activity and PI3K/Akt Signaling Network," *International Journal of Molecular Sciences*, vol. 20, no. 12, p. 2902, 2019.
- [47] K. Kim, O. Kwon, T. Y. Ryu et al., "Propionate of a microbiota metabolite induces cell apoptosis and cell cycle arrest in lung cancer," *Molecular Medicine Reports*, vol. 20, no. 2, pp. 1569–1574, 2019.
- [48] N. Nie, C. Bai, S. Song, Y. Zhang, B. Wang, and Z. Li, "Bifidobacterium plays a protective role in TNF- $\alpha$ -induced inflammatory response in Caco-2 cell through NF- $\kappa$ B and p38MAPK pathways," *Molecular and Cellular Biochemistry*, vol. 464, no. 1-2, pp. 83–91, 2020.
- [49] C. J. Li and T. H. Elsasser, "Butyrate-induced apoptosis and cell cycle arrest in bovine kidney epithelial cells: involvement of caspase and proteasome pathways1," *Journal of Animal Science*, vol. 83, no. 1, pp. 89–97, 2005.
- [50] L. F. Iannucci, J. Sun, B. K. Singh et al., "Short chain fatty acids induce UCP2-mediated autophagy in hepatic cells," *Biochemical and biophysical research communications*, vol. 480, no. 3, pp. 461–467, 2016.
- [51] C. M. Qiao, M. F. Sun, X. B. Jia et al., "Sodium butyrate causes  $\alpha$ -synuclein degradation by an Atg5-dependent and PI3K/Akt/mTOR-related autophagy pathway," *Experimental Cell Research*, vol. 387, no. 1, article 111772, p. 111772, 2020.
- [52] J. Liu, Y. Wang, H. Meng et al., "Butyrate rather than LPS subverts gingival epithelial homeostasis by downregulation of intercellular junctions and triggering pyroptosis," *Journal of Clinical Periodontology*, vol. 46, no. 9, pp. 894–907, 2019.
- [53] P. Wu, J. Chen, J. Chen et al., "Trimethylamine N-oxide promotes apoE(-/-) mice atherosclerosis by inducing vascular endothelial cell pyroptosis via the SDHB/ROS pathway," *Journal of Cellular Physiology*, vol. 235, no. 10, pp. 6582–6591, 2020.
- [54] J. Gu, W. Huang, W. Zhang et al., "Sodium butyrate alleviates high-glucose-induced renal glomerular endothelial cells damage via inhibiting pyroptosis," *International Immunopharmacology*, vol. 75, article 105832, 2019.
- [55] H. Cohen, N. Baram, L. Edry-Botzer, A. Munitz, D. Salomon, and M. Gerlic, "Vibriopore-forming leukocidin activates pyroptotic cell death via the NLRP3 inflammasome," *Emerging Microbes & Infections*, vol. 9, no. 1, pp. 278–290, 2020.
- [56] R. S. Chapkin, S. L. Navarro, M. Hullar, and J. W. Lampe, "Diet and gut microbes act coordinately to enhance programmed cell death and reduce colorectal cancer risk," *Digestive Diseases and Sciences*, vol. 65, no. 3, pp. 840–851, 2020.
- [57] P. Lavtar, G. Rudolf, A. Maver et al., "Association of circadian rhythm genes ARNTL/BMAL1 and CLOCK with multiple sclerosis," *PloS One*, vol. 13, no. 1, article e0190601, 2018.
- [58] Z. Liu, Z. Y. Wei, J. Chen et al., "Acute sleep-wake cycle shift results in community alteration of human gut microbiome," *Mosphere*, vol. 5, no. 1, 2020.
- [59] A. Segers, L. Desmet, T. Thijs, K. Verbeke, J. Tack, and I. Depoortere, "The circadian clock regulates the diurnal levels of microbial short-chain fatty acids and their rhythmic effects on colon contractility in mice," *Acta Physiologica*, vol. 225, no. 3, article e13193, 2019.
- [60] F. Z. Marques, E. Nelson, P. Y. Chu et al., "High-fiber diet and acetate supplementation change the gut microbiota and prevent the development of hypertension and heart failure in hypertensive mice," *Circulation*, vol. 135, no. 10, pp. 964–977, 2017.

- [61] X. Wu, L. Chen, F. Zeb et al., "Clock-Bmal1 mediates MMP9 induction in acrolein-promoted atherosclerosis associated with gut microbiota regulation," *Environmental Pollution*, vol. 252, no. Part B, pp. 1455–1463, 2019.
- [62] I. H. Page, "The mosaic theory of arterial hypertension—its interpretation," *Perspectives in Biology and Medicine*, vol. 10, no. 3, pp. 325–333, 1967.
- [63] J. Li, F. Zhao, Y. Wang et al., "Gut microbiota dysbiosis contributes to the development of hypertension," *Microbiome*, vol. 5, no. 1, p. 14, 2017.
- [64] J. Huart, J. Leenders, B. Taminiau et al., "Gut microbiota and fecal levels of short-chain fatty acids differ upon 24-hour blood pressure levels in men," *Hypertension*, vol. 74, no. 4, pp. 1005–1013, 2019.
- [65] S. Kim, R. Goel, A. Kumar et al., "Imbalance of gut microbiome and intestinal epithelial barrier dysfunction in patients with high blood pressure," *Clinical Science*, vol. 132, no. 6, pp. 701–718, 2018.
- [66] J. L. Pluznick, R. J. Protzko, H. Gevorgyan et al., "Olfactory receptor responding to gut microbiota-derived signals plays a role in renin secretion and blood pressure regulation," *Proceedings of the National Academy of Sciences*, vol. 110, no. 11, pp. 4410–4415, 2013.
- [67] I. R. Barrows, A. Ramezani, and D. S. Raj, "Inflammation, immunity, and oxidative stress in hypertension—partners in crime?," *Advances in Chronic Kidney Disease*, vol. 26, no. 2, pp. 122–130, 2019.
- [68] A. W. C. Man, H. Li, and N. Xia, "Resveratrol and the interaction between gut microbiota and arterial remodelling," *Nutrients*, vol. 12, no. 1, p. 119, 2020.
- [69] X. Ge, L. Zheng, R. Zhuang et al., "The gut microbial metabolite trimethylamine N-oxide and hypertension risk: a systematic review and dose-response meta-analysis," *Advances in Nutrition*, vol. 11, no. 1, pp. 66–76, 2020.
- [70] J. Liu, T. Li, H. Wu et al., "Lactobacillus rhamnosus GG strain mitigated the development of obstructive sleep apnea-induced hypertension in a high salt diet via regulating TMAO level and CD4(+) T cell induced-type I inflammation," *Biomedicine & Pharmacotherapy*, vol. 112, article 108580, 2019.
- [71] M. Ufnal, R. Jazwiec, M. Dadlez, A. Drapala, M. Sikora, and J. Skrzypecki, "Trimethylamine-N-oxide: a carnitine-derived metabolite that prolongs the hypertensive effect of angiotensin II in rats," *Canadian Journal of Cardiology*, vol. 30, no. 12, pp. 1700–1705, 2014.
- [72] X. Yan, J. Jin, X. Su et al., "Intestinal flora modulates blood pressure by regulating the synthesis of intestinal-derived corticosterone in high salt-induced hypertension," *Circulation Research*, vol. 126, no. 7, pp. 839–853, 2020.
- [73] O. Koren, A. Spor, J. Felin et al., "Human oral, gut, and plaque microbiota in patients with atherosclerosis," *Proceedings of the National Academy of Sciences*, vol. 108, Supplement\_1, pp. 4592–4598, 2011.
- [74] Z. Jie, H. Xia, S. L. Zhong et al., "The gut microbiome in atherosclerotic cardiovascular disease," *Nature Communications*, vol. 8, no. 1, p. 845, 2017.
- [75] S. D. Wright, C. Burton, M. Hernandez et al., "Infectious agents are not necessary for murine atherogenesis," *The Journal of Experimental Medicine*, vol. 191, no. 8, pp. 1437–1442, 2000.
- [76] Z. Wang, E. Klipfell, B. J. Bennett et al., "Gut flora metabolism of phosphatidylcholine promotes cardiovascular disease," *Nature*, vol. 472, no. 7341, pp. 57–63, 2011.
- [77] K. Kasahara, K. A. Krautkramer, E. Org et al., "Interactions between Roseburia intestinalis and diet modulate atherogenesis in a murine model," *Nature Microbiology*, vol. 3, no. 12, pp. 1461–1471, 2018.
- [78] Y. Du, X. Li, C. Su et al., "Butyrate protects against high-fat diet-induced atherosclerosis via up-regulating ABCA1 expression in apolipoprotein E-deficiency mice," *British Journal of Pharmacology*, vol. 177, no. 8, pp. 1754–1772, 2020.
- [79] Z. He, W. Hao, E. Kwek et al., "Fish oil is more potent than flaxseed oil in modulating gut microbiota and reducing trimethylamine-N-oxide-exacerbated atherogenesis," *Journal of Agricultural and Food Chemistry*, vol. 67, no. 49, pp. 13635–13647, 2019.
- [80] R. A. Hasan, A. Y. Koh, and A. Zia, "The gut microbiome and thromboembolism," *Thrombosis Research*, vol. 189, pp. 77–87, 2020.
- [81] K. Huynh, "Novel gut microbiota-derived metabolite promotes platelet thrombosis via adrenergic receptor signalling," *Nature Reviews Cardiology*, vol. 17, no. 5, p. 265, 2020.
- [82] A. Lassiger-Herfurth, G. Pontarollo, A. Grill, and C. Reinhardt, "The gut microbiota in cardiovascular disease and arterial thrombosis," *Microorganisms*, vol. 7, no. 12, p. 691, 2019.
- [83] Z. X. Wu, S. F. Li, H. Chen et al., "The changes of gut microbiota after acute myocardial infarction in rats," *PLoS One*, vol. 12, no. 7, article e0180717, 2017.
- [84] X. Zhou, J. Li, J. Guo et al., "Gut-dependent microbial translocation induces inflammation and cardiovascular events after ST-elevation myocardial infarction," *Microbiome*, vol. 6, no. 1, p. 66, 2018.
- [85] A. Haghikia, X. S. Li, T. G. Liman et al., "Gut microbiota-dependent trimethylamine N-oxide predicts risk of cardiovascular events in patients with stroke and is related to proinflammatory monocytes," *Arteriosclerosis, Thrombosis, and Vascular Biology*, vol. 38, no. 9, pp. 2225–2235, 2018.
- [86] T. Tang, H. C. Chen, C. Y. Chen et al., "Loss of gut microbiota alters immune system composition and cripples postinfarction cardiac repair," *Circulation*, vol. 139, no. 5, pp. 647–659, 2019.
- [87] V. Lam, J. Su, S. Koprowski et al., "Intestinal microbiota determine severity of myocardial infarction in rats," *The FASEB Journal*, vol. 26, no. 4, pp. 1727–1735, 2011.
- [88] P. Ponikowski, A. A. Voors, S. D. Anker et al., "016 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure: The Task Force for the diagnosis and treatment of acute and chronic heart failure of the European Society of Cardiology (ESC) Developed with the special contribution of the Heart Failure Association (HFA) of the ESC," *European Heart Journal*, vol. 18, no. 8, pp. 891–975, 2016.
- [89] Y. Heianza, W. Ma, J. E. Manson, K. M. Rexrode, and L. Qi, "Gut microbiota metabolites and risk of major adverse cardiovascular disease events and death: a systematic review and meta-analysis of prospective studies," *Journal of the American Heart Association*, vol. 6, no. 7, 2017.
- [90] J. Peng, X. Xiao, M. Hu, and X. Zhang, "Interaction between gut microbiome and cardiovascular disease," *Life Sciences*, vol. 214, pp. 153–157, 2018.

- [91] E. Pasini, R. Aquilani, C. Testa et al., "Pathogenic gut flora in patients with chronic heart failure," *JACC: Heart Failure*, vol. 4, no. 3, pp. 220–227, 2016.
- [92] Q. Jia, H. Li, H. Zhou et al., "Role and effective therapeutic target of gut microbiota in heart failure," *Cardiovascular Therapeutics*, vol. 2019, Article ID 5164298, 10 pages, 2019.
- [93] T. T. Tang, J. Yuan, Z. F. Zhu et al., "Regulatory T cells ameliorate cardiac remodeling after myocardial infarction," *Basic Research in Cardiology*, vol. 107, no. 1, pp. 232–232, 2012.
- [94] Y. Nagatomo and W. H. Tang, "Intersections between microbiome and heart failure: revisiting the gut hypothesis," *Journal of Cardiac Failure*, vol. 21, no. 12, pp. 973–980, 2015.
- [95] A. Zabell and W. H. Tang, "Targeting the microbiome in heart failure," *Current Treatment Options in Cardiovascular Medicine*, vol. 19, no. 4, p. 27, 2017.
- [96] M. Savi, L. Bocchi, L. Bresciani et al., "Trimethylamine-N-oxide (TMAO)-induced impairment of cardiomyocyte function and the protective role of urolithin B-glucuronide," *Molecules*, vol. 23, no. 3, p. 549, 2018.
- [97] B. Jin, F. Ji, A. Zuo et al., "Destructive role of TMAO in T-tubule and excitation-contraction coupling in the adult cardiomyocytes," *International Heart Journal*, vol. 61, no. 2, pp. 355–363, 2020.
- [98] C. L. Organ, H. Otsuka, S. Bhushan et al., "Choline diet and its gut microbe-derived metabolite, trimethylamine N-oxide, exacerbate pressure overload-induced heart failure," *Circulation: Heart Failure*, vol. 9, no. 1, article e002314, 2016.
- [99] K. Schuett, M. E. Kleber, H. Scharnagl et al., "Trimethylamine-N-oxide and heart failure with reduced versus preserved ejection fraction," *Journal of the American College of Cardiology*, vol. 70, no. 25, pp. 3202–3204, 2017.
- [100] G. Wang, B. Kong, W. Shuai, H. Fu, X. Jiang, and H. Huang, "3,3-Dimethyl-1-butanol attenuates cardiac remodeling in pressure-overload-induced heart failure mice," *The Journal of Nutritional Biochemistry*, vol. 78, p. 108341, 2020.
- [101] J. Herrmann, "Adverse cardiac effects of cancer therapies: cardiotoxicity and arrhythmia," *Nature Reviews Cardiology*, vol. 17, no. 8, pp. 474–502, 2020.
- [102] C. A. Vahdatpour, J. J. Luebbert, and H. I. Palevsky, "Atrial arrhythmias in chronic lung disease-associated pulmonary hypertension," *Pulmonary Circulation*, vol. 10, no. 1, article 204589402091068, 2020.
- [103] K. Zuo, J. Li, K. Li et al., "Disordered gut microbiota and alterations in metabolic patterns are associated with atrial fibrillation," *GigaScience*, vol. 8, no. 6, 2019.
- [104] K. Zuo, J. Li, P. Wang et al., "Duration of persistent atrial fibrillation is associated with alterations in human gut microbiota and metabolic phenotypes," *Msystems*, vol. 4, no. 6, 2019.
- [105] K. Zuo, X. Yin, K. Li et al., "Different types of atrial fibrillation share patterns of gut microbiota dysbiosis," *Msphere*, vol. 5, no. 2, 2020.
- [106] G. Svingen, H. Zuo, P. M. Ueland et al., "Increased plasma trimethylamine-N-oxide is associated with incident atrial fibrillation," *International Journal of Cardiology*, vol. 267, pp. 100–106, 2018.
- [107] D. Gong, L. Zhang, Y. Zhang, F. Wang, Z. Zhao, and X. Zhou, "Gut microbial metabolite trimethylamine N-oxide is related to thrombus formation in atrial fibrillation patients," *The American Journal of the Medical Sciences*, vol. 358, no. 6, pp. 422–428, 2019.
- [108] K. Shivkumar, O. A. Ajijola, I. Anand et al., "Clinical neurocardiology defining the value of neuroscience-based cardiovascular therapeutics," *The Journal of Physiology*, vol. 594, no. 14, pp. 3911–3954, 2016.
- [109] G. Meng, X. Zhou, M. Wang et al., "Gut microbe-derived metabolite trimethylamine N-oxide activates the cardiac autonomic nervous system and facilitates ischemia-induced ventricular arrhythmia via two different pathways," *EBioMedicine*, vol. 44, pp. 656–664, 2019.
- [110] L. Yu, G. Meng, B. Huang et al., "A potential relationship between gut microbes and atrial fibrillation: trimethylamine N-oxide, a gut microbe-derived metabolite, facilitates the progression of atrial fibrillation," *International Journal of Cardiology*, vol. 255, pp. 92–98, 2018.
- [111] H. Antushevich, "Fecal microbiota transplantation in disease therapy," *Clinica Chimica Acta*, vol. 503, pp. 90–98, 2020.
- [112] F. Zhang, T. Zhang, H. Zhu, and T. J. Borody, "Evolution of fecal microbiota transplantation in methodology and ethical issues," *Current Opinion in Pharmacology*, vol. 49, pp. 11–16, 2019.
- [113] P. F. de Groot, M. N. Frissen, N. C. de Clercq, and M. Nieuwdorp, "Fecal microbiota transplantation in metabolic syndrome: History, present and future," *Gut Microbes*, vol. 8, no. 3, pp. 253–267, 2017.
- [114] X. F. Hu, W. Y. Zhang, Q. Wen et al., "Fecal microbiota transplantation alleviates myocardial damage in myocarditis by restoring the microbiota composition," *Pharmacological Research*, vol. 139, pp. 412–421, 2019.
- [115] L. P. Smits, R. S. Kootte, E. Levin et al., "Effect of vegan fecal microbiota transplantation on carnitine- and choline-derived trimethylamine-N-oxide production and vascular inflammation in patients with metabolic syndrome," *Journal of the American Heart Association*, vol. 7, no. 7, 2018.
- [116] C. Chi, C. Li, D. Wu et al., "Effects of probiotics on patients with hypertension: a systematic review and meta-analysis," *Current Hypertension Reports*, vol. 22, no. 5, 2020.
- [117] I. Robles-Vera, M. Toral, N. la Visitación et al., "Probiotics prevent dysbiosis and the rise in blood pressure in genetic hypertension: role of short-chain fatty acids," *Molecular Nutrition & Food Research*, vol. 64, no. 6, article 1900616, 2020.
- [118] S. M. de Almeida, F. E. Mowry, S. C. Peaden, T. U. Andrade, and V. C. Biancardi, "Kefir ameliorates hypertension via gut-brain mechanisms in spontaneously hypertensive rats," *The Journal of Nutritional Biochemistry*, vol. 77, 2020.
- [119] Y. Fang, H. Q. Chen, X. Zhang et al., "Probiotic administration of lactobacillus rhamnosus GR-1 attenuates atherosclerotic plaque formation in ApoE<sup>-/-</sup> mice fed with a high-fat diet," *European Review for Medical and Pharmacological Sciences*, vol. 23, no. 8, pp. 3533–3541, 2019.
- [120] L. Qiu, X. Tao, H. Xiong, J. Yu, and H. Wei, "Lactobacillus plantarum ZDY04 exhibits a strain-specific property of lowering TMAO via the modulation of gut microbiota in mice," *Food & Function*, vol. 9, no. 8, pp. 4299–4309, 2018.
- [121] Y. Ou, C. Zhang, M. Yao, and L. Wang, "Gut Flora: Novel therapeutic target of Chinese medicine for the treatment of cardiovascular diseases," *Evidence-Based Complementary and Alternative Medicine*, vol. 2019, Article ID 3719596, 7 pages, 2019.
- [122] W. Anlu, C. Dongcheng, Z. He et al., "Using herbal medicine to target the "microbiota-metabolism-immunity" axis as possible therapy for cardiovascular disease," *Pharmacological Research*, vol. 142, pp. 205–222, 2019.

- [123] M. L. Chen, L. Yi, Y. Zhang et al., “Resveratrol attenuates trimethylamine-N-oxide (TMAO)-induced atherosclerosis by regulating TMAO synthesis and bile acid metabolism via remodeling of the gut microbiota,” *MBio*, vol. 7, no. 2, pp. e02210–e02215, 2016.
- [124] S. S. Ghosh, J. Bie, J. Wang, and S. Ghosh, “Oral supplementation with non-absorbable antibiotics or curcumin attenuates western diet-induced atherosclerosis and glucose intolerance in LDLR<sup>-/-</sup> mice—role of intestinal permeability and macrophage activation,” *PLoS One*, vol. 9, no. 9, article e108577, 2014.
- [125] S. Anwar, U. Bhandari, B. P. Panda, K. Dubey, W. Khan, and S. Ahmad, “Trigonelline inhibits intestinal microbial metabolism of choline and its associated cardiovascular risk,” *Journal of Pharmaceutical and Biomedical Analysis*, vol. 159, pp. 100–112, 2018.