

⁶ Tanigaki, N., Nakamura, K., Natori, T., Kreiter, V. P. & Pressman, D. *Transplantation* **18**, 74-78 (1974).
⁷ Sanderson, A. R. & Welsh, K. I. *Transplant. Proc.* **5**, 471-475 (1973).
⁸ Sanderson, A. R. & Welsh, K. I. *Transplantation* **16**, 304-312 (1973).
⁹ Metzgar, R. S. & Miller, J. L. *Transplantation* **13**, 467-471 (1972).
¹⁰ Farr, R. S. *J. infect. Dis.* **103**, 239-251 (1958).
¹¹ Rajewsky, K. & Rottländer, E. *Cold Spring Harb. Symp. quant. Biol.* **32**, 547-555 (1967).
¹² Mitchison, N. A. *Cold Spring Harb. Symp. quant. Biol.* **32**, 431-440 (1967).
¹³ Welsh, K. I., Burgos, H. & Batchelor, J. R. *Eur. J. Immun.* (in the press).
¹⁴ Lake, P. & Mitchison, N. A. *Cold Spring Harb. Symp. quant. Biol.* (in the press).
¹⁵ Nisbet, N. W., Simonsen, M. & Zaleski, M. *J. exp. Med.* **129**, 459-467 (1969).
¹⁶ Wilson, D. B., Blyth, J. L. & Nowell, P. C. *J. exp. Med.* **128**, 1157-1181 (1968).
¹⁷ Sanderson, A. R. & Welsh, K. I. *Transplant. Proc.* **6**, 163-167 (1974).
¹⁸ Peck, A. B., Alter, B. J. & Lindahl, K. F. *Transplant. Rev.* **29**, 189-221 (1976).
¹⁹ Zinkernagel, R. M. & Doherty, P. C. *Nature* **248**, 701-702 (1974).

In a radiation chimaera, host H-2 antigens determine immune responsiveness of donor cytotoxic cells

CELL membrane structures controlled by genes in the major histocompatibility complex (H-2 in mice) are involved in most immune interactions between T lymphocytes and other cells¹. Cytotoxic T lymphocytes (CTL) immunised against viruses², haptens³, minor histocompatibility antigens⁴ or tumour antigens⁵, are specific for self H-2 antigens as well as for the foreign antigen. But CTL are not restricted to recognising antigens in combination with only self H-2. H-2^d homozygous CTL which have matured in an irradiated H-2^d/H-2^k host can respond to antigen plus H-2^k in addition to antigen plus H-2^d (refs 6-8). It is not known whether the H-2 environment in which T cells mature influences their range of specificity, that is, whether CTL from a normal mouse can respond quantitatively as well to antigen plus foreign H-2 as they do to antigen plus self H-2. These experiments were designed to test this influence. The results suggest that host H-2 antigens do exert an effect on the specificity of T-cell responses.

A single suspension of bone marrow cells from F₁(BALB/c × BALB.B) (F₁(C × C.B), H-2^d/H-2^b) mice was used to reconstitute groups of lethally irradiated parental mice, C(H-2^d) ([F₁ → C] chimaeras) and C.B(H-2^b) ([F₁ → C.B] chimaeras). Eight weeks later these chimaeric mice and normal F₁(C × C.B) mice were primed against minor H antigens by injecting 8 × 10⁶ F₁(B10 × B10.D2) (H-2^b/H-2^d) spleen cells. The B10 background offers more than 30 minor histocompatibility antigenic differences that can be recognised by BALB mice⁹⁻¹⁰. Some weeks later the primed spleen cells were boosted in culture with irradiated F₁(B10 × B10.D2) stimulator cells and assayed for cytotoxicity 5 d later.

Following this immunisation procedure, cells from normal F₁(C × C.B) mice lysed B10 targets and B10.D2 targets almost equally (Table 1). (The two activities are mediated by separate

pools of CTL²⁻⁴) The chimaeras responded differently. In the same conditions of immunisation with F₁(B10 × B10.D2) cells, they responded preferentially to the minor antigens in association with the H 2 antigens of the host. CTL from the [F₁ → C] chimaeras killed B10.D2 targets better than B10 targets, whereas CTL from [F₁ → C.B] chimaeras lysed B10 targets better than B10.D2 targets (Table 1).

Spleen cells from five chimaeras were assayed for their content of host and donor cells at time of killing. Complement-mediated lysis with H-2^b anti-H-2^d serum and with H-2^d anti-H-2^b serum indicated that in all cases at least 85% of the cells were of F₁ (donor) origin. The cytotoxic effector cells were also lysed with anti-H-2 serum and complement just before the ⁵¹Cr-release assay (Table 2). Here, the [F₁ → C] chimaera cells lysed B10.D2 targets ninefold more efficiently than they lysed B10 targets (data not shown). The killer cells were treated with antiserum plus complement, washed, and assayed for lysis of labelled B10.D2. Table 2 shows, most importantly, that BALB/c anti-C57BL/6 (anti-H-2^b) serum reduced the cytotoxic activity 86% compared with controls. This antiserum does not lyse BALB/c effector cells. Therefore, at least 86% of the CTL were of F₁ bone marrow origin.

These experiments show that H-2^d/H-2^b cytotoxic cells which mature in an irradiated H-2^d host respond preferentially to antigens plus H-2^d, whereas H-2^d/H-2^b cells which mature in an irradiated H-2^b host respond preferentially to the same antigens in conjunction with H-2^b gene products. The experiments were designed to test the 1971 Jerne hypothesis¹⁴, or a modified version of it^{4,15}. The hypothesis accepts that a somatic theory of generation of receptor diversity is correct and proposes that self-H-2 antigens drive the diversity. Immature T cells first express an anti-self-H-2 receptor, leading to proliferation and to accumulation of V gene mutations until there is no significant reaction with self-H-2. According to this hypothesis, the receptor repertoire of A strain T cells which had matured in an A environment would be quite different from that of A strain T cells which had matured in a B environment. The results presented here are compatible with this hypothesis and with another theory of 'adaptive differentiation'¹.

There is an alternative explanation of the results. It may be that the host haplotype preference seen at the level of effector CTL does not reflect a bias in specificity at the level of precursor CTL. The H-2^d/H-2^b precursor CTL in the H-2^d host may have exactly the same range of reactivity as those in the H-2^b host. The haplotype preference of the effector CTL would then be due to the way antigen is presented to CTL precursors. Even though the immunogen (B10 minor antigens) was introduced on H-2 heterozygous cells, the antigen which was responsible for priming CTL may have been processed antigen presented on radiation resistant host cells⁴. In the [F₁ → C] chimaera such radiation-resistant antigen-present-

Table 1 Specificity of H-2^d/H-2^b cytotoxic cells from normal and chimaeric mice

Responder*	Immunised with†	% Specific lysis of targets‡			F ₁ (C × C.B) H-2 ^d /H-2 ^b	Ratio of lytic activity on§ B10/B10.D2
		B10 H-2 ^b	B10.D2 H-2 ^d	B10.BR H-2 ^k		
Experiment 1						
Normal F ₁ (C × C.B)(H-2 ^d /H-2 ^b)	F ₁ (B10 × B10.D2)(H-2 ^b /H-2 ^d)	56.2	62.2	1.5	ND	0.7
Chimaera [F ₁ → C]	F ₁ (B10 × B10.D2)(H-2 ^b /H-2 ^d)	19.3	72.7	2.9	ND	0.02
Chimaera [F ₁ → C.B]	F ₁ (B10 × B10.D2)(H-2 ^b /H-2 ^d)	51.8	29.1	0.1	ND	5.2
Normal F ₁ (C × C.B)	C3H(H-2 ^k)	4.5	3.8	71.5	ND	—
Experiment 2						
Normal F ₁ (C × C.B)	F ₁ (B10 × B10.D2)	54.8	68.4	ND	0.9	0.5
Chimaera [F ₁ → C]	F ₁ (B10 × B10.D2)	4.8	80.5	ND	1.5	<0.02
Chimaera [F ₁ → C.B]	F ₁ (B10 × B10.D2)	61.3	8.8	ND	0.1	>43.0

*Chimaeras were prepared as follows: BALB/c(C(H-2^d)) and BALB.B(C.B(H-2^b)) mice were irradiated with 850 R and reconstituted on the same day with 13.4 × 10⁶ anti-Thy-1 plus complement (C)-treated F₁(C × C.B) bone marrow cells. Eight weeks later they were immunised.

†Primed against minor H antigens by injecting 8 × 10⁶ viable F₁(B10 × B10.D2) spleen cells intraperitoneally. Spleen cell suspensions were prepared 4 weeks later (experiment 1) or 6 weeks later (experiment 2), and boosted for 5 d in culture with an equal number of 1,000 R irradiated F₁(B10 × B10.D2) or C3HeB/F₁ spleen cells as described previously¹¹.

‡2-d con A (conavalin A) blasts were labelled with ⁵¹Cr-sodium chromate and used as targets as described previously¹¹. Serial dilutions of the killers were assayed against a constant number of targets, and the figures for specific lysis presented here are for a killer: target ratio of 100:1. Spontaneous release of ⁵¹Cr varied from 18.1-22% in experiment 1 and from 9.6-13.7% in experiment 2.

§Calculated from the titrations of killers: targets^{12,13}. For example, in experiment 1, the [F₁ → C] cells lysed B10.D2 targets 50 times better than they lysed B10 since a 100:1 ratio caused 19.3% specific release from ⁵¹Cr-B10, whereas the same amount of specific lysis of ⁵¹Cr-B10.D2 was obtained at a ratio of 2:1. ND. Not determined.

Table 2 Sensitivity of chimaera cytotoxic cells to anti-H-2 serum plus C'

Cytotoxic cells*	Treatment of effector cells†	% Specific lysis of ⁵¹ Cr-B10.D2‡	% Reduction in lytic activity§
[F ₁ →C] anti-F ₁ (B10×B10.D2)	Medium	34.0	—
	Normal B10 serum + C'	34.3	0
	Anti-H-2 ^b + C'	9.8	86
	Anti-H-2 ^d + C'	5.1	94

*[F₁→C] chimaeras were primed *in vivo* 8 weeks after reconstitution, their spleen cells boosted in culture 20 weeks later and assayed for cytotoxicity on day 5 of culture.

†BALB/c anti-C57BL/6 and C57BL/6 anti-BALB/c sera were prepared by hyperimmunisation with spleen cells. Effector cells were incubated with mouse sera 1:2, washed, and incubated in guinea pig serum 1:9 as a source of complement (C'). Cells were re-suspended to the same volume and assayed.

‡Con A blasts from B10.D2 mice were labelled with ⁵¹Cr and used as target. Data presented are for original number of responder spleen cells:target cells of 7:1. Spontaneous release of ⁵¹Cr was 20.6%.

§Calculated from the titrations of killer:targets as in Table 1.

ing cells would be homozygous H-2^d and would naturally stimulate only anti-B10.D2(H-2^d) CTL, not anti-B10(H-2^b) CTL. Experiments to decide between these interpretations are in progress.

This work was supported by USPHS (grants A114269 and CA14051).

MICHAEL J. BEVAN

Center for Cancer Research and Department of Biology,
Massachusetts Institute of Technology,
Cambridge, Massachusetts 02139

Received 6 June; accepted 22 July 1977.

- Katz, D. H. *Lymphocyte Differentiation, Recognition and Function* (Academic, New York, 1977).
- Doherty, P. C., Blanden, R. V. & Zinkernagel, R. M. *Transplant. Rev.* **29**, 89-124 (1976).
- Shearer, G. M., Rehm, T. G. & Schmitt-Verhulst, A. M. *Transplant. Rev.* **29**, 222-248 (1976).
- Bevan, M. J. *Cold Spring Harbor Symp. Quant. Biol.* **41**, 519-527 (1976).
- Blank, K. J., Freedman, H. A. & Lilly, F. *Nature* **260**, 250-252 (1976).
- Pfeiffer, K. *Exp. Med.* **143**, 999-1004 (1976).
- Zinkernagel, R. M. *Nature* **261**, 139-141 (1976).
- von Boehmer, H. & Haas, W. *Nature* **261**, 141-142 (1976).
- Bailey, D. W. *Immunogenetics* **2**, 249-256 (1975).
- Staats, J. *Cancer Res.* **36**, 4333-4377 (1976).
- Bevan, M. J. *J. exp. Med.* **142**, 1349-1365 (1975).
- MacDonald, H. R., Phillips, R. A. & Miller, R. G. *J. Immun.* **111**, 565-574 (1973).
- Bevan, M. J. *J. Immun.* **114**, 316-319 (1975).
- Jernse, N. K. *Eur. J. Immun.* **1**, 9 (1973).
- Nahholz, M. & Migasawa, Y. in *Band T cells in Immune Recognition* (eds Loo, F. & Roelants, G. E.) (Wiley, Chichester, UK, 1977).

Cytotoxic T lymphocytes induced in mice by inactivated influenza virus vaccine

THERE is abundant evidence that T cell-mediated lysis of virus-infected target cells generally requires H-2K or H-2D region histocompatibility between cytotoxic T lymphocytes (CTL) and target cells¹⁻⁶. The biological basis of this requirement is still uncertain. Studies using chimaeric mice suggest that T lymphocytes are selectively sensitised by the viral and H-2 antigens expressed on the infected cell and that killing only occurs when the target cell shares H-2K or H-2D identity with the immunising virus-infected cell⁷. It is important to determine whether virus infection of the host tissue is, in fact, necessary for specific CTL immunity to develop. If this were so, the use of inactivated virus or purified viral antigens would not be suitable as vaccines against diseases in which CTL were required to afford optimal protection. We report here that inactivated influenza virus vaccines are, in fact, quite capable of evoking haemagglutinin specific CTL.

Epidemiological and experimental evidence indicate that protective immunity against influenza virus is principally directed against the virus haemagglutinin^{8,9}. We have recently described the development of haemagglutinin-specific CTL in influenza-infected mice¹⁰. This finding is consistent with these cells playing an important *in vivo* function in influenza immunity. Our finding contrasts with




Fig. 1 Specific immune lysis by spleen cells of BALB/c mice immunised intraperitoneally with 5,000 chick cell haemagglutinating units (HAU), (○), 500 HAU (▲) and 50 HAU (□) of A/Port Chalmers formalin-inactivated whole virus influenza vaccine. The spleen cells were tested at a 100:1 ratio on A/Port Chalmers-infected syngeneic kidney-derived cells in an 18-h ⁵¹Cr-release cytotoxicity assay as described previously¹⁰. The cytotoxic cells generated in vaccine-immunised mice were shown to be T cells by the removal of cytotoxic activity using anti-θ antiserum plus complement. Thus, whereas untreated spleen cells and spleen cells treated with complement alone achieved 42.3 and 37.0% specific lysis respectively, spleen cells treated with anti-θ antiserum plus complement achieved 7.9% specific lysis, which was not significant ($P > 0.5$).

recent studies which failed to demonstrate haemagglutinin-specific CTL in influenza-infected mice¹¹⁻¹³. These studies differed from our study, however, in the use as target cells of relatively non-permissive influenza-infected transformed cell lines rather than productively-infected normal tissue-derived target cells¹⁴. In the present study we have similarly tested mice inoculated with influenza vaccine, for haemagglutinin-specific CTL reactive with productively-infected target cells. The virus vaccines used contained an early H3N2 virus strain⁹, A/Aichi/68; a later virus of the same H3N2 subtype, but with a serologically distinguishable haemagglutinin, A/Port Chalmers/73; and an unrelated influenza B virus, B/Hong Kong/72. The target cells were syngeneic kidney-derived infected cells¹⁰. The experiment was carried out in both BALB/c mice (H-2^d) and C3H mice (H-2^k).

Figure 1 depicts the CTL response we observed in BALB/c mice inoculated intraperitoneally with various doses of A/Port Chalmers vaccine (virus vaccine supplied by Merrell-National Laboratories). Significant cytotoxic T-cell response was observed with each of the doses tested. To determine whether the response was restricted to target cells histocompatible with the immunised mice, we tested spleen cells from BALB/c and C3H immunised mice on syngeneic and allogeneic virus-infected target cells. As indicated in Table 1, the cytotoxic response was detectable only on the syngeneic virus-infected target cells. Specificity of the CTL for the haemagglutinin antigen of the immunising vaccine virus was shown in the experiments recorded in Table 2. Thus, CTL-distinguished target cells infected with the immunising strain H3N2 influenza virus, from target cells infected with a serologically different but related

Table 1 Virus specificity of cell-mediated lysis by CTL in influenza vaccine immunised mice

Vaccine virus used as immunogen	Specific immune lysis of target cells*		
	A/Port Chalmers-infected	A/Aichi-infected	B/Hong Kong-infected
A/Port Chalmers/73	96.6	6.2	8.8
A/Aichi/68	5.2	67.0	2.7
B/Hong Kong/72	-6.9	3.2	63.4

*Splenic lymphocytes were tested 8 d after intraperitoneal immunisation.